(A)

ZNF217 Mutations by HER2 Status

(B)

ZNF703 Mutations by HER2 Status

(C)

ZNF703 Mutations by HER2 Status

(A)

(B)

(C)

(A)

(B)

(A)

Expression of ZNF703 gene per cancer stage

(B)

ZNF703 Expression per HER2 Status by Cancer Stage

(A)

(B)

(A)

Correlation of ZNF217 Expression and Mutations

(B)

Correlation of ZNF703 Expression and Mutations

(C)

Correlation of ZNF750 Expression and Mutations

Sample numbers	AGE	Systolic	Diastolic	heart rate	Weight	BMI	Cancer Stage	HER2
201	65	142	94	94	59	22.5	2	Positive
203	30	101	64	75	70.66	27.8	3	Positive
205	51	137	92	79	75	23.68	3	Positive
207	72	172	85	71	50	23.14	2	Negative
209	66	147	82	100	70	30.30	2	Negative
211	45	133	96	82	84	32.81	1	Positive
213	43	150	87	64	84	31.42	2	Negative
215	44	129	83	96	75.5	25.52	3	Positive
217	41	116	60	86	74.4	29.06	2	Positive
219	36	114	72	75	67.3	26.26	3	Positive
221	53	123	86	81	78	31.96	2	Negative
223	58	113	67	98	73.8	27.95	2	Negative
225	53	147	83	96	70	28.8	1	Positive
227	60	150	92	75	79	30.9	1	Negative
229	54	119	81	80	101	37.8	2	Positive
231	72	134	91	74	71	27.7	2	Negative
233	44	144	94	88	63	25.6	2	Positive
235	49	120	84	90	98.5	34.9	3	Negative
237	66	124	55	78	63	24.9	3	Positive
239	53	112	67	78	61	21.1	1	Negative
241	58	133	88	71	90		2	Negative
243	51	137	83	119	70	27.7	3	Positive
245	66	153	87	118	66	24.8	3	Negative

Patient ID	Position	Substitution /Indel	Mutation Type	Impact Level	Amino Acid
201	37,697,439	C/T	Missense	MODERATE	Ser 177' Phe
	37,698,452	T/C	Synonymous	LOW	His 517'
205	37,698,165	C/G	Missense	MODERATE	Pro 422' Ala
215	37,698,266	G/T	Synonymous	LOW	Pro 455'
223	37,697,290	C/A	Missense	MODERATE	Ala 130' Asp
233	37,698,467	G/C	Synonymous	LOW	Pro 455'
	37,697,422	C/T	Missense	Moderate	Ser 174' Leu
237	37,698,415	GCGC CGCCGC/G	In-frame deletion	HIGH	Ala 514' del

Sample ID	Position	Mutation	Type	Impact	Amino acid
217	82,831,023	C/T	Missense	MODERATE	Val 478'> Ile
231	82,830,234	C/T	Missense	MODERATE	Asp 694' > Asn
241	82,831,982	C/T	Missense	MODERATE	Arg 158' Gln
243	82,831,023	C/T	Missense	MODERATE	Val 478' Ile

Sample ID	Position	Mutation	Type	Impact	Amino acid
203	82,830,589	T/C	Synonymous	LOW	Ala 575'
	82,831,280	T/C	Missense	MODERATE	Gln 392' > Arg
205	82,830,589	T/C	Synonymous	LOW	Ala 575'
207	82,830,589	T/C	Synonymous	LOW	Ala 575'
215	82,830,589	T/C	Synonymous	LOW	Ala 575'
	82,831,752	T/C	Missense	MODERATE	Met 235' >Val
217	82,830,589	T/C	Synonymous	LOW	Ala 575'
	82,831,280	T/C	Missense	MODERATE	Gln 392' > Arg
221	82,830,589	T/C	Synonymous	LOW	Ala 575'
225	82,830,589	T/C	Synonymous	LOW	Ala 575'
227	82,830,589	T/C	Synonymous	LOW	Ala 575'
	82,830,589	T/C	Synonymous	LOW	Ala 575'
237	82,830,589	T/C	Synonymous	LOW	Ala 575'
	82,831,752	T/C	Missense	MODERATE	Met 235' > Val
	82,832,178	T/C	Missense	MODERATE	Asn 93' Asp
239	82,830,589	T/C	Synonymous	LOW	Ala 575'
241	82,830,589	T/C	Synonymous	LOW	Ala 575'
	82,831,280	T/C	Missense	MODERATE	Gln 392' > Arg
243	82,830,589	T/C	Synonymous	LOW	Ala 575'
	82,831,280	T/C	Missense	MODERATE	Gln 392' Arg
245	82,830,589	T/C	Synonymous	LOW	Ala 575'

Sample ID	Position	Mutation	Type	Impact	Amino Acid
201	82,831,393	G/A	Synonymous	LOW	Thr 354'
205	82,832,218	G/A	Synonymous	LOW	Pro 566'
217	82,832,090	G/A	Synonymous	LOW	Val 165'
233	82,831,393	G/A	Synonymous	LOW	Thr 354'
	82,831,429	G/A	Synonymous	LOW	Thr 342'
243	82,831,393	G/A	Synonymous	LOW	Thr 354'

Sample ID	Position	Mutation	Type	Impact	Amine Acid
203	82,830,616	A/G	Synonymous	LOW	Pro 566'
205	82,830,616	A/G	Synonymous	LOW	Pro 566'
207	82,830,616	A/G	Synonymous	LOW	Pro 566'
215	82,830,616	A/G	Synonymous	LOW	Pro 566'
217	82,830,616	A/G	Synonymous	LOW	Pro 566'
221	82,830,616	A/G	Synonymous	LOW	Pro 566'
227	82,830,616	A/G	Synonymous	LOW	Pro 566'
237	82,830,616	A/G	Synonymous	LOW	Pro 566'
239	82,830,616	A/G	Synonymous	LOW	Pro 566'
241	82,830,616	A/G	Synonymous	LOW	Pro 566'
243	82,830,616	A/G	Synonymous	LOW	Pro 566'
245	82,830,616	A/G	Synonymous	LOW	Pro 566'

Sample ID	Position	Mutation	Type	Impact	Amino acid
201	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
207	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
209	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
213	53,575,869	C/T	Synonymous	LOW	Ala 965'
	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
215	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
217	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
	53,582,818	C/T	Synonymous	LOW	Ser 3'
223	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
225	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
227	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
235	53,576,549	C/T	Missense	MODERATE	Val 739' > Ile
239	53,582,354	C/T	Missense	MODERATE	Cys 158' Tyr
245	53,576,549	C/T	Missense	MODERATE	Cys 158' Tyr

Sample ID	Position	Mutation	Type	Impact	Amino acid
205	53,582,438	C/A	Missense	MODERATE	Cys 130' > Phe
217	53,582,080	C/A	Synonymous	LOW	Ala 249'
229	53,582,400	C/A	Missense	MODERATE	Val 143' > Phe
239	53,576,873	C/A	Missense	MODERATE	Ala 631' >Ser
241	53,582,080	C/A	Synonymous	LOW	Ala 249'
245	53,577,058	C/A	Missense	MODERATE	Gly 569' >Val
	53,582,169	C/A	Missense	MODERATE	Val 220' >Phe

Sample ID	Position	Mutation	Type	Impact	Amino acid
239	53,582,352	TA/T	Frameshift	High	Cys 158' Ter

Mutational and Expression Profile of ZNF217, ZNF750, ZNF703 Zinc Finger Genes in Kenya Women diagnosed with Breast Cancer

Michael Kitoi[13*], John Gitau[2], Godfrey Wagutu[3], Kennedy Mwangi[4], Florence Ngonga[1], Francis Makokha[3]

[1] Jomo Kenyatta University of Agriculture and Technology (JKUAT)

[2] Département de Chimie, Université du Quebéc à Montréal, Montréal, Québec, Canada

[3] Directorate of Research and Innovation, Mount Kenya University, Thika, Kenya

[4] International Livestock Research Institute, Nairobi, Kenya

* Correspondence:
Michael Kitoi
Michael.o.kitoi@gmail.com

ABSTRACT

Objectives: To characterize the somatic mutational spectrum and transcriptomic expression of the zinc-finger genes ZNF217, ZNF703, and ZNF750 in Kenyan women with breast cancer, and to explore their associations with clinicopathologic features.

Methods: Whole-exome sequencing and RNA-sequencing were performed on paired tumor and adjacent normal tissues from 23 consented patients treated at two Kenyan referral hospitals. Variants were called with Mutect2 using a study-specific panel of normal; functional consequences were annotated with VEP. After featureCounts quantification, differential expression was analyzed in DESeq2 (fold-change $\geq$ 1.5, $p < 0.05$). Two-sample t-tests (mutations) and ANOVA (expression) evaluated relationships with HER2 status and clinical stage.

Results: A total of 358 somatic mutations were detected: 170 in ZNF217, 24 in ZNF70*3* and 164 in ZNF750. Single-nucleotide substitutions (319 SNPs) dominated, with C→T and A→G changes most common; 27 deletions and 2 insertions were also observed. Frameshift events in ZNF217 and ZNF703 introduced premature stop codons predicted to truncate protein function. All three genes were significantly up-regulated in tumors versus normal (ZNF217 p = 0.0004), with the greatest expression in HER2-positive tumors and in stages 2–3 disease. Mutation burden for each gene did not differ by HER2 status ($p > 0.56$) or by stage ($p > 0.32$).

Conclusions: Kenyan breast tumors harbor frequent, functionally relevant mutations and marked over-expression of ZNF217, ZNF703, and ZNF750. These alterations, especially the pronounced up-regulation of ZNF217*,* highlight the trio's potential as diagnostic or prognostic biomarkers and warrant larger studies to validate their clinical utility and suitability as therapeutic targets in sub-Saharan African populations.

Keywords: ZNF217; ZNF703; ZNF750; Kenyan breast cancer; whole-exome sequencing

INTRODUCTION

The global incidence of cancer reported approximately 20 million new cases in 2022 including 11.6% female breast cancer case which ranked as the second most common cancer worldwide (Bray et al., 2024). Cancer research indicates that breast cancer represented 6.7 percent of total cancer-related fatalities among the 9.7 million deaths (Bray et al., 2024). The GLOBOCAN 2022 report shows breast cancer takes the lead position as the most common cancer affecting Kenyan females with a rate of 25.6% for new cancer cases (Global Cancer Observatory, 2022). Breast cancer stands as the main cause of fatalities in Keenya where the annual new cases reached 7,243 (16.2% of all cases) while mortality rates showed 3,398 deaths (Global Cancer Observatory, 2022).

Recent genomic studies have proven that Zinc Finger proteins (ZNF) play an essential role in the development of breast cancer (Stradella et al., 2022). Breast cancer pathogenesis shows ZNF217, ZNF750 and ZNF703 as potential contributors because several mutations and dysregulated expression patterns during development have been linked to disease progression. ZNF217 functions as a normal cellular repressor that controls both cell survival and growth dynamics (Krig et al. (2010). The overexpression of ZNF217 occurs when amplifications or mutations take place causing gene expression disturbances as well as aggressive breast cancer development that affects African ancestry populations (Ansari-Pour et al., 2021). Breast cancer in the African population undergoes more aggressive development due to ZNF217 non-coding mutations (Ansari-Pour et al., 2021). Scientific studies have demonstrated that ZNF217 frame-shift mutations damage protein structures which results in improper gene expression patterns that create cancer development (Krig et al., 2010a). Breast cancer cells express ZNF217 at 20q13 amplification regions that drive the transcription and protein synthesis of Erb-B2 Receptor Tyrosine Kinase 3 (ErbB3) (Krig et al., 2010).

ZNF750 gene plays a regulatory role in epidermal differentiation through its control of differentiation-specific genes located on chromosome 17q25.3 (Butera et al., 2020). ZNF750 functions as a tumor suppressor in normal tissue because it blocks cell multiplication and facilitates cell differentiation. Genomic abnormalities in ZNF750 disrupt crucial cell functions that then increase cancer development probabilities according to (Butera et al., 2020) across various aggressive breast cancer variants. Similarly, ZNF750 gene fusions promote cancer-related signal pathways for breast cancer progression. Butera et al. (2020) demonstrates that ZNF750 gene mutations connect with two types of breast cancer: aggressive triple-negative breast cancer (TNBC) and lymph node metastasis. Mutations in ZNF750 lead to dysfunctional cellular processes in breast tissue because they generate both the condition of hyper-expression and hypo-expression of ZNF750 (Cassandri et al., 2020).

The transcription factor ZNF703 functions normally by controlling developmental processes as well as cellular differentiation pathways when present on chromosome 8p12 (Klæstad et al., 2021). The study of ZNF703 in breast cancer shows that it activates oncogenic signaling through excessive protein production which advances tumor growth by increasing cell survival and growth potential (Zhang et al., 2022). The poor clinical outcome and aggressive features of breast cancer are strongly associated with high ZNF703 expression levels especially in patients with luminal B breast cancer subtypes (Marzbany et al., 2019). The research on ZNF703 gene-

copy associations with breast cancer proliferation and outcome and Luminal subtype revealed
that tumor cells with high copy numbers showed increased proliferation and aggressive traits in 7
percent of cases most frequently within the Luminal B subgroup (Klæstad et al., 2021).

In the current study, therefore, whole genomic exome and RNA-sequencing data of tumor and
adjacent normal tissues of 23 Kenyan women with breast cancer were analyzed. We aimed (i) to
define the somatic-mutation map of the three genes, ZNF217, ZNF703 and ZNF750, (ii) to
measure their level of dysregulation in cancer and normal breast tissue, and (iii) to investigate
links between alterations of these genes and clinicopathologic parameters. Verifying the
molecular phenotypes of these ZNF genes in a Kenyan population will progress precision-
oncology studies in sub-Saharan Africa and provide the foundations of subsequent biomarker
validation and identification of new drug targets.

MATERIALS AND METHODS

Patients and Samples

This research collected its data from a pilot study examining breast cancer progression in Kenyan patients {Institutional Ethics Review Committee (IERC), Ref: 2018/REC-80 (v7); License No: NACOSTI/P/22/21374}. After patients granted their consent at AIC Kijabe Hospital and Aga Khan University Hospital, Nairobi, purposive sampling enrolled 23 patients with malignant breast cancer. Fresh tumor tissue together with adjacent normal tissue were extracted from 23 patients while 46 patient samples were sent to National Cancer Institute at Bethesda MD USA for sequencing procedures.

Whole-exome sequencing and RNA-sequencing

Tang et al. (2023) provided the methods for processing tumors together with normal samples. Whole-exome sequencing together with RNA-sequencing was done using a Novaseq system which produced paired end reads of 150bp at a total read depth of 30 million according to Tang et al. (2023). Tumor sample analysis was done at 250x depth while normal adjacent tissue analysis reached 150x depth (Tang et al., 2023).

Reads mapping and variant calling

The workflow proceeded with quality control evaluations of raw whole exome data (WED) and RNA-seq data from twenty-three samples through implementation of FastQC (ref). MultiQC (Ewels et al., 2016) generated the summary of results. The reads underwent trimming for low quality and adapter elements through Trimmomatic before undergoing FASTQC and MultiQC quality screening. The hg38 human reference genome received mapped BWA MEM (ref) reads from those samples successful during the quality control phase. The analysis report demonstrated that mapping fully succeeded for greater than 95% of input reads. Picard tool performed read duplication followed by addition of read-groups into the deduped BAM files. The base quality recalibration process utilized GATK according to G. Zhang et al. (2015). The somatic variant calling process in paired tumor normal mode occurred through MuTect2 (ref) (Chen et al., 2020).

The analysis made use of the panel of normal (PON) derived from normal reads obtained through the study. Normal samples entered Mutect2 with tumor-only mode in order to produce the panel of normal according to Chen et al. (2020). The CreateSomaticPanelOfNormals tool (O'Donovan, 2019) performed variant call aggregation of individual results from separate processing of each normal sample. The somatic variant detection accuracy increased because this method helped identify and eliminate technical artifacts and germline variants that normally exist in normal samples. The PON accompanied variant detection to remove artifacts together with germline variants while ensuring proper identification of somatic mutations. The variant tool set normalization (VT) (Bayat et al., 2017) served as the variant normalization method which involved GATK filtering while variant effect predictor (VEP) (Salz et al., 2023) delivered functional/consequence annotations. Each annotation score (whether, moderate, high or low) originated from the gene effect analysis. The variants linked to ZNF genes can be viewed using the Integrative Genomic Viewer (IGV) (ref) (Robinson et al., 2017).

The panel of normal (PON) used for analysis stemmed from normal reads which were collected
for this purpose. Mutect2 with tumor-only mode processing generated the panel of normal from
test samples according to Chen et al., 2020. Each individual normal sample received independent
processing for variant calling before the CreateSomaticPanelOfNormals tool combined the
output (O'Donovan, 2019). The somatic variant detection accuracy increased because this
method helped identify and eliminate technical artifacts and germline variants that normally exist
in normal samples. The PON operated as a filtering tool during variant calling to delete technical
artifacts together with germline variants which ensured precise somatic mutation identification.
The variant tool set normalization (VT) (Bayat et al., 2017) served as the variant normalization
method which involved GATK filtering while variant effect predictor (VEP) (Salz et al., 2023)
delivered functional/consequence annotations. The annotation process determined the impact
levels (whether, moderate, high or low) of each gene. The variants linked to ZNF genes can be
viewed using the Integrative Genomic Viewer (IGV) (ref) (Robinson et al., 2017).

Gene Expression Analysis

Quality assessment through FASTQC with MultiQC tools was conducted before performing the
RNA-seq analysis for the 46 processed samples. Testing of sample quality yielded satisfactory
results so researchers performed human reference genome mapping of the samples using the
STAR aligner. The program calculated the reference genome comparisons with each read
sequence by determining the number of mismatches (NM). The featureCounts software (Liao et
al., 2014) determined the quantitative count of reads per gene. DESeq2 software executed
normalization procedures on the gene expression data (Malick et al., 2022). Each sequencing-
read proceeded to produce raw gene count data after the quantification process. The raw counts
measured the sequencings' numbers that aligned to each gene and served as the starting point for
normalization procedures. DESeq2 applied size factor normalization which generated unique
size factors for each sample to normalize the varying sequencing depths in data (Malick et al.,
2022). Size factors were determined by finding the median value of dividing gene counts with
their corresponding geometric mean across all measured genes. The normalization procedure
ended by producing a count matrix which contained normalized gene expression measurements
for every sample.

Differential Gene Expression (DGE) Analysis and Principal Component Analysis (PCA)

The analysis of Differential Gene Expression (DGE) through DESeq2 occurred in R (Malick et
al., 2022). The biological variance calculation for each experimental condition combines
replicated results to establish expression distinctions among two specific conditions. The
identified differentially expressed genes were identified through a combination of $\geq$ 1.5-fold
change and statistical significance $\leq$ 0.05 (Gitau et al., 2024). Principal Component Analysis
(PCA) using Python processed normalized gene expression data in order to detect expression
profile patterns and reduce dataset dimensions of ZNF217, ZNF750, and ZNF703 genes
(Greenacre et al., 2022). R was used for detailed visualization of ZNF217, ZNF750, and ZNF703
gene differential expression through ggplot2 boxplot creation.

Statistical Significance

175   Simple t-test established the expression level differences' significance between normal and tumor
176   samples through a statistical significance analysis.

 RESULTS

 Patients and sample characteristics

The research has detailed the demographic and clinical/pathological characteristics of the 23
studied patients in Table 1. Participation groups included cancer stages together with human
epidermal growth factor receptor 2 (HER2) expression results. Table 1 reveals that 52.2% of
patients showed HER2+ status while 8.6% were in cancer stage I and 34.7% had stage III and
47.8% were at stage II of their condition.

ZNF703 Gene Mutations

A total of 24 gene mutations were detected in the 23 patient samples including 20 SNPs
combined with 3 deletions and a single insertion (Figure 1A). A total of twelve patients among
the twenty-three samples (12) exhibited ZNF703 gene mutations while eleven patients did not
show any mutations in the gene (Figure 1B). Patient 237 demonstrated the greatest mutating
activity with four different mutations yet 14 patients presented one mutation and 8 patients did
not have mutagenic changes (Figure 1C). The C/T substitutions appeared as the dominant
substitution type whereas G to C occurred least frequently (Figure 1C). Study results showed
four missense types among the classified variants and patient 237 held the sole frameshift
mutation in ZNF703 gene (Table 2). The variants affecting this gene generated moderate effects
across samples except for patient 237 who experienced a high impact due to frameshift mutation
(Table 2). Missense types resulted in moderate consequences for ZNF703 genes of patient 201
and 205 and were also observed in patient 223 and 233 but to a lower extent. A high-degree of
influence resulted from the in-frame deletion which replaced Alanine 517. The in-frame deletion
demonstrated homozygous removal affecting both alleles of this gene when visualized through
IGV (Figure 1D).

ZNF750 Gene Mutations

A total of 164 mutations were identified among the 23 patients who exhibited 154 SNPs and 9
deletions together with 1 insertion (Figure 2A). About three samples contained 15 to 12 and 11
SNP mutations while two other samples showed less than three SNPs in total (Figure 2B).
Vertical transitions based on A/G substitutions constituted 40 mutations in the gene pool yet A/C
substitutions and A/T substitutions together with C/A and G/T substitutions totaled fewer than 10
counts (Figure 2C). The highest occurring indel mutation is TTTTG/T which occurred five times
and all other mutations occurred only once during the analysis (Figure 2D).

The C to T DNA change at positions 82,831,023 and 82,831,023 in two specimens led to a
coding sequence mutation transforming amino acid valine 478' to isoleucine which shares
comparable hydrophobic properties (Table 3). A genetic alteration at position 82,830,234 in
sample 231 modified Asp to Asn according to the information contained in Table 3. At position
82,831,982 of sample 241 the C/T change transformed arginine (Arg) into glutamine (Gln at
position 158' as listed in Table 3.

The majority of T/C SNPs resulted in no change of the amino acid sequence as Ala 575' remains
constant in the protein chain (Figure 2E) (Table 4). Three specimens demonstrated T to C
substitution at position 82,831,280 of the gene leading to missense mutations which changed
glutamine (Gln) at position 392 into arginine (Table 4 and Figure 2E). The T/C substitution at
position 82,831,752 generated a missense mutation altering methionine (Met) at position 235' to
valine (Val) based on the data in Table 4.

These gene alterations at position 354 produced no impact on the ZNF750 protein because they
maintained amino acid threonine (Thr) while position 165' retained valine (Val) resulting in an
unmodified subsequent protein structure (Table 5). At position 82,831,752 of the gene, twelve
annotated A/G were found to be synonymous because proline (Pro) at specific positions
maintained the same amino acid sequence without affecting protein functionality (Table 6).

ZNF217 Gene Mutations

Investigators uncovered 170 mutations along with 145 SNPS and 24 deletion and one insertion
throughout the 23 samples (Figure 3A). A total of 105 C to T substitutions were the most
frequent mutations among ZNF217 changes while G to C substitutions only happened less than
two times (Figure 3B). GAGAC/C in-frame deletion represented the most often detected
mutation in patients with eight mutations among all samples (Figure 3B). A total of 36 mutations
among all identified changes have received annotations. Recorded mutations comprised of C/T,
C/A and TA/T variants with 22 mutations classified as missense and 13 as synonymous
mutations and one classified as a frameshift disturbance (Figure 3D).

Twelve mutations identified turned out to be missense variations and two mutations were defined
as synonymous. Ten C to T mutations found across the samples resulted in the creation of
ZNF217 gene variants with isoleucine (Ile) at position 739 instead of valine (Val) due to a
change in base coding (Figure 3E). Two analyzed samples showed a C/T base substitution at
position 53,582,354 that equated to a synonymous substitution of Cys to Tyr at position 158
(Figure 3E and Table 7).

The nucleotide mutation C/A at position 53,582,438 within sample 205 caused a phenotypic
variation where cysteine (Cys) at position 130 was mutated into phenylalanine (Phe) (Table 8).
Through its synonymous mutation at position 53,582,080 Sample 217 maintains an identical
amino acid sequence for alanine located at position 249 according to Table 8. A C/A substitution
found at position 53,582,400 in sample 229 produces a missense mutation that changes amino
acid valine at position 143 into phenylalanine (Table 8).

At nucleotide position 53,576,873 the C/A substitution leads to a missense mutation that
switches alanine to serine at position 631 (Table 8). Two C/A mutations occur in Sample 245
where the first substitution at position 53,577,058 leads to Val in position 569 and another
mutation at position 53,582,169 converts Val to Phe found at position 220 (Table 8). The
position 53,582,352 TA/T mutation produces a frameshift type mutation (Figure 4E). The
frameshift mutation leads to a premature translation termination because it replaces position 158
cysteine (Cys) with a termination codon (Ter) (Table 9).

The research team evaluated the mutation rates between the three genes while comparing results
with HER2 status. Researchers applied two-sample t-tests for evaluating significance in the data
relationship. Mutation counts of the ZNF217 gene displayed similar median counts between
HER2-positive cases and those that were HER2-negative during analysis. However, the
distribution of mutation levels was more variable in HER2-positive samples. Results from the t-
test yielded no remarkable statistical difference (t = -0.022, df = 21, p = 0.9825) as shown in
Figure 4A. The relationship between ZNF703 mutation counts and HER2 status failed to show
any statistical significance (t = -0.587, df = 21, p = 0.5637) at the same time that the data
exhibited elevated variability in HER2-positive samples. Mutations of ZNF750 produced
identical distribution patterns between HER2-positive and HER2-negative groups since their
interquartile ranges cross and statistical tests showed no significant differences (t = -0.587, df =
21, p = 0.5637) (Figure 4C). Data shows that HER2 status has no impact on the mutation
burdens found for the three genes ZNF217, ZNF703 and ZNF750 within this patient cohort.

Analysis of the three gene mutations showed no significant relationship with cancer staging.
ANOVA statistical analyses established the evaluation of significance levels between Stage 1, 2
and 3 data points. Stage 2 exhibited minimal variations in total mutations but statistical analysis
showed no significant difference (p > 0.32 for all stage comparisons) compared to other stages
(Figure 5A). ZNF703 mutation counts demonstrated an upward trend between stage 1 and 3
according to ANOVA analysis results but there was no statistical significance (p > 0.35 in all
pairwise tests). ZNF750 mutation data displayed an upward trend (Figure 5C) despite significant
range association yet demonstrated no statistically meaningful differences between categories (p
> 0.35 for all analyses). The stage of cancer does not create a clear link between mutation counts
for ZNF217, ZNF703 or ZNF750.

Expression Profile for 23 Patients

Analysis of the three ZNF gene samples for the 23 paired breast tissues was conducted in R by
using Principal Component Analysis (PCA). Three ZNF gene cluster analysis for 23 paired
breast specimen samples occurred within R through the utilization of Principal Component
Analysis (PCA) methods (Figure 6). The normal tissue group aligned closely in the cluster
indicating uniform expression levels but the tumor group spread out widely consistent with
differences between various cancer stages (Figure 6). The Stage 1 tumors exhibited blue clusters
that stayed compressed together while Stage 2 and Stage 3 tumors expanded in dispersion
revealing elevated progression-related heterogeneity (Figure 6). The two principal components
account for 58.46% and 24.77% of the variance independently so specific changes in gene
expression in tumors explain most of the profile variability (Figure 6).

The evaluation of differentially-expressed genes (DGE) occurred with normalized data. The
research analysis indicated that all three target genes displayed important expression intensities
(P <0.05) in examined tumor tissues. The experiment results confirmed ZNF217 genes as
significantly active (P= 0.0004) while ZNF750 genes showed significant expression (p= 0.001)
and ZNF703 genes also exhibited significant expression (0.002) in tumor samples (Figure 7).

The research evaluated ZNF750 gene expression based on patient cancer progression levels. The
expression levels of this gene were enhanced in tumor samples obtained from patients in stages 2

and 3. Analysis of stage 1 tissue samples yielded five results which two of them indicated
elevated ZNF750 gene activity in healthy tissues. Some patients diagnosed with stage 1 breast
cancer show elevated ZNF750 gene expression in their healthy breast tissue which suggests the
gene might play a regulatory part during the first stages of cancer advancement (Figure 8A). The
results from the Kruskal-Wallis test showed no significant differences between cancer stages
based on $\chi^2 = 5.6918$ (df = 2) at p = 0.05808. This study examined how the gene expression
levels changed when considering both HER2 protein levels and cancer development stage. The
analysis showed HER2-positive stage 2 patients had the highest ZNF750 expression levels yet
stage 3 showed higher expression levels in HER2-negative patients compared to HER2-positive
patients (Figure 8B). The stage 1 results demonstrated that HER2-positive breast tumors
demonstrated greater gene expression than their HER2-negative counterparts (Figure 8B). The
results from paired t-test revealed no significant statistical difference between the two groups (t =
0.18044, df = 20.998, p = 0.8585) when comparing HER2-positive and HER2-negative
expression. Stage 2 cancer cells with elevated HER2 expression likely support tumor progression
during intermediate stages but stage 3 cancer cells with HER2-negative status may compensate
their more advanced condition through increased gene expression.

Researchers evaluated ZNF703 gene expression data to match it against the cancer stage
evaluations of the specimens under study. The gene exhibited elevated expression levels
throughout tumor tissue at stages 2 and 3 based on Figure 9A. Within stage 1 and the five
analyzed samples two demonstrated elevated ZNF703 gene expression at the normal tissue level
(Figure 9A). The expression level of ZNF703 showed no statistical difference between different
cancer stages based on the Kruskal-Wallis test ($\chi^2 = 1.2261$, df = 2, p = 0.5417). Analysis of
ZNF703 gene expression took place based on HER2 status and cancer stage distribution. In
stages 1 and 3 ZNF703 showed its highest expression in HER2-negative samples than in HER2-
positive samples (Figure 9B). According to a paired t-test analysis the expression levels between
HER2-positive and HER2-negative groups matched up with no significant statistical difference
at t = 0.36584, df = 18.554, and p = 0.7186. The ZNF703 gene shows higher importance for
HER2-negative breast cancer development in stages one and three along with advanced cases
because it promotes tumor progression beyond HER2 signaling mechanisms.

Most research samples documented enhanced ZNF217 gene expression levels in normal tissues
with five specimens showing increased gene activity specifically in normal tissue (Figure 10A).
Stage 1 saw three out of five cases where the gene had elevated expression levels in normal
breast tissue according to Figure 10A. The Kruskal-Wallis analysis demonstrated that cancer
stages did not affect ZNF217 expression levels because the results indicated $\chi^2 = 2.9136$, df = 2,
p = 0.233. Analysis of ZNF217 gene expression involved evaluation through normal and HER2
status comparison across different cancer stages. The gene expression levels in stage 1 and 2
samples with HER2 positivity outpaced HER2-negative results (Figure 10B). The analysis of
Figure 10B revealed greater ZNF217 gene expression in cancer samples that were HER2-
negative during stage 3. A paired t-test analysis revealed no significant difference in expression
patterns between HER2-positive and HER2-negative groups (t = -0.62653, df = 20.973, p =
0.5377). The ZNF217 gene functions differently according to HER2 status when it accelerates
early-stage HER2-positive cancer progression before taking precedence in late-stage HER2-
negative cancer development.

Gene regulation patterns follow distinct routes based on the effects of genetic mutations on
ZNF217 together with ZNF750 and ZNF703. Structural modification from frameshift mutations
linked to ZNF217 expression reductions yet these defects have probable functional failures
(Figure 11A). Conversely, missense mutations had positive correlations with ZNF217 expression
because they facilitated this elevation. Total SNPs in ZNF703 demonstrated robust positive
association with expression alterations and missense mutations showed moderate contribution to
expression elevation through known oncogenic functions (Figure 11B). Total SNP burden in
ZNF750 showed a direct correlation with gene expression variations because elevated mutational
load powerfully manipulates regulatory patterns (Figure 11C). The analysis showed synonymous
mutations presented minimal effect on gene expression across all sequences in the three genes
studied. Breast cancer gene expression differences primarily result from functional mutations
that lead to missense alterations yet frameshift mutations can reduce expression levels when they
affect protein stability.

DISCUSSION

The research analyzed the ZNF217, ZNF750, and ZNF703 genomic alterations and mRNA levels within 23 Kenyan breast cancer patients. The examination revealed 359 genetic alterations that primarily exhibited C>T nucleotide changes in each assessed gene. The transitions identified through genomic analysis become essential in cancer research because Alexandrov et al. (2020) explains they commonly appear as DNA damage markers from oxidative stress in cancerous cells. Tissues from tumor areas demonstrated substantial elevation of ZNF217 together with ZNF703 and ZNF750 transcriptional levels compared to normal tissues (P<0.05), yet ZNF217 exhibited the greatest detected difference. The Principal Component Analysis (PCA) data effectively distinguished tumor sample data from normal sample data. The first two principal components in the PCA plot explain 58.46% and 24.77% of the variance which evidences that tumor-specific alterations in the three gene expressions drive most variations in measurement profiles (Gitau et al., 2024). The expression patterns differed according to HER2 status together with cancer stage progression indicating contextual roles for each gene in tumor developments. Studies showed that ZNF750 expression levels rose progressively from stage 2 through stage 3 tumor samples yet certain stage 1 specimens displayed elevated expression in healthy tissues thus indicating that ZNF750 works as an early regulatory factor (Gitau et al., 2024).

ZNF217 mutations included an in-frame GAGAC/C deletion which presented a strong potential to disrupt protein structure. An in-frame GAGAC/C deletion in the gene sequence is relevant because it may disrupt protein functioning which can advance breast cancer through modified EMT and TGF-β signaling mechanisms (Vendrell et al., 2012). A frameshift mutation of TA/T during analysis for sample 239 transformed the amino acid cysteine at position 158 into a termination codon (Ter) leading to early translation termination and functional damage to its cell cycle and apoptosis regulatory actions. Evidences indicate truncating ZNF217 mutations can cause deregulated cell proliferation through destruction of its gene-controlling abilities on apoptosis regulation (Krig et al., 2010b). Our research supports data showing how the frameshift mutation might push cells toward malignancy because it abets uncontrolled cell growth while demonstrating the importance of this mechanism due to ZNF217 serving as an ErbB3 pathway activator which commonly plays a role in breast cancer development (Beroukhim et al., 2007). The ZNF217 gene showed multiple missense mutations which contained exchanges at cysteine (C) 158 position that substituted the amino acid with tyrosine (Y). The conversion of a polar amino acid into a nonpolar type leads to substantial structural modifications of ZNF217 protein that could modify its DNA-binding features essential for regulatory activity (Briest et al., 2023). The aggressive cancer phenotype may result from this breast cancer-specific mutation of ZNF217 because ZNF217 frequently shows amplification in breast cancer tissues (Vendrell et al., 2012). Research results indicated that ZNF217 expression levels were elevated in most examined normal tissue samples where five samples displayed quantitatively higher ZNF217 protein levels. The research indicates ZNF217 regulates early cancer development stages before oncogenesis completely manifests (Tang et al., 2023).

A frameshift mutation found in ZNF703 displayed an alanine deletion at position 514 which was an in-frame variant. The scientific literature shows that ZNF703 frameshift mutations produce increased cell proliferation in breast cancer by modifying MAPK and PI3K/AKT signal pathways (X. Zhang et al., 2013). ZNF703 overexpression links to higher proliferation rates in

HER2-negative specimens because such mutations prove significant for both moderate-impact
and HER2-negative cases (X. Zhang et al., 2022). The altered ZNF703 gene demonstrates how it
promotes aggressive cancer growth in HER2-negative tumors which makes it a possible
therapeutic target for those breast cancers. Twelve breast cancer cases provided evidence of
increased ZNF703 expression in HER2-negative tumors across stage 1 and stage 3 samples
showing its role in tumor expansion that bypasses HER2 pathways (Stradella et al., 2022).

Scientists believe the ZNF750 gene mutation profile acknowledges missense mutations affecting
tumor-suppressive functions through DNA-binding domain amino acid substitutions at position
478 from valine (V) to isoleucine (I) (Butera et al., 2020). The mutations found in ZNF750 are
highly important for triple-negative breast cancers since the loss of ZNF750 function remains
linked to progressive disease features (Cassandri et al., 2020). The study demonstrates that
ZNF750 mutations stand out as a biomarker that indicates aggressive breast cancer types within
African populations along with their distinct mutational patterns compared to other research
populations (Brewster et al., 2014). The different types of mutations found within ZNF217
ZNF703 and ZNF750 gene sequences within the Kenyan cohort seem to result from the
combination of genetic and environmental elements that affect how breast cancer evolves in
Africans. ZNF750 demonstrates increased expression specifically in HER2-positive stage 2
tumors according to expression analysis findings (Klæstad et al., 2021). Its oncogenic role seems
to play a greater part in HER2-negative tumors that appear later in progression thus suggesting a
compensatory mechanism.

REFERENCES

Aganezov, S., Yan, S. M., Soto, D. C., Kirsche, M., Zarate, S., Avdeyev, P., Taylor, D. J., Shafin, K., Shumate, A., Xiao, C., Wagner, J., McDaniel, J., Olson, N. D., Sauria, M. E. G., Vollger, M. R., Rhie, A., Meredith, M., Martin, S., Lee, J., … Schatz, M. C. (2022). A complete reference genome improves analysis of human genetic variation. *Science (New York, N.Y.)*, *376*(6588), eabl3533. https://doi.org/10.1126/science.abl3533

Alessio, B., Matteo, C., Rugolo, F., Massimiliano, A., & Gerry, M. (2020). The ZNF750–RAC1 axis as potential prognostic factor for breast cancer. *Cell Death Discovery*, *6*(1). https://search.proquest.com/openview/1b6425cdbc87baf718b409d5533928ee/1?pq-origsite=gscholar&cbl=2041962

Alexandrov, L. B., Kim, J., Haradhvala, N. J., Huang, M. N., Tian Ng, A. W., Wu, Y., Boot, A., Covington, K. R., Gordenin, D. A., Bergstrom, E. N., Islam, S. M. A., Lopez-Bigas, N., Klimczak, L. J., McPherson, J. R., Morganella, S., Sabarinathan, R., Wheeler, D. A., Mustonen, V., Getz, G., … Stratton, M. R. (2020). The repertoire of mutational signatures in human cancer. *Nature*, *578*(7793), 94–101. https://doi.org/10.1038/s41586-020-1943-3

Ansari-Pour, N., Zheng, Y., Yoshimatsu, T. F., Sanni, A., Ajani, M., Reynier, J.-B., Tapinos, A., Pitt, J. J., Dentro, S., Woodard, A., Rajagopal, P. S., Fitzgerald, D., Gruber, A. J., Odetunde, A., Popoola, A., Falusi, A. G., Babalola, C. P., Ogundiran, T., Ibrahim, N., … Olopade, O. I. (2021). Whole-genome analysis of Nigerian patients with breast cancer reveals ethnic-driven somatic evolution and distinct genomic subtypes. *Nature Communications*, *12*(1), 6946. https://doi.org/10.1038/s41467-021-27079-w

Bayat, A., Gaëta, B., Ignjatovic, A., & Parameswaran, S. (2017). Improved VCF normalization for accurate VCF comparison. *Bioinformatics*, *33*(7), 964–970. https://doi.org/10.1093/bioinformatics/btw748

Bray, F., Laversanne, M., Sung, H., Ferlay, J., Siegel, R. L., Soerjomataram, I., & Jemal, A. (2024). Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians*, *74*(3), 229–263. https://doi.org/10.3322/caac.21834

Brewster, A. M., Chavez-MacGregor, M., & Brown, P. (2014). Epidemiology, biology, and treatment of triple-negative breast cancer in women of African ancestry. *The Lancet Oncology*, *15*(13), e625–e634. https://doi.org/10.1016/S1470-2045(14)70364-X

Butera, A., Cassandri, M., Rugolo, F., Agostini, M., & Melino, G. (2020). The ZNF750–RAC1 axis as potential prognostic factor for breast cancer. *Cell Death Discovery*, *6*(1), 1–9. https://doi.org/10.1038/s41420-020-00371-2

Cassandri, M., Butera, A., Amelio, I., Lena, A. M., Montanaro, M., Mauriello, A., Anemona, L., Candi, E., Knight, R. A., Agostini, M., & Melino, G. (2020). ZNF750 represses breast cancer

invasion via epigenetic control of prometastatic genes. *Oncogene*, *39*(22), 4331–4343.
https://doi.org/10.1038/s41388-020-1277-5

Chen, Z., Yuan, Y., Chen, X., Chen, J., Lin, S., Li, X., & Du, H. (2020). Systematic comparison
of somatic variant calling performance among different sequencing depth and mutation
frequency. *Scientific Reports*, *10*(1), 3501. https://doi.org/10.1038/s41598-020-60559-5

Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048.
https://doi.org/10.1093/bioinformatics/btw354

Gitau, J., Kinyori, G., Sayed, S., Saleem, M., Makokha, F. W., & Kirabo, A. (2024). *The*
Association between the JAK-STAT Pathway and Hypertension among Kenyan Women
Diagnosed with Breast Cancer (p. 2024.06.07.597892). bioRxiv.
https://doi.org/10.1101/2024.06.07.597892

GLOBOCAN (2022). *Global cancer statistics 2022- GLOBOCAN estimates of incidence and*
mortality worldwide for 36 cancers in 185 countries. https://doi.org/10.3322/caac.21834

Greenacre, M., Groenen, P. J. F., Hastie, T., D'Enza, A. I., Markos, A., & Tuzhilina, E. (2022).
Principal component analysis. *Nature Reviews Methods Primers*, *2*(1), 1–21.
https://doi.org/10.1038/s43586-022-00184-w

Klæstad, E., Sawicka, J. E., Engstrøm, M. J., Ytterhus, B., Valla, M., & Bofin, A. M. (2021).
ZNF703 gene copy number and protein expression in breast cancer; associations with
proliferation, prognosis and luminal subtypes. *Breast Cancer Research and Treatment*, *186*(1),
65–77. https://doi.org/10.1007/s10549-020-06035-0

Krig, S. R., Miller, J. K., Frietze, S., Beckett, L. A., Neve, R. M., Farnham, P. J., Yaswen, P. I.,
& Sweeney, C. A. (2010a). ZNF217, a candidate breast cancer oncogene amplified at 20q13,
regulates expression of the ErbB3 receptor tyrosine kinase in breast cancer cells. *Oncogene*,
29(40), 5500–5510. https://doi.org/10.1038/onc.2010.289

Krig, S. R., Miller, J. K., Frietze, S., Beckett, L. A., Neve, R. M., Farnham, P. J., Yaswen, P. I.,
& Sweeney, C. A. (2010b). ZNF217, a candidate breast cancer oncogene amplified at 20q13,
regulates expression of the ErbB3 receptor tyrosine kinase in breast cancer cells. *Oncogene*,
29(40), 5500–5510. https://doi.org/10.1038/onc.2010.289

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: An efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–930.

Malick, S. B. A., Conteh, F., & Sawo, M. (2022). *Bioinformatics Analysis of Differentially*
Expressed Gene's in Breast Cancer Using DESeq2 [Thesis, Department of Computer Science
and Engineering(CSE), Islamic University of Technology(IUT), Board Bazar, Gazipur,
Bangladesh]. http://localhost:8080/xmlui/handle/123456789/1866

Miller, J. K., Farnham, P. J., Yaswen, P., & Sweeney, C. (2010). ZNF217, a candidate breast cancer oncogene amplified at 20q13, regulates expression of the ErbB3 receptor tyrosine kinase... *Oncogene*, *29*(40), 5500–5510.

O'Donovan, P. (2019). *Identification of driver mutations and tumour evolution in HER2 positive breast cancer* [PhD Thesis, Royal College of Surgeons in Ireland]. https://repository.rcsi.com/articles/thesis/Identification_of_driver_mutations_and_tumour_evolut ion_in_HER2_positive_breast_cancer/10764350

Robinson, J. T., Thorvaldsdóttir, H., Wenger, A. M., Zehir, A., & Mesirov, J. P. (2017). Variant Review with the Integrative Genomics Viewer. *Cancer Research*, *77*(21), e31–e34. https://doi.org/10.1158/0008-5472.CAN-17-0337

Salz, R., Saraiva-Agostinho, N., Vorsteveld, E., van der Made, C. I., Kersten, S., Stemerdink, M., Allen, J., Volders, P.-J., Hunt, S. E., Hoischen, A., & 't Hoen, P. A. C. (2023). SUsPECT: A pipeline for variant effect prediction based on custom long-read transcriptomes for improved clinical variant annotation. *BMC Genomics*, *24*(1), 305. https://doi.org/10.1186/s12864-023-09391-5

Stradella, A., Gargallo, P., Cejuela, M., Petit, A., Bosch-Schips, J., Carbonell, P., Recalde, S., Vethencourt, A., Fernandez-Ortega, A., Falo, C., Gil-Gil, M., Vázquez, S., Obadia, V., Villanueva-Vázquez, R., Soler-Monsó, T., Calabria, I., & Pernas, S. (2022). Genomic characterization and tumor evolution in paired samples of metaplastic breast carcinoma. *Modern Pathology*, *35*(8), 1066–1074. https://doi.org/10.1038/s41379-022-01017-7

Tang, W., Zhang, F., Byun, J. S., Dorsey, T. H., Yfantis, H. G., Ajao, A., Liu, H., Pichardo, M. S., Pichardo, C. M., Harris, A. R., Yang, X. R., Figueroa, J. D., Sayed, S., Makokha, F. W., & Ambs, S. (2023). Population-specific Mutation Patterns in Breast Tumors from African American, European American, and Kenyan Patients. *Cancer Research Communications*, *3*(11), 2244–2255. https://doi.org/10.1158/2767-9764.CRC-23-0165

Vendrell, J. A., Thollet, A., Nguyen, N. T., Ghayad, S. E., Vinot, S., Bièche, I., Grisard, E., Josserand, V., Coll, J.-L., Roux, P., Corbo, L., Treilleux, I., Rimokh, R., & Cohen, P. A. (2012). ZNF217 Is a Marker of Poor Prognosis in Breast Cancer That Drives Epithelial–Mesenchymal Transition and Invasion. *Cancer Research*, *72*(14), 3593–3606. https://doi.org/10.1158/0008-5472.CAN-11-3095

Voutsadakis, I. A. (2020). 8p11.23 Amplification in Breast Cancer: Molecular Characteristics, Prognosis and Targeted Therapy. *Journal of Clinical Medicine*, *9*(10), Article 10. https://doi.org/10.3390/jcm9103079

Wang, S., Wang, C., Liu, O., Hu, Y., Li, X., & Lin, B. (2022). miRNA-651-3p regulates EMT in ovarian cancer cells by targeting ZNF703 and via the MEK/ERK pathway. *Biochemical and Biophysical Research Communications*, *619*, 76–83. https://doi.org/10.1016/j.bbrc.2022.06.005

Wang, Y., Ma, C., Yang, X., Gao, J., & Sun, Z. (2024). ZNF217: An Oncogenic Transcription
Factor and Potential Therapeutic Target for Multiple Human Cancers. *Cancer Management and*
Research. https://www.tandfonline.com/doi/abs/10.2147/CMAR.S431135

Zhang, G., Wang, J., Yang, J., Li, W., Deng, Y., Li, J., Huang, J., Hu, S., & Zhang, B. (2015).
Comparison and evaluation of two exome capture kits and sequencing platforms for variant
calling. *BMC Genomics*, *16*(1), 581. https://doi.org/10.1186/s12864-015-1796-6

Zhang, X., Mu, X., Huang, O., Wang, Z., Chen, J., Chen, D., & Wang, G. (2022). ZNF703
promotes triple-negative breast cancer cells through cell-cycle signaling and associated with poor
prognosis. *BMC Cancer*, *22*(1), 226. https://doi.org/10.1186/s12885-022-09286-w

Zhang, X., Mu, X., Huang, O., Xie, Z., Jiang, M., Geng, M., & Shen, K. (2013). Luminal Breast
Cancer Cell Lines Overexpressing ZNF703 Are Resistant to Tamoxifen through Activation of
Akt/mTOR Signaling. *PLOS ONE*, *8*(8), e72053. https://doi.org/10.1371/journal.pone.0072053

536