1

AAV delivery of RNA editing machinery rescues SUDEP and seizure phenotype in a mouse model of Dravet Syndrome

Ellie M. Chilcott[1], Juan F. Antinao Diaz[1], Amanda Almacellas Barbanoj[2], Marc Moore[3], Anna Keegan[1], Zakary Waddington[1], Vicky Fang[1], Stephanie Schorge[4], Gabriele Lignani[2], Simon N Waddington[1] & Rajvinder Karda[1*]

Affiliations

[1] EGA Institute for Women's Health, University College London, 86-96 Chenies Mews, London, WC1E 6HX, UK

[2] Queen Square Institute of Neurology, Queens Square, University College London, London, WC1N 3BG, UK

[3] School of Health and Life Sciences, National Horizon Centre, Teesside University, 38 John Dixon Ln, Darlington, DL1 1HG, UK

[4] Division of Biosciences, Medical Science Building, University College London, London WC1E 6BT

*Corresponding author: Dr Rajvinder Karda

      Maternal and Fetal Medicine Department

      Institute For Women's Health

      University College London

      86-96 Chenies Mews

      WC1E 6HX

      London

      r.karda@ucl.ac.uk

ABSTRACT

Dravet syndrome (DS) is a severe childhood genetic epilepsy, caused by *de novo* heterozygous mutations in *SCN1A*, resulting in a loss-of-function of the voltage-gated sodium ion channel, Nav1.1. Nav1.1 is expressed in the brain, and at a lower level, in the heart. DS manifests in the first year of life. Patients exhibit tonic-clonic seizures, febrile seizures, cognitive decline, developmental delays, ataxia, and sudden unexpected death from epilepsy (SUDEP). We have developed a novel AAV-F mediated CRISPR-Cas-inspired RNA targeting system (CIRTS) preclinical treatment to increase endogenous *Scn1a* and ameliorate the disease phenotype in a clinically-relevant heterozygous loss-of-function mouse model of DS. We designed novel guide RNAs (gRNAs) to target the long non-coding RNA, (or natural antisense transcript) of *Scn1a* to increase the expression of *Scn1a* mRNA in DS mice. We show that intracerebroventricular and intravenous administration of AAV-F-CIRTS-gRNA9 to target the brain and the heart to neonatal *Scn1a*$^{+/-}$ mice resulted in a significant increase in survival, and a reduction in SUDEP, febrile seizures and seizure duration. These findings provide proof-of-concept evidence that an AAV-F-CIRTS mediated therapy hold promise as a potential treatment for DS.

INTRODUCTION

Dravet syndrome (DS) is a severe childhood genetic epilepsy, with an incidence rate of 1 in 15,400-40,900 live births worldwide[1, 2]. The disease typically manifests within the first year of life accompanied by tonic-clonic seizures, febrile and afebrile seizures, episodes of status epilepticus, cognitive impairment, ataxia and unfortunately, ~15% of patients experience sudden unexpected death in epilepsy (SUDEP)[3–5]. >90% of patients harbour *de novo* loss-of-function variants in the *SCN1A* gene which encodes an α-subunit of a voltage-gated ion channel, Nav1.1[6]. In humans, Nav1.1 expression has been detected in the hippocampus (including the

dentate gyrus, CA3, and CA2 regions), layers V and VI of the cortex, the granular and molecular

layers, as well as in the deep nuclei and Purkinje cells of the cerebellum*(7)* and the

heart. Previous studies have explored altered cardiac function in individuals with DS and in

mouse models, suggesting a potential contribution to SUDEP*(8–11)*.

Current anti-seizure medications, including valproate, stiripentol, fenfluramine and

cannabidiol can reduce the seizure frequency*(12)*, but the disease manifestation continues. Thus,

there remains a significant unmet need to establish an effective long-term treatment for DS.

Adeno-associated viral (AAV) vectors, specifically AAV9, have been used extensively

for many preclinical*(13, 14)* and clinical trials*(15, 16)*. However, the high doses of AAV9

required to achieve efficacious peripheral delivery has also caused liver toxicity and thrombotic

microangiopathy*(16, 17)*. As such, there is a need to develop novel capsids that enable efficient

therapeutic delivery at a lower dose, avoiding immunotoxicity associated with the gene therapy

vector. Novel AAV capsids targeting the central nervous system (CNS) have been generated*(18,*

19). One example of this is AAV-F, a novel capsid derived from AAV9, which has shown a

greater CNS transduction efficiency relative to the parental AAV9 capsid*(18)* and has been used

successfully in a preclinical study to treat a severe mitochondrial disorder*(20)*.

RNA targeting technologies have emerged as therapeutic, inspired by successes in

oligonucleotide based therapies*(21, 22)*. Targeting RNA is advantageous over DNA as it can

influence gene expression without permanent alterations to the host genome and modulate

protein expression regardless of the specific mutation. Furthermore, DNA and RNA targeting

systems such as Cas9 and Cas13 are built from bacterial proteins and have shown to induce an

immune response *in vivo(23)*. A new RNA editing technology has been developed to target RNA

sequences, referred to as CRISPR-Cas-inspired RNA targeting system (CIRTS)*(24)*. CIRTS is

built from human proteins and previous validation studies *in vitro* and *in vivo* have shown the

CIRTS system to modulate protein expression by RNA editing *(24–26)*, although until now this

has not been used in a therapeutic setting. In this study, we used a CIRTS construct consisting of

a ssRNA binding protein (β-defensin 3), RNA hairpin-binding protein (TBP6.7) and effector

protein (YTHDF2)*(24)* to down-regulate target RNA expression.

Here, we used the AAV-F-mediated CIRTS technology and designed novel guide RNAs

(gRNAs) to edit a *Scn1a* specific long non-coding RNA (lncRNA) of the natural antisense

transcript (NAT), which normally functions to negatively regulate *Scn1a* expression*(27)*. In

doing so, we aimed to 'inhibit the inhibitor', leading to an upregulation of *Scn1a* mRNA (Fig.

1A). We screened 21 gRNAs *in vitro* and selected two lead candidate gRNAs (gRNA9 & 21)

which significantly increased endogenous *Scn1a* mRNA and shared highest homology to the

human lncRNA sequence. We used a clinically-relevant DS heterozygous (*Scn1a$^{+/-}$*) model,

which recapitulates much of the human disease, including spontaneous and febrile seizures, and

SUDEP*(28, 29)*. We administered our candidates AAV-F-CIRTS-gRNA 9 & 21 vectors to

neonatal DS mice via intracerebroventricular (ICV) and intravenous (IV) administration to target

the brain and the heart. Candidate AAV-F-CIRTS-gRNA9 resulted in significant increase in

survival over 100 days post-injection. In addition, febrile and spontaneous seizure assessment of

DS mice injected with AAV-F-CIRTS-gRNA9 showed a significant reduction in seizure severity

and seizure duration. Molecular analysis revealed a significant up-regulation of endogenous

Scn1a mRNA in treated AAV-F-CIRTS-gRNA9 DS mice compared to untreated DS control

mice. Our work demonstrates for the first time, a potential AAV-F mediated RNA editing

approach to treat patients with DS.

RESULTS

Design and *in vitro* validation of novel gRNAs targeting *Scn1a* lncRNA

We firstly validated the CIRTS machinery and consistent with previous studies*(24)*, AAV-CIRTS-gRNAluc was able to significantly reduce luciferase expression (Fig. S1). We designed 21 novel gRNA sequences (20-30mer) to target the *Scn1a* lncRNA (Fig. 1A) (here referred to as *Scn1anat*), which were cloned into the AAV-CIRTS plasmid (Fig. 1B). We validated our gRNA designs *in vitro*, in differentiated mouse Neuro2a cells and assessed the effects on endogenous *Scn1a* mRNA expression. Transient transfection of AAV-CIRTS-gRNA constructs onto differentiated Neuro2a cells resulted in a significant increase of endogenous *Scn1a* mRNA by 4 out of 21 AAV-CIRTS-gRNA constructs (Fig. 1C). Constructs containing gRNAs 9, 11, 15 and 21 led to average *Scn1a* mRNA fold changes of 1.89 (P=0.0016), 2.15 (P=0.0015), 2.12 (P=0.022) and 2.56 (P<0.0001), respectively. gRNAs 9 and 21 were selected for the *in vivo* study as they showed the highest homology to the *SCN1A* lncRNA (*SCN1ANAT*; 93.3% and 100%, respectively) (Fig. 1C).

AAV-F-mediated RNA editing in DS mice significantly improves survival

In this study we crossed the males from 129Sv-*Scn1a^{tm1Kea}*/Mmjax model*(28, 29)* with wild-type CD1 females to generate F2 129Sv-*Scn1a^{tm1Kea}*/Mmjax x CD1 heterozygote females, which were then bred to wild-type C57BL/6J males to obtain experimental litters (*Scn1a^{+/-}*). This enabled a larger litter size. *Scn1a^{+/-}* mice exhibit spontaneous seizures and SUDEP with an overall survival of 38.2% over 100 days of development (Fig. S2A), febrile seizures (Fig. S2D) and a reduction in *Scn1a* gene expression, particularly in the cortex at P50 (P=0.0078; Fig. S2E),

consistent with previous studies*(22, 28, 30)*. RNA-seq of *Scn1a*$^{+/+}$ and *Scn1a*$^{+/-}$ cortical tissue at

P50 confirmed 140 up- and 41 down-regulated genes (Fig. S3A and B), with significant down-

regulation of *Scn1a* (-0.77Log2FC, P=5.85x10^{-6}; Fig. S2C). Molecular function gene ontology

(GO) analysis of the differentially expressed genes in *Scn1a*$^{+/-}$ mice reveal up-regulation in terms

related to neuropeptide receptor activity, monoatomic ion transport and voltage-gated sodium

channel activity (Fig. S3D).

Using this DS mouse model, we sought to determine the optimal capsid and gRNA

combination by generating AAV-F and AAV9 vectors for AAV-CIRTS-gRNA9 and AAV-

CIRTS-gRNA21, using AAV9 as a gold-standard comparison. At P0, *Scn1a*$^{+/-}$ mice received an

intracerebroventricular (ICV) and intravenous (IV) delivery of either AAV-F (7x10^{10} vg/pup) or

AAV9 (7x10^{11} vg/pup), each encoding either gRNA9 or 21 to determine the effects on survival.

This study was fully blinded and randomised (Fig. 2A). AAV-F vectors were injected at 10-fold

lower dose than AAV9, we have previously demonstrated a greater CNS transduction of AAV-F

at a 10-fold lower dose after neonatal ICV delivery*(20)*. Furthermore, here in this study, we

demonstrated a superior biodistribution in the brain and visceral organs with AAV-F compared

to titre matched AAV9 (Fig S4) after neonatal IV delivery. In addition, AAV-F displayed a

greater transduction of neurons and astrocytes compared to AAV9 (Fig. S5 & S6).

PBS-injected *Scn1a*$^{+/+}$ and *Scn1a*$^{+/-}$ mice served as controls. *Scn1a*$^{+/-}$ mice treated with AAV-F-

CIRTS-gRNA9 had a significant (P=0.0235) increase in overall survival to 70.4% at P100

compared to 36.4% survival at P100 in PBS-treated *Scn1a*$^{+/-}$ controls (Fig. 2A). In contrast,

AAV9-CIRTS-gRNA9 only showed an overall survival of 26.5% (P=0.8439, Fig. 2A). The

AAV-F-CIRTS-gRNA21 treatment group also appeared less efficient, showing only a 12.5%

survival (P=0.0509, Fig. S8A), while AAV9-CIRTS-gRNA21 had a 50% survival at P100

(P=0.3715, Fig. S8A). Weights and open field analysis of neither AAV-F- or AAV9-CIRTS-

gRNA9/21 treated mice differed from PBS $Scn1a^{+/+}$ and $Scn1a^{+/-}$ control group (Fig. 2D, Fig.

S7A & Fig. S8C and D). From these data, AAV9 RNA editing at a 10-fold higher titre than

AAV-F failed to improve survival we therefore, proceeded with AAV-F-CIRTS-gRNA9 for the

remaining experiment.

We interrogated dose-range efficacy of AAV-F-CIRTS-gRNA9 with one higher and one

lower dose. High dose (1.75×10^{11} vg/pup) resulted in 65.5% survival (P=0.0732, Fig. 3A), whilst

low dose (1.75×10^{10} vg/pup) resulted in 44.4% survival, compared to PBS-treated $Scn1a^{+/-}$ mice

over 100 days (P=0.5815 and P=0.8556, respectively, Fig. 3A). Neither of the additional AAV-

F-CIRTS-gRNA9 doses were superior to our originally tested dose of 7×10^{10} vg/pup.

Furthermore, no doses showed difference in weight or open field assessment compared to PBS

$Scn1a^{+/+}$ control group (Fig. 3B, C and Fig. S9A).

AAV-F-CIRTS-gRNA9 reduces seizure burden in DS mice

To assess the effects of this optimal dose of AAV-F-CIRTS-gRNA9 therapy on febrile

and spontaneous seizures in $Scn1a^{+/-}$ mice, we administered a separate cohort of $Scn1a^{+/-}$ mice

with neonatal ICV and IV delivery (7×10^{10} vg/pup). We also administered control groups AAV-

F-CIRTS-gRNAluc (7×10^{10} vg/pup) and PBS to neonatal $Scn1a^{+/-}$ mice via ICV and IV delivery

(Fig. 4A). Febrile seizure assessment revealed a significant improvement in seizure severity

determined by the Racine scale*(31)* (Fig. 4B), in AAV-F-CIRTS-gRNA9 treated $Scn1a^{+/-}$ mice

compared to PBS $Scn1a^{+/-}$ control group (P<0.0001; Fig. 4B), and AAV-F-CIRTS-gRNAluc

$Scn1a^{+/-}$ control group (P=0.0111; Fig. 4B).

Electroencephalogram (EEG) recordings of spontaneous seizures were recorded for 15

days between P30-45 and the results revealed a trend to reduction of seizure burden in AAV-F-

CIRTS-gRNA9 mice, compared to a control AAV-F-CIRTS-gRNAluc. Total numbers of

seizures (9.7±13 vs 4.1±7.3; P=0.4121; Fig. S10A) and average numbers of seizures per day

(0.78±0.93 vs 0.33±0.48; P=0.43; Fig. S10B) were reduced. In the mice which displayed

seizures, the average seizure duration was significantly reduced in the treated AAV-F-CIRTS-

gRNA9 $Scn1a^{+/-}$ group compared to control AAV-F-CIRTS-gRNAluc $Scn1a^{+/-}$ group (P=0.0279;

Fig. 4D), as well as a trend for reduction in cumulative seizure time (P=0.0640; Fig. S10C).

AAV-F-CIRTS-gRNA9 normalises molecular signature of DS

To determine the effects of AAV-F-CIRTS-gRNA9 on endogenous $Scn1a$ mRNA, we

assessed gene expression in the cortex and the heart of treated $Scn1a^{+/-}$ mice. Raw wild-type

$Scn1a$ gene expression levels were restored in both the cortex and heart of AAV-F-CIRTS-

gRNA9 mice (P=0.0072 and P=0.0002 vs PBS $Scn1a^{+/-}$; Fig. 5A and B).

Vector copy number (VCN) assessing the presence of the effector protein $YTHDF2$

genomic sequence, showed an increase in both $Scn1a^{+/-}$ and $Scn1a^{+/+}$ mice treated with AAV-F-

CIRTS-gRNA9 treated mice in the cortex at P100 compared to both PBS (P=0.0014) and AAV-

F-CIRTS-gRNAluc (P=0.0146) groups $Scn1a^{+/+}$ (VCN 0.35±0.44; Fig. 5C). No difference was

observed in the heart and the liver (VCN 0.19±0.14 and 0.04±0.05, respectively; Fig. 5C).

$YTHDF2$ mRNA expression mimicked the pattern of vector copy number with highest

expression in the cortex in AAV-F-CIRTS-gRNA9 $Scn1a^{+/-}$ mice. No significance was observed

in the heart and the liver (Fig. S11A).

In vivo safety of AAV-F-CIRTS-gRNA9

Both AAV-F-CIRTS-gRNA9 and AAV-F-CIRTS-gRNAluc were injected to *Scn1a*$^{+/+}$

mice at the two highest dosages 1.75×10^{11} vg/pup and 7×10^{10} vg/pup via neonatal ICV and IV to

determine any toxicological findings. No adverse effects on survival or weight were seen (Fig.

S12A and B). Open field analyses revealed hyperactivity in the total distance of *Scn1a*$^{+/+}$ AAV-

F-CIRTS-gRNAluc mice treated with dose 1.75×10^{11} vg/pup at P18 and P55 (P=0.0003,

P=0.0217; Fig. S12C), also with an increase in maximum speed at P55 (P=0.0099; Fig. S12D)

compared to PBS *Scn1a*$^{+/+}$ control group.

Immunohistochemical analysis to detect astrocytes (GFAP) and macrophages, including

kupffer cells, in the brain and liver respectively was performed post-AAV-F-CIRTS-gRNA9

treatment. No increases in reactive astrocytes were seen in the brains of treated *Scn1a*$^{+/+}$ or

Scn1a$^{+/-}$ mice (Fig. 6A and B). An increase in the presence of macrophages/kupffer cells in the

liver was seen in only the *Scn1a*$^{+/+}$ mice treated with AAV-F-CIRTS-gRNAluc (P=0.0025), this

was not observed in mice of either genotype treated with the same dose of AAV-F-CIRTS-

gRNA9 (Fig. 6A and C).

In vivo editing specificity of AAV-F-CIRTS-gRNA9

To identify gRNA-specific changes in gene expression, we conducted RNA-seq on

cortical tissue 50 days post-injection and compared AAV-F-CIRTS-gRNAluc and AAV-F-

CIRTS-gRNA9 *Scn1a*$^{+/-}$ mice. Only 9 differentially expressed genes were found; 3 up-regulated

and 6 down-regulated (Fig. 7A and B). Of note, inflammation-associated genes were down-

regulated including *Igkc* (-7.27Log2FC, P=0.0417), *Cxcl13* (-7.35Log2FC, P=0.0417) and *Adgre*

(-1.86Log2FC, P=0.0019) and the only driver GO term changed across the comparison, was the

down-regulation of CXCR5 chemokine receptor binding (Fig. 7C; P=0.0499). None of the *Scna*

genes, which encode other voltage-gated sodium ion channels, were differentially expressed in

the gRNA9 group compared to gRNAluc.

DISCUSSION

Here, we provide preclinical proof-of-concept data demonstrating that CIRTS-mediated

RNA editing can mitigate the phenotype of a severe neurological disorder, Dravet syndrome. A

single administration of AAV-F-CIRTS-gRNA9 by combined ICV and IV delivery to *Scn1a*$^{+/-}$

DS mice induces destruction of *Scn1anat* and completely rescues endogenous *Scn1a* mRNA

expression in both the cortex and the heart. This was shown to improve disease manifestation by

reducing febrile seizure severity and significantly reducing SUDEP, with minimal off-target

effects; whilst being compromised entirely of human components.

We designed the gRNAs used in this study with the aim of restoring wild-type *Scn1a*

expression levels, without leading to supraphysiological expression. Notably, modest 2.6- and

1.9-fold increase of *Scn1a* expression *in vitro* were achieved with gRNA21 and gRNA9

respectively. We sought to combine these with AAV-F, an engineered capsid, shown to have a

more efficient CNS transduction profile than AAV9 in rodents[32], human neurons[32] and

NHPs[19]. Furthermore, we have validated this with neonatal delivery both by ICV[20] and IV

delivery. By packaging AAV-CIRTS-gRNA9 into AAV-F and AAV9 we demonstrated vastly

different therapeutic effects with 70.4% and 26.5% survival, respectively, strikingly this was

achieved with AAV-F being administered at a 10-fold lower dose. However, gRNA21 delivered

by either AAV-F or AAV9 did not reveal a significant improvement in survival. We tested a

varying dose of AAV-F-CIRTS-gRNA9 and determined that 7×10^{10} vg/pup achieved a

significant survival of 70.4% at P100, interestingly increasing the dose further to 1.75×10^{11}

vg/pups showed a reduced survival of 65.5% over 100 days. These findings suggest a narrow

therapeutic dose ranging window for a successful treatment for DS. This is particularly important

in the context of *SCN1A*, as gain-of-function *SCN1A* variants, which are known to cause familial

hemiplegic migraine type 3 and a severe infantile developmental and encephalopathy (distinct

from DS) with associated movement disorder*(33)*. Therefore, future studies require a more

refined dose ranging evaluation of AAV-F with gRNA9 or gRNA21 to determine the optimal

therapeutic dose, with appropriate levels of physiological *Scn1a* and no behavioural side effects.

250   As a main symptom of DS is seizures, we sought to assess the seizure activity in our

AAV-F-CIRTS-gRNA9 treated *Scn1a*$^{+/-}$ DS mice via febrile seizure assessment and EEG

monitoring for 15 days (P30-45). We delivered a vehicle and control vector AAV-F-CIRTS-

gRNAluc to *Scn1a*$^{+/-}$ DS mice as a comparators. Febrile seizure assessment revealed a

significant reduction in seizure severity determined by the Racine scale in AAV-F-CIRTS-

gRNA9 treated *Scn1a*$^{+/-}$ mice compared to PBS and AAV-F-CIRTS-gRNAluc *Scn1a*$^{+/-}$ control

group. EEG seizure monitoring was performed for AAV-F-CIRTS-gRNA9 and AAV-F-CIRTS-

gRNAluc treated *Scn1a*$^{+/-}$ DS mice. Our results showed that we could significantly reduce the

seizure duration in our treated *Scn1a*$^{+/-}$ DS mice. From these data, we observed a variability in

the AAV-F-CIRTS-gRNAluc *Scn1a*$^{+/-}$ DS mice, as some of these mice did not display seizures

during the EEG recording. Which may partly be due to the variable penetrance of spontaneous

seizures in this model*(34)*. This was a limitation in our study and future studies, will require

further power calculations to establish the appropriate number of *Scn1a*$^{+/-}$ mice required for EEG

seizure monitoring.

Molecular assessment revealed a significant increase in endogenous *Scn1a* mRNA in the

cortex and the heart of treated AAV-F-CIRTS-gRNA9 *Scn1a*$^{+/-}$ mice. Immunohistochemical

analysis in the brain of AAV-F-CIRTS-gRNA9 *Scn1a*$^{+/-}$ and *Scn1a*$^{+/+}$ mice showed no signs of

toxicity. Liver toxicity was only observed in AAV-F-CIRTS-gRNAluc *Scn1a*$^{+/+}$ treated group,

suggesting potential guide-specific side-effects of AAV-F-CIRTS-gRNAluc vector.

To determine the off-target effects of our gRNA9, we conducted RNA-seq on cortical

brain tissues for AAV-F-CIRTS-gRNAluc and AAV-F-CIRTS-gRNA9 *Scn1a*$^{+/-}$ treated mice.

The data revealed inflammation-associated genes including *Igkc(35)*, *Cxcl13(36)* and *Adgre(37)*

were down-regulated in the AAV-F-CIRTS-gRNA9 treated group. Previously, up-regulation of

the CXCL13-CXCR5 signalling pathway has been associated with intractable epilepsy*(36)*,

suggesting that our therapy may be able to reduce some molecular signatures of epilepsy

pathogenesis. Furthermore, our data suggest no off-target effects of the gRNA9 coupled with the

CIRTS machinery.

Several approaches in preclinical development have used genetic therapies specifically in

inhibitory interneurons using *SCN1A* gene replacement with split-intein dual AAVs*(38)* and

Adenoviral vectors*(39)*, CRISPR/dCas9 activation of *SCN1A(40, 41)* and transcription factor-

mediated *SCN1A* up-regulation*(30)*. While these approaches target the primary pathology of

Dravet syndrome (DS), our strategy complements these alternative therapeutic interventions by

also considering effects in less well-studied cell populations, such as cardiomyocytes*(10)*. Here,

we have shown that our AAV-F-CIRTS-gRNA9 can increase *Scn1a* mRNA in the cortex and the

heart.

SCN1ANAT has been detected in surgical brain tissue specimens from patients (1-19

years of age) undergoing resection to treat refractory, non-DS epilepsy*(42)*. Previous evidence

has shown that targeting *Scn1anat* using small oligonucleotides termed AntagoNATs were able

to reduce seizures and normalise neuronal activity in DS knock-in mice*(27)*. Furthermore, we

have also shown that these AntagoNATs can be vectorised with AAV9, which also demonstrates

a significant improvement in DS phenotype*(43)*. Here, we also are targeting *Scn1anat*, but

instead of blocking the interaction between this and the *Scn1a* mRNA, we are inducing the

degradation of *Scn1anat*, with similar phenotypic improvement. These findings demonstrate the

feasibility of using AAV mediated RNA editing to modulate *Scn1a* expression, highlighting its

potential as a therapeutic for DS.

Genetic therapies for DS have progressed to clinical evaluation. These include the

antisense oligonucleotide (ASO) zorevunersen, currently under investigation in Phase 1/2a

MONARCH and ADMIRAL trials, which involve repeated intrathecal administration*(44)*.

Encoded Therapeutics has developed an AAV9-based transcription factor*(30)* approach

specifically targeting GABAergic interneurons, now being assessed in a Phase 1/2 clinical trial

utilizing ICV delivery alone*(45)*. Similar to Encoded Therapeutics' strategy, our AAV-F-CIRTS-

gRNA9 platform offers a one-time treatment; however, it confers the additional advantage of

enabling broader targeting of CNS cell populations expressing Nav1.1. Furthermore, the

inclusion of IV administration is intended to achieve efficient cardiac transduction, addressing

evidence implicating altered cardiac electrophysiology*(8, 11)* in contributing to SUDEP.

Our study presented limitations. Some of the *Scn1a*$^{+/-}$ DS mice did not present seizures

during EEG recording and therefore, for future studies we will require further power

calculations. The future studies required have been detailed in this discussion.

In conclusion, we have developed and characterized an AAV-F-CIRTS RNA editing

therapeutic strategy for the treatment of DS. Incorporation of gRNA targeting *Scn1anat*

successfully modulated *Scn1a* expression and significantly improved survival, seizure severity

and duration in *Scn1a*$^{+/-}$ mice. These findings provide compelling proof-of-concept evidence

supporting the potential of this approach as a therapeutic intervention for DS.

MATERIALS AND METHODS

Study design

This study was designed to investigate if AAV-CIRTS-mediated RNA editing to induce

the destruction of *Scn1anat* could lead to therapeutic benefit in a clinically relevant DS mouse

model. The following design elements were included in this study. N=3 independent experiments

were conducted for *in vitro* gRNA screening experiments with at least n=3 technical replicates

per condition. Mice were assigned randomly to study groups, with at least one control (either

PBS or AAV-F-CIRTS-gRNAluc-treated) *Scn1a*$^{+/-}$ mouse per litter and remaining *Scn1a*$^{+/+}$ and

Scn1a$^{+/-}$ littermates mice treated with AAV-F-CIRTS-gRNA9/21 or AAV9-CIRTS-gRNA9/21

treated mouse. Prospectively selected endpoints of these studies were survival, weight, open

field, longitudinal gene expression and neuropathological analyses. Sample sizes for *in vivo*

experiments were not pre-defined before the outset of data collection as no preliminary data were

available to estimate effect sizes and thus be able to complete power analyses. *In vivo* studies

analysing seizures used *Scn1a*$^{+/-}$ littermates treated with AAV-F-CIRTS-gRNAluc as a non-

targeting guide as a control for AAV-F-CIRTS-gRNA9. At termination of the in-life period for

all *in vivo* experiments, mice were assigned a numeric code and investigators were blinded to

treatment and genotype whilst analysing tissue.

gRNA design and cloning

333   The *Scn1anat* target sequence was inputted into sOligo software

334 (https://sfold.wadsworth.org/cgi-bin/soligo.pl), varying the parameter for gRNA length. Folding

335 temperature was set to 37°C and ionic conditions set to 1M NaCl with no divalent ions.

336 Resulting outputs were screened for those with less than -8.0 Gibbs free energy with positional

337 clusters identified. Potential gRNA sequences were manually mapped to RNA secondary

338 structures produced using mFold software (http://www.unafold.org/mfold/applications/rna-

339 folding-form.php) to determine qualities of each gRNA such as the number of termini and

340 percentage of gRNA sequence in open conformation. gRNAs were cloned into the AAV-CIRTS

341 plasmid (Addgene plasmid #132544), using gBlocks from Integrated DNA Technologies and

342 InFusion cloning (Takara Bio). The AAV-CIRTS plasmid contains a cytomegalovirus promoter

343 (CMV), driving ssRNA binding protein β-defensin 3, RNA hairpin-binding protein (TBP6.7),

344 effector protein (YTHDF2) and U6 promoter driving gRNA (referred to as AAV-CIRTS in the

345 main text, Fig. 1B).

346 ***In vitro* analyses**

347   For luciferase assays; HEK293T cells ($5x10^4$ per well of 24-well plate) were transfected

348 with 0.8μg of an AAV plasmid containing an Spleen focus forming virus ubiquitous promoter

349 (SFFV) driving luciferase linked to green fluorescent reporter gene with a T2A linker

350 peptide*(46)* (referred to as AAV-SFFV-*luciferase*-T2A-*eGFP* in the main text) ± 0.8μg AAV-

351 CIRTS-gRNAluc using Lipofectamine 2000 (Thermo Fisher Scientific). Cells were lysed in

352 Luciferase Cell Culture Lysis Reagent (Promega) at 72h post-transfection, with half the lysate

353 used to determine protein concentration using the DC protein assay (BioRad). The remaining

354 half of the lysate was used for luminometry measured in a FLUOstar® Omega microplate

reader (BMG Labtech). Luciferase expression was normalised to protein concentration within each sample.

For *Scn1anat* gRNA screening; Neuro2a cells ($3x10^4$ per well of 24-well plate) were differentiated in DMEM (Thermo Fisher Scientific), 2% FCS (Sigma-Aldrich), 0.5mM cAMP (Merck) and 20μM Retinoic Acid (Merck). At 6DIV, cells were transfected with 0.8μg AAV-SFFV-*luciferase*-T2A-*eGFP* or AAV-CIRTS-gRNA1-21 using Lipofectamine 2000 (Thermo Fisher Scientific). Cells were lysed in TRIzol reagent (Thermo Fisher Scientific) at 5d post-transfection. RNA was extracted and qPCR used to determine changes in *Scn1a* gene expression (protocols described below).

AAV vector production

Recombinant AAV vectors were produced by triple-transfection[47] of AAVpro 293T cells (Takara Bio) with AAV-CIRTS, adenovirus helper plasmid (Harvard University) and either AAV9 (University of Pennsylvania; Addgene #112865) or AAV-F (Harvard University, Addgene #166921) capsid plasmids using PEImax (Polysciences Inc.). HPLC using POROS CaptureSelect AAVX Resin (Thermo Fisher Scientific) was used to purify resulting vector before being titrated by qPCR.

Animals

All procedures were performed in accordance with UK Home Office Animals (Scientific Procedures) Act 1986 under PPLs 70/14300, PP8023218 & PP9223137. All mice were housed in IVCs with enrichment, under a 12 hour:12 hour light–dark cycle and had access to food and water *ad libitum*. The mouse model of Dravet syndrome used in this project is derived from the 129Sv-*Scn1a*[tm1Kea]/Mmjax model[28, 29] (Jackson Laboratory) that has been extensively used in

DS pre-clinical research. Males from this strain were bred with wild-type CD1 females to obtain

a F1 129Sv-*Scn1a^{tm1Kea}*/Mmjax x CD1 generation. F1xF1 crosses were bred to obtain F2 129Sv-

Scn1a^{tm1Kea}/Mmjax x CD1 heterozygote females which were bred to wild-type C57BL/6J males

to obtain experimental litters (here referred to as *Scn1a^{+/-}* in the text). Humane endpoints of

>15% weight loss, observation of two seizures and/or clear signs of illness (piloerection,

hunched posture and laboured breathing).

At post-natal day 0 (P0), *Scn1a^{+/-}* pups were injected by bilateral intracerebroventricular

(ICV; 5 μL per hemisphere, using previously published coordinates*(48)*) and intravenous (IV; 25

μL via superficial temporal vein*(49)*) routes of administration with either PBS, AAV9 (total dose

7×10^{11} vg/pup (2×10^{11} vg ICV + 5×10^{11} vg IV)) or AAV-F (total dose 1.75×10^{10} vg/pup (5×10^9

vg ICV + 1.25×10^{10} vg IV), 7×10^{10} vg/pup (2×10^{10} vg ICV + 5×10^{10} vg IV) and 1.75×10^{11}

vg/pup (5×10^{10} vg ICV + 1.25×10^{11} vg IV)) using a 33-gauge Hamilton needle (VWR). For the

biodistribution study AAV9 (1.9×10^{10} and 1.9×10^{11} vg/pup) and AAV-F (1.9×10^{10} vg/pup) was

delivered intravenously via superficial temporal vein*(49)*). After the procedure, mice were

returned to their dams. Weight and survival were monitored until endpoint.

Open field analysis with 15 minutes free locomotion was completed at P18, P38, P55 and

P75 using SMART v3 software (Harvard Panlab). At endpoint, tissues were collected by snap-

freezing or fixing in 4% PFA.

EEG recordings

AAV-F-CIRTS-gRNAluc and AAV-F-CIRTS-gRNA9 treated *Scn1a^{+/-}* mice underwent

EEG recordings from P30-P45 using wireless electrocorticogram (ECoG) transmitters (Cat. No.

A3048P2-AA-C37-D, single-channel transmitter, Open Source Instruments), as previously

described*(43)*.

RNA/DNA extraction, gene expression and VCN

RNA extraction was completed using the PureLink kit (Thermo Fisher Scientfic), DNA

extraction was completed using DNeasy Blood & Tissue kit (Qiagen), as previously

described*(43)*. The following primers and probes were used to quantify gene expression, titrate

AAV vectors and perform vector copy number analyses. *Scn1a;* forward:

TCAGAGGGAAGCACAGTAGAC, reverse: TTCCACGCTGATTTGACAGCA, probe:

CCAGAAGAAACCCTTGAGCCCGAA (fluorophore: ABY, quencher: QSY, Thermo Fisher

Scientific). *Gapdh*; forward: ACGGCAAATTCAACGGCAC, reverse:

TAGTGGGGTCTCGCTCCTGG, probe: TTGTCATCAACGGGAAGCCCATCA (fluorophore:

VIC, quencher: QSY, Thermo Fisher Scientific). *YTHDF2*; forward:

CCTCCATTGGCTTCTCCTATTC, reverse: GTTGCTCAGCTGTCCATAAGA, probe:

TTAAGTAGGGCATGGCTGTGTCACC (fluorophore: FAM, quencher: ZEN / Iowa Black™

FQ, Integrated DNA Technologies).

The same sequence *Gapdh* primers and probe, and *Scn1a* primers were used for ddPCR

analysis, but with the *Scn1a* probe fluorophore was adapted from ABY to FAM. 5μL of each

cDNA (equivalent to 85 pg) was loaded per reaction with 1X ddPCR Multiplex Supermix (Bio-

Rad) and 4 mM DTT. ddPCR analysis was performed using the BioRad QXOne in the UCL

NeuroGTx Vector Core Facility. The Facility is supported by the NIHR Great Ormond Street

Hospital Biomedical Research Centre. The views expressed are those of the author(s) and not

necessarily those of the NHS, the NIHR or the Department of Health.

Immunoperoxide staining

40 μm coronal sections cut using a Micron HM 430 freezing microtome (Thermo Fisher

Scientific) were analysed using previously published protocols[43]. Neuropathological analyses

used the following primary antibodies; anti-GFAP (Proteintech 16825-1-AP; 1:2000) and anti-

CD68 (BioRad MCA1957; 1:100). Images were acquired using a Leica Microsystems

stereoscopic microscope (DM4000), Leica DFC7000T camera and Leica LAX X software (all

Leica Microsystems UK). Images were background corrected using FIJI (ImageJ) and

thresholding software (Image-Pro Premier v10; Media Cybernetics) was used to quantify

expression.

Immunofluorescence staining

40 μm coronal sections were used[43]. Immunolabeling was performed on free-floating

sections using antibodies against GFP (Abcam; ab290), NeuN (Merek; MAB377) and GFAP

(Proteintech; 16825-1-AP)[20]. Images were assessed using fluorescence microscope (Leica

DFC7000 7) using LAS X Microscope Software (Leica Microsystems). The percentage of GFP

positive neurons and astrocytes were analysed using the CellProfiler Software (Broad Institute

2021). The images were overlayed using ImageJ.

RNA-seq

Total RNA was extracted from the cortex of $Scn1a^{+/+}$ and $Scn1a^{+/-}$ P50 mice that were

controls, or had been treated with, AAV-F-CIRTS-gRNAluc or AAV-F-CIRTS-gRNA9.

Untreated control samples were processed by UCL Genomics using a KAPA mRNA

HyperPrep and NextSeq2000 P2 100c configuration with single-end reads at 25M reads/sample.

N=8 mice per genotype were submitted, but due to a pooling error, $Scn1a^{+/+}$ Untreated_5 could

not be included in the analysis.

AAV-CIRTS-treated samples (n=4 per group) were processed by Azenta Life Sciences

using Illumina NovaSeq with paired-end reads in 2x150 bp configuration. Trimmomatic v.0.36

was used to trim sequence reads to remove adapter sequences before reads were mapped to *Mus*

musculus GRCm38 reference genome using the STAR aligner v.2.5.2b. Unique gene counts

were extracted using featureCounts (Subread package v.1.5.2).

Both RNA-seq analyses then used DESeq2*(50)* to determine differential gene expression

analysis. Untreated samples were subjected to an absolute Log2 fold change > 0.7, whilst AAV-

CIRTS comparisons used an absolute Log2 fold change > 1. In both analyses, genes with an

adjusted P value < 0.05 were called differentially expressed genes. A gene ontology analysis was

performed on the statistically significant set of genes by implementing enrichGO and

clusterProfiler R packages*(51)* to cluster the set of genes based on their molecular function and

determine their statistical significance.

Statistical analyses

GraphPad Prism (version 10.3.1, Boston) was used for statistical analyses. All continuous

variables are presented as mean±SD in graphs. *In vitro* data were analysed using an unpaired,

two-tailed t-test for luciferase assay or one-way ANOVA with Dunnett's multiple comparisons

for gRNA screening. Survival was determined using a Log-rank (Mantel-Cox) test. Weights

were initially screened for differences with one-way ANOVA with Dunnett's multiple

comparisons for group-group comparisons and a mixed-effect analysis was completed with

analysis at each individual timepoint. Groups were excluded in mixed-effect analyses from the

timepoint that only a single mouse remained. Open field was analysed using two-way ANOVA

with Dunnett's multiple comparisons. Racine scaling of febrile seizures were contingency

analysed with Fisher's exact test. A mixed-effects analysis was used to compare overall EEG

seizure burden, with either Mann-Whitney or unpaired, two-tailed t-test for individual seizure

readouts. One-way and two-way ANOVAs were used to analyse all molecular data.

Supplementary Materials

PDF file includes;

Fig S1 to S12 supplementary figures

REFERENCES

1. E. Chilcott, J. A. Díaz, C. Bertram, M. Berti, R. Karda, Genetic therapeutic advancements for Dravet Syndrome.
Epilepsy & Behavior **132**, 108741 (2022).

2. A. Strzelczyk, L. Lagae, J. Wilmshurst, A. Brunklaus, P. Striano, F. Rosenow, S. Schubert-Bast, Dravet
syndrome: a systematic literature review of the illness burden. *Epilepsia Open* (2023).

3. A. Brunklaus, S. M. Zuberi, Dravet syndrome—From epileptic encephalopathy to channelopathy. *Epilepsia* **55**,
979–984 (2014).

4. M. Wolff, C. Cassé-Perrot, C. Dravet, Severe myoclonic epilepsy of infants (Dravet syndrome): natural history
and neuropsychological findings. *Epilepsia* **47 Suppl 2**, 45–48 (2006).

5. A. Brunklaus, E. Pérez-Palma, I. Ghanty, J. Xinge, E. Brilstra, B. Ceulemans, N. Chemaly, I. de Lange, C.
Depienne, R. Guerrini, D. Mei, R. S. Møller, R. Nabbout, B. M. Regan, A. L. Schneider, I. E. Scheffer, A.-S.
Schoonjans, J. D. Symonds, S. Weckhuysen, M. W. Kattan, S. M. Zuberi, D. Lal, Development and Validation of a
Prediction Model for Early Diagnosis of SCN1A-Related Epilepsies. *Neurology* **98**, e1163–e1174 (2022).

6. S. Gataullina, O. Dulac, From genotype to phenotype in Dravet disease. *Seizure* **44**, 58–64 (2017).

7. W. R. J. Whitaker, R. L. M. Faull, H. J. Waldvogel, C. J. Plumpton, P. C. Emson, J. J. Clare, Comparative
distribution of voltage-gated sodium channel proteins in human brain. *Molecular Brain Research* **88**, 37–53 (2001).

8. A. T. Berg, K. Coffman, D. Gaebler-Spira, Dysautonomia and functional impairment in rare developmental and
epileptic encephalopathies: the other nervous system. *Developmental Medicine & Child Neurology* **63**, 1433–1440
(2021).

9. F. Kalume, R. E. Westenbroek, C. S. Cheah, F. H. Yu, J. C. Oakley, T. Scheuer, W. A. Catterall, Sudden
unexpected death in a mouse model of Dravet syndrome. *J Clin Invest* **123**, 1798–1808 (2013).

10. S. Mishra, V. Reznikov, V. A. Maltsev, N. A. Undrovinas, H. N. Sabbah, A. Undrovinas, Contribution of
sodium channel neuronal isoform Nav1.1 to late sodium current in ventricular myocytes from failing hearts. *The*
Journal of Physiology **593**, 1409–1427 (2015).

11. D. S. Auerbach, J. Jones, B. C. Clawson, J. Offord, G. M. Lenk, I. Ogiwara, K. Yamakawa, M. H. Meisler, J. M.
Parent, L. L. Isom, Altered Cardiac Electrophysiology and SUDEP in a Model of Dravet Syndrome. *PLOS ONE* **8**,
e77843 (2013).

12. A. Strzelczyk, S. Schubert-Bast, A Practical Guide to the Treatment of Dravet Syndrome with Anti-Seizure
Medication. *CNS Drugs* **36**, 217–237 (2022).

13. G. Massaro, C. N. Z. Mattar, A. M. S. Wong, E. Sirka, S. M. K. Buckley, B. R. Herbert, S. Karlsson, D. P.
Perocheau, D. Burke, S. Heales, A. Richard-Londt, S. Brandner, M. Huebecker, D. A. Priestman, F. M. Platt, K.
Mills, A. Biswas, J. D. Cooper, J. K. Y. Chan, S. H. Cheng, S. N. Waddington, A. A. Rahim, Fetal gene therapy for
neurodegenerative disease of infants. *Nature Medicine* (2018), doi:10.1038/s41591-018-0106-7.

14. J. Ng, S. Barral, C. de la Fuente Barrigon, G. Lignani, F. A. Erdem, R. Wallings, R. Privolizzi, G. Rossignoli, H.
Alrashidi, S. Heasman, E. Meyer, A. Ngoh, S. Pope, R. Karda, D. Perocheau, J. Baruteau, N. Suff, J. A. Diaz, S.
Schorge, J. Vowles, L. R. Marshall, S. A. Cowley, S. Sucic, M. Freissmuth, J. R. Counsell, R. Wade-Martins, S. J.
R. Heales, A. A. Rahim, M. Bencze, S. N. Waddington, M. A. Kurian, Gene therapy restores dopamine transporter
expression and ameliorates pathology in iPSC and mouse models of infantile parkinsonism. *Science Translational*
Medicine **13**, 1–13 (2021).

15. J. R. Mendell, S. Al-Zaidy, R. Shell, W. D. Arnold, L. R. Rodino-Klapac, T. W. Prior, L. Lowes, L. Alfano, K.
Berry, K. Church, J. T. Kissel, S. Nagendran, J. L'Italien, D. M. Sproule, C. Wells, J. A. Cardenas, M. D. Heitzer,
A. Kaspar, S. Corcoran, L. Braun, S. Likhite, C. Miranda, K. Meyer, K. D. Foust, A. H. M. Burghes, B. K. Kaspar,
Single-Dose Gene-Replacement Therapy for Spinal Muscular Atrophy. *New England Journal of Medicine* **377**,
1713–1722 (2017).

16. K. A. Strauss, M. A. Farrar, F. Muntoni, K. Saito, J. R. Mendell, L. Servais, H. J. McMillan, R. S. Finkel, K. J.
Swoboda, J. M. Kwon, C. M. Zaidman, C. A. Chiriboga, S. T. Iannaccone, J. M. Krueger, J. A. Parsons, P. B. Shieh,
S. Kavanagh, S. Tauscher-Wisniewski, B. E. McGill, T. A. Macek, Onasemnogene abeparvovec for presymptomatic
infants with two copies of SMN2 at risk for spinal muscular atrophy type 1: the Phase III SPR1NT trial. *Nat Med* **28**,
1381–1389 (2022).

17. D. Chand, F. Mohr, H. McMillan, F. F. Tukov, K. Montgomery, A. Kleyn, R. Sun, S. Tauscher-Wisniewski, P.
Kaufmann, G. Kullak-Ublick, Hepatotoxicity following administration of onasemnogene abeparvovec (AVXS-101)
for the treatment of spinal muscular atrophy. *Journal of Hepatology* **74**, 560–566 (2021).

18. K. S. Hanlon, J. C. Meltzer, T. Buzhdygan, M. J. Cheng, M. Sena-Esteves, R. E. Bennett, T. P. Sullivan, R.
Razmpour, Y. Gong, C. Ng, J. Nammour, D. Maiz, S. Dujardin, S. H. Ramirez, E. Hudry, C. A. Maguire, Selection
of an Efficient AAV Vector for Robust CNS Transgene Expression. *Molecular Therapy - Methods and Clinical*
Development **15**, 320–332 (2019).

19. A. Beharry, Y. Gong, J. C. Kim, K. S. Hanlon, J. Nammour, K. Hieber, F. Eichler, M. Cheng, A. Stemmer-
Rachamimov, K. M. Stankovic, D. B. Welling, C. Ng, C. A. Maguire, The AAV9 Variant Capsid AAV-F Mediates
Widespread Transgene Expression in Nonhuman Primate Spinal Cord After Intrathecal Administration. *Hum Gene*
Ther **33**, 61–75 (2022).

20. A. Keegan, O. Cetin, E. M. Chilcott, J. F. Antinao Diaz, S. Eaton, S. N. Waddington, J. R. Counsell, S. Rahman,
R. Karda, Next Generation AAV-F Capsid gene therapy rescues disease pathology in a model of Pyruvate
Dehydrogenase Complex Deficiency. *bioRxiv* , 2025.07.30.667478 (2025).

21. L. E. Burbano, M. Li, N. Jancovski, P. Jafar-Nejad, K. Richards, A. Sedo, A. Soriano, B. Rollo, L. Jia, E.
Gazina, S. Piltz, F. Adikusuma, P. Q. Thomas, F. Rigo, C. A. Reid, S. Maljevic, S. Petrou, Antisense
oligonucleotide therapy for KCNT1 encephalopathy, 2020.11.12.379164 (2020).

22. Z. Han, C. Chen, A. Christiansen, S. Ji, Q. Lin, C. Anumonwo, C. Liu, S. C. Leiser, I. Aznarez, G. Liau, L. L.
Isom, *Antisense oligonucleotides increase Scn1a expression and reduce seizures and SUDEP incidence in a mouse*
model of Dravet syndrome (2020; http://stm.sciencemag.org/), p. 6100.

23. D. W. Morgens, M. Wainberg, E. A. Boyle, O. Ursu, C. L. Araya, C. K. Tsui, M. S. Haney, G. T. Hess, K. Han,
E. E. Jeng, A. Li, M. P. Snyder, W. J. Greenleaf, A. Kundaje, M. C. Bassik, Genome-scale measurement of off-
target activity using Cas9 toxicity in high-throughput screens. *Nat Commun* **8**, 15178 (2017).

24. S. Rauch, E. He, M. Srienc, H. Zhou, Z. Zhang, B. C. Dickinson, Programmable RNA-Guided RNA Effector
Proteins Built from Human Parts. *Cell* **178**, 122-134.e12 (2019).

25. S. Rauch, K. A. Jones, B. Dickinson, Small Molecule-Inducible RNA-Targeting Systems for Temporal Control
of RNA Regulation in Vivo (2020), doi:10.26434/chemrxiv.11673711.v1.

26. S. U. Madugalle, W.-S. Liau, Q. Zhao, X. Li, H. Gong, P. R. Marshall, A. Periyakaruppiah, E. L. Zajaczkowski,
L. J. Leighton, H. Ren, M. R. B. Musgrove, J. W. A. Davies, G. Kim, S. Rauch, C. He, B. C. Dickinson, B.
Fulopova, L. N. Fletcher, S. R. Williams, R. C. Spitale, T. W. Bredy, Synapse-Enriched m6A-Modified Malat1
Interacts with the Novel m6A Reader, DPYSL2, and Is Required for Fear-Extinction Memory. *J. Neurosci.* **43**,
7084–7100 (2023).

27. J. Hsiao, T. Y. Yuan, M. S. Tsai, C. Y. Lu, Y. C. Lin, M. L. Lee, S. W. Lin, F. C. Chang, H. Liu Pimentel, C.
Olive, C. Coito, G. Shen, M. Young, T. Thorne, M. Lawrence, M. Magistri, M. A. Faghihi, O. Khorkova, C.
Wahlestedt, Upregulation of Haploinsufficient Gene Expression in the Brain by Targeting a Long Non-coding RNA
Improves Seizure Phenotype in a Model of Dravet Syndrome. *EBioMedicine* **9**, 257–277 (2016).

28. A. R. Miller, N. A. Hawkins, C. E. Mccollom, J. A. Kearney, Mapping genetic modifiers of survival in a mouse
model of Dravet syndrome. *Genes, Brain and Behavior* **13**, 163–172 (2014).

29. A. M. Mistry, C. H. Thompson, A. R. Miller, C. G. Vanoye, A. L. George,, J. A. Kearney, Strain- and age-
dependent hippocampal neuron sodium currents correlate with epilepsy severity in Dravet syndrome mice.
Neurobiology of Disease **65**, 1–11 (2014).

30. A. Tanenhaus, T. Stowe, A. Young, J. McLaughlin, R. Aeran, I. W. Lin, J. Li, R. Hosur, M. Chen, J. Leedy, T.
Chou, S. Pillay, M. C. Vila, J. A. Kearney, M. Moorhead, A. Belle, S. Tagliatela, Cell-Selective Adeno-Associated
Virus-Mediated SCN1A Gene Regulation Therapy Rescues Mortality and Seizure Phenotypes in a Dravet Syndrome
Mouse Model and Is Well Tolerated in Nonhuman Primates. *Hum Gene Ther* **33**, 579–597 (2022).

31. J. Van Erum, D. Van Dam, P. P. De Deyn, PTZ-induced seizures in mice require a revised Racine scale.
Epilepsy & Behavior **95**, 51–55 (2019).

32. K. S. Hanlon, J. C. Meltzer, T. Buzhdygan, M. J. Cheng, M. Sena-Esteves, R. E. Bennett, T. P. Sullivan, R.
Razmpour, Y. Gong, C. Ng, J. Nammour, D. Maiz, S. Dujardin, S. H. Ramirez, E. Hudry, C. A. Maguire, Selection
of an Efficient AAV Vector for Robust CNS Transgene Expression. *Mol Ther Methods Clin Dev* **15**, 320–332
(2019).

33. A. Brunklaus, T. Brünger, T. Feng, C. Fons, A. Lehikoinen, E. Panagiotakaki, M.-A. Vintan, J. Symonds, J.
Andrew, A. Arzimanoglou, S. Delima, J. Gallois, D. Hanrahan, G. Lesca, S. MacLeod, D. Marjanovic, A. McTague,
N. Nuñez-Enamorado, E. Perez-Palma, M. Scott Perry, K. Pysden, S. J. Russ-Hall, I. E. Scheffer, K. Sully, S. Syrbe,
U. Vaher, M. Velayutham, J. Vogt, S. Weiss, E. Wirrell, S. M. Zuberi, D. Lal, R. S. Møller, M. Mantegazza, S.
Cestèle, The gain of function SCN1A disorder spectrum: novel epilepsy phenotypes and therapeutic implications.
Brain **145**, 3816–3831 (2022).

34. N. A. Hawkins, L. L. Anderson, T. S. Gertler, L. Laux, A. L. George Jr., J. A. Kearney, Screening of
conventional anticonvulsants in a genetic mouse model of epilepsy. *Annals of Clinical and Translational Neurology*
4, 326–339 (2017).

35. J. P. Pandey, P. J. Nietert, A. M. Namboodiri, C. Kimball, P. A. Flume, Roles of immunoglobulin GM and KM
allotypes and Fcγ receptor 2 A genotypes in humoral immunity to a conserved microbial polysaccharide in
pulmonary diseases. *Genes Immun* **26**, 91–95 (2025).

36. R. Li, L. Ma, H. Huang, S. Ou, J. Yuan, T. Xu, X. Yu, X. Liu, J. Yang, Y. Chen, X. Peng, Altered Expression of
CXCL13 and CXCR5 in Intractable Temporal Lobe Epilepsy Patients and Pilocarpine-Induced Epileptic Rats.
Neurochem Res **42**, 526–540 (2017).

37. H.-H. Lin, C.-C. Hsiao, C. Pabst, J. Hébert, T. Schöneberg, J. Hamann, in *Advances in Immunology*, G Protein-
Coupled Receptors in Immune Response and Regulation. A. K. Shukla, Ed. (Academic Press, 2017), vol. 136, pp.
163–201.

38. J. K. Mich, J. Ryu, A. D. Wei, B. B. Gore, R. Guo, A. M. Bard, R. A. Martinez, E. M. Luber, J. Liu, Y. M.
Bishaw, R. J. Christian, L. M. Oliveira, N. Miranda, J.-M. Ramirez, J. T. Ting, E. S. Lein, B. P. Levi, F. K. Kalume,
Interneuron-specific dual-AAV SCN1A gene replacement corrects epileptic phenotypes in mouse models of Dravet
syndrome. *Science Translational Medicine* **17**, eadn5603 (2025).

39. A. Ricobaraza, M. Bunuales, M. Gonzalez-Aparicio, S. Fadila, M. Rubinstein, I. Vides-Urrestarazu, J. Banderas,
N. Sola-Sevilla, R. Sanchez-Carpintero, J. L. Lanciego, E. Roda, A. Honrubia, P. Arnaiz, R. Hernandez-Alcoceba,
Preferential expression of SCN1A in GABAergic neurons improves survival and epileptic phenotype in a mouse
model of Dravet syndrome. *J Mol Med* **101**, 1587–1601 (2023).

40. G. Colasante, G. Lignani, S. Brusco, C. Di Berardino, J. Carpenter, S. Giannelli, N. Valassina, S. Bido, R. Ricci,
V. Castoldi, S. Marenna, T. Church, L. Massimino, G. Morabito, F. Benfenati, S. Schorge, L. Leocani, D. M.
Kullmann, V. Broccoli, dCas9-Based Scn1a Gene Activation Restores Inhibitory Interneuron Excitability and
Attenuates Seizures in Dravet Syndrome Mice. *Molecular Therapy* **28**, 235–253 (2020).

41. T. Yamagata, M. Raveau, K. Kobayashi, H. Miyamoto, T. Tatsukawa, I. Ogiwara, S. Itohara, T. K. Hensch, K.
Yamakawa, CRISPR/dCas9-based Scn1a gene activation in inhibitory neurons ameliorates epileptic and behavioral
phenotypes of Dravet syndrome model mice. *Neurobiology of Disease* **141**, 104954 (2020).

42. M. F. Schneider, M. Vogt, J. Scheuermann, V. Müller, A. H. L. Fischer-Hentrich, T. Kremer, S. Lugert, F.
Metzger, M. Kudernatsch, G. Kluger, T. Hartlieb, S. Noachtar, C. Vollmar, M. Kunz, J. C. Tonn, R. Coras, I.
Blümcke, C. Pace, F. Heinen, C. Klein, H. Potschka, I. Borggraefe, Brain expression profiles of two SCN1A
antisense RNAs in children and adolescents with epilepsy. *Transl Neurosci* **15**, 20220330 (2024).

43. J. Antinao Diaz, E. Chilcott, A. Almacellas Barbanoj, A. Keegan, A. McTague, J. H. Cross, S. Schorge, G.
Lignani, S. N. Waddington, R. Karda, AAV9-mediated targeting of natural antisense transcript as a novel treatment
for Dravet Syndrome. *bioRxiv* , 2024.09.27.615323 (2024).

44. Z. Han, C. Chen, A. Christiansen, S. Ji, Q. Lin, C. Anumonwo, C. Liu, S. C. Leiser, Meena, I. Aznarez, G. Liau,
L. L. Isom, Antisense oligonucleotides increase *Scn1a* expression and reduce seizures and SUDEP incidence in a
mouse model of Dravet syndrome. *Sci. Transl. Med.* **12**, eaaz6100 (2020).

45. Encoded Therapeutics Announces US IND Clearance and Australian CTA Approval for Dravet Syndrome Gene
Therapy Candidate ETX101*Encoded Therapeutics, Inc.* (available at https://encoded.com/press-releases/encoded-
therapeutics-announces-us-ind-clearance-and-australian-cta-approval-for-dravet-syndrome-gene-therapy-candidate-
etx101/).

46. Z. Liu, O. Chen, J. B. J. Wall, M. Zheng, Y. Zhou, L. Wang, H. Ruth Vaseghi, L. Qian, J. Liu, Systematic
comparison of 2A peptides for cloning multi-genes in a polycistronic vector. *Sci Rep* **7**, 2193 (2017).

47. X. Xiao, J. Li, R. J. Samulski, Production of high-titer recombinant adeno-associated virus vectors in the absence
of helper adenovirus. *J Virol* **72**, 2224–2232 (1998).

48. J.-Y. Kim, S. D. Grunke, Y. Levites, T. E. Golde, J. L. Jankowsky, Intracerebroventricular Viral Injection of the
Neonatal Mouse Brain for Persistent and Widespread Neuronal Transduction. *Journal of Visualized Experiments
(JoVE)* , e51863 (2014).

49. S. E. Gombash Lampe, B. K. Kaspar, K. D. Foust, Intravenous Injections in Neonatal Mice. *J Vis Exp* , 52037
(2014).

50. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. *Genome Biology* **15**, 550 (2014).

51. T. Wu, E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, L. Zhou, W. Tang, L. Zhan, X. Fu, S. Liu, X. Bo, G.
Yu, clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The Innovation* **2**, 100141 (2021).

Acknowledgments: We would like to thank A. Rahim and J. Baruteau for the use of their Home
Office PPL. We would like to thank, R.Towns, M. Bandol, N. Foster, N. Marzouki and the staff
from Biological Services for their help with maintenance of the DS colony at University College
London. Figures 1A and 2A were created in BioRender. Antinao Diaz, J. (2025)
https://BioRender.com/k81r069.

Funding: This work was supported by Great Ormond Street Hospital Children Charity and

Dravet Syndrome UK Charity (V4720 and V4919 to R.K), Medical Research Council

Development Pathway Funding Scheme (MR/Z505201/1 to R.K), LifeArc (P2020-

0008 and P2023-0011 to R.K), Therapeutic Acceleration Support (TAS), UCL (to R.K),

GOSH/Spark Research Grant V4019 (to G.L.), Medical Research Council Programme Grant

(MR/V034758/1 to G.L. and S.S.), Epilepsy Research UK Emerging Leader Fellowship (F1701

to G.L.), Medical Research Council New Investigator Project Grant (MR/S011005/1 to G.L.).

Author contributions: Conceptualization: R.K and S.N.W. Methodology: R.K, M.M and

E.M.C. Investigation: R.K, E.M.C, J.F.A.D, A.K, Z.W, V.F, A.A.B, G.L. Visualization: R.K and

E.M.C. Funding acquisition: R.K, S.S, M.M, S.N.W and G.L. Project administration: R.K.

Writing – original draft: R.K and E.M.C. Writing – review & editing: all authors.

Competing interests: RK, EMC, MM are co-inventors on patent application "Therapy for

Dravet Syndrome". RK, SS & SNW have consultancy agreements with Rocket Pharma. G.L. and

S.S. have equity in a company that aims to bring epilepsy gene therapy to the clinic.

Data and materials availability: All data associated with this study are present in the paper in

Supplementary Material. Requests for data should be addressed to R.K.

Figures

Fig. 1. **Novel gRNAs increase in endogenous *Scn1a in vitro.*** (A) Schematic illustrating

mechanism of action of CIRTS to increase endogenous *Scn1a* expression through targeting of

the lncRNA. (B) Schematic diagram of the AAV-CIRTS-gRNA construct. (C) 4 novel AAV-

CIRTS-gRNA constructs significantly increased endogenous *Scn1a* mRNA after transient

transfection of differentiated mouse Neuro2a cells. The homology of these gRNAs to the human

SCN1ANAT sequence are indicated on the graph as a %, with gRNAs ranked from least to most

homologous. One-way ANOVA with Dunnett's multiple comparisons.

Fig. 2. Neonatal AAV-F-CIRTS-gRNA9 RNA editing *in vivo* increases survival of *Scn1a*$^{+/-}$

DS mice. *Scn1a*$^{+/-}$ mice were administered either AAV-F-CIRTS-gRNA9 (7×10^{10} vg/pup) or

AAV9-CIRTS-gRNA9 (7×10^{11} vg/pup) at P0 by intracerebroventricular (ICV) and intravenous

(IV) injection, with PBS-treated *Scn1a*$^{+/-}$ and *Scn1a*$^{+/+}$ littermate controls. (A) Experimental

timeline. (B) Kaplan-Meier survival curves of treated mice with percentage survival shown. Log-

rank (Mantel-Cox) test. (C) Weights of treated mice. One-way ANOVA with Dunnett's multiple

comparison. (D) Total distance read-outs from open field analyses at 4 different timepoints.

Two-way ANOVA with Dunnett's multiple comparisons.

Fig. 3. Dosage study of AAV-F-CIRTS-gRNA9 in *Scn1a*$^{+/-}$ mice. AAV-F-CIRTS-gRNA9 was

administered at 2 additional doses (1.75×10^{10} and 1.75×10^{11} vg/pup) and was compared to the

original 7×10^{10} vg/pup cohort presented in Fig. 2. (A) Kaplan-Meier survival curves of treated

mice with percentage survival shown. Log-rank (Mantel-Cox) test. (B) Weights of treated mice.

AAV-F-CIRTS-gRNA9 1.75×10^{11} vg/pup mice were significantly different from $Scn1a^{+/-}$ PBS

mice at P70 & P80-95 (P=0.0214-0.0451). Mixed-effects analysis with Geisser-Greenhouse

correction. (C) Total distance read-outs from open field analyses at 4 different timepoints. Two-

way ANOVA with Dunnett's multiple comparisons.

Fig. 4. Reduction of seizure burden following AAV-F-CIRTS-gRNA9 treatment. (A) Febrile

seizure probability curves. (B) Severity of observed seizures ranked by Racine scale. Fisher's

exact test. (C) Heatmap with coloured squares indicating seizure activity, monitored by EEG.

Crosses indicate the death of an animal during EEG recording. (D) Average seizure duration in

mice which presented with seizures, during EEG. Unpaired, two-tailed t-test.

Fig. 5. Restoration of *Scn1a* gene expression in the cortex and heart of AAV-F-CIRTS-

gRNA9 treated *Scn1a*$^{+/-}$ mice. *Scn1a* gene expression in the (A) cortex and (B) heart between

$Scn1a^{+/+}$ and $Scn1a^{+/-}$ mice (P20-100 tissues) treated with non-targeting AAV-F-CIRTS-

gRNAluc vector compared to AAV-F-CIRTS-gRNA9 $Scn1a^{+/-}$ mice. One-way ANOVA with

Dunnett's multiple comparisons. (C) Vector copy number analysis in P100 cortex, heart and

liver. Two-way ANOVA with Tukey's multiple comparisons.

Fig. 6. AAV-F-CIRTS-gRNA9 does not lead to toxicity *in vivo*. (A) Representative brain and

liver immunohistochemistry images. Scale bar = 100 μm. (B) Quantification of reactive

astrocytes (GFAP immunohistochemistry) in the brain of treated mice; CA1 = CA1 region of the

hippocampus, CA3 = CA3 region of the hippocampus, VPM = Ventral posteromedial nucleus of

the thalamus. Two-way ANOVA with Šídák's multiple comparisons. (C) Quantification of

macrophages (CD68 immunohistochemistry) in the liver of treated mice. One-way ANOVA with

Dunnett's multiple comparisons.

Fig. 7. RNA-seq depicting differential gene expression following AAV-F-CIRTS-gRNA9

treatment. $Scn1a^{+/-}$ cortical tissue analysed at P50. Comparison of AAV-F-CIRTS-gRNAluc

and AAV-F-CIRTS-gRNA9 mice. (A) Volcano plot showing the main differentially expressed

genes (DEGs). (B) Heatmap showing top up- and down-regulated genes per comparison,

presented as z-score, ranked by P value. Driver gene ontology (GO) term classification of the

molecular function of (C) down-regulated DEGs. No molecular function GO terms for up-

regulated genes were identified.

Fig. S1. *In vitro* validation of AAV-CIRTS machinery. AAV-CIRTS-gRNAluc significantly

down-regulated luciferase expression in HEK293T cells after transient co-transfection with

AAV-SFFV-*luciferase*-T2A-*eGFP* plasmid. Luciferase expression normalised to protein

concentration within lysate. Unpaired, two-tailed t-test.

Fig. S2. Characterisation of untreated *Scn1a*$^{+/-}$ DS mice. (A) Kaplan-Meier survival curves of

untreated mice over 100 days of development with percentage survival shown. Log-rank

(Mantel-Cox) test. (B) Weights of untreated mice. (C) Total distance and maximum speed read-

outs from open field analyses at 4 different timepoints. Two-way ANOVA with Dunnett's

multiple comparisons. (D) Induction of febrile seizures with temperature of seizure onset. (E)

Scn1a gene expression in 4 brain regions. Two-way ANOVA with Šídák's multiple comparisons

within each brain region. (F) Quantification of reactive astrogliosis (GFAP

immunohistochemistry) in the brain of untreated mice; CA1 = CA1 region of the hippocampus,

CA3 = CA3 region of the hippocampus, VPM = Ventral posteromedial nucleus of the thalamus.

Fig. S3. RNA-Seq analysis of untreated *Scn1a*$^{+/-}$ DS mice. *Scn1a*$^{+/+}$ and *Scn1a*$^{+/-}$ cortical tissue

analysed at P50. (A) Volcano plot showing the main differentially expressed genes. (B) Heatmap

showing top 10 up- and down-regulated genes, presented as z-score, ranked by P value. (C)

Confirmation of *Scn1a* down-regulation in *Scn1a*$^{+/-}$ mice with normalised read count, (-

0.77Log2FC P=5.85×10^{-6}). (D) Driver gene ontology (GO) term classification of the molecular

function of up-regulated DEGs. No molecular function GO terms for down-regulated genes were

identified.

Fig. S4. Increased biodistribution and transgene expression of AAV-F-CAG-GFP vector

compared to AAV9. Wild-type C57BL/6J mice were administered either AAV-F-CAG-GFP

(1.9×10^{10} vg/pup) or AAV9- CAG-GFP (1.9×10^{10} vg/pup or 1.9×10^{11} vg/pup) at P0 by

intravenous (IV) injection, with uninjected littermate controls. (A) Experimental timeline. (B) *Ex*

vivo images of organs from injected mice. Scale bar = 2mm. (C) Quantification of vector copy

number in discreet brain regions and visceral organs. One-way ANOVA with Dunnett's multiple

comparisons.

Fig. S5. GFP biodistribution in the brain of AAV-F and AAV9-CAF-GFP vectors.
Immunofluorescence stain showing GFP positive cells in the olfactory bulb (OFB), cortex, hippocampus and cerebellum. Scale bar shown on the images.

Fig. S6. Co-localisation and quantification of neuron and astrocyte targeting of AAV-F. (A) Representative images co-localisation of GFP and NeuN. Images were taken at 400X magnification. Scale bar, 100µm. **(B)** Quantification of GFP and NeuN positive cells. One-way ANOVA with Dunnett's multiple comparisons. **(C)** Representative images co-localisation of GFP and GFAP. Images were taken at 400X magnification. Scale bar = 100µm. **(D)** Quantification of GFP and GFAP positive cells. One-way ANOVA with Dunnett's multiple comparisons.

Fig. S7. Maximum speed of AAV-F/9-CIRTS-gRNA9 mice and survival curve for AAV-F-CIRTS-gRNA9. (A) Open field maximum speed assessment at 4 different timepoints. Two-way ANOVA with Dunnett's multiple comparisons. (B) Kaplan-Meier survival curves of treated mice with percentage survival shown. Log-rank (Mantel-Cox) test.

Fig. S8. Neonatal AAV-F/9-CIRTS- gRNA21 treatment. $Scn1a^{+/-}$ mice were administered either AAV-F-CIRTS-gRNA21 at $7x10^{10}$ vg/pup or AAV-9-CIRTS-gRNA21 at $7x10^{11}$ vg/pup, with PBS-treated $Scn1a^{+/-}$ and $Scn1a^{+/+}$ littermate controls. (A) Kaplan-Meier survival curves of treated mice with percentage survival shown. Log-rank (Mantel-Cox) test. (B) Weights of treated mice. AAV-9-CIRTS-gRNA21 was significantly different from $Scn1a^{+/-}$ PBS mice at P90 only (P=0.0390), but only female mice were alive in the treatment group at this time (100% female at

P100 vs 25% female in *Scn1a*$^{+/-}$ PBS). Mixed-effects analysis with Geisser-Greenhouse

correction. (C) Total distance and (D) maximum speed read-outs from open field analyses at 4

different timepoints. Two-way ANOVA with Dunnett's multiple comparisons.

Fig. S9. Maximum speed of AAV-F-CIRTS-gRNA9 dosage. 4 different timepoints. Two-way

ANOVA with Dunnett's multiple comparisons.

Fig. S10. Seizure monitoring in AAV-F-CIRTS-gRNA9 treated DS mice. (A) Total seizures

measured in 15 days. Mann-Whitney test. (B) Average number of seizures per day. Unpaired,

two tailed t-test (C) Cumulative seizure duration. Unpaired, two tailed t-test.

Fig. S11. *YTHDF2* gene expression analysis. P100 cortex, heart and liver. Two-way ANOVA

with Tukey's multiple comparisons.

Fig. S12. AAV-F-CIRTS-gRNA9/luc administration to healthy mice. *Scn1a*$^{+/+}$ mice were

administered either AAV-F-CIRTS-gRNA9 or AAV-F-CIRTS-gRNAluc at the optimum

therapeutic dose 7×10^{10} vg/pup, or 1 dose higher at 1.75×10^{11} vg/pup to assess signs of toxicity,

compared to PBS littermates. (A) Kaplan-Meier survival curves of treated mice with percentage

survival shown. Log-rank (Mantel-Cox) test. (B) Weights of treated mice. One-way ANOVA

with Dunnett's multiple comparison. (C) Total distance and (D) maximum speed read-outs from

open field analyses at 4 different timepoints. Two-way ANOVA with Dunnett's multiple

comparisons.

Figure 1

Figure 2

Figure 3

A

Percentage Survival vs **Day**

100%
70.4%
65.5%
44.4%
36.4%

P=0.0235

B

Weight (g) vs **Day**

▲ Scn1a⁺/⁺ PBS (n=14)
■ Scn1a⁺/⁻ AAVF-CIRTS-gRNA9 (1.75x10¹⁰ vg; n=9)
▲ Scn1a⁺/⁻ AAVF-CIRTS-gRNA9 (1.75x10¹¹ vg; n=14)

◆ Scn1a⁺/⁻ PBS (n=16)
▼ Scn1a⁺/⁻ AAVF-CIRTS-gRNA9 (7x10¹⁰ vg; n=27)

C

Total distance (cm) vs **P18, P38, P55, P75**

P = 0.0107
P = 0.0325

● Scn1a⁺/⁺ PBS (n=14)   ● Scn1a⁺/⁻ PBS (n=15)
● Scn1a⁺/⁻ AAV-F-CIRTS-gRNA9 (1.75x10¹⁰ vg; n=9)   ● Scn1a⁺/⁻ AAV-F-CIRTS-gRNA9 (7x10¹⁰ vg; n=22)
● Scn1a⁺/⁻ AAV-F-CIRTS-gRNA9 (1.75x10¹¹ vg; n=12)

Figure 4

Figure 5

Figure 6

Figure 7

A

Scn1a$^{+/-}$ gRNAluc vs gRNA9

B

Top up- and down-regulated genes

C

Down-regulated molecular function GO terms

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

A — *Scn1a*$^{+/+}$ vs *Scn1a*$^{+/-}$

B — Top up- and down-regulated genes

C — *Scn1a* expression

D — Up-regulated molecular function GO terms

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

Supplementary Figure 7

A

- ● *Scn1a*$^{+/+}$ PBS (n=14)
- ● *Scn1a*$^{+/-}$ PBS (n=15)
- ● *Scn1a*$^{+/-}$ AAV-F-CIRTS-gRNA9 (n=22)
- ● *Scn1a*$^{+/-}$ AAV9-CIRTS-gRNA9 (n=10)

B

- ⊢ *Scn1a*$^{+/+}$ PBS
- ⊢ *Scn1a*$^{+/-}$ PBS
- ⊢ *Scn1a*$^{+/-}$ AAV-F-CIRTS-gRNAluc (n=18)
- ⊢ *Scn1a*$^{+/-}$ AAV-F-CIRTS-gRNA9 (n=27)

Supplementary Figure 8

- ● Scn1a[+/+] PBS (n=14)　── Scn1a[+/-] PBS (n=16)　── Scn1a[+/-] AAVF-CIRTS-gRNA21 (n=8)　── Scn1a[+/-] AAV9-CIRTS-gRNA21 (n=11)

- ● Scn1a[+/+] PBS (n=14)
- ● Scn1a[+/-] PBS (n=15)
- ● Scn1a[+/-] AAV-F-CIRTS-gRNA21 (n=8)
- ● Scn1a[+/-] AAV9-CIRTS-gRNA21 (n=12)

Supplementary Figure 9

- ● *Scn1a*$^{+/+}$ PBS (n=14)
- ● *Scn1a*$^{+/-}$ PBS (n=15)
- ● *Scn1a*$^{+/-}$ AAV-F-CIRTS-gRNA9 (1.75x10^{10} vg; n=9)
- ● *Scn1a*$^{+/-}$ AAV-F-CIRTS-gRNA9 (7x10^{10} vg; n=22)
- ● *Scn1a*$^{+/-}$ AAV-F-CIRTS-gRNA9 (1.75x10^{11} vg; n=12)

Supplementary Figure 10

Supplementary Figure 11

Legend:
- *Scn1a*$^{+/+}$ PBS (n=13)
- *Scn1a*$^{+/+}$ AAV-F-CIRTS-gRNAluc (n=4)
- *Scn1a*$^{+/-}$ AAV-F-CIRTS-gRNA9 (n=9-10)

Supplementary Figure 12

A

B

Legend:
- ▲ *Scn1a*$^{+/+}$ PBS (n=14)
- ▼ *Scn1a*$^{+/+}$ AAV-F-CIRTS-gRNAluc (7x10^{10} vg; n=4)
- ◆ *Scn1a*$^{+/+}$ AAV-F-CIRTS-gRNA9 (7x10^{10} vg; n=10)
- ■ *Scn1a*$^{+/+}$ AAV-F-CIRTS-gRNAluc (1.75x10^{11} vg; n=8)
- ■ *Scn1a*$^{+/+}$ AAV-F-CIRTS-gRNA9 (1.75x10^{11} vg; n=7)

C

Legend:
- ● *Scn1a*$^{+/+}$ PBS (n=14)
- ▼ *Scn1a+/+* AAV-F-CIRTS-gRNAluc (7x10^{10} vg; n=4)
- ■ *Scn1a*$^{+/+}$ AAV-F-CIRTS-gRNA9 (7x10^{10} vg; n=5)
- ◆ *Scn1a+/+* AAV-F-CIRTS-gRNAluc (1.75x10^{11} vg; n=8)
- ▲ *Scn1a*$^{+/+}$ AAV-F-CIRTS-gRNA9 (1.75x10^{11} vg; n=7)

D