Litter biota and quality as drivers of litter decomposition in mature oak forests subject to drought or elevated CO2

Matthew L. Meehan[1,8*], Mathilde Chomel[1,2], Zhabiz Vilkiji[1], Katy J. Faulkner[1,3], Tancredi Caruso[4], A. Robert MacKenzie[5,6], Virginie Baldy[7], Richard D. Bardgett[1,9], David Johnson[1,9]

[1]Department of Earth and Environmental Sciences, The University of Manchester, Manchester, United Kingdom

[2]FiBL France, Research Institute of Organic Agriculture, Eurre, France

[3]Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, United Kingdom

[4]School of Biology and Environmental Science, University College Dublin, Dublin, Ireland.

[5]School of Geography, Earth & Environmental Sciences, University of Birmingham, Birmingham, United Kingdom

[6]Birmingham Institute of Forest Research, University of Birmingham, Birmingham, United Kingdom

[7]IMBE, Aix-Marseille Université, Univ Avignon, CNRS, IRD, Marseille, France

[8]Current address: Natural Resources Canada, Canadian Forest Service, Laurentian Forestry Centre, Québec, Canada

[9]Current address: Lancaster Environment Centre, Lancaster University, Bailrigg, Lancaster, United Kingdom

*Corresponding author's email: matthew.meehan@nrcan-rncan.gc.ca

Submitted as a *Full paper* to New Phytologist

Introduction:1,110 words, Methods: 2,525 words, Results: 1,357 words, Discussion: 1,929 words
Combined word count of sections listed of 6,921 words
Five figures (all published in colour)

Summary (Word count: 195 words)

- Increasing atmospheric CO_2 concentrations alongside more frequent and severe
droughts are key global change factors impacting litter decomposition, and global
carbon and nitrogen cycles. Yet, we have a poor understanding of how these
perturbations impact interactions between litter chemistry and decomposer food
webs.

- We tested how drought and elevated CO_2 concentrations modify litter
decomposition *via* litter chemistry and the decomposer communities using two
separate, long-term manipulative drought or elevated CO_2 field experiments in
mature oak woodlands. Litter bags were deployed in a reciprocal transplant design
whereby control and treated litter was incubated in control and treated plots. We
measured litter mass loss, quality and stoichiometry, moisture content, and
microbial and animal community composition.

- Elevated CO_2 and drought affected litter stoichiometry and quality which in turn
affected decomposition: litter in droughted plots decomposing slower than controls,
while litter from elevated CO_2 plots decomposed slower over the first three harvests.
In addition, we found that litter mass loss and decomposers were related both
directly and indirectly to litter C:N.

- Drought and elevated CO_2 have distinct pathways of influence on litter quality,
decomposers and decomposition which challenges our ability to predict how
combinations of factors influence litter decomposition.

Keywords: BIFoR FACE, Decomposition, Drought, Elevated CO_2, Mesofauna, Litter, Microbial,
O_3HP

Plain Language Summary

Global change factors like drought and elevated atmospheric CO_2 concentrations can impact litter

decompositions, an important ecosystem process, *via* changes to litter properties and the

decomposer community. Examining links between litter properties, decomposers and

decomposition is therefore critical to understand how both drought and elevated CO_2 will affect

nutrient release and cycling of belowground environments.

Introduction (Word count: 1,110)

Increasing atmospheric CO_2 concentrations often modifies leaf anatomy, chemistry and

physiology (Pritchard et al., 1999; Ainsworth and Long, 2005), and at larger geographic scales,

significantly alters global climatic patterns, including the frequency and intensity of extreme

weather events such as droughts (IPCC, 2021). For instance, in 2022, summer drought resulted in

nearly half of Europe's soil moisture levels becoming critically low (Toreti et al., 2022). Both

elevated atmospheric CO_2 and drought have the potential to affect critical components of terrestrial

biogeochemical cycles, including through changes in root inputs and litter decomposition (Chomel

et al., 2019; Deng et al., 2021; Dong et al., 2021; Norby et al., 2024; Xu et al., 2024).   For

decomposition, the impact of elevated atmospheric CO_2 and drought begins prior to litterfall,

because the C:N ratio of leaves and shoots can increase when grown under drought and elevated

atmospheric CO_2. These effects can stem from increased sugar accumulation, decreasing nitrogen

(N) content, and changes to primary and secondary plant metabolites due to water stress (drought)

(Chen et al., 2015; Piper et al., 2017; Orians et al., 2019). For elevated CO_2, N dilution and

increased assimilation of carbon (C) can alter the C:N within leaves (Taub and Wang, 2008; Welti

et al., 2020), although this effect can be dependent on soil nutrient availability (Gardner et al.,

2022). As a result, decomposition is affected by drought and elevated atmospheric CO_2 *via*

changes to leaf and litter quality (Cotrufo and Ineson, 1996; Kasurinen et al., 2006; Quer et al.,

2022; Wilson et al., 2022). Forest ecosystems dominated by mature, established trees play a

critical role in global carbon (C) cycling (Harris et al., 2021) and exhibit cellular- to ecosystem-

level responses to drought and elevated atmospheric CO_2 (Blaschke et al., 2001; Cech et al., 2003;

Jiang et al., 2020; Lee et al., 2021; Arain et al., 2022; Martínez-Sancho et al., 2022). As such, there

is a need to better understand the mechanisms by which these environmental stressors impact litter

quality and decomposition in mature forest ecosystems.

Dead organic matter, such as leaf litter, constitutes a significant portion of below-ground

C (Lonsdale, 1988). Through decomposition, leaf litter is converted and released as smaller

compounds, consumed by soil organisms, and taken up by plants largely in mineral forms (Krishna

and Mohan, 2017). Both C are released into the atmosphere as by-products of decomposition, but

also incorporated into various soil C pools (Lavallee et al., 2020), making litter decomposition a

critical component of ecosystem C storage. Globally, climatic factors such as average yearly

temperature and annual precipitation, along with C:N ratio of litter (Zhang et al., 2008), are

strongly correlated with decomposition rate. Decomposer communities, consisting of soil animals

(e.g., mites and collembolans), bacteria, and fungi, drive decomposition through fragmentation of

litter by fauna (Seastedt, 1984; Coleman et al., 2018) and microbial secretion of enzymes, such as

lignin peroxidase (Janusz et al., 2017) that further breakdown detritus. As a result, litter

decomposition is regulated through an interaction between biota, litter quality, and environmental

factors.

Drought and elevated atmospheric CO_2 can directly (e.g., through increased susceptibility

to desiccation) or indirectly (e.g., through changes to litter quality) affect microbial and animal

communities. For example, fungi are generally more resistant to drought than bacteria (Yuste et

al., 2011; de Vries et al., 2012, 2018; Manzoni et al., 2012, but see Santonja et al., (2017)) as

drought can also be an environmental filter for soil and litter animals as moisture content strongly

shapes their community structure (Kardol et al., 2011; Lindo et al., 2012; Barreto et al., 2023), and

alter interaction strength between trophic levels (Aupic-Samain, et al., 2021a) that may indirectly

affect litter decomposition (Lang et al., 2014). Conversely, the effect of elevated atmospheric CO_2

on litter decomposition and associated decomposer communities are more subtle and highly

variable (Blankinship et al., 2011), and potentially, weaker than drought. This effect may be due

to belowground systems containing greater CO_2 concentrations than the atmosphere (Enoch and

Dasberg, 1971). Generally, microbial and animal communities respond to changing litter quality

(Ilieva-Makulec et al., 2006; Gergócs and Hufnagel, 2016; Liu et al., 2016), often stemming from

fungi favouring recalcitrant litter (relative to bacteria) and the subsequent available feeding

resources for animals. Higher C:N of litter under elevated atmospheric CO_2 (Vestergård et al.,

2015; Holmstrup et al., 2017) can subsequently shift associated animal and microbial communities

(Cha et al., 2017; Holmstrup et al., 2017).

Changes to microbial and animal community structure and biomass under environmental

stress can strongly affect litter decomposition. For example, decreasing litter moisture content can

shift microbial composition, resulting in lower mass loss under warming and reduced rainfall

(Prieto et al., 2019). Microbes and animals show a consistent, but highly variable ability to

accelerate decomposition of 'home' litter relative to 'away' litter (i.e. the home field advantage;

HFA) (Ayres et al., 2009; Veen et al., 2015). This variation is partly driven by litter quality, where

recalcitrant litter is subject to stronger HFA than labile litter (Wallenstein et al., 2013). Potentially,

drought or elevated atmospheric CO_2 (along with other environmental stressors) may alter HFA,

thereby affecting litter mass moss. For example, Wang et al. (2020) found generally weaker HFA

effects under years of environmental stress (i.e., drier and colder climate) stemming from changes

to decomposer communities. Both drought and elevated atmospheric CO_2 have potential to affect

litter decomposition through changing interactions between litter quality and stoichiometry and

decomposers, and ultimately C cycling within ecosystems. But it remains unknown how long-

term effects of drought and elevated atmospheric CO_2 impact interactions among litter quality and

stoichiometry and the biotic drivers of decomposition *in situ*, especially in mature forests.

Resolving this uncertainty requires large-scale experimental infrastructure in combination with

multi-variate analysis of litter properties and other factors.

We investigated how drought and elevated atmospheric CO_2 independently affect litter

decomposition in mature forests by conducting a reciprocal transplant across two, large-scale

manipulative field experiments: the University of Birmingham Institute of Forest Research Free-

Air CO_2 Enrichment (BIFoR FACE) experiment and the Oak Observatory at the Observatoire de

Haute-Provence (O_3HP) drought experiment. We tested the hypotheses that elevated atmospheric

CO_2 and drought (generated *via* a reduction in seasonal rainfall) would alter litter decomposition

by acting *via* both litter quality and stoichiometry and the decomposer community. Although

experiments are established in contrasting climates (cooler and wetter vs. warmer and drier), both

drought and elevated CO_2 can shift leaf and litter properties in a similar manner *via* changes to

carbon and nitrogen concentrations, and also impact the decomposer community. By using the

same experimental methodology, we tested how significant correlates of decomposition (i.e., litter

quality / stoichiometry and the decomposer community) are shaped by these key global change

factors.

Methods (Word count: 2,525)

Overview

We collected control and experimental treated litter from the respective field experiments

and transplanted the two litter types into both plots to tease apart and isolate how litter quality and

stoichiometry and decomposers influence rates of litter decomposition under of long-term elevated

CO_2 concentrations and drought. We harvested litterbags across four time points and measured

litter decomposition (as mass loss), litter chemistry and metabolomics, and microbial

(phospholipid fatty acid analysis; PLFA) and animal community structure (the abundance of mites,

collembolans and other larger macrofauna). We then constructed structural equation models

(SEMs) to test for significant pathways derived from well-documented relationships between

abiotic (litter moisture and quality) and biotic (decomposer community) parameters with litter

decomposition to determine how these relationships are directly and indirectly affected by drought

and elevated atmospheric CO_2.

Experimental sites

We conducted a reciprocal transplant study at BIFoR FACE (52.801°N, 2.301°W), United

Kingdom and O_3HP (43.750°N, 5.299°E) in southern France (Supplementary Figure S1). The

BIFoR FACE facility is within a *Quercus robur* L. forest that was planted around 1850 (Hart et

al., 2020; MacKenzie et al., 2021) containing coppice *Corylus avellana* L. and abundant self-set

Acer pseudoplatanus L. understory. The CO_2 treatment began in April 2017 and comprises three

non-infrastructure arrays (that were not used in this experiment) along with six infrastructure

arrays (or plots) (elevated CO_2, n = 3, ambient CO_2, n = 3). Each array is approximately 30 m in

diameter and comprises 16 peripheral towers and one central tower that extend ~ 1m above the

canopy. In the elevated CO_2 plots, an additional 150 ppm of CO_2 above ambient (so, ~36 %

increase over ambient) is emitted into the forest canopy, while ambient CO_2 arrays mimic the

treatment arrays but only receive ambient air. The towers release CO_2 from ~April 1 to October

31 each year, coinciding with oak bud burst and last leaf fall (Supplementary Figure S2).

The O_3HP drought experiment started in 2012 on *Quercus pubescens* Wild forest,

containing *Acer monspessulanum* L. (75% vs. 25% coverage, respectively) and *Cotinus coggygria*

Scop understory. A rain exclusion device, measuring 15 m x 20 m excludes ~30% of precipitation,

relative to the nearby control plot. In our study, the rain exclusion device was operational between

the months of May to September and the total precipitation was 40% lower in rain exclusion plots

over the course of the experiment (total precipitation, control: 1652 mm, drought: 997 mm;

Supplementary Information Figure S3) (see also Gauquelin et al., 2011, Santonja et al., 2017 and

Aupic-Samain, et al., 2021b).

Litterbag preparation and sampling

Quercus litter was sampled from spatially separated areas of each site in BIFoR FACE and

O₃HP in October 2019 and shipped in coolers to The University of Manchester, air dried and used

for analysis of %C, %N, C:N and metabolomic signatures (Quer et al., 2022; see *Metabolomic*

signature of initial litter in Supplementary Information for more details on methods). Litterbags

comprised 1.8 mm mesh with two separate compartments (Supplementary Information Image S1),

each receiving 3 g of dried litter. One compartment was used for chemical analysis of litter and

assessment of microbial community composition, decomposition, and moisture content. The other

compartment was used solely to characterize soil fauna communities. A total of 480 total litterbags

were used (2 sites (BIFoR FACE and O₃HP) x 2 treatments (ambient and drought/elevated CO_2 x

2 litter origins (control and experimental, i.e., from either drought or elevated CO_2 plots) x 3 plots

x 4-time points x 5 replicates = 480). The experiment began January 2020 and litterbags were

harvested across four time points at both experimental sites. At BIFoR FACE, litterbags were

harvested September 2020 (8 months; Harvest one; H1), January 2021 (12 months; Harvest two;

H2), June 2021 (17 months; Harvest three, H3) and December 2021 (23 months; Harvest four, H4),

corresponding to seasons 4 and 5 of the elevated CO_2 treatment. Four litter bags could not be

retrieved from BIFoR FACE at H4 due to a tree limb falling and disturbing litter bags. Also, one

litterbag was removed from H2 because the final dry weight of the litter was not recorded. At

O₃HP, litterbags were harvested October 2020 (10 months; H1), January 2021 (13 months; H2),

September 2021 (21 months; H3) and October 2022 (34 months; H4). The total time for

decomposition varied between our two experiments due to differences in the carbon and nitrogen

concentrations (and subsequent C:N ratio) within our initial litter (Supplementary Information

Table S1). Here, the C:N ratio in our elevated CO_2 experiment was far lower than the drought

experiment. We initially aimed for ~24 months of decomposition, but to ensure litter in the drought

experiment had at least loss half their initial values by the end of the drought experiment, we

continued incubation for an additional year. For the structural equation models (for more details,

see later sections) we elected to use data from H3 where mass loss was on average 40-55% from

the drought experiment and 60-70% for the elevated CO_2 experiment (Figure 1), ensuring that

decomposition was on-going within our study and had not yet plateaued.

At each sampling time, litterbags were shipped to The University of Manchester for

processing. From the 'chemical analysis compartment', remaining litter was removed from the bag

and leaves roughly cleaned by hand from soil residuals. The fresh weight (g) of the litter was taken

before litter was frozen at -20 °C overnight and 48 hours of freeze-drying for determination of dry

weight. For soil fauna litter, litter was removed from the compartment and placed on Tullgren

funnels to extract individual animals for seven days into absolute ethanol. After extraction, litter

was placed into ovens at 65 °C for two days before weighing to measure litter dry weight.

Litter decomposition, quality, stoichiometry, and moisture content

Litter decomposition was calculated as litter mass loss using the following equation:

$$Mass\ loss\ (\%) = \frac{Initial\ litter\ (g) - harvested\ litter\ (g)}{Initial\ litter} * 100$$

We also calculated the exponential decomposition rate coefficient (k_t) at each harvest point

using the equation from (Olson, 1963):

$$k_t = \frac{-\ln\left(\frac{X_H}{X_I}\right)}{t}$$

Where X_H is the weight (in g) of harvested litter, X_I is the weight of initial litter (in g), and

t is time in years since the beginning of the experiment. Litter decomposition was measured from

all litterbags across all four time points. We measured the %C, %N and C:N ratio with a Elementar

Vario EL element analyser (Hanau, Germany) on 5 mg of ball-milled litter (Retsch Ball Mill MM

400, Haan, Germany) to gauge litter quality in H3 and H4 time points, at both sites.

In addition, we measured litter stoichiometry on H3 and H4 litter from both sites using

Fourier transform infrared (FTIR) spectroscopy, following Bell et al. (2018). Ball-milled leaves

were plaved onto a Universal Attenuated Total Reflection (UATR) spectrometer and the

absorbance (A cm^{-1}) measured within a 650 to 4000 A cm^{-1} range, at 4 cm^{-1} resolution and 16

scans per sample. We also ran a background correction after every ten samples measured. After

samples were processed, we applied a baseline correction for each spectra using the modpolyfit

method in the R package 'baseline'. To standardize values across our samples, we Z-transformed

each spectra, where values were transformed to a mean of 0 and SD of 1 (Jardine et al., 2017; Bell

et al., 2018). We then calculated the area of the curve for each respective functional group using

a shoelace formula (Wang et al., 2023). The assignment of wave number regions to leaf litter

functional groups was based on (Liu et al., 2016) who used FTIR to examine functional groups

across four deciduous litter origins, including *Quercus robur* (Supplementary Information Table

S2).

Finally, we also calculated leaf litter moisture content for H3 and H4 litter and both sites

using the fresh (or wet) weight of the harvested litter and freeze-dried litter from the chemical

compartment of the litterbag.

Phospholipid fatty acid analysis (PLFA)

We used PLFAs to analyse microbial community composition in H3 leaf litter (at both sites)

using methods adapted from (Buyer and Sasser, 2012). We quantified microbial communities with

~0.4-0.5 g of ball milled, freeze-dried litter. A single litter bag (from the elevated CO_2 experiment)

contained 0.29 g after freeze-drying that PLFAs were measured. Four additional litterbags were

removed from analysis from the elevated CO_2 experiment due to either insufficient mass of litter

needed to measure PLFAs or no PLFAs were detected. Four ml of Bligh-Dyer extractant

containing C19:0 of an internal standard was added to each sample to extract lipids. Once

extracted, we used a gas chromatograph (Agilent 7890A Gas chromatograph, Santa Clara, CA,

USA) to assign and calculate peak height, where PLFA group abundances were expressed as nmol

g dry weight of litter^{-1}. We used 18 fatty acid markers to characterise microbial communities (Liu

et al., 2016; Dhaou et al., 2022; Biryol et al., 2024) with 6 and 4 markers for gram positive (i14:0,

i15:0, a15:0, i16:0, i17:0, a17:0) and gram negative (16:1ω7c, cy17:0, 18:1ω7, and cy19:0)

bacteria, respectively. In addition, we used 14:0, 15:0, 16:0 and 17:0 as general bacteria marks,

and 10Me16:0, 10Me17:0, and 10Me18:0 as markers for actinobacteria. The marker 18:2ω6,9 was

the sole fungal marker in our experiments. With these markers, we analyzed treatment and litter

origin effects on total PLFA concentrations (biomass), F:B ratio (by dividing 18:2ω6,9 abundance

by the summed total of bacteria markers) and PLFA community composition across both

experiments.

Soil fauna community

Soil fauna community composition was quantified on H3 and H4 litter at both sites and

was extracted using Tullgren funnels over seven days into absolute ethanol. Soil fauna was sorted

and counted into higher taxonomic or mixed groups, comprising: Collembola (Hexapoda),

Oribatida (Arachnida: Sarcoptiformes), Astigmata (Arachnida: Sarcoptiformes), Prostigmata

including the mite family Nanorchestidae (hereafter, called Prostigmata, Arachnida:

Trombidiformes, Sarcoptiformes), Mesostigmata (Arachnida: Parasitiformes), and 'other' fauna,

which included Annelida, Isopoda, and Insecta. Soil fauna abundances were standardized by the

dry weight of litter (individuals g dwt^{-1}), to account for differences in sample mass prior to analysis.

We examined the effect of treatment and litter origin on total fauna and fauna group abundance

and composition.

Statistical analysis

All statistical analyses were performed with the program R (version 4.3.0) (R Core Team,

2023) and figures produced with the package 'ggplot2' (Wickham, 2016). To analyze for

differences in initial litter chemistry between our litter types (litter from control and experimental

plots) prior to the experiment beginning, we used one-way Analysis of Variance (ANOVA) tests

and Permutation Multivariate ANOVAs (PERMANOVAs) to examine for differences between

C%, N% and C:N ratio (ANOVA) and leaf metabolomic profile (PERMANOVA, with

Euclidean distance) between our treatments (ambient vs. drought or elevated CO_2). To do this,

we used the aov function in base R and the adonis2 function in the package 'vegan' (Oksanen et

al., 2022), respectively. Here, we partitioned the data by sites (BIFoR FACE and O_3HP) and ran

separate models for both.

We tested the effect of treatment (ambient vs. drought or elevated CO_2) and litter origin

(control vs. experimental) on our parameters, analyzing the sites (BIFoR FACE and O_3HP) and

harvest time points (H1-4) separately (excluding litter decomposition, for more details see

below). For univariate variables, we used linear mixed models where the fixed factors were

treatment, litter origin, and their interaction, with a random factor of experimental plot. We used

linear mixed models for %C, %N, C : N ratio, total PLFA biomass (nmol g dwt^{-1}), fungi :

bacteria ratio, animal total and group abundances (individuals g dwt^{-1}), which includes

Collembola, Oribatida, Astigmata, Mesostigmata, Prostigmata, and 'other' fauna. Note, we did

not analyze Astigmata abundances in the drought experiment as low abundances were present in

samples. For litter decomposition (measured as mass loss) we included all time points together

into a single model, where the fixed factors were treatment, litter origin, harvest, and their

subsequent interactions with a random factor of experimental plot. To run linear mixed models,

we used the lmer function in the package 'lme4' (Bates et al., 2015) where we calculated F-

statistics and p-values with Satterthwaite's method using the anova function in the package

'lmerTest' (Kuznetsova et al., 2017). If model residuals were not normally distributed, we ln or

arscine-transformed the dependent variable and re-ran the model. At times, we found singularity

within our linear mixed models as the variance within our random factor terms was extremely

low (essentially zero) and the model was overfit. When this occurred, we removed the random

factor, and we re-ran the model without the random term.

For multivariate variables, including leaf litter functional groups (FTIR), microbial

community composition, and animal community composition, we ran PERMANOVAs using the

adonis2 function in the package 'vegan' (Oksanen et al., 2022). For leaf litter functional groups,

we used Euclidean distances within our models. But for microbial and animal communities we

used Bray-Curtis dissimilarity as it it better suited for compositional data (like the abundances of

each PLFA peak). Each model was fitted with the same fixed factors, consisting of treatment,

litter origin and their interaction. To visualize leaf litter functional groups and microbial

communities we used a Principal Component Analysis (PCA) ordination plot using the rda

function in the package 'vegan' (Oksanen et al., 2022). For microbial communities, we

Hellinger-transformed datasets using the decostand function in the 'vegan' package prior to the

PCA.

Finally, we used structural equation models (SEMs) to construct a global model to

analyze the direct and indirect pathways of the effect of drought and elevated CO_2 on litter

decomposition on H3 data. We constructed three *a priori* models for the drought and elevated

CO_2 experiment (Supplementary Information Figure S4) to detect significant pathways ($p <$

0.05) between parameters (Garrido et al., 2022). Here, we wanted to test whether abiotic and

biotic factors, abiotic factors only (litter C:N and moisture content) or biotic factors only (animal

and microbial communities) best explain litter mass loss in our experiments. Within our SEMs,

we included treatment and litter origin as binary variables where experimental litter (litter origin)

and elevated CO_2 and drought (treatment) where coded as 1 (with the respective controls coded

as 0) to reduce the number of pathways in our model. As well, we Z-transformed litter mass loss,

C:N ratio, litter moisture content, along with animal and microbial abundance, but not PCA axes,

to standardize parameters. To include parameters on microbial and animal communities in our

SEM that best described litter mass loss, we constructed four candidate models (linear mixed

models with experimental plot as random factor) that contained either total animal abundance or

animal community PC1 (first axis from PCA), along with total microbial abundance or microbial

PC1. We compared models using AIC (Akaike Information Criterion) with maximum likelihood

and chose the candidate model with the lowest AIC value to be included in the SEM. To

construct SEM models, we used the psem function in the 'piecewiseSEM' package (Lefcheck,

2016) where individual models within the global SEM were fitted with linear mixed models

(with plot as random factor) with maximum likelihood, using the lme function from the 'nlme'

package (Pinheiro and Bates, 2000; Pinheiro et al., 2023). To evaluate our SEM models, we used

AIC, $\chi 2$ goodness of fit test and Fisher's C, where p-values > 0.05 represents acceptable data fit

for our model.

Results (Word count: 1,357)

Initial litter chemistry

We found strong differences between the C:N ratio of the initial control and droughted

litter ($F_{1,8} = 13.92$, $p = 0.006$), with the C:N ratio of control litter exceeding that of the droughted

litter, but not between litter from ambient and elevated CO_2 treatments ($F_{1,8} = 0.001$, $p = 0.977$)

(Supplementary Information Table S1). Furthermore, strong differences were detected between

the metabolomic profiles between the initial control and droughted litter ($F_{1,8} = 3.91$, $p = 0.009$)

but not between litter from ambient and elevated CO_2 treatments ($F_{1,8} = 1.04$, $p = 0.336$)

(Supplementary Information Figure S5). Control litter from the drought experiment was mainly

composed of ellagitannins (from $C_{34}H_{24}O_{22}$ to $C_{56}H_{40}O_{31}$), catechin ($C_{15}H_{14}O_6$), and ellagic

hexoside ($C_{20}H_{16}O_{13}$) that were overexpressed within litter. Other metabolites such as quercetin

hexosides (or rhamnoside), kaempferol hexosides, and isorhamnetin hexosides at similar quantities

between control and drought treatment.

Elevated CO$_2$ experiment

During the first eighteen months of the decomposition experiment, litter from elevated CO_2

plots (i.e., experimental litter) decomposed slower than litter from control plots (litter origin:

$F_{1,215.09} = 5.932$, $p = 0.016$) through H1-3 (time: $F_{3,215.46} = 155.97$, $p < 0.001$) until H4. Whereafter,

mass loss varied little between the two treatments and litter origins (Figure 1, Supplementary

Information Table S3 contains exponential decomposition rate coefficient, k_t, values). Similarly,

the C:N ($F_{1,56} = 6.03$, $p = 0.02$) of litter was higher in experimental litter (i.e., initial litter that was

harvested from elevated CO_2 plots) at H3, but not at H4 (Supplementary Information Table S4 and

S5), suggesting that litter characteristics converged over time. This finding is also supported by

FTIR spectroscopy analysis, which showed significant, but limited differences in litter functional

groups at H3 between litter origins ($F_{1,56} = 4.33$, $p = 0.03$). Litter functional group composition

between control and experimental litter was no longer significant at H4 ($F_{1,52} = 0.26$, $p = 0.81$)

(Figure 2A-B). Neither treatment or litter origin affected litter moisture content at either H3 or H4

(Supplementary Information Table S4 and S5).

Treatment and litter origin did not influence PLFA abundance, the fungi : bacteria (F:B)

ratio, or microbial community composition at H3 (Supplementary Information Table S4,

Supplementary Information Figure S4). Likewise, no individual or total animal group abundances,

along with entire community structure, were affected by treatment or litter origin in H3. In H4,

however, the abundance of Mesostigmata ($F_{1,52} = 5.04$, $p = 0.03$) and Prostigmata ($F_{1,52} = 7.30$, p

$= 0.009$) was significantly greater in control litter relative to experimental litter, but animal

abundance composition did not shift with litter origin ($F_{1,48} = 1.24$, $p = 0.28$) (Supplementary

Information Table S5).

The SEM revealed several significant pathways between parameters in response to elevated

CO_2 (Figure 5) with strong goodness-of-fit where model 1 (containing both abiotic and biotic

factors) best explained mass loss. Litter moisture content was negatively correlated with microbial

community structure. On PC1, microbial composition was driven by fungal abundance (PLFA

marker C18:2ω6,9) (Supplementary Information Figure S6); the negative association between

moisture and microbial community composition suggests that drier litter supported greater fungal

abundance. In addition, microbial community structure was correlated with mass loss, meaning

lower litter moisture content indirectly shaped mass loss through changes to microbial community

composition. Litter origin indirectly affected mass loss *via* litter C:N, as experimental litter (litter

derived from elevated CO_2 plots) led to higher C:N in litter, which corresponded with slower

decomposition. In addition, higher C:N ratio was negatively associated with animal abundance

within litter bags, with greater animal abundance being associated with faster decomposition.

Treatment had no significant direct or indirect effect in the SEM, further supporting that elevated

CO_2 did not directly affect litter decomposition or associated microbial or animal communities in

our study.

Drought experiment

By contrast to elevated CO_2, there were stronger effects of both treatment and litter origin

on measured parameters in response to drought. First, litter decomposition was slower in

droughted plots (relative to ambient moisture plots) (treatment: $F_{1,4} = 147.91$, $p < 0.001$) as

remaining litter mass decreased over time ($F_{1,220} = 494.76$, $p < 0.001$) and at varying rates

(treatment x time: $F_{1,220} = 4.56$, $p = 0.004$) (Figure 1; Supplementary Information Table S3). Litter

had lower C:N ratios in ambient plots (H3: $F_{1,4} = 21.92$, $p = 0.009$; H4: $F_{1,56} = 37.85$, $p < 0.001$)

compared to drought plots. In addition, experimental litter (i.e., litter derived from drought plots)

had lower C:N ratio, relative to the control litter (H3: $F_{1,52} = 32.53$, $p < 0.001$; H4: $F_{1,56} = 9.54$, p

$= 0.003$) suggesting both treatment and litter origin affected litter quality during the experiment

(C:N but also %C and %N; Figure 3; Supplementary Information Table S6 and S7).

Similarly, we observed a strong effect of treatment on leaf litter functional compounds

(Figure 2C-D). FTIR spectroscopy revealed that drought (H3: $F_{1,56} = 6.73$, $p = 0.006$; H4: $F_{1,56} =$

15.32, $p = 0.002$) shifted litter C-biochemistry; hydroxyl groups (B1) and amides (I and II, B4 and

B6) were grouped together along PC1 in drought litter at both H3 and H4 time points. Conversely,

carboxylates (B5), sulfones and esters (B10), along with aromatics (B15) were correlated with

ambient litter and both H3 and H4. The functional groups B11 (phenols) and B14 (carbohydrates)

were correlated in both H3 and H4 litter.

We found divergent effects of the drought treatment with litter moisture content in the

drought experiment. At H3, litter moisture content significantly decreased (by around 35%) in

drought plots ($F_{1,4}$ = 85.96, p < 0.001; Supplementary Information Table S6). However, at H4,

when litter moisture content in ambient conditions was already low, there was no significant

difference between the ambient moisture or drought plots ($F_{1,56}$ = 1.086, p = 0.30; Supplementary

Information Table S7).

After 21 months of decomposition (H3), experimental litter (i.e., litter from the droughted

plots) had greater concentrations of PLFAs ($F_{1,52}$ = 9.06, p < 0.004, Figure 4A), but the drought

treatment reduced total PLFA concentrations ($F_{1,4}$ = 15.69, p = 0.02) in litter regardless of its origin.

The F:B ratio was also greater in the experimental litter ($F_{1,52}$ = 8.43, p = 0.005, Figure 4B). Both

litter origin ($F_{1,56}$ = 6.64, p < 0.004) and treatment ($F_{1,56}$ = 23.59, p < 0.001; Figure 4C) shifted

PLFA composition. Microbial composition in response to treatments was largely explained by PC1

whereas microbial composition in response to litter origin was explained more by PC2. These axes

reflected distinct PLFAs, with the fungal-related PLFA C18:2ω6,9 driving separation in both PC1

and PC2 (Fig. 4C). In addition, animal communities responded more to the drought treatment than

litter origin (Supplementary Information Table S6 and S7). At both H3 and H4, animal

communities shifted under drought (H3: $F_{1,56}$ = 8.07, p < 0.001; H4: $F_{1,56}$ = 6.70, p < 0.001):

mesostigmatid (H3: $F_{1,56}$ = 31.18, p < 0.001) and prostigmatid mite (H4: $F_{1,56}$ = 7.65, p = 0.007)

abundances significantly decreased, and there was a trend for drought to reduce total animal

abundances at H4 ($F_{1,4}$ = 6.04, p = 0.07). Notably, there was a significant interaction of treatment

and litter origin on 'other' fauna abundances at H3 (treatment x litter origin: $F_{1,52}$ = 6.51, p = 0.01)

where abundances were greatest in the drought treatment with control litter.

The SEM for the drought experiment yielded several significant pathways and had

acceptable goodness-of-fit where model 2 (abiotic factors only) best explained mass loss (Figure

5). First, litter origin affected litter mass loss directly, but also indirectly through litter C:N, with

higher C:N ratio of litter corresponding to slower mass loss (which also matched the result from

the SEM describing pathways under elevated CO_2). Greater litter moisture also led to lower C:N

ratio but moisture content did not directly affect litter mass loss. Similar to mass loss, microbial

community structure was also negatively associated with litter C:N. Here, PC1 represented a

gradient from fungal dominated to bacterial dominated communities (Figure 4C), where higher

C:N ratio corresponded with fungal dominated litter communities. The SEM provides robust

evidence that litter mass loss was shaped strongly by litter C:N rather than a shift in the

composition of microbial and animal communities in the drought experiment.

Discussion (Word count: 1,929)

Our results reveal that long-term *in situ* manipulations of drought and increased

atmospheric CO_2 concentrations to mature oak forests have contrasting impacts on rates of litter

decomposition. We found that after experimental summer drought, representing an average 40%

reduction in rainfall, litter had greater C:N ratio and reduced mass loss compared to controls. The

SEM revealed that the main drivers of litter mass loss were the origin of litter, i.e., whether it was

derived from plants subjected to drought or not. Litter origin had clear effects on both the C:N

ratio of the litter and on the composition of secondary metabolites, and both of the these factors

likely led to litter origin having both direct and indirect effects on mass loss. The reduction of

seasonal rainfall influenced mass loss mainly by reducing the water content of the litter which also

affected the C:N of litter. By contrast, elevated CO_2 mainly affected the C:N of the litter, which

was associated with a shift in the abundance of soil fauna, while the environment itself was not a

significant factor driving decomposition.

Our findings generally support those from previous studies conducted with other plant litter

species and often more artificial experimental set-ups (Cotrufo and Ineson, 1996; Norby et al.,

2001; Santonja et al., 2015; Vestergård et al., 2015; Holmstrup et al., 2017; Prieto et al., 2019;

Quer et al., 2022) that drought and elevated CO_2 concentrations will impact litter decomposition.

In general, both drought and eCO_2 treatments had substantial impacts on the 'quality' of litter

entering and residing in the decomposer subsystem. We found that the C:N ratio, a general

indicator of litter quality, is critical at determining decomposition rate. Higher C:N ratios generally

corresponded with slower mass loss (Zhang et al., 2008), and our results highlight the robustness

of this relationship in the mature forests. Of note is the observation that our initial litter from

control plots (at O$_3$HP) had a higher C:N than the litter from the droughted plot, which carried

over into subsequent harvests, and which stemmed from the litter from droughted plots having

relatively high nitrogen concentration, at least compared to past work at this site (Aupic-Samain

et al., 2021b). Nitrogen concentration within leaves can be affected by both nutrient availability

in soil, climatic factors like precipitation, and their interaction (Cheaib et al., 2025), but we do not

know why nitrogen concentrations differed in our experiment compared to past work. Our SEM

revealed decomposition in response to drought was driven indirectly via litter C:N and also through

other, albeit unknown, direct pathways.

The SEM from BIFoR FACE showed that both animal abundances and microbial

community structure were correlated with mass loss, along with litter C:N, providing further

evidence that animals and microbes are important contributors to litter decomposition

(Hättenschwiler et al., 2005; Rouifed et al., 2010; Handa et al., 2014; Tan et al., 2021; Njoroge et

al., 2022). By contrast, litter quality (litter C:N, directly) and moisture content (indirectly) best

explained mass loss in the drought experiment. Notable climate differences exist between the two

sites; BIFoR FACE is in a cool, wet temperate climate (MacKenzie et al., 2021), while O$_3$HP is in

a seasonally hot-dry Mediterranean climate (Gauquelin et al., 2011) (Figures S2 and S3). Joly et

al. (2023) re-affirmed the role of climate in regulating decomposition, and strong climate x

decomposer community interactions underlie mass loss (Glassman et al., 2018; Yin et al., 2019).

For instance,Wall et al. (2008) found that soil fauna enhanced decomposition in temperate and

wetter climates, but had limited effects on decomposition when constrained by relatively cooler

temperatures and/or lower precipitation inputs. When not as constrained by climate, like at BIFoR

FACE, animal and microbial communities were strongly associated with decomposition rates.

Furthermore, we found that the decomposer community was strongly correlated to mass loss at

later stages of decomposition when around half of the original litter mass remained. Understanding

the interactions between biota and litter quality on decomposition is crucial to predict how

environmental stressors will impact decomposition and biogeochemical cycling. Our findings

demonstrate that the influence of drought and elevated CO$_2$ on decomposition was strongly

dictated by indirect effects acting via litter chemistry (notably C:N), while direct influences on

decomposer abundance and community composition were apparent only when not constrained by

temperature or moisture.

Past studies have demonstrated drought can lead to fungal-dominated communities due to

their ability to persist under water stress (Yuste et al., 2011; Biryol et al., 2024) as the SEM model

showed fundamental linkages between moisture content, litter C:N, and microbial communities.

Here changes to litter moisture content was critical as it indirectly affected microbes *via* litter C:N.

Specifically, drought impacted carbon and nitrogen concentrations in litter by limiting moisture,

inducing water stress conditions that shifted the decomposer community and was indirectly

correlated with fungal-rich communities. By contrast, the SEM from the elevated CO_2 experiment

found a significant relationship between litter C:N and animal abundances, with a higher C:N ratio

corresponding with fewer animals. This result was found alongside the preferential feeding of

defoliator insects on elevated CO_2 leaves in 2019 prior to litter collection (Norby et al., 2024),

further highlighting how elevated CO_2 will affect multiple dimensions of the ecosystem.

Holmstrup et al. (2017) found two years of elevated CO_2 to a heathland changed litter C:N that

affected animal diversity, while microbes (denoted by fungi : bacteria) were unaffected. In our

study, microbial and animal communities differed significantly between the two experiments;

foodwebs in the drought experiment were dominated by fungi with few animals, while in the

elevated CO_2 experiment foodwebs were dominated by bacteria with many animals. The

contrasting relationships among animal and microbial communities with litter quality between our

experiments may partly be attributed to litter C:N being far higher in the drought experiment

compared to the elevated CO_2 experiment. Although the mechanisms underpinning these

contrasting relationships is unknown, we posit that climate and environmental stressors, or

differences in litter quality, may modify linkages between litter and their decomposers.

Home-field advantage (HFA) stipulates that decomposer communities are more adept to

decomposing litter from their own environment ('home') relative to litter from other locations

('away') (Ayres et al., 2009; Veen et al., 2015). We did not find strong evidence for home-field

effects, as decomposition of home and away litter were similar as no significant interaction

between treatment and litter type was observed. Higher litter quality dissimilarity leads to stronger

HFA effects (Veen et al., 2015). We found weak or no differences in litter C:N between control

and experimental litter in the elevated CO_2 experiment (both in the initial litter and across two

harvests), but strong treatment and litter origin effects in the drought experiment. Notably, the

metabolic profiles of initial leaf litter differed strongly from control and droughted litter, while

litter leaf functional groups (measured by Fourier transform infrared spectroscopy) also differed

between litter incubated in droughted plots and removed at H3 and H4. However, our data suggests

that differences in litter properties between the experimental treatments were potentially not large

enough to observe a HFA effect which has often been found in situations where litter properties

vary considerably, such as between coniferous and deciduous litter (Chomel et al., 2015), or more

generally, recalcitrant versus labile litter (Wallenstein et al., 2013). Individual and group microbial

and animal abundances shifted strongly under drought, but HFA may have been negligible as litter

quality, rather than direct feeding and processing of litter by decomposer communities, was the

main correlate of decomposition rates within the O_3HP experiment. Potentially, if environmental

stressors or climate weaken linkages between decomposers and litter mass loss, HFA may lessen

within a given system, thereby affecting the efficiency of biotic communities on litter

decomposition and nutrient release.

We found few significant effects of elevated CO_2 experimental litter on microbial

communities and most animal groups, but elevated CO_2 experimental litter was associated with

fewer mesostigmatid and prostigmatid mites at H4. Most mesostigmatid mites, and some

prostigmatid mite species, are predators, feeding on arthropods and nematodes (Lindquist et al.,

2009; Walter et al., 2009). Lower abundances within experimental litter but also the drought

treatment suggest that changes to prey composition (but likely not group abundances in our

experiments) hindered predators in our litterbags. Changing mesofauna and macrofauna predator

composition can indirectly influence litter decomposition (Lang et al., 2014; Koltz et al., 2018)

and may have impacted decomposition in our study as well. For example, predators can reduce

competion amongst lower trophic levels resulting in higher litter decomposition (Melguizo-Ruiz

et al., 2020), highlighting the complexity of these trophic interactions on litter decomposition.

Although not statistically significant, oribatid mite abundance was, on average, less under drought

than control conditions, which aligns with past studies (Aupic-Samainet al., 2021b; Santonja et al.,

2017). But overall, experimental manipulation had greater effects on predator abundances

compared to other animal groups.

Evaluating environmental stressors across time is critical to understand their impact on C

cycling and the release of other mineral nutrients during decomposition because the relative effects

of climate and litter traits on decomposition changes temporally. Initial decomposition (e.g., 1-2

years) is correlated relatively more with litter traits, but climate regulates mass loss at later stages

(Currie et al., 2010; Canessa et al., 2021). Similarly, the biotic contributions to decomposition also

diminishes with time (García-Palacios et al., 2016). Because we conducted separate experiments

(within contrasting climates and varying litter quality), we cannot fully tease apart the temporal

and climatic effects that drove experimental results. However, by conceptually merging the SEM

models, our results indicate litter mass loss was slower in the warmer Mediterranean climate where

litter C:N was higher, as litter quality consistently drives mass loss rates (within the first 2-3 years

at least). While in the cooler, wetter temperate climate of BIFoR FACE, the decomposer

community was correlated with mass loss when half the litter remained. Future research at both

these on-going experimental sites (of BIFoR FACE and O_3HP) and elsewhere, should incorporate

multiple stressors (Rillig et al., 2019) and recognize that climate x temporal interactions may be

critical to disentangling the contributions of litter quality and and local enviromental conditions

on biota and subsequently litter decomposition.

Conclusions

Our findings reveal how elevated CO_2 and drought can impact decomposition through their

effects on litter properties, decomposer communities, and the environment. Drought and elevated

CO_2 independently impacted litter mass loss by shifting litter C:N, which was a significant

determinant of decomposition in both SEMs. Moreover, litter quality influenced animal

abundances in the elevated CO_2 experiment and microbial communities in the drought experiment.

The decomposer community was only correlated with mass loss in the elevated CO_2 experiment,

which might be because the climate being cooler and wetter than in southern France where the

drought experiment was located. Our findings help interpret the broader impacts of drought and

elevated CO_2 on ecosystem dynamics. Future climate scenarios of increased atmospheric CO_2

concentractions will alter litter decomposition primarily with alterations to litter quality,

specifically, through higher C:N ratio. Reduced seasonal rainfall will both strongly shape litter

properties and associated decomposer communities. Non-additive responses between treatment

and litter type were largely absent within the drought experiment, as prior exposure of leaves to

reduced rainfall did not distinctly alter litter decomposition. Here, drought will have a continuous,

pervasive effect on litter decomposition, specifically, changing litter quality, decomposer

communities, and dampening litter mass loss rates and nutrient release. Overall, these findings

provides us with a more complete understanding of how elevated CO_2 and drought will impact C

inputs to forests, but highlights a need to disentangle the individual and interactive effects of these

perturbations on litter inputs, properties and decomposition.

Acknowledgments

This study was carried out as part of UK Natural Environment Research Council (NERC) grant

NE/S002189/1. ARMK gratefully acknowledges additional support from NERC through grant

NE/S015833/1 (QUINTUS). TC acknowledges support by Research Ireland (20/FFP-P/8584).

The BIFoR-FACE facility acknowledges generous underpinning support from the JABBS

Foundation, Norbury Park Estate, The John Horseman Trust, Ecological Continuity Trust, and

the University of Birmingham. We thank Dr Kris Hart and the operations team at BIFoR FACE,

Dr. Ully Kritzler, Deborah Ashworth, Dr. Thomas Bishop and Dr. Benjamin Bell for technical

support. We thank Dr. Stéphane Greff (Aix-Marseille Université) for analyzing leaf litter and

Jean-Phillipe Orts (Aix-Marseille Université) for collecting litter from the O_3HP experimental

site.

Conflict of Interest Statements

The authors have no conflict of interest to declare.

Data availability statement

Data will be submitted to the Natural Environment Research Council (NERC) data repository

(Environmental Information Data Center) upon acceptance of the manuscript.

Orcid accounts

Matthew L. Meehan: https://orcid.org/0000-0001-5427-8241

Mathilde Chomel: https://orcid.org/0000-0001-5110-2355

Katy J. Faulkner: https://orcid.org/0000-0002-9204-6502

Tancredi Caruso: https://orcid.org/0000-0002-3607-9609

A. Robert MacKenzie: https://orcid.org/0000-0002-8227-742X

Virginie Baldy: https://orcid.org/0000-0002-4048-5132

Richard D. Bardgett: https://orcid.org/0000-0002-5131-0127

David Johnson: https://orcid.org/0000-0003-2299-2525

Author Contributions

622   MC, ARB, VB, RDB, and DJ conceptualized the study and attained funding. MLM, MC, ZV and

623   KJF collected data. MLM and MC curated data, and MLM performed the data analysis with

624   support from TC. MLM wrote the first version of the manuscript and MC, KJF, TC, ARB, VB,

625   RDB, and DJ provided feedback and contributed to finalizing the manuscript.

626   .

References

Ainsworth, E.A., Long, S.P., 2005. What have we learned from 15 years of free-air CO_2 enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO_2. New Phytologist 165, 351–372. https://doi.org/10.1111/j.1469-8137.2004.01224.x

Arain, M.A., Xu, B., Brodeur, J.J., Khomik, M., Peichl, M., Beamesderfer, E., Restrepo-Couple, N., Thorne, R., 2022. Heat and drought impact on carbon exchange in an age-sequence of temperate pine forests. Ecological Processes 11, 7. https://doi.org/10.1186/s13717-021-00349-7

Aupic-Samain, A., Baldy, V., Delcourt, N., Krogh, P.H., Gauquelin, T., Fernandez, C., Santonja, M., 2021a. Water availability rather than temperature control soil fauna community structure and prey–predator interactions. Functional Ecology 35, 1550–1559. https://doi.org/10.1111/1365-2435.13745

Aupic-Samain, A., Santonja, M., Chomel, M., Pereira, S., Quer, E., Lecareux, C., Limousin, J.M., Ourcival, J.M., Simioni, G., Gauquelin, T., Fernandez, C., Baldy, V., 2021b. Soil biota response to experimental rainfall reduction depends on the dominant tree species in mature northern Mediterranean forests. Soil Biology and Biochemistry 154, 108122. https://doi.org/10.1016/j.soilbio.2020.108122

Ayres, E., Steltzer, H., Simmons, B.L., Simpson, R.T., Steinweg, J.M., Wallenstein, M.D., Mellor, N., Parton, W.J., Moore, J.C., Wall, D.H., 2009. Home-field advantage accelerates leaf litter decomposition in forests. Soil Biology and Biochemistry 41, 606–610. https://doi.org/10.1016/j.soilbio.2008.12.022

Barreto, C., Conceição, P.H.S., de Lima, E.C.A., Stievano, L.C., Zeppelini, D., Kolka, R.K., Hanson, P.J., Lindo, Z., 2023. Large-scale experimental warming reduces soil faunal biodiversity through peatland drying. Frontiers in Environmental Science 11, 1153683. https://doi.org/10.3389/fenvs.2023.1153683

Bates, D., Maechler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Bell, B.A., Fletcher, W.J., Ryan, P., Seddon, A.W.R., Wogelius, R.A., Ilmen, R., 2018. UV-B-absorbing compounds in modern *Cedrus atlantica* pollen: The potential for a summer UV-

B proxy for Northwest Africa. Holocene 28, 1382–1394.
https://doi.org/10.1177/0959683618777072

Blankinship, J.C., Niklaus, P.A., Hungate, B.A., 2011. A meta-analysis of responses of soil biota
to global change. Oecologia 165, 553–565. https://doi.org/10.1007/s00442-011-1909-0

Biryol, C., Aupic-Samain, A., Lecareux, C., Gauquelin, T., Baldy, V., Santonja, M., 2024.
Interactive effects of soil moisture, air temperature and litter nutrient diversity on soil
microbial communities and *Folsomia candida* population. Oikos 2024, e10345.
https://doi.org/10.1111/oik.10345

Blaschke, L., Schulte, M., Raschi, A., Slee, N., Rennenberg, H., Polle, A., 2001. Photosynthesis,
soluble and structural carbon compounds in two Mediterranean oak species (*Quercus*
pubescens and *Q. ilex*) after lifetime growth at naturally elevated CO_2 concentrations. Plant
Biology 3, 288–298. https://doi.org/10.1055/s-2001-15203

Buyer, J.S., Sasser, M., 2012. High throughput phospholipid fatty acid analysis of soils. Applied
Soil Ecology 61, 127–130. http://dx.doi.org/10.1016/j.apsoil.2012.06.005

Canessa, R., van den Brink, L., Saldaña, A., Rios, R.S., Hättenschwiler, S., Mueller, C.W., Prater,
I., Tielbörger, K., Bader, M.Y., 2021. Relative effects of climate and litter traits on
decomposition change with time, climate and trait variability. Journal of Ecology 109, 447–
458. https://doi.org/10.1111/1365-2745.13516

Cech, P.G., Pepin, S., Körner, C., 2003. Elevated CO_2 reduces sap flux in mature deciduous forest
trees. Oecologia 137, 258–268. https://doi.org/10.1007/s00442-003-1348-7

Cha, S., Chae, H.-M., Lee, S.-H., Shim, J.-K., 2017. Effect of elevated atmospheric CO_2
concentration on growth and leaf litter decomposition of *Quercus acutissima* and *Fraxinus*
rhynchophylla. PLoS ONE 12, e0171197. https://doi.org/10.1371/journal.pone.0171197

Cheaib, A., Waring, E.F., McNellis, R., Perkowski, E.A., Martina, J.P., Seabloom, E.W., Borer,
E.T., Wilfahrt, P.A., Dong, N., Prentice, I.C., Wright, I.J., Power, S.A., Hersch-Green, E.I.,
Risch, A.C., Caldeira, M.C., Nogueira, C., Chen, Q., Smith, N.G., 2025. Soil nitrogen
supply exerts largest influence on leaf nitrogen in environments with the greatest leaf
nitrogen demand. Ecology Letters 28, e70015. https://doi.org/10.1111/ele.70015

Chen, D., Wang, S., Xiong, B., Cao, B., Deng, X., 2015. Carbon/nitrogen imbalance associated
with drought-induced leaf senescence in *Sorghum bicolor*. PLoS ONE 10, e0137026.
https://doi.org/10.1371/journal.pone.0137026

Chomel, M., Guittonny-Larchevêque, M., DesRochers, A., Baldy, V., 2015. Home field advantage
of litter decomposition in pure and mixed plantations under boreal climate. Ecosystems 18,
1014–1028. https://doi.org/10.1007/s10021-015-9880-y

Chomel, M., Lavallee, J.M., Alvarez-Segura, N., de Castro, F., Rhymes, J.M., Caruso, T., de Vries,
F.T., Baggs, E.M., Emmerson, M.C., Bardgett, R.D., Johnson, D., 2019. Drought decreases
incorporation of recent plant photosynthate into soil food webs regardless of their trophic
complexity. Global Change Biology 25, 3549–3561. https://doi.org/10.1111/gcb.14754

Coleman, D.C., Callaham Jr., M.A., Crossley Jr., D.A., 2018. Fundamentals of soil ecology.
Elsevier Inc., Burlington.

Cotrufo, M.F., Ineson, P., 1996. Elevated CO_2 reduces field decomposition rates of *Betula pendula*
(Roth.) leaf litter. Oecologia 106, 525–530. https://doi.org/10.1007/BF00329711

Currie, W.S., Harmon, M.E., Burke, I.C., Hart, S.C., Parton, W.J., Silver, W., 2010. Cross-biome
transplants of plant litter show decomposition models extend to a broader climatic range
but lose predictability at the decadal time scale. Global Change Biology 16, 1744–1761.
https://doi.org/10.1111/j.1365-2486.2009.02086.x

de Vries, F.T., Griffiths, R.I., Bailey, M., Craig, H., Girlanda, M., Gweon, H.S., Hallin, S.,
Kaisermann, A., Keith, A.M., Kretzschmar, M., Lemanceau, P., Lumini, E., Mason, K.E.,
Oliver, A., Ostle, N., Prosser, J.I., Thion, C., Thomson, B., Bardgett, R.D., 2018. Soil
bacterial networks are less stable under drought than fungal networks. Nature
Communications 9, 3033. https://doi.org/10.1038/s41467-018-05516-7

de Vries, F.T., Liiri, M.E., Bjørnlund, L., Bowker, M.A., Christensen, S., Setälä, H.M., Bardgett,
R.D., 2012. Land use alters the resistance and resilience of soil food webs to drought.
Nature Climate Change 2, 276–280. https://doi.org/10.1038/nclimate1368

Deng, L., Peng, C., Kim, D.G., Li, J., Liu, Y., Hai, X., Liu, Q., Huang, C., Shangguan, Z.,
Kuzyakov, Y., 2021. Drought effects on soil carbon and nitrogen dynamics in global
natural ecosystems. Earth-Science Reviews 214, 103501.
https://doi.org/10.1016/j.earscirev.2020.103501

Dhaou, D., Gros, R., Baldy, V., Adotévi, A., Gaboriau, M., Estevez, Y., Lecareux, C., Dupouyet,
S., Fernandez, C., Bousquet-Mélou, A., 2022. Comparison of leaf litter decomposition and
microbial decomposer communities in fringe and riverine mangroves in French Guiana.
Regional Environmental Change 22, 102. https://doi.org/10.1007/s10113-022-01956-6

Dong, J., Hunt, J., Delhaize, E., Zheng, S.J., Jin, C.W., Tang, C., 2021. Impacts of elevated CO_2

721   on plant resistance to nutrient deficiency and toxic ions via root exudates: A review.

722   Science of the Total Environment 754, 142434.

723   https://doi.org/10.1016/j.scitotenv.2020.142434

724   Enoch, H., Dasberg, S., 1971. The occurrence of high CO_2 concentrations in soil air. Geoderma 6,

725   17–21. https://doi.org/10.1016/0016-7061(71)90048-6

726   García-Palacios, P., Prieto, I., Ourcival, J.-M., Hättenschwiler, S., 2016. Disentangling the litter

727   quality and soil microbial contribution to leaf and fine root litter decomposition responses

728   to reduced rainfall. Ecosystems 19, 490–503. https://doi.org/10.1007/s10021-015-9946-x

729   Gardner, A., Ellsworth, D.S., Crous, K.Y., Pritchard, J., MacKenzie, A.R., 2022. Is photosynthetic

730   enhancement sustained through three years of elevated CO_2 exposure in 175-year-old

731   *Quercus robur*? Tree Physiology 42, 130–144. https://doi.org/10.1093/treephys/tpab090

732   Garrido, M., Hansen, S.K., Yaari, R., Hawlena, H., 2022. A model selection approach to structural

733   equation modelling: A critical evaluation and a road map for ecologists. Methods in

734   Ecology and Evolution 13, 42–53. https://doi.org/10.1111/2041-210X.13742

735   Gauquelin, T., Boer, M., Baldy, V., Fernandez, C., Montes, N., Santonja, M., Orts, J., Reiter, I.,

736   2011. L'O3HP, Oak Observatory at OHP Un site expérimental pour l'étude du

737   fonctionnement et de la biodiversité de la chênaie pubescente face aux changements

738   climatiques. Forêt Méditerranéenne XXXII, 127–132.

739   Gergócs, V., Hufnagel, L., 2016. The effect of microarthropods on litter decomposition depends

740   on litter quality. European Journal of Soil Biology 75, 24–30.

741   https://doi.org/10.1016/j.ejsobi.2016.04.008

742   Glassman, S.I., Weihe, C., Li, J., Albright, M.B.N., Looby, C.I., Martiny, A.C., Treseder, K.K.,

743   Allison, S.D., Martiny, J.B.H., 2018. Decomposition responses to climate depend on

744   microbial community composition. Proceedings of the National Academy of Sciences of

745   the United States of America 115, 11994–11999. https://doi.org/10.1073/pnas.1811269115

746   Handa, I.T., Aerts, R., Berendse, F., Berg, M.P., Bruder, A., Butenschoen, O., Chauvet, E., Gessner,

747   M.O., Jabiol, J., Makkonen, M., McKie, B.G., Malmqvist, B., Peeters, E.T.H.M., Scheu,

748   S., Schmid, B., Van Ruijven, J., Vos, V.C.A., Hättenschwiler, S., 2014. Consequences of

749   biodiversity loss for litter decomposition across biomes. Nature 509, 218–221.

750   https://doi.org/10.1038/nature13247

Harris, N.L., Gibbs, D.A., Baccini, A., Birdsey, R.A., de Bruin, S., Farina, M., Fatoyinbo, L.,
Hansen, M.C., Herold, M., Houghton, R.A., Potapov, P.V., Suarez, D.R., Roman-Cuesta,
R.M., Saatchi, S.S., Slay, C.M., Turubanova, S.A., Tyukavina, A., 2021. Global maps of
twenty-first century forest carbon fluxes. Nature Climate Change 11, 234–240.
https://doi.org/10.1038/s41558-020-00976-6

Hart, K.M., Curioni, G., Blaen, P., Harper, N.J., Miles, P., Lewin, K.F., Nagy, J., Bannister, E.J.,
Cai, X.M., Thomas, R.M., Krause, S., Tausz, M., MacKenzie, A.R., 2020. Characteristics
of free air carbon dioxide enrichment of a northern temperate mature forest. Global Change
Biology 26, 1023–1037. https://doi.org/10.1111/gcb.14786

Hättenschwiler, S., Tiunov, A.V., Scheu, S., 2005. Biodiversity and litter decomposition in
terrestrial ecosystems. Annual Review of Ecology, Evolution, and Systematics 36, 191–
218. https://doi.org/10.1146/annurev.ecolsys.36.112904.151932

Holmstrup, M., Damgaard, C., Schmidt, I.K., Arndal, M.F., Beier, C., Mikkelsen, T.N., Ambus,
P., Larsen, K.S., Pilegaard, K., Michelsen, A., Andresen, L.C., Haugwitz, M., Bergmark,
L., Priemé, A., Zaitsev, A.S., Georgieva, S., Dam, M., Vestergård, M., Christensen, S.,
2017. Long-term and realistic global change manipulations had low impact on diversity of
soil biota in temperate heathland. Scientific Reports 7, 41388.
https://doi.org/10.1038/srep41388

Ilieva-Makulec, K., Olejniczak, I., Szanser, M., 2006. Response of soil micro- and mesofauna
to diversity and quality of plant litter. European Journal of Soil Biology 42, S244–S249.
https://doi.org/10.1016/j.ejsobi.2006.07.030

IPCC, 2021. Summary for Policymakers, in: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors,
S.L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.I., Haung, M.,
Leitzell, K., Lonnoy, E., Matthews, J.B.R., Maycock, T.K., Waterfield, T., Yelekçi, O., Yu,
R., Zhou, B. (Eds.), Climate Change 2021: The Physical Science Basis. Contribution of
Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on
Climate Change. Cambridge University Press, Cambridge, pp. 3–32.
https://doi.org/10.1017/9781009157896.001

Janusz, G., Pawlik, A., Sulej, J., Świderska-Burek, U., Jarosz-Wilkołazka, A., Paszczyński, A.,
2017. Lignin degradation: Microorganisms, enzymes involved, genomes analysis and

evolution. FEMS Microbiology Reviews 41, 941–962.

https://doi.org/10.1093/femsre/fux049

Jardine, P.E., Abernethy, F.A.J., Lomax, B.H., Gosling, W.D., Fraser, W.T., 2017. Shedding light

on sporopollenin chemistry, with reference to UV reconstructions. Review of Palaeobotany

and Palynology 238, 1–6. https://doi.org/10.1016/j.revpalbo.2016.11.014

Jiang, M., Medlyn, B.E., Drake, J.E., Duursma, R.A., Anderson, I.C., Barton, C.V.M., Boer, M.M.,

Carrillo, Y., Castañeda-Gómez, L., Collins, L., Crous, K.Y., De Kauwe, M.G., dos Santos,

B.M., Emmerson, K.M., Facey, S.L., Gherlenda, A.N., Gimeno, T.E., Hasegawa, S.,

Johnson, S.N., Kännaste, A., Macdonald, C.A., Mahmud, K., Moore, B.D., Nazaries, L.,

Neilson, E.H.J., Nielsen, U.N., Niinemets, Ü., Noh, N.J., Ochoa-Hueso, R., Pathare, V.S.,

Pendall, E., Pihlblad, J., Piñeiro, J., Powell, J.R., Power, S.A., Reich, P.B., Renchon, A.A.,

Riegler, M., Rinnan, R., Rymer, P.D., Salomón, R.L., Singh, B.K., Smith, B., Tjoelker,

M.G., Walker, J.K.M., Wujeska-Klause, A., Yang, J., Zaehle, S., Ellsworth, D.S., 2020.

The fate of carbon in a mature forest under carbon dioxide enrichment. Nature 580, 227–

231. https://doi.org/10.1038/s41586-020-2128-9

Joly, F.X., Scherer-Lorenzen, M., Hättenschwiler, S., 2023. Resolving the intricate role of climate

in litter decomposition. Nature Ecology and Evolution 7, 214–223.

https://doi.org/10.1038/s41559-022-01948-z

Kardol, P., Reynolds, W.N., Norby, R.J., Classen, A.T., 2011. Climate change effects on soil

microarthropod abundance and community structure. Applied Soil Ecology 47, 37–44.

https://doi.org/10.1016/j.apsoil.2010.11.001

Kasurinen, A., Riikonen, J., Oksanen, E., Vapaavuori, E., Holopainen, T., 2006. Chemical

composition and decomposition of silver birch leaf litter produced under elevated CO_2 and

O_3. Plant and Soil 282, 261–280. https://doi.org/10.1007/s11104-005-6026-6

Koltz, A.M., Classen, A.T., Wright, J.P., 2018. Warming reverses top-down effects of predators

on belowground ecosystem function in Arctic tundra. Proceedings of the National

Academy of Sciences of the United States of America 115, E7541–E7549.

https://doi.org/10.1073/pnas.1808754115

Krishna, M.P., Mohan, M., 2017. Litter decomposition in forest ecosystems: a review. Energy,

Ecology and Environment 2, 236–249. https://doi.org/10.1007/s40974-017-0064-9

Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest Package: Tests in Linear
Mixed Effects Models. Journal of Statistical Software 82, 1–26.
https://doi.org/10.18637/jss.v082.i13

Lang, B., Rall, B.C., Scheu, S., Brose, U., 2014. Effects of environmental warming and drought
on size-structured soil food webs. Oikos 123, 1224–1233. https://doi.org/10.1111/j.1600-
0706.2013.00894.x

Lavallee, J.M., Soong, J.L., Cotrufo, M.F., 2020. Conceptualizing soil organic matter into
particulate and mineral-associated forms to address global change in the 21st century.
Global Change Biology 26, 261–273. https://doi.org/10.1111/gcb.14859

Lee, H., Jeon, J., Kang, M., Cho, S., Park, J., Lee, M., Lee, H., Kim, D., Kim, H.S., 2021. The
resilience of the carbon cycles of temperate coniferous and broadleaved forests to drought.
Forest Ecology and Management 491, 119178.
https://doi.org/10.1016/j.foreco.2021.119178

Lefcheck, J., 2016. piecewiseSEM: Piecewise structural equation modeling in R for ecology,
evolution, and systematics. Methods in Ecology and Evolution 7, 573–579.
https://doi.org/10.1111/2041-210X.12512

Lindo, Z., Whiteley, J., Gonzalez, A., 2012. Traits explain community disassembly and trophic
contraction following experimental environmental change. Global Change Biology 18,
2448–2457. https://doi.org/10.1111/j.1365-2486.2012.02725.x

Lindquist, E.E., Krantz, G.W., Walter, D.E., 2009. Order Mesostigmata, in: A manual of
Acarology. Texas Tech University Press, Lubbock, pp. 124–232.

Liu, D., Keiblinger, K.M., Leitner, S., Mentler, A., Zechmeister-Boltenstern, S., 2016. Is there a
convergence of deciduous leaf litter stoichiometry, biochemistry and microbial population
during decay? Geoderma 272, 93–100. https://doi.org/10.1016/j.geoderma.2016.03.005

Lonsdale, W.M., 1988. Predicting the amount of litterfall in forests of the world. Annals of Botany
61, 319–324. https://doi.org/10.1093/oxfordjournals.aob.a087560

MacKenzie, A.R., Krause, S., Hart, K.M., Thomas, R.M., Blaen, P.J., Hamilton, R.L., Curioni, G.,
Quick, S.E., Kourmouli, A., Hannah, D.M., Comer-Warner, S.A., Brekenfeld, N., Ullah,
S., Press, M.C., 2021. BIFoR FACE: Water–soil–vegetation–atmosphere data from a
temperate deciduous forest catchment, including under elevated CO2. Hydrological
Processes 35, e14096. https://doi.org/10.1002/hyp.14096

Manzoni, S., Schimel, J.P., Porporato, A., 2012. Responses of soil microbial communities to water
stress: Results from a meta-analysis. Ecology 93, 930–938. https://doi.org/10.1890/11-
0026.1

Martínez-Sancho, E., Treydte, K., Lehmann, M.M., Rigling, A., Fonti, P., 2022. Drought impacts
on tree carbon sequestration and water use – evidence from intra-annual tree-ring
characteristics. New Phytologist 236, 58–70. https://doi.org/10.1111/nph.18224

Melguizo-Ruiz, N., Jiménez-Navarro, G., De Mas, E., Pato, J., Scheu, S., Austin, A.T., Wise, D.H.,
Moya-Laraño, J., 2020. Field exclusion of large soil predators impacts lower trophic levels
and decreases leaf-litter decomposition in dry forests. Journal of Animal Ecology 89, 334–
346. https://doi.org/10.1111/1365-2656.13101

Njoroge, D.M., Chen, S.-C., Zuo, J., Dossa, G.G.O., Cornelissen, J.H.C., 2022. Soil fauna
accelerate litter mixture decomposition globally, especially in dry environments. Journal
of Ecology 110, 659–672. https://doi.org/10.1111/1365-2745.13829

Norby, R.J., Cotrufo, M.F., Ineson, P., O'Neill, E.G., Canadell, J.G., 2001. Elevated CO_2, litter
chemistry, and decomposition: A synthesis. Oecologia 127, 153–165.
https://doi.org/10.1007/s004420000615

Norby, R.J., Loader, N.J., Mayoral, C., Ullah, S., Curioni, G., Smith, A.R., Reay, M.K., Van
Wijngaarden, K., Amjad, M.S., Brettle, D., Crockatt, M.E., Denny, G., Grzesik, R.T.,
Hamilton, R.L., Hart, K.M., Hartley, I.P., Jones, A.G., Kourmouli, A., Larsen, J.R., Shi,
Z., Thomas, R.M., MacKenzie, A.R., 2024. Enhanced woody biomass production in a
mature temperate forest under elevated CO_2. Nature Climate Change 14, 983–988.
https://doi.org/10.1038/s41558-024-02090-3

Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara, R., Solymos,
P., Stevens, M., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard,
D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., Evangelista, H., FitzJohn, R.,
Friendly, M., Furneaux, B., Hannigan, G., Hill, M., Lahti, L., McGlinn, D., Ouellette, M.,
Ribeiro, C.E., Smith, T., Stier, A., Ter Braak, C., Weedon, J., 2022. vegan: Community
Ecology Package. R package version 2.6-4. https://CRAN.R-project.org/package=vegan

Olson, J.S., 1963. Energy storage and the balance of producers and decomposers in ecological
systems. Ecology 44, 322–331. https://doi.org/10.2307/1932179

Orians, C.M., Schweiger, R., Dukes, J.S., Scott, E.R., Müller, C., 2019. Combined impacts of
prolonged drought and warming on plant size and foliar chemistry. Annals of Botany 124,
41–52. https://doi.org/10.1093/aob/mcz004

Pinheiro, J.C., Bates, D.M., 2000. Mixed-Effects Models in S and S-PLUS. Springer, New York.
https://doi.org/10.1007/b98882

Pinheiro, J., Bates, D., R Core Team, 2023. nlme: Linear and Nonlinear Mixed Effects Models. R
package version 3.1-162. https://CRAN.R-project.org/package=nlme

Piper, F.I., Fajardo, A., Hoch, G., 2017. Single-provenance mature conifers show higher non-
structural carbohydrate storage and reduced growth in a drier location. Tree Physiology 37,
1001–1010. https://doi.org/10.1093/treephys/tpx061

Prieto, I., Almagro, M., Bastida, F., Querejeta, J.I., 2019. Altered leaf litter quality exacerbates the
negative impact of climate change on decomposition. Journal of Ecology 107, 2364–2382.
https://doi.org/10.1111/1365-2745.13168

Pritchard, S.G., Rogers, H.H., Prior, S.A., Peterson, Curt.M., 1999. Elevated CO_2 and plant
structure: A review. Global Change Biology 5, 807–837. https://doi.org/10.1046/j.1365-
2486.1999.00268.x

Quer, E., Pereira, S., Michel, T., Santonja, M., Gauquelin, T., Simioni, G., Ourcival, J.M., Joffre,
R., Limousin, J.-M., Aupic-Samain, A., Lecareux, C., Dupouyet, S., Orts, J.-P., Bousquet-
Mélou, A., Gros, R., Sagova-Mareckova, M., Kopecky, J., Fernandez, C., Baldy, V., 2022.
Amplified drought alters leaf litter metabolome, slows down litter decomposition, and
modifies home field (dis)advantage in three Mediterranean forests. Plants 11, 2582.
https://doi.org/10.3390/plants11192582

R Core Team, 2023. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Rillig, M.C., Ryo, M., Lehmann, A., Aguilar-Trigueros, C.A., Buchert, S., Wulf, A., Iwasaki, A.,
Roy, J., Yang, G., 2019. The role of multiple global change factors in driving soil functions
and   microbial   biodiversity.   Science   366,   886–890.
https://doi.org/10.1126/science.aay2832

Rouifed, S., Handa, I.T., David, J.-F., Hättenschwiler, S., 2010. The importance of biotic factors
in predicting global change effects on decomposition of temperate forest leaf litter.
Oecologia 163, 247–256. https://doi.org/10.1007/s00442-009-1528-1

Santonja, M., Fernandez, C., Gauquelin, T., Baldy, V., 2015. Climate change effects on litter
decomposition: intensive drought leads to a strong decrease of litter mixture interactions.
Plant and Soil 393, 69–82. https://doi.org/10.1007/s11104-015-2471-z

Santonja, M., Fernandez, C., Proffit, M., Gers, C., Gauquelin, T., Reiter, I.M., Cramer, W., Baldy,
V., 2017. Plant litter mixture partly mitigates the negative effects of extended drought on
soil biota and litter decomposition in a Mediterranean oak forest. Journal of Ecology 105,
801–815. https://doi.org/10.1111/1365-2745.12711

Seastedt, T.R., 1984. The role of microarthopods in decomposition and mineralization process.
Annual Review of Entomology 29, 25–46. https://doi.org/10.1146/annurev.ento.29.1.25

Tan, B., Yin, R., Zhang, J., Xu, Z., Liu, Y., He, S., Zhang, L., Li, H., Wang, L., Liu, S., You, C.,
Peng, C., 2021. Temperature and moisture modulate the contribution of soil fauna to litter
decomposition via different pathways. Ecosystems 24, 1142–1156.
https://doi.org/10.1007/s10021-020-00573-w

Taub, D.R., Wang, X., 2008. Why are nitrogen concentrations in plant tissues lower under elevated
CO_2? A critical examination of the hypotheses. Journal of Integrative Plant Biology 50,
1365–1374. https://doi.org/10.1111/j.1744-7909.2008.00754.x

Toreti, A., Bavera, D., Acosta Navarro, J., Cammalleri, C., de Jager, A., Di Ciollo, C., Hrast
Essenfelder, A., Maetens, W., Magni, D., Masante, D., Mazzeschi, M., Niemeyer, S.,
Spinoni, J., 2022. Drought in Europe August 2022, Publications Office of the European
Union. Luxembourg. https://doi.org/10.2760/264241

Veen, G.F. (Ciska), Freschet, G.T., Ordonez, A., Wardle, D.A., 2015. Litter quality and
environmental controls of home-field advantage effects on litter decomposition. Oikos 124,
187–195. https://doi.org/10.1111/oik.01374

Vestergård, M., Dyrnum, K., Michelsen, A., Damgaard, C., Holmstrup, M., 2015. Long-term
multifactorial climate change impacts on mesofaunal biomass and nitrogen content.
Applied Soil Ecology 92, 54–63. https://doi.org/10.1016/j.apsoil.2015.03.002

Wall, D.H., Bradford, M.A., St. John, M.G., Trofymow, J.A., Behan-Pelletier, V., Bignell, D.E.,
Dangerfield, J.M., Parton, W.J., Rusek, J., Voigt, W., Wolters, V., Gardel, H.Z., Ayuke,
F.O., Bashford, R., Beljakova, O.I., Bohlen, P.J., Brauman, A., Flemming, S., Henschel,
J.R., Johnson, D.L., Jones, T.H., Kovarova, M., Kranabetter, J.M., Kutny, L., Lin, K.-C.,
Maryati, M., Masse, D., Pokarzhevskii, A., Rahman, H., Sabará, M.G., Salamon, J.A.,

Swift, M.J., Varela, A., Vasconcelos, H.L., White, D., Zou, X., 2008. Global
decomposition experiment shows soil animal impacts on decomposition are climate-
dependent. Global Change Biology 14, 2661–2677. https://doi.org/10.1111/j.1365-
2486.2008.01672.x

Wallenstein, M.D., Haddix, M.L., Ayres, E., Steltzer, H., Magrini-Bair, K.A., Paul, E.A., 2013.
Litter chemistry changes more rapidly when decomposed at home but converges during
decomposition–transformation. Soil Biology and Biochemistry 57, 311–319.
https://doi.org/10.1016/j.soilbio.2012.09.027

Walter, D.E., Lindquist, E.E., Smith, I.M., Cook, D.R., Krantz, G.W., 2009. Order
Trombidiformes, in: A Manual of Acarology. Texas Tech University Press, Lubbock, pp.
233–420.

Wang, T., Bell, B.A., Fletcher, W.J., Ryan, P.A., Wogelius, R.A., 2023. Influence of common
palynological extraction treatments on ultraviolet absorbing compounds (UACs) in sub-
fossil pollen and spores observed in FTIR spectra. Frontiers in Ecology and Evolution 11,
1096099. https://doi.org/10.3389/fevo.2023.1096099

Wang, X., Gossart, M., Guinet, Y., Fau, H., Lavignasse-Scaglia, C.-D., Chaieb, G., Michalet, R.,
2020. The consistency of home-field advantage effects with varying climate conditions.
Soil Biology and Biochemistry 149, 107934. https://doi.org/10.1016/j.soilbio.2020.107934

Welti, E.A.R., Roeder, K.A., De Beurs, K.M., Joern, A., Kaspari, M., 2020. Nutrient dilution and
climate cycles underlie declines in a dominant insect herbivore. Proceedings of the
National Academy of Sciences of the United States of America 117, 7271–7275.
https://doi.org/10.1073/pnas.1920012117

Wickham, H., 2016. Elegant Graphics for Data Analysis. Springer-Verlag, New York.

Wilson, A.M., Burtis, J.C., Goebel, M., Yavitt, J.B., 2022. Litter quality and decomposition
responses to drought in a northeastern US deciduous forest. Oecologia 200, 247–257.
https://doi.org/10.1007/s00442-022-05263-z

Xu, S., Wang, J., Sayer, E.J., Lam, S.K., Lai, D.Y.F., 2024. Precipitation change affects forest soil
carbon inputs and pools: A global meta-analysis. Science of the Total Environment 908,
168171. https://doi.org/10.1016/j.scitotenv.2023.168171

Yin, R., Eisenhauer, N., Auge, H., Purahong, W., Schmidt, A., Schädler, M., 2019. Additive effects
of experimental climate change and land use on faunal contribution to litter decomposition.

Soil Biology and Biochemistry 131, 141–148.

https://doi.org/10.1016/j.soilbio.2019.01.009

Yuste, J.C., Peñuelas, J., Estiarte, M., Garcia-Mas, J., Mattana, S., Ogaya, R., Pujol, M., Sardans,

J., 2011. Drought-resistant fungi control soil organic matter decomposition and its response

to temperature. Global Change Biology 17, 1475–1486. https://doi.org/10.1111/j.1365-

2486.2010.02300.x

Zhang, D., Hui, D., Luo, Y., Zhou, G., 2008. Rates of litter decomposition in terrestrial ecosystems:

Global patterns and controlling factors. Journal of Plant Ecology 1, 85–93.

https://doi.org/10.1093/jpe/rtn002

976
977
Figure 1. Litter decomposition (shown as % remaining litter) for the elevated CO_2 experiment (top)
and drought experiment (bottom). The x-axis indicates the number of months since initiation of
the experiment and differs for upper and lower panels. Larger points represent mean ($\pm$ SE) while
smaller points are the raw data values. * = $p < 0.05$, ** = $p < 0.01$, and *** = $p < 0.001$. We have
only provided factor p-values that were statistically significant. Litter bags in the elevated CO_2
experiment were harvested after 8 (H1), 12 (H2), 17 (H3) and 23 (H4) months, while litter bags in
the drought experiment were harvested after 10 (H1), 13 (H2), 21 (H3), and 34 (H4) months. To
improve plot interpretation, points were positioned with greater spacing on the x-axis. Note the
differences in y axis-es between the two experiments if comparing results with one another.
987
988

989
990
991 Figure 2. Principal component analysis (PCA) ordinations of leaf litter functional groups measured by Fourier transform infrared (FTIR)
992 spectroscopy for the elevated CO_2 experiment (top) and drought experiment (bottom) at harvest three (A and C, left) and four (B and D,
993 right). Corresponding functional groups for band numbers (B1, B2, B3, etc.) can be found in Supplementary Information Table S2. *
994 = p < 0.05, ** = p < 0.01, and *** = p < 0.001. We have only provided factor p-values that were statistically significant.

995
996
Figure 3. Litter quality indices of %carbon (%C; top), %nitrogen (%N; middle) and C:N (bottom)
of drought experiment litter at harvest three (left) and four (right). Larger points represent mean
(± SE) while smaller points are the raw data values. * = p < 0.05, ** = p < 0.01, and *** = p <
0.001. We have only provided factor p-values that were statistically significant. T = treatment
and LT = litter origin. Note the differences in y axis-es between the two harvests if comparing
results between each other.
1003
1004

1005

1006
1007
Figure 4. Microbial community indices as measured by phospholipid fatty acid analysis (PLFA)
in litter for the drought experiment after 21 months (harvest 3). For A) (total PLFA) and B) (fungi :
bacteria) larger points represent mean (± SE) while smaller points are the raw data values. C)
Principal component analysis (PCA) ordination of microbial community structure. * = p < 0.05,
** = p < 0.01, and *** = p < 0.001.   We have only provided factor p-values that were statistically
significant.

1014
1015
Figure 5. Structural equation models (SEMs) showing significant pathways between parameters from the elevated CO_2 experiment (n =
56, left) and drought experiment (n = 60, right). Data to parameterize models were derived solely from thethird harvest in both
experiments. Four litterbags were removed from analysis from the elevated CO_2 model as microbial community structure was not
measured. Blue arrows indicate positive, significant pathways, while red pathways indicate negative, significant pathways where the
standardized path estimates are provided. Parameters are coloured by categories: experiment manipulations (treatment and litter origin,
red), litter chemistry and environmental factors (litter moisture and C:N, yellow), biota (animals and microbes, blue) and mass loss
(green). Asterisks indicate significance of path, * = p < 0.05, and *** = p < 0.001. We have only provided parameter p-values that
were statistically significant. Numbers above individual parameters indicate conditional R-square (R^2_c) of models. In the top-right
corner of each figure, goodness-of-fit tests are provided (χ2 and Fisher's C). Please see Supplementary Information Table S8 for AIC
values and goodness-of-fit metrics for all models tested.