Title

The yEvo Mutation Browser: Enhancing student understanding of experimental evolution and genomics through interactive data visualization

Authors

Leah Anderson[1], Jasmine Schoch[1,2], Eleftheria Anastasia[2], Virginia Wang[1,2], Zilong Zeng[1,2], Sayeh Gorjifard[1,3], Maitreya J. Dunham[1,*]

[1]Department of Genome Sciences, University of Washington; Seattle, WA 98195, USA
[2]Department of Computer Science, University of Washington; Seattle, WA 98195, USA
[3]Current address: Department of Genetics, Stanford University School of Medicine; Palo Alto, CA 94305, USA

*Corresponding author
Correspondence: maitreya@uw.edu

Abstract

Experimental evolution is a powerful method for studying the relationship between genotype and phenotype by observing how populations genetically adapt to controlled selective pressures. In educational settings, this approach also offers a dynamic way for students to engage with molecular genetics. One such educational effort, known as "yEvo" (yeast evolution), introduces experimental evolution into high school classrooms, allowing students to evolve the baker's yeast *Saccharomyces cerevisiae* under various stressors and investigate the resulting *de novo* genetic changes. While the hands-on experiments have been successful in fostering student interest and understanding of evolution, the downstream data analysis—interpreting whole-genome sequencing results of evolved yeast compared to the ancestor—remains a challenge. Students often struggle to grasp the significance of their mutated genes and lack the broader context to determine which mutations are most phenotypically relevant. To address these issues, we developed the yEvo Mutation Browser, an intuitive web tool designed to assist students and researchers alike in visualizing and contextualizing genome sequencing data. Developed using R Shiny, this tool features an interactive chromosome map displaying mutated genes, graphs categorizing mutation types, and a gene viewer illustrating specific mutation sites within genes. The app also features an option for non-yEvo-affiliated users to upload their own experimental evolution or genetic screen datasets and compare them with all yEvo data collected since 2018. The yEvo Mutation Browser streamlines data interpretation, helping students understand how organisms employ diverse genetic strategies to adapt to environmental stress. In the future, this framework could be adapted for use with other model organisms, offering a valuable resource for both genetics research and education.

Summary

The yEvo project, a collaboration between high school classrooms and university yeast research laboratories, allows students to experimentally evolve yeast and observe the effects of mutations and natural selection in real time. The yEvo Mutation Browser, a new web tool,

enables users to visualize the evolved strains' mutations at both the gene and chromosome levels, compare their findings with other classrooms using various conditions, and even upload their own data, benefiting both educators and the yeast genetics community.

Introduction

Experimental evolution is a powerful tool for studying the genotype-phenotype relationship by subjecting populations to selective pressures that drive adaptation over time via genetic mutations. In microbiology, experimental evolution is conducted by propagating microbes under suboptimal conditions such as nutrient limitation, drug exposure, or chemical stress, which restricts their growth. Under these selective conditions, rare spontaneous mutations that improve survival and growth become enriched in the population. These mutations can then be detected through whole genome sequencing, allowing researchers to pinpoint genetic changes associated with the selected trait. Follow-up experiments can then verify whether these mutations are causative for the observed phenotypic changes. This approach is not only a robust method for studying molecular adaptation but also provides a broader understanding of the role of genetic variation on meaningful phenotypic changes. In contrast to classical genetic screens—which most typically are designed to find single, large effect mutations—experimental evolution proceeds via the combined effect of multiple mutations with quantitative effects.

Experimental evolution is a valuable tool not only for research but also for education, offering a hands-on approach to understanding key concepts in evolution and genetics (Bennett *et al.* 2021; Cooper *et al.* 2019; Dahan *et al.* 2019; Ratcliff *et al.* 2014; Smith *et al.* 2016; Taylor *et al.* 2024; Matela *et al.* 2025). One example of this is the "yEvo" (yeast evolution) project, a series of laboratory modules designed for high school biology students to experimentally evolve the baker's yeast *Saccharomyces cerevisiae*. Through this approach, students gain practical laboratory skills, such as sterile technique and liquid handling, while directly observing evolutionary processes in action. The yEvo curriculum deepens students' understanding of microbial evolution, genetic variation, and the genetic basis of adaptation, while also boosting confidence in designing experiments and fostering interest in STEM careers (Taylor and Warwick *et al.* 2024). In addition to its educational impact, the data generated in yEvo classrooms has contributed to published scientific studies, offering insights into yeast adaptation and demonstrating the broader applicability of this teaching model (Geck *et al.* 2024; Moresi *et al.* 2023; Taylor *et al.* 2022). The success of yEvo lies in its use of *S. cerevisiae* as an ideal classroom model organism. Yeast is easy to grow, has a well-characterized genome, and adapts to new environments in just a few weeks. Students work in lab groups to expose yeast populations to stressors such as clotrimazole, caffeine, or other agents, simulating real-world evolutionary pressures like exposure to antimicrobial drugs. After the experiments, samples are analyzed for genomic mutations, and students use resources like the *Saccharomyces* Genome Database (SGD) (Engel *et al.* 2024) to hypothesize how specific mutations may drive adaptation.

Despite the success of the hands-on experimental evolution portion of the yEvo project, the downstream analysis and hypothesis-building steps have proven to be more challenging for students. Many students report that the final stages, which involve interpreting whole genome sequencing data and making connections between their mutated gene lists and the stress conditions, feel much less interactive and engaging than the evolution experiments themselves. Historically, our mutation data analysis module had students review lists of mutated genes from their lab groups' samples and conduct independent online research using resources like SGD and primary literature. However, through this approach, students often found it difficult to understand the significance of their mutated genes and how to put the data into context. Additionally, students were lacking the broader context to determine which of their identified genes are most likely to be causative for the observed phenotypic changes, as they only have access to their group's data without knowing how reproducible their results might be. To keep students engaged through the entirety of the project and help them see the big picture, there has been a need for a user-friendly, visually appealing approach to data analysis that makes complex genomic data easier to interpret.

To meet this need, we have developed the "yEvo Mutation Browser" (available at yevo.org/mutation-browser). This web-browser based application (web app) was written using the R Shiny package, and offers an accessible and engaging way for students to interact with their data. The browser was specifically designed to be user-friendly for students while capturing the full scope of their experimental evolution projects, including linking them to the larger yEvo network of classrooms. The tool also allows users to upload their own data locally, expanding its possible use to other yeast geneticists who would like to interact with their own experimental evolution or genetic screen data. We have made the code available such that other users could also create their own instances of the browser populated with the data of their choice. Here, we outline the key features and applications of the yEvo Mutation Browser.

Methods

Classroom experiments and sequencing

Classroom evolution protocols, sample preparation, and sequencing were performed as previously described (Geck et al 2024, Moresi et al 2023, Taylor et al 2022, Taylor and Warwick *et al.* 2024). In high school classrooms, students evolved *Saccharomyces cerevisiae* under various selective pressures including caffeine, clotrimazole, echinocandins, acetic acid, sodium bicarbonate, wood hydrolysate, and tea tree oil. After the experimental evolution phase, university collaborators isolated three individual yeast clones from each lab group's evolved population. One clone per group showing a clear phenotypic improvement was selected for Illumina whole-genome sequencing. The sequencing data were analyzed using a custom variant-calling computational pipeline (available at: https://github.com/leahmarieanderson/exp_evo_variant_calling). Sequencing reads were aligned to the *S. cerevisiae* S288C reference genome using Burrows-Wheeler Aligner (BWA) (Li and Durbin 2009). Mutations were called using three variant callers: gatk4, lofreq, and

freebayes. Quality filters were applied to the variant calling files (VCFs) via bcftools, and ancestor variants were removed from evolved VCFs using bedtools. Variant annotations were then added via a custom annotation python script (also available at the git repository above). Annotations include coding (such as nonsense, missense, synonymous, frameshift) and non-coding mutations (such as intergenic, intronic, non-coding RNA, and retrotransposons). Variants within 200 base pairs upstream of a start codon were labeled 5' upstream mutations. All mutations passing the quality thresholds were manually reviewed in the Integrative Genomics Viewer (IGV) (Robinson *et al.* 2017) to confirm that the variant was both high quality and present only in the evolved sample.

yEvo Mutation Browser R Shiny app

The yEvo Mutation Browser was developed in R and leverages the Shiny package to build the app's user interface (UI). The plotly package was used to create interactive visualizations either as a wrapper around a static ggplot graph or implemented directly. The Shiny web app is deployed on the shinyapps.io platform. When a user accesses the yEvo Mutation Browser, a new session is spun up, loading a comma-separated values (CSV) file storing the yEvo Cumulative Database, which contains all yEvo mutation data dating back to 2018. These data were bundled into the web app during deployment to shinyapps.io, and each new session loads its own isolated copy of the data. The Shiny framework assigns each user a unique session, enabling multiple users to explore the app concurrently without interference. Within a session, users can interactively filter the default dataset or upload their own mutation data in CSV format; uploaded data is then appended only to that user's session-specific copy and exists solely in memory for the duration of that session. These temporary uploads do not modify the original yEvo Cumulative Database or affect other users' sessions.

The mutation browser's UI is split into 2 main sections: The filtering panel and the visualization panel. The filtering panel allows the user to upload their own data and select key attributes to narrow the points being visualized for more thorough analysis and customization. At the top of the visualization panel, the user can select their desired method (Chromosome Map, Variant Pie Chart, SNP Counts, or Gene View) to view the filtered data. The filtered data can also be viewed under the "Table" tab, and downloaded in CSV format.

The yEvo Mutation Browser can be accessed via Shiny apps (https://leahmarieanderson.shinyapps.io/yEvoMutBrowser/) or on the yEvo website (https://yevo.org/mutation-browser/). The web app can also be installed and run locally as per the instructions provided on the GitHub repository.

Software availability and requirements

The yEvo mutation browser is freely available on GitHub at https://github.com/dunhamlab/yEvoMutBrowser as an R application. The installation requires R (>= 4.3.0). We have tested local versions on both macOS and Windows operating systems. Please refer to the Github repository for more information on installation.

<u>Large language model usage</u>

ChatGPT (versions 4o and 4.5) was used to assist with coding sections of the yEvo Mutation Browser by suggesting methods and formatting of code to improve efficiency and readability. It was also employed to support the writing process by offering syntax and wording suggestions aimed at enhancing clarity and coherence, without introducing new content or altering the intended meaning. All authors carefully reviewed and edited the text following the use of GPT to ensure the accuracy and integrity of the manuscript's content.

Results

yEvo Mutation Browser usage and visualizations

The yEvo Mutation Browser is built using R Shiny (https://shiny.posit.co/), featuring a user-friendly interface that allows users to toggle between different data options while the underlying server processes and displays the selected information reactively. This functionality enables users to explore various datasets interactively, generating visualizations tailored to their specific selections. The Mutation Browser has now been used in all 2024-2025 yEvo classrooms, with teacher and student feedback incorporated into its design. Prior to this app, no tool existed for our students to visualize their own data, and the Mutation Browser fills a major gap in the broader yEvo project. The source code and datasets used to develop this tool are publicly available at https://github.com/dunhamlab/yEvoMutBrowser. In the following sections, we describe the application's usage, its user interface, and the visualizations users encounter while navigating the app.

<u>Application usage</u>

Upon launch, all cumulative yEvo data collected since 2018 is loaded into memory. This data exists in a comma-separated-values (CSV) file which is then used to generate all interactive visualizations (see Methods). Users begin by selecting the data subset they wish to explore. One option is to view data by classroom, where users can choose either the entire class dataset or individual lab group data using a dropdown menu after specifying their class (Figure S1A); this is particularly useful for students who have completed the experimental evolution module of yEvo with their specific classroom. Alternatively, users can select "View by Selection Condition," which displays all the cumulative yEvo data generated under specific selection conditions (Figure S1B). Users are able to see all the selection conditions used in yEvo since 2018: the antifungal agents clotrimazole, caspofungin, micafungin, and tea tree oil, and the industrial/environmental conditions acetic acid, caffeine, sodium bicarbonate, and wood hydrolysate. While most data in the app is from evolution of the haploid lab strain S288C, some experiments began with different backgrounds, including a diploid version of S288C, a baking

strain, or genetically modified knockout strains. If the selected dataset contains multiple strain options, users can specify which strain(s) they want to view using the "Ancestor Strain" dropdown menu (Figure S1C and D). For simplicity, results from all strains are viewed against the S288C reference genome, though this could be modified in the future (see discussion).

The app also allows users to upload their own Variant Call Format (VCF) files in CSV format, which enables any user to visualize their own data (see Methods). Instructions for uploading datasets are provided in both the "Tutorial" tab within the app and on the README on our GitHub repository. When uploading, users are prompted to input their name and year, which then become accessible via the "View by Class" dropdown menu (Figure S1E). This new data is then temporarily appended to the yEvo dataset available on the app for that user's session. If the selection condition in the uploaded dataset matches pre-existing yEvo data on the server, the new data is added to the cumulative dataset for that condition, allowing users to see how their results compare with patterns observed in our yEvo experiments. This feature is beneficial not only for educators but also for researchers seeking a comparison of their data with existing experimental evolution datasets.

Once the desired dataset is selected, users can access all of the available data visualizations. Users can also modify their dataset selections while viewing any of the visual plots, which are described in the following sections.

Figure 1: Visualization examples of a single lab group vs cumulative classroom data.
Visualizations of data from the Brewer 2022 classroom, where yeast was evolved in caffeine. (A) Chromosome plot showing genes mutated in a single sample (left) or the whole class (right). The scale bar on the right shows the number of samples that had that gene mutate. (B) Pie chart showing mutation types. (C) Gene view displaying where each coding mutation affected the gene, by amino acid position.

<u>Visualizations</u>

1. Chromosome plot

The first visualization users encounter in the yEvo Mutation Browser is the chromosome plot, which provides a linear representation of the 16 nuclear chromosomes along with the mitochondrial genome of *S. cerevisiae*. This plot displays the locations of all mutated genes in the selected dataset, marked at their chromosomal positions. For simplicity of viewing, only mutations in coding regions are shown on the chromosome plot. Tick marks along the linear chromosomes indicate these sites, and hovering the mouse over each tick mark reveals the name of the mutated gene, as well as how many times it was mutated in repeat experiments, which is also indicated visually using a color scale. Users can also zoom in on specific regions by clicking and dragging, offering a closer look at mutations in a given area (Figure S2).

The chromosome plot allows students to directly compare their results with those from other students within their own classrooms as well as to previous years and other yEvo sites performing similar experiments, providing a more robust and visually intuitive way to assess the significance of genetic mutations. Figure 1A highlights this feature. The left panel shows data from a single lab group who evolved yeast in increasing concentrations of caffeine, where four mutated genes are mapped: *SVF1, PDR1, NGL3,* and *SKY1*. Once a user switches the view to "View by Condition" and selects "caffeine," the chromosome map is labeled with how recurrently each gene in the dataset was mutated in the cumulative caffeine yEvo data, with a white-to-red gradient displaying frequency (Figure 1A, right panel). Using the "Selection condition" view, it can be seen that *PDR1* and *SKY1* were mutated in 21 and 7 samples respectively, while *NGL3* and *SVF1* were unique to the single sample. This reinforces the likelihood that *PDR1* and *SKY1* play a key role in adaptation to caffeine, which has now been experimentally verified since the high school experiments were carried out (Geck *et al.* 2024).

2. Pie chart of mutation types

The second visualization provided by the yEvo Mutation Browser is a pie chart that displays the distribution of mutation types identified by the sequencing analysis pipeline. Unlike the chromosome plot, which focuses solely on the locations of mutated genes, the pie chart encompasses all variants, including those in non-coding regions (such as intergenic, intronic, tRNA, rRNA, noncoding RNA, and autonomously replicating sequence regions). This gives students a broader overview of the types of genetic changes that occurred in their evolution experiments. One key insight students gain from this visualization is that, regardless of the selection condition, the majority of mutations in the evolved samples are those that alter protein sequences. For instance, the left panel of Figure 1B shows the mutation types for one lab

group's data, while the right panel shows cumulative data from multiple classrooms that evolved yeast in caffeine. In both cases, the vast majority of mutations are missense mutations. This becomes a valuable teaching tool, helping students understand that while mutations occur randomly across the genome, those that lead to phenotypic changes are primarily found in coding sequences, which are then selected for during the course of the experiment.

3. Mutation spectrum

The third visualization offered by the yEvo Mutation Browser is the mutation spectrum, depicted in Figure 3C and D. This plot illustrates the distribution of single nucleotide polymorphisms (SNPs) and small insertions and deletions (indels). By categorizing the different types of SNPs, the mutation spectrum provides insights into the underlying biochemical processes that drive DNA mutations. Students can observe that transitions—mutations where a purine is replaced with another purine (A to G / G to A) or a pyrimidine with another pyrimidine (C to T / T to C)—are more frequent than transversions, which involve purine-to-pyrimidine exchanges and vice versa. This plot displays that SNPs and indels can occur at different frequencies during evolution, and that different mutagen types or background genotypes can cause altered mutation spectra compared to the natural mutation rate. Recognizing the prevalence of transitions over transversions reinforces the concept that certain mutations are inherently more likely to occur based on the chemical stability and replication mechanisms of DNA. Further applications of this visualization, such as using it to explore how yeast strains with different mutation rates exhibit distinct mutation spectra, will be described later in this paper.

4. Gene view

The final visualization is "Gene view," a linear lollipop plot that displays the position of mutations along the protein product and 200 bp upstream of the start codon for each mutated gene (Figure 1C). In this view, users can select a mutated gene from a dropdown menu based on the selected dataset. Once a gene is selected, the plot reveals the length of the amino acid sequence and color-coded lollipops indicating where mutations occurred. The colors differentiate between mutation types, such as missense, nonsense, frameshift, synonymous, or 5'-upstream mutations. The height of each lollipop reflects how frequently that particular site in the protein was mutated within the dataset. For example, the left panel of Figure 1C shows the gene view for an evolved yeast strain, where the dropdown menu displays the four genes that were mutated. In contrast, the right panel illustrates the cumulative caffeine data, with all genes mutated in that condition in the dropdown menu and the lollipop heights showing the recurrence of mutations at specific sites. Additionally, hovering the mouse over a lollipop provides information about the mutation, including the amino acid site number, residue type, and the resulting amino acid caused by the DNA mutation (if any).

Furthermore, the gene view includes a "Learn about gene" button, which redirects users to the SGD locus page for the selected gene. This feature allows students to explore the gene's function in yeast and use this information to generate hypotheses about how mutations in that gene might contribute to observed phenotypic changes. Overall, the gene view visualization is designed to help students better conceptualize where mutations occur within genes and to encourage critical thinking about why certain regions may be more frequently mutated and by what types of changes. By connecting mutation sites to their potential impact on protein function, students gain deeper insights into the molecular mechanisms driving adaptation in their experiments.

Further educational applications

While we have outlined the foundational educational use cases for the yEvo Mutation Browser, there are additional benefits that can offer valuable teaching points for more advanced biology classrooms.

Signatures of mutation effects

As yEvo students explore gene functions and hypothesize which mutations might be phenotypically causative, predicting how specific mutations alter gene function remains a challenge. Does increased survival arise from the loss of gene function, or from a gain or alteration of function? Historically, students could only make educated guesses based on the mutation types in their genes of interest (synonymous, missense, nonsense, etc). With the yEvo Mutation Browser, students can now make more informed predictions about the nature of mutations by analyzing mutation signatures from cumulative data. For instance, when viewing cumulative data in Gene View for a particular condition, a student might observe that mutations in a particular gene are clustered in one region and are exclusively missense. A tight clustering of mutations at a single site, especially if only missense, suggests a potential gain-of-function (GOF) mutation, as there are limited ways to generate a GOF mutation that enhances fitness. Conversely, if a gene exhibits mutations scattered throughout, with a mix of missense and nonsense mutations, this pattern likely indicates a loss-of-function (LOF) mutation, as many different types of mutations can disrupt gene function. These mutation signatures have been extensively studied and are seen in various human diseases, including cancer (Gerasimavicius et al. 2022; Prior et al. 2012; Sivley et al. 2018; Stehr et al. 2011; Turner et al. 2015). This offers teachers a valuable opportunity to connect the mutation patterns observed in the classroom to real-world medical applications, such as cancer and other genetic disorders, making the lessons more relatable and impactful for students.

An example of these mutation signatures is depicted in Figure 2, which illustrates cumulative data for micafungin. Students may notice using Chromosome View that two genes are particularly enriched for mutations, *FKS1* and *GSC2.* In Gene View, *FKS1* shows mutations clustered at two distinct locations, all of which are missense (Figure 2A). This suggests that mutations in *FKS1* may lead to a gain or alteration of gene function, a pattern consistent with prior research in both pathogenic yeast and *Saccharomyces* species (Katiyar and Edlind 2009; Johnson and Edlind 2012; Katiyar *et al.* 2006; Laverdiere *et al.* 2006; Park *et al.* 2005; Thompson *et al.* 2008; Balashov *et al.* 2006; Cleary *et al.* 2008; Garcia-Effron *et al.* 2008; Khan *et al.* 2007; Johnson *et al.* 2011; Perlin 2007). In contrast, *GSC2* shows mutations spread across the gene, with many being nonsense and frameshift mutations (Figure 2B), suggesting that loss of function may be responsible for the observed phenotype. In fact, this aligns with prior work which has shown that *gsc2Δ* can confer echinocandin resistance in *S. cerevisiae* (Lesage *et al.* 2004). Armed with this information, students can generate more informed hypotheses. They may discover through SGD and primary literature that *FKS1* and *GSC2* are homologous genes involved in β-1,3-glucan synthesis, a critical component of fungal cell wall integrity. Moreover, they may learn that micafungin targets the fungal cell wall by inhibiting the enzyme β-1,3-glucan synthase (Denning 2002), which is composed of catalytic subunits encoded by *FKS1* and *GSC2* (Lesage and Bussey 2006). From this, students could hypothesize that a GOF mutation in *FKS1* may alter the enzyme's structure in a way that reduces the binding affinity of micafungin, allowing the enzyme to retain functionality. In the same vein, loss of *GSC2* function (which the app shows almost always co-occurs with an *FKS1* missense mutation) could enable the yeast to prioritize *FKS1*-based glucan synthesis, providing resistance to micafungin without completely impairing cell wall biosynthesis. While this idea is still hypothetical, (and we are currently exploring this hypothesis in our lab), we believe that involving students in hypothesis building on a concept that has not yet been explored can be critical for their learning and interest in science. This example also displays how leveraging the power of cumulative data visualizations in the yEvo Mutation Browser can help students think more deeply about gene function and gain a better understanding of the molecular mechanisms driving adaptive evolution.

Figure 2: Gene view can reveal mutation patterns for specific genes. Mutations in *FKS1* (A) and *GSC2* (B) display differences in mutation patterns for yeast samples evolved in caspofungin.

<u>Visualizing the effects of a mutator</u>

While most yEvo classrooms conduct experimental evolution by relying on the natural mutation rate of *S. cerevisiae*, one class was interested in exploring how a higher mutation rate might affect the evolutionary process. Since many mutagenesis methods are not safe for classroom use, we provided them with a strain of *S. cerevisiae* with a deletion of the *MSH2* gene, which encodes the Msh2 protein involved in DNA mismatch repair. Loss of *MSH2* leads to an 85-fold increase in the spontaneous mutation rate (Alani *et al.* 1995; Reenan and Kolodner 1992) and mutations in the human homolog elevate the risk of cancers (Fishel *et al.* 1993, Bridge *et al.* 2014, Win *et al.* 2017, Bowen *et al.* 2025). The class performed experimental evolution in micafungin side-by-side with both the *msh2Δ* mutator strain and the wt S288C lab control strain that is normally used for yEvo experiments. While increasing the mutation rate might seem advantageous for rapidly observing adaptation, it can significantly complicate the sequencing data analysis. For example, instead of observing clear, repeated mutations concentrated in a handful of genes consistently linked to antifungal resistance, the mutator strain accumulates numerous additional "hitchhiker" mutations across the genome, which could be completely unrelated to the selected phenotype. Consequently, experimentally evolved mutator strains often have dozens of mutated genes, most of which do not contribute to the adaptive phenotype, thereby complicating the identification and verification of causative mutations. This concept can be challenging for students who are just learning about evolutionary biology, but the visualizations provided by the yEvo Mutation Browser clearly illustrate how employing a higher mutation rate during experimental evolution can affect the results.

To see the effects of strain backgrounds, we added a feature in the user interface that allows users to toggle between strain backgrounds when a dataset contains more than one. Therefore, when selecting the "micafungin" condition, users can choose between the S288C strain and the *msh2del* strain (Figure S1C). By comparing the chromosome plots of these two strains, users can observe that the mutator strain shows far more mutated genes, most of which are color coded white, indicating that they were only mutated in a single sample. Conversely, the S288C strain has fewer mutated genes overall, but a larger proportion of these genes were recurrently mutated across multiple samples (Figure 3A and B). Interestingly, in both mutator and non-mutator datasets, *FKS1* and *GSC2* remained the most frequently mutated genes. This indicates that, in this particular example, the mutator strain did not provide additional insights into new antifungal drug resistance mechanisms, as it recapitulated the same key mutations seen in the S288C control strain.

The mutation spectrum visualization also reveals striking differences between these parallel experiments. Specifically, the mutator strain exhibits a significant bias towards indels compared to SNPs, a finding consistent with previous research (Reenan and Kolodner 1992; Alani *et al.* 1994; Greene and Jinks-Robertson, 2001; Lang *et al.* 2013). For more advanced classrooms, this provides an opportunity to delve deeper into the molecular mechanisms driving these biases, promoting critical thinking about mutation processes. Additionally, students can use this data to reinforce the concept of hitchhiker mutations, as the mutator strain displays many more mutations across the genome, most of which are likely not significant but are carried along with beneficial mutations elsewhere in the genome. Since these results are available to all via the

Mutation Browser, teachers could refer to them instead of performing the experiment themselves in order to demonstrate why relying on the natural mutation rate is sufficient for experimental evolution, while also reinforcing key concepts such as hitchhiker mutations and the complexities of mutation analysis.

Figure 3: Visualizations reveal differences in mutation frequency and spectra between strains.
When compared to data from an evolved wt S288C strain (A and C), mutations in a mutator strain
missing the *MSH2* mismatch repair gene occur in more locations across the genome (A), and show
a bias towards indel mutations (D).

Discussion

Here we describe the yEvo Mutation Browser, an application built in R Shiny, designed to provide comparative visualizations of *S. cerevisiae* experimental evolution data. We were motivated to create this application to enhance yEvo students' understanding of the evolutionary experiments they conducted in the classroom. Previously, our students were limited to reviewing lists of mutated genes from their lab groups' samples and conducting independent research using data available online. With the yEvo Mutation Browser, students can now not only visualize the genomic locations of their mutations but also explore how frequently these mutations recur in repeat experiments under the same or similar conditions. This added layer of interactivity allows students to engage more deeply with their data and recognize patterns that may not have been apparent before.

Each visualization in the yEvo Mutation Browser contributes uniquely to student learning by offering insights that were not previously accessible through static gene lists. The chromosome plot, for instance, helps students better grasp the likelihood that specific genes are causative of observed phenotypes. By enabling a direct comparison of their results with those from other classrooms and experiments, students can see how certain genes are commonly mutated in repeated experiments, increasing the likelihood that mutations in these genes can affect the observed phenotype. This helps students gain broader context, which is more effective than the previous approach of hypothesizing gene significance through isolated online searches. The pie chart visualization breaks down mutation types, helping students understand why certain mutations, such as missense mutations, occur at a higher frequency. This becomes a valuable teaching tool, helping students understand that while mutations occur randomly across the genome, those that lead to phenotypic changes are primarily found in coding sequences, which are then selected for during the course of the experiment. For more advanced classrooms, the pie chart can also be used to explore why certain mutation types, such as 5'-UTR and nonsense mutations, are also frequently observed in evolved yeast samples. This also provides an opportunity to introduce students to the concept of hitchhiker mutations—mutations that may not directly contribute to adaptation but are carried along with beneficial mutations. This concept can be reinforced by comparing individual samples containing intergenic or synonymous mutations to their corresponding chromosome plot, helping students identify whether these mutations co-occur with coding mutations, which are more likely to be phenotypically significant. The mutation spectrum provides an opportunity to explore the biochemical mechanisms behind mutation events, distinguishing between transitions and transversions. It also provides the opportunity to display how different mutagens or genetic backgrounds can lead to specific mutation signatures. Finally, the gene view allows students to connect the presence of mutations with their possible functional effects, displaying "hotspot" mutation sites and the corresponding amino acid changes. This offers teachers a valuable opportunity to connect the mutation patterns observed in the classroom to real-world medical applications, such as cancer, antibiotic resistance, viral evolution, and other genetic processes, making the lessons more relatable and impactful for students.The integration of the "Learn about gene" button further streamlines the learning process by directing students to detailed information on SGD,

facilitating better learning, engagement, and hypothesis development. This feature allows students to explore the gene's function in yeast and use this information to generate hypotheses about how mutations in that gene might contribute to observed phenotypic changes. Finally, beyond its use in the classroom, the yEvo Mutation Browser is a tool that can benefit the broader yeast research community by allowing users to upload and compare their data with existing yEvo datasets, enabling quick and easy comparisons to the results of our yeast evolution experiments.

A key advantage of the app is its ability to combine data from all classrooms using the same experimental evolution conditions, enabling students to see which of the genes in their groups' data are recurrently mutated and therefore more likely to be phenotypically causative. Students can also compare mutations observed under one condition to those identified in other evolution conditions, helping them hypothesize about why certain mutations may be beneficial in some environments but not others. For instance, when students explore cumulative data from multiple yEvo experiments within the browser, they might notice that almost all of the conditions lead to mutations in one or more *PDR* genes, particularly *PDR5, PDR3,* and *PDR1.* This observation may prompt students to investigate what makes this gene family so commonly mutated in various environments. Through further exploration, they might learn that the *PDR5* gene encodes a drug transporter, its expression is controlled by the transcription factors Pdr1 and Pdr3*,* and that variants in all three genes can be involved in adaptive responses to drug exposure (Katzmann *et al.* 1996, Kolaczkowski *et al.* 1996 and 1998)**.** In this example, by identifying the prevalence of mutations in the *PDR* family across multiple conditions, students gain insight into how cells mechanistically respond to different stressors. This serves as a concrete example of how cells utilize drug pumps and related mechanisms to handle diverse environmental challenges, reinforcing broader lessons about cellular adaptation and stress tolerance.

While the yEvo Mutation Browser offers an interactive and user-friendly platform for exploring SNPs and indels in evolved yeast samples, a key limitation is its inability to display other important genomic changes, such as copy number variants (CNVs) or transposon insertions. These types of mutations, although significant, have not been included in this initial version of the app. However, there is potential to expand the tool to visualize these mutations in the future, particularly for advanced students or researchers. In published studies using yEvo data, university collaborators have conducted CNV and transposon analyses, but these were not a part of classroom discussions. This is a notable limitation, as these mutations often play a critical role in adaptation under specific conditions. For instance, in clotrimazole-evolved clones, 14 out of 99 contained transposon insertions (Taylor *et al.* 2022). Additionally, while *PDR5* was frequently mutated in both clotrimazole and caffeine-evolved clones, clotrimazole-evolved samples often involved copy number gains, whereas caffeine-evolved clones had mostly point mutations (Geck *et al.* 2024). These distinctions are currently missing from the Mutation Browser, which focuses solely on SNPs and small indels. Nonetheless, the app provides a valuable foundation for visualizing simpler mutation types, and we are actively exploring ways to

incorporate CNVs and transposons to provide a more comprehensive view of genomic changes during experimental evolution.

Ongoing development of the yEvo Mutation Browser is focused on optimizing it based on feedback from students and teachers in yEvo classrooms. The reference genome used in the app can also be modified to better represent diverse *S. cerevisiae* strain backgrounds, such as the baking strain used for some prior experiments. For researchers or educators interested in adapting the browser for other strains or species, our detailed GitHub README provides step-by-step instructions. We encourage users to reach out to the authors for assistance if needed.

We believe that yEvo is an incredibly rewarding and effective way to introduce students to scientific inquiry by teaching them microbiology laboratory skills, practicing scientific thinking, and providing an opportunity to participate in real research. The yEvo Mutation Browser enhances this educational experience by providing students with a more meaningful way to visualize and understand the data they generate in the classroom. Through this tool, students can see that while mutations may occur randomly across the genome, the genes that are frequently mutated independently are the ones most likely driving adaptation. This insight gives students a clearer picture of genetic causality in evolutionary contexts, deepening their understanding of how mutations shape phenotypic outcomes. In conclusion, the yEvo Mutation Browser has allowed us to fill a long-standing gap in our program, bringing the full scope of the yEvo project into clearer focus for students by providing interactive visualizations of students' data. As we continue to expand yEvo into additional classrooms, we believe that this resource will grow into an invaluable asset for educators, students, and researchers alike, further strengthening the bridge between research and education.

Data availability

All cumulative yEvo data used in the Mutation Browser as well as the scripts used to generate the app are available at our GitHub repository: https://github.com/dunhamlab/yEvoMutBrowser.git.

Acknowledgements

We are deeply grateful to Renee Geck for her early input and guidance during the initial brainstorming stages of the project at the 2022 UW Genome Sciences Hackathon, her feedback during development of the app, and her thoughtful review of this manuscript. We also thank all the participants of the 2022 and 2023 GS Hackathon yEvo teams: Nicole Wang (especially for her efforts to rewrite the Python annotation script used in our analysis pipeline), Dennis Godin, Sophie Gibson, Molly Perchlik, Luana Paleologu, Rachel Powell, Conor Camplisson, Matthew Chaw, and Lucas Kampman.

We are also grateful to the yEvo teachers who implemented the app in their classrooms during the 2024-2025 school year and to their students who completed the evolution modules and provided user feedback. In particular, we thank: David Maxwell (The Pingry School), Kelsey Kovarik and Frances Cheong (The Downtown School), Ryan Skophammer (Westridge School), Kenneth Berger, Jonathan Schaper, and Lee Anne Eareckson (Moscow High School), and Michael Delmont (Frontier Senior High School). We are also grateful to Rebecca Brewer and her students at Troy High School for their work on the caffeine evolution experiments, whose data was used for Figure 1.

We acknowledge Sarah McClymont and Brenda Andrews at University of Toronto, along with their undergraduate students, who analyzed over 40 of the evolved samples from Moscow High. Their data was integrated into the cumulative yEvo data presented in the browser. We'd also like to thank Paul Rowley and Claire Warren from University of Idaho for their support in sample curation, phenotype testing, and teacher communication, as well as their continued dedication to the yEvo project as a whole. We are especially grateful to M. Bryce Taylor for spearheading the yEvo project at its inception and for generating and analyzing much of the initial data that served as the foundation for the Mutation Browser, and we also appreciate his feedback on the project when the Mutation Browser was first getting started. Finally, we thank Faheem Azeemi and Adrianna Matos-Nieves at the Broad Institute for joining the yEvo Mutation Browser team and for their valuable ideas in brainstorming features for future iterations of this tool.

Funding

We acknowledge support from the National Institute of General Medical Sciences grant R25GM154336, National Science Foundation grant 1817816, and the National Human Genome Research Institute T32 HG000035. The 2023 Genome Sciences Hackathon was supported by a UW Wellness Grant.

Literature cited

Alani E, Reenan RA, Kolodner RD. Interaction between mismatch repair and genetic recombination in *Saccharomyces cerevisiae. Genetics*. 1994;137(1):19-39. doi:10.1093/genetics/137.1.19

Alani E, Chi NW, Kolodner R. 1995. The *Saccharomyces cerevisiae* Msh2 protein specifically binds to duplex oligonucleotides containing mismatched DNA base pairs and insertions. Genes Dev. 9(2):234–247. doi:10.1101/gad.9.2.234.

Balashov SV, Park S, Perlin DS. 2006. Assessing Resistance to the Echinocandin Antifungal Drug Caspofungin in *Candida albicans* by Profiling Mutations in *FKS1*. Antimicrob Agents Chemother. 50(6):2058–2063. doi:10.1128/AAC.01653-05.

Bennett KF, Arriola PE, Marsh TL, Mineo PM, Raimondi SL, Shaffer CL. 2021. CURE in a Box: An Online CURE for Introductory Biology Majors That Incorporates *Vision and Change*. J Microbiol Biol Educ. 22(1):ev22i1.2325. doi:10.1128/jmbe.v22i1.2325.

Botstein D, Fink GR. 2011. Yeast: An Experimental Organism for 21st Century Biology. Genetics. 189(3):695–704. doi:10.1534/genetics.111.130765.

Bowen MB, Melendez B, Zhang Q, et al. Mitochondrial defects and metabolic vulnerabilities in Lynch syndrome–associated MSH2-deficient endometrial cancer. *JCI Insight*. 2025;10(6). doi:10.1172/jci.insight.185946

Bridge G, Rashid S, Martin S. DNA Mismatch Repair and Oxidative DNA Damage: Implications for Cancer Biology and Treatment. *Cancers*. 2014;6(3):1597-1614. doi:10.3390/cancers6031597

Cleary JD, Garcia-Effron G, Chapman SW, Perlin DS. 2008. Reduced *Candida glabrata* Susceptibility Secondary to an *FKS1* Mutation Developed during Candidemia Treatment. Antimicrob Agents Chemother. 52(6):2263–2265. doi:10.1128/AAC.01568-07.

Cooper VS, Warren TM, Matela AM, Handwork M, Scarponi S. 2019. EvolvingSTEM: a microbial evolution-in-action curriculum that enhances learning of evolutionary biology and biotechnology. Evo Edu Outreach. 12(1):12. doi:10.1186/s12052-019-0103-4.

Dahan O, Dorfman B-S, Sayin S, Rosener B, Hua T, Yarden A, Mitchell A. 2019. Harnessing robotic automation and web-based technologies to modernize scientific outreach. PLoS Biol. 17(6):e3000348. doi:10.1371/journal.pbio.3000348.

Denning DW. 2002. Echinocandins: a new class of antifungal. Journal of Antimicrobial Chemotherapy. 49(6):889–891. doi:10.1093/jac/dkf045.

Desnos-Ollivier M, Bretagne S, Raoux D, Hoinard D, Dromer F, Dannaoui E. 2008. Mutations in the *fks1* Gene in *Candida albicans* , *C. tropicalis* , and *C. krusei* Correlate with Elevated Caspofungin MICs Uncovered in AM3 Medium Using the Method of the European Committee on Antibiotic Susceptibility Testing. Antimicrob Agents Chemother. 52(9):3092–3098. doi:10.1128/AAC.00088-08.

Engel SR, Aleksander S, Nash RS, Wong ED, Weng S, Miyasato SR, Sherlock G, Cherry JM (2025) *Saccharomyces* Genome Database: Advances in Genome Annotation, Expanded Biochemical Pathways, and Other Key Enhancements. Genetics. 2024 Nov 12:iyae185. doi: 10.1093/genetics/iyae185. PMID:39530598

Fishel R, Lescoe MK, Rao MRS, et al. The human mutator gene homolog *MSH2* and its association with hereditary nonpolyposis colon cancer. *Cell*. 1993;75(5):1027-1038. doi:10.1016/0092-8674(93)90546-3

Garcia-Effron G, Kontoyiannis DP, Lewis RE, Perlin DS. 2008. Caspofungin-Resistant *Candida tropicalis* Strains Causing Breakthrough Fungemia in Patients at High Risk for Hematologic Malignancies. Antimicrob Agents Chemother. 52(11):4181–4183. doi:10.1128/AAC.00802-08.

Geck RC, Moresi NG, Anderson LM, yEvo Students, Addepalli I, Aggarwal D, Agnihotri P, Ali AA, Amorosi CJ, Anand A, et al. 2024. Experimental evolution of *Saccharomyces cerevisiae* for

caffeine tolerance alters multidrug resistance and target of rapamycin signaling pathways. Steinmetz L, editor. G3: Genes, Genomes, Genetics. 14(9):jkae148. doi:10.1093/g3journal/jkae148.

Gerasimavicius L, Livesey BJ, Marsh JA. 2022. Loss-of-function, gain-of-function and dominant-negative mutations have profoundly different effects on protein structure. Nat Commun. 13(1):3895. doi:10.1038/s41467-022-31686-6.

Greene CN, Jinks-Robertson S. Spontaneous Frameshift Mutations in *Saccharomyces cerevisiae*: Accumulation During DNA Replication and Removal by Proofreading and Mismatch Repair Activities. *Genetics*. 2001;159(1):65-75. doi:10.1093/genetics/159.1.65

Johnson ME, Edlind TD. 2012. Topological and Mutational Analysis of Saccharomyces cerevisiae Fks1. Eukaryot Cell. 11(7):952–960. doi:10.1128/EC.00082-12.

Johnson ME, Katiyar SK, Edlind TD. 2011. New Fks Hot Spot for Acquired Echinocandin Resistance in *Saccharomyces cerevisiae* and Its Contribution to Intrinsic Resistance of *Scedosporium* Species. Antimicrob Agents Chemother. 55(8):3774–3781. doi:10.1128/AAC.01811-10.

Kahn JN, Garcia-Effron G, Hsu M-J, Park S, Marr KA, Perlin DS. 2007. Acquired Echinocandin Resistance in a *Candida krusei* Isolate Due to Modification of Glucan Synthase. Antimicrob Agents Chemother. 51(5):1876–1878. doi:10.1128/AAC.00067-07.

Katiyar S, Pfaller M, Edlind T. 2006. *Candida albicans* and *Candida glabrata* Clinical Isolates Exhibiting Reduced Echinocandin Susceptibility. Antimicrob Agents Chemother. 50(8):2892–2894. doi:10.1128/AAC.00349-06.

Katiyar SK, Edlind TD. 2009. Role for Fks1 in the Intrinsic Echinocandin Resistance of *Fusarium solani* as Evidenced by Hybrid Expression in *Saccharomyces cerevisiae*. Antimicrob Agents Chemother. 53(5):1772–1778. doi:10.1128/AAC.00020-09.

Katzmann DJ, Hallstrom TC, Mahé Y, Moye-Rowley WS. 1996. Multiple Pdr1p/Pdr3p Binding Sites Are Essential for Normal Expression of the ATP Binding Cassette Transporter Protein-encoding Gene. Journal of Biological Chemistry. 271(38):23049–23054. doi:10.1074/jbc.271.38.23049.

Kolaczkowski M, Kolaczkowska A, Luczynski J, Witek S, Goffeau A. 1998. *In Vivo* Characterization of the Drug Resistance Profile of the Major ABC Transporters and Other Components of the Yeast Pleiotropic Drug Resistance Network. Microbial Drug Resistance. 4(3):143–158. doi:10.1089/mdr.1998.4.143.

Kolaczkowski M, Michel VDR, Cybularz-Kolaczkowska A, Soumillion J-P, Konings WN, André G. 1996. Anticancer Drugs, Ionophoric Peptides, and Steroids as Substrates of the Yeast Multidrug Transporter Pdr5p. Journal of Biological Chemistry. 271(49):31543–31548. doi:10.1074/jbc.271.49.31543.

Lang GI, Parsons L, Gammie AE. 2013. Mutation Rates, Spectra, and Genome-Wide Distribution of Spontaneous Mutations in Mismatch Repair Deficient Yeast. G3: Genes, Genomes, Genetics. 3(9):1453–1465. doi:10.1534/g3.113.006429.

Laverdière M, Lalonde RG, Baril J-G, Sheppard DC, Park S, Perlin DS. 2006. Progressive loss of echinocandin activity following prolonged use for treatment of *Candida albicans* oesophagitis. Journal of Antimicrobial Chemotherapy. 57(4):705–708. doi:10.1093/jac/dkl022.

Lesage G, Bussey H. 2006. Cell Wall Assembly in *Saccharomyces cerevisiae*. Microbiol Mol Biol Rev. 70(2):317–343. doi:10.1128/MMBR.00038-05.

Lesage G, Sdicu AM, Ménard P, Shapiro J, Hussein S, Bussey H. Analysis of β-1,3-Glucan Assembly in *Saccharomyces cerevisiae* Using a Synthetic Interaction Network and Altered Sensitivity to Caspofungin. *Genetics*. 2004;167(1):35-49. doi:10.1534/genetics.167.1.35

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform. Bioinformatics. 25(14):1754–1760. doi:10.1093/bioinformatics/btp324.

Matela AM, Siatkowski CW, Yan C, Thiagarajan S, Cooper VS. Student-led experimental evolution reveals novel biofilm regulators of adaptation to multiple niches. Published online June 6, 2025. doi:10.1101/2025.06.06.658356

Moresi NG, Geck RC, Skophammer R, Godin D, Students yEvo, Taylor MB, Dunham MJ. 2023. Caffeine-tolerant mutations selected through an at-home yeast experimental evolution teaching lab. MicroPubl Biol. 2023. doi:10.17912/micropub.biology.000749.

Olivier M, Hollstein M, Hainaut P. 2010. *TP53* Mutations in Human Cancers: Origins, Consequences, and Clinical Use. Cold Spring Harbor Perspectives in Biology. 2(1):a001008–a001008. doi:10.1101/cshperspect.a001008.

Park S, Kelly R, Kahn JN, Robles J, Hsu M-J, Register E, Li W, Vyas V, Fan H, Abruzzo G, et al. 2005. Specific Substitutions in the Echinocandin Target Fks1p Account for Reduced Susceptibility of Rare Laboratory and Clinical *Candida* sp. Isolates. Antimicrob Agents Chemother. 49(8):3264–3273. doi:10.1128/AAC.49.8.3264-3273.2005.

Perlin DS. 2007. Resistance to echinocandin-class antifungal drugs. Drug Resistance Updates. 10(3):121–130. doi:10.1016/j.drup.2007.04.002.

Prior IA, Lewis PD, Mattos C. 2012. A Comprehensive Survey of Ras Mutations in Cancer. Cancer Research. 72(10):2457–2467. doi:10.1158/0008-5472.CAN-11-2612.

Ratcliff WC, Raney A, Westreich S, Cotner S. 2014. A Novel Laboratory Activity for Teaching about the Evolution of Multicellularity. The American Biology Teacher. 76(2):81–87. doi:10.1525/abt.2014.76.2.3.

Reenan RA, Kolodner RD. 1992a. Characterization of insertion mutations in the *Saccharomyces cerevisiae MSH1* and *MSH2* genes: evidence for separate mitochondrial and nuclear functions. Genetics. 132(4):975–985. doi:10.1093/genetics/132.4.975.

Reenan RA, Kolodner RD. 1992b. Isolation and characterization of two *Saccharomyces cerevisiae* genes encoding homologs of the bacterial HexA and MutS mismatch repair proteins. Genetics. 132(4):963–973. doi:10.1093/genetics/132.4.963.

Robinson JT, Thorvaldsdóttir H, Wenger AM, Zehir A, Mesirov JP. Variant Review with the Integrative Genomics Viewer. *Cancer Research*. 2017;77(21):e31-e34. doi:10.1158/0008-5472.can-17-0337

Sivley RM, Dou X, Meiler J, Bush WS, Capra JA. 2018. Comprehensive Analysis of Constraint on the Spatial Distribution of Missense Variants in Human Protein Structures. The American Journal of Human Genetics. 102(3):415–426. doi:10.1016/j.ajhg.2018.01.017.

Smith JJ, Johnson WR, Lark AM, Mead LS, Wiser MJ, Pennock RT. 2016. An Avida-ED digital evolution curriculum for undergraduate biology. Evo Edu Outreach. 9(1):9. doi:10.1186/s12052-016-0060-0.

Stehr H, Jang S-HJ, Duarte JM, Wierling C, Lehrach H, Lappe M, Lange BM. 2011. The structural impact of cancer-associated missense mutations in oncogenes and tumor suppressors. Mol Cancer. 10(1):54. doi:10.1186/1476-4598-10-54.

Taylor MB, Skophammer R, Warwick AR, Geck RC, Boyer JM, yEvo Students, Walson M, Large CRL, Hickey AS-M, Rowley PA, et al. 2022. yEvo: experimental evolution in high school classrooms selects for novel mutations that impact clotrimazole resistance in *Saccharomyces cerevisiae*. Gresham D, editor. G3 Genes|Genomes|Genetics. 12(11):jkac246. doi:10.1093/g3journal/jkac246.

Taylor MB, Warwick AR, Skophammer R, Boyer JM, Geck RC, Gunkelman K, Walson M, Rowley PA, Dunham MJ. 2024. yEvo: A modular eukaryotic genetics and evolution research experience for high school students. Ecology and Evolution. 14(1):e10811. doi:10.1002/ece3.10811.

Thompson GR, Wiederhold NP, Vallor AC, Villareal NC, Lewis JS, Patterson TF. 2008. Development of Caspofungin Resistance following Prolonged Therapy for Invasive Candidiasis Secondary to *Candida glabrata* Infection. Antimicrob Agents Chemother. 52(10):3783–3785. doi:10.1128/AAC.00473-08.

Turner TN, Douville C, Kim D, Stenson PD, Cooper DN, Chakravarti A, Karchin R. 2015. Proteins linked to autosomal dominant and autosomal recessive disorders harbor characteristic rare missense mutation distribution patterns. Hum Mol Genet. 24(21):5995–6002. doi:10.1093/hmg/ddv309.

Win AK, Jenkins MA, Dowty JG, et al. Prevalence and Penetrance of Major Genes and Polygenes for Colorectal Cancer. *Cancer Epidemiology, Biomarkers & Prevention*. 2017;26(3):404-412. doi:10.1158/1055-9965.epi-16-0693

Figure S1: (A) When users select "View By Class", they will then be provided with dropdown menus for Instructor, Year, and Sample Name. Once Instructor and Year are selected, all samples will be the student groups from that particular classroom. (B) When users select "View By Condition", all yEvo conditions that have been used for classroom evolution experiments are available in the dropdown menu. Selecting any of these will filter for all yEvo data collected in that condition. (C) Some classrooms used alternative genetic backgrounds of *S. cerevisiae* in their evolution experiments. When this is the case, an additional dropdown menu option will appear, and users can select the desired background. (D) Users also have the option to upload their own CSV of variant data. Once they add their own instructor name and year, the new teacher will appear in the dropdown menu when selecting "View By Class".

Figure S2: Users can click and drag a box around sites on the chromosome plot to see genes in clustered regions.