Microbial consortia in salt marsh sediments are sequentially buried over millennia and genomic complementarity analysis indicates an important role in complex carbon decomposition

Joseph H. Vineis[1,2], Ashley N. Bulseco[1,3], Zoe G. Cardon[2], and Jennifer L. Bowen[1]

[1]Marine Science Center, Northeastern University, Nahant, Massachusetts, USA

[2]The Ecosystems Center, Marine Biological Laboratory, Woods Hole, Massachusetts, USA

[3]Department of Biological Sciences, University of New Hampshire, Durham, New Hampshire, USA

Abstract

Salt marsh sediments contain high levels of microbial diversity and functions that reflect the heterogeneity of available nutrients, dynamic hydrology, and layers of organic matter developed over millennia. However, the microbially mediated processes involved in the cycling of complex carbon are still largely undescribed. We used genome reconstruction, co-occurrence networks, and genome scale metabolic modeling to identify the functional capacity for organic matter decomposition and transformation among groups of co-occurring microbes from 5-240 cm of sediment in a *Spartina patens* salt marsh, representing over 3000 years of sediment accumulation. We identified four consortia with similar taxonomic and functional profiles, but distinctly different distributions within the sedimentary layers. Microbial members common to each of these consortia included novel members of Bathyarchaeia BA1, Desulfatiglandales, Chloroflexota, Caldatribacteriota (JS1), Planctomycetota, and WOR3. The collective metabolic potential of these core consortia contained the capacity to decompose complex carbon and aromatics through syntrophic interactions. The composition and

functional potential of these co-occurring members are most similar to communities

inhabiting sediment hundreds of meters beneath the surface of deep-sea sediments.

Our results suggest that the assembly of communities within the salt marsh sediment

are similarly governed by burial and nutrient limitation. The burial processes and

putative syntrophy identified here provide a mechanistic understanding of how microbial

life can persist under energy limited conditions and contribute to the transformation of

carbon within salt marsh sediment.

Keywords: *Spartina alterniflora*, salt marsh sediment, syntrophy, consortia, acetogens,

aromatic carbon, Wood-Ljungdahl**,** benzoyl-CoA

Introduction

Salt marshes are globally important to carbon storage and have the potential to

influence both the changing climate and the stability of coastal systems. Salt marsh

sediments contain between 400 – 6500 Tg of organic carbon (Duarte et al. 2013;

Mcleod et al. 2011) and store an additional 10.2 - 44.6 Mt annually, the equivalent of

0.5-1.0% of anthropogenic C emissions (Ouyang and Lee 2014). However, carbon

storage in salt marshes is declining and estimates indicate that over a recent nine-year

period, salt marsh habitat loss resulted in emission of 16.3 Tg of CO_2 and a reduction of

0.045 Tg of CO_2 burial per year (Campbell et al. 2022).

Rapidly increasing sea level rise and warming climate could destabilize current

stocks of organic carbon with unpredictable consequences (Fontaine et al. 2007; Spivak

et al. 2019). Microbial communities within salt marsh sediments are compositionally and

metabolically diverse and their interactions influence pools of soil organic carbon (SOC)

(Bulseco et al. 2019; Erb et al. 2024). However, there are important knowledge gaps in

our understanding of the metabolic capacity and distribution of microbial communities in

2this system often lead to the identification of previously unidentified carbon cycling

among novel taxa (Payne et al. 2025; Seitz et al. 2016; Vineis et al. 2023). Identifying

how the pools of carbon are influenced by microbial communities will enhance our

understanding of the fate of blue carbon stocks, how they are formed, and how they

change over time.

The decomposition of organic material is strongly influenced by plant

communities that blanket the surface of the marsh in nearly homogenous stands. The

regions of the marsh inhabited by plants are highly dynamic zones that contain hot

spots of microbial activity due to plant-microbial interactions and feedbacks that are

sensitive to seasonal and tidal influence (Kuzyakov and Blagodatskaya 2015). *Spartina*

patens is a highly productive salt marsh macrophyte with roots that release organic

carbon and aerate the sediment, thereby influencing carbon sequestration of salt

marshes through the "microbial carbon pump" (Liang et al. 2017; Spivak et al. 2019).

Live roots of *S. patens* are dominantly 0-10cm below the sediment surface (Gallagher

and Plumley 1979) and plant inputs could influence the SOC pool up to 100 cm (Liu et

al. 2017).

The patchy distribution of resources and dynamic processes in marsh sediments

present challenges for unraveling the underlying rules for community assembly and

metabolism. Identifying the distribution and putative functions of the microbial genomes

within the dynamic zone influenced by plants and tidal inputs remains one of the most

challenging but critical aspects to understanding the carbon and nutrient cycling in salt

marsh systems. While decomposition within the salt marsh is conducted primarily by

sulfate reducers (Howes et al. 1984), most of the alternative pathways for anaerobic

respiration, fermentation, and carbon fixation pathways have also been identified in

marsh sediments (Baker et al. 2015; Vineis et al. 2023; Bulseco et al. 2020) and we are

only beginning to understand how many of the carbon cycling processes are linked

among organisms and within individual genomes.

Below the rooting zone and beyond the influence of living plants, salt marshes

have accumulated vast stores of organic peat that date back, in some cases, millennia

(Kirwan et al. 2023). These older sediments are likely to contain microbes adapted to

survive in the deep, low energy environment where prior oxidation of organic material

has led to a decline in the bioavailability of organic carbon with sediment age and depth

(Arndt et al. 2013). Microbial communities below the influence of living plants therefore

might mirror other marine low energy sediment communities supported by complex

organic matter derived from compounds such as lignin, cellulose, and phenolic

polymers (Jørgensen and Boetius 2007). Decomposition of these compounds within

energy limited sediment is often dependent on syntrophic interactions among microbial

consortia (Peng et al. 2023) The specific mechanism of microbial decomposition has

important consequences for carbon stability because microbial turnover can result in the

production of carbon that is protected from further decomposition (Liang et al. 2017).

Therefore, identifying the capacity for microbial decomposition is important to

understanding the susceptibility of long-term carbon storage within salt marsh

sediments.

The physical and biogeochemical differences between the dynamic upper layers

of the sediment, the energy poor deeper layers, and the transition between the two

environments, are likely to support distinctly different microbial communities. We

hypothesize that communities within dynamic upper layers will contain genomes with

diverse functional potential. Additionally, we hypothesize that the energy limited deeper

regions will contain microbial consortia assembled to efficiently metabolize complex

carbon. We aim to identify the boundaries of both metabolically flexible genomes and

those relying on metabolic handoffs. We also consider the possibility that compositional

changes in these low energy sediments will mirror the burial processes observed in

marine environments that contain similar energetic limitations (Kirkpatrick et al. 2019;

Walsh et al. 2016).

To test these hypotheses, we evaluated the structure and functional potential of

microbial communities within 5-240 cm depth fractions from two *S. patens* salt marshes,

using a metagenomics and genome reconstruction. Initially, we evaluated the

phylogeny, distribution, and co-occurrence of reconstructed draft genomes. We also

characterized the potential genomic capacity to decompose organic carbon. Our initial

observations of co-occurrence and functional potential prompted us to employ genome-

scale metabolic models (GEMs) among co-occurring microbial genomes to identify

potential functional complementarity important to carbon cycling.

Methods

1. Sampling

Six sediment cores were taken using a Russian peat corer from the salt marsh platform

at two sites in Rowley MA, USA (42.759 N, 70.891 W) primarily vegetated by *Spartina*

patens. The sites are part of the Plum Island Ecosystem (PIE) Long-Term Ecological

Research site (LTER). Three cores were collected from the high marsh *Spartina patens*

platform at two adjacent creeks, West and Sweeney Creek (Fig. 1A). The two creeks

are approximately 2 km apart and cores collected within each creek were separated by

less than 50 meters. Cores were sampled up to the point of resistance, reaching a

maximum depth of 240 cm. Once extracted, subsections of the core were collected

approximately every 10 cm and a 2 cm sub-section was homogenized in a sterile 50 mL

falcon tube, flash frozen using liquid nitrogen and then stored at -80°C for nucleic acid

analysis. A detailed description of sample handling and analysis, including

biogeochemistry of the cores has been previously described (Bulseco 2018).

Figure 1. A) Map of the coring locations within the Plum Island Estuary Long Term

Ecological Research (PIE LTER) site and B) Details of core sampling and the

bioinformatic approach used to identify metabolic complementarity among co-occurring

MAGs within the sediment layers.

2. Metagenomic Sequencing

The input DNA for sequencing library construction was extracted from homogenized

sediment from each 2 cm thick fraction (0.25 g wet weight) using a Qiagen Power Soil

DNA extraction kit (Qiagen, Germantown, MD) according to the manufacturer's

recommendations. We evaluated the quality and the quantity of DNA using a Nanodrop

and Quant-IT DNA assay (Invitrogen, USA) respectively. Purified DNA was shipped

overnight to the Joint Genome Institute (JGI) on dry ice. At JGI, libraries were created

using paired-end (PE) library construction with an insert size of 250 bp. Each library was

uniquely barcoded and sequenced according to PE 2x151 chemistry on an Illumina

NovaSeq. Sequences were quality filtered using the default settings of BBTools v38.26

at JGI (https://jgi.doe.gov/data-and-tools/bbtools/). Quality filtered sequences of each

sample were independently assembled using SPAdes assembler 3.12.0 (Bankevich et

al. 2012).

3. MAG reconstruction, dereplication, phylogenomics and metagenomic read

recruitment

We used assemblies and quality filtered sequences generated by JGI/IMG for each of

the 55 samples to reconstruct metagenome-assembled genomes (MAGs) for each

independent sample (Fig. 2B). The assembled MAGs represent a population of cells.

Therefore, we refer to MAGs as populations or population genomes. Genome

reconstruction efforts can be improved by co-assembly of multiple samples in a gradient

or time series, especially when read counts are not sufficient to provide the minimum

amount of coverage necessary for assembly (Stewart et al. 2018). However, the

detection of redundant genome populations across samples requires that genome

reconstruction be carried out for each sample independently. Additionally, the number of

high-quality genomes recovered can be improved by reconstructing genomes from

individual samples compared to co-assembly of multiple samples (Olm et al. 2017).

We reconstructed draft genomes for each sample using CONCOCT (Alneberg et

al. 2014) as an automated guide for genome binning, followed by manual curation of

each CONCOCT MAG in Anvi'o, using contig tetranucleotide frequency, coverage,

taxonomic assignment, completion, and contamination to guide this process (Eren et al.

2015; 2020). In cases where the placement of contigs could not be resolved among a

group of similar MAGs, we used anvi-refine to provide increased resolution on this

subset of similar MAGs. This allowed for the separation of closely related MAGs and/or

the improvement of draft genome bins, as determined by Anvi'o real-time completion

and contamination scores.

Dereplication of the MAGs and selection of the representative MAG from the

dereplicated collections were conducted using the default settings for dRep (Olm et al.

2017). Default settings in dRep include dereplication of genomes with an average

nucleotide identity (ANI) of 0.95 and a minimum overlap of 10% between the two

genomes. dRep incorporates additional essential methods, including CheckM (Parks et

al. 2015) and FastANI (Jain et al. 2018). dRep uses completion and contamination

based on checkM, which were never at or above the 75% threshold for any of the

Patescibacteria. Patescibacteria MAGs were dereplicated separately without a checkM

quality score filter. We used the GTDB-Tk to assign taxonomy to each of the MAGs

recovered using default settings (Chaumeil et al. 2020).

To identify the phylogenomic relationships among the collection of dereplicated

MAGs, we first exported the amino acid sequence of the best hit for 71 single copy

genes (Bacteria_71). The amino acid sequences for these genes were aligned using

MUSCLE and the alignment was trimmed using trimAL v1.4 (Capella-Gutiérrez et al.

2009). We reconstructed the phylogenetic tree using the concatenated sequence

alignment according to anvi-gen-phylgenomic-tree which employs the FastTree tree

building algorithm (Price et al. 2009). The fasta file of single copy gene sequences,

alignment, tree, and additional details of phylogenetic reconstruction are available in this

github repository https://github.com/jvineis/DEEP-CORE-MANUSCRIPT.git

We mapped the metagenomic reads from each sample to the collection of

dereplicated MAGs using bowtie2 v2.5.3 (Langmead and Salzberg 2012). Unmapped

reads were filtered from the results and samtools v1.19.2 (Li et al. 2009) was used to

tabulate the number of reads recruited to each MAG. The relative abundance of each

MAG was estimated by dividing the number of reads mapped to the MAG by the

number of reads recruited to the non-redundant collection of MAGs.

4. Identifying co-occurring MAGs using network reconstruction

To identify groups of co-occurring genome populations that may be indicative of a

shared niche space and/or burial trajectory, we constructed a molecular ecological

network (MEN) using random matrix theory (RMT) according to the procedures

recommended by iNAP (Feng et al. 2022). RMT is more commonly used for functional

or phylogenetic marker genes but is broadly designed to identify co-occurrence patterns

among complex datasets that contain large numbers of variables (species occurrence,

microarray data, marker genes) (Gibson et al. 2013). The count matrix of reads

recruited to each of the 377 dereplicated MAGs was used as the input for RMT MEN

construction. Following log transformation of the count matrix, an adjacency matrix was

calculated according to Pearson correlation coefficients. To separate significant co-

occurrence from noise, RMT estimates the nearest neighbor spacing distribution

(NNSD) within the entire correlation matrix. A random matrix of NNSD values will follow

a Gaussian distribution and a non-random matrix will follow a Poisson distribution (Luo

et al. 2007). To identify the threshold where the correlation estimates transition from

Gaussian to Poisson, we ran iterations of successively smaller correlations starting at 1

with a step size of 0.01. A Chi-square test was used to evaluate transition from a

Poisson to a Gaussian distribution at each step and iNAP created a plot of resulting p-

values to evaluate each step. Visual inspection of the cutoff indicated that correlations

below 0.92 were more likely due to random noise (Fig. S1). All correlations below this

0.92 cutoff were set to zero and a network was constructed. RMT network properties

were calculated for the resulting filtered network, including R-square of the power-law

relationship. A network containing few MAGs with large numbers of connections with

other MAGs will have connectivity that follows a power-law distribution. If the

connectivity doesn't fit the power-law distribution, this indicates that there are many

MAGs with high connectivity to other MAGs (a very dense network). Module separation

and identification of module hubs were calculated according to a greedy modularity

optimization (Newman 2004). Within-module (Z) and among-module connectivity (P)

were used to classify each MAG as a network hub (Z > 2.5 and P > 0.62), module hub

(Z > 2.5 and P < 0.62), connector (Z < and P > 0.62), or peripheral (Z < 2.5 and P <

0.62) (Deng et al. 2012). Based on previous ecological interpretation of networks, the

peripheral, connector hub, module hub, and network hub might represent a continuum

from specialist to generalist in that order (Olesen et al. 2006). For our purposes of

identifying co-occurring genomes, we focused on the identification of modules, which

are interlinked subsets of MAGs, and connector hubs that could be important to the

stability of the different modules (Olesen et al. 2006).

5. Metabolic potential and distribution of MAGs within identified network

modules.

To estimate the differences in functional potential among the RMT modules, we

constructed a matrix composed of KEGG functional IDs obtained using anvi-estimate-

metabolism. MAGs that were included in modules containing fewer than four members

or not assigned to a module were also removed from the matrix. We constructed an

Atchinson distance matrix and built an NMDS ordination using the vegdist and

metaMDS functions respectively. We tested the influence of RMT module and

taxonomic classification at the phylum level on the community turnover using the envfit

function. All diversity analysis was conducted in R (Core Team 2024) with functions

contained in the vegan package (Oksanen et al. 2024). Relative abundance of MAGs

within each module were visually inspected to identify their distribution within the vertical

sediment profile.

6. Connector hub (Bathyarchaeia BA1) subnetwork genome scale metabolic networks

Visualization of the complete RMT network (described above) using Cytoscape (Shannon et al. 2003) led to the identification of four modules with similar taxonomy, functional profiles, and high relative abundance within deeper sediment layers. The MAGs identified as connector hubs common to all four modules, Bathyarchaeia BA1, was used to identify a subnetwork of MAGs that were positively connected and therefore contained the potential for metabolic complementarity. Focusing on MAGs in this subnetwork, we estimated the metabolic potential for key pathways related to decomposition of organic compounds, aromatics, acetate, hydrogenases, and other pathways important to carbon cycling within low energy sediment environments using METABOLIC according to the default parameters (Zhou et al. 2022). We reconstructed genome-scale metabolic networks (GSMNs) using gapseq (Zimmermann et al. 2021) and miscoto (Frioux et al. 2018) was employed to screen GSMNs for key members of minimal communities required to produce the metabolites of the entire community. Analysis of GSMNs was employed using Metag2Metabo (M2M) (Belcour et al. 2020). This approach identifies putative metabolites produced by the co-occurring members of the network module, the key microbial taxa required to produce these metabolites, and interchangeable (redundant) members of the network module.

Results

1. Diverse MAG populations were broadly distributed within salt marsh

sedimentary layers and both sites

We generated 645.9 Gbp sequences from 55 samples collected up to ~240 cm of

salt marsh sediment from six cores (Fig. 1B). The manual genome binning and

refinement effort for each of the 55 samples recovered a total of 2,346 MAGs.

Dereplication resulted in a collection of 377 MAGs that included 44 phyla, with novel

MAGs identified at the class, family, order and genus level (Table S1). The dereplicated

MAGs were an average of 87% of expected single copy genes (completion) and 2.7%

of these genes found more than once (contaminated) (Fig. 2, Table S1).

Most of the MAGs recovered represented novel families within Bacteria and

Archaea and were well distributed within the layers of sediment (Fig. 2) and

dereplication revealed that many of the MAGs were independently recovered from

multiple metagenomic samples (Fig. 3). Phyla containing the largest number of MAGs

included Chloroflexota, Planctomycetota, Alphaproteobacteria, Gammaproteobacteria,

and Desulfobacterota. A large fraction of the Chloroflexota MAGs were

Dehalococcoidia, which were primarily detected in sediment layers below 70 cm and

Anaerolineae which had a shallower distribution (Fig. 2). Nearly all 41 MAGs in the

Planctomycetota were primarily located below 40 cm. MAGs classified as

Desulfobacterota were primarily found in shallow samples, but a distinct clade was

detected primarily within shallow sediment. The 39 Proteobacteria MAGs were split

evenly between Alphaproteobacteria with MAGs that span the entire depth of sampling

and Gammaproteobacteria that were mostly detected in sediment above 40 cm (Fig. 2).

Similarly, the 28 MAGs within the phylum Bacteroidota were partitioned according to

depth, although not as strictly as Alpha and Gammaproteobacteria. Within the

Bacteroidota, families Vadin HA17, UBA10428, and UBA5072 were generally found

below 30-40 cm while Ignavibacteriaceae were found at 10 cm.

Figure 2. The tree at the center of the display shows the phylogenetic relationship
among all MAGs. The innermost ring is a histogram the single copy gene completion
estimate (range = 0-100%), followed by redundancy (range = 0-10%), total length of the
genome (range = 0-7.6Mbp). The "Depth proportion" ring shows the proportion of the
total reads mapped to a MAG from each depth. The outermost ring shows the phylum

The number of MAGs contained in each dereplicated cluster ranged from 1 to 30 (Fig. 3). Some examples of genome populations represented by a large cluster of dereplicated genomes include 25 independently reconstructed *Ca. Aminicenans* that were reduced to two unique MAGs, 63 Actinobacteriota reduced to 4 MAGs, 52 *Bipolaricaulia* reduced to 6 MAGs, 85 WOR3-B reduced to 11 MAGs, and 24 TA06 reduced to 1 unique MAG (Fig. 3). Many of the MAGs were recovered independently from more than one depth and often from different cores at locations adjacent to different creeks (Fig. 3). In some instances, MAGs from the same dereplicated cluster were independently reconstructed from the top 10-20 cm and below 200 cm. For example, a dereplicated WOR3 MAG contained nearly 30 independently assembled MAGs from samples that spanned the entire 10-240cm vertical profile (Fig. 3). The Desulfobacterota, Bacteroidota, Actinobacteriota, Planctomycetota, and Chloroflexota each contained several MAGs that were recovered from all depths. Sparsely detected Proteobacteria, Gemmatimonadota, and Asgardarchaeota were recovered predominantly from shallow sediment layers, 90cm and above (Fig. 3).

The proportion of metagenomic sequences recruited (mapped) to our collection of nonredundant MAGs was generally greater in sediment samples collected below 40 cm (Fig. S2). Read recruitment below 40 cm was at least 35% and reached nearly 60% in most of the samples below 90 cm. Conversely, metagenomic sequences from sediment samples collected from above 20 cm recruited less than 20% of sequences per sample (Fig. S2) and the number of MAGs recovered was generally lower.

Figure 3. Summary of redundant genome recovery per sample. The line plot at the top
of the figure (dark gray and black) shows how many genomes were grouped into the
dereplicated genome cluster. The second layer from the top represents the phylum level
classification for the representative MAG in the cluster. The heat map layers show the
proportion of samples where a MAG in the dereplicated genome cluster was recovered.
The key indicating how many of the same MAGs (identified by dRep) were collected per
depth is shown on the right. The MAGs are organized according to the phylogenetic
reconstruction outlined in the methods section. The phylum color key is shown at the
bottom of the figure and the gray triangles above the phylum color align with the
heatmap.

2. Co-occurring MAGs

2.1 Strong positive correlations exist among groups of functionally similar MAGs in the

deeper sediment layers

Based on MAG relative abundance, we identified a connected network containing

1547 positive connections (edges) among 226 of the 377 MAGs and zero negative

edges (Fig. 4A). Most of MAGs were peripheral nodes in the network, but one MAG

(Dehalococcoidales – UBA5760) was identified as a module hub and 15 MAGs were

classified as connector hubs (Fig. S3). Among the connector hubs were two

Bathyarchaeia classified as BA1, two Dehalococcoidales, Desulfatiglandales, and GIF9

MAGs. The R^2 of the power-law was 0.97, indicating a scale-free network where most

MAGs in the network have few connections. We chose to focus our functional

analysis on the top seven modules identified in the network because the remaining 11

modules contained fewer than 5 MAGs (Table S1).

Modules 1 and 2 contained the largest number of MAGs with a high degree of

connectivity within and between the two modules (Fig. 4A). Modules 1 and 2 were also

diverse, containing MAGs from 14 and 23 phyla respectively. MAGs within module 7

contained many positive links with modules 1 and 2 and the functional potential among

MAGs within these three linked modules was very similar (Fig. 4B, Fig. S4). However,

modules 1 and 2 contained larger dispersion in the ordination than module 7 due to the

presence of Patescibacteria, which were functionally distinct from other MAGs, largely

due to their limited functional potential. Most of the Patescibacteria contained fewer than

800 genes (Table S1). A relatively large proportion of MAGs in modules 1, 2, and 7 (>

25%) contained pathways for complex carbon decomposition, fermentation, amino acid

metabolism, and several C1 pathways (Fig. S4). However, pathways for carbon fixation,

nitrogen and sulfur reduction, and oxidative phosphorylation were identified in a

relatively lower proportion relative to modules 3 and 6 (Fig. S4). Connectivity among

modules 4 and 5 was sparse in comparison to the large number of positive connections

among modules 1, 2, and 7. However, the overall functional potential was similar among

all five modules (1,2,4,5,7) (Fig. 4B, S4).

Figure 4. A.) The random matrix theory derived co-occurrence network, based on MAG
relative abundance. The point size (genome size), color (phylum), and shape (module)
are used to describe each MAG in the network. All connections are significant and
positive. Ellipses are used to clarify the modules. B) The NMDS ordination of MAGs
included in the network based on KEGG functional potential. Both panels show the
same ordination, with the left panel colors indicating the phylum classification and right,
the module assignment. The key adjacent to panel A applies to the colors used for taxa
and modules shown in panel B.

2.2 Modules containing widespread and abundant MAGs in sediment layers.

The relative abundance of MAGs at the phylum level within a module was

consistent across sediment layers (Fig. 5). For example, the top four most abundant

phyla in module 1 included Chloroflexota, Planctomycetota, Bacteroidota, and

Actinobacteriota. In module 2, the cumulative relative abundance of Choloroflexota,

Planctomycetota, TA06, and WOR3-B were also among the most abundant phyla within

this module and were found in similar relative proportions in the sediment layers.

Despite the diversity of MAGs assigned to a particular phylum within a module, a

small number of MAGs were often the most abundant at all sediment depths. For

example, 12 of the 70 MAGs in module 1 represented the majority of reads recruited

(Fig. S5). The abundant MAGs in the module were distributed across 6 of the 13 phyla

in the module. Within module 1, 5 Chloroflexota MAGs were among the most abundant

(Fig. S5). Other phyla within the module were low diversity, such as TA06 in module 2,

where a single MAG was the sole contributor to the phylum level abundance patterns.

Although often detected in the entire depth profile, MAGs within modules 1, 2, 4

and 7 were especially abundant below 20 cm (Fig. 5, S4). Modules 1 and 2 contained

several Chloroflexota MAGs that collectively recruited at least 15% of the mapped

reads. Average coverage of these MAGs was over 50X in the majority of samples below

40 cm. Module 2 contained two of the most abundant MAGs below 40 cm, TA06 and

Bipolaricaulota, which recruited nearly 15% of mapped reads in many samples below

100 cm. The peak relative abundance of MAGs in modules 1, 2, 5, and 7 were distinct,

with module 1 peaking in abundance at 70-110 cm, module 2 peaking in all samples

below 100 cm, module 5 peaking at 30 - 40 cm, and module 7 peaking at 150 – 250 cm

(Fig S4).

Figure 5. Bubble plots for the cumulative phylum level relative abundance of MAGs contained in each of the network modules. Relative abundance (size of the bubble) indicates the percentage of reads mapped to the MAGs glommed at the phylum level relative to the number of reads mapped to the sample. The colors for each phylum are congruent across all four panels. The box-whisker relative abundance plots nested within each box describe the relative abundance of MAGs in the module at each depth.

2.3 Unconnected MAGs are dominant in the upper 40 cm and potentially have a niche in the deepest sediment

In addition to MAGs that were associated with the modules described above, there were numerous unconnected MAGs that were predominantly located in sediments at or above 40 cm (Fig. S4). Proteobacteria (Pseudolabrys and Methyloceanibacter) and Crenarchaeota (Nitrososphaeraceae and Bathyarchaeia BA1) MAGs were among the most abundant unconnected MAGs. However, we also observed MAGs within many sediment layers that were abundant both in the upper 40 cm and below 200 cm within the core collected from West Creek (WE3) (Fig. S6). Although they maintained a patchy distribution in the surface, the relative abundance was consistent at 200-240 cm (Fig. S6). *Methyloceanibacter* MAGs identified in this "bimodal" group are putative bacterial methylotrophs (Vekeman et al. 2016), but none of the mags contained the methane monooxygenase or co-methyltransferase genes. Bimodal Nitrososphaeraceae MAGs contained ammonia oxidation potential (*amo*) and the *Pseudolabrys* contained sulfur oxidation (*sox*), sulfur reduction (*dsr, apr, sat*), and nitrate reduction potential (*narG*) (Fig. S6). These functions suggest a mix of aerobic and anaerobic respiration potential within the deepest layers sampled in the West creek sediment core.

2.4 Bathyarchaeia BA1 are potentially important facilitators of abundant deep sediment communities

Because of the potentially important ecological role of Bathyarchaeia BA1 MAGs in the network as connector hubs (Fig. S3) and global significance of this family to carbon cycling (Hou et al. 2023), we focused on these MAGs and the connections to other MAGs in the network to understand the potential relevance to carbon cycling. Our network analysis identified 2 separate Bathyarchaeia BA1 MAGs that were connector hubs in the RMT network (Fig. S3). To explore the potential role of these Bathyarchaeia BA1 we created a subnetwork of MAGs that are directly connected to the Bathyarchaeia MAGs, resulting in a total of 33 MAGs in the subnetwork (Fig 5A). Among the 12 different orders contained in the subnetwork were members of Bipolaricaulia, GIF9, WOR3, Desulfatiglandales, and Caldatribacteriota and each of the four modules (1, 2, 5, and 7) were represented (Fig. 6A). The relative abundance of members at the phylum level in this subnetwork was also consistent across many of the samples (Fig. 6B).

3. Putative functional complementarity among Bathyarchaeia subnetwork MAGs suggests potential decomposition of hydrocarbon and aromatics

3.1 Consistent relative abundance among members of the Bathyarchaeia subnetwork

Within the Bathyarchaeia subnetwork, GIF9 MAGs were the most abundant, totaling 8 MAGs, all of which are putative acetogens based on their functional profile. Six GIF9 MAGs were located in the same module (module 2), indicating a close ecological connection and yet they were not identified as functionally redundant members of the community according to the M2M analysis. The collective GIF9 MAG

relative abundance reached 15% in many of the samples below 20 cm, but like all

MAGs in the subnetwork, they were not excluded from the upper 20 cm (Fig. 6B).

Desulfatiglandales, Dehalococcoidales, and Planctomycetota were also among the

most abundant MAGs in the subnetwork and occurred at similar cumulative relative

abundances. The high correlation among these MAGs detected in the network is

reflected in the consistent relative abundance of the group members within a core and

generally across all six cores (Fig. 6B).

3.2 Bathyarchaeia BA1 subnetwork MAGs use cooperative metabolism to degrade

aromatics

M2M identified the putative functions of each community member and the added

value metabolites produced by minimal communities. An added value metabolite is one

produced through complementary metabolism among the microbes in a minimal

community, and a minimal community is defined as a cohort of MAGs required to

produce the entire set of metabolites in the collection of genomes. We found 82

metabolites that were producible by each of the 33 MAGs in the subnetwork (core

functions) and 1098 total metabolites could be produced by the individual organisms all

together (collective functions). The added value of cooperation within the minimal

communities required 22 bacteria and 20 essential symbionts to produce an additional

228 compounds. These 228 compounds were only possible when all members of a

minimal community had the potential to exchange metabolites. Redundant members of

the community that were identified in at least one, but not all minimal communities

included Dehalococcoidia – RBG-16-60-22, Caldatribacteriota JS1, Dehalococcoidia

AB-539-J10, and Dehalococcoidia UBA5760.

Among the 228 compounds that could be formed solely if organisms exchanged

compounds, key metabolites involved in anaerobic decomposition of aromatic

compounds were identified and the presence of the genes conferring capacity for these

putative functions were confirmed by METABOLIC. Benzoyl-CoA reductase (*bcr*) was

identified in the genomes of three Desulfatiglandales, a novel order of Cloroflexota, and

Bathyarchaeia. Genes to produce benzoyl-CoA from phenol (*ubiX*/*bsdC*) occurred in

two of the Desulfatiflandales, the Bathyarchaeia and Bacteroidales MAGs (Fig. 6C).

Phenylacetyl-CoA was also identified as a metabolite putatively produced through

functional complementarity within the subnetwork.

3.3 Genomic potential for acetogenesis and metabolic versatility is common among the

Bathyarchaeia BA1 subnetwork

Genes for heterotrophic and autotrophic acetogenesis, primarily through the

Wood-Ljungdahl (WL) pathway, were detected in 25 of the 33 genomes in the

Bathyarchaeia B26-1 subnetwork (Fig. 6C). These genes included acetyl-CoA

dehydrogenase (*acdA*), acetate kinase (*ack*), and phosphate acetyltransferase (*pta*).

acdA was the most commonly detected gene indicating acetogenic metabolism, and *acs*

(Acetyl-CoA synthetase) was commonly detected within the same MAG (Table S2). The

co-occurrence of genes encoding a putative WL pathway within most genomes and a

formate oxidation gene (*fae*) required to convert CO_2 into formate provides additional

evidence for synthesis of acetate from H_2 and CO_2. The genomic potential for the

production of acetate enabled by carbon fixation via the WL pathway was observed in

Desulfatiglandales, Bathyarchaeia, Dehalococcoidales, Chloroflexota, GIF9,

Bipolaricaulia, and Planctomycetota MAGs in the subnetwork (Fig. 6C). A WL pathway

was not detected in Caldatribacteriota, WOR3, and Bacteroidales. Instead, these MAGs

might produce acetate through fermentation of pyruvate or lactate. Most acetogens in

the network contain genes indicative of a flexible metabolic capacity through autotrophic

and/or fermentative acetate production (Fig. 6C).

We detected several additional pathways potentially related to energy

conservation among the members of the subnetwork. The presence of methane

monooxygenase regulatory protein (*mmoBD*) in Bipolaricaulia and Desulfatiglandales

(Fig. 6C) suggests methane oxidation potential. Additionally, aerobic carbon monoxide

dehydrogenase *coxSM* was detected in members of Bipolaricaulia, Caldatribacteriota,

Bathyarchaeia TCS64, and Desulfatiglandales. Additional pathways include selenate or

arsenate reduction potential in most of the subnetwork MAGs and one of the GIF9

MAGs contained genes for both pathways.

Figure 6. A) The submodule of tightly correlated MAGs with connectivity among the four subsurface modules. The shape in the network indicates module origin of each MAG, the size represents the number of genes in the MAG, and the color is used for taxonomic classification. Gray lines connecting the MAGs represent significant positive correlations. B.) Bubble plot of relative abundance for MAG phylum level relative abundance within the vertical profile of the sediment. The color of the bubble is based on the phylum classification and matches the color in panel A. C) Select functional annotation for all MAGs in the network. Functional potential of each MAG in the network is described as presence (black bar) or absence (gray bar) for each function. Major functional categories are shown on the left of the panel, and the taxonomic classification of each MAG is shown at the top. MAGs are organized according to presence-absence of the genes in the display and the hierarchical clustering tree is displayed at the bottom of the panel.

Discussion

Salt marsh sediments contain diverse microbial communities with vast numbers of uncultivated microbial organisms that are important to carbon storage and biogeochemical cycling. Within the complex microbial communities inhabiting these sediments, we identified metabolic complementarity important to cycling pools of aromatic carbon among co-occurring groups of microbes. Bathyarchaeia BA1 was a key member of each group of MAGs that were uniquely distributed within the sediment profile, and they were primarily located below the dynamic upper layers of the sediment. Another key finding was that similar genome populations exist throughout the entire depth profile of multiple cores, spanning nearly 3000 years of sediment accumulation (Bulseco 2018). This pattern may be the result of a burial process followed by environmental selection and slow growth rates. Conversely, the upper layers of sediment contain additional diverse MAGs with greater functional potential representing the canonical pathways commonly associated with salt marsh microbial communities (Baker et al. 2015; Tobias and Neubauer 2019). Our results are important for

understanding how the pools of organic carbon are shaped by microbial communities

within the less well characterized subsurface because we identify which microbes are

co-located, what forms of carbon may be made accessible due to metabolic

complementarity, and the underlying biogeochemical processes that preserve these

relationships.

1. Potential metabolic flexibility is a key feature among MAGs in the zone of plant

and tidal influence.

Plant roots and rhizomes along with bioturbating organism and other structural features

can increase permeability in the upper layers of the sediment (Guimond et al. 2020;

Moffett et al. 2012; Xiao et al. 2019) and these dynamic conditions potentially lead to

observed high diversity of microbial communities in the upper sediment layers. Redox

conditions and carbon availability can change rapidly in the upper sediment layers,

creating conditions where metabolic flexibility would be favored. As a result, there is a

decreased need for metabolic handoffs among members of the community.

Most of the MAGs that were unconnected in our network analysis were localized

in the dynamic upper layers of the sediment and did not occur below 90 cm. The

unconnected MAGs contained striking genomic versatility including the capacity to

utilize multiple electron acceptors for oxidation of diverse carbon sources, conduct

fermentation, and fix inorganic carbon. Many of the unconnected Proteobacteria and

Desulfobacterota MAGs contain the genomic potential for each of these metabolic

pathways in a single organism which was observed previously in similar sediments

(Vineis et al. 2023). This metabolic flexibility allows each of the members of the

community to inhabit a variety of conditions and this strategy makes sense given the

dynamic nature of these sediments. Members of these communities contain canonical

pathways that are commonly associated with carbon cycling in salt marsh sediments

including fermentation, denitrification, and sulfate reduction (Baker et al. 2015).

2. Organic carbon decomposition in deeper sediment layers is potentially

facilitated by co-occurring community members with complimentary metabolic

pathways

The consistency in relative abundance among members within the core

Bathyarchaeia subnetwork is a potential result of metabolic handoffs. Of particular

significance is the detection of benzoyl-CoA as a metabolite that could only be produced

by Bathyarchaeia, Dehalococcoidia, and Desulfatiglandales through metabolic handoffs

in the community. Benzoyl-CoA is a known intermediate involved in the anaerobic

decomposition of aromatic compounds toluene, phenol, ethylbenzene and benzoate

(Carmona et al. 2009; Porter and Young 2014). Under anaerobic conditions, reduction

of benzoyl-CoA leads to the production of acetyl-CoA which can enter into central

metabolic pathways for growth and energy conservation (Porter and Young 2014).

Phenylacetyl-CoA was also putatively produced through metabolic complementarity and

is similar to benzoyl-CoA, leading to the potential production acetyl-CoA/succinyl-CoA.

These metabolic pathways, central to aromatic decomposition are only possible due to

the potential metabolic handoffs among the subnetwork members. Although it is

possible that other combinations of MAGs could lead to the decomposition of aromatics,

co-occurrence provides additional support that handoffs among the genome populations

identified here are likely to occur in the environment.

Syntrophic interactions between a bacterial fermenter and CO_2 reduction are

often cited as important to the decomposition of aromatic carbon (Dai et al. 2024; Qiu et

al. 2006). The collective functional profile of the Bathyarchaeia subnetwork suggests

that the initial fermentation of complex organic carbon is conducted by members of the

Planctomycetota, WOR3, JS1 and others, followed by secondary decomposition of

aromatics by Bathyarchaeia, Deslufatiglandales, Chloroflexota. The products of

secondary fermentation, including H_2 and formate, usually used by methanogens, are

instead potentially being used by a diverse group of acetogens. Acetogens grow on

organic materials derived from cellulose, lignin, alcohols, hemicellulose, and chitin

within energy depleted sediments (Lever 2012; Yu et al. 2018), and there is evidence for

the use of lignin as an energy source in Bathyarchaeota (Yu et al. 2018).

These results suggest a central mechanism for carbon cycling among closely

coupled consortia within the sediment layers of salt marsh sediment. While this process

may be very slow, we have evidence from amplicon analysis of cDNA, that these

communities are active (Bulseco 2018). Conversion of these aromatics into microbial

biomass suggests that there may be a relevant cycling and/or recycling of carbon

stabilized by microbial activity even within these energy depleted sediments.

3. Burial as a mechanism for aggregating putative consortia

Burial as a mechanism for co-occurrence patterns has been suggested in energy

starved deep marine sediments where selection on the microbial community is thought

to result in low diversity (Kirkpatrick et al. 2019; Walsh et al. 2016). Previous studies of

microbial turnover within salt marshes described seasonal and transect community

compositional changes (Tebbe et al. 2022; Yao et al. 2019) or succession along a

chronosequence (Dini-Andreote et al. 2014; Wang et al. 2017). However, the

composition and stability of microbial communities within salt marsh sediments remains

relatively uncharacterized at the depths examine here.

Our study of the vertical sediment profile recovered two pieces of evidence that

suggests burial is a likely mechanism for the distribution of closely associated consortia.

The first observed pattern suggestive of burial was the recovery of MAGs assembled

independently from multiple depths in the same core that were dereplicated into a single

genome population. One of the strongest examples of this was observed in the

candidate phylum TA06, where a single population was recovered from 24 of the 55

samples, including every depth fraction except 10 cm. For dereplication to occur at this

scale (> 95% identity among MAGs in the dereplicated cluster), genome populations

would have experienced limited evolution over the nearly 3000 yrs of separation that

exists between the surface and bottom of the core. The second observed pattern was

the consistent relative abundance of MAGs within each module, relative to each other at

the phylum level (Fig. 5). The consistent relative abundance patterns among MAGs

within a module was observed both within the vertical sampling of an individual core and

across multiple cores. This is not to say that the relative abundance of members within

a module are the same at every depth across all cores. However, when the MAGs were

identified in a sample, they generally appear in similar ratios. The ratios of microbial

taxa might reflect the proportions required to support syntrophic interactions.

Similar compositions of sediment communities in marine systems have been

recovered from geographically distant communities (although at high taxonomic levels)

and were presumed to be subseafloor specialists (Inagaki et al. 2006; Petro et al. 2017;

2019). However, our data suggest that the same populations were buried within the

sediment, essentially preserving the population through high selection coupled with slow

growth.

The low porosity, small particle size, and low circulation observed in salt marsh

sediments on the East Coast of the US (Guimond and Tamborski 2021) provide

additional support to co-occurrence shaped by burial that we observed in our genomic

data. The turnover of water within the sediment can require 30-90 years (Guimond and

Tamborski 2021), depending on the marsh system, indicating that water flow from

groundwater or surface sources may be limited. These physical characteristics present

significant barriers to the dispersal of subsurface communities, leaving burial as a likely

mechanism. One caveat to the low porosity and water flow are the observations in the

deepest sediment layers collected from West Creek where we observed members of the

community that require oxygen or terminal electron acceptors commonly associated

with anaerobic shallow sediments. We speculate that the groundwater may potentially

"well up" from below the bedrock or sand located beneath the salt marsh platform at this

site.

Conclusion

The metabolic connections identified among correlated members of the microbial

community provide key insights into how carbon is cycled by microbial communities

within salt marsh sediments. Additionally, these results provide evidence for the

mechanisms important to understanding the assembly and distribution of these

communities. Our analysis identifies syntrophy among members of a putative core

consortia that provides a potential mechanistic understanding of organic matter

decomposition. The process begins with decomposers of complex carbon, followed by

aromatic decomposition, coupled with acetogenic energy conservation and finally

oxidation of acetate through arsenate or selenate reduction. These findings represent a

significant step forward in our understanding of how microbial communities survive

within these energy limited systems and contribute to the stability of soil organic carbon

in this blue carbon system. The persistent conversion of soil organic carbon by the

microbial community indicates that the pools of carbon potentially originating from plant

material undergo decomposition and transformation even under energy depleted

conditions.

Data Availability

The data described here are described in the U.S. Department of Energy (DOE)

Facilities Integrating Collaborations for User Science (FICUS) grant titled "Combining

high resolution organic matter characterization and microbial meta-omics to assess the

effects of nutrient loading on salt marsh carbon sequestration" (proposal ID = 503576,

https://genome.jgi.doe.gov/portal/Comhiguestration/Comhiguestration.info.html).

Supplemental data can be accessed using this link to a publicly available github page

https://github.com/jvineis/DEEP-CORE-MANUSCRIPT.git

Acknowledgements

Samples were collected from the Plum Island Ecosystems LTER, which is

supported by NSF (OCE 2224608) and from the NSF funded TIDE project (DEB

1902712). Sequencing was conducted at the Joint Genome Institute through a funded

Facilities Integrating Collaborations for User Science (FICUS) grant (proposal ID

503576). Additional funding for this project was provided by an NSF PRFB to ANB

(1907285), Simons Foundation (Grant #1247132 to ZGC, and Grant #01248016 to

JLB), and Department of Energy (DOE DE-SC0024270).

References

Alneberg, Johannes, Brynjar Smári Bjarnason, Ino De Bruijn, et al. 2014. "Binning
Metagenomic Contigs by Coverage and Composition." *Nature Methods* 11 (11):
1144–46. https://doi.org/10.1038/nmeth.3103.

Arndt, Sandra, B.B. Jørgensen, D.E. LaRowe, J.J. Middelburg, R.D. Pancost, and P.
Regnier. 2013. "Quantifying the Degradation of Organic Matter in Marine
Sediments: A Review and Synthesis." *Earth-Science Reviews* 123 (August): 53–
86. https://doi.org/10.1016/j.earscirev.2013.02.008.

Baker, Brett J, Cassandre Sara Lazar, Andreas P Teske, and Gregory J Dick. 2015.
"Genomic Resolution of Linkages in Carbon, Nitrogen, and Sulfur Cycling among
Widespread Estuary Sediment Bacteria." *Microbiome* 3 (1): 14.
https://doi.org/10.1186/s40168-015-0077-6.

Bankevich, Anton, Sergey Nurk, Dmitry Antipov, et al. 2012. "SPAdes: A New Genome
Assembly Algorithm and Its Applications to Single-Cell Sequencing." *Journal of
Computational Biology* 19 (5): 455–77. https://doi.org/10.1089/cmb.2012.0021.

Belcour, Arnaud, Clémence Frioux, Méziane Aite, Anthony Bretaudeau, Falk Hildebrand,
and Anne Siegel. 2020. "Metage2Metabo, Microbiota-Scale Metabolic
Complementarity for the Identification of Key Species." *eLife* 9 (December):
e61968. https://doi.org/10.7554/eLife.61968.

Bulseco. 2018. "Nitrate as an Electron Acceptor in Microbial Decomposition of Salt
Marsh Sediment Organic Matter and Implications for Carbon Storage."
Northeastern University. https://doi.org/10.17760/D20293211.

Bulseco, Ashley N., Anne E. Giblin, Jane Tucker, et al. 2019. "Nitrate Addition
Stimulates Microbial Decomposition of Organic Matter in Salt Marsh Sediments."
Global Change Biology 25 (10): 3224–41. https://doi.org/10.1111/gcb.14726.

Bulseco, Ashley N., Joseph H. Vineis, Anna E. Murphy, et al. 2020. "Metagenomics
Coupled with Biogeochemical Rates Measurements Provide Evidence That
Nitrate Addition Stimulates Respiration in Salt Marsh Sediments." *Limnology and
Oceanography* 65 (S1). https://doi.org/10.1002/lno.11326.

Campbell, Anthony D., Lola Fatoyinbo, Liza Goldberg, and David Lagomasino. 2022.
"Global Hotspots of Salt Marsh Change and Carbon Emissions." *Nature* 612
(7941): 701–6. https://doi.org/10.1038/s41586-022-05355-z.

Capella-Gutiérrez, Salvador, José M. Silla-Martínez, and Toni Gabaldón. 2009. "trimAl:
A Tool for Automated Alignment Trimming in Large-Scale Phylogenetic Analyses."
Bioinformatics 25 (15): 1972–73. https://doi.org/10.1093/bioinformatics/btp348.

Carmona, Manuel, María Teresa Zamarro, Blas Blázquez, et al. 2009. "Anaerobic
Catabolism of Aromatic Compounds: A Genetic and Genomic View." *Microbiology*
and Molecular Biology Reviews 73 (1): 71–133.
https://doi.org/10.1128/MMBR.00021-08.

Chaumeil, Pierre-Alain, Aaron J Mussig, Philip Hugenholtz, and Donovan H Parks.
2020. "GTDB-Tk: A Toolkit to Classify Genomes with the Genome Taxonomy
Database." *Bioinformatics* 36 (6): 1925–27.
https://doi.org/10.1093/bioinformatics/btz848.

Core Team, R. 2024. *R: A Language and Environment for Statistical Computing*.
Released. https://www.R-project.org/.

Dai, Shixiang, Falk Harnisch, Micjel Chávez Morejón, Nina Sophie Keller, Benjamin
Korth, and Carsten Vogt. 2024. "Microbial Electricity-Driven Anaerobic Phenol
Degradation in Bioelectrochemical Systems." *Environmental Science and*
Ecotechnology 17 (January): 100307. https://doi.org/10.1016/j.ese.2023.100307.

Deng, Ye, Yi-Huei Jiang, Yunfeng Yang, Zhili He, Feng Luo, and Jizhong Zhou. 2012.
"Molecular Ecological Network Analyses." *BMC Bioinformatics* 13 (1): 113.
https://doi.org/10.1186/1471-2105-13-113.

Dini-Andreote, Francisco, Michele De Cássia Pereira E Silva, Xavier Triadó-Margarit,
Emilio O Casamayor, Jan Dirk Van Elsas, and Joana Falcão Salles. 2014.
"Dynamics of Bacterial Community Succession in a Salt Marsh Chronosequence:
Evidences for Temporal Niche Partitioning." *The ISME Journal* 8 (10): 1989–
2001. https://doi.org/10.1038/ismej.2014.54.

Duarte, Carlos M., Iñigo J. Losada, Iris E. Hendriks, Inés Mazarrasa, and Núria Marbà.
2013. "The Role of Coastal Plant Communities for Climate Change Mitigation
and Adaptation." *Nature Climate Change* 3 (11): 961–68.
https://doi.org/10.1038/nclimate1970.

Erb, Hailey, Ashley Keiser, and Kristen M DeAngelis. 2024. "Salt Marsh Soil Organic
Carbon Is Regulated by Drivers of Microbial Activity." *Sustainable Microbiology* 1
(1): qvae009. https://doi.org/10.1093/sumbio/qvae009.

Eren, A. Murat, Özcan C. Esen, Christopher Quince, et al. 2015. "Anvi'o: An Advanced
Analysis and Visualization Platform for 'omics Data." *PeerJ* 3: e1319.

Eren, A. Murat, Evan Kiefl, Alon Shaiber, et al. 2020. "Community-Led, Integrated,
Reproducible Multi-Omics with Anvi'o." *Nature Microbiology* 6 (1): 3–6.
https://doi.org/10.1038/s41564-020-00834-3.

Feng, Kai, Xi Peng, Zheng Zhang, et al. 2022. "iNAP: An Integrated Network Analysis
Pipeline for Microbiome Studies." *iMeta* 1 (2): e13.
https://doi.org/10.1002/imt2.13.

Fontaine, Sébastien, Sébastien Barot, Pierre Barré, Nadia Bdioui, Bruno Mary, and
Cornelia Rumpel. 2007. "Stability of Organic Carbon in Deep Soil Layers
Controlled by Fresh Carbon Supply." *Nature* 450 (7167): 277–80.
https://doi.org/10.1038/nature06275.

Frioux, Clémence, Enora Fremy, Camille Trottier, and Anne Siegel. 2018. "Scalable and
Exhaustive Screening of Metabolic Functions Carried out by Microbial Consortia."
Bioinformatics 34 (17): i934–43. https://doi.org/10.1093/bioinformatics/bty588.

Gallagher, John L., and F. Gerald Plumley. 1979. "Underground Biomass Profiles and
Productivity in Atlantic Coastal Marshes." *American Journal of Botany* 66 (2):
156–61. https://doi.org/10.1002/j.1537-2197.1979.tb06208.x.

Gibson, Scott M., Stephen P. Ficklin, Sven Isaacson, Feng Luo, Frank A. Feltus, and
Melissa C. Smith. 2013. "Massive-Scale Gene Co-Expression Network
Construction and Robustness Testing Using Random Matrix Theory." *PLoS ONE*
8 (2): e55871. https://doi.org/10.1371/journal.pone.0055871.

Guimond, Julia A, Angelia L Seyfferth, Kevan B Moffett, and Holly A Michael. 2020. "A
Physical-Biogeochemical Mechanism for Negative Feedback between Marsh
Crabs and Carbon Storage." *Environmental Research Letters* 15 (3): 034024.
https://doi.org/10.1088/1748-9326/ab60e2.

Guimond, Julia A, and Joseph Tamborski. 2021. "Salt Marsh Hydrogeology: A Review."
Water 13 (4): 543. https://doi.org/10.3390/w13040543.

Howes, Brian L., John W. H. Dacey, and Gary M. King. 1984. "Carbon Flow through
Oxygen and Sulfate Reduction Pathways in Salt Marsh Sediments1." *Limnology*
and Oceanography 29 (5): 1037–51. https://doi.org/10.4319/lo.1984.29.5.1037.

Inagaki, Fumio, Takuro Nunoura, Satoshi Nakagawa, et al. 2006. "Biogeographical
Distribution and Diversity of Microbes in Methane Hydrate-Bearing Deep Marine
Sediments on the Pacific Ocean Margin." *Proceedings of the National Academy*
of Sciences 103 (8): 2815–20. https://doi.org/10.1073/pnas.0511033103.

Jain, Chirag, Luis M. Rodriguez-R, Adam M. Phillippy, Konstantinos T. Konstantinidis,
and Srinivas Aluru. 2018. "High Throughput ANI Analysis of 90K Prokaryotic
Genomes Reveals Clear Species Boundaries." *Nature Communications* 9 (1):
5114. https://doi.org/10.1038/s41467-018-07641-9.

Jørgensen, Bo Barker, and Antje Boetius. 2007. "Feast and Famine — Microbial Life in
the Deep-Sea Bed." *Nature Reviews Microbiology* 5 (10): 770–81.
https://doi.org/10.1038/nrmicro1745.

Kirkpatrick, John B., Emily A. Walsh, and Steven D'Hondt. 2019. "Microbial Selection
and Survival in Subseafloor Sediment." *Frontiers in Microbiology* 10 (May): 956.
https://doi.org/10.3389/fmicb.2019.00956.

Kirwan, Matthew L., J. Patrick Megonigal, Genevieve L. Noyce, and Alexander J. Smith.
2023. "Geomorphic and Ecological Constraints on the Coastal Carbon Sink."
Nature Reviews Earth & Environment 4 (6): 393–406.
https://doi.org/10.1038/s43017-023-00429-6.

Kuzyakov, Yakov, and Evgenia Blagodatskaya. 2015. "Microbial Hotspots and Hot
Moments in Soil: Concept & Review." *Soil Biology and Biochemistry* 83 (April):
184–99. https://doi.org/10.1016/j.soilbio.2015.01.025.

Langmead, Ben, and Steven L Salzberg. 2012. "Fast Gapped-Read Alignment with
Bowtie 2." *Nature Methods* 9 (4): 357–59. https://doi.org/10.1038/nmeth.1923.

Lever, Mark Alexander. 2012. "Acetogenesis in the Energy-Starved Deep Biosphere – A
Paradox?" *Frontiers in Microbiology* 2. https://doi.org/10.3389/fmicb.2011.00284.

Li, Heng, Bob Handsaker, Alec Wysoker, et al. 2009. "The Sequence Alignment/Map
Format and SAMtools." *Bioinformatics* 25 (16): 2078–79.
https://doi.org/10.1093/bioinformatics/btp352.

Liang, Chao, Joshua P. Schimel, and Julie D. Jastrow. 2017. "The Importance of
Anabolism in Microbial Control over Soil Carbon Storage." *Nature Microbiology* 2
(8): 17105. https://doi.org/10.1038/nmicrobiol.2017.105.

Liu, Jin-e, Rui-Ming Han, Hai-Rong Su, et al. 2017. "Effects of Exotic Spartina
Alterniflora on Vertical Soil Organic Carbon Distribution and Storage Amount in
Coastal Salt Marshes in Jiangsu, China." *Ecological Engineering* 106
(September): 132–39. https://doi.org/10.1016/j.ecoleng.2017.05.041.

Luo, Feng, Yunfeng Yang, Jianxin Zhong, et al. 2007. "Constructing Gene Co-
Expression Networks and Predicting Functions of Unknown Genes by Random
Matrix Theory." *BMC Bioinformatics* 8 (1): 299. https://doi.org/10.1186/1471-
2105-8-299.

Mcleod, Elizabeth, Gail L Chmura, Steven Bouillon, et al. 2011. "A Blueprint for Blue
Carbon: Toward an Improved Understanding of the Role of Vegetated Coastal
Habitats in Sequestering CO_2." *Frontiers in Ecology and the Environment* 9 (10):
552–60. https://doi.org/10.1890/110004.

Moffett, Kevan B., Steven M. Gorelick, Robert G. McLaren, and Edward A. Sudicky.
2012. "Salt Marsh Ecohydrological Zonation Due to Heterogeneous Vegetation–

Groundwater–Surface Water Interactions." *Water Resources Research* 48 (2): 2011WR010874. https://doi.org/10.1029/2011WR010874.

Newman, M. E. J. 2004. "Fast Algorithm for Detecting Community Structure in Networks." *Physical Review E* 69 (6): 066133. https://doi.org/10.1103/PhysRevE.69.066133.

Oksanen, Jari, Gavin Simpson, Guillaume Blanchet, et al. 2024. *Vegan: Community Ecology Package*. Version 2.6-10. Released. https://CRAN.R-project.org/package=vegan.

Olesen, Jens M., Jordi Bascompte, Yoko L. Dupont, and Pedro Jordano. 2006. "The Smallest of All Worlds: Pollination Networks." *Journal of Theoretical Biology* 240 (2): 270–76. https://doi.org/10.1016/j.jtbi.2005.09.014.

Olm, Matthew R, Christopher T Brown, Brandon Brooks, and Jillian F Banfield. 2017. "dRep: A Tool for Fast and Accurate Genomic Comparisons That Enables Improved Genome Recovery from Metagenomes through de-Replication." *The ISME Journal* 11 (12): 2864–68. https://doi.org/10.1038/ismej.2017.126.

Ouyang, X., and S. Y. Lee. 2014. "Updated Estimates of Carbon Accumulation Rates in Coastal Marsh Sediments." *Biogeosciences* 11 (18): 5057–71. https://doi.org/10.5194/bg-11-5057-2014.

Parks, Donovan H., Michael Imelfort, Connor T. Skennerton, Philip Hugenholtz, and Gene W. Tyson. 2015. "CheckM: Assessing the Quality of Microbial Genomes Recovered from Isolates, Single Cells, and Metagenomes." *Genome Research* 25 (7): 1043–55. https://doi.org/10.1101/gr.186072.114.

Payne, Paige E., Loren N. Knobbe, Patricia Chanton, Julian Zaugg, Behzad Mortazavi, and Olivia U. Mason. 2025. "Uncovering Novel Functions of the Enigmatic, Abundant, and Active *Anaerolineae* in a Salt Marsh Ecosystem." *mSystems* 10 (1): e01162-24. https://doi.org/10.1128/msystems.01162-24.

Peng, Qiannan, Lu Lin, Qichao Tu, et al. 2023. "Unraveling the Roles of Coastal Bacterial Consortia in Degradation of Various Lignocellulosic Substrates." *mSystems*, July 7, e01283-22. https://doi.org/10.1128/msystems.01283-22.

Petro, Caitlin, Piotr Starnawski, Andreas Schramm, and Kasper U. Kjeldsen. 2017. "Microbial Community Assembly in Marine Sediments." *Aquatic Microbial Ecology* 79 (3): 177–95. https://doi.org/10.3354/ame01826.

Petro, Caitlin, Birthe Zäncker, Piotr Starnawski, et al. 2019. "Marine Deep Biosphere Microbial Communities Assemble in Near-Surface Sediments in Aarhus Bay." *Frontiers in Microbiology* 10 (April): 758. https://doi.org/10.3389/fmicb.2019.00758.

Porter, Abigail W., and Lily Y. Young. 2014. "Benzoyl-CoA, a Universal Biomarker for
Anaerobic Degradation of Aromatic Compounds." In *Advances in Applied*
Microbiology, vol. 88. Elsevier. https://doi.org/10.1016/B978-0-12-800260-
5.00005-X.

Price, M. N., P. S. Dehal, and A. P. Arkin. 2009. "FastTree: Computing Large Minimum
Evolution Trees with Profiles Instead of a Distance Matrix." *Molecular Biology and*
Evolution 26 (7): 1641–50. https://doi.org/10.1093/molbev/msp077.

Qiu, Yan-Ling, Yuji Sekiguchi, Satoshi Hanada, et al. 2006. "Pelotomaculum
Terephthalicum Sp. Nov. and Pelotomaculum Isophthalicum Sp. Nov.: Two
Anaerobic Bacteria That Degrade Phthalate Isomers in Syntrophic Association
with Hydrogenotrophic Methanogens." *Archives of Microbiology* 185 (3): 172–82.
https://doi.org/10.1007/s00203-005-0081-5.

Seitz, Kiley W, Cassandre S Lazar, Kai-Uwe Hinrichs, Andreas P Teske, and Brett J
Baker. 2016. "Genomic Reconstruction of a Novel, Deeply Branched Sediment
Archaeal Phylum with Pathways for Acetogenesis and Sulfur Reduction." *The*
ISME Journal 10 (7): 1696–705. https://doi.org/10.1038/ismej.2015.233.

Shannon, Paul, Andrew Markiel, Owen Ozier, et al. 2003. "Cytoscape: A Software
Environment for Integrated Models of Biomolecular Interaction Networks."
Genome Research 13 (11): 2498–504. https://doi.org/10.1101/gr.1239303.

Spivak, Amanda C., Jonathan Sanderman, Jennifer L. Bowen, Elizabeth A. Canuel, and
Charles S. Hopkinson. 2019. "Global-Change Controls on Soil-Carbon
Accumulation and Loss in Coastal Vegetated Ecosystems." *Nature Geoscience*
12 (9): 685–92. https://doi.org/10.1038/s41561-019-0435-2.

Stewart, Robert D., Marc D. Auffret, Amanda Warr, et al. 2018. "Assembly of 913
Microbial Genomes from Metagenomic Sequencing of the Cow Rumen." *Nature*
Communications 9 (1): 870. https://doi.org/10.1038/s41467-018-03317-6.

Tebbe, Dennis Alexander, Simone Geihser, Bernd Wemheuer, Rolf Daniel, Hendrik
Schäfer, and Bert Engelen. 2022. "Seasonal and Zonal Succession of Bacterial
Communities in North Sea Salt Marsh Sediments." *Microorganisms* 10 (5): 859.
https://doi.org/10.3390/microorganisms10050859.

Tobias, Craig, and Scott C. Neubauer. 2019. "Salt Marsh Biogeochemistry—An
Overview." In *Coastal Wetlands*. Elsevier. https://doi.org/10.1016/B978-0-444-
63893-9.00016-2.

Vineis, Joseph H, Ashley N Bulseco, and Jennifer L Bowen. 2023. "Microbial
Chemolithoautotrophs Are Abundant in Salt Marsh Sediment Following Long-
Term Experimental Nitrate Enrichment." *FEMS Microbiology Letters* 370
(January): fnad082. https://doi.org/10.1093/femsle/fnad082.

Walsh, Emily A, John B Kirkpatrick, Scott D Rutherford, David C Smith, Mitchell Sogin,
and Steven D'Hondt. 2016. "Bacterial Diversity and Community Composition
from Seasurface to Subseafloor." *The ISME Journal* 10 (4): 979–89.
https://doi.org/10.1038/ismej.2015.175.

Wang, Miao, Erqin Li, Chen Liu, Alexandre Jousset, and Joana F. Salles. 2017.
"Functionality of Root-Associated Bacteria along a Salt Marsh Primary
Succession." *Frontiers in Microbiology* 8 (October): 2102.
https://doi.org/10.3389/fmicb.2017.02102.

Xiao, Kai, Alicia M. Wilson, Hailong Li, and Carolyn Ryan. 2019. "Crab Burrows as
Preferential Flow Conduits for Groundwater Flow and Transport in Salt Marshes:
A Modeling Study." *Advances in Water Resources* 132 (October): 103408.
https://doi.org/10.1016/j.advwatres.2019.103408.

Yao, Zhiyuan, Shicong Du, Chunling Liang, et al. 2019. "Bacterial Community Assembly
in a Typical Estuarine Marsh with Multiple Environmental Gradients." *Applied and*
Environmental Microbiology 85 (6): e02602-18.
https://doi.org/10.1128/AEM.02602-18.

Yu, Tiantian, Weichao Wu, Wenyue Liang, Mark Alexander Lever, Kai-Uwe Hinrichs,
and Fengping Wang. 2018. "Growth of Sedimentary Bathyarchaeota on Lignin as
an Energy Source." *Proceedings of the National Academy of Sciences* 115 (23):
6022–27. https://doi.org/10.1073/pnas.1718854115.

Zhou, Zhichao, Patricia Q. Tran, Adam M. Breister, et al. 2022. "METABOLIC: High-
Throughput Profiling of Microbial Genomes for Functional Traits, Metabolism,
Biogeochemistry, and Community-Scale Functional Networks." *Microbiome* 10
(1): 33. https://doi.org/10.1186/s40168-021-01213-8.

Zimmermann, Johannes, Christoph Kaleta, and Silvio Waschina. 2021. "Gapseq:
Informed Prediction of Bacterial Metabolic Pathways and Reconstruction of
Accurate Metabolic Models." *Genome Biology* 22 (1): 81.
https://doi.org/10.1186/s13059-021-02295-1.