**Advantages of outcrossing in *Plasmodium falciparum*: insights from genetic**

**crosses using fluorescent labelled parasites**

Xue Li [1], Kathrin S. Jutzeler [1], Biley A. Abatiyow [2], Nastaran Rezakhani [2], Meseret T. Haile [2],

Amanda S. Leeb [2], Hardik Patel [2], Stefan H.I. Kappe [2, 3], François Nosten [4, 5], Ian H. Cheeseman

[6], Michael T. Ferdig [7], Timothy J. C. Anderson [1, *], Ashley M. Vaughan [2, 3, *], Sudhir Kumar [2, 8, *]

[1] Disease Intervention and Prevention Program, Texas Biomedical Research Institute, San

Antonio, TX, USA.

[2] Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle,

WA, USA.

[3] Department of Pediatrics, University of Washington, Seattle, WA, USA.

[4] Shoklo Malaria Research Unit, Mahidol-Oxford Tropical Medicine Research Unit, Faculty of

Tropical Medicine, Mahidol University, Mae Sot, Thailand

[5] Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine Research

building, University of Oxford Old Road campus, Oxford, UK

[6] Host Pathogen Interactions Program, Texas Biomedical Research Institute, San Antonio, TX,

USA

[7] Eck Institute for Global Health, Department of Biological Sciences, University of Notre Dame,

Notre Dame, IN, USA.

[8] Current affiliation: Department of Biomedical Sciences, Iowa State University, Ames, Iowa, USA

* Corresponding authors: Timothy J. C. Anderson (tanderso@txbiomed.org), Ashley M. Vaughan

(ashley.vaughan@seattlechildrens.org) and Sudhir Kumar (sudhirk@iastate.edu).

**Keywords:** oocyst, sporozoite, genetic load, outcrossing, inbreeding.

**ABSTRACT**

Malaria parasites are obligately sexual hermaphrodite protozoans with gamete fusion occurring

in the mosquito midgut, followed by meiosis and recombination. Malaria parasite populations

show a spectrum of populations structures ranging from predominantly selfing to highly

outcrossed. However, the fitness consequences of selfing and outcrossing for malaria parasites

are poorly understood. This project was designed to investigate the dynamics of gamete fusion

within the mosquito midgut and the relative fitness of selfed and outcrossed zygotes. We

generated florescent-labelled clones of NF54 (mCherry), an African parasite, and NHP4026

(GFP), a Thai parasite, crossed these parasites, and scored genotypes of 8540 oocysts from 435

mosquitoes sampled from 7 to 14 days post infection. We observed decreasing proportions of

outcrossed oocysts and increasing levels of inbreeding over the course of the infection in two

independently replicated crosses. These results are consistent with the faster maturation of

transmissible sporozoites derived from outcrossed compared with selfed oocysts. Our results

suggest a substantial outcrossing advantage, perhaps because this allows for the removal of

deleterious mutations accumulated during asexual parasite replication in the vertebrate host. We

also found that selfed NF54 oocysts were larger than outcrossed or selfed NHP4026 oocysts,

which may influence production of sporozoites and onward transmission. We conclude that

fluorescent labelled parasites provide clear resolution of mating patterns, temporal dynamics and

transmission potential of malaria parasites in mosquitoes. Importantly, faster maturation of

outcrossed parasites can maximize levels of recombination in transmitted malaria parasite

populations.

**INTRODUCTION**

Many animals and plants are simultaneous hermaphrodites that produce both male and female

gametes and have the capacity for both inbreeding and outbreeding (Goodwillie, et al. 2005; Jarne

and Auld 2006). Self-fertilization has obvious benefits, because reproduction is possible when partners are not available (reproductive assurance). Furthermore, the offspring are 100% related to parents. However, inbreeding can also result in accumulation of deleterious mutations and declining fitness (Charlesworth and Willis 2009). Outcrossing allows purging of deleterious mutations and restoration of fitness, but offspring share only 50% relatedness to their parents. Nevertheless, the advantages of outcrossing are evident from the elaborate mechanisms used by many hermaphrodite plants to avoid inbreeding, including self-incompatibility and physical barriers to pollen transfer (Charlesworth 2006; Zhang, et al. 2024; Shang, et al. 2025). Furthermore, many controlled experiments have directly compared fitness of inbred and outbred plants and animals, revealing substantial fitness advantages to outbreeding (Willis 1993; Coltman, et al. 1999; Meagher, et al. 2000; Christen and Milinski 2003; Goodwillie, et al. 2005).

Malaria parasites are obligately sexual hermaphrodite protozoans that produce both male and female gametes. These parasites have a mixed mating system, and the contribution of inbreeding and outbreeding varies dramatically in different locations (Anderson, Haubold, et al. 2000). Haploid blood stages differentiate into immature male or female sexual stages (gametocytes) in the blood stream. These are ingested by mosquitoes and male and female gametes mature, fuse in the mosquito midgut to form a diploid zygote which develops into a short-lived motile stage, the ookinete, where meiosis and recombination occur. The ookinete crosses the mosquito midgut and attaches to the basal lamina and dedifferentiates into an extracellular oocyst which contains the four haploid products of meiosis. These divide mitotically and after 10-16 days the oocyst ruptures, releasing one to five thousand (Rosenberg and Rungsiwongse 1991) haploid sporozoites into the hemolymph which ultimately invade the salivary gland. Salivary gland sporozoites can be transmitted to the next vertebrate host when the mosquito takes a bloodmeal. During fertilization, if male and female gametes are from the same parasite genotype – which can

happen when a mosquito feeds on a person infected with a single parasite clone – then gamete fusion is inbred. While meiosis and recombination occur, there is no change in parasite genotype from the parent. On the other hand, when gametocytes of two or more genotypes are present in the blood meal, both inbreeding and outcrossing can occur. In the case of outcrossing, recombination results in the reshuffling of chromosomal segments from different parental parasites and the formation of recombinant progeny.

Several controlled genetic crosses have been conducted in the human malaria parasite *Plasmodium falciparum*. This is achieved by growing parasite clones in *in vitro* sexual culture conditions and allowing mosquitoes to feed on blood containing male and female gametocyte stage parasites from two different clones. Vertebrate hosts are needed for mammalian stage parasite development and splenectomized chimpanzees were initially used (Su, et al. 2007). Genetic crosses are now conducted using humanized mice seeded with human hepatocytes that support parasite liver stage development and transition to asexual blood stage replication (Vaughan, et al. 2015; Vendrely, et al. 2020; Button-Simons, et al. 2021; Mok, et al. 2023). Cloning and sequencing of haploid parasites from the mammalian host from prior crosses conducted in chimpanzees or humanized mice have frequently revealed a surprising deficit of inbred parasites (Ranford-Cartwright, et al. 1993; Button-Simons, et al. 2021; Amambua-Ngwa, et al. 2023). For example, genotyping of 376 blood stage cloned progeny derived from three independent *P. falciparum* crosses between NF54 and NHP4026 (or NF54[mCherry] and NHP4026[GFP]) recovered only six selfed progeny (1.6%) (**Table 1**). Assuming random mating and equal frequencies of gametes from the two parents, the expectation would be to recover 50% of inbred parasites (25% of each parental genotype); when gamete frequencies are unbalanced frequencies of inbred parasites should be >50%. Preferential outcrossing of gametes in the mosquito midgut, or

selection of outcrossed parasites during development may explain the deficits of selfed parasites

observed.

We designed experiments to better understand mating patterns of malaria parasites within

mosquitoes in controlled genetic crosses where both inbred and outcrossed mating are possible.

Previous work on parasite mating patterns in the mosquito have utilized PCR genotyping of

polymorphic markers in individual oocysts dissected from the mosquito midgut (**Table 2**)

(Ranford-Cartwright, et al. 1993; Babiker, et al. 1994; Paul, et al. 1995; Razakandrainibe, et al.

2005; Morlais, et al. 2015; Soontarawirat, et al. 2017). This is labor intensive limiting the sample

size possible; studies to date have examined 35 – 623 oocysts. Furthermore, because oocysts

are small and difficult to dissect without rupture, PCR methods are prone to PCR artifacts –

contamination from ruptured oocysts may result in over estimation of heterozygosity, while null

alleles or allelic drop out can result in underestimation of heterozygosity (Anderson, Paul, et al.

2000). Previous studies  have revealed patterns ranging from random mating and good fit to

Hardy-Weinberg expectations (Ranford-Cartwright, et al. 1993), to high levels of inbreeding with

deficits of heterozygous oocysts and positive inbreeding coefficients (Babiker, et al. 1994; Paul,

et al. 1995; Razakandrainibe, et al. 2005; Soontarawirat, et al. 2017) (**Table 2**).

In the current study, we introduced fluorescent markers into two parental parasites at the same

genetic locus, so oocysts resulting from the selfing of parental parasites are labelled with mCherry

("red"), in the case of the African parasite NF54 or GFP ("green"), in the case of the Thai parasite

NHP4026, and heterozygous oocysts, which carry both markers, are labeled with both

fluorophores and are thus "orange". With these marked parasites we could rapidly score genotype

and size of thousands of oocysts from mosquitoes between 7- and 14-days post infection with

gametocytes, allowing a detailed quantification of the dynamics of the parasite mating patterns

and oocyst development. Our data reveal (i) decreasing proportions of heterozygous oocysts and thus (ii) increasing inbreeding coefficients over the course of midgut infections in two independent crosses, consistent with increased development rates of outbred oocysts and consequently, an outbreeding advantage. We also observed a significant impact of genotype on oocyst size. NF54 oocysts were 7-40% bigger than the other oocyst genotypes (NHP4026 selfed and NF54 $\times$ NHP4026 outcrossed), suggesting that oocyst size can play a role in infective stage production and transmission.

**RESULTS**

**Generation of genetic crossing using fluorescently labelled parental parasites**

Transgenic parasites expressing fluorescent proteins are a powerful tool in parasitology research (Hoshizaki, et al. 2022). In this study, we used two parasite strains to generate fluorescently labeled transgenic lines. NF54 is an African strain, isolated in the 1980's, while NHP4026 was obtained in 2007 from a patient at the Shoklo Malaria Research Unit (SMRU) clinic on the Thailand–Myanmar border. We engineered NF54 to express mCherry (red) and NHP4026 to express GFP (green), generating two distinct transgenic lines: NF54$^{mCherry}$ and NHP4026$^{GFP}$ (**Fig. S1 and Table S1**). The fluorophore was expressed constitutively and replaced the *Pfs47* locus in both parasite parents. These edited parasite lines were then used to perform genetic crosses.

**Genetic diversity during oocyst development within the mosquito vector**

Mosquito sampling was conducted across two independent crosses, with midgut dissections and oocyst collections occurring between 8 - 13 days post-infection (DPI) for *Cross 1* and 7 - 14 DPI for *Cross 2* (**Fig. 1**). Following mosquito infections, a total of 8,540 oocysts were counted and analyzed from 509 mosquitoes, including 435 that were infected (**Table S2**). This included 156 infected mosquitoes from *Cross 1* (infection rate: 0.79) and 279 from *Cross 2* (infection rate: 0.89).

The mean oocyst count per mosquito was 23 in *Cross 1* (range: 1 - 137; median: 14) and 17 in

*Cross 2* (range: 1 - 62; median: 13). In both crosses, oocyst burden per mosquito declined

significantly over time (**Fig. 1B**, **Fig. S2**). Prior to DPI 10, the average oocyst burden per mosquito

was more than 20 oocysts; however, a significant reduction was observed after DPI 10 (*Cross 1*:

$p = 4.36 \times 10^{-5}$; *Cross 2*: $p = 1.42 \times 10^{-32}$; two-tailed Welch's *t*-test) (**Table 3**), consistent with

oocyst maturation resulting in sporozoite release.

Reduction in heterozygosity proportion among oocysts over time: We observed a significant

decline in the proportion of heterozygous (outcrossed) oocysts isolated from the mosquito midgut

over time, coinciding with a reduction in overall oocyst counts (**Fig. 2A, Table 3, Fig. S3**). In *Cross*

*1*, the frequency of heterozygous (orange) oocysts decreased from 0.54 on DPI 8 to 0.33 on DPI

13 ($p = 4.84 \times 10^{-4}$, linear regression). A similar trend was observed in *Cross 2*, where

heterozygous oocysts dropped from a frequency of 0.36 on DPI 7 to 0.13 on DPI 14 ($p = 3.10 \times$

$10^{-4}$).

For both crosses, the total proportion of homozygous oocysts (red plus green) decreased over

time, consistent with the increased proportion of heterozygous (orange) oocysts. However, the

proportion of red or green oocysts differed between the two genetic crosses. Higher proportions

of NF54$^{mCherry}$ selfed (red) oocysts were observed in *Cross 2* but not in *Cross 1*.  In *Cross 1*, the

proportion of red oocysts remained relatively stable across time points (frequency from 0.26 to

0.19, $p = 0.854$), while the proportion of NHP4026$^{GFP}$ selfed (green) oocysts increased

significantly from 0.19 to 0.48 ($p = 0.011$). In *Cross 2*, NF54$^{mCherry}$ selfed oocysts consistently

dominated the population, with a non-significant increase from 0.53 to 0.68 over time ($p = 0.093$).

Meanwhile, NHP4026$^{GFP}$ selfed oocysts increased significantly from 0.11 to 0.19 ($p = 0.019$).

<u>Increased inbreeding coefficient over time in mosquito midgut</u>: To quantify the observed decline

in heterozygosity proportion, we calculated hierarchical Wright's F-statistics for each cross and

time point (**Fig. 2B, Table 3**). The inbreeding coefficient ($F_{IS}$), which reflects deviations from

random mating within mosquito midguts, increased significantly over time for both crosses. In

*Cross 1*, $F_{IS}$ rose from –0.110 on DPI 8 (95% CI: –0.240 to 0.030) to 0.322 on DPI 13 (95% CI: –

0.067 to 0.548), with a strong linear trend ($p = 8.19 \times 10^{-4}$, R² = 0.954, linear regression). Similarly,

in *Cross 2*, $F_{IS}$ increased from 0.111 on DPI 7 (95% CI: 0.032 to 0.153) to 0.657 on DPI 14 (95%

CI: 0.437 to 0.762) ($p = 6.76 \times 10^{-4}$, R² = 0.873).

In both crosses, the among-host differentiation ($F_{ST}$) remained close to zero across time points,

indicating little genetic structure among individual mosquitoes. Consequently, the total inbreeding

coefficient ($F_{IT}$) followed a similar trajectory to $F_{IS}$, reinforcing the conclusion that temporal

increases in homozygosity were primarily driven by within-mosquito processes rather than

among-mosquito differentiation.

**Oocyst size dynamics**

To investigate the factors influencing oocyst size, we performed multiple linear regression using

DPI, oocyst genotype, and oocyst burden (total oocyst number per mosquito) as predictors. The

analysis was conducted separately for each genetic cross (**Fig. 3, Fig. S4, Table S3**).

Across both crosses, DPI was positively and significantly associated with oocyst size (volume),

which is to be expected as oocysts grow larger as they mature over time. In *Cross 1*, the effect

was substantial ($p = 2.38 \times 10^{-26}$, multiple linear regression), while in *Cross 2* the growth rate was

more modest but still highly significant ($p = 1.60 \times 10^{-8}$). These results confirm a strong temporal

component in oocyst development.

Oocyst genotypes also had a significant impact on size. NF54$^{mCherry}$ selfed (red) oocysts were

consistently and significantly larger than NHP4026$^{GFP}$ selfed (green) in both crosses (*Cross 1*:

32.9% larger, *p* = 8.44 × 10$^{-15}$; *Cross 2*: 40.1% larger, *p* = 3.15 × 10$^{-23}$). Orange (heterozygous)

oocysts were significantly larger than green in *Cross 1* (24.6% larger, *p* = 1.17 × 10$^{-13}$) but showed

no significant difference in *Cross 2* (*p* = 0.239). This pattern reinforces the conclusion that the

NF54 (red) genotype is associated with increased oocyst size, while the effect of the heterozygous

(orange) genotype may be context dependent.

Total oocyst burden showed a negative relationship with oocyst volume in both crosses. In *Cross*

*2*, this association was statistically significant (*p* = 1.08 × 10$^{-4}$), suggesting a crowding effect where

high parasite loads lead to reduced oocyst size, likely due to competition for space or resources.

In *Cross 1*, the trend was similar but not statistically significant (*p* = 0.094).

**DISCUSSION**

Fluorescent labelling of parental parasites enabled high-throughput, real-time analysis of oocyst

genotype (selfed vs. outcrossed) without PCR-based artifacts in thousands of oocysts. We

observed a significant temporal decline in the proportion of heterozygous oocysts within the

mosquito midgut, accompanied by a corresponding increase in the inbreeding coefficient ($F_{IS}$).

This pattern was consistent in two independent crosses and suggests that outcrossed

(heterozygous) oocysts mature and rupture earlier than homozygous (selfed) oocysts (**Fig. 2C**).

Consequently, outcrossed oocysts become progressively underrepresented in the midgut at later

time points due to more rapid development. Based on these dynamics, we infer that sporozoites

derived from heterozygous oocysts are likely to reach the salivary glands earlier than those from

homozygous oocysts, thereby gaining a temporal advantage for transmission. Faster migration of

sporozoites to the salivary glands is likely to be important for transmission, given that many

mosquitoes die before taking a second (infectious) blood meal. This makes mosquito longevity

critical for malaria transmission models (Smith and McKenzie 2004). The observed >20%

reduction in the frequency of outcrossed oocysts between days 7 and 14 post-infection

corresponds to an estimated selection coefficient of ~3 - 4% per day (**Table 3**), highlighting a

substantial advantage for outcrossed oocysts in the transition to infectious sporozoites. Our data

is consistent with a strong outcrossing advantage in the maturation of oocysts in the NF54$^{mCherry}$

× NHP4026$^{GFP}$ genetic cross.

Although our current study focused on the mosquito stages of the malaria parasite life cycle, there

is evidence that the advantages of outcrossing extend to other developmental stages. In our

previous work on *P. falciparum* genetic crosses between MKK2835 and NHP1337 (Li, et al. 2019),

we observed strong selection against selfed progeny in pooled samples during early blood stage

development, with recombinant progeny outcompeting inbred parasites carrying parental

genotypes. These findings suggest that outcrossed progeny may survive and propagate more

successfully in the vertebrate host. Combined with our current observation that the heterozygous

oocysts in our crosses mature earlier and likely invade the salivary glands sooner, outcrossing

may confer broader fitness advantages across the parasite's life cycle. Given that liver stage

development represents a critical bottleneck, future work could investigate whether similar

selection pressures favor outcrossed parasites during hepatic schizogony, potentially shaping

host colonization dynamics and transmission efficiency.

We suggest two possible explanations for the difference in development and maturation of inbred

and outcrossed oocysts in the mosquito midgut. First, outcrossed oocysts may more effectively

evade the mosquito immune response than inbred oocysts leading to faster development and

rupture. Mosquito immune responses against oocysts are mediated by midgut stem cells and enteroblasts or by hemocytes (Barletta, et al. 2025) and the specific oocyst molecular signals used by the mosquito to recognize oocysts are not currently known. The mechanisms that might underlie differential survival of inbred and outbred oocysts are therefore unclear. Second, outcrossing may be advantageous because recombination between different parasite genotypes allows removal of "genetic load" (Bertorelle, et al. 2022) caused by deleterious mutations accumulated during asexual multiplication. These deleterious mutations cannot be removed during selfing because they are present in both male and female gametes. The burden of deleterious mutations may be considerable. After initial infection by mosquito inoculation, sporozoites establish infection in liver cells, where they divide rapidly. Blood stage infection in the human host is initiated by the emergence of > 90,000 clonal merozoites from infected liver cells (Vaughan, et al. 2012; Vaughan and Kappe 2017). Populations of malaria-infected red blood cells within an infected individual can reach $10^{11}$ -$10^{12}$. This equates to a minimum of 37-40 mitotic cell divisions if we assume geometrical cell division; in reality cell division is asynchronous (Arnot, et al. 2011) so actual numbers of cell divisions may be considerably higher. Given SNP mutation rates of 2.10-5.23 × $10^{-10}$ per site per asexual cycle in *P. falciparum* (Bopp, et al. 2013; Hamilton, et al. 2017; McDew-White, et al. 2019), indel and microsatellite mutations that are 3-4 orders of magnitude higher (Hamilton, et al. 2017; McDew-White, et al. 2019), and conservative estimates of 37-40 cell divisions in the human host, we estimate that on average, in mature infections, each blood stage parasite contains at least 0.45 ± 0.007 SNPs and 3,433 ± 1,591 indel/microsatellite mutations.

Single cell approaches now allow direct quantification of mutations within infections (Nkhoma, et al. 2020; Dia and Cheeseman 2021; Dia, et al. 2021). Dia *et al.* (Dia, et al. 2021) used single cell genome sequencing of natural *P. falciparum* and *P. vivax* infections. In 15 *P. falciparum* infections

they identified 203 *de novo* mutations in 181 genes, while in 11 *P. vivax* infections they identified 159 *de novo* mutations in 90 genes. In both parasite species, Dia *et al.* (Dia, et al. 2021) observed strong evidence for selection, with excess non-synonymous mutations and enrichment of mutations in some gene families, including the *ApiAP2* gene family. These data are consistent with substantial genetic load resulting from accumulated mutations. Furthermore, the strong signatures of selection and non-random distribution of mutations are consistent with short term adaptation (Didelot, et al. 2016) for blood stage growth, which may have deleterious impact on parasite development in the mosquito or elsewhere in the parasite lifecycle. Outcrossing allows parasites to remove mutations that have a deleterious effect on oocyst growth and maturation, while inbreeding retains such mutations. The ookinete and oocysts are the only diploid stages of the mosquito lifecycle. Outcrossed ookinetes or oocysts are therefore more likely than inbred parasites to have fully functional gene product from at least one parent for genes critical for parasite development in the mosquito. While we suspect that removal or masking of genetic load is the principal driver of the outcrossing advantage observed in these laboratory crosses, it is also the case that in nature recombination has additional benefits for parasites. For example, recombination generates diversity in surface antigens that are key to immune evasion and allows evolution of multidrug resistance.

Malaria parasite populations range from predominantly outcrossed, to highly inbred, and this scales with levels of transmission (Anderson, Haubold, et al. 2000; Nkhoma, et al. 2013; Schaffner, et al. 2023). In countries with low levels of transmission and elevated inbreeding (e.g., South America and Southeast Asia), parasites are expected to accumulate substantial levels of genetic load and inbreeding depression, due to serial inbreeding across multiple parasite generations. Population genomic studies reveal parasites sampled 14 years apart that are identical-by-descent (IBD) in Colombia (Taylor, et al. 2020), 5 years apart in Myanmar (Li, et al.

2025) or 6 years apart in Senegal (Redmond, et al. 2018). Furthermore, in the Senegalese study (Redmond, et al. 2018) numbers of mutations increase within clonal lineages over time. Outcrossing may be help restore fitness in parasite lineages where serial sefling and inbreeding predominates. Observations that are consistent with this include: (i) multiple clone infections typically contain related parasites from recent recombination events, even in low transmission regions (Nkhoma, et al. 2012; Nkhoma, et al. 2020) and (ii) studies of genomic epidemiology in low transmission regions often contain large numbers of closely related parasites, generated from recombination (Wasakul, et al. 2023; Li, et al. 2025).

Prior studies based on PCR genotyping of oocysts in field collected mosquitoes have typically revealed mild to severe deficits of heterozygotes (**Table 2**). This pattern is consistent with our observations of heterozygote deficit in laboratory crosses and suggests that the heterozygote advantage may also drive patterns observed in the field. It is not possible to determine time since infection in field collected mosquitoes. This may influence the wide range of heterozygote deficits observed in prior studies, with older infections showing greater heterozygote deficit than young infections. Other factors may contribute to deviations from Hardy-Weinberg expectations. Hardy-Weinberg genotype ratios are expected when mating is random. However, imbalances in male/female gametocyte ratios between parental strains may skew mating outcomes away from the Hardy-Weinberg equilibrium (**Fig. S5**) (Waples 2015). Indeed, there is an extensive literature describing deviations from one-to-one gamete sex ratios in malaria parasites, and changes in sex ratio during the course of infection (Reece, et al. 2008).

Our results demonstrate a strong genetic influence on oocyst size, a key indicator of parasite developmental fitness and transmission potential (Qi, et al. 2015; Oke, et al. 2024). Contrary to expectations that outcrossed oocysts might exhibit superior growth, we observed no consistent

size advantage among heterozygotes. Instead, oocysts derived from the NF54 parent were significantly larger than those from NHP4026 with heterozygous oocysts displaying intermediate sizes. Our findings are consistent with previous studies in rodent malaria parasites *Plasmodium yoelii* (Qi, et al. 2015) and *Plasmodium berghei* (Balakrishnan, et al. 2025), which also demonstrated that oocyst size is a genotype-dependent trait. In *P. yoelii*, disruption of the D-type small-subunit rRNA gene (*D-ssu*) resulted in significantly smaller oocysts and a complete loss of sporozoite development. In *P. berghei*, a chloroquine-resistance-transporter-like (CRTL) protein was shown to be essential for both oocyst growth and sporogony. These studies underscore the importance of specific genetic loci in regulating mosquito-stage parasite development. Furthermore, recent work has shown that the nutritional status of the mosquito does not fully account for variation in oocyst size, further supporting the primacy of genotype in determining this trait (Oke, et al. 2024). Together, these results reinforce the conclusion that oocyst size is a heritable, genotype-dependent phenotype shaped by parasite genomic background. The large size of NF54-derived oocysts may reflect natural variation in reproductive strategy among *P. falciparum* isolates. With over 237 genes estimated to be expressed during the oocyst stage (Howick, et al. 2019; Real, et al. 2021), future genetic mapping efforts could help identify the loci underlying this variation and provide insight into the evolutionary forces shaping transmission-related traits.

Our study has some limitations. To visualize and precisely quantify parasite genotypes, we examined a single parasite cross (NF54 × NHP4026) for which fluorescent markers were developed. Examining levels of heterozygote advantage in additional crosses to better understand the importance of this phenomenon is ultimately required. Also, while informative, fluorescent color does not capture more subtle genomic contributions (e.g., recombination patterns, epistasis). Given the unusual pattern of maternal inheritance observed in prior crosses

(Li, et al. 2019; Button-Simons, et al. 2021), fluorescent markers for monitoring mitochondria and

apicoplast genomes could also be informative for future work. Nevertheless, the current study

clearly demonstrates that outcrossed parasites develop faster than selfed parasites and dominate

early stage infections. Our results from mosquito stage infections parallel the selective advantage

of outcrossed blood stage parasites relative to selfed parasites and indicate advantages to

outcrossing in both the vertebrate and invertebrate host (Li, et al. 2019). These dynamics

maximize the impact of limited recombination and maintain genetic diversity and adaptive

potential in natural *P. falciparum* populations.

**MATERIALS AND METHODS**

**Ethics approval and consent to participate**

The study was performed in strict accordance with the recommendations in the Guide for the Care

and Use of Laboratory Animals of the National Institutes of Health (NIH), USA. To this end, the

Seattle Children's Research Institute (SCRI) has an Assurance from the Public Health Service

(PHS) through the Office of Laboratory Animal Welfare (OLAW) for work approved by its

Institutional Animal Care and Use Committee (IACUC). All the work carried out in this study was

specifically reviewed and approved by the SCRI IACUC.

**Generation of fluorescent-labelled parasites**

We generated transgenic *P. falciparum* parasites by integrating fluorescent markers into NF54

and NHP4026 strains using a CRISPR/Cas9-based transgenesis (**Fig. S1**). Parasites were

maintained as asexual blood-stage cultures following standard protocols in complete RPMI

medium supplemented with either 0.5% AlbuMAX™ II (Thermo Scientific) or 10% (v/v) type O+

human serum, with daily medium changes. Fluorescent labeling was achieved through double-

crossover homologous recombination, using the plasmids pLf0115 (a modified version of pLf0116

(Mogollon, et al. 2016) with EF1α-GFP replaced by EF1α-mCherry) and pLf0019 (Mogollon, et al.

2016). Transfected NF54 and NHP4026 parasites were selected with WR99210 (8 nM) and

blasticidin (2.5 µg/mL) for five days, followed by cloning by limiting dilution. Successful integration

was confirmed by genotyping PCRs, demonstrating disruption of the *Pfs47* gene and insertion of

EF1α-mCherry in NF54 and EF1α-GFP in NHP4026 (**Fig. S1**). Oligonucleotide primers used for

transfection and genotyping are listed in **Table S1**. Clones NF54[mCherry] 2B5 and NHP4026[GFP] 1F

were used for subsequent experiments.

**Preparation of genetic crosses**

We used *Anopheles stephensi* mosquitoes and human liver-chimeric FRG huHep mice, following

the protocol described by Vaughan et al. (Vaughan, et al. 2015), to generate two genetic crosses

(*Cross 1* and *Cross 2*) between NF54[mCherry] and NHP4026[GFP] parasites. Gametocytes from each

parental strain were diluted to 0.5% gametocytemia in a human serum/erythrocyte mixture to

prepare infectious blood meals (IBMs). The IBMs were mixed at an approximate 1:1 ratio and

used to infect ~450 mosquitoes per cross.

**Fluorescence microscopy**

Midguts were dissected from *Anopheles* mosquitoes at 7–14 days post-infection in phosphate-

buffered saline (PBS). Wet mounts were prepared on glass slides, and live-cell fluorescence

microscopy was performed to detect and differentiate oocysts labeled with red (mCherry) and

green (GFP) fluorescent markers. Images were acquired in red, green, and brightfield channels

using a Keyence fluorescence microscope equipped with a 10× air objective. Oocysts were

manually counted based on fluorescence: green fluorescence indicated GFP-positive (NHP4026)

oocysts, red fluorescence indicated mCherry-positive (NF54) oocysts, and co-expression of both

signals (appearing orange in merged images) was used to identify heterozygous oocysts. Counts

were recorded per midgut and classified by color to quantify the proportions of homozygous and

heterozygous oocysts.

**Oocyst size quantification**

We quantified oocyst size using QuPath v0.2.3 (Bankhead, et al. 2017). Fluorescence microscopy

images of dissected mosquito midguts were imported into the software and individual oocysts

were initially annotated manually using the polygon tool to train the program for automated

analysis. Subsequent automated batch analyses were then performed to classify oocysts by

fluorescent label (red, green, or merged) and to calculate their area (in μm²) based on the

calibrated image scale. The measured areas were then converted to volume estimates to

approximate oocyst size (volume, in μm³), assuming that oocysts are spherical. All measurements

were exported for downstream statistical analysis. Consistent imaging settings and scale

calibrations were maintained across all samples to ensure accuracy and comparability.

**Statistical analysis and measurement of deviations from Hardy-Weinberg Equilibrium**

To investigate the population genetic structure of oocysts, we estimated Wright's *F*-statistics,

which quantify deviations from expected genetic diversity under random mating. Oocyst

genotypes were analyzed across three hierarchical levels: **I** – individual oocysts, **S** –

subpopulations of oocysts within individual mosquitoes, and **T** – the total population of oocysts

across all mosquitoes.

The *F*-statistics were calculated as follows:

• $F_{IS} = 1 - \frac{H_I}{H_S}$

*F*$_{IS}$ ranges from -1 to 1 and measures inbreeding coefficient of individual oocysts relative to

individual mosquitoes. A value of 0 indicates random mating, negative values indicate an

excess of heterozygotes consistent with outcrossing, and positive values indicate a deficit of

heterozygotes, suggesting selfing or inbreeding.

• $F_{ST} = 1 - \frac{H_S}{H_T}$

$F_{ST}$ ranges from 0 to 1 and measures genetic differentiation among subpopulations (individual

mosquitoes). A value of 0 indicates no differentiation, while higher values reflect increasing

population genetic divergence. In rare cases, $F_{ST}$ values may fall below zero due to statistical

noise and are generally interpreted as indicating no genetic differentiation between

populations.

• $F_{IT} = 1 - \frac{H_I}{H_T}$

$F_{IT}$ ranges from −1 to 1 and quantifies the inbreeding coefficient of individuals relative to the

total population. A value of 0 indicates random mating across the entire population. Negative

$F_{IT}$ values reflect an excess of heterozygotes, suggesting high levels of outcrossing, while

positive values indicate reduced heterozygosity, consistent with inbreeding or population

structure.

Where:

• $H_I$ = observed heterozygosity at the individual oocyst level

• $H_S$ = expected heterozygosity within mosquitoes (subpopulations)

• $H_T$ = expected heterozygosity in the total oocyst population

These statistics were computed using the R packages *HardyWeinberg* (Graffelman 2015) and

*hierfstat* (Goudet 2005). The 95% confidence intervals (CIs) were calculated using a bootstrap

approach with 1,000 resampling replicates. Significance testing of the fixation indices was

performed using non-parametric permutation tests to assess whether observed values significantly deviated from those expected under random mating.

**DATA AVAILABILITY**

All data supporting the conclusions of this study are included in the main text and/or Supplementary Materials. Genotype data for cloned progeny are available in the following GitHub repository: https://github.com/emilyli0325/P01-cloned.progeny. Additional data related to this paper may be requested from the authors.

**AUTHOR CONTRIBUTIONS**

XL, TJCA, AMV, and SK conceived and designed the study. BAA, NR, MTH, ASL, HP, AMV and SK generated the fluorescent-labeled *Plasmodium falciparum* parental lines. BAA, NR and MTH and SK carried out the genetic crosses, dissected mosquito midguts, and quantified oocysts. KSJ performed oocyst size measurements. XL conducted statistical analyses and data visualization, with input from TJCA. XL, TJCA, AMV, and SK drafted the initial manuscript. All authors contributed to critical revision of the manuscript, had full access to all data, and approved the final version for submission.

**ACKNOWLEDGMENTS**

This work was supported by National Institutes of Health (NIH) program project grant P01 AI127338 (to MTF and AMV) and by NIH grant R37 AI048071 (to TJCA). Work at Texas Biomedical Research Institute was conducted in facilities constructed with support from Research Facilities Improvement Program grant C06 RR013556 from the National Center for Research Resources. The authors thank Drs. Shinya Miyazaki and Chris Janse from Leiden university for providing plasmids pLf0115, pLf0116 and pLf0019 for the generation of fluorescent parasites.

**REFERENCES**

Amambua-Ngwa A, Button-Simons KA, Li X, Kumar S, Brenneman KV, Ferrari M, Checkley LA, Haile MT,
Shoue DA, McDew-White M, et al. 2023. Chloroquine resistance evolution in Plasmodium falciparum is
mediated by the putative amino acid transporter AAT1. Nat Microbiol 8:1213-1226.
Anderson TJ, Haubold B, Williams JT, Estrada-Franco JG, Richardson L, Mollinedo R, Bockarie M, Mokili J,
Mharakurwa S, French N, et al. 2000. Microsatellite markers reveal a spectrum of population structures
in the malaria parasite Plasmodium falciparum. Mol Biol Evol 17:1467-1482.
Anderson TJ, Paul RE, Donnelly CA, Day KP. 2000. Do malaria parasites mate non-randomly in the
mosquito midgut? Genet Res 75:285-296.
Arnot DE, Ronander E, Bengtsson DC. 2011. The progression of the intra-erythrocytic cell cycle of
Plasmodium falciparum and the role of the centriolar plaques in asynchronous mitotic division during
schizogony. Int J Parasitol 41:71-80.
Babiker HA, Ranford-Cartwright LC, Currie D, Charlwood JD, Billingsley P, Teuscher T, Walliker D. 1994.
Random mating in a natural population of the malaria parasite Plasmodium falciparum. Parasitology 109
( Pt 4):413-421.
Balakrishnan A, Hunziker M, Tiwary P, Pandey V, Drew D, Billker O. 2025. A CRISPR homing screen finds a
chloroquine resistance transporter-like protein of the Plasmodium oocyst essential for mosquito
transmission of malaria. Nat Commun 16:3895.
Bankhead P, Loughrey MB, Fernandez JA, Dombrowski Y, McArt DG, Dunne PD, McQuaid S, Gray RT,
Murray LJ, Coleman HG, et al. 2017. QuPath: Open source software for digital pathology image analysis.
Sci Rep 7:16878.
Barletta ABF, Barillas-Mury C, Molina-Cruz A. 2025. Mosquito immune responses to Plasmodium
parasites that limit malaria transmission. Cell Mol Life Sci 82:143.
Bertorelle G, Raffini F, Bosse M, Bortoluzzi C, Iannucci A, Trucchi E, Morales HE, van Oosterhout C. 2022.
Genetic load: genomic estimates and applications in non-model animals. Nat Rev Genet 23:492-503.
Bopp SE, Manary MJ, Bright AT, Johnston GL, Dharia NV, Luna FL, McCormack S, Plouffe D, McNamara
CW, Walker JR, et al. 2013. Mitotic evolution of Plasmodium falciparum shows a stable core genome but
recombination in antigen families. PLoS Genet 9:e1003293.
Button-Simons KA, Kumar S, Carmago N, Haile MT, Jett C, Checkley LA, Kennedy SY, Pinapati RS, Shoue
DA, McDew-White M, et al. 2021. The power and promise of genetic mapping from Plasmodium
falciparum crosses utilizing human liver-chimeric mice. Commun Biol 4:734.
Charlesworth D. 2006. Evolution of plant breeding systems. Curr Biol 16:R726-735.
Charlesworth D, Willis JH. 2009. The genetics of inbreeding depression. Nat Rev Genet 10:783-796.
Christen M, Milinski M. 2003. The consequences of self-fertilization and outcrossing of the cestode
Schistocephalus solidus in its second intermediate host. Parasitology 126:369-378.
Coltman DW, Pilkington JG, Smith JA, Pemberton JM. 1999. Parasite-Mediated Selection against Inbred
Soay Sheep in a Free-Living Island Populaton. Evolution 53:1259-1267.
Dia A, Cheeseman IH. 2021. Single-cell genome sequencing of protozoan parasites. Trends Parasitol
37:803-814.
Dia A, Jett C, Trevino SG, Chu CS, Sriprawat K, Anderson TJC, Nosten F, Cheeseman IH. 2021. Single-
genome sequencing reveals within-host evolution of human malaria parasites. Cell Host Microbe
29:1496-1506 e1493.

Didelot X, Walker AS, Peto TE, Crook DW, Wilson DJ. 2016. Within-host evolution of bacterial pathogens. Nat Rev Microbiol 14:150-162.

Goodwillie C, Kalisz S, Eckert CG. 2005. The Evolutionary Enigma of Mixed Mating Systems in Plants: Occurrence, Theoretical Explanations, and Empirical Evidence. Annual Review of Ecology, Evolution, and Systematics 36:47-79.

Goudet J. 2005. Hierfstat, a package for R to compute and test variance components and F-statistics. Molecular Ecology Notes 5:184-186.

Graffelman J. 2015. Exploring Diallelic Genetic Markers: The HardyWeinberg Package. Journal of statistical software 64:1 - 23.

Hamilton WL, Claessens A, Otto TD, Kekre M, Fairhurst RM, Rayner JC, Kwiatkowski D. 2017. Extreme mutation bias and high AT content in Plasmodium falciparum. Nucleic Acids Res 45:1889-1901.

Hoshizaki J, Jagoe H, Lee MCS. 2022. Efficient generation of mNeonGreen Plasmodium falciparum reporter lines enables quantitative fitness analysis. Front Cell Infect Microbiol 12:981432.

Howick VM, Russell AJC, Andrews T, Heaton H, Reid AJ, Natarajan K, Butungi H, Metcalf T, Verzier LH, Rayner JC, et al. 2019. The Malaria Cell Atlas: Single parasite transcriptomes across the complete Plasmodium life cycle. Science 365.

Jarne P, Auld JR. 2006. Animals mix it up too: the distribution of self-fertilization among hermaphroditic animals. Evolution 60:1816-1824.

Li X, Arya GA, Thu AM, Landier J, Parker DM, Delmas G, Reyes A, Lwin KM, Sriprawat K, Nosten F, et al. 2025. Malaria parasite population genomics during an elimination program in Eastern Myanmar.2025.2005.2021.655408.

Li X, Kumar S, Brenneman KV, Anderson TJC. 2022. Bulk segregant linkage mapping for rodent and human malaria parasites. Parasitol Int 91:102653.

Li X, Kumar S, McDew-White M, Haile M, Cheeseman IH, Emrich S, Button-Simons K, Nosten F, Kappe SHI, Ferdig MT, et al. 2019. Genetic mapping of fitness determinants across the malaria parasite Plasmodium falciparum life cycle. PLoS Genet 15:e1008453.

McDew-White M, Li X, Nkhoma SC, Nair S, Cheeseman I, Anderson TJC. 2019. Mode and Tempo of Microsatellite Length Change in a Malaria Parasite Mutation Accumulation Experiment. Genome Biol Evol 11:1971-1985.

Meagher S, Penn DJ, Potts WK. 2000. Male-male competition magnifies inbreeding depression in wild house mice. Proc Natl Acad Sci U S A 97:3324-3329.

Mogollon CM, van Pul FJ, Imai T, Ramesar J, Chevalley-Maurel S, de Roo GM, Veld SA, Kroeze H, Franke-Fayard BM, Janse CJ, et al. 2016. Rapid Generation of Marker-Free P. falciparum Fluorescent Reporter Lines Using Modified CRISPR/Cas9 Constructs and Selection Protocol. PLoS One 11:e0168362.

Mok S, Yeo T, Hong D, Shears MJ, Ross LS, Ward KE, Dhingra SK, Kanai M, Bridgford JL, Tripathi AK, et al. 2023. Mapping the genomic landscape of multidrug resistance in Plasmodium falciparum and its impact on parasite fitness. Sci Adv 9:eadi2364.

Morlais I, Nsango SE, Toussile W, Abate L, Annan Z, Tchioffo MT, Cohuet A, Awono-Ambene PH, Fontenille D, Rousset F, et al. 2015. Plasmodium falciparum mating patterns and mosquito infectivity of natural isolates of gametocytes. PLoS One 10:e0123777.

Nkhoma SC, Nair S, Al-Saai S, Ashley E, McGready R, Phyo AP, Nosten F, Anderson TJ. 2013. Population genetic correlates of declining transmission in a human pathogen. Mol Ecol 22:273-285.

Nkhoma SC, Nair S, Cheeseman IH, Rohr-Allegrini C, Singlam S, Nosten F, Anderson TJ. 2012. Close kinship within multiple-genotype malaria parasite infections. Proc Biol Sci 279:2589-2598.

Nkhoma SC, Trevino SG, Gorena KM, Nair S, Khoswe S, Jett C, Garcia R, Daniel B, Dia A, Terlouw DJ, et al.
2020. Co-transmission of Related Malaria Parasite Lineages Shapes Within-Host Parasite Diversity. Cell
Host Microbe 27:93-103.e104.
Oke CE, O'Donnell AJ, Schneider P, Reece SE. 2024. Plasticity in malaria parasite development: mosquito
resources influence vector-to-host transmission potential.  Volume 2 - 2024.
Paul RE, Packer MJ, Walmsley M, Lagog M, Ranford-Cartwright LC, Paru R, Day KP. 1995. Mating patterns
in malaria parasite populations of Papua New Guinea. Science 269:1709-1711.
Qi Y, Zhu F, Eastman RT, Fu Y, Zilversmit M, Pattaradilokrat S, Hong L, Liu S, McCutchan TF, Pan W, et al.
2015. Regulation of Plasmodium yoelii oocyst development by strain- and stage-specific small-subunit
rRNA. mBio 6:e00117.
Ranford-Cartwright LC, Balfe P, Carter R, Walliker D. 1993. Frequency of cross-fertilization in the human
malaria parasite Plasmodium falciparum. Parasitology 107 ( Pt 1):11-18.
Razakandrainibe FG, Durand P, Koella JC, De Meeus T, Rousset F, Ayala FJ, Renaud F. 2005. "Clonal"
population structure of the malaria agent Plasmodium falciparum in high-infection regions. Proc Natl
Acad Sci U S A 102:17388-17393.
Real E, Howick VM, Dahalan FA, Witmer K, Cudini J, Andradi-Brown C, Blight J, Davidson MS, Dogga SK,
Reid AJ, et al. 2021. A single-cell atlas of Plasmodium falciparum transmission through the mosquito. Nat
Commun 12:3196.
Redmond SN, MacInnis BM, Bopp S, Bei AK, Ndiaye D, Hartl DL, Wirth DF, Volkman SK, Neafsey DE. 2018.
De Novo Mutations Resolve Disease Transmission Pathways in Clonal Malaria. Mol Biol Evol 35:1678-
1689.
Reece SE, Drew DR, Gardner A. 2008. Sex ratio adjustment and kin discrimination in malaria parasites.
Nature 453:609-614.
Rosenberg R, Rungsiwongse J. 1991. The number of sporozoites produced by individual malaria oocysts.
Am J Trop Med Hyg 45:574-577.
Schaffner SF, Badiane A, Khorgade A, Ndiop M, Gomis J, Wong W, Ndiaye YD, Diedhiou Y, Thwing J, Seck
MC, et al. 2023. Malaria surveillance reveals parasite relatedness, signatures of selection, and correlates
of transmission across Senegal. Nat Commun 14:7268.
Shang L, Gad K, Lenhard M. 2025. Converging on long and short: The genetics, molecular biology and
evolution of heterostyly. Curr Opin Plant Biol 85:102731.
Smith DL, McKenzie FE. 2004. Statics and dynamics of malaria infection in Anopheles mosquitoes. Malar
J 3:13.
Soontarawirat I, Andolina C, Paul R, Day NPJ, Nosten F, Woodrow CJ, Imwong M. 2017. Plasmodium
vivax genetic diversity and heterozygosity in blood samples and resulting oocysts at the Thai-Myanmar
border. Malar J 16:355.
Su X, Hayton K, Wellems TE. 2007. Genetic linkage and association analyses for trait mapping in
Plasmodium falciparum. Nat Rev Genet 8:497-506.
Taylor AR, Echeverry DF, Anderson TJC, Neafsey DE, Buckee CO. 2020. Identity-by-descent with
uncertainty characterises connectivity of Plasmodium falciparum populations on the Colombian-Pacific
coast. PLoS Genet 16:e1009101.
Vaughan AM, Kappe SHI. 2017. Malaria Parasite Liver Infection and Exoerythrocytic Biology. Cold Spring
Harb Perspect Med 7.
Vaughan AM, Mikolajczak SA, Wilson EM, Grompe M, Kaushansky A, Camargo N, Bial J, Ploss A, Kappe
SH. 2012. Complete Plasmodium falciparum liver-stage development in liver-chimeric mice. J Clin Invest
122:3618-3628.

Vaughan AM, Pinapati RS, Cheeseman IH, Camargo N, Fishbaugher M, Checkley LA, Nair S, Hutyra CA,
Nosten FH, Anderson TJ, et al. 2015. Plasmodium falciparum genetic crosses in a humanized mouse
model. Nat Methods 12:631-633.
Vendrely KM, Kumar S, Li X, Vaughan AM. 2020. Humanized Mice and the Rebirth of Malaria Genetic
Crosses. Trends Parasitol 36:850-863.
Waples RS. 2015. Testing for Hardy-Weinberg proportions: have we lost the plot? J Hered 106:1-19.
Wasakul V, Disratthakit A, Mayxay M, Chindavongsa K, Sengsavath V, Thuy-Nhien N, Pearson RD,
Phalivong S, Xayvanghang S, Maude RJ, et al. 2023. Malaria outbreak in Laos driven by a selective sweep
for Plasmodium falciparum kelch13 R539T mutants: a genetic epidemiology analysis. Lancet Infect Dis
23:568-577.
Willis JH. 1993. Effects of Different Levels of Inbreeding on Fitness Components in Mimulus Guttatus.
Evolution 47:864-876.
Zhang D, Li YY, Zhao X, Zhang C, Liu DK, Lan S, Yin W, Liu ZJ. 2024. Molecular insights into self-
incompatibility systems: From evolution to breeding. Plant Commun 5:100719.

 **FIGURES AND FIGURE CAPTIONS**

**Fig. 1.** Oocysts genotype and developmental dynamics within the mosquito vector. (A)
Representative fluorescence images of mosquito midguts showing oocyst genotypes. Oocysts
expressing mCherry appear red, those expressing GFP appear green, and dual-labeled oocysts
appear orange in the merged image. (B) Temporal dynamics of oocyst burden per mosquito
across days post-infection (DPI). Panels from left to right show the number of red (mCherry-only),
green (GFP-only), and orange (dual-labeled) oocysts. Top panels show data from *Cross 1*, and
bottom panels show data from *Cross 2*. See Fig. S2 for plots showing total numbers of oocysts.

**Fig. 2.** Genetic diversity during oocyst development within the mosquito vector. (A) Temporal
dynamics of oocyst genotypes, shown as the proportions of red (mCherry-only, top bars), green
(GFP-only, middle bars), and orange (dual-labeled, bottom bars) oocysts across days post-
infection (DPI). The number of infected mosquitoes and the total oocyst count for each DPI are
indicated above the corresponding panel. We observed significant reduction in proportions of
outcrossed oocysts over time in both crosses (Cross 1: $R^2$ = 0.964, $p$ = 4.84E-4; Cross 2: $R^2$ =
0.902, $p$ = 3.10E-4; see also Fig. S3). (B) Regression analysis of inbreeding coefficient ($F_{IS}$) over
time, with 95% confidence intervals indicated by vertical lines. (C) Model illustrating how faster
maturation of heterozygous oocysts may lead to an outcrossing advantage.

A

| cross | term | estimate | se | p value |
|---|---|---|---|---|
| | Intercept | -18,342.80 | 5,501.85 | 8.68E-04 |
| | DPI | 5,820.76 | 542.20 | 2.38E-26 |
| *Cross1* | oocyst type:Orange | 9,542.15 | 1,279.22 | 1.17E-13 |
| | oocyst type:Red | 11,429.44 | 1,464.27 | 8.44E-15 |
| | oocyst No. per mosquito | -24.28 | 14.49 | 0.094 |
| | Intercept | 22,334.18 | 3,926.39 | 1.39E-08 |
| | DPI | 2,033.18 | 358.93 | 1.60E-08 |
| *Cross2* | oocyst type:Orange | -2,183.03 | 1,852.59 | 0.239 |
| | oocyst type:Red | 12,490.58 | 1,248.80 | 3.15E-23 |
| | oocyst No. per mosquito | -159.60 | 41.16 | 1.08E-04 |

**Fig. 3.** Analysis of factors influencing oocyst size. (A) Multiple regression analysis assessing the effects of oocyst genotype, days post-infection (DPI), and oocyst burden per mosquito on oocyst size across genetic crosses. (B) Box plots showing oocyst size variation by DPI and oocyst genotype across genetic crosses. For each DPI, boxes are ordered from left to right as green (NHP4026$^{GFP}$ selfed), orange (outcrossed), and red (NF54$^{mCherry}$ selfed). Asterisks indicate significant differences compared to the green group ($p < 0.05$).

**Table 1.** Summary of previous genotyping of cloned blood stage progeny. Cloned progeny were sequenced as part of the Malaria Crosses Collaboration project (https://mcc.nd.edu/) and sequences are available in GitHub repository (https://github.com/emilyli0325/P01-cloned.progeny) (Li, et al. 2022). Progeny were classified as follows: (1) those with ≥90% IBD with one parent were considered selfed; (2) those with <90% IBD with both parents were classified as recombinants; and (3) recombinant progeny sharing ≥90% IBD with each other were defined as non-unique clones.

| Study | Cross | No. of Clones | Progeny Types | | | No. of Unique Recombinant |
|---|---|---|---|---|---|---|
| | | | NF54 Selfed | NHP4026 Selfed | Recombinant | |
| Button-Simons et al., 2021 | NF54 x NHP4026 (before 2018) [a] | 146 | 1 | 0 | 145 | 88 |
| Amambua-Ngwa et al., 2023 | NF54 x NHP4026 (2019) | 158 | 2 | 0 | 156 | 113 |
| Unpublished | NF54$^{mCherry}$ x NHP4026$^{GFP}$ (2021) | 72 | 2 | 1 | 69 | 69 |

a. Multiple cloning attempts (between 2013 and 2018) of the same genetic cross conducted in 2013.

**Table 2.** Summary of previous genotyping studies of oocysts (mosquito stage).

| Study | Location | No. of Oocyst | Study details | $F_{IS}$ (inbreeding in mosquitoes) | $F_{IT}$ (inbreeding in total population | Heterozygote deficit |
|---|---|---|---|---|---|---|
| Ranford-Cartwright et al., 1992 | Laboratory | 98 | Cross between 3D7 and HB3 | nd | -0.143 | No |
| Babiker et al., 1994 | Tanzania | 71 | Field caught mosquitoes | nd | 0.389 | Yes |
| Paul et al., 1995 | New Guinea | 128 | Field caught mosquitoes | nd | 0.915 (0.835-0.966) [a] | Yes |
| Anthony et al., 2007 | Tanzania | 35 | Field caught mosquitoes | 0.16 (0.10-0.22) | nd | Yes |
| Annan et al., 2007 | Cameroon and Kenya | 221 | Field caught mosquitoes | 0.18 (0.14-0.23) for *A. gambiae*, 0.23 (0.17-0.28) for *A. funestus* | 0.47 (0.43–0.51) for *A. gambiae*, 0.60 (0.57–0.64) for *A. funestus* | Yes |
| Razakandrainibe et al., 2005 | Kenya | 553 | Field caught mosquitoes | 0.147 (0.062-0.209) | 0.43 (0.37-0.48) | Yes |
| Morlais et al., 2015 | Laboratory | 623 | Mosquitoes infected with blood donors | 0.28 (0.234 - 0.339) | nd | nd |
| Soontarawirat et al., 2017 | Thailand | 131 | Mosquitoes infected with *P. vivax* infected blood through membrane feeder | 0.55 | 0.89 | Yes |

a. Technical issues may result in underestimation of heterozygote frequency and overestimation of $F_{IS}$. *nd*, no data. Numbers in brackets indicate 95% confidence intervals.

**Table 3. Summary of mosquito infection data from genetic crosses between NF54$^{mCharry}$ and NHP4026$^{GFP}$.**

| Cross | DPI | Infected/Total Mosquitoes | Total Oocyst No. | Oocyst No. per Infected Mosquito Median (Range) | HWE *p-value* | *F*-statistics (95% CI) | | |
|---|---|---|---|---|---|---|---|---|
| | | | | | | $F_{IS}$ | $F_{ST}$ | $F_{IT}$ |
| Cross 1 | 8 | 15/17 | 342 | 21(3-68) | 0.085 | -0.110 ([-0.240]-[-0.030]) | 0.019 (0.007-0.073) | -0.089 ([-0.199]-0.013) |
| | 9 | 48/62 | 1374 | 20.5(1-137) | 0.388 | -0.034 ([-0.102]-0.002) | 0.011 (0.014-0.040) | -0.022 ([-0.074]-0.030) |
| | 10 | 34/36 | 1164 | 29.5(1-124) | 4.82E-06 | 0.130 (0.058-0.176) | 0.006 (0.007-0.037) | 0.135 (0.077-0.192) |
| | 11 | 28/35 | 486 | 11.5(1-79) | 3.49E-03 | 0.115 ([-0.003]-0.182) | 0.023 (0.020-0.091) | 0.135 (0.039-0.225) |
| | 12 | 22/31 | 195 | 9(1-19) | 8.65E-05 | 0.273 (0.078-0.351) | 0.015 (0.021-0.157) | 0.284 (0.164-0.411) |
| | 13 | 9/17 | 52 | 4(1-17) | 0.039 | 0.322 ([-0.067]-0.548) | -0.046 ([-0.066]-0.222) | 0.291 (0.004-0.577) |
| Cross 2 | 7 | 44/45 | 1230 | 28(4-53) | 1.83E-05 | 0.111 (0.032-0.153) | 0.013 (0.016-0.051) | 0.123 (0.063-0.180) |
| | 8 | 38/41 | 1025 | 24.5(7-57) | 2.23E-07 | 0.162 (0.082-0.213) | 0.000 (0.005-0.038) | 0.162 (0.100-0.229) |
| | 9 | 36/37 | 862 | 22(2-53) | 1.88E-05 | 0.146 (0.054-0.199) | 0.001 (0.008-0.045) | 0.146 (0.072-0.219) |
| | 10 | 34/35 | 845 | 23.5(4-62) | 5.74E-19 | 0.291 (0.190-0.348) | 0.023 (0.026-0.079) | 0.307 (0.231-0.382) |
| | 11 | 36/39 | 439 | 8(1-48) | 2.03E-06 | 0.214 (0.062-0.280) | 0.018 (0.030-0.121) | 0.228 (0.123-0.337) |
| | 12 | 34/41 | 261 | 7(1-27) | 1.03E-12 | 0.421 (0.238-0.505) | 0.039 (0.045-0.184) | 0.444 (0.329-0.560) |
| | 13 | 31/39 | 119 | 3(1-11) | 2.15E-08 | 0.490 (0.132-0.577) | 0.053 (0.090-0.350) | 0.517 (0.335-0.675) |
| | 14 | 26/34 | 146 | 4(1-28) | 1.67E-15 | 0.657 (0.437-0.762) | 0.011 (0.012-0.285) | 0.661 (0.517-0.802) |