TITLE
Activation of PP2A-B56α leads to aberrant EGFR signaling and proliferative phenotypes in PDAC

AUTHORS AND AFFILIATIONS

Claire M. Pfeffer[1, 2], Sydney J. Clifford[1], Saadia A. Karim[3], Ella Rose D. Chianis[1], Brittany N. Heil[1, 2], Lauren E. Gartenhaus[1, 2], Emma F. Kay[1], Garima Baral[1], Elizabeth G. Hoffman[1], Sagar M. Utturkar[4], Harish Kothandaraman[4], Whitney Smith-Kinnaman[5], Kasi Hansen[5], Emily G. Smith[1], Elisabeth Porter[6], Fuping Zhang[7], Nadia Atallah Lanman[4,8], Jukka Westermarck[9], Emma H. Doud[5,10], Amber L. Mosley[5,10], Jennifer P. Morton[3,11], Brittany L. Allen-Petersen[1, 2, 4] *

[1]Department of Biological Sciences, Purdue University, West Lafayette, IN USA
[2]Purdue University Interdisciplinary Life Sciences Program (PULSe), West Lafayette, IN USA
[3]Cancer Research UK Scotland Institute, Glasgow, UK
[4]Purdue Institute for Cancer Research, West Lafayette, IN USA
[5]Center for Proteome Analysis, Indiana University School of Medicine (IUSM), Indianapolis, IN, USA
[6]Department of Health and Human Sciences, Purdue University, West Lafayette, IN USA
[7]Turku Center for Disease Modeling (TCDM), Institute of Biomedicine, University of Turku, Turku, Finland
[8]Department of Comparative Pathobiology, Purdue University, West Lafayette, IN, USA
[9]Turku Bioscience Centre, University of Turku and Åbo Akademi University, Turku, Finland
[10]Biochemistry and Molecular Biology, Indiana University School of Medicine, (IUSM), Indianapolis, IN, USA
[11]School of Cancer Sciences, University of Glasgow, Glasgow, UK

*Corresponding Author

Brittany L. Allen-Petersen, Purdue University. 201 S. University St, West Lafayette, IN 47907.
Phone: 765-496-1896 Email: ballenpe@purdue.edu

RUNNING TITLE
PP2A regulates EGFR in PDAC

KEYWORDS
PDAC, PP2A, EGFR, AREG

ABSTRACT

Pancreatic ductal adenocarcinoma (PDAC) stands to become the second most deadly cancer by 2030. The small GTPase, KRAS, is mutated in over 90% of PDAC patients and considered the primary driver mutation. Despite being an almost ubiquitous event, KRAS mutations have been difficult to target therapeutically, particularly KRAS^{G12D}, the most common mutation in PDAC. In addition to these pharmacological challenges, KRAS mutations have been shown to drive signaling plasticity and therapeutic resistance through phosphorylation cascades in most cancers. Protein phosphatases are master regulators of kinase signaling, however the contribution of phosphatase deregulation to mutant KRAS cancer phenotypes is poorly understood. Protein phosphatase 2A (PP2A) inhibits effectors downstream of KRAS, placing this family of enzymes as key regulators of PDAC oncogenic signaling. However, our previous studies utilizing small molecule activating compounds of PP2A show a heterogeneous response in PDAC, with some cell lines displaying increased oncogenic signaling despite induction of phosphatase activity. Similarly, specific PP2A subunits exhibit both tumor suppressive and oncogenic functions depending on the cellular context. Therefore, understanding the role of PP2A in regulating cancer phenotypes is critical for the future development of therapeutic strategies that leverage this phosphatase. Here, we determined the impact of the specific PP2A subunit, B56α, on PDAC phenotypes using both genetic and pharmacological activation strategies in human PDAC cell lines and genetic mouse models. We demonstrate that while PP2A-B56α suppresses specific oncogenic pathways, B56α activation exacerbates PDAC proliferative phenotypes and decreases overall survival *in vivo*, potentially through increased epidermal growth factor receptor (EGFR) signaling. EGFR is a critical signaling node in PDAC as inhibition or loss of EGFR prevents KRAS-driven tumorigenesis and increased EGFR activity is associated with poor patient outcome. The activation of EGFR by PP2A-B56α is in part mediated through increased expression and processing of EGFR ligands, specifically amphiregulin, HB-EGF, and epiregulin. Furthermore,

pharmacological PP2A activation in combination with EGFR inhibitors mitigates this signaling and

increases cell death. Together, these studies implicate a previously undescribed non-canonical

role for PP2A-B56α in EGFR signaling that contributes to PDAC progression.

INTRODUCTION

Pancreatic cancer is the 3[rd] leading cause of cancer-related deaths in the United States and has a five-year survival rate of just 13%[1]. Over 90% of pancreatic ductal adenocarcinoma (PDAC) patients have driver mutations in the small GTPase, KRAS[2]. The aberrant activation of KRAS signaling has been shown to alter a wide range of signaling cascades leading to highly aggressive and therapeutically refractory disease[2–4]. As there are limited therapeutic options for PDAC patients, the identification of signaling mechanisms that contribute or regulate KRAS-driven PDAC is of the utmost importance.

Protein phosphatases are master regulators of cellular signaling cascades but their role in cancer remains poorly understood. Protein phosphatase 2A (PP2A) is a large family of Ser/Thr phosphatases that negatively regulate many oncogenic pathways including KRAS[5–7]. PP2A is a heterotrimeric holoenzyme which is composed of an A (scaffolding) subunit, B (regulatory) subunit, and C (catalytic) subunit. Therefore, the function and targets of PP2A vary depending on which specific B subunit is incorporated into the complex. There are four distinct PP2A B subunit families (B/B55, B'/B56, B''/PR72, B'''/STRN) with multiple genes and splice variants in each family. While there is some functional redundancy amongst the B subunits, the impact of individual B subunits is context specific. The B56α subunit has been implicated to have tumor suppressor function with loss of B56α driving tumorigenic phenotypes[8,9]. *In vivo*, we have shown that the loss of B56α accelerates the formation of KRAS[G12D]-driven PDAC precursor lesions suggesting that B56α plays a critical role in suppressing PDAC initiation[10]. Based on these findings, we hypothesize that activation of PP2A-B56α may negatively regulate PDAC-associated oncogenic pathways and attenuate PDAC tumor progression. In support of this hypothesis, pharmacological activation of PP2A decreases tumor burden in multiple preclinical models, including mutant KRAS lung and pancreatic cancer[11–17]. The small molecule activator of PP2A (SMAP), DT-061, globally

activates PP2A but displays a preferential stabilization of B56α-containing heterotrimeric PP2A complexes[18].

Here, we sought to understand the role of PP2A-B56α in regulating late-stage PDAC progression. In contrast to the tumor suppressive roles of B56α, we have identified a novel feedback mechanism in a subset of PDAC cell lines in which PP2A-B56α activation drives Epidermal Growth Factor Receptor (EGFR) signaling through upregulation of EGFR ligands. This activation exacerbates proliferation in PDAC cell lines which can be attenuated with loss of EGFR. Similarly, the conditional loss of the PP2A B56 inhibitor, Cip2a, in an aggressive PDAC mouse model significantly decreases overall survival and drives increased EGFR phosphorylation. Finally, the combination of PP2A activators and EGFR inhibition is synergistic *in vitro* and *in vivo*. Together these findings support a potentially adverse feedback loop through EGFR in response to PP2A-B56α activation in PDAC.

RESULTS

Loss of B56α reduces proliferation in PDAC cell lines

As B56α has been implicated as a B subunit contributing to PP2A's tumor suppressive functions, we generated B56α knockout PDAC cell lines (AsPc1, MiaPaCa2, and Panc1) and assessed the impact on cell proliferation. Cells were first transduced and selected for stable expression of Cas9. Single cell Cas9 clones were then transduced with two independent CRISPR guide RNAs (sgB56α1 and sgB56α2) or a non-targeting (sgNT) guide. Single cell clones from selected cells were then screened for B56α loss by western blot (Figure 1A) and proliferation and clonogenic colony formation were then evaluated. Unexpectedly, loss of B56α led to a significant reduction in proliferation over 96 hours compared to sgNT control (Figure 1A). Similarly, we found that B56α loss resulted in a reduced ability to grow in a clonogenic colony formation assay relative to sgNT control (Figure 1B). Together, these results suggest that the loss of B56α attenuates the

proliferative capacity of PDAC cell lines. Given B56α's known tumor suppressive roles, these findings highlight a potentially contradictory role for B56α in PDAC.

Overexpression of B56α increases tumorigenic phenotypes in PDAC cell lines

Given that loss of B56α reduced proliferation, we then wanted to determine if increased B56α expression would result in the converse phenotype. AsPc1, MiaPaCa2, and Panc1 cell lines with stable overexpression of B56α (B56αOE) or empty vector (EV) were generated and B56α expression was confirmed by RT-qPCR and western blot (Supplemental Figure 1A, B). In contrast to B56α knockout phenotypes, B56αOE PDAC cells displayed a significant increase in proliferation, clonogenic colony growth, and soft agar colony growth relative to the EV control (Figure 2A-C). Given that there is functional redundancy amongst the PP2A B subunits, we asked if overexpression of the other subunits would recapitulate B56α overexpression phenotypes. To this end, we generated stable overexpression cell lines in AsPc1 with three B56 family subunits (B56γ, B56δ, and B56ϵ) as well as B55α (Supplemental Figure 1C). In contrast to B56α, the overexpression of B55α, B56γ, B56δ, or B56ϵ did not significantly alter proliferation or clonogenic colony growth compared to EV controls (Figure 2D, E), suggesting that these phenotypes are specific to B56α in PDAC. Finally, to determine if the increased tumorigenic phenotypes with B56α overexpression leads to similar tumor phenotypes in cancers driven by similar mutations, we created stable B56α overexpression lines in KRAS-mutant non-small cell lung cancer (NSCLC) cell lines (Supplemental Figure 1D). In the three NSCLC lines assayed (A549, H23, Calu6), overexpression of B56α failed to increase proliferation relative to EV control (Supplemental Figure 1E), suggesting that cancer cells do not respond uniformly to B56α expression. Together, these results contrast the known tumor suppressive roles of PP2A-B56α, and potentially points to a unique role for B56α in late-stage PDAC.

PP2A-B56α activation enriches for cell cycle and EGFR signaling

To identify the impact of PP2A activation on transcriptional programs, we treated a panel of PDAC cell lines (AsPc1, Panc1, and HPAFII) with DT-061 for 24 hours and performed RNAseq. PP2A-B56α is a direct negative regulator of the oncoprotein c-MYC (MYC), with PP2A-B56α-driven dephosphorylation of MYC at S62 resulting in MYC degradation[17,19,20]. Therefore, as a read-out of PP2A activity, we analyzed the expression of MYC target genes using gene set enrichment analysis (GSEA)[21–23]. Within the genes that become downregulated in response to DT-061, there was a significant enrichment of MYC target genes across all three cell lines, consistent with PP2A activation (Supplemental Figure 2A). In contrast to MYC, there were several datasets that were significantly enriched within genes that become upregulated in response to DT-061 treatment (Supplemental Table 1). Of the top 20 significantly enriched gene sets, seven were common amongst the three cell lines, including genes that are upregulated in response to EGFR activation (EGFR_UP.V1_UP)(Figure 3A), suggesting that PP2A activation may exacerbate EGFR signaling. EGFR expression is known to increase during PDAC initiation and progression, and is overexpressed in PDAC compared to normal healthy pancreas[24,25]. Although EGFR is upstream of constitutively active mutant KRAS, loss of EGFR in pancreas-specific mutant KRAS PDAC mouse models (KRAS[G12D] and KRAS[G12V]) prevents pancreatic tumorigenesis[24,25], implicating EGFR signaling as a critical signaling node in PDAC. Furthermore, EGFR overexpression occurs in 40-90% of patients[25–27] and correlates with poor patient surviva[28,29] highlighting the importance of EGFR signaling in PDAC. DT-061 preferentially stabilizes PP2A complexes that contain the specific B subunit, B56α[30]. Therefore, to confirm that the enrichment of EGFR transcriptional programs is due to B56α signaling, HPAFII cells were transfected with an siRNA pool to B56α (siB56α) or non-targeting siRNA (siNT) and subsequently treated with DT-061. Knockdown of B56α prevented the DT-061-driven increase in EGFR

activation genes, suggesting that the induction of EGFR signaling predominately functions through PP2A-B56α (Figure 3B, Supplemental Table 2).

As pharmacological activation of PP2A altered EGFR transcriptional programs, we then performed proteomics and phospho-proteomics on AsPc1 EV and B56αOE cells, as well as AsPc1 cells treated with DT-061 or vehicle. In accordance with the role of PP2A as a phosphatase, DT-061 treatment predominately decreased phosphorylation with 306 phospho-sites significantly downregulated compared to 38 upregulated phospho-sites (Supplemental Figure 2B). Similar to previous reports, proteins with decreased phosphorylation in response to DT-061 were enriched for Myc targets, mitotic spindle, and mRNA processing[11,31,32] (Supplemental Figure 2C). Similarly, overexpression of B56α resulted in the downregulation of 280 phospho-sites with similar enrichment (Supplemental Figure 2D, E). In contrast, 141 phospho-sites were upregulated in B56αOE cells. Consistent with the increased proliferation observed in B56αOE cells, proteins with increased phosphorylation were enriched for cell cycle programs and targets of cyclin dependent kinases (CDKs)(Supplemental Figure 2F, G).

Analysis of total proteomic changes identified that overexpression of B56α resulted in a significant downregulation of 371 proteins and upregulation of 378 proteins, while DT-061 treatment resulted in a significant downregulation of 202 proteins and upregulation of 163 proteins (Figure 3C, D). Similar to our RNAseq results, activation/overexpression of B56α resulted in a significant negative enrichment of MYC and E2F datasets (Supplemental Figure 2H,I). However, gene set enrichment analysis of oncogenic signature gene sets (C6) revealed a number of pathways upregulated in response to PP2A activation (Figure 3E,F). Within the top 17 significant datasets, five were found to be enriched in both B56αOE and DT-061 conditions, including EGFR and ERBB2 activation signatures. Together, these results suggest that while activation of PP2A-

B56α is able to suppress known PP2A regulated pathways such as MYC, it also aberrantly increases EGFR signaling.

PP2A-B56α increases EGFR ligand expression and availability to drive EGFR signaling

In PDAC, EGFR is rarely mutated and instead is commonly amplified or upregulated[29,33]. Therefore, EGFR ligands play a critical role in the activation of EGFR signaling during PDAC progression. Highlighting the necessity of ligand-driven EGFR signaling, loss of A Disintegrin and Metalloprotease 17 (ADAM17), which is required for processing five of the seven EGFR ligands, prevents mutant KRAS pancreatic tumorigenesis[25]. Furthermore, pancreata from KRAS mutant mice have increased mRNA expression of the EGFR ligands, transforming growth factor alpha (TGFα) and amphiregulin (AREG), relative to wildtype mice[25]. Given the critical roles of EGFR ligands in activating EGFR in PDAC, we asked if activation of PP2A-B56α increased EGFR ligand availability. AsPc1 cells were treated with DT-061 or vehicle for 6 hours, and the media was subsequently changed to remove the unbound drug and allowed to condition for 48 hours. Conditioned media was then filtered, added to AsPc1 parental cells for 24 hours, and then cell lysates were analyzed via western blot (Figure 4A). Cells treated with conditioned media from DT-061 treated cells had increased levels of phosphorylation at EGFR Tyr1068 (Y1068), a critical posttranslational modification that mediates downstream signaling, relative to vehicle conditioned media (Figure 4B, C). Similar results were obtained with conditioned media derived from AsPc1 B56αOE or EV cells added to AsPc1 EV cells for 1.5 hours (Figure 4D-F). Together, these findings indicate that both pharmacological and genetic activation of PP2A-B56α results in increased soluble factors capable of activating EGFR. To identify which EGFR ligands may contribute to PP2A activation, a panel of PDAC cell lines was treated with DT-061 for 12 hours and mRNA expression of EGFR ligands was measured. While DT-061 increased the expression of several

EGFR ligands, AREG, epiregulin (EREG), and Heparin Binding EGF-like Growth Factor (HB-EGF) displayed the greatest fold increase relative to vehicle treated controls (Figure 4G).

Of all the EGFR ligands, AREG displayed the greatest increase in mRNA expression in response to PP2A activation. Increased expression of AREG has been shown to correlate with poor patient outcome, aggressive tumor phenotypes, and therapeutic resistance across several cancer types [34–36]. In PDAC, AREG expression is increased in primary pancreatic tumors relative to healthy tissue, and in KRAS mutant PDAC mouse models compared to wild-type mice[25,37]. Additionally, AREG has been proposed as a diagnostic marker for distinguishing if pancreatic cysts are benign or malignant[38]. Furthermore, it has been recently identified that AREG signaling in PDAC cancer associated fibroblasts (CAFs) promotes a premetastatic environment[39]. Therefore, to identify if PP2A activation results in increased AREG shedding, AREG was quantified by ELISA from media derived from PDAC cells with either B56αOE or treated with DT-061. In AsPc1 and MiaPaCa2 cells, both genetic overexpression and pharmacological activation of PP2A resulted in a significant increase in AREG shedding (Figure 4H, I; Supplemental Figure 3A, B). In contrast, PP2A activation was unable to alter AREG shedding in either Capan2 or HPAFII cells (Supplemental Figure 3C-E), suggesting that there are subsets of PDAC cell lines that are more sensitive to alterations in PP2A signaling. Concurrently, EGF, HB-EGF, and EREG ELISAs were performed on the same conditioned media; however, all three ligands were below detectable values (<6.25pg/ml EREG, 3.9pg/ml EGF, and 16pg/ml HB-EGF). Together, these data suggest that PP2A-B56α activation induces AREG expression and processing.

Increased proliferation in response to B56α overexpression is driven by AREG-EGFR signaling

Our data suggest that B56α overexpression in PDAC cells increases proliferation potentially through the activation of the AREG-EGFR signaling axis (Figure 1-4). To determine if

EGFR is necessary for the B56α-dependent increase in PDAC proliferation, EGFR was knocked down using siRNA (Supplemental Figure 4A, B), and proliferation and migration were assessed. Over the course of 96 hours, knockdown of EGFR had no effect on the proliferation rates of EV cells (EV siEGFR vs EV)(Figure 5A). In contrast, EGFR inhibition was able to prevent the increased proliferation rates seen in B56αOE AsPc1 cells (B56αOE siEGFR vs B56αOE), with no significant difference observed between B56αOE siEGFR and EV cells (Figure 5A). Similar results were observed in a transwell migration assay, where B56αOE increased the migratory capacity of AsPc1 cells, which was significantly attenuated with knockdown of EGFR relative to siNT controls (Figure 5B, C). These data suggest that the B56α overexpression mediates tumorigenic phenotypes in part through EGFR signaling. Given that AREG shedding was increased with B56α overexpression (Figure 4), we then assessed the contribution of AREG to B56αOE proliferation rates. Similar to siEGFR, knockdown of AREG (siAREG) had no significant effect on AsPC1 EV cells. However, the suppression of AREG prevented the increased proliferation seen in B56αOE cells, returning proliferation rates back to baseline EV levels (Figure 5D, Supplemental Figure 4C). Knockout of B56α reduced PDAC proliferation (Figure 1). Therefore, we determined if the addition of exogenous AREG could restore the growth of the B56α knockout cells. Previous studies determined that 300pg/ml AREG potentially represents a diagnostic threshold for malignant PDAC cysts[38]. The addition of 300pg/ml AREG[28] restored proliferation of AsPc1 B56α knockout cells back to levels seen in sgNT control conditions but had no effect on EV cells (Figure 5E). Together, these data suggest that the AREG-EGFR axis functions as a critical signaling pathway underlying the increased tumorigenic phenotypes that occur in response to B56α overexpression.

Genetic loss of the PP2A inhibitor, CIP2A, reduces overall survival in a model of PDAC *in vivo*

Previous studies have determined that knockdown of Cancerous inhibitor of PP2A (CIP2A), an endogenous cellular inhibitor of the PP2A B56 family[40], decreases PDAC tumor growth and MYC signaling in Panc1 and Capan1 cells[41]. These results are consistent with our findings that PP2A activation suppresses MYC transcriptional programs (Figure 3). However, given that B56αOE cells also display an increase in EGFR signaling, we asked whether the chronic activation of PP2A-B56 in response to CIP2A loss would alter PDAC progression *in vivo*. For these studies, we generated a conditional CIP2A knockout mouse model of PDAC by crossing *PDX1-Cre*, *Kras^{LSL-G12D}*, *Tp53^{LSL-R172H}* (KPC) mice with *Cip2a^{fl/fl}* (Cip2a^{fl/fl}) mice to generate KPCC (*PDX1-Cre*, *Kras^{LSL-G12D}*, *Tp53^{LSL-R172H}*, *Cip2a^{fl/fl}*) mice. KPC mice are considered the gold standard genetically engineered mouse model for PDAC, with KPC mice recapitulating both metastatic tumor progression and desmoplastic microenvironment found in PDAC patient tumors[42–44]. The genetic loss of this potent PP2A-B56 inhibitor accelerated PDAC tumorigenesis leading to a significant decrease in overall survival, with a median KPC survival of 148 days compared to 105 days in KPCC mice (Figure 6A). Furthermore, consistent with our *in vitro* findings, there was a significant increase in phosphorylated Y1068 EGFR in KPCC pancreatic tissues compared to KPC (Figure 6B, C). These results suggest that the chronic activation of PP2A exacerbates PDAC progression potentially through elevated EGFR signaling.

Pharmacological PP2A activation functions synergistically with EGFR inhibition

EGFR signaling significantly contributes to the activation of oncogenic cascades that drive proliferation and survival in PDAC[24,25,45–47]. Therefore, it is possible that increased EGFR signaling in response to PP2A activation may attenuate the tumor suppressor functions of PP2A. In the current studies, we have determined that EGFR signaling is necessary for B56α-driven increase in PDAC cell proliferation. Therefore, we hypothesized that EGFR inhibitors may suppress the feedback loop between PP2A-B56α and EGFR, decreasing PDAC viability. To identify if the

combination of PP2A activation and EGFR inhibition is efficacious, a drug matrix with DT-061 and the HER family inhibitor, Afatinib, was performed in 5 PDAC cell lines. SynergyFinder+ was used to calculate synergy between DT-061 and Afatinib and bliss synergy model was used[48]. The combination of DT-061 and Afatinib was synergistic in AsPc1, Panc1, and MiaPaCa2, but not in Capan2 or HPAFII (Figure 7A, B, Supplemental Figure 5A-C). We then tested if the combination of DT-061 and Afatinib was able to reduce tumor growth $in\ vivo$. Briefly, 2M MiaPaCa2 cells were injected subcutaneously into NOD.Cg-Rag1tm1Mom Il2rgtm1Wjl/SzJ (NRG mice). Once the average tumor size reached 100mm^3, mice were randomly enrolled into four treatment arms: Vehicle, DT-061 (15mg/kg BID), Afatinib (15mg/kg QD), or Combination. Supporting our $in\ vitro$ results, the combination of DT-061 and Afatinib significantly reduced tumor burden (Figure 7A) and endpoint tumor weight (Figure 7B), no significant impact on mouse body weight (Supplemental Figure 5D). In whole cell tumor lysates, DT-061 treated tumors had a significant increase in phospho-Y1068 EGFR (pEGFR) expression levels compared to vehicle (Figure 7E). As expected, Afatinib effectively inhibited pEGFR, with no detectable phosphorylation at Y1068, and was able to prevent DT-061-driven pEGFR in combination with DT-061 (Figure 7E). These results indicate that inhibition of EGFR increases the tumor suppressive capabilities of PP2A.

DISCUSSION

PP2A represents a critical and complex regulator of cellular signaling cascades. Given the diverse functions of this large phosphatase family, a deep mechanistic understanding of the roles that PP2A plays during oncogenesis is currently lacking. Here, we have identified a novel mechanism by which activation of the specific PP2A subunit, B56α, results in increased signaling through the AREG-EGFR axis to drive proliferative phenotypes in PDAC. These studies highlight the context specific functions of PP2A and support further exploration into defining patient populations that exhibit either tumor suppressive and tumor promotional PP2A phenotypes.

To our knowledge, this is the first report of increased B56α expression exacerbating oncogenic phenotypes and represents a critical shift in the way we approach phosphatase biology. However, other subunits have been reported as having contradictory roles in different cancer cell types[49–52] . B55α has been described as a tumor suppressive subunit in lung and breast cancer, but a previous report concluded that B55α was pro-tumorigenic in PDAC cell lines through shRNA-mediated knockdown experiments[53]. We were unable to recapitulate these findings with overexpression of B55α; however, differences in methods/approach could be a contributing factor. The increase in tumorigenic phenotypes in PDAC cell lines with B56α overexpression was not replicable in three KRAS-mutant NSCLC cell lines. While further studies are necessary in order to compare cancer specific phenotypes, these results support a potential differential role for B56α in NSCLC and PDAC, and broadly suggest that different type types or subsets of patients may display unique PP2A phenotypes. This concept becomes critically important in the context of therapeutic strategies that target PP2A.

Dynamic shifts in PP2A holoenzyme composition are poorly understood given the large number of B subunits and rapid signaling changes, and most likely play a significant role in determining the functional consequence of PP2A perturbations. As B56α expression increases, there may be other B subunits that are outcompeted or no longer incorporated into the PP2A holoenzyme. Conversely, it is unknown which B subunits are preferentially incorporated into the PP2A holoenzyme in response to B56α loss, if any. Given that our results for B56α overexpression and loss are complementary and coupled with pharmacological activation strategies give strength that B56α signaling alters EGFR pathways and tumorigenic phenotypes.

There was a significant increase in the amount of AREG in the media from DT-061 treated cells and B56αOE cell lines (Figure 4). We were unable to detect EGF, HB-EGF, or EREG from the same media so we are unable to conclude if other ligands are participating in activating EGFR with PP2A activation. Given that knockdown of AREG prevented the increased proliferation in

B56αOE cells, AREG may be necessary for the increased proliferation that occurs with B56α overexpression. AREG is a low affinity ligand and, therefore, be available in the media for longer amounts of time. Furthermore, the fate of EGFR downstream of AREG is poorly understood. Future studies interrogating the dynamics of EGFR internalization and cellular fate in response to PP2A activation will shed light on the differential responses observed between cell lines and cell types. Additionally, as EGFR can function in a homodimer or heterodimer, understanding the contribution of other HER family proteins to our phenotypes is critical.

In summary, our studies suggest that cellular context plays a significant role in determining whether PP2A-B56α effectively functions as a tumor suppressor. In contrast to global PP2A functions, our findings identify EGFR signaling as a critical mediator of PDAC proliferation in response to PP2A-B56α activation and support therapeutic strategies to mitigate these phenotypes.

CONFLICT OF INTEREST: All authors have no potential conflict of interest.

ACKNOWLEDGEMENTS: We would like to thank all the members of the BAP lab for their contributions to this work, editing of the manuscript, and providing constructive feedback. The Cip2a$^{fl/fl}$ mouse model was generated by Turku Center for Disease Modeling (www.tcdm.fi) supported by the Biocenter Finland. This study was supported by Purdue EVPRP NIH New R01 Program (BLA-P PI), Grace M. Showalter Research Trust (BLA-P); Pancreatic Cancer Action Network (BLA-P; 22-20-ALLE), Concern Foundation (BLA-P), and American Cancer Society (BLA-P; RSG-24-1259215-01-MM). BNH, ERDC, SJC, and EGS was supported by the Purdue Institute for Cancer Research (NIH grant P30 CA023168). CMP was supported by the Purdue Bilsland Dissertation Fellowship. The bioinformatics analyses were supported by the Purdue Institute for Cancer Research and the Walther Cancer Foundation. The mass spectrometry work performed in this manuscript was done by the Indiana University School of Medicine Center for

Proteome Analysis. Acquisition of the IUSM Center for Proteome Analysis instrumentation used for this project was provided by the Indiana University Precision Health Initiative. The proteomics work was supported, in part, by the Indiana Clinical and Translational Sciences Institute (Award Number UL1TR002529 from the National Institutes of Health, National Center for Advancing Translational Sciences, Clinical and Translational Sciences Award) and the P30 IU Simon Comprehensive Cancer Center Support Grant (Award Number P30CA082709 from the National Cancer Institute).

AUTHOR CONTRIBUTION: CMP and BLA-P designed experiments. CMP, SJC, SK, ERDC, BNH, LEG, EFK, GB, EGH, EGS, and EP performed and analyzed experiments. SMU, HK, and NAL performed RNA-seq data analysis. WS-K, KH, EHD, and ALM performed mass spectrometry sample preparation, data analysis, and methods Write-up. FZ and JW generated the Cip2a$^{fl/fl}$ mouse. SK and JPM generated KPC and KPCC survival curves and maintained the mouse colony. BLA-P secured funds and provided supervision. BLA-P and CMP wrote and revised the manuscript. All authors reviewed and finalized the manuscript.

METHODS

Cell culture

AsPc1, MiaPaCa2, HPAFII, Panc1, CfPac1, Capan1, Capan2, Colo357, and HEK293T were obtained from ATCC. PDAC cell lines were maintained in DMEM (Gibco 11995065) + 10% Fetal Bovine Serum (FBS) (Fisher Scientific FB12999102) at 37°C and 5% CO_2. NSCLC cell lines (A549, H23, and Calu6) were maintained in RPMI (Cytiva SH30027.01) + 10% FBS at 37°C and 5% CO_2. Lines were regularly tested for Mycoplasma using a PCR strategy. Viral particles were generated by transfecting HEK293T cells with the vector of interest and two lentiviral vectors (psPAX2 Addgene #12260 and pMD2.G Addgene #12259) using lipofectamine 3000 (Invitrogen L3000015). B56α knockout cell lines were generated by first generating Cas9 (Addgene #125592) expressing cells through clonal selection. Then the sgRNAs (sgNT: Horizon Discovery

GSG11817; sgB56α1: Horizon Discovery GSGH11838-246527026; sgB56α2: Horizon Discovery GSGH11838-246527024) were transduced into the Cas9 expressing cells, selected and then clonal selected and screened for B56α loss by western blot. Overexpression of each of the B subunits were generated by cloning the gene of interest into a modified pSin-PURO empty vector. The following vectors were used for the B subunits: 4HA-B56α (Addgene #14532), 4HA-B56γ1 (Addgene #14534), 4HA-B56δ (Addgene #14536), 4HA-B56ε (Addgene #14537), Flag-B55α (Addgene #13804). For transient knockdown experiments, cells were transfected using lipofectamine 3000 with 35nM siRNA (NT (Horizon Discovery D-001810-20), EGFR (Horizon Discovery L-003114-00-005), AREG (Horizon Discovery L-017435-00-005), or B56α (Horizon Discovery L-009352-00-0020).

Western blot

Cells were lysed and protein was quantified using DC Protein Assay Kit II (Bio-Rad 50000112). Equally loaded samples were separated on a 4-12% Criterion XT Bis-Tris Protein Gel (Bio-Rad 3450123) and transferred onto a PVDF membrane (Millipore IPFL00010). Membrane was blocked using Intercept (PBS) Blocking Buffer (Licor 927-70001) for 1 hour at room temperature. Primary antibodies were incubated overnight at 4°C and secondary antibodies were incubated at room temperature for 1 hour (Table 1). Membranes were imaged with a LI-COR Odyssey and analyzed using Image Studio Lite.

RNA isolation/quantitative PCR

RNA was isolated using GeneJet RNA Purification Kit (ThermoScientific K0732) according to manufacturer's protocol. cDNA was generated using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems 4368814). Quantitative PCR was run using PowerUp SYBR Green Master Mix (Applied Biosystems A25741) using primers in Table 2 on a QuantStudio3. Fold change was calculated using the delta delta ct method and normalized to 18S.

Tumorigenic phenotypes

For proliferation, cells were seeded in technical triplicate and counted every 24 hours. Area under the curve was calculated and used for statistical significance. For soft agar assays, 12 well plates were coated with 500µL of a 1:1 mixture of 1.4% noble agar and 2X DMEM + 20% FBS. Based on the parental cell line, 30,000-50,000 cells were resuspended in a 1:1 mixture of 0.7% noble agar and 2X DMEM + 20% FBS and seeded on top of the 1.4% layer. After 14 days, cells were stained with 0.1% crystal violet and imaged using an EVOS M7000. Number of colonies was quantified using ImageJ. For migration, transwell migration assay was used. 1 hour prior to seeding, 8um pore inserts (Corning 3464) were soaked in serum-free DMEM. 20,000 cells were seeded in 8um inserts in serum-free DMEM and DMEM + 10% FBS was added to the bottom of the well. After 48 hours, cells were fixed and stained with 0.1% crystal violet. Cells that remained at the top of the insert and had not migrated were removed. Migrated cells were imaged with an EVOS M7000 and number of cells migrated was quantified using ImageJ. For clonogenic colony formation, 100-500 cells (depending on the parental cell line) were seeded in a 6-well plate. After 10-14 days (depending on parental cell line), cells were stained with 0.1% crystal violet and scanned on an EVOS M7000. Percent area occupied by colonies was quantified using ImageJ.

RNA sequencing

AsPc1, Panc1, and HPAFII cells were plated in DMEM +2% FBS in triplicate. 48 hours later, cells were treated with 10µM DT-061 or vehicle (DMSO) for 24 hours. RNA was isolated as described above. RNA-seq libraries were generated and Illumina sequenced by Novogene. Datasets were trimmed using fastp (v0.23.2) to remove bases with a Phred quality score lower than Q30 and adaptor sequences. Reads were aligned to GRCh38.p14 using STAR version 2.7.9a[55], and genes were quantified with subread featureCounts version 1.6.1[56] using the annotation from ensembl release 104. Initial exploratory analyses were performed with DESeq2 version 1.26.0[57] and differential gene expression analyses were performed with edgeR version 3.28.1[58] on the R Statistical Computing Project version 3.6.1[59] . Ranked lists were generated for each cell line and used for Gene Set Enrichment Analysis (GSEA) version 4.3.2.[22,23].

<u>Total and phospho proteomics</u>

$5X10^6$ AsPc1 cells were plated onto a 15cm plate in DMEM + 2% FBS. 48 hours later, cells were treated with 5mM DT-061 or vehicle for 24 hours. Cells were lifted, washed, and then pelleted and flash frozen. Sample preparation, mass spectrometry analysis, bioinformatics, and data evaluation for quantitative proteomics experiments were performed in collaboration with the Indiana University Proteomics Center for Proteome Analysis at the Indiana University School of Medicine similarly to previously published protocols[60,61].

Protein extraction and digestion

Cell pellets were resuspended in 100 μL 8 M Urea, 100 mM Tris pH 8.5. The resuspended cell pellets were transferred to Diagenode Bioruptor tubes (Cat No: C30010010). Cells were lysed via Diagenode Bioruptor, 30s on/30s off, for 30 cycles. Samples were incubated for 1 hour at 35 ºC with 2 units of benzonase (EMD Millipore Cat NO. 70746-4). Samples were then clarified by centrifuging for 20 min at 12,000 rcf. Supernatants were analyzed in a Bradford assay (Biorad Cat No: 5000002) to determine protein concentration. 520 μg of each sample was treated with 5 mM tris(2-carboxyethyl)phosphine hydrochloride (Sigma-Aldrich Cat No: C4706) to reduce disulfide bonds and the resulting free cysteine thiols were alkylated with 10 mM chloroacetamide (Sigma Aldrich Cat No: C0267). Samples were diluted with 50 mM Tris.HCl pH 8.5 (Sigma-Aldrich Cat No: 10812846001) to a final urea concentration of 2 M for overnight Trypsin Platinum digestion at 35 ºC (1:25 protease:substrate ratio, Mass Spectrometry grade, Promega Corporation, Cat No: VA9000).

Peptide purification and phosphopeptide enrichment

Digestions were quenched with formic acid (Fluka Cat No: 94318) and desalted on Waters Sep-Pak® Vac cartridges (Waters™ Cat No: WAT054955) with a wash of 1 mL 0.1% TFA followed by elution in 3x 0.2 mL of 70% acetonitrile 0.1% formic acid (FA). Peptides were dried by speed vacuum. Samples were resuspended in phosphopeptide binding buffer and phosphopeptides were enriched using Thermo Fisher Scientific High Select TiO2 tips (Cat No A32993) according

to manufacturer's instructions. Flow through (non-phosphopeptides) and phosphopeptides were dried down by speed vacuum.

TMTpro labeling

An equivalent of 50µg of the global (non-phospho) peptides and all of each phosphopeptide enrichment were resuspended in 100 mM triethylammonium bicarbonate (TEAB, pH 8.5 from 1 M stock). Each sample was then labeled overnight at room temperature, with 0.5 mg of Tandem Mass Tag Pro (TMTpro™) reagent (16-plex kit, manufactures instructions Thermo Fisher Scientific, TMTpro™ Isobaric Label Reagent Set; Cat No: 44520, lot no. ZA382395 see Table X below). Reactions were quenched with 0.3 % hydroxylamine (v/v) at room temperature for 10 minutes. Labeled peptides were then mixed and dried by speed vacuum.

High pH basic fractionation

Half of the combined global sample and all of the phosphopeptide sample were resuspended in 0.5% TFA and fractionated on a Waters Sep-Pak® Vac cartridge (Waters™ Cat No: WAT054955) with a 1 mL wash of water, 1 mL wash of 5% acetonitrile, 0.1% triethylamine (TEA) followed by elution for the global sample in 8 fractions of 12.5%, 15%, 17.5%, 20%, 22.5%, 25%, 30%, and 70% acetonitrile, all with 0.1% TEA).

Nano-LC-MS/MS

Mass spectrometry was performed utilizing an EASY-nLC 1200 HPLC system (SCR: 014993, Thermo Fisher Scientific) coupled to Eclipse™ mass spectrometer with FAIMSpro interface (Thermo Fisher Scientific). Each multiplex was run on a 25 cm Aurora Ultimate column (Ion Opticks Cat No: AUR3-25075C18) in a 50 °C column oven with a 180-minute gradient or 130-minute gradient. For each global fraction, 6.67% of the sample was loaded and run at 350 nl/min with a gradient of 10-40%B over 160 minutes; 40-90% B over 10 mins; held at 90% for 2 minutes; and dropping from 90-8% B over the final 5 min (Mobile phases A: 0.1% formic acid (FA), water; B: 0.1% FA, 80% Acetonitrile (Thermo Fisher Scientific Cat No: LS122500)). The mass spectrometer was operated in positive ion mode, default charge state of 2, advanced peak

determination on, and EasyIC on. Three FAIMS CVs were utilized (-45 CV; -55 CV; -65CV) each with a cycle time of 2 s and with identical MS and MS2 parameters. Precursor scans (m/z 400-1600) were done with an orbitrap resolution of 120000, RF lens% 30, 50 ms maximum inject time, standard automatic gain control (AGC) target, minimum MS2 intensity threshold of 2.5e4, MIPS mode to peptide, including charges of 2 to 6 for fragmentation with 60 sec dynamic exclusion shared across the cycles excluding isotopes. MS2 scans were performed with a quadrupole isolation window of 0.7 m/z, 34% HCD collision energy, 50000 resolution, 200% AGC target, dynamic maximum IT, fixed first mass of 100 m/z.

For each phosphopeptide fraction 25% of the sample was loaded and run at 350 nl/min with a gradient of 10-35%B over 120 minutes; 35-95% B over 5 mins; and dropping from 95- 6% B over the final 5 min (Mobile phases A: 0.1% formic acid (FA), water; B: 0.1% FA, 80% Acetonitrile (Thermo Fisher Scientific Cat No: LS122500)) with identical mass spectrometry parameters to the global proteomics samples.

Mass spectrometry data analysis

Resulting RAW files were analyzed in Proteome Discover™ 2.5.0.400 (Thermo Fisher Scientific)[62] with a *Mus musculus* UniProt reference proteome FASTA (downloaded 051322) plus common laboratory contaminants (73 sequences). SEQUEST HT searches were conducted with full trypsin digest, 3 maximum number missed cleavages; precursor mass tolerance of 10 ppm; and a fragment mass tolerance of 0.02 Da. Static modifications used for the search were: 1) TMTpro label on peptide N-termini 2) TMTpro label on lysine (K) and 3) carbamidomethylation on cysteine (C) residues. Dynamic modifications used for the search were 1) oxidation on M (2) phosphorylation on S, T or Y, 3) acetylation on protein N-termini, 4) methionine loss on protein N-termini or 5) acetylation with methionine loss on protein N-termini. A maximum of 3 dynamic modifications were allowed per peptide. Percolator False Discovery Rate was set to a strict setting of 0.01 and a relaxed setting of 0.05. IMP-ptm-RS node was used for all modification site localization scores. Values from both unique and razor peptides were used for quantification. In

the consensus workflows, peptides were normalized by total peptide amount with no scaling. Unique and razor peptides were used, and all peptides were used for protein normalization and roll-up. Quantification methods utilized TMTpro isotopic impurity levels available from Thermo Fisher Scientific. Reporter ion quantification filters were set to an average S/N threshold of 5 and co-isolation threshold of 30%. Resulting grouped abundance values for each sample type, abundance ratio values; and respective p-values (Protein Abundance based with ANOVA individual protein based) from Proteome Discover were exported to Microsoft Excel. Pathway enrichment was performed using Gene Set Enrichment Analysis (GSEA) version 4.3.2[21–23].

Conditioned media

2.5M AsPc1 B56αOE or EV cells were plated in a 10cm plate in DMEM +10% FBS. 48 hours later, media was changed to 5.5ml serum free DMEM and allowed to condition for 48 hours. Conditioned media was filtered through a 0.45μM filter and stored at 4°C until use. 2ml of conditioned media was added to AsPc1 parental cells 48 hours after seeding 200k on 35mm plates. After 90 minutes, cells receiving conditioned media were lysed and protein was analyzed as described in western blotting section. The same process was used with DT-061 treated cells with the following modifications: 2.5M AsPc1 cells were plated in a 10cm plate in DMEM + 10% FBS. 24 hours later, media was changed to serum free DMEM and 7.5μM DT-061 was added for 6 hours, then drug was washed out and serum free DMEM was allowed to condition for 48 hours. After 24 hours, cells receiving conditioned media were lysed and protein was analyzed as described in western blotting section.

ELISA

To collect media, 300,000 cells were seeded in a 6-well plate in DMEM + 10% FBS. 24 hours after seeding, cells were serum starved for 24 hours prior to adding DMEM to condition for 24 hours. Media was stored at -20°C until quantified for AREG (Invitrogen EHAREG), HB-EGF (Invitrogen EHHBEGF), EGF (Invitrogen KHG0061), or EREG (Abcam ab277077) according to

manufacturer's instructions. For the KPC/KPCC cell lines, AREG (Invitrogen EMAREG) was quantified and normalized to cell number for each biological replicate.

Rescue experiments

2M AsPc1 EV or B56αOE cells were seeded onto 10cm plate. 24 hours later, lipofectamine 3000 (Invitrogen, L3000015) was used to transfect 35nM siRNA (NT (Horizon Discovery D-001810-10-20), EGFR (Horizon Discovery L-003114-00-005), or AREG (Horizon Discovery L-017435-00-005)). After 24 hours later, the cells were lifted, and the rescue experiments (proliferation and migration) were plated as described above. Cells were taken at all time points to validate knockdown of target gene via qPCR.

Drug curves and matrices

7,000 cells were seeded and immediately dosed with Afatinib and DT-061. 72 hours later, cell viability was assessed using crystal violet[63]. Absorbance (570nm) was read on a Synergy H1 Microplate Reader (BioTek). Viability was calculated relative to vehicle control. 4 biological replicates were averaged, and synergy was calculated using the bliss synergy model in SynergyFinder+[48].

Animal studies

Animal experiments were performed in compliance with Purdue University Institutional Animal Care and Use Committee, University of Glasgow Animal Welfare and Ethics Review Body, and the Finnish ethical committee for experimental animals, complying with international guidelines on the care and use of laboratory animals.

Xenograft Studies: MiaPaCa2 cells (2×10^6) were injected subcutaneously into the flanks of NRG (NOD-*Rag1*[null] *IL2Rg*[null]) mice in a 1:1 ratio with Matrigel (Corning 356231). Once palpable, tumors were calipered every other day and tumor volume was calculated by Volume = (length x width^2)/2. Once average tumor volume was 100mm^3, mice were randomly enrolled into the four treatment arms. 15mg/kg DT-061 or vehicle (1:1:8 N,N dimethylacetamide (DMA): Kolliphor HS 15: PBS) were dosed two times daily via oral gavage. 15mg/kg Afatinib or vehicle (1:1:18 N,N

dimethylacetamide (DMA): Kolliphor HS 15: PBS) were dosed once daily by intraperitoneal injection. Mice were treated for 21 days and then mice were sacrificed. Tumors were halved and fixed in 10% formalin or snap frozen for qPCR and western blot. For histology, tumors were paraffin embedded and 6um sections were used H&E and immunofluorescence.

Cip2a^{fl/fl} mice: Generation of Cip2a^{fl/fl} mice was done in collaboration with the Turku Center for Disease Modeling (TCDM) at the University of Turku (Finland). LoxP sites were inserted into introns 1 and 2 using the following strategy: BAC clones containing mouse Cip2a gene (ENSMUSG00000033031) were obtained from BACPAC Resources Center (Children's Hospital Oakland Research Institute, Oakland (CHORI), California, USA). A 9696 bp genomic DNA fragment containing exons 1 to 8 of Cip2a gene was subcloned into the minimal vector, pACYC177 (New England Biolabs, MA, USA). To insert a single loxP site, a floxed PGK-Neo cassette (Gene Bridges) containing 50 bp Cip2a homology arms amplified by PCR was inserted into the intron 1 of Cip2a gene by Red/ET recombination method. Primers were 5'-TAGAGAAGGAAGAACTCTATATATGGGCAGAATGAACAAAAGATGTGAATGAAGCAGGGATT CTGCAAAC-3' and 5'-AAACCAATTTTCTAAATTGCTACTTAGCCCTTTCTTTCCCAGTTTAGA AAGGCGGATTTGTCCTACTCAGG-3'. Site-specific recombination was carried out *in vitro* using Cre recombinase in E.coli. (Gene Bridges). A second LoxP site was introduced into the intron 2 of Cip2a gene with Frt-PGK-gb2-Neo-Frt-loxP cassette (Gene Bridges) containing 50 bp Cip2a homology arms amplified by PCR. The primers were 5'- ATAGATAGAATTATA AAATTTCTTCAACAATGATTTACATTCAAAAGATTAATTAACCCTCACTAAAGGGCGG-3' and 5'-GTAGGAACTTTGAGGTATCTGTAATACTAACTCTAAAAGGTTTTACAATATAATACGACTCA CTATAGGG-3'. G4 embryonic stem cells (derived from mouse 129S6/C57BL/6Ncr) were electroporated with 30 μg of linearized targeting construct. In order to delete Neo cassette in the targeted ES cells, cells were re-electroporated with plasmid, pCAGGS-Flpe. Targeted ES cells

with Neo deletion were then injected into C57BL/6 (Charles River Laboratories) mouse blastocysts to generate chimeric mice. Germline transmission was achieved by cross-breeding male chimeras with C57BL/6 females. Cip2a$^{fl/fl}$ mice were maintained in a specific pathogen free stage at Central Animal Laboratory at the University of Turku before being transferred to the University of Glasgow. To generate KPCC mice, Cip2a$^{fl/fl}$ were backcrossed 10 generations prior to breeding with *PDX1-CRE; Kras$^{LSL-G12D}$; Trp53$^{LSL-R172H}$* mice. Both KPC and KPCC mice were monitored weekly and sacrificed when mice began to show signs of being moribund.

Immunofluorescence

Paraffin embedded tissue slides were incubated at 60^{oC} for 1 hour prior to progressive rehydration in xylene twice for 10 minutes, then 100% ethanol twice for 5 minutes, then 95% ethanol, 70% ethanol, PBS, and water for 5 minutes each. Antigen retrieval was performed with slides first covered with 1x high pH antigen retrieval solution (Invitrogen 00-4956-58) and cooked in a pressure cooker on high pressure for 10 minutes. The slides were then allowed to cool to room temperature and rinsed in water before being covered with 1x low pH antigen retrieval solution (Invitrogen 00-4955-58) and cooked as described above. After cooling to room temperature, the tissues were covered with 0.025% tween in PBS for five minutes. Each tissue was blocked with 3% BSA in PBS for 1 hour at room temperature before the addition of primary antibodies diluted in 3% BSA in PBS and incubation overnight at 4ºC. Tissues underwent three 5-minute washes with 0.025% tween in PBS before the addition of secondary antibodies which were incubated at room temperature for 1 hour. Secondary antibodies diluted in 3% BSA in PBS were followed with another series of three 5-minute washes in 0.025% tween in PBS. Dilutions of antibodies used are listed in (TABLE 2). DAPI was diluted in 3% BSA in PBS before incubation on the tissues for 5 minutes at room temperature. After an additional 5-minute wash in 0.025% tween in PBS, a glass coverslip was mounted to each slide with Prolong Gold (Invitrogen P36934). Coverslips were allowed to harden for at least 12 hours before microscopy. A Nikon Ni-U Upright Microscope

was used to take 20x images of each tissue. From each tissue, mean fluorescence intensity of

phospho-EGFR (pY1068) from four to eight fields of view was quantified using ImageJ.

Statistical Analysis

All experiments were repeated for at least three biological replicates unless otherwise noted. Error

bars represent standard deviation unless otherwise noted. For two groups, a student's two-tailed

unpaired t-test was performed using GraphPad Prism with 95% confidence interval. For groups

of three or more, a 1-way ANOVA

Table 1. Antibodies used.

Protein	Company	Catalog number	Lot number	Dilution
Western Blot				
EGFR	Cell Signaling	2646S	8	1:500
p-EGFR (Y1068)	Cell Signaling	2234L	22	1:500
AKT	Cell Signaling	2920S	8	1:1000
p-AKT (S473)	Cell Signaling	4058S	30	1:1000
HA	Cell Signaling	3724S	10	1:1000
c-MYC	Abcam	AB32072	1053331-47	1:1000
GAPDH	Invitrogen	AM4300	3035436	1:20,000
B56α	Made in Ogris lab			1:500
CIP2A	Santa Cruz	SC-80659	H2522	1:500
B55α	Cell Signaling	5689T	3	1:500
IRDye 680RD Goat Anti-Rabbit IgG Secondary	Licor	926-68071	D20809-05	1:5000

IRDye 800CW Goat Anti-Mouse IgG Secondary	Licor	926-32210	D20802-25	1:5000
Immunofluorescence				
p-EGFR (Y1068)	Cell Signaling	2234L	22	1:100
Pan Cytokeratin C11	Santa Cruz	sc-8018	J0721	1:1000
Goat anti-Rabbit IgG (H+L) Alexa Fluor Plus 647	Invitrogen	A32733	ZJ405179	1:200
Goat anti-Mouse IgG (H+L) Alexa Fluor Plus 488	Invitrogen	A32723	ZH395844	1:200

Table 2. Primers used for qPCR.

Gene	Species	Forward Sequence (5' → 3')	Reverse Sequence (5' → 3')
18S	Human	CACCAACATCgATgggCgg	CACACgTTCCACCTCATCCTCAg
PPP2R5A	Human	CTTTgCATTgCCACTgAAAg	CAgCAgTCCTCTgATCAC
PPP2R5B	Human	CTCCTCTCAgTTCCgCTATCA	gCCACACAgTCCAAgAAgTCAA
PPP2R5C	Human	gAAgATCCTCgggAgAgAgATT	TCTgCTATgCCATTATgATgCTC
PPP2R5D	Human	CCTTCATCgAATCCCACAggg	TgTTTggCTggAAATCAggAg
PPP2R5E	Human	TAggTCTCAAggCAAgCCTAT	CTgTCAAACAgCCTCTgCTTATT
B55a	Human	CATACCAggTgCATgAATACCTC	gggTTATgTCTCgCTTTgTgTTT
Areg	Human	gAgCCgACTATgACTACTCAgA	TCACTTTCCgTCTTgTTTTggg
BTC	Human	CCTgggTCTAgTgATCCTTCA	CTTTCCgCTTTgATTgTgTgg
EGF	Human	gTACTCTCgCAggAAATggg	TCCACCACCAATTGCTCATA
Ereg	Human	ATgTggCTTTgACCgTgATTC	TCCCCTgAggTAACTCTCTCATA
Epigen	Human	CTTTgggAgTTCCAATATCAgTC	ggCTATgggTCCTTCTATgTTg
HBEGF	Human	ATCgTggggCTTCTCATgTTT	TTAgTCATgCCCAACTTCACTTT
TGFα	Human	ggTgCAggAggACAAgCC	CACgATggAgACCACCACC

EGFR	Human	AggCACgAgTAACAAgCTCAC	ATgAggACATAACCAgCCACC

1. Siegel, R.L., Kratzer, T.B., Giaquinto, A.N., Sung, H., and Jemal, A. (2025). Cancer statistics, 2025. CA: A Cancer J. Clin. *75*, 10–45. https://doi.org/10.3322/caac.21871.

2. Yousef, A., Yousef, M., Chowdhury, S., Abdilleh, K., Knafl, M., Edelkamp, P., Alfaro-Munoz, K., Chacko, R., Peterson, J., Smaglo, B.G., et al. (2024). Impact of KRAS mutations and co-mutations on clinical outcomes in pancreatic ductal adenocarcinoma. npj Precis. Oncol. *8*, 27. https://doi.org/10.1038/s41698-024-00505-0.

3. Huang, L., Guo, Z., Wang, F., and Fu, L. (2021). KRAS mutation: from undruggable to druggable in cancer. Signal Transduct. Target. Ther. *6*, 386. https://doi.org/10.1038/s41392-021-00780-4.

4. Wang, S., Zheng, Y., Yang, F., Zhu, L., Zhu, X.-Q., Wang, Z.-F., Wu, X.-L., Zhou, C.-H., Yan, J.-Y., Hu, B.-Y., et al. (2021). The molecular biology of pancreatic adenocarcinoma: translational challenges and clinical perspectives. Signal Transduct. Target. Ther. *6*, 249. https://doi.org/10.1038/s41392-021-00659-4.

5. Arnold, H.K., and Sears, R.C. (2006). Protein Phosphatase 2A Regulatory Subunit B56α Associates with c-Myc and Negatively Regulates c-Myc Accumulation†. Molecular and Cellular Biology *26*. https://doi.org/10.1128/mcb.26.7.2832-2844.2006.

6. Arnold, H.K., Zhang, X., Daniel, C.J., Tibbitts, D., Escamilla-Powers, J., Farrell, A., Tokarz, S., Morgan, C., and Sears, R.C. (2009). The Axin1 scaffold protein promotes formation of a degradation complex for c-Myc. EMBO J. *28*, 500–512. https://doi.org/10.1038/emboj.2008.279.

7. Peris, I., Romero-Murillo, S., Vicente, C., Narla, G., and Odero, M.D. (2023). Regulation and role of the PP2A-B56 holoenzyme family in cancer. Biochim. Biophys. Acta (BBA) - Rev. Cancer *1878*, 188953. https://doi.org/10.1016/j.bbcan.2023.188953.

8. Sablina, A.A., Hector, M., Colpaert, N., and Hahn, W.C. (2010). Identification of PP2A Complexes and Pathways Involved in Cell Transformation. Cancer Res *70*, 10474–10484. https://doi.org/10.1158/0008-5472.can-10-2855.

9. Janghorban, M., Langer, E.M., Wang, X., Zachman, D., Daniel, C.J., Hooper, J., Fleming, W.H., Agarwal, A., and Sears, R.C. (2017). The tumor suppressor phosphatase PP2A-B56α regulates stemness and promotes the initiation of malignancies in a novel murine model. Plos One *12*, e0188910. https://doi.org/10.1371/journal.pone.0188910.

10. Tinsley, S.L., Chianis, E.R.D., Shelley, R.A., Mall, G.K., Dhiman, A., Baral, G., Kothandaraman, H., Thoma, M.C., English, I.A., Daniel, C.J., et al. (2024). KRAS-mediated upregulation of CIP2A promotes suppression of PP2A-B56α to initiate pancreatic cancer development. Oncogene *43*, 3673–3687. https://doi.org/10.1038/s41388-024-03196-w.

11. Allen-Petersen, B.L., Risom, T., Feng, Z., Wang, Z., Jenny, Z.P., Thoma, M.C., Pelz, K.R., Morton, J.P., Sansom, O.J., Lopez, C.D., et al. (2018). Activation of PP2A and inhibition of mTOR synergistically reduce MYC signaling and decrease tumor growth in pancreatic ductal adenocarcinoma. Cancer Res *79*, canres.0717.2018. https://doi.org/10.1158/0008-5472.can-18-0717.

12. Sangodkar, J., Perl, A., Tohme, R., Kiselar, J., Kastrinsky, D.B., Zaware, N., Izadmehr, S., Mazhar, S., Wiredja, D.D., O'Connor, C.M., et al. (2017). Activation of tumor suppressor protein PP2A inhibits KRAS-driven tumor growth. J Clin Investigation *127*, 2081–2090. https://doi.org/10.1172/jci89548.

13. Risom, T., Wang, X., Liang, J., Zhang, X., Pelz, C., Campbell, L.G., Eng, J., Chin, K., Farrington, C., Narla, G., et al. (2019). Deregulating MYC in a model of HER2+ breast cancer mimics human intertumoral heterogeneity. J Clin Invest *130*, 231–246. https://doi.org/10.1172/jci126390.

14. Tohmé, R., Izadmehr, S., Gandhe, S., Tabaro, G., Vallabhaneni, S., Thomas, A., Vasireddi, N., Dhawan, N.S., Ma'ayan, A., Sharma, N., et al. (2019). Direct activation of PP2A for the treatment of tyrosine kinase inhibitor–resistant lung adenocarcinoma. Jci Insight *4*, e125693. https://doi.org/10.1172/jci.insight.125693.

15. Jayappa, K.D., Tran, B., Gordon, V.L., Morris, C.G., Saha, S., Farrington, C.C., O'Connor, C.M., Zawacki, K.P., Isaac, K.M., Kester, M., et al. (2023). PP2A modulation overcomes multidrug resistance in chronic lymphocytic leukemia via mPTP-dependent apoptosis. J. Clin. Investig. *133*. https://doi.org/10.1172/jci155938.

16. Avelar, R.A., Armstrong, A.J., Carvette, G., Gupta, R., Puleo, N., Colina, J.A., Joseph, P., Sobeck, A.M., O'Connor, C.M., Raines, B., et al. (2023). Small molecule mediated stabilization of PP2A modulates the Homologous Recombination pathway and potentiates DNA damage-induced cell death. Mol Cancer Ther. https://doi.org/10.1158/1535-7163.mct-21-0880.

17. Allen-Petersen, B.L., Risom, T., Feng, Z., Wang, Z., Jenny, Z.P., Thoma, M.C., Pelz, K.R., Morton, J.P., Sansom, O.J., Lopez, C.D., et al. (2018). Activation of PP2A and inhibition of mTOR synergistically reduce MYC signaling and decrease tumor growth in pancreatic ductal adenocarcinoma. Cancer Res *79*, canres.0717.2018. https://doi.org/10.1158/0008-5472.can-18-0717.

18. Leonard, D., Huang, W., Izadmehr, S., O'Connor, C.M., Wiredja, D.D., Wang, Z., Zaware, N., Chen, Y., Schlatzer, D.M., Kiselar, J., et al. (2020). Selective PP2A Enhancement through Biased Heterotrimer Stabilization. Cell. https://doi.org/10.1016/j.cell.2020.03.038.

19. Arnold, H.K., and Sears, R.C. (2008). A tumor suppressor role for PP2A-B56α through negative regulation of c-Myc and other key oncoproteins. Cancer Metastasis Rev. *27*, 147–158. https://doi.org/10.1007/s10555-008-9128-9.

20. Arnold, H.K., and Sears, R.C. (2006). Protein Phosphatase 2A Regulatory Subunit B56α Associates with c-Myc and Negatively Regulates c-Myc Accumulation. Mol Cell Biol 26, 2832–2844. https://doi.org/10.1128/mcb.26.7.2832-2844.2006.

21. Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J.P., and Tamayo, P. (2015). The Molecular Signatures Database Hallmark Gene Set Collection. Cell Syst. 1, 417–425. https://doi.org/10.1016/j.cels.2015.12.004.

22. Mootha, V.K., Lindgren, C.M., Eriksson, K.-F., Subramanian, A., Sihag, S., Lehar, J., Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., et al. (2003). PGC-1α-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. Nat. Genet. 34, 267–273. https://doi.org/10.1038/ng1180.

23. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proc. Natl. Acad. Sci. 102, 15545–15550. https://doi.org/10.1073/pnas.0506580102.

24. Navas, C., Hernández-Porras, I., Schuhmacher, A.J., Sibilia, M., Guerra, C., and Barbacid, M. (2012). EGF Receptor Signaling Is Essential for K-Ras Oncogene-Driven Pancreatic Ductal Adenocarcinoma. Cancer Cell 22, 318–330. https://doi.org/10.1016/j.ccr.2012.08.001.

25. Ardito, C.M., Grüner, B.M., Takeuchi, K.K., Lubeseder-Martellato, C., Teichmann, N., Mazur, P.K., DelGiorno, K.E., Carpenter, E.S., Halbrook, C.J., Hall, J.C., et al. (2012). EGF Receptor Is Required for KRAS-Induced Pancreatic Tumorigenesis. Cancer Cell 22, 304–317. https://doi.org/10.1016/j.ccr.2012.07.024.

26. Cook, N., Frese, K.K., and Moore, M. (2014). Assessing the role of the EGF receptor in the development and progression of pancreatic cancer. Gastrointest. Cancer: Targets Ther. 4, 23–37. https://doi.org/10.2147/gictt.s58925.

27. Valsecchi, M.E., McDonald, M., Brody, J.R., Hyslop, T., Freydin, B., Yeo, C.J., Solomides, C., Peiper, S.C., and Witkiewicz, A.K. (2012). Epidermal growth factor receptor and insulinlike growth factor 1 receptor expression predict poor survival in pancreatic ductal adenocarcinoma. Cancer 118, 3484–3493. https://doi.org/10.1002/cncr.26661.

28. Fujita, H., Ohuchida, K., Mizumoto, K., Itaba, S., Ito, T., Nakata, K., Yu, J., Kayashima, T., Hayashi, A., Souzaki, R., et al. (2011). High EGFR mRNA expression is a prognostic factor for reduced survival in pancreatic cancer after gemcitabine-based adjuvant chemotherapy. Int J Oncol 38, 629–641. https://doi.org/10.3892/ijo.2011.908.

29. Grapa, C.M., Mocan, T., Gonciar, D., Zdrehus, C., Mosteanu, O., Pop, T., and Mocan, L. (2019). Epidermal Growth Factor Receptor and Its Role in Pancreatic Cancer Treatment Mediated by Nanoparticles. Int. J. Nanomed. 14, 9693–9706. https://doi.org/10.2147/ijn.s226628.

30. Leonard, D., Huang, W., Izadmehr, S., O'Connor, C.M., Wiredja, D.D., Wang, Z., Zaware, N., Chen, Y., Schlatzer, D.M., Kiselar, J., et al. (2020). Selective PP2A Enhancement through Biased Heterotrimer Stabilization. Cell *181*, 688-701.e16. https://doi.org/10.1016/j.cell.2020.03.038.

31. Avelar, R.A., Gupta, R., Carvette, G., Leprevost, F. da V., Jasti, M., Colina, J., Teitel, J., Nesvizhskii, A.I., O'Connor, C.M., Hatzoglou, M., et al. (2024). Integrated stress response plasticity governs normal cell adaptation to chronic stress via the PP2A-TFE3-ATF4 pathway. Cell Death Differ. *31*, 1761–1775. https://doi.org/10.1038/s41418-024-01378-3.

32. McClinch, K., Avelar, R.A., Callejas, D., Izadmehr, S., Wiredja, D., Perl, A., Sangodkar, J., Kastrinsky, D.B., Schlatzer, D., Cooper, M., et al. (2017). Small-Molecule Activators of Protein Phosphatase 2A for the Treatment of Castration-Resistant Prostate Cancer. Cancer Res *78*, 2065–2080. https://doi.org/10.1158/0008-5472.can-17-0123.

33. Barton, C.M., Hall, P.A., Hughes, C.M., Gullick, W.J., and Lemoine, N.R. (1991). Transforming growth factor alpha and epidermal growth factor in human pancreatic cancer. J. Pathol. *163*, 111–116. https://doi.org/10.1002/path.1711630206.

34. Wang, L., Wang, L., Zhang, H., Lu, J., Zhang, Z., Wu, H., and Liang, Z. (2020). AREG mediates the epithelial-mesenchymal transition in pancreatic cancer cells via the EGFR/ERK/NF-κB signalling pathway. Oncol Rep *43*, 1558–1568. https://doi.org/10.3892/or.2020.7523.

35. Nakanishi, T., Koma, Y., Miyako, S., Torigoe, R., Yokoo, H., Omori, M., Yamanaka, K., Ishihara, N., Tsukamoto, S., Kodama, T., et al. (2024). AREG Upregulation in Cancer Cells via Direct Interaction with Cancer-Associated Fibroblasts Promotes Esophageal Squamous Cell Carcinoma Progression Through EGFR-Erk/p38 MAPK Signaling. Cells *13*, 1733. https://doi.org/10.3390/cells13201733.

36. Wang, L., Wu, H., Wang, L., Lu, J., Duan, H., Liu, X., and Liang, Z. (2016). Expression of amphiregulin predicts poor outcome in patients with pancreatic ductal adenocarcinoma. Diagn Pathol *11*, 60. https://doi.org/10.1186/s13000-016-0512-4.

37. Carpenter, E.S., Elhossiny, A.M., Kadiyala, P., Li, J., McGue, J., Griffith, B.D., Zhang, Y., Edwards, J., Nelson, S., Lima, F., et al. (2023). Analysis of donor pancreata defines the transcriptomic signature and microenvironment of early neoplastic lesions. Cancer Discov. *13*, 1324–1345. https://doi.org/10.1158/2159-8290.cd-23-0013.

38. Tun, M.T., Pai, R.K., Kwok, S., Dong, A., Gupta, A., Visser, B.C., Norton, J.A., Poultsides, G.A., Banerjee, S., Dam, J.V., et al. (2012). Diagnostic accuracy of cyst fluid amphiregulin in pancreatic cysts. BMC Gastroenterol. *12*, 15. https://doi.org/10.1186/1471-230x-12-15.

39. Mucciolo, G., Henríquez, J.A., Jihad, M., Teles, S.P., Manansala, J.S., Li, W., Ashworth, S., Lloyd, E.G., Cheng, P.S.W., Luo, W., et al. (2024). EGFR-activated myofibroblasts promote

metastasis of pancreatic cancer. Cancer Cell *42*, 101-118.e11.
https://doi.org/10.1016/j.ccell.2023.12.002.

40. Pavic, K., Gupta, N., Omella, J.D., Derua, R., Aakula, A., Huhtaniemi, R., Määttä, J.A., Höfflin, N., Okkeri, J., Wang, Z., et al. (2023). Structural mechanism for inhibition of PP2A-B56α and oncogenicity by CIP2A. Nat. Commun. *14*, 1143. https://doi.org/10.1038/s41467-023-36693-9.

41. Farrell, A.S., Allen-Petersen, B., Daniel, C.J., Wang, X., Wang, Z., Rodriguez, S., Impey, S., Oddo, J., Vitek, M.P., Lopez, C., et al. (2014). Targeting Inhibitors of the Tumor Suppressor PP2A for the Treatment of Pancreatic Cancer. Mol Cancer Res *12*, 924–939. https://doi.org/10.1158/1541-7786.mcr-13-0542.

42. Hruban, R.H., Adsay, N.V., Albores-Saavedra, J., Anver, M.R., Biankin, A.V., Boivin, G.P., Furth, E.E., Furukawa, T., Klein, A., Klimstra, D.S., et al. (2006). Pathology of Genetically Engineered Mouse Models of Pancreatic Exocrine Cancer: Consensus Report and Recommendations. Cancer Res. *66*, 95–106. https://doi.org/10.1158/0008-5472.can-05-2168.

43. Olive, K.P., Jacobetz, M.A., Davidson, C.J., Gopinathan, A., McIntyre, D., Honess, D., Madhu, B., Goldgraben, M.A., Caldwell, M.E., Allard, D., et al. (2009). Inhibition of Hedgehog Signaling Enhances Delivery of Chemotherapy in a Mouse Model of Pancreatic Cancer. Science *324*, 1457–1461. https://doi.org/10.1126/science.1171362.

44. Hingorani, S.R., Wang, L., Multani, A.S., Combs, C., Deramaudt, T.B., Hruban, R.H., Rustgi, A.K., Chang, S., and Tuveson, D.A. (2005). Trp53R172H and KrasG12D cooperate to promote chromosomal instability and widely metastatic pancreatic ductal adenocarcinoma in mice. Cancer Cell *7*, 469–483. https://doi.org/10.1016/j.ccr.2005.04.023.

45. Diersch, S., Wirth, M., Schneeweis, C., Jörs, S., Geisler, F., Siveke, J.T., Rad, R., Schmid, R.M., Saur, D., Rustgi, A.K., et al. (2016). KrasG12D induces EGFR-MYC cross signaling in murine primary pancreatic ductal epithelial cells. Oncogene *35*, 3880–3886. https://doi.org/10.1038/onc.2015.437.

46. Schneeweis, C., Wirth, M., Saur, D., Reichert, M., and Schneider, G. (2017). Oncogenic KRAS and the EGFR loop in pancreatic carcinogenesis—A connection to licensing nodes. Small Gtpases *9*, 00–00. https://doi.org/10.1080/21541248.2016.1262935.

47. Liu, H., Zhang, B., and Sun, Z. (2020). Spectrum of EGFR aberrations and potential clinical implications: insights from integrative pan-cancer analysis. Cancer Commun. *40*, 43–59. https://doi.org/10.1002/cac2.12005.

48. Zheng, S., Wang, W., Aldahdooh, J., Malyutina, A., Shadbahr, T., Tanoli, Z., Pessia, A., and Tang, J. (2022). SynergyFinder Plus: Toward Better Interpretation and Annotation of Drug Combination Screening Datasets. Genom., Proteom. Bioinform. *20*, 587–596. https://doi.org/10.1016/j.gpb.2022.01.004.

49. Cheng, Y., Liu, W., Kim, S.-T., Sun, J., Lu, L., Sun, J., Zheng, S.L., Isaacs, W.B., and Xu, J. (2011). Evaluation of PPP2R2A as a prostate cancer susceptibility gene: a comprehensive germline and somatic study. Cancer Genet-ny *204*, 375–381. https://doi.org/10.1016/j.cancergen.2011.05.002.

50. Kalev, P., Simicek, M., Vazquez, I., Munck, S., Chen, L., Soin, T., Danda, N., Chen, W., and Sablina, A. (2012). Loss of PPP2R2A Inhibits Homologous Recombination DNA Repair and Predicts Tumor Sensitivity to PARP Inhibition. Cancer Res *72*, 6414–6424. https://doi.org/10.1158/0008-5472.can-12-1667.

51. Goguet-Rubio, P., Amin, P., Awal, S., Vigneron, S., Charrasse, S., Mechali, F., Labbé, J.C., Lorca, T., and Castro, A. (2020). PP2A-B55 Holoenzyme Regulation and Cancer. Biomol *10*, 1586. https://doi.org/10.3390/biom10111586.

52. Curtis, C., Shah, S.P., Chin, S.-F., Turashvili, G., Rueda, O.M., Dunning, M.J., Speed, D., Lynch, A.G., Samarajiwa, S., Yuan, Y., et al. (2012). The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. Nature *486*, 346–352. https://doi.org/10.1038/nature10983.

53. Hein, A.L., Seshacharyulu, P., Rachagani, S., Sheinin, Y.M., Ouellette, M.M., Ponnusamy, M.P., Mumby, M.C., Batra, S.K., and Yan, Y. (2016). PR55α Subunit of Protein Phosphatase 2A Supports the Tumorigenic and Metastatic Potential of Pancreatic Cancer Cells by Sustaining Hyperactive Oncogenic Signaling. Cancer Res. *76*, 2243–2253. https://doi.org/10.1158/0008-5472.can-15-2119.

54. Kauko, O., O'Connor, C.M., Kulesskiy, E., Sangodkar, J., Aakula, A., Izadmehr, S., Yetukuri, L., Yadav, B., Padzik, A., Laajala, T.D., et al. (2018). PP2A inhibition is a druggable MEK inhibitor resistance mechanism in KRAS-mutant lung cancer cells. Sci Transl Med *10*, eaaq1093. https://doi.org/10.1126/scitranslmed.aaq1093.

55. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2012). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15–21. https://doi.org/10.1093/bioinformatics/bts635.

56. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–930. https://doi.org/10.1093/bioinformatics/btt656.

57. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550. https://doi.org/10.1186/s13059-014-0550-8.

58. Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2009). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics *26*, 139–140. https://doi.org/10.1093/bioinformatics/btp616.

59. Ihaka, R., and Gentleman, R. (1996). R: A Language for Data Analysis and Graphics. J. Comput. Graph. Stat. *5*, 299–314. https://doi.org/10.1080/10618600.1996.10474713.

60. Grecco, G.G., Huang, J.Y., Muñoz, B., Doud, E.H., Hines, C.D., Gao, Y., Rodriguez, B., Mosley, A.L., Lu, H.-C., and Atwood, B.K. (2022). Sex-Dependent Synaptic Remodeling of the Somatosensory Cortex in Mice With Prenatal Methadone Exposure. Adv. Drug Alcohol Res. *2*, 10400. https://doi.org/10.3389/adar.2022.10400.

61. McCracken, N.A., Liu, H., Runnebohm, A.M., Wijeratne, H.R.S., Wijeratne, A.B., Staschke, K.A., and Mosley, A.L. (2023). Obtaining Functional Proteomics Insights From Thermal Proteome Profiling Through Optimized Melt Shift Calculation and Statistical Analysis With InflectSSP. Mol. Cell. Proteom. *22*, 100630. https://doi.org/10.1016/j.mcpro.2023.100630.

62. Orsburn, B.C. (2021). Proteome Discoverer—A Community Enhanced Data Processing Suite for Protein Informatics. Proteomes *9*, 15. https://doi.org/10.3390/proteomes9010015.

63. Marx-Blümel, L., Marx, C., Kühne, M., and Sonnemann, J. (2016). HDAC/HAT Function Assessment and Inhibitor Development, Methods and Protocols. Methods Mol. Biol. *1510*, 23–45. https://doi.org/10.1007/978-1-4939-6527-4_3.

Figure legends.

Figure 1. PDAC cell lines with loss of B56α have reduced proliferation. (A) Representative western blot from AsPc1, MiaPaca2, and Panc1 cells showing loss of B56α protein in B56α knockout cell lines (sgB56α1 and sgB56α2) compared to control (sgNT). Proliferation across 96 hours in the above cell lines comparing loss of B56α to control cell line (sgNT). Area under the curve was calculated from 3 biological replicates and a one-way ANOVA was run on the area under the curve to determine significance. (B) Representative images of clonogenic colony growth and quantification of percent area of the well covered in in AsPc1 and Panc1 cells comparing loss of B56α to control cell line (sgNT). Fold change was calculated from at least 3 biological replicates relative to sgNT. Significance was determined using a one-way ANOVA. Error bars represent standard deviation *p<0.05, **p<0.01, ****p<0.0001

Figure 2. Overexpression of B56α increases proliferative phenotypes. (A) Proliferation across 96 hours in AsPc1, MiaPaca2, and Panc1 with overexpression of B56α (B56αOE) compared to control cell line (EV). Area under the curve was calculated from 3 biological replicates and two-tailed unpaired student's t test was run on the area under the curve to determine significance. (B) Representative images of clonogenic colony growth in AsPc1, Panc1, and MiaPaca2 with B56αOE after 14 days. Fold change was calculated from at least 3 biological replicates relative to EV. Significance was determined using a two-tailed unpaired student's t test. (C) Representative images of growth in soft agar after 14 days in Panc1, and MiaPaca2 with overexpression of B56α (B56αOE). Fold change in number of colonies was calculated across 3 biological replicates relative to EV control and two-tailed unpaired student's t test was used to determine significance. (D) Proliferation over 96 hours in AsPc1 cells with overexpression of B56ɣ (B56ɣOE), B56δ (B56δOE), B56ε (B56εOE), B55α (B55αOE), or B56α (B56αOE) compared to EV control. 3 biological replicates were used to calculate area under the curve and a one-way ANOVA on the area under the curve was used to determine significance. (D) Representative images and quantification of clonogenic colony growth in the above cell lines. Fold change of percent of well covered was calculated relative to EV control from 3 biological replicates and one-way ANOVA was used to calculate significance. Error bars represent standard deviation. Significance: *p<0.05, **p<0.01, ***p<0.001

Figure 3. B56α overexpression or pharmacological activation of PP2A increases EGFR signaling. (A) 10μM DT-061 treatment significantly enriched for genes associated with activated EGFR

signaling based on transcriptomics. (B) Heat map of mRNA expression of EGFR gene set from HPAFII cells treated with DT-061 with knockdown of B56α (siB56α) or non-targeting siRNA (siNT). (C) Volcano plot of differentially expressed total proteins with B56αOE compared to EV cells. (D) Volcano plot of differentially expressed proteins with DT-061 treatment compared to vehicle. (E) Significantly enriched oncogenic signature data sets of total proteomics with B56αOE. (F) Oncogenic signature data sets significantly enriched with DT-061 compared to vehicle. Sets are separated by oncogenic function (top) and tumor suppressive function (bottom). Gene sets are arranged by normalized enrichment score (NES), sized by size of the gene set, and colored according to nominal p-value (NOM p-val).

Figure 4. PP2A-B56α activation increases ligands which can activate EGFR. (A) Schematic for conditioned media (CM) experiment with DT-061 treatment. (B) Representative western blot of phosphorylated (Y1068) and total EGFR from AsPc1 cells treated with conditioned media (CM) from AsPc1 cells treated with 7.5µM DT-061 compared to vehicle CM. (C) Quantification of B of phosphorylated EGFR relative to total EGFR normalized to GAPDH. (D) Schematic for CM experiment with B56αOE or EV cells. (E) Representative western blot of phosphorylated (Y1068) and total EGFR from AsPc1 cells treated with CM from AsPc1 cells with B56αOE or EV. (F) Quantification of E of phosphorylated EGFR relative to total EGFR normalized to GAPDH. (G) mRNA expression of EGFR ligands from human PDAC cells lines in DMEM + 2% FBS treated with 10µM DT-061 or vehicle (DMSO) for 12 hours. Fold change calculated relative to 18S expression and the vehicle condition for each cell line. (H) Processed AREG from AsPc1 cells treated with 5µM DT-061 or vehicle for 24 hours. (I) Processed AREG from B56αOE or EV AsPc1 cells. Significance *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by two-tail unpaired student's t test

Figure 5. AREG and EGFR are necessary for increased proliferation with B56αOE. (A) Proliferation across 96 hours in AsPc1 B56αOE compared to control cell line (EV) with knockdown of EGFR (siEGFR) or control (siNT). Area under the curve was calculated from 3 biological replicates and one-way ANOVA was run on the area under the curve to determine significance. (B) Representative images of transwell migration of AsPc1 B56αOE cells compared to EV cells with knockdown of EGFR (siEGFR) or control (siNT). (C) Quantification of B, fold change of number of cells migrated through transwell relative to EV siNT. (D) Proliferation across 96 hours in AsPc1 B56αOE compared to control cell line (EV) with knockdown of AREG (siAREG) or control (siNT). Area under the curve was calculated from 3 biological replicates and one-way ANOVA

was run on the area under the curve to determine significance. (E) 300pg/ml AREG added to B56α knockout cells rescues proliferation at 96 hours. Significance *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by one-way ANOVA

Figure 6. Loss of CIP2A decreases survival of KPC mice and increases EGFR phosphorylation. (A) Kaplan-Meier survival curve of KPC PDAC mouse model showing decreased survival with loss of CIP2A (fl/fl, n=32, pink line) compared to wildtype (+/+, n=29, purple line) or heterozygous (fl/+, n=29, blue line). KPC =PDX1-Cre, KRAS$^{LSL-G12D}$, Tp53$^{LSL-R172H}$ (B) Representative images of endpoint tissue stained with PanCK to identify regions of interest and phosphorylated EGFR. Scale bars represent100mm (C) Quantification of mean fluorescent intensity of phospho-EGFR averaged from 5 fields of view from 6 tumors per genotype. *p<0.05 by two-tailed unpaired student's t test

Figure 7. DT-061 and Afatinib treatment are a synergistic combination *in vitro* and *in vivo*. (A) Drug matrix between Afatinib and DT-061 in MiaPaCa2. Cell viability (left) was read through crystal violet 72 hours after drug treatment. Bliss synergy (right) across the matrix. (B) Drug matrix between Afatinib and DT-061 in AsPc1. Cell viability (left) was read through crystal violet 72 hours after drug treatment. Bliss synergy (right) across the matrix. (C) MiaPaCa2 cells were grown subcutaneously in NRG mice and mice were treated with vehicle, afatinib (15mg/kg), DT-061 (15mg/kg), or combination for 3 weeks. Mean ± SEM are shown. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by 2way ANOVA, significance of DT-061 + Afatinib compared to: vehicle (black stars), Afatinib (teal stars), DT-061 (pink stars) B by one-way ANOVA. (D) Final tumor weight from C. *p<0.05 by 1-way ANOVA (E) Western blot from whole cell lysate from tumors from C (left). Quantification of phosphorylated EGFR (Y1068) relative to total EGFR normalized to GAPDH. *p<0.05, **p<0.01 by 1-way ANOVA.

Supplementary Figure 1. Overexpression of B56α in KRAS mutant NSCLC does not increase proliferation. (A) mRNA of B56α (*PPP2R5A*) in the overexpression cell lines (B56αOE) relative to the control (EV) normalized to 18S. (B) Representative western blots and quantification of B56α protein levels relative to control cells (EV) normalized to GAPDH. (C) mRNA expression of the overexpressed subunit relative to the EV from AsPc1 B56γ, B56δ, B56ε, or B55α overexpression cell lines. (D) Representative western blots showing exogenous expression of the HA tagged B56 subunit or B55α overexpression. (E) mRNA of B56α (*PPP2R5A*) in the overexpression cell lines (B56αOE) relative to the control (EV) normalized to 18S in Calu6, H23, and A549. (F)

Representative western blots and quantification of B56α protein levels relative to control cells (EV) normalized to GAPDH in Calu6 and H23. (C) Proliferation over 96 hours with overexpression of B56α (B56αOE) to control cell line (EV). Area under the curve was calculated to determine significance. Error bars represent standard deviation. Significance *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by two-tail unpaired student's t test.

Supplemental Figure 2. B56α overexpression or pharmacological activation of PP2A inhibits MYC targets. (A) AsPc1, Panc1, and HPAFII cells treated with 10µM DT-061 in DMEM +2% FBS for 24 hours show significant negative enrichment of genes regulated by MYC, indicating activity of PP2A. (B) Volcano plot of differentially phosphorylated phospho-sites with DT-061 treatment compared to vehicle. (C) Datasets from Human Molecular Signatures Database or Reactome pathways that are enriched with proteins that have significantly downregulated phospho-sites with DT-061. (D) Volcano plot of differentially phosphorylated phospho-sites with B56αOE compared to EV. (E) Datasets from Human Molecular Signatures Database or Reactome pathways that are enriched with proteins that have significantly downregulated phospho-sites with B56αOE. (F) Kinase enrichment analysis (KEA) of significantly upregulated phospho-sites with B56αOE. (G) Significantly enriched hallmarks of significantly upregulated phospho-sites with B56αOE. (H) Hallmark gene sets significantly enriched from AsPc1 B56αOE cells compared to EV cells based on total proteomics. (I) Hallmark gene sets significantly enriched from DT-061 treated cells compared to vehicle from total proteomics. Gene sets are arranged by normalized enrichment score (NES), sized by size of the gene set, and colored according to nominal p-value (NOM p-val).

Supplemental Figure 3. PP2A-B56α activation increases AREG shedding (A) Processed AREG from MiaPaCa2 cells treated with 5µM DT-061 or vehicle for 24 hours. (B) Processed AREG from B56αOE or EV MiaPaCa2 cells. (C) Processed AREG from Capan2 cells treated with 5µM DT-061 or vehicle for 24 hours. (D) Processed AREG from B56αOE or EV Capan2 cells. (E) Processed AREG from HPAFII cells treated with 10µM DT-061 or vehicle for 24 hours. Significance *p<0.05, **p<0.01, by two-tail unpaired student's t test

Supplemental Figure 4. mRNA expression from rescue experiments. (A) mRNA expression of EGFR over the proliferation assay. (B) mRNA expression of EGFR and B56α (*PPP2R5A*) at the time of migration harvest. (C) mRNA expression of AREG over the time course of the proliferation assay. Significance **p<0.01, ***p<0.001, ****p<0.0001 by one-way ANOVA

Supplemental Figure 5. (A) Relative cell viability in drug matrix of DT-061 and Afatinib. Cell viability was read using crystal violet 72 hours after drug treatment. (B) Bliss synergy scores of matrices from A. (C) Body weight remained steady and similar amongst treatment and vehicle arms *$p < 0.05$ by 1-way ANOVA.

Figure 1

A

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

KPC Cip2A KO (KPCC)
Overall Survival

KPC
KPC Cip2a$^{fl/+}$
KPC Cip2a$^{fl/fl}$

Log-rank Mantel-Cox: *p=0.0005*
Gehan-Breslow-Wilcoxon test: *p=0.0002*

Probability of Survival

Time

C

Mean fluorescence intensity

KPC   KPCC

B

DAPI   PanCK   p-EGFR (Y1068)   Merge

KPC

KPCC

Figure 7