1 **Slowing down: A macroevolutionary approach to the hypometabolic strategies of**
2 **amphibians**
3
4 **Abstract**
5

The ability to survive harsh environmental conditions has probably been a key factor in the
evolutionary success of organisms that cannot migrate long distances, such as amphibians. We
expect that having a hypometabolic strategy (HS) —aestivation or hibernation— to deal with
severe climates, would be a plesiomorphic trait. We 1) inferred the ancestral state of a HS, using
two phylogenies for amphibians, 2) tested if species with a HS have larger distributional ranges,
and 3) explored how a HS may affect amphibian assemblage resilience using multiple models of
climate change. Ancestral state reconstruction for the most recent common ancestor (MRCA) of
Class Amphibia showed ~50% probability of a HS. The probability was higher for the MRCA of
each Order (>70%), suggesting a widespread HS in the ancestors of modern amphibians.
Phylogenetic regressions showed no relation between the probability of having a HS and the
distribution range size. Climate analyses predict that tropical zones will have the greatest change
in climate, involving novel harsh seasonality. Since tropical amphibian assemblages have the
lowest proportion of species with HS, they may be more vulnerable to climate change. It is
probable that HS have been key for the evolutionary success in amphibians, and they will likely
impact their future survival in the face of climate change. Despite the potential importance of the
HS for amphibians, information was available for a diverse but only a small subset of species;
we urge researchers to report data on aestivation or hibernation in amphibians to facilitate future
studies.

Keywords (4-10): ancestral states**,** aestivation, conservation, hibernation, hypometabolism,
environmental seasonality.

Introduction
Most animals seek shelter when environmental conditions turn harsh. For instance, amphibians
use shelters to hide during their inactive hours on a daily basis (Wells, 2007) in order to hide
from predators and conserve water. Amphibians are required to keep their skin moist because it
is crucial for breathing, and, therefore, their evaporative water loss is normally high (Lillywhite,
1975). Consequently, to survive long periods (weeks, months, or years) during adverse
environmental conditions, amphibians resort to a type of dormancy or latency in a state of
reduced or depressed metabolism (hereafter, hypometabolism) when they hide underground or
underwater. The lengths of these hypometabolic periods vary considerably across species and
populations because it seems to be driven by the local environmental conditions (Figure 1).
Here we will use the most common terminology for the different states of dormancy: we
will refer to summer dormancy as aestivation and winter dormancy as hibernation (see Box 1).
We also make the assumption that both strategies imply a hypometabolic state. Thus, we will

refer to them collectively as hypometabolic strategies (HS). Here we will not include cryptobiosis (from the Greek *crypto* and *bios* meaning 'hidden life'), an extreme form of hypometabolic state, where metabolism is reduced to an undetectable but non-zero level (e.g., cryobiosis, when organisms freeze in order to survive the winter), although it might be considered an extreme case of hypometabolism (Figure 1). Nonetheless, whether in the hottest and driest season of the year (i.e., aestivation) or in the coldest (i.e., hibernation), the end result is the same, some amphibians "slow their lives" (namely, lower their metabolism) in order to get through the unfavorable conditions that the environment imposes. In some cases, this pattern is so extreme that amphibians are active only a few weeks each year, like some desert amphibians (McClanahan, 1972; Cowan & Storey, 1999; Withers & Cooper, 2010). For example, the maximum aestivation period recorded in amphibians is five years in the Australian striped burrowing frog *Cyclorana platycephala* (Van Beurden, 1980). It is unknown if these extremely long periods of hypometabolism can be repeated several times throughout the animal's life or if there are limits. In some cases, aestivating amphibians produce a cocoon from several layers of shed skin (Withers, 1995); the cocoon provides extra insulation to prevent water loss and might increase the duration of aestivation periods in these species (Withers, 1998). For example, estimations show that it would take more than eight years for a *Cyclorana* frog with a cocoon and an empty bladder to lose 30% of its body mass by evaporation (Sadowski-Fugitt *et al.*, 2012).

According to Storey & Storey (2010), three requirements must be satisfied in order for a state to be considered hypometabolic: 1) an overall strong suppression of metabolic rate (at least a 70–80% reduction); 2) differential control over the rates of various metabolic processes so that energy use is reprioritized to favor core vital cell functions; and 3) implementation of actions that protect cells and preserve viability over long periods of time. There are several mechanisms that are activated during metabolic rate depression, and it is proposed that these mechanisms, their interactions, and their regulatory signals, form a common molecular basis for all states of metabolic depression including aestivation, hibernation, and some types of anoxybiosis (metabolic depression due to a lack of oxygen; Storey & Storey, 2004). Although the relationship between environmental condition and hypometabolic state is well documented (see below), we still lack assessment of this idea in amphibians across a large taxonomic scale.

The ability to tolerate unfavorable or extremely unfavorable environmental conditions (such as very high or very low temperatures or droughts) has probably been a key factor in the evolutionary success of organisms that cannot escape such harsh conditions. As a group, amphibians have been on Earth for more than 300 Myr and there have been many periods of unfavorable conditions during that long time (Vitt & Caldwell, 2014; Jetz & Pyron, 2018; Judd *et al.*, 2024), including several mass extinctions. The ability to sustain a hypometabolic state might have also been an important factor in the colonization of land, given that lungfish also aestivate and produce cocoons (Jiang *et al.*, 2023). In this paper we first aim to infer the ancestral state and evolutionary trajectory of hypometabolic strategies in all living amphibians. Our approach is based on the observation that many groups of frogs with aestivation evolved in arid

or semiarid environments, such as Ceratophryidae (Faivovich *et al.*, 2014). All ceratophryids are
capable of prolonged aestivation and cocoon formation, even the ones that no longer live in
seasonal environments (Zimicz *et al.*, 2024). Paleozoic amphibians appear to have burrowed in
response to seasonal droughts (Hembree *et al.*, 2004), and likely other climate crises. For
example, radiation of fossorial amphibians and increase in abundance and complexity of burrows
coincide with the Cisuralian aridification and end-Permian extinction events (Marchetti *et al.*,
2024). The ability of early Tertiary pelobatids to avoid drought by estivating in burrows is
thought to be a preadaptation to survive the global cooling of the middle Eocene (Henrici &
Haynes, 2006). Therefore, we would expect that hypometabolic strategies evolved or were
retained in places with extreme environmental conditions, specifically with aestivation occurring
in semiarid conditions and hibernation in cold conditions (Hembree *et al.*, 2004; Henrici &
Haynes, 2006; Faivovich *et al.*, 2014; Marchetti *et al.*, 2024; Zimicz *et al.*, 2024). As part of this
aim, we then inferred the ancestral state and evolutionary trajectory of amphibian HS; given the
role of amphibians colonizing land and inhabiting harsh or seasonal environments, we expected
to find that the capacity for hypometabolism is a plesiomorphic trait—an ancestral trait inherited
from a common ancestor and shared by all members of a clade that in some lineages may be lost
subsequently— in amphibians.
As a second aim, we explore the association between the presence of hypometabolic
strategies and distributional ranges. We suggest that if a species has the possibility to avoid
stressful environments, it should be able to invade novel areas without negative fitness
consequences. Moreover, having a hypometabolic strategy may allow a species to survive over a
broad range of environmental conditions. Therefore, we expect that species with a
hypometabolic strategy will have larger distribution ranges.
Finally, the presence or absence of a hypometabolic strategy has possible implications for
how amphibians might respond to future changes in climate, and thus may be a concern for
conservation biologists. Indeed, climate change will impact amphibian distributions and ranges
(Ochoa-Ochoa *et al.*, 2012; Mokhatla *et al.*, 2015; Velasco *et al.*, 2021), will vary in speed across
regions (Ochoa-Ochoa & Velasco, 2024; García-Rodríguez et al., 2025), and produce climatic
debt—a disequilibrium between species communities' climatic tolerances and the magnitude of
climatic change (He *et al.*, 2023; Smith *et al.*, 2025). As a third aim, we explore the vulnerability
of amphibian assemblages due to climate change in relation to the presence of hypometabolic
strategies. Species that can enter dormancy or a sustained hypometabolic state may be able to
deal better with environmental changes such as desertification, severe temperature/precipitation
changes or extreme cold conditions. Therefore, we propose that climatic resilience—the ability
to cope with climatic stressors—of amphibian assemblages increases with increasing proportions
of species with hypometabolism (Figure 2). Based on this hypothesis we make predictions
regarding amphibian assemblage vulnerability across the globe. Amphibian hypometabolic
strategies remain an understudied phenotype and here we take a macroevolutionary perspective
to improve our current state of knowledge.
120

Methods

Hypometabolism data collection. We compiled information about hypometabolic strategies, particularly aestivation and hibernation, through a systematic literature review (see *Systematic literature review*). A comparative dataset for hypometabolic states was assembled including aestivation/hibernation status (present/possible/absent), and range size in km^2 (Table S1 in the Supplementary Information). For scoring aestivation/hibernation, we considered them as present in a species if it was reported in at least one population or subspecies in the literature. We recorded absences (tentatively) when published studies or researchers noted its absence from personal experiences in fieldwork. However, we note that these cases will likely require further validation, and therefore, we did not include them as part of the training dataset (see *HS probability*) for subsequent models. Rather, an estimated probabilistic value based on a model was used instead.

Systematic literature review. In order to compile the dataset of amphibians that present a hypometabolic state, an exhaustive bibliographic search was carried out. The systematic search was performed using the advanced search tool of each of the following search engines: Scopus, PubMed, and Web of Science, until December 2024. We used the words (aestivat* AND amphibian*) and (estivat* AND amphibian*) for aestivation, and (hibernat* AND amphibian*) for hibernation. Studies were selected according to the following criteria: 1) original and review manuscripts including not just peer-reviewed literature, but also theses; 2) original articles in which some biological, ecological, physiological, or genetic aspect of the aestivation process were mentioned; and 3) manuscripts where the focus was not related to dormancy processes, but an aestivating or hibernating species was mentioned. We also performed a search on AmphibiaWeb to review and complete our resulting list of species that aestivate, hibernate, or have a cryptobiotic state. Scientific names follow AmphibiaWeb (2024) nomenclature.

Phylogenetic and geographic data. Because our subsequent analyses require a phylogenetic framework, we used the amphibian supertrees featured in Jetz and Pyron (2018) and in Portik *et al.* (2023), although these last ones are just for the Anura clade. Given that the first set of trees were generated with a mixed methodology incorporating a backbone for the species with available genetic information and the imputation of the species without data, the final consensus is highly polytomic. Therefore, in order to account for the inherent phylogenetic uncertainty (Rangel *et al.*, 2015), we subsampled 100 randomly resolved trees from each dataset. In addition, we downloaded the available distribution polygons for amphibians from the International Union for Conservation of Nature Red List (IUCN, 2024; https://www.iucnredlist.org) in its 2023-1 update. Because the trees and polygons may differ in the taxonomic treatment of some species, we harmonized the names between data sources to maximize the number of species in the final dataset. To do so, we manually validated the non-matching species against their synonym list available in the Amphibian Species of the World (https://amphibiansoftheworld.amnh.org) looking for alternative names. From the 857 species with hypometabolism data, 774 from Jetz

and Pyron's dataset (approx. 90%) and 477 from Portik *et al.*'s dataset (83.52%) were shared between the phylogeny and geographic data. Although the non-matching species were not included in the phylogenetic comparative analyses, they were included to generate the hypometabolic probabilities (see *HS probability*) and to illustrate the summary information on the global status of hypometabolism in amphibians.

HS probability. Since we have literature information for only 857 species out of the 8,823 amphibians currently described, fewer than 10% of the total number of species (AmphibiaWeb, 2024), an important issue in this study is the large amount of missing data. Thus, we calculated the probabilities of the presence of a hypometabolic strategy in the tips with unknown states by assuming that hypometabolic states are expressed in response to environmental conditions and that species are adapted to the environments they inhabit (Hembree *et al.*, 2004; Henrici & Haynes, 2006; Faivovich *et al.*, 2014; Zimicz *et al.*, 2024). Importantly, we highlight that this approach makes the assumption that species have tracked their niches over evolutionary time. To conduct this analysis, we first extracted for each species the maximum values for the 19 WorldClim bioclimatic variables (https://www.worldclim.org) at 10 arcmin resolution (approx. 18.5 km at the equator) based on 7,949 distribution polygons. To address multicollinearity, we thinned our dataset by excluding highly correlated variables using the Variance Inflation Factor (VIF) with the *vifstep* function as implemented in 'usdm' v.2.1 (Naimi *et al.*, 2014). We then fit a One-Class Support Vector Machine (OC-SVM) model using a subsampled training dataset that included only species with confirmed presence separately for aestivation and hibernation.

OC-SVM is suitable for presence-only data, allowing us to avoid model biasing given the uncertainty in the absence data (false negatives). For this, we employed the *svm* function in the 'e1071' v.1.7-16 package (Meyer *et al.*, 2024), using a radial kernel with a custom gamma (shape of the kernel) for each trait (aestivation = 0.25, hibernation = 0.15), nu = 0.1 (sensitivity to outliers), and a 10 k-fold cross-validation strategy. We manually tuned the shape of the kernel to optimize the model's performance by minimizing the number of false negatives (true presences with low estimated probability) while increasing the model's accuracy (Figures S1–S2). Posteriorly, decision values were scaled between 0 and 1 and transformed into probabilities using a sigmoid function. Subsequently, for species with confirmed presence we kept the probability as 1, but replaced the other ones in our literature review (i.e., possible state of aestivation or hibernation) with their inferred probability in order to account for the uncertainty in defining true absences. We repeated this estimation twice, once for aestivation and once for hibernation, for each phylogeny since the number of species in the model change, thus will the probabilities. For the species assigned by the literature review as "possible", two outcomes were possible: 1) they were assigned a low probability, equivalent to an absence; 2) they were assigned a high probability (> 80%), equivalent to a presence under our assumption that climate is triggering such hypometabolic strategies (and our initial coding from the literature is just a reflection of the absence of data).

Afterwards, in order to use these probabilities in the combined hypometabolism analysis,
we assigned the maximum probability of either aestivation or hibernation to each species (i.e., if
the species is predicted to estivate but not hibernate it would still be counted as a "presence" for
having a hypometabolic state). Although we calculated the missing data for all the amphibians in
the trees using probabilities, we performed subsequent ancestral state reconstructions (see
Reconstruction of ancestral states) in trees with twice the number of species with hypometabolic
data, by using the 774 species from the literature (fixed species) and randomly subsampling
another 774 species for each tree with Jetz and Pyron (2018) phylogeny and 477 species for
Portik *et al*. (2023). Thus including 50% of species with literature hypometabolic state and 50%
of species with inferred hypometabolic state. The aim of this last is: 1) to increase the taxonomic
representativeness of the different amphibian taxa; 2) to reduce the bias in the results by
accounting only for species with presence or potential presence data; and, finally, 3) to balance
the effect of using the entire estimated dataset.
213
Reconstruction of ancestral states. Here, we modeled the evolutionary transitions between the
presence and inferred absence of aestivation, hibernation, and hypometabolism (either
aestivation or hibernation), the last following the assumption that all states of metabolic
depression have a common molecular basis (Storey & Storey, 2004) but we stress that the
environmental cues to enter aestivation or hibernation are very likely distinct. Subsequent
analyses were performed using 'phytools' v.2.3 (Revell, 2024), and a script can be found as part
of the Supplementary Files. First, we employed the Markov k model (hereafter Mk model) for
discrete character evolution of characters with k unordered states (Lewis, 2001) to find the best
state transition model, as implemented in the function *fitMk*. As a Markovian model, it assumes
that transitions between states depend only on the current state and not on previous ones. We
tested two different models of transition rates: equal (ER) and all different (ARD), allowing the π
parameter to estimate the prior distribution of the root state jointly with the other model
parameters rather than defining it *a priori* (Fitzjohn *et al.*, 2009). Since we are using a collection
of trees rather than a single one, we ran the analyses 100 times and chose the best model based
on its likelihood and Akaike information criterion (AIC) values. For both phylogenies the best
transition model for aestivation and hypometabolism was inferred to be the ARD, while for
hibernation, there were no differences between the ARD and the ER (Figures S3–S4). Thus, the
reconstructions were estimated using the ARD models.
Given that the presence of a hypometabolic state for many species may be ambiguous
given the paucity of data, we accounted for uncertainty in trait states as probabilities. Then, we
mapped possible scenarios of trait evolution using stochastic character mapping over the 100
trees for aestivation, hibernation, and hypometabolism separately. We employed the
make.simmap function with 100 simulations per tree using the ARD transition model, π as in the
fitMk, a non-fixed transition matrix (Q) estimated with a Markov Chain Monte Carlo (MCMC)
approach, empirical priors from observed data, a variance of 0.1 (vQ), and a sampling frequency
of 10. Thus, we ran a total 1000 MCMC iterations, sampling every 10 for a total of 100

independent histories per tree. Subsequently, we calculated the average number of transitions
across 100 simulations for each of the 100 trees using the countSimmap function. Since the Q
matrices were estimated independently for each simulation and tree, the MCMC algorithm may
get stuck in suboptimal peaks, yielding extremely low or high numbers of transitions in some
cases and increasing the uncertainty in the transition counts. Therefore, the average number of
gains and losses was calculated excluding outlier values (those above or below 1.5 times the
interquartile range).
Because each tree differs in topology, it is not possible to summarize the nodes across the
entire set of reconstructions (10,000 different sampled histories). Thus, for the purpose of
visualizing the results, we selected one of the 100 trees. Instead of choosing a random tree, we
selected the one whose likelihood in the evaluation of the evolutionary models was closest to the
median of the distribution frequency of all trees (Figures S3–S4). These particular trees were
only used for graphical purposes, the results are based on 100 trees from each phylogeny. It is
important to highlight that for the Jetz and Pyron phylogeny, given the method used for inferring
the base topology, such differences (i.e., polytomies) are expected to be more pervasive at lower
taxonomic levels (e.g., species or genus). Thus, to overcome topological differences while
accounting for uncertainty, we identified the MRCA for superfamilies and orders, for both set of
analyses, using the fixed species on each tree (i.e., those non-randomly selected), and averaged
the posterior probabilities of those nodes. It is important to note that, those MRCA are not
necessarily the same in all the trees but the most inclusive monophyletic groups encompassing
our fixed species. Because this kind of reconstruction requires discrete traits at tips, we
considered values equal or greater than 80% as strong evidence for presence, given that the
probability for all true presences from the literature was estimated with the same threshold (see
Figures S1 and S2). Moreover, in order to explore the transition of the states in the nodes along
time, we also performed a joint ancestral state reconstruction over the 100 random trees from
each phylogeny using the function *ancr*, to quantify the proportion of nodes with aestivation or
hibernation on 5-Myr bins and its possible association with the mean Earth surface temperature
(Judd *et al.*, 2024).
268
Geographic distribution of species with hypometabolism. To explore geographic patterns in
our data, we generated maps of species richness with aestivation, hibernation, and
hypometabolism by overlapping the IUCN species polygons (IUCN, 2024) with a grid of $1° \times 1°$
(ca. 111 km at the equator); we also generated maps using the proportion of species with a
hypometabolic state compared to the assemblage. We generated maps using only the species
with data from the literature review, including any species with presence or possibly presence of
a hypometabolic state as presence. Then, we repeated the calculation of species richness with all
the species in a grid cell to calculate the proportions of the total species in each cell with a HS.
277
Distributional range analyses. In order to explore whether the species with a hypometabolic
strategy had larger geographic ranges and account for phylogeny we used a phylogenetic

generalized least squares (PGLS) regression. We ran five different evolutionary models
(OURandom, OUFixed, BM, Pagel's Lambda and Pagel's Kappa) to estimate the relationship
between the continuous probability estimates for aestivation, hibernation, and hypometabolism
with distributional range size, specifically log1p(area), which computes $\log(1 + \text{area})$. These
models were run for each phylogeny. The best performance models, for both phylogenies, were
Pagel's Lambda and Pagel's Kappa in both normality of the residuals (visually checked through
histograms) and lowest AIC. Since Pagel's Lambda was the model with the lowest values of AIC
we chose this model to discuss in the manuscript. We used the 'phylolm' package (Ho & Ane,
2014). Finally, we counted the threatened category of the species in the literature dataset and
discussed the implications for conservation, particularly regarding the upcoming environmental
changes.
291
Amphibian assemblage climatic resilience. According to our hypothesis, the greater the
proportion of species with hypometabolism, the higher climatic resilience the assemblage would
have. To test this idea, we quantified dissimilarities between the baseline (1990–2020) and future
climate scenarios (2070–2100) by calculating the standardized local anomalies (i.e., how atypical
is a value compared with the historical variability) per raster pixel (~ 1 km^2), using the
standardized Euclidean distance (SED) (Williams *et al.*, 2007). In other words, the temporal
differences for each climate parameter are standardized using the local inter-annual standard
deviation for that parameter. This method is able to provide information on the extent of the
change in each location in absolute terms, therefore it gives direct information on the magnitude
of climate change to which the assemblage might be exposed in a particular grid cell (Williams
et al., 2007; Taheri *et al.*, 2024a).
We focused on the monthly values of the multi-model mean of precipitation, and
maximum and minimum of temperature from the experiments from KNMI 2023 standard set of
CMIP6 General Circulation Models. Historical and future change scenarios were obtained from
the Climate Explorer (https://climexp.knmi.nl). We choose a middle shared socioeconomic
pathway: the regional rivalry climate change scenario ssp370 or the "rocky road", where
countries prioritize national interests over global cooperation thus leading to an increase in
greenhouse gases. We use the mean of all 24 models for historical and future scenarios of
climate change (CCESS-CM2, ACCESS-ESM1-5, AWI-CM-1-1-MR, BCC-CSM2-MR,
CMCC-CM2-SR5, CNRM-CM6-1-HR-f2, CNRM-CM6-1-f2, CNRM-ESM2-1-f2, CanESM5-
CanOE-p2, CanESM5, EC-Earth3-Veg, EC-Earth3, FGOALS-g3, GFDL-ESM4, GISS-E2-1-G-
p3, INM-CM4-8, INM-CM5-0, IPSL-CM6A-LR, MIROC-ES2L-f2, MIROC6, MPI-ESM1-2-
HR, MPI-ESM1-2-LR, MRI-ESM2-0, and UKESM1-0-LL-f2). In order to estimate the   SED
, we employed the *sed* function in the 'climetrics' package v.1.0-15 (Taheri *et al.*, 2024b),
analyses were performed using the platform R version 4.4.2 (R Core Team, 2024).
To explore where would be located the most vulnerable amphibian assemblages we first
averaged climatic standardized distance per grid (1° × 1°) to match the scale of the species
richness maps. We generated a heat map of the climate change magnitude through SED against

the proportion of species with a HS (only literature data) under the assumption that the higher the
proportion of species with a HS, the higher the expected resilience to climatic change. In order to
localize the most vulnerable assemblages we selected the sites with highest increase in climate
variance (> 2 of SED) and low proportion of species with hypometabolism (≤ 0.5). Afterwards,
we plotted the selected assemblages on a map, we also included grid cells with the 10th highest
percentile of the anomaly scores which corresponds to large changes in temperature and
precipitation.
327
Results
The systematic search rendered 280 articles for aestivation and 650 for hibernation. We found
information about a hypometabolic strategy for 857 amphibian species (Table S1). From these
species, 648 had some information about aestivation, 342 with presence, 235 with the possibility
of presenting that strategy, and 52 as non-aestivating species. In the case of hibernation, we
found information for 479 species, 308 with presence, 62 species with possible hibernation, and
48 that showed no hibernating strategy. The representativity of this data set is as follows:
Gymnophiona, one of nine families (11%); Caudata, ten of ten (100%); and Anura, 28 of 59
(48%). Most of these species that present aestivation (389) are categorized as Least Concern
(LC), 20 species are classified as Endangered (EN), 9 species from *Litoria* and *Xenopus* genera,
are in the category of Critically Endangered (CR) and only *Anaxyrus baxteri* is classified as
Extinct in the Wild (EW). In the case of hibernation, the proportion of species is similar with
most of the species (242) categorized as LC, 19 species classified as EN, 3 as CR, two species,
Cynops wolterstorffi and *Rheobatrachus silus*, as Extinct (EX) and *A. baxteri* as EW (Table S1;
Figure S5).
343
Ancestral states. The accuracy of the traits inferred for the test dataset using OC-SVM
compared to data from the literature was high, with accuracies of 73% and 88 % for aestivation
and 72% and 88 % for hibernation, for Jetz & Pyron and Portik *et al.* phylogenies, respectively
(Figures S1–S2); in addition, all confirmed aestivation and hibernation presences based on the
literature were predicted as presences by OC-SVM with $> 80\%$ probability.
The ancestral state reconstruction analyses for the 100 Jetz & Pyron (2018) trees showed
a high uncertainty regarding the state for the MRCA of all amphibians, with an estimate close to
50/50 (presence/absence) for each strategy: aestivation 48.33/51.67 (SD $\pm$ 9.07), hibernation
51.51/48.49 (SD $\pm$ 14.70), and hypometabolism 48.27/51.73 (SD $\pm$ 9.83) (Figure 3; Table 1). It
is worth mentioning that the standard deviation is the same for presence and absence since the
values for each tree node are complementary. With the Jetz and Pyron (2018) phylogeny, the
probability of the presence of a hypometabolic state for either aestivation or hypometabolism
increases rapidly towards the present, being the most probable state for the MRCA of each
amphibian Order: probabilities for Anura (frogs) of aestivation 62.79 (SD $\pm$ 11.23) and
hypometabolism 73.09 (SD $\pm$ 13.73); for Caudata (salamanders) 68.34 (SD $\pm$ 11.97) and 78.66
(SD $\pm$ 12.30); and for Gymnophiona (caecilians) 71.72 (SD $\pm$ 16.22) and 77.26 (SD $\pm$ 14.74)

(Table 1). But this was not the case for hibernation where Anura (53.97 ± 14.47) and Caudata
(67.23 ± 18.49) tended for a presence, while in Gymnophiona hibernation seems more likely to
be absent (64.74 ± 15.44) than present (35.26 ± 15.44). Analyses using the Portik *et al.* (2023)
phylogeny also support the presence of a HS in the ancestor of Anura (70.9/29.1 ± 14.6), but
provided more mixed results for hibernation (47.9/52.1 ± 9.90) and aestivation (50.2/49.8 ±
2.30), specifically. Nonetheless, the internal nodes for both phylogenies tended to be predicted to
have presence of hibernation even in the superfamilies (Table 1). On average, 362 (SD ± 86)
gains were inferred for aestivation and 262 (SD ± 62) for hibernation; 327 (SD ± 80) and 248
(SD ± 84) losses were inferred for aestivation and hibernation, respectively. In the combined
hypometabolism, 283 (SD ± 76) gains and 250 (SD ± 69) losses were inferred. In all three
strategies, there were more gains than losses (Figure S6). For the Portik *et al.* (2023) phylogeny,
on average, 200 (SD ± 36.1) gains were inferred for aestivation and 163 (SD ± 34.2) for
hibernation; 150 (SD ± 41.7) and 170 (SD ± 35.4) losses were inferred for aestivation and
hibernation, respectively. For hypometabolism, 161 (SD ± 45.1) gains and 151 (SD ± 32.7)
losses were inferred (Figure S7).
For aestivation it is possible to observe a complex pattern of gains and losses of the trait
(Figure 4), where most of the clades have representative lineages of both, although there are a
few groups exhibiting only presence such as the Pelodryadinae subfamily (i.e., *Cyclorana*), and
some genera within the subfamilies Hyloxalinae and Mantellinae, and the Ceratophryidae family.
The transitional evolution of aestivation is likely to be complex, for example being the most
likely ancestral state in groups like Myobatrachoidea (> 80% with Jetz & Pyron phylogeny and >
60% with Portik *et al.* phylogeny), but for Ranoidea although probabilities are above 50 and 80%
there seems to be multiple independent losses (Figure 4). For hibernation there are only a few
groups that show a high probability for the presence of hibernation as an ancestral state, such as
Caudata, as well as in some anurans, including Myobatrachoidea (~ 80% and 78%),
Paleobatoidea (~ 50% and 80%) and Pipoidea (~ 64% and 55%), while Hylodea 15% and 25%
seems to have a very low probability of presence of hibernation (Table 1, Figures 4 and S8).
Meanwhile groups such as Sirenoidea, Salamandroidea, Myobatrachoidea, and Bombinatoroidea
showed remarkably high probabilities of presence of hypometabolism (Table 1). In the case of
the Portik *et al.* phylogeny all anuran superfamilies showed > 60% probability of
hypometabolism.
At a broad scale, based on the selected tree only for the Jetz & Pyron phylogeny, the
proportion of ancestral nodes with aestivation decreased during the Jurassic when the Earth's
surface temperature was transitioning to a cool-house (ca. 175–160 Myr), to posteriorly recover
again in the transition to a warm-house (ca. 160–150 Myr). Unexpectedly, aestivation's greatest
depression was inferred during a warmer period between 150–125 Myr, with a posterior recovery
and fluctuating but steady trend to the present. On the other hand, the hibernation trend was
stable during the cooled period in the Jurassic, with a drop during the transitional period to
warmer temperatures 160 Myr, with a posterior recovery but reduction again during the

remaining Mesozoic, and a recovery during the transition to a cooler Earth in the past 50 Myr
(Figure 5).
For Portik *et al.* phylogeny and based on the selected tree only, a similar pattern as with
Jetz & Pyron phylogeny occurred during the Jurassic but with the interesting fact that the
average of presence decreases to zero, to posteriorly recover again in the transition to a warm-
house (ca. 150 Myr). Another aestivation depression was inferred during a warmer period
between ~ 95–85 Myr, with a posterior recovery, and another depression ~ 70 Myr and finally a
recovery fluctuating but steady trend to the present. While aestivation was in depression during
the Jurassic the proportion of ancestral nodes with hibernation increased. The hibernation trend
showed gains and losses (but never as deep as in aestivation) during the remaining Mesozoic,
and a stable trend during the transition to a cooler Earth in the past 50 Myr (Figure S9).
410
Geographic distribution of species with hypometabolism. As mentioned in the methods, here
we are only using literature data. The greatest number of aestivating species are located in the
Eastern United States, Northern and Eastern Australia, parts of Indonesia, the Dry Diagonal in
South America, and in South Africa and Zimbabwe (Figure S10). In terms of the proportion of
species in a community with aestivation, the pattern is much different, highlighting regions of the
world with arid climates such as southwestern US, Botswana, parts of the Middle East and
northern Africa, New Zealand and much of Australia (Figure S11). Hibernating species richness
is predicted to be highest in the eastern United States, parts of China and Europe (Figure S12),
yet communities with the highest proportion of species that hibernate clearly predominate in
northern latitudes and in the Himalayas (Figure S13).
421
422 **Distributional range analyses.** The results of the phylogenetic generalized least squares (PGLS)
423 regressions based on the Lambda model showed no relation between the probability of any
424 hypometabolic state and distribution range size on both phylogenies. This can be observed in the
425 really low values of the adjusted R^2 (Table 2), meaning that the unexplained variance and error
426 of the model is high.
427
Amphibian assemblage climatic resilience. The results of the standardized climate anomalies
showed higher magnitudes of climate change occur in the tropics (Figure 6). However, most sites
will experience low climate change; these sites have a wide range of proportions of species with
a HS (Figure 6a). We consider that most vulnerable assemblages are sites with high climate
variance (> 2 of SED). If we take the subset of these that contain a low proportion of species
with hypometabolism (≤ 0.5), there are a total of 238 assemblages, all of which are located in the
tropics (Figure 6b). A few of these sites do have a high proportion of species with a
hypometabolic state, and, therefore, a climatically resilient assemblage.
436
Discussion

With more than 8,800 described amphibian species, we only recovered information for fewer
than 10%, clearly indicating that more information about the ecology of amphibian species is
needed. This is troublesome to say the least, not only because it increases the uncertainty in the
analyses (see below) but also because it means that for the majority of the species, we do not
have the minimum information to establish effective conservation plans and we will need to rely
on assumptions derived from < 10% of each group or class. Nevertheless, we note that these 857
species include a variety of distinct lineages from caecilians to salamandrids, pipids, and hylids
to name just a few. Another aspect to take into account is the inherent geographical bias of the
data, where most of the data is from the USA, Australia, New Zealand, and the European region.
Although, it is worth mentioning that other regions also stand out, such as South America and
South Africa, probably due to the extreme seasonality and the number of species that are known
to estivate in these regions.
450
Reconstruction of ancestral states. At the root of the tree for all amphibians the ancestral state
reconstruction showed high uncertainty for the presence of any hypometabolic state (~ 50%).
However, at the order and superfamily levels, the ancestral state reconstruction analyses provide
support for the presence of hypometabolism, > 70% for all three orders in Jetz & Pyron and in
Anura for Portik *et al.* phylogenies. Support was low for the presence of hibernation as an
ancestral state in Gymnophiona (35%) and moderate to low for the other two orders, 67% in
Caudata, and 54% and 49% for Anura in the Jetz & Pyron and Portik *et al.* phylogenies,
respectively. Support for the presence of aestivation was moderate to high in all three orders,
ranging from 63% and 50% for Anura (on each phylogeny, respectively), 68% for Caudata, and
71% for Gymnophiona.
Based on the probability of presence at the Order level, the HS seem to have been
secondarily lost multiple times within the amphibian tree; the complex pattern presented in the
different reconstruction analyses prevents us from drawing a general inference for the whole
group. As we acknowledged, there is very limited data regarding the absence of either aestivation
or hibernation, so additional data and analyses will be necessary to confidently assign losses of
the trait. For example, our OC-SVM model yielded several "false positive" species. In this
regard, if the climate is indeed highly influencing the presence of a hypometabolic strategy, and
it is somehow predictable, those "false positive" could be "true positive" reflecting a bias in the
lack of knowledge about the species' ecological traits (Hortal *et al.*, 2015).
There are differences in the reconstruction of ancestral states between aestivation and
hibernation, and these are likely a reflection of their different evolutionary histories, even though
aestivation and hibernation are both cases of an adaptive dormancy (Navas & Carvalho, 2010),
the mechanisms that trigger each of the different states of hypometabolism are completely
different. On the one hand, aestivation seems to be triggered by lack of water and excessive heat,
on the other, hibernation seems to be triggered by low temperatures, and it is commonly
associated with accessible water. Nonetheless, it is possible that the regulatory mechanisms
responsible for depressing metabolisms are the same between aestivation and hibernation (e.g.,

methylation of DNA) (Hawkins & Storey, 2020; Bell & Hellmann, 2025) but the triggering event
is different in each case. So, our results seem to agree partially with the proposal of Storey &
Storey (1990) in that if having an hypometabolic state is an ancestral state, therefore the
mechanisms and regulatory signals that are activated during metabolic depression have a
common molecular basis for all states of metabolic depressions, including aestivation,
hibernation, and some types of anoxybiosis, probably even cryobiosis. It might be possible to go
even further and predict that torpor also relies on the same mechanisms; however, little is known
in this aspect for amphibians since most of the literature about torpor is actually referring to
winter torpor, here classified as hibernation (Boutilier *et al.*, 1997; Wilkinson *et al.*, 2017) but
see Castanho and De Luca (2001).
488
Distributional range analyses. The results of the phylogenetic generalized least squares (PGLS)
regressions showed no relation between the probability of having a HS and the distribution range
size.   We expected that species with a hypometabolic strategy would have significantly larger
distribution ranges because having a hypometabolic strategy allows individuals and populations
to overcome harsh environments since they do not have to 'suffer' those conditions. Therefore,
this ability could allow them to disperse to stressful or unfavorable environments because they
are not limited by narrow environmental constraints. Nevertheless, when accounting for
phylogeny no relationship was shown.
497
Geographic distribution of species with hypometabolism.
The pattern of aestivating species richness (Figure S9) is really striking in some parts of the
world such as in the east coast of the United States where there is a high number of species that
present aestivation. In other regions such as Australia, South Africa, the arid zones of North
America and the Arid Diagonal in South America, the high number of species that present
aestivation is unsurprising due the high precipitation seasonality in those regions. The pattern of
the proportion of species that present aestivation (Figure S10) follows very closely the arid zones
of the world (Zomer et al., 2022), with obvious exceptions where no amphibians are found such
as the Sahara, Arabian and Gobi Deserts which are deemed as hyper-arid zones (Yang &
Williams, 2015).
The pattern of hibernating species richness is also interesting in some parts of the world
such as in the east coast of the United States and southeast China, where there is a high number
of species that present hibernation (Figure S12). In other parts such as the rest of North America,
Europe, and northern Asia, the high number of species that present hibernation is expected due
the high seasonality in temperature in those regions as they coincide with Köppen cold climates
map (Beck et al., 2023).
514
Amphibian assemblage climatic resilience. It is interesting that when accounting for the
variability along the periods tested (i.e., using the SED method), the results showed that the
largest differences between current and future climate is going to occur in the tropics, in other

words where the most atypical values will occur. These sites are going to experience the greatest
change in climate, according to the models, and therefore amphibian populations living there are
going to experience a substantial pressure to adapt, possibly using hypometabolic states to help
them to survive. It does make sense because much of the tropics have little variability in
temperature and precipitation compared to the higher latitudes where the inter-annual variation is
huge. In addition, most of the species with a hypometabolic strategy currently occur outside the
tropics (Figures. S10-S13). Consequently, according to our definition, the most vulnerable
assemblages are located in the tropics, specifically in the lowlands. For example, soil
temperature variation in tropical mountains like the Andes may reach almost 50 °C in one day
(Navas, 1996), and we expect that these species are likely to demonstrate some form of
hypometabolism. For example, *Atelopus carrikeri* lives at 4,500 m a.s.l. where temperatures can
be high during the day but it can be trapped occasionally in ice sheets and survives (Navas,
1997). It is important to stress that high-elevation Neotropical amphibians do not belong to a
single taxonomic group. Species from several divergent families such as Bufonidae,
Leptodactylidae, Dendrobatidae, and Hylidae are represented at elevations over 3,000 m a.s.l. in
the tropical Andes, and these high-elevation species are apparently derived from low elevation
stocks (e.g., Fouquet *et al.*, 2013; Mendoza *et al.*, 2015). Thus, given the results of the ancestral
state reconstruction, it might be possible to assume that these species were able to persist in those
conditions because they have converged on the mechanisms to survive harsh conditions.

Another aspect that is worth highlighting is the temperature spectrum of activity. Navas
(1996) showed that high-elevation anurans are capable of performing activities across a wide
range of temperatures, however low-elevation species can be active in narrower ranges of
temperature. This supports our results of assemblage climatic resilience in the sense that most
vulnerable assemblages were found in tropical lowlands. Plastic responses to past environments
shape adaptation to novel selection pressures (Coates *et al.*, 2025), therefore it is logical to
assume that if there is geographical variation in the response to climate across population, that is,
not necessarily all populations aestivate or hibernate, this plasticity may give said species a
chance to adapt to novel climate. It is important to stress that given the scarcity of the data it
might result in a less worrisome scenario for the tropical assemblages of amphibians given the
probability that more species could aestivate.

548
Limitations of the analyses. Even though we have tried to account for missing data, it is
important to stress that the results presented here have to be taken with caution as the
information used in the analyses represents fewer than 10% of the total amphibian species
described (Frost, 2024; AmphibiaWeb, 2024). Two aspects emerge from this lack of data, the
first is the high uncertainty that was generated by inferring the probabilities of a hypometabolic
strategy based only on climate. Although the inferences did correctly predict all of the species
known to have hypometabolic strategies, such uncertainty may have contributed to noise in the
dataset. The second aspect regarding uncertainty in our analyses is the assumption that species
not observed to have a HS in the wild cannot present a HS. For example, *Rhinella marina* does

not normally aestivate under natural conditions but when subjected to experimental conditions of
water stress and temperature it does aestivate (Boutilier *et al.*, 1979), but see Hilje & Arévalo-
Huezo (2012). Solving this dilemma will be crucial in the years to come when planning effective
conservation strategies, either by collecting more information in the field or by performing
experiments. It would be ideal to have information for all genera.
563
Future research directions. Aside from a call for more data on hypometabolism in amphibians,
we propose a few research themes that would be fruitful to pursue in the future. First, how long
can species aestivate or hibernate? A few accounts suggest that amphibians can survive five or
more years while estivating (Van Beurden, 1980) and estimations suggest even more (Sadowski-
Fugitt *et al.*, 2012), which conflicts with common narratives that amphibians are sensitive and
fragile organisms. Persisting in a hypometabolic state for years would provide impressive
resiliency to sudden or sustained environmental change and may start to explain how amphibians
have survived several mass extinction events. Aestivation may also help amphibians survive
wildfires. Second, could resorting to aestivation or hibernation affect longevity in populations?
Third, do amphibians have different levels of hypometabolism, for example, very shallow (days),
shallow (weeks) or deep (months to years)? In this sense should we classify torpor as an actual
hypometabolic state? Are the molecular mechanisms the same? Fourth, what are the
physiological mechanisms of the species that live in mountains, such as the Andes, where they
can experience an oscillation of temperature of more that 40 °C in one day? Finally, in the era of
huge environmental changes, will the limits of hypometabolic ability be reached in the near
future (Macip-Ríos *et al.*, 2023)? Would having a hypometabolic strategy give populations
resilience to face coming changes in climate?
Clearly the major drawback of the analyses and results presented here are the number of
species used for the analyses. Therefore, we call on our colleagues to report whether or not a
species goes into a hypometabolic state, and if possible, to test them. A review of amphibian
phenology may also identify additional cases of aestivation or hibernation.
585
586 **Supplementary material**
587
588 **File S1. Additional methods and results.** This file contains additional results.
589
590 **File S2. R scripts used for the analyses performed**.
591
592 **Table S1. Dataset of species with hypometabolic strategy information.** Compiled dataset
593 containing the species found in literature regarding hypometabolic strategy (HS): aestivation or
594 hibernation. HS was classified as **present** = confirmed hypometabolic state, **possible** = authors
595 mention the possibility but it was not confirmed, **absent** = the reference clearly mentions that the
596 species/population does not present the hypometabolic state**, unknown** = when the authors

clearly mention that it is unknown. The table also lists IUCN category (2024), name in the trees
(Jetz & Pyron, 2018 and Portik *et al.*, 2023), and the calculated probabilities for the states.

References

AmphibiaWeb. 2024. <https://amphibiaweb.org> University of California, Berkeley, CA, USA.

Beck, H.E., McVicar, T.R., Vergopolan, N., Berg, A., Lutsko, N.J., Dufour, A., *et al.* 2023.
High-resolution (1 km) Köppen-Geiger maps for 1901–2099 based on constrained CMIP6
projections. *Sci Data* **10**: e724.

Bell, A.M. & Hellmann, J.K. 2025. An Integrative Framework for Understanding the
Mechanisms and Multigenerational Consequences of Transgenerational Plasticity. *Annu
Rev Ecol Evol Syst* **50**: 97–118.

Borror, D.J. 1960. *Dictionary of word roots and combining forms*. Mayfield Publishing
Company, Mountain View, California.

Boutilier, R.G., Donohoe, P.H., Tattersall, G.J. & West, T.G. 1997. Hypometabolic homeostasis
in overwintering aquatic amphibians. *J Exp Biol* **200**: 387–400.

Boutilier, R.G., Randall, D.J., Shelton, G. & Toews, D.P. 1979. Acid-Base Relationships in the
Blood of the Toad, *Bufo marinus*: I. The Effects of Environmental CO2. *J Exp Biol* **82**:
331–344.

Castanho, M.L. & De Luca, I.M.S. 2001. Moulting Behavior in Leaf-Frogs of the Genus
Phyllomedusa (Anura: Hylidae). *Zool Anz* **240**: 3–6.

Coates, S.E.R., Comeault, A.A., Wood, D.P., Fay, M.F., Creer, S., Osborne, O.G., *et al.* 2025.
Plastic responses to past environments shape adaptation to novel selection pressures. *PNAS*
122: e2409541122.

Cowan, K.J. & Storey, K.B. 1999. Reversible phosphorylation control of skeletal muscle
pyruvate kinase and phosphofructokinase during estivation in the spadefoot toad,
Scaphiopus couchii. *Mol Cell Biochem* **195**: 173–181.

Faivovich, J., Nicoli, L., Blotto, B.L., Pereyra, M.O., Baldo, D., Barrionuevo, J.S., *et al.* 2014.
Big, bad, and beautiful: Phylogenetic relationships of the horned frogs (Anura:
Ceratophryidae). *South Am J Herpetol* **9**: 207–227.

Fitzjohn, R.G., Maddison, W.P. & Otto, S.P. 2009. Estimating trait-dependent speciation and
extinction rates from incompletely resolved phylogenies. *Syst Biol* **58**: 595–611.

Forti, L.R., Pontes, M.R., Augusto-Alves, G., Martins, A., Hepp, F. & Szabo, J.K. 2022. Data
collected by citizen scientists reveal the role of climate and phylogeny on the frequency of
shelter types used by frogs across the Americas. *Zoology* **155**: e126052.

Fouquet, A., Cassini, C.S., Haddad, C.F.B. *et al.* 2013 Species delimitation, patterns of
diversification and historical biogeography of the Neotropical frog genus *Adenomera*
(Anura, Leptodactylidae). *J Biogeogr* **41**:855–870

Frost, D.R. 2024. Amphibian Species of the World: an Online Reference. Version 6.2. [Accessed
October 10, 2024]

García-Rodríguez, A., Lenzner, B., Velasco, J.A., Schertler, A., Omer, A., Seebens, H., *et al.*
2025. The global distribution patterns of alien vertebrate richness in mountains. *Nature
Commun* **16**: e1977.

Geiser, F. 2020. Seasonal expression of avian and mammalian daily torpor and hibernation: not a
simple summer-winter affair. *Front Physiol*, **11**: 436.

Geiser, F. 2021. *Ecological physiology of daily torpor and hibernation* (Vol. 317). Springer, Berlin, Germany.

Hawkins, L.J. & Storey, K.B. 2020. Advances and applications of environmental stress adaptation research. *Comp Biochem Physiol A Mol Integr Physiol* **240**: e110623.

He, Y., Liao, Z., Chen, X. & Chen, Y. 2023. Climatic debts for global amphibians: Who, where and why? *Biol Conserv* **279**: e109929.

Hembree, D.I., Martin, L.D. & Hasiotis, S.T. 2004. Amphibian burrows and ephemeral ponds of the Lower Permian Speiser Shale, Kansas: Evidence for seasonality in the midcontinent. *Palaeogeogr Palaeoclimatol Palaeoecol* **203**: 127–152.

Henrici, A.C. & Haynes, S.R. 2006. *Elkobatrachus brocki*, a new pelobatid (Amphibia: Anura) from the Eocene Elko Formation of Nevada. *Ann Carnegie Mus* **75**: 11–35.

Hortal, J., De Bello, F., Diniz-Filho, J.A.F., Lewinsohn, T.M., Lobo, J.M. & Ladle, R.J. 2015. Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity. *Annu Rev Ecol Evol Syst* **46**: 523–549.

IUCN. 2024. The IUCN Red List of Threatened Species. Version Version 2023-1. http://www.iucnredlist.org. Downloaded October 11, 2024.

Jetz, W. & Pyron, R.A. 2018. The interplay of past diversification and evolutionary isolation with present imperilment across the amphibian tree of life. *Nat Ecol Evol* **2**: 850–858.

Jiang, C., Storey, K.B., Yang, H. & Sun, L. 2023. Aestivation in Nature: Physiological Strategies and Evolutionary Adaptations in Hypometabolic States. *Int J Mol Sci* **24**: e14093.

Judd, E.J., Tierney, J.E., Lunt, D.J., Montañez, I.P., Huber, B.T., Wing, S.L., *et al.* 2024. A 485-million-year history of Earth's surface temperature. *Science (1979)* **385**: eadk3705.

Lewis, P.O. 2001. A likelihood approach to estimating phylogeny from discrete morphological character data. *Syst Biol* **50**: 913–925.

Lynch, J.D. & Suárez-Mayorga, Á.M. 2002. Análisis biogeográfico de los anfibios paramunos. *Caldasia* **24**: 471–480.

Lillywhite, H.B. 1975. Physiological correlates of basking in amphibians. *Comp. Biochem. Physiol. A* **52**: 323–330.

Macip-Ríos, R., Butterfield, T. & Raya-García, E. 2023. How Aestivation Evolved in Turtles: A Macroevolutionary and Morphological Approach. *Evol Biol* **50**: 381–394.

Marchetti, L., MacDougall, M.J., Buchwitz, M., Canoville, A., Herde, M., Kammerer, C.F., *et al.* 2024. Origin and early evolution of vertebrate burrowing behaviour. *Earth Sci Rev* **250**: e104702.

Mayhew, W.W. 1965. Adaptations of the Amphibian, *Scaphiopus couchi*, to Desert Conditions'. *Am Midl Nat* **74**: 95–109.

McClanahan, L. 1972. Changes in body fluids of burrowed Spadefoot toads as a function of soil water potential. *Copeia* **1972**: 209–216.

Mendoza, Á.M., Ospina, O.E., Cárdenas-Henao, H., & García-R, J.C. 2015. A likelihood inference of historical biogeography in the world's most diverse terrestrial vertebrate genus: Diversification of direct-developing frogs (Craugastoridae: *Pristimantis*) across the Neotropics. *Mol Phylogenetics Evol* **85**: 50-58.

Meyer, D., Dimitriadou, E., Hornik, K., Weingessel, A. & Leisch, F. 2024. e1071: Misc Functions of the Department of Statistics, Probability Theory Group (Formerly: E1071), TU Wien. https://CRAN.R-project.org/package=e1071.

Mokhatla, M.M., Rödder, D. & Measey, G.J. 2015. Assessing the effects of climate change on
distributions of Cape Floristic Region amphibians. *S Afr J Sci* **111**: 7.
Naimi, B., Hamm, N.A.S., Groen, T.A., Skidmore, A.K. & Toxopeus, A.G. 2014. Where is
positional uncertainty a problem for species distribution modelling? *Ecography* **37**: 191–
203.
Navas, C.A. 1996. The Effect of Temperature on the Vocal Activity of Tropical Anurans: A
Comparison of High and Low-Elevation Species. *J Herpetol* **30**: 488–497.
Navas, C.A. 1997. Thermal extremes at high elevations in the Andes: Physiological ecology of
frogs. *J Therm Biol* **22**: 467–477.
Navas, C.A. & Carvalho, J.E. 2010. *Aestivation Molecular and Physiological Aspects* (C. A.
Navas & J. E. Carvalho, eds). Springer-Verlag, Berlin Heidelberg.
Ochoa-Ochoa, L.M., Rodríguez, P., Mora, F., Flores-Villela, O., Whittaker, R.J., Rodriguez, P.,
et al. 2012. Climate change and amphibian diversity patterns in Mexico. *Biol Conserv* **150**:
94–102.
Ochoa-Ochoa, L.M. & Velasco, J.A. 2024. Long-term stability in protected-areas? A vision from
American/New World amphibians. *Geogr Sustain* **5**: 673–683.
Pinder, A.W., Storey, K.B. & Ultsch, G.R. 1992. Estivation and Hibernation. In: *Environmental
Physiology of the Amphibians* (M. E. Feder & W. W. Burggren, eds), pp. 250–274. The
University of Chicago Press, Chicago.
Portik, D.M., Streicher, J.W., & Wiens, J.J. 2023. Frog phylogeny: a time-calibrated, species-
level tree based on hundreds of loci and 5,242 species. *Mol Phylogenet Evol* **188**: 107907.
R Core Team. 2024. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna Austria.
Rangel, T.F., Colwell, R.K., Graves, G.R., Fučíková, K., Rahbek, C. & Diniz-Filho, J.A.F. 2015.
Phylogenetic uncertainty revisited: Implications for ecological analyses. *Evolution* **69**:
1301–1312.
Revell, L.J. 2025. Ancestral State Reconstruction of Phenotypic Characters. *Evol Biol* **52**: 1–25.
Revell, L.J. 2024. phytools 2.0: an updated R ecosystem for phylogenetic comparative methods
(and other things). *PeerJ* **12**: e16505.
Reynolds, J.A. 2019. Noncoding rna regulation of dormant states in evolutionarily diverse
animals. *Biological Bulletin* **237**: 192–209.
Sadowski-Fugitt, L.M., Tracy, C.R., Christian, K.A. & Williams, J.B. 2012. Cocoon and
epidermis of australian *Cyclorana* frogs differ in composition of lipid classes that affect
Water loss. *PBZ* **85**: 40–50.
Seymour, R.S. 1973. Gas Exchange in Spadefoot Toads beneath the Ground. *Copeia* **1973**: 452–
460.
Smith, C., Ochoa-Ochoa, L.M. & Velasco, J.A. 2025. Using bioclimatic indicators to analyse the
presence of climatic mismatches amongst local amphibian assemblages of America. *Rev
Lat Herp* **8**: 114–128.
Storey, K.B. & Storey, J.M. 2012. Aestivation: Signaling and hypometabolism. *J Exp Biol* **215**:
1425–1433.
Storey, K.B. & Storey, J.M. 1990. Metabolic Rate Depression and Biochemical Adaptation in
Anaerobiosis, Hibernation and Estivation. *Q Rev Biol* **65**: 145–174.
Storey, K.B. & Storey, J.M. 2004. Metabolic rate depression in animals: Transcriptional and
translational controls. *Biol. Rev. Camb. Philos. Soc.* **79**: 207–233.

Storey, K.B. & Storey, J.M. 2010. Metabolic Regulation and Gene Expression During
Aestivation. In: *Aestivation: Molecular and Physiological Aspects* (C. A. Navas & J. E.
Carvalho, eds), pp. 25–45. Springer.
Taheri, S., Naimi, B. & Araújo, M.B. 2024a. climetrics: an R package to quantify multiple
dimensions of climate change. *Ecography* **2024**: e07176.
Taheri, S., Naimi, B. & Araujo, M.B. 2024b. climetrics R package. https://CRAN.R-
project.org/package=climetrics.
Tan, W.C., Herrel, A. & Rödder, D. 2023. A global analysis of habitat fragmentation research in
reptiles and amphibians: what have we done so far? *Biodivers Conserv* **32**: 439–468.
Van Beurden, E.K. 1980. Energy Metabolism of Dormant Australian Water-Holding Frogs
(*Cyclorana platycephalus*). *Copeia* **1980**: 787–799.
Velasco, J.A., Estrada, F., Calderón-Bustamante, O., Swingedouw, D., Ureta, C., Gay, C., *et al.*
2021. Synergistic impacts of global warming and thermohaline circulation collapse on
amphibians. *Commun Biol* **4**: e141. Sp
Vitt, L.J. & Caldwell, J.P. 2014. *Herpetology. An Introductory Biology of Amphibians and*
Reptiles, Fourth Edition. Elsevier.
Wells, K. D. 2007. *The ecology and behavior of amphibians*. University of Chicago press.
Wilkinson, A., Hloch, A., Mueller-Paul, J. & Huber, L. 2017. The effect of brumation on
memory retention. *Sci Rep* **7**: e40079.
Williams, J.W., Jackson, S.T., Kutzbach, J.E. & Schneider, S.H. 2007. Projected distributions of
novel and disappearing climates by 2100 AD. *PNAS* **104**: 5738–5742.
Withers, P.C. 1995. Cocoon Formation and Structure in the Aestivating Australian Desert Frogs,
Neobatrachus and *Cyclorana*. *Aust. J. ZooL* **43**: 429–470.
Withers, P.C. 1998. Evaporative water loss and the role of cocoon formation in Australian frogs.
Aust J Zool **46**: 405–418.
Withers, P.C. & Cooper, C.E. 2010. Metabolic depression: a historical perspective. *Prog Mol*
Subcell Biol **49**: 1–23.
Yang, X. & Williams, M. 2015. Landforms and processes in arid and semi-arid environments.
Catena (Amst) **134**: 1–3.
Zimicz, N., Fabrezi, M., Aramayo, A., Bianchi, C., Hongn, F. & Montero-López, C. 2024.
Ceratophryid frogs in the late Miocene of central Andes of Argentina: insights on the
paleoenvironment of Palo Pintado Formation. *Hist Biol* **30 Sep 2024**: e2403590.
Zomer, R.J., Xu, J. & Trabucco, A. 2022. Version 3 of the Global Aridity Index and Potential
Evapotranspiration Database. *Sci Data* **9**: e409.

Box 1. The use of different terms of dormancy.

Hypometabolism is a general phenomenon that is not unique to amphibians and includes a wide range of metabolic depressions such as hibernation, aestivation, dormancy, brumation and torpor. In general, dormancy is considered to be a term that encompasses all hypometabolic states, and the key aspect is that dormancy is preventive and anticipates stressful physiological conditions to avoid harmful changes in physiology (Navas & Carvalho, 2010; Storey & Storey, 2012; Reynolds, 2019). Torpor, from the Latin *torpōr* (Borror, 1960), means to be numb, inactive, or dull, and is referred to as a daily rest; it was commonly used as synonym of hibernation (i.e., winter torpor; Pinder *et al.*, 1992; Boutilier *et al.*, 1997; Withers

& Cooper, 2010; Wilkinson *et al.*, 2017), yet it should not be considered a sustained hypometabolic state such as aestivation or hibernation (Castanho & De Luca, 2001; Navas & Carvalho, 2010 but see Geiser, 2020, 2021). Aestivation or estivation, both derived from the Latin *aestās* that mean "pertaining to summer" (Borror, 1960), is defined as a state of reduced metabolism occurring in hot and dry conditions. Nevertheless, Pinder et al. (1992) consider that aestivation occurs in both hot and cold conditions and that its main driver is the lack of water. For example, toads of the genus *Scaphiopus* aestivate during winter (Seymour, 1973). Aestivation can occur when the amphibian burrows itself into the soil or seeks shelter in crevices or below logs and rocks, but it can also occur out in the open either on land or on plants, although it is much less common (Navas & Carvalho, 2010). Finally, brumation and hibernation both come from Latin words meaning "related to winter" (Borror, 1960), the former from *bruma* and the latter from *hībern*. The use of the terms brumation and hibernation may be sometimes ambiguous, as they refer to the lethargic state that some animals assume during cold conditions. However, brumation was proposed to refer to obligate hibernators that cannot control their body temperature during winter dormancy rather than those that can (Mayhew, 1965).

Tables

Table 1. Comparative table from the ancestral state reconstruction for aestivation, hibernation and hypometabolism for all amphibians with Jetz and Pyron (2018) phylogeny and for just Anura with Portik *et al.* (2023) phylogeny.

	Aestivation						Hibernation						Hypometabolism					
	Jetz & Pyron (2018)			Portik *et al.* (2023)			Jetz & Pyron (2018)			Portik *et al.* (2023)			Jetz & Pyron (2018)			Portik *et al.* (2023)		
	Presence	Absence	SD	Presence	Absence	SD	Presence	Absence	SD	Presence	Absence	SD	Presence	Absence	SD	Presence	Absence	SD
Amphibia	0.483	0.517	0.091				0.515	0.485	0.147				0.483	0.517	0.098			
Anura	0.628	0.372	0.112	0.502	0.498	0.230	0.540	0.460	0.145	0.479	0.521	0.099	0.731	0.269	0.137	0.709	0.291	0.146
Caudata	0.683	0.317	0.120				0.672	0.328	0.185				0.787	0.213	0.123			
Gymnophiona	0.717	0.283	0.162				0.353	0.647	0.154				0.773	0.227	0.147			
Bombinatoroidea	0.642	0.358	0.114	0.519	0.481	0.265	0.572	0.429	0.175	0.486	0.514	0.112	0.788	0.212	0.060	0.736	0.264	0.124
Cryptobranchoidea	0.641	0.359	0.120				0.702	0.298	0.164				0.795	0.205	0.121			
Gymnophiona	0.717	0.283	0.162				0.353	0.647	0.154				0.773	0.227	0.147			
Hyloidea	0.644	0.356	0.199	0.634	0.366	0.387	0.151	0.849	0.215	0.258	0.743	0.350	0.429	0.571	0.221	0.661	0.339	0.406
Leiopelmatoidea	0.660	0.341	0.147	0.494	0.506	0.249	0.486	0.515	0.148	0.458	0.542	0.108	0.798	0.202	0.129	0.680	0.320	0.154
Myobatrachoidea	0.803	0.197	0.142	0.604	0.397	0.402	0.786	0.214	0.218	0.325	0.675	0.298	0.842	0.159	0.128	0.731	0.269	0.313
Paleobatoidea	0.625	0.375	0.112	0.540	0.460	0.376	0.711	0.289	0.127	0.549	0.451	0.139	0.798	0.202	0.124	0.752	0.248	0.143
Pipoidea	0.648	0.352	0.119	0.448	0.552	0.305	0.550	0.450	0.171	0.425	0.575	0.107	0.761	0.239	0.130	0.715	0.285	0.142
Ranoidea	0.541	0.460	0.143	0.820	0.180	0.256	0.467	0.533	0.270	0.383	0.617	0.197	0.727	0.273	0.172	0.618	0.382	0.194
Salamandroidea	0.712	0.288	0.123				0.726	0.274	0.191				0.804	0.196	0.124			
Sirenoidea	0.855	0.145	0.062				0.881	0.119	0.102				0.903	0.097	0.051			

Table 2. Summary of the phylogenetic generalized least squares (PGLS) regressions based on the Lambda model. Colums represent the average and standard deviation of each resulting parameter.

	σ^2	Lambda	logLik	Adjusted R^2
Jetz & Pyron (2018)				
Aestivation	0.070 ± 0.004	0.670 ± 0.016	-15299.414 ± 10.069	0.071 ± 0.001
Hibernation	0.064 ± 0.004	0.644 ± 0.016	-15181.902 ± 8.571	0.106 ± 0.001
Hypometabolism	0.066 ± 0.004	0.652 ± 0.016	-15254.762 ± 9.189	0.084 ± 0.001
Portik *et al*. (2023)				
Aestivation	0.118 ± 0.008	0.856 ± 0.010	-9431.548 ± 4.733	0.115 ± 0.001
Hibernation	0.122 ± 0.008	0.842 ± 0.011	-9603.590 ± 4.116	0.034 ± 0.001
Hypometabolism	0.103 ± 0.007	0.829 ± 0.012	-9380.125 ± 3.979	0.139 ± 0.001

Figure legends

Figure 1. Representation of the different levels of metabolic depressions currently reported in amphibians. The ovals represent the range in which each state would be denominated as such. For example, there would be a wide range of aestivation and hibernation levels. Illustrations were generated using the Adobe Illustrator generative AI.

Figure 2. Resilience of amphibian assemblages to cope with climate change. Highly resilient assemblages are expected to have a high proportion of species with presence of hypometabolism. In contrast, highly vulnerable assemblages would be where very few species have hypometabolism. Hypothetical lines are drawn to represent two types of vulnerability transitions: abrupt threshold (dashed) and elastic threshold (dotted).

Figure 3. Probabilities from the ancestral state reconstruction and age node estimation with the selected from Jetz and Pyron (2018) phylogeny for A) aestivation, B) hibernation and C) hypometabolism. Result from the MCMC approach to sample character histories from their posterior probability distribution (multiSIMMAP analyses) for 100 trees.

Figure 4. Stochastic character mapping for the hypometabolic strategies in amphibians (red = presence; blue absence) based on a single tree. The internal colored ring depicts the current state of each tip, while the external ring indicates major groups as featured in Jetz and Pyron (2018): 1. Gymnophiona, 2. Cryptobranchoidea, 3. Sirenoidea, 4. Salamandroidea, 5. Leiopelmatoidea, 6. Bombinatoroidea, 7. Pipoidea, 8. Paleobatoidea, 9. Myobatrachoidea, 10. Hyloidea, and 11. Ranoidea.

Figure 5. Average proportion of nodes that were inferred as presence as the most likely state for one of the two hypometabolic strategies, according to the joint ancestral state reconstruction for selected tree for the Jetz & Pyron (2018) phylogeny. The temperature bar is the same as featured in Judd *et al.* (2024). Curves were smoothed using the smoothing splines method and shaded areas are derived from the confidence interval (95%). Dotted lines with the fish skeleton indicate major extinction events, the Triassic–Jurassic (Tr-J) 200 Myr, and the Cretaceous–Paleogene (K–Pg) 66 Myr.

Figure 6. Standardized Euclidean distance based on temporal differences for each climate parameter which are standardized using the local inter-annual standard deviation for that parameter in both periods of time, in this case t1 (1990-2020) and t2 (2070-2100), this can be interpreted as magnitude of climate change. The higher the standardized local anomaly scores the larger the predicted changes in temperature and precipitation. The most vulnerable amphibian assemblages based on the low proportion of species with a HS, only literature data (bottom right in A) are highlighted with a red star in B.

823 **Figures**

824 **Figure 1**

825
826
827 **Figure 2**

829
830
831
832   **Figure 3**

833
834   **Figure 4**

846 **Figure 5**

868 **Figure 6**
869