Collective bacterial condensation is fundamentally constrained by the emergence
of active turbulence.

Nir Livne and Ady Vaknin*

The Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

Running title: active turbulence constrains bacterial condensation.

*Corresponding author:

Mailing address: The Racah Institute of Physics,
11   The Hebrew University,
12   Edmond J. Safra campus
13   Jerusalem 91904, Israel
14
Phone: (972) 658-4266
FAX: (972) 658-6168
E-mail: *avaknin@mail.huji.ac.il*

Keywords: bacterial chemotaxis | collective condensation | correlated swimming | active turbulence

Collective bacterial condensation arises from positive feedback between the ability of

bacteria to generate chemical gradients and their chemotactic ability to track those

gradients, resulting in the spontaneous formation of local cell accumulations. Here, by

following the bacterial condensation dynamics in uniform acidic environments, we show

that condensation is critically constrained by the spontaneous emergence of correlated

bacterial swimming. Under the unique conditions prevailing within the condensate—

characterized by sharp cell-density gradients and the absence of physical boundaries—

correlated bacterial motion leads to vortex-like cell motion with an additional pronounced

radial component directed down the cell-density gradient. This correlated motion

induces active turbulence behavior associated with local fluid motion that broadens the

condensate and expels non-chemotactic bacteria. However, when cells are embedded in

a porous environment, correlated motion is suppressed, allowing spontaneous

condensation to proceed further and form exceptionally dense bacterial accumulations.

Overall, these findings highlight the fundamental interplay between self-generated

bacterial condensation and correlated swimming.

Significance statement

Self-induced condensation and active turbulence are two prominent forms of collective bacterial

behaviors. However, their interplay has not been experimentally studied. Here, we show that

collective bacterial condensation gives rise to correlated swimming, which in turn fundamentally

limits further condensation. Furthermore, we demonstrate that in dilute porous environments—

common in natural bacterial habitats—correlated swimming is strongly suppressed, enabling the

spontaneous formation of extremely dense cell condensates. These condensates may serve as

a basis for the development of more structured bacterial communities.

Introduction

Motile bacteria navigate their environment by tracking chemical gradients toward regions they deem favorable (1-4). This behavior, termed chemotaxis, influences many aspects of bacterial ecology (5, 6) and human health (7-9). Even in spatially uniform environments, bacteria can collectively follow self-generated gradients, allowing them to accelerate their expansion into resource-rich regions (10, 11), or, alternatively, condense into focal points by either secreting beneficial (attractant) effectors (12, 13) or neutralizing harmful (repellent) effectors (14). Notably, in contrast to the inherently dynamic bacterial distributions triggered by exogenous chemoeffector release (15) or in expanding bacterial bands (16), bacterial condensation can lead to sharp but quasi-static bacterial accumulation (14). Condensation relies on positive feedback between self-generated gradients and the chemotactic responses of the bacteria to those gradients (Fig. 1A) that, in principle, can lead to a divergence in the local cell density (17). Studies of low cell-density and transient cell accumulations, limited by oxygen depletion, suggested that a fundamental limit on such cell condensation arises from the fact that chemotaxis itself is a non-local behavior (18). However, the constraints on bacterial condensations, under conditions of oxygen abundance, remain unclear. Here, we show that under these conditions, bacterial condensation is, in fact, limited by the emergence of correlated bacterial swimming and active turbulence.

Escherichia coli (E. coli) cells use rotating helical filaments (flagella) that bundle at the back of the cell and propel it forward. However, during stochastic episodes of counter-rotation the bundle comes apart, causing random reorientation of the cell. These two swimming modes—forward swimming ('run') and random reorientation ('tumble')—together generate an active random walk (19) with a typical run length (~10–30 µm). In the presence of effector gradients, the probability of tumbling is modulated to bias the random walk toward beneficial effectors and away from harmful ones. In particular, bacteria can migrate towards neutral pH (20, 21), which along with their ability to neutralize the environment, can lead to their

spontaneous, self-induced, condensation (14). Notably, this behavior is readily observed only

under conditions that the bacterial suspension is coupled to a large and cell-free environment

that keeps the pH level within the dynamic range of the sensory detection and prevents oxygen

depletion.

On the other hand, at high cell density, bacteria tend to exhibit correlated swimming and

flocking-like behavior (22-24), which can lead to turbulent-like motion often termed 'active

turbulence' (25, 26). However, the interplay between bacterial condensation and correlated

swimming was scarcely studied and remains unclear. Bacterial correlated swimming arises from

direct steric interactions and hydrodynamic interactions between cells. With notable exceptions

(27, 28), this behavior has mostly been studied in uniformly distributed populations confined by

physical boundaries. In contrast, bacterial condensation leads to highly non-uniform cell

distributions in the absence of explicit physical boundaries. Numerical studies have suggested

that dilute self-attracting run-and-tumble particles with rather weak chemotactic strength

aggregate less efficiently when self-induced fluid motion is considered (29). In addition,

experiments have suggested that cell alignment may constrain the angular rotation of individual

cells and thus reduce their chemotactic efficiency in uniform cell suspensions and shallow

effector gradients (30). However, other studies have shown that the tendency of cells to align

may, in fact, enhance their taxis efficiency toward a local source (31). Thus, how correlated

bacterial swimming may influence their collective condensation remains unclear.

Here, we investigate the limits of bacterial condensation under conditions where oxygen

deprivation and effector-saturation are not limiting factors. As a model, we used the recently

described assay for bacterial condensation in uniform acidic environments (14). We show that

stable bacterial condensation is sharply constrained by the emergence of correlated bacterial

swimming during cell accumulation. This correlated swimming gives rise to substantial

turbulent-like fluid motion, with averaged velocity characterized by vortex-like motion and with an

additional velocity component in the outward direction aligned with the downward cell-density

gradient. This fluid motion effectively exerts outward physical forces on the cells that counter the

inward chemotaxis force. Interestingly, at very high cell densities the condensate becomes ring-

shaped; yet, the radial component of the collective swimming was still aligned with the cell-

density gradient. However, the characteristic vorticity in these condensates reverted between

the outer and inner regions of the ring. Finally, we show that in porous environments, bacterial

correlated motion is diminished without significantly affecting the chemotactic performance of

cells. Consequently, bacterial condensation is considerably enhanced in porous media,

approaching the low cell-density limit but with significantly higher cell densities.

Results

Bacterial condensation is constrained by the emergence of correlated swimming. To test

the constraints on bacterial condensation in a controlled environment, we used the setup

illustrated in Fig. 1B (see ref. 14). In this setup, free-swimming cells are confined within a

shallow liquid layer (typically 120 μm thick), sandwiched between a glass slide and a larger

hydrogel layer, which together form a uniformly low-pH environment (pH 5.2). Under these

conditions, bacteria tend to collectively condense due to inherent feedback between the ability of

the bacteria to neutralize their local pH and their chemotactic ability to swim towards neutral pH

(14). While cells cannot penetrate the dense hydrogel mesh (3 % agarose), protons and oxygen

readily diffuse across the thin bacterial layer and into the gel, preventing their rapid depletion

inside the bacterial condensate and thereby enabling quasi-stable, high-density cell

condensation (Fig. 1B, right plot).

By following the peak cell-density (ρ_{max}) and the width at half-maximum (*W*) of the cell

distribution we identified two distinct temporal phases of the condensation (Fig. 1C). Initially, in

phase 1, the peak cell-density ρ_{max} increased while the width *W* decreased, reflecting strong

bacterial condensation. Then, beyond a critical time point (t_{sat}), in phase 2, the width *W* abruptly

plateaued while ρ_{max} continued to rise, reflecting continued cell accumulation without further

narrowing of the condensate. These trends are further demonstrated in Fig. 1D, where the width

of the condensation is plotted against the maximum cell density for several repetitions. Notably,

the saturation value of the condensate width was substantially larger than the low-cell-density

limit previously reported in ref. 18. The spatial profile of the bacterial accumulation also reflected

this dynamical transition (Fig. 1D, inset). By normalizing each profile by ρ_{max} and the width by

W, the profiles collapsed onto a single curve in each phase, but these cumulative curves were

distinct between the two phases.

The dynamic transition described in Fig. 1D, was clearly associated with the appearance

of significant density fluctuations within the condensate (Fig. 1E). Initially (phase 1), the

bacterial condensate was smooth and stable (Fig. S1). However, beyond the critical time point

t_{sat}, pronounced spatial non-uniformity developed, with dynamic cell-density fluctuations on a

scale of approximately 20-100 μm (Fig. 1E, see also Supplementary Movie 1). These

fluctuations indicate correlated swimming of large groups of cells. Analysis of these dynamic cell

density fluctuations using particle image velocimetry (PIV; see Materials and Methods) revealed

that, on-average (~ 30 sec), cell motion is characterized by a vortex-like clockwise rotation of the

entire condensate with an additional radial component (Fig. 1E). Interestingly, the emergence of

the cell-density fluctuations and correlated swimming coincided with the dynamic transition at

t_{sat} (Fig. 1E).

Fluid motion associated with the bacterial correlated swimming. To characterize the fluid-

motion caused by the bacterial correlated swimming, we mixed passive green fluorescent beads

(1 μm in diameter) within the bacterial population, which in this case expressed red fluorescent

proteins (Fig. 2), and analyzed their motion by particle tracking (Materials and Methods).

Typically, the beads' trajectories included periods of local motion interrupted by approximately

directional movements, termed 'jets' (Fig. 2A). To elucidate the effect of the bacterial motion on

the beads, we evaluated for each jet the ratio $Pe = \upsilon \cdot L/D_{beads}$ (Péclet number), where υ and L are

the characteristic velocity and length of the jet, respectively, and D_{beads} is the diffusion constant

of the beads. This Pe characterizes the time required for a bead to traverse a typical distance

by fluid flow compared to the typical diffusion time associated with the same distance. The

cumulative distribution of Pe is shown in Fig. 2A (right plot) for trajectories within (green) or

outside (blue) the condensate region. Clearly, the beads' motion was substantially expedited

within the condensate.

Turbulent-like fluid motion can potentially also modulate the effector (H^+) distribution. To

evaluate the potential significance of this effect, we evaluated the Pe parameter for the protons

($D_{protons} \sim 10^3 \cdot D_{beads}$), assuming protons follow the local fluid motion. Fig. 2B presents the

cumulative number of jets per bead, integrated over the diffusion time of protons across the

condensate $t_D = W^2/4D_{H^+}$, where W is the width of the condensate. These data show that, on

average, each bead in the condensate experiences at least one substantial jet with $Pe > 0.5$

over the time t_D. We therefore conclude that, while cell-induced fluid motion can influence the

effector distribution, it is most likely not dominant.

As with the bacterial collective motion, averaging the beads' motion in time (90 sec)

reveals a vortex like fluid flow in the condensate (Fig. 2C–D). The spatial distributions of the

velocity components are shown in Fig. 2C, and the corresponding profile of the angular velocity

is shown in Fig. 2D, revealing a coordinated vortex like motion. Notably, in addition to the

tangential velocity contributing to vorticity, the flow field exhibited a substantial outward radial

component peaking at ~ 0.85 $\mu m \cdot s^{-1}$ near its half-width and extended throughout the

condensate. This outward radial velocity is comparable to $E.\ coli's$ chemotactic drift velocity (19,

30). Additionally, we noted that above the condensed cells, which tended to occupy the lower-

half region of the fluid layer (~ 60 µm), the tracer beads exhibit an inward flow (Fig. 2E;

Supplementary Movie 2). Thus, the fluid motion in the vicinity of the condensate seem to exhibit

a three-dimensional toroidal pattern (Fig. 2F).

Correlated swimming leads to expulsion of non-chemotactic cells. Given the observed

fluid motion, and particularly the outward radial component, we tested its effect on the cells

within the condensate. We repeated the condensation experiments using a mixture of

chemotactic (wild-type, red-labeled) and low density non-chemotactic, smooth-swimming cells

(green-labeled). Because non-chemotactic cells are not subjected to the inward chemotactic

force, we expected them to be more sensitive to influence by the fluid motion. We found that as

the wild-type cells condensed, the non-chemotactic cells were expelled from the condensate

(Fig. 3). Additional run-and-tumble but non-chemotactic strain was tested (32), also showing

significant depletion, see Fig. S2 and related text. We expect this outward force to act also on

chemotactic cells; however, in their case, this force is balanced by an inward chemotactic force

that drives condensation. Still, such an outward force may influence the width of the

condensate.

Porous environment suppresses correlated swimming and enhances condensation.

Porous media is expected to dampen fluid flows due to the added drag with the porous scaffold

(33) and potentially inhibit correlations between bacteria. For example, porous media have been

suggested to inhibit bioconvection (34). Thus, to further test the effect of correlated swimming

on bacterial condensation, we conducted experiments in which cells were immersed in a thin

porous environment formed by a dilute agarose hydrogel instead of liquid (Fig. 4A). The rest of

the setup was unchanged. Using the channel chemotaxis assay described in ref. 21, we verified

that the bacterial chemotaxis performance in this porous environment was comparable to that in

liquid (Fig. S3). Moreover, by introducing opposing chemoeffectors at one end of the channel,

we triggered low cell-density accumulation. The agarose hydrogel had no detectable effect on

the cell-diffusion-driven broadening of this accumulation (Fig. S3).

In contrast, the porous environment had a dramatic effect on bacterial condensation and

correlated swimming (Fig. 4). Although the condensation proceeded at a rate similar to that in

liquid, in hydrogel it saturated only when the width of the condensate approached the low-

density limit of ~180 µm (18) (Fig. 4A-B). Correspondingly the cell density in these tightly

condensed bacterial populations became extremely high. As in liquid, in the porous

environment the shape of the bacterial condensate was modified when it reached its limit

(Fig. S4). Finally, the saturated condensates were also more compact in the vertical direction,

spanning only ~30 µm. Overall, in porous environments, the bacterial population spontaneously

condensed to its fundamental limit, reaching exceptionally high cell densities (Fig. S5), while the

cell population remained largely dynamic (Supplementary Video 3).

The enhanced condensation observed in porous environments was indeed associated

with a suppression of correlated swimming and associated fluid motion (Fig. S6 and Fig. 4C).

Furthermore, when we repeated the experiments with mixed populations of chemotactic and

non-chemotactic cells in the hydrogel environment, we found that, despite the strong

condensation of the chemotactic cells, the expulsion of the non-chemotactic cells was

significantly weaker (Fig. 4D). Taken together, these results further support the notion that

bacterial condensation in liquid is limited by the emergence of correlated swimming, rather than

merely the locally elevated density of the chemotactic cells.

Ring-like condensation and the potential role of oxygen-taxis. To test the condensation at

still higher cell densities, we repeated the experiments while increasing the initial (uniform) cell

density (Fig. 5). Since the well diameters used here were chosen such that only one

condensate is formed (14), we expected the overall cell density in the condensate to increase

accordingly. Yet, we find that in both liquid and porous media, following the initial condensation,

once enough cells accumulate, the peak density ceases to increase while the condensate

broadens and turns into a stable ring (Fig. 5). A likely cause for such ring formation is the

development of oxygen gradients around the condensate that induce an outward chemotactic

force (aerotaxis) (35). Compared with the liquid medium, in the porous environment ring

formation occurred at much higher cell density but at significantly lower total number of cells

(Fig. 5A). This difference may result from the fact that turbulent swimming in liquid weakens the

net inward force and can be more easily be dominated by aerotaxis. Additionally, in liquid,

aeration of the condensate by bacteria-mediated convection may support the accumulation of

higher cell numbers prior to ring formation.

Interestingly, we found that in liquid, following ring formation, the vorticity reverses

between the inner and outer regions of the 'aerotaxis' ring (Fig. 5C), including both the radial

and tangential components. We note, however, that the reversal of both components is

consistent with them being determined relative to the cell density gradient: in both cases the

collective motion flows radially down the cell density gradient and tends to exhibit also a

rightward lateral deflection (illustrated in Fig. 5C).

Simplified model for condensate broadening. To test whether the observed cell-induced fluid

motion (Fig. 2) can indeed lead to repulsion of non-chemotactic cells (Fig. 3) and broaden the

condensation of wild-type cells, we used Keller-Segel type chemotaxis model (36) that describes

the bacterial condensation in such acidic environment (14, 37):

240
$$\frac{\partial \rho}{\partial t} = \vec{\nabla} \cdot \left(D\vec{\nabla}\rho - \rho\vec{v}_D \right) \quad (1)$$
$$\frac{\partial c}{\partial t} = D_c \nabla^2 c - r_H \rho c \quad (2)$$

Here, D is the active diffusion of the cells, v_D is their drift velocity, D_c is the effective proton

diffusion coefficient, and r_H is the proton-removal rate. Note that bacteria are not growing in

these experiments. Following ref. 38, the chemotaxis-induced drift velocity was modeled as

$\vec{v}_D = \beta\vec{\nabla}\ln\left(c + K_{off}/c + K_{on}\right)$ where K_{on} and K_{off} are the effective dissociation constants for the

chemoreceptors in the ON or OFF states, and β is the chemotaxis coefficient. The proton-

removal rate was given the empirical form $r_H = \gamma c^{1/2}$ (14). The cell density dynamics was

assumed to be two dimensional and radially-symmetric, occupying the top layer of a three-

dimensional proton field. For more details see Supplementary Information. This model

accurately describes the shape of condensates in phase 1 (Fig. 6A). However, it predicts much

sharper condensation and a different shape than those observed experimentally in phase 2

(Fig. 6B, dashed blue line).

Thus, considering the random fluid motion at short time scales with the added outward

averaged drift component observed in Fig. 2, we suggest that bacteria subjected to such fluid

motion experience not only the inward chemotactic force, but also an additional outward drift.

Since this radial velocity was found to be proportional to the cell-density gradient (Figs. 2C, 5C,

and S7), we added a drift velocity of the form:

257                 $$\vec{U} = -\eta \cdot \vec{\nabla}(\rho/\rho_0) \qquad\qquad (3)$$

with $\eta = 10^{-7}$ cm^2s^{-1} and ρ_0 the initial (uniform) cell density ($\rho_0 \sim 0.7\cdot10^6$ cells·mm^2). Indeed,

adding this fluid-induced velocity component to the model can strictly limit the condensation and

account for the broadening and shape of mature condensates during phase 2 (Fig. 6A-B, black

lines). Notably, adding the drift velocity from the beginning of the simulation resulted in a

gradual transition between phase 1 and phase 2, which does not agree with the sharp transition

observed in the experiments (Figs. 1C and 6B). However, adding this velocity only above a

certain threshold cell density clearly accounts for the observed transition (Fig. 6B). This is

consistent with the emergence of the correlated motion only above a certain cell density (Fig. 1).

The added drift velocity can also account for the expulsion of non-chemotactic cells from the

condensate (Fig. 6C, see also Fig. 3 and Supplementary Information).

Finally, we also considered the potential influence of correlated swimming on the

chemotaxis efficiency (30, 39). First, we note that for such influence to limit condensation, it

must reduce the chemotaxis efficiency. To incorporate this into the model, we allowed the

chemotactic coefficient (β) to depend on the local cell-density according to the findings in ref. 30

(Fig. S8). Clearly, such an effect can also contribute to limiting the condensation (Fig. 5B,

orange line). However, modulating the chemotaxis efficiency cannot lead to repulsion of non-

chemotactic cells.

The overall relationship between the initial cell density and condensation is summarized

in Fig. 6D. When the initial cell density is below approximately $0.4 \cdot 10^6$ cells/mm^2, no

condensation is observed (14, 37). At higher densities, the initially uniform distribution becomes

unstable, and condensation emerges. In the absence of corelated swimming, the bacteria can

condense until the local cell density becomes sufficient to raise the local pH beyond the sensory

dynamic range of the cells (blue line, see also ref. 14). However, because correlated swimming

readily emerges, bacterial condensation is severely restricted (simulations: dark-gray line; data:

red circles). In contrast, in porous media, condensation proceeds further (gray circles), though

still limited below the value set by the dynamic range (blue line), likely due to the non-local

nature of bacterial gradient sensing (18). At even higher cell densities, condensation becomes

limited by ring formation, likely due to aerotaxis (Fig. 5).

Discussion

Chemotaxis-mediated condensation and correlated swimming are two prominent collective

behaviors exhibited by motile bacteria. Here, we describe a fundamental interplay between

these behaviors. We show that bacterial condensation is fundamentally constrained by the

emergence of correlated swimming (Figs. 1 and 6). In liquid, bacterial condensation proceeds in

two dynamic phases: initially, the bacterial population condenses and the local cell density

increases; then, upon the onset of correlated swimming, the width of the condensate plateaus

while cell density continues to rise. These bacterial condensates —characterized by steep cell-

density gradients and lack of physical boundaries— give rise to a distinct form of active

turbulence, with a radial component aligned down the cell-density gradient (Figs. 1 and 5).

Furthermore, we find that correlated swimming is suppressed in porous environments, enabling

exceptionally robust bacterial condensation with extremely high cell density (Fig. 4).

By imposing active turbulence, correlated swimming not only constrains bacterial

condensation but also expels non-chemotactic bacteria from the condensate, effectively leading

to segregation of the initially mixed population into chemotactic and non-chemotactic

subcommunities (Fig. 3). Previous experiments demonstrated bacterial expulsion from a local

vortex generated by rotating magnetic microspheres at 2–40 Hz (27). Here, however, the

condensates exhibit much weaker vorticity ($\sim 10^{-3}$ Hz). Moreover, the expulsion effect is

substantially reduced in a porous environment, where the local cell density in the condensate is

approximately 10-fold higher, but correlated swimming is absent (Fig. 4). Taken together, these

observations suggest that the expulsion of non-chemotactic cells is not merely due to the

imposed vorticity or the accumulation of the chemotactic cells. Instead, we suggest that this

expulsion is driven by the outward fluid flow generated by correlated swimming, which imposes

an outward velocity bias (Eq. 3 and Fig. 6). Notably, this radial velocity bias, generically aligns

with the bacterial density gradient, even in ring-shaped condensates (Figs. 1,2, 5 and S7),

consistent with the observation that correlated swimming is enhanced at high cell densities (28)

but fades as cells move down the density gradient.

Correlated bacterial swimming and active turbulence can influence condensation through

two additional mechanisms. First, by promoting advection of chemotactic effectors (in this case,

protons), which can limit the buildup of steep gradients. We find that over a time period

corresponding to the diffusion time across the condensate, each probe particle typically

experiences at least one significant event that enhances its motion beyond simple diffusion, i.e.,

with $Pe > 0.5$ (Fig. 2B). This suggests that while advection may not dominate over diffusion, it

can be significant. Second, correlated swimming may directly affect chemotaxis efficiency by

altering run-and-tumble dynamics. This effect, however, been shown to enhance bacterial

accumulation near local attractive sources (31) but has also been shown to either improve or

impair chemotaxis efficiency in shallow gradients, depending on cell density (30). Although this

effect could potentially contribute to the observed broadening of the condensation (Fig. 6), it

remains unclear to what extent such modulation of chemotaxis efficacy is effective under sharp

gradients of both the effector distribution and cell-density. In any case, these effects cannot

account for the observed expulsion of non-chemotactic cells.

Bacterial suspensions confined in small cylindrical wells (~ 45 µm diameter) have been

shown to exhibit vortex-like motion with a random direction, whereas in larger wells, chaotic

motion develops without coherent vorticity (40). In contrast, the correlated swimming observed

here is consistently characterized by clockwise vorticity of the entire condensate, (~ 600 µm

diameter), as viewed from the glass side (Figs. 1–2). At higher cell densities, when the

condensate develops a ring-like structure the vorticity reverses between the inner and outer

regions of the ring (Fig. 5C). Because the radial velocity also reverses across the ring, a global

rightward swimming tendency is maintained (Fig. 5C), consistent with the circular motion

observed when non-interacting individual cells swim near glass surfaces (< 1 µM; Fig. S9) (41).

However, bacterial condensates extend up to 60 µm from the surface and are far denser. We

suggest that interactions with the nearby surface, while unable to generate strong vorticity on

their own, may nevertheless break the symmetry and promote clockwise bias to otherwise

chaotic motion.

The distinct swimming behaviors observed here may also be related to the unique

properties of the bacterial condensate. In general, but with notable exceptions (28, 42), active

turbulence has been studied in uniform cell populations confined by well-defined lateral

boundaries, either solid (43) or at liquid–air or liquid–oil interfaces (43-46). In such settings, the

bacterial behavior near the lateral physical boundaries may play an important role in shaping the

observed swimming patterns (44, 47). In contrast, the bacterial condensate examined here

represents a unique system: a self-sustained yet highly non-uniform cell suspension lacking

lateral physical boundaries. The fluid motion in the vicinity of the condensate exhibits a three-

dimensional toroidal pattern (Fig. 2F), reminiscent of the flow patterns observed in bioconvection

(34, 48). However, in bioconvection it arises due to instability caused by upward bacterial taxis

opposing the downward gravitational force. More complex behavior has been observed when

metallic particles were also driven up and down by the applied voltage of the sample (49). In

contrast, the bacterial condensate is overall stable, and the fluid motion arises directly from local

bacterial motion combined with the strong cell density gradients imposed by the inward

chemotaxis force.

Correlated bacterial swimming was clearly suppressed in porous environments, leading

to exceptionally dense cell condensates (Fig. 4). Notably, experiments conducted in mucin-

containing media at neutral pH still exhibit correlated swimming, albeit with larger vortex size

(50). Interestingly, however, the agarose-based hydrogel used here more closely resembles

mucin-containing media at low pH (51), which may be more relevant to certain physiological

conditions than neutral pH. Moreover, the enhanced condensation observed under these

conditions may facilitate the formation of more structured biofilm communities.

Materials and Methods

Strains and plasmids. We used *E. coli* MG1655 (VF6 (32, 52)) and its derivatives harboring

deletion of *cheY* and *cheZ* (MK3) or deletions of all chemoreceptors and a dysfunctional *cheZ*

gene (non-chemotactic run-and-tumble cells (AVE4 (32)). Cells expressed also a fluorescent

mGFP (75 µM IPTG) or mCherry (5 mM) from a pTrc99a vector (pSA11).

Cell growth and motility medium. Cells were grown over night in Tryptone Broth (TB, tryptone

extract 10 g/l and 5 g/l NaCl) supplemented with Ampicillin, diluted 100-fold into fresh medium

with IPTG, and allowed to grow at 33.5 °C with shaking. Cells were harvested at an OD_{600} of 0.5

by centrifugation and gently washed twice in motility medium and resuspended in motility

medium with final cell concentration of OD_{600} 0.4 (unless otherwise mentioned). The motility

medium (0.1 mM potassium phosphate, 0.94 mM lactic acid, 85.5 mM sodium chloride, 0.1 mM

EDTA, and 1 µM methionine) titrated to the desired pH with NaOH. Notably, this medium

supports long-term bacterial motility but not cell proliferation. Experiments in porous medium

(Fig. 4) were done gently mixing the cell suspension with agarose hydrogel at 37 °C to a final

agarose concentration of 0.08%.

The condensation setup. The setup used here is schematically illustrated in Fig. 1a. For more

details see ref. 14. An agarose gel (3% in motility medium) was casted in a cylindrical chamber

made from grade-2 titanium (6 mm height, 10.7mm diameter) and sealed at the top with a mold

with a thin cylindrical protrusion (~120 μm high, and radius of ~ 3 or 5 mm), which will form the

bacterial layer; fabricated using a resin printer (Elegoo Mars 3). After the agarose solidified, the

mold and the bottom cover were removed and the agarose was dialyzed overnight in motility

buffer and a second time, for an additional 4 h, in a fresh motility buffer just before the

experiment (final dialysis ratio of 10^4:1). Bacterial cells were then introduced into the shallow

well, suspended in either liquid or soft agarose (0.08%), and the bottom and top of the chamber

were covered with BSA-treated glass coverslips (1% bovine serum albumin in double-distilled

water for 30 minutes) and secured in position. Experiments were performed at room

temperature (24 °C).

Fluorescence microscopy. The bacterial distribution was followed by using a Nikon Ti

microscope equipped with a 20× (0.5 NA) or 4× (0.2 NA) objective, a decoded controlled stage

and a 'perfect-focus' system. Large images were obtained by stitching smaller images

(approximately 2mm× 2mm each) with 20% overlap using a built-in function in NIS Elements

software (Nikon). The bacterial distribution inside the well was extracted from the images by

removing the measured background intensity and normalizing to the average intensity within

each well. Radial averaging was done by measuring the fluorescence intensity around the

center of focal points (Radial Profile Angle plugin for ImageJ).

Cell density estimates. Due to inhomogeneity of the cell distribution in the vertical direction

(see Fig. 2F), cell distributions are shown in cells/mm^2. Calibration of OD_{600} to cells/mm^3 was

done by plating serial dilution of cells after growth (OD_{600} 0.5), yielding OD_{600} 0.5 $\approx$ 7.2·10^9

cfu/mm^3. Surface density was then estimated by multiplying the cell density (ρ_{3D}), averaged

over the vertical direction (z), by the well height (h): $\rho = h\langle\rho_{3D}\rangle_z$.

Non-uniformity estimates. To determine the non-uniformity of the cell distribution within

condensates from still images, where temporal averaging is unavailable, a smooth radially-

averaged version of each image was subtracted from the original ones. A high-pass filter was

then applied to remove residual, large-scale intensity fluctuations (0-140 µm). Non-uniformity

was then measured as the difference between the standard deviation of the intensity within a

disk around the condensate center (600 µm diameter) and away from it.

Fluorescent spheres. In the experiments presented in Fig. 2, we used green fluorescent

spheres (beads) with a 1µm diameter (FluoSpheres, ThermoFisher Scientific). Before use,

spheres were thoroughly washed in motility medium to which Tween-20 (0.01%) was added.

The spheres were then added to a cell suspension, which also contained Tween-20 (0.01%).

The added Tween-20 prevented self-aggregation of the fluorescent spheres but had no

detectable effect on the bacterial condensation.

Particle image velocimetry (PIV). To capture velocity fields, time-lapse fluorescent images (2-

10 frames per second) were captured and analyzed by PIV using PIVLab (53). To analyze the

collective motion of cells, a filtering procedure was first applied to remove the large-scale

bacterial condensation (see Non-uniformity estimates) and a nearest-neighbors smoothing

scheme was applied to remove residual noise. PIV analysis proceeded using multipass FTT

window deformation (4 passes) with the final interrogation window being 16-18 pixels (64–

72 µm). To avoid false detection, only vectors with a correlation coefficient larger than 0.5 were

kept. The PIV parameters were chosen to balance between the resolution of the extracted

velocity field and the signal-to-noise ratio; changing them did not modify the results significantly.

To analyze the motion of the fluorescent beads, an initial mask was applied to ensure only

particles within the focal plane were analyzed. This was possible since beads within the focal

plane seem smaller and brighter than beads in other planes (see SI Video 3). Despite the BSA

treatment, beads slowly adhered to the glass coverslip. By subtracting the median image from

the time lapse, only non-adhered beads were analyzed. Analysis then proceeded in PIVLab as

described for the bacteria.

Particle tracking and detection of jets. Time lapse images (2 frames per second) of the

fluorescent beads were masked to ensure only particles within the focal plane are analyzed (see

Particle image velocimetry (PIV)). Tracking was then performed using the TrackMate plugin for

ImageJ. Detected tracks were exported and analyzed separately. Each track was smoothed

using a Savgol-Golay filter (5 frames window) and velocities were estimated numerically. Jets

were detected as temporal peaks in the velocity magnitude with a mean prominence of more

than 0.5 µm/s. The length of each track (L) and it's mean velocity (v) were captured. If

neighboring beads, less than L apart, exhibited a jet in the same direction ($\Delta\theta < 0.15$), their jets

were considered as correlated; the velocity of such jets were averaged over and considered as

one. Due to the low density of beads, such events were rare (7 of the 2362 detected jets).

Numerical solution of the model. Solutions and parameters were based on (14) and full

details can be found in the Supp. Text. Briefly, the dynamics of cells was modeled in 2D

assuming radial symmetry and the dynamics of protons was modeled in 3D assuming cylindrical

symmetry. Proton uptake in the bacterial layer was accounted for by imposing absorbing

boundary conditions for the protons at the bacterial layer interface. Numerical solutions were

obtained by calculating spatial derivatives with 2nd order accuracy and using a 4th order Runge-

Kutta method for time integration.

Author Contributions

NL and AV designed the experiments; NL performed the experiments and theoretical calculations; NL and

AV analyzed the data and wrote the manuscript.

Acknowledgments

We thank Alexander Feigel (Hebrew University) and Igor Aronson (Penn. State University) for

help full conversations. We thank the Milner Foundation for partly funding NL's fellowship. This

work was supported by the Israeli Foundation of Sciences and Humanities and by a research

grant from the U.S.-Israel Binational Science Foundation.

Competing Interest Statement: The authors declare no competing interests.

Data Availability Statement: All data is available upon request.

References

1. Berg HC (2004) *E. coli in Motion* (Springer, NY).
2. Parkinson JS, Hazelbauer GL, & Falke JJ (2015) Signaling and sensory adaptation in *Escherichia coli* chemoreceptors: 2015 update. *Trends Microbiol.* 23(5):257-266.
3. Colin R & Sourjik V (2017) Emergent properties of bacterial chemotaxis pathway. *Curr. Opin. Microbiol.* 39:24-33.
4. Moore JP, Kamino K, & Emonet T (2021) Non-genetic diversity in chemosensing and chemotactic behavior. *Int. J. Mol. Sci.* 22(13):6960.
5. Keegstra JM, Carrara F, & Stocker R (2022) The ecological roles of bacterial chemotaxis. *Nat. Rev. Microbiol.*
6. Raina J-B*, et al.* (2022) Chemotaxis shapes the microscale organization of the ocean's microbiome. *Nature* 605(7908):132-138.
7. Matilla MA & Krell T (2018) The effect of bacterial chemotaxis on host infection and pathogenicity. *FEMS Microbiol Rev* 42(1).
8. Butler SM & Camilli A (2005) Going against the grain: chemotaxis and infection in *Vibrio cholerae*. *Nat. Rev. Microbiol.* 3(8):611-620.
9. Zhou B, Szymanski CM, & Baylink A (2023) Bacterial chemotaxis in human diseases. *Trends Microbiol.* 31(5):453-467.
10. Cremer J*, et al.* (2019) Chemotaxis as a navigation strategy to boost range expansion. *Nature* 575(7784):658-663.
11. Adler J (1966) Chemotaxis in bacteria. *Science* 153:708-716.
12. Budrene EO & Berg HC (1991) Complex patterns formed by motile cells of *Escherichia coli*. *Nature* 349(6310):630-633.
13. Park S*, et al.* (2003) Motion to Form a Quorum. *Science* 301(5630):188-188.
14. Livne N, Koler M, & Vaknin A (2024) Collective condensation and auto-aggregation of *Escherichia coli* in uniform acidic environments. *Comm. Biol.* 7(1):1028.
15. Brumley DR*, et al.* (2019) Bacteria push the limits of chemotactic precision to navigate dynamic chemical gradients. *Proc. Natl. Acad. Sci.* 116(22):10792-10797.
16. Fu X*, et al.* (2018) Spatial self-organization resolves conflicts between individuality and collective migration. *Nat. Comm.* 9(1):2177.
17. Childress S & Percus JK (1981) Nonlinear aspects of chemotaxis. *Mathematical Biosciences* 56(3):217-237.
18. Mittal N, Budrene EO, Brenner MP, & Van Oudenaarden A (2003) Motility of *Escherichia coli* cells in clusters formed by chemotactic aggregation. *Proc. Natl. Acad. Sci.* 100(23):13259-13263.
19. Berg HC & Brown DA (1972) Chemotaxis in *Escherichia coli* analysed by three-dimensional tracking. *Nature* 239(5374):500-504.
20. Yang Y & Sourjik V (2012) Opposite responses by different chemoreceptors set a tunable preference point in Escherichia coli pH taxis. *Mol. Microbiol.* 86(6):1482-1489.
21. Livne N & Vaknin A (2022) Collective responses of bacteria to a local source of conflicting effectors. *Sci. Rep.* 12(1):4928.
22. Toner J & Tu Y (1998) Flocks, herds, and schools: A quantitative theory of flocking. *Phys. Rev. E* 58(4):4828-4858.

23. Wensink HH*, et al.* (2012) Meso-scale turbulence in living fluids. *Proc. Natl. Acad. Sci.*
109(36):14308-14313.
24. Koch DL & Subramanian G (2011) Collective hydrodynamics of swimming
microorganisms: Living fluids. *Annl. Rev. Fluid Mechanics* 43(Volume 43, 2011):637-
659.
25. Aranson IS (2022) Bacterial active matter. *Rep. Prog. Phys.* 85(7).
26. Alert R, Casademunt J, & Joanny J-F (2022) Active turbulence. *Annl. Rev. Cond. Matt.*
Phys. 13(Volume 13, 2022):143-170.
27. Sokolov A & Aranson IS (2016) Rapid expulsion of microswimmers by a vortical flow.
Nat. Commun. 7(1):11114.
28. Sokolov A, Aranson IS, Kessler JO, & Goldstein RE (2007) Concentration dependence of
the collective dynamics of swimming bacteria. *Phys. Rev. Lett.* 98(15):158102.
29. Lushi E, Goldstein RE, & Shelley MJ (2012) Collective chemotactic dynamics in the
presence of self-generated fluid flows. *Phys. Rev. E* 86(4):040902.
30. Colin R, Drescher K, & Sourjik V (2019) Chemotactic behaviour of *Escherichia coli* at
high cell density. *Nat. Commun.* 10(1):5329.
31. Kai T*, et al.* (2024) Collective gradient sensing by dilute swimming bacteria without
clustering. *Phys. Rev. Res.* 6(3):L032061.
32. Tamar E, Koler M, & Vaknin A (2016) The role of motility and chemotaxis in the bacterial
colonization of protected surfaces. *Sci. Rep.* 6:19616.
33. Brinkman HC (1949) A calculation of the viscous force exerted by a flowing fluid on a
dense swarm of particles. *Flow, Turbulence and Combustion* 1(1):27-34.
34. Bees MA (2020) Advances in bioconvection. *Annl. Rev. Fluid Mech.* 52(Volume 52,
2020):449-476.
35. Taylor BL, Zhulin IB, & Johnson MS (1999) Aerotaxis and other energy-sensing behavior
in bacteria. *Annu. Rev. Microbiol.* 53:103-128.
36. Keller EF & Segel LA (1971) Traveling bands of chemotactic bacteria: a theoretical
analysis. *J. Theor. Biol.* 30(2):235-248.
37. Livne N, Vaknin A, & Agam O (2025) Pattern formation in *E. coli* through negative
chemotaxis: Instability, condensation, and merging. *Phys. Rev. Res.* 7(2):023095.
38. Si G, Wu T, Ouyang Q, & Tu Y (2012) Pathway-based mean-field model for *Escherichia*
coli chemotaxis. *Phys. Rev. Lett.* 109(4):048101.
39. Miller M & Toner J (2024) Following your nose: Autochemotaxis and other mechanisms
for spinodal decomposition in flocks. *Phys. Rev. Lett.* 132(12):128301.
40. Nishiguchi D, Shiratani S, Takeuchi KA, & Aranson IS (2025) Vortex reversal is a
precursor of confined bacterial turbulence. *Proc. Natl. Acad. Sci. USA*
122(11):e2414446122.
41. Lauga E, DiLuzio WR, Whitesides GM, & Stone HA (2006) Swimming in circles: motion
of bacteria near solid boundaries. *Biophys J* 90(2):400-412.
42. Sokolov A, Rubio LD, Brady JF, & Aranson IS (2018) Instability of expanding bacterial
droplets. *Nat. Commun.* 9(1):1322.
43. Nishiguchi D, Aranson IS, Snezhko A, & Sokolov A (2018) Engineering bacterial vortex
lattice via direct laser lithography. *Nat. Commun.* 9(1):4486.

44. Wioland H, Woodhouse FG, Dunkel J, Kessler JO, & Goldstein RE (2013) Confinement
stabilizes a bacterial suspension into a spiral vortex. *Phys. Rev. Lett.* 110(26):268102.
45. Vincenti B*, et al*. (2019) Magnetotactic bacteria in a droplet self-assemble into a rotary
motor. *Nat. Commun.* 10(1):5082.
46. Nishiguchi D, Shiratani S, Takeuchi KA, & Aranson IS (2025) Vortex reversal is a
precursor of confined bacterial turbulence. *Proc Natl Acad Sci U S A*
122(11):e2414446122.
47. Lushi E, Wioland H, & Goldstein RE (2014) Fluid flows created by swimming bacteria
drive self-organization in confined suspensions. *Proc. Natl. Acad. Sci. (USA)*
111(27):9733-9738.
48. Abe T, Nakamura S, & Kudo S (2017) Bioconvection induced by bacterial chemotaxis in
a capillary assay. *Biochem. Biophys. Res. Comm.* 483(1):277-282.
49. Sapozhnikov MV, Tolmachev YV, Aranson IS, & Kwok WK (2003) Dynamic self-assembly
and patterns in electrostatically driven granular media. *Phys. Rev. Let.* 90(11):114301.
50. Liao W & Aranson IS (2023) Viscoelasticity enhances collective motion of bacteria.
PNAS Nexus 2(9).
51. Celli J*, et al*. (2005) Viscoelastic properties and dynamics of porcine gastric mucin.
Biomacromolecules 6(3):1329-1333.
52. Frank V, Piñas GE, Cohen H, Parkinson JS, & Vaknin A (2016) Networked
chemoreceptors benefit bacterial chemotaxis performance. *mBio* 7(6).
53. Thielicke W & Sonntag R (2021) Particle Image Velocimetry for MATLAB: Accuracy and
enhanced algorithms in PIVlab. *J. Open Res. Soft.*

Figure 1. Bacterial condensation is constrained by correlated swimming. (A) Schematic of bacterial condensation in acidic environments (see ref. 14). Bacteria neutralize their vicinity and generate pH gradients in an otherwise uniform acid medium. At the same time, cells are chemotactically attracted toward neutral pH. These behaviors form reinforcing feedback that derives condensation. **(B)** The experimental setup consists of a thin GFP-expressing cell suspension (5 mm diameter) confined in a circular well made by 3% agarose gel. Both the cell suspension and the surrounding gel initially form a uniform acidic environment (pH 5.2). Condensation was monitored using fluorescence microscopy. Shown on the right is a typical mature condensate 2 h after initiation (scale bar: 300 μm), along with the radially averaged cell-density profile. The peak cell density (ρ_{max}) and the width at half maximum (w) are marked. **(C)** A typical condensation dynamic, characterized by increasing ρ_{max} and decreasing w until w abruptly saturates at time t_{sat}. **(D)** The condensate width (w) is plotted against peak cell density (ρ_{max}) for 7 independent experiments. The solid gray line reparents a power-law slope of -2/3. The saturation value of w (dark-gray dashed line) and the corresponding value measured at low cell density (light-gray dashed line, ref. 18) are indicated. *Inset* – normalized cell density profiles observed at different time points for 3 independent experiments: profiles taken at $t < t_{sat}$ (phase 1) are labeled green and profiles taken at $t > t_{sat}$ (phase 2) are labeled red. Each profile was normalized by subtracting the background intensity and dividing by the peak intensity; the radial coordinate was normalized by $w/2$. **(E)** *Upper-panel* – A typical filtered fluorescence image of the condensate (left, see Materials and Methods and SI Video 1; scale bar: 300 μm) and the corresponding time-averaged (30 s) velocity field obtained by particle image velocimetry (right, grey arrows are a guide to the eye). These are demonstrating the dynamic non-uniformity developed at high cell density. *Lower panel* – The non-uniformity within condensates (Δ), evaluated within the area r ≤ 300 μm, is plotted over time for 4 independent experiments (see Materials and Methods). *Inset* – The relative non-uniformity ($\Delta /\sqrt{I_{peak}}$) plotted over time.

Figure 2. Fluid motion within the condensate. Fluid motion was assessed by tracking 1 μm beads (green fluorescent) mixed in the bacterial suspension. In this case, cells were labeled with mCherry (red). (**A**) Traces of individual beads (over 50 sec) within a mature condensate, using particle tracking analysis (Materials and Methods). The dashed white circle indicates the contour of the condensate width ($w \sim 550$ μm). A typical trace is highlighted on the right, illustrating intermittent motion composed of periods of local random motion (gray symbols) and periods of directed motion, termed 'jets' (green symbols). The length (L) and average speed (v) of each jet were used to calculate $Pe = v \cdot L/D_{beads}$ (with $D_{beads} = 5 \cdot 10^{-9}$ cm^2 s^{-1}). The cumulative distribution of Pe is shown on the right for beads inside or outside the area defined by the width of the condensate (dashed line). (**B**) Comparison of the beads motion (using 90 sec traces) with the expected proton diffusion. First, the number of jets per bead with a Pe value larger than a given threshold (Pe_0) were found and then multiplied by the typical diffusion time of protons across the condensate $t_D \sim w^2/4D_{H+} \sim 4$min (using $D_{H+} = 5 \cdot 10^{-6}$ cm^2 s^{-1}). (**C**) Time-averaged velocity field (averaged over 90 sec) in the condensate plane, extracted using PIV (Materials and Methods). Profiles of the radial and tangential velocity components are shown on the right alongside the corresponding (normalized) cell density profile. The red arrow marks the condensate width (w). (**D**) The angular velocity profile, revealing correlated vortex-like motion. Clockwise direction was chosen as positive. (**E**) Measured velocity field above the condensate (h $\sim$ 100 μm) showing a radially-inward flow. (**F**) Schematic summarizing the inferred fluid motion in the vicinity of the condensate (see also Supplementary Video 2).

Figure 3. Non-chemotactic cells are forced out of the condensate. Chemotactic cells (mCherry labeled) and non-chemotactic cells ($\Delta cheY/Z$; GFP labeled) were initially mixed in a 100:1 ratio, respectively, and applied to the setup shown in Fig. 1B. The cell-density profiles of the non-chemotactic cells in a mature condensate are shown (green symbols; 3 independent experiments), along with the corresponding profile of the condensate (red dashed line). Profiles were normalized to the background density and divided by a factor of 3000 (red) and 1 (green).

Figure 4. Bacterial condensation in porous media. Experiments similar to those described thus far were conducted, except that the cell suspension was mixed with low concentration of agarose (0.08% w/v), resulting in cells being embedded within a dilute hydrogel. General chemotactic behavior in this hydrogel medium is shown in Fig. S3. (**A**) Fluorescence image of a typical mature condensates condensate in porous medium (scale bar: 300 μm) along with the corresponding radially-averaged density profile. For comparison, a typical profile of a condensate observed in liquid is also shown (blue). A schematic description of the revised setup is also shown on the right. Because condensations in porous medium were significantly narrower, the diameter of the cell suspension was reduced to 3 mm to avoid the formation of multiple focal points (see ref. 14 and Materials and Methods). (**B**) The condensation dynamics in porous medium (left plot, black symbols). The saturation value of the condensate width in liquid is indicated by a dashed blue line, and the corresponding value at low cell-density (from ref. 18) is indicated by the gray dashed lines. *Inset* – The saturation width of the condensate in porous medium is plotted against the peak cell density for 17 experiments. Shaded gray region represents a linear fit to the data. (**C**) Fluid motion in porous medium. The 2D velocity field within the plane of the condensate was measured and analyzed as in Fig. 2. The time averaged velocity field (30 s) is shown (right) along with the radial and tangential velocity components plotted as a function of the distance from the center of the condensate (left). (**D**) Expulsion of non-chemotactic cells from the condensate was measured in porous media in experiments similar to those described in Fig. 3. Expulsion efficiency is defined as the ratio of the fluorescence intensity at the center of the condensation to that at the periphery.

Figure 5. Condensation at even higher initial cell density. (**A**) The peak cell density is plotted against the total number of cells in the condensate for experiments performed with the cells in liquid (red symbols) or in porous medium (blue symbols). The total number of cells was modulated by the varying the initial (uniform) cell density. The decrease in the peak cell density observed at high cell numbers was correlated with the formation of ring-shaped condensation, demonstrated in (B) and (C). (**B**) *Porous medium*: A representative of a mature ring-shaped condensate. Evolution of the condensate profile is shown (left plot), along with a florescence image taken at 50 min (scale bar: 150 µm). (**C**) *Liquid medium*: A typical fluorescence image of a mature condensate at high cell number. Time-averaged bacterial velocity field, extracted using PIV, are overlaid on the image (red arrows), and also shown separately on the right (green arrows). Below the image, the radial and tangential velocity components are plotted as function of the distance from the condensate center, along with the corresponding cell-density gradient. A schematic shown on the right illustrates the correlated reversal of both velocity components across the ring, consistently producing a rightward swimming tendency in both the inner and outer regions.

Figure 6. A model for condensate broadening by collective motion. Numerical solutions for the model described by Eqs. 1–3 were used to compare its predictions with the data obtained in liquid medium. Chemotaxis is described by Eqs. 1 and 2, and an additional outward velocity drift, $\vec{U} \sim -\eta \cdot \vec{\nabla}\rho$ (Eq. 3) is assumed be imposed by the active turbulence. The parameter η is given hereafter in units of 10^{-7} cm^2 s^{-1}. (**A**) *Condensation shape*. Measured condensate profiles from phase 1 (three independent experiments; see also Fig. 1) are well captured by the model with $\eta = 0$ (right plot; blue dashed line). However, a value of $\eta = 1.7$ was required to match the data from phase 2 (left plot, black line). We also tested a model with $\eta = 0$, but in which the chemotaxis efficiency (Eq. 1) depends on the local cell-density as described in ref. 30 (orange line; Fig. S8). (**B**) *Condensation dynamics*. The condensate width is plotted against time (gray symbols; 4 independent experiments, see Fig. 1). To capture the onset of active turbulence at a critical peak density ρ_c, we modeled allowed η to depend on the peak cell density ρ_c: $\eta \sim 1/(1 + \exp(-(k/\rho_0)(\rho_{max} - \rho_c)))$, with ρ_0 being the initial (uniform) density, $k = 3$ and $\rho_c = 25 \cdot \rho_0$. (**C**) *Expulsion of non-chemotactic cells*. Data from Fig. 3 (green symbols) is replotted together with the corresponding distribution predicted by the model $\rho = \exp\{-(\eta/D_{NC}) \cdot (\rho_{WT}(r)/\rho_0)\}$ with $\eta = 1.7$, $\rho_{WT}(r)$ taken from the experimental data in (A), and $D_{NC} = 2.5 \cdot 10^{-6}$ cm^{2}s^{-1} representing the effective diffusion of the non-chemotactic cells (see derivation in Supp. text). (**D**) *Summary plot*. The saturated condensation level (defined as inverse width: $1/w$) is plotted as a function of the initial (uniform) cell density. Data from experiments in liquid and porous media (empty circles) and the corresponding simulations (lines) are shown (as labeled). The vertical black line ('aerotaxis') indicates the density above which ring-shaped condensates appear. The low-density limit (18) is plotted as a dashed gray line ('non-locality'). The shaded gray region on the left ('no condensation') marks a range of concentrations that do not support condensation (14, 37).