Exosome Trafficking Is a Key Regulator of Adipocyte Thermogenesis

Devesh Kesharwani[1], Michele Karolak[1], Chad Doucette[1], Rachel Mendola[1], Summer Pray[1], Raghu Bhardwaj[1], Edward Jachimowicz[1], Su Su[1], Anne Harrington[1], Victoria DeMambro[1], Clifford Rosen[1,2,3], Lucy Liaw[1,2,3], Aaron C. Brown[1,2,3*]

1- Center for Molecular Medicine, MaineHealth Institute for Research, 81 Research Drive, Scarborough, ME 04074
2- Graduate School of Biomedical Science and Engineering, The University of Maine, Orono, Maine 04469
3- Tufts University School of Medicine, 145 Harrison Ave, Boston, MA 02111

*Corresponding author. Aaron C. Brown. Email: aaron.c.brown@mainehealth.org

Author Contributions: A.C.B., L.L., and C.R. designed research; D.K., M.K., C.D., R.M., S.P., R.B., E.J., S.S., and V.M. performed research; M.K., A.H., and L.L. contributed new reagents/analytic tools; D.K., M.K., C.D., S.S., V.D., and A.C.B. analyzed data; A.C.B and D.K. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Classification: Biological Sciences, Physiology.

Keywords: beige adipocytes, thermogenesis, exosomes, Rab27a, metabolic disease.

This PDF file includes:

   Main Text

   Figures 1 to 4
   Supplemental Figures 1 to 3
   Tables 1 and 2

Abstract
Activation of beige adipocytes enhances energy expenditure and promotes metabolic health, presenting a promising approach for combating obesity and diabetes. As part of this process, thermogenesis, fueled by uncoupled mitochondrial respiration, plays a central role in converting calories into thermal energy, thereby preventing their storage as fat. Here, we identify a role for exosome trafficking as an intrinsic regulator of beige adipocyte thermogenesis. Exosomes are small extracellular vesicles that mediate cell-cell and intracellular communication by transporting regulatory cargo, including microRNAs, proteins, and lipids. Using both human cells and mouse models, we show that thermogenic activation of beige adipocytes promotes the rapid release of exosomes enriched in microRNAs known to suppress thermogenic programs. Genetic or pharmacological blockade of exosome secretion attenuates thermogenesis, whereas enhancing exosome release amplifies thermogenic output. Mice deficient in the exosome secretion regulator *Rab27a* exhibit reduced energy expenditure in response to both cold exposure and β3-adrenergic stimulation. These findings establish exosome trafficking as a key contributor to beige adipocyte thermogenic capacity, highlighting an intracellular mechanism that may be leveraged to enhance energy expenditure and treat obesity-related metabolic diseases.

Significance statement
Thermogenic adipocytes, including beige fat cells, help maintain energy balance by converting excess nutrients into heat, thereby reducing fat storage and supporting metabolic function. Although these cells are known to promote energy expenditure, the intracellular processes that enable their full thermogenic response are not well defined. Here, we show that exosome secretion is required for beige adipocytes to reach their full thermogenic potential. Blocking exosome release dampens this response, while boosting it amplifies thermogenic output. These findings point to exosome release as an essential part of thermogenic regulation and a potential target for improving metabolic health.

Main Text

Introduction

Obesity, defined by excessive lipid accumulation in white adipose tissue, leads to chronic inflammation, insulin resistance, and dysregulated adipokine secretion, which together drive systemic metabolic dysfunction and increase the risk of type 2 diabetes, cardiovascular disease, and certain cancers[1]. Lifestyle interventions such as diet and exercise are first-line treatments for obesity, but long-term adherence is often poor and weight regain is common. Although bariatric surgery and GLP-1 receptor agonists are effective treatments for obesity, both are costly, can cause adverse effects, and are frequently associated with weight regain after initial success[2,3]. Given the limitations of current therapies, thermogenic adipocytes, including brown and beige fat cells, have emerged as compelling targets for obesity and diabetes treatment due to their ability to enhance energy expenditure and improve metabolic function through increased lipid and glucose utilization[4-7].

Brown and beige adipocytes play a crucial role in maintaining body temperature and supporting systemic metabolism by burning nutrients to produce heat upon activation. Brown adipocytes reside in dedicated depots such as the interscapular, cervical, and spinal regions and are developmentally programmed for stable thermogenic activity. In contrast, beige adipocytes are inducible thermogenic cells within subcutaneous white fat that acquire reversible thermogenic activity in response to changes in temperature[4,8]. Upon cold exposure, norepinephrine released from sympathetic nerves activates brown and beige adipocytes through α_1- and β_3-adrenergic signaling[9,10]. Both brown and beige adipocytes contain multilocular lipid droplets, are rich in mitochondria, and express uncoupling protein-1 (UCP1), which drives heat generation by uncoupling mitochondrial respiration[4]. In addition to UCP1-dependent thermogenesis, several UCP1-independent energy-dissipating mechanisms, including calcium flux, creatine turnover, and fatty acid cycling, consume ATP without driving cellular activity and release that energy as heat[11]. Recent studies indicate that beige adipocytes can be subdivided into distinct populations based on their reliance on these various pathways[12,13].

Upon cold exposure or β-adrenergic activation, brown and beige adipocytes enhance glucose uptake and fatty acid oxidation to fuel thermogenesis, a process that supports thermal homeostasis and also improves insulin sensitivity, facilitates triglyceride clearance, and contributes to systemic glucose regulation[4,6,9]. The discovery of beige fat and its inducible nature has sparked interest in its therapeutic potential, with efforts underway to pharmacologically recruit subcutaneous beige fat to combat obesity and metabolic disease. However, current approaches are limited: β-adrenergic receptor agonists can carry cardiovascular risks, while prolonged cold exposure is often impractical and may pose safety concerns for individuals with elevated cardiovascular risk[14]. These challenges underscore the need for alternative strategies to safely and effectively harness thermogenic adipocytes for metabolic benefit.

Thermogenic adipocytes also act as endocrine cells that influence distant organs and systemic metabolism by secreting adipokines and other bioactive factors, including lipids and peptides, which help regulate glucose homeostasis, lipid metabolism, and insulin sensitivity[15]. Among these secreted factors, exosomes, small extracellular vesicles that carry microRNAs, proteins, lipids, and RNAs, have emerged as important mediators of intercellular communication with systemic metabolic effects[16,17].

Adipose-derived exosomes can exert both beneficial and detrimental effects depending on physiological state and tissue origin. In obesity, white adipose tissue-derived exosomes are often enriched with proinflammatory cargo that promotes insulin resistance and metabolic dysfunction[17]. However, they may also serve adaptive functions. For example, exosomes from adipose tissue of obese mice deliver proteins to pancreatic β-cells, enhancing insulin secretion and glucose control, suggesting they signal insulin resistance in adipose tissue to boost insulin production when needed[18]. In contrast, exosomes from thermogenic depots support metabolic homeostasis. Brown adipose tissue-derived exosomal miR-132-3p, regulated by norepinephrine during cold exposure, inhibited hepatic lipogenesis by targeting *Srebf1*, thereby facilitating the body's adaptation to cold stress[19]. Similarly, cold exposure induced secretion of miR-99b-5p-enriched exosomes from brown fat that target the liver to regulate expression of fibroblast growth factor 21 (FGF21), a key metabolic hormone involved in energy expenditure and glucose homeostasis[20]. Circulating levels of exosomal miR-92a inversely correlated with brown fat activity in humans, suggesting its potential use as a non-invasive biomarker of thermogenic capacity[21]. Exosomes from adipose tissue macrophages of lean mice also improved glucose tolerance and insulin sensitivity when administered to obese recipients[22]. While these findings highlight the complex endocrine roles of adipose-derived exosomes, it remains unclear whether exosome secretion also plays a direct, cell-autonomous role in thermogenic adipocyte activation.

Here, we show that beige adipocyte activation is associated with increased exosome release containing suppressive cargo, leading us to hypothesize that exosome secretion is linked to thermogenesis. Supporting

110 this, inhibition of exosome secretion reduced UCP1 expression and thermogenic output, whereas enhanced
111 secretion boosted thermogenesis. Loss of the exosome trafficking protein *Rab27a* diminished exosome
112 release and impaired thermogenic capacity, resulting in reduced energy expenditure in response to cold or β_3-
113 adrenergic stimulation. These findings identify exosome secretion as a cell-intrinsic mechanism that facilitates
114 beige adipocyte thermogenesis and supports systemic energy balance.
115
116 **Results**
117
Exosome release during beige adipocyte activation is associated with thermogenic potential. In adult
mammals, the generation of beige adipocytes from adipose-resident precursor cells of stromal, perivascular,
and smooth muscle origin contributes modestly to overall beige adipocyte mass and occurs primarily after
prolonged cold exposure.[23]. In contrast, most beige adipocytes arise in response to acute cold exposure[24] via
rapid reprogramming of "whitened" beige adipocytes into a brown-like state (browning), which is defined by
acquisition of markers of thermogenic adipocytes, including *UCP1*. To examine this adaptive process in greater
detail, we employed a human iPSC-derived beige adipocyte model to investigate exosome release during
browning of mature adipocytes and to characterize their microRNA cargo[25]. iPSC-derived beige adipocyte
precursors were first differentiated into mature beige adipocytes over 12 days using a defined adipogenic
cocktail[25]. These mature beige adipocytes were then cultured in non-adipogenic basal medium for 4 days to
induce a "whitening" process, which resulted in a marked reduction in UCP1 expression[25]. To mimic cold-
induced browning, whitened cells were treated for 6 hours with forskolin, a β-adrenergic analog that activates
adenylyl cyclase to elevate intracellular cAMP. This short-term stimulation increased the expression of *UCP1*
and *PGC1α,* markers of mitochondrial uncoupling and biogenesis, respectively (Fig. 1A). Nanoparticle tracking
analysis revealed that forskolin activation triggered an >8-fold increase in exosome release into the culture
medium (Fig. 1B-C). Based on published literature, we compiled a panel of 98 microRNAs previously linked to
adipogenesis, metabolism, and obesity (Fig. S1A)[26-30]. We then developed targeted qPCR assays to quantify
these microRNAs in exosomes secreted by beige adipocytes under both whitened and forskolin-stimulated
conditions (Fig. 1D, $P < 0.02$; see Fig. S1A for the full data set).
Several microRNAs were significantly up- or downregulated in exosomes following beige adipocyte
browning, suggesting that thermogenic activation alters the selective packaging and secretion of microRNAs.
Among the most enriched was miR-99b-5p (>1200-fold), consistent with prior reports showing its secretion
from brown adipose tissue (BAT) following cold exposure and its role in targeting *Fgf21* in the liver (Fig. 1D)[20].
FGF21 is an adipokine secreted by both beige and brown adipocytes and plays a key role in promoting
browning and enhancing thermogenesis in white fat[31]. The most highly enriched microRNA in exosomes from
beige adipocytes undergoing browning was miR-27b-3p (~3500-fold). Its paralog miR-27a-3p, which shares an
identical seed sequence, was also significantly enriched (~46-fold), consistent with coordinated regulation
within the miR-27 family. Previous studies have shown that cellular levels of miR-27a and miR-27b are
downregulated during beige and brown adipocyte differentiation *in vitro*, as well as in subcutaneous and brown
adipose depots following acute cold exposure[32]. The miR-27 family targets and suppresses translation of
multiple transcription factors critical for UCP1-dependent thermogenesis, and *in vivo* knockdown of miR-27b
enhances browning and improves metabolic outcomes in obese mice[32,33]. Loss of miR-27, therefore, is
predicted to be required for the thermogenic adipocyte phenotype. Other microRNAs enriched in exosomes
from forskolin-activated beige adipocytes included known inhibitors of adipogenesis (miR-130b-3p, miR-454-
3p), thermogenesis (miR-133a-5p), and insulin signaling (miR-126-3p)[34-37]. Cellular levels of thermogenesis-
associated microRNAs (miR-99b-5p, miR-27a/b-3p, and miR-133a-5p) were found to decrease rapidly (within
1 hour) following forskolin stimulation, consistent with their selective export during browning (Fig. 1E). To test
whether this exosome secretion impacts thermogenesis, beige adipocytes were stimulated with forskolin in the
presence of the exosome secretion inhibitors manumycin A or GW4869[38,39]. Both inhibitors partially blocked
forskolin-induced *Ucp1* expression (Fig. 1F). Furthermore, oxygen consumption assays revealed that proton
leak-linked respiration was suppressed in the presence of manA or GW, indicating reduced mitochondrial
uncoupling and thermogenic activity (Fig. 1G). Together, these findings are consistent with a role for exosome
secretion in beige adipocyte thermogenesis, potentially by promoting the export of anti-thermogenic
microRNAs or other exosomal cargo that may influence thermogenic regulation.

Exosome-associated thermogenesis is conserved between humans and mice. Similar to human cells,
whitened beige adipocytes generated from the stromal vascular fraction of mouse inguinal adipose tissue
exhibited a substantial increase in exosome secretion following 6 hours of forskolin treatment, which was
reversed by co-treatment with the exosome secretion inhibitors manumycin A and GW4869 (Fig. 1H-I). qPCR

analysis of exosomal RNA revealed enrichment of miR-27a/b, miR-99b-5p, and miR-133a-5p in exosomes
from forskolin-activated beige adipocytes (Fig. 1J), accompanied by a corresponding decrease in their
intracellular levels (Fig. 1K). Inhibition of exosome secretion with manumycin A and GW4869 partially
attenuated the forskolin-induced upregulation of *Ucp1* transcript levels during beige adipocyte browning (Fig.
1L), and suppressed UCP1 protein levels in a dose-dependent manner (Fig. 1M). Comparable results were
observed when beige adipocytes were activated with cAMP instead of forskolin, indicating that the effects were
not due to off-target actions of this plant-derived adenylyl cyclase activator (Fig. S1B). Furthermore, exosome
secretion inhibition significantly reduced heat production during beige adipocyte browning, as measured by
high-resolution microcalorimetry (Fig. 1N-O). These results indicate that the role of exosome secretion in beige
adipocyte thermogenesis is conserved across species and that mouse models can provide valuable insight into
exosome-regulated browning processes relevant to human physiology.

Genetic enhancement of exosome secretion increases beige adipocyte thermogenesis. To determine
whether increasing exosome secretion enhances beige adipocyte thermogenesis, we generated transgenic
doxycycline-inducible mice expressing an exosome secretion booster cassette (Tg-ExoBooster; see schematic
in Fig. 2A). This cassette includes the genes STEAP3, SDC4, and NadB, which promote exosome biogenesis
and multivesicular body (MVB) formation, thereby enhancing exosome release[40]. Following 72 hours of
doxycycline treatment in both wild-type and Tg-ExoBooster inguinal SVF-derived beige adipocytes, we
confirmed transgene expression by RT-PCR (Fig. 2B) and observed a ~13-fold increase in exosome secretion
in the transgenic group (Fig. 2C–D). This increase was partially reversed by co-treatment with exosome
secretion inhibitors manumycin A and GW4869, suggesting pathway overlap (Fig. 2D). Beige adipocytes
expressing the ExoBooster cassette exhibited elevated *Ucp1* transcript levels and increased UCP1 protein
following forskolin stimulation compared to wild-type controls, at 4 (RNA) and 24 (protein) hours following
treatment, indicating a hypersensitized thermogenic response (Fig. 2E-F). Furthermore, Tg-ExoBooster-
positive beige adipocytes maintained higher *Ucp1* expression after whitening in non-adipogenic basal medium
compared to wild-type adipocytes (Fig. S2A). These cells also produced significantly more heat than wild-type
beige adipocytes, as measured by microcalorimetry (Fig. 2G-H). To test whether this enhanced thermogenic
phenotype was retained *in vivo*, Tg-ExoBooster mice were housed at thermoneutrality (29°C) for 7 days to
normalize basal energy expenditure, treated with doxycycline from days 2–7, and then subjected to a 3-day
cold challenge. Inguinal adipose tissue isolated from Tg-ExoBooster mice exhibited elevated *Ucp1* expression
(Fig. 2I) and generated more heat *ex vivo* than tissue from wild-type mice (Fig. 2J-K), as determined by
microcalorimetry. In contrast, no significant changes in *Ucp1* expression or heat production were observed in
brown adipose tissue under the same temperature conditions (Fig. S2B–D). Collectively, these data
demonstrate that genetic enhancement of exosome secretion can promote beige adipocyte thermogenesis
both *in vitro* and *in vivo*.

RAB27A deficiency impairs exosome secretion and thermogenesis in beige adipocytes. RAB27A is
critical for the fusion of multivesicular bodies (MVBs) with the plasma membrane and the subsequent release
of exosomes[41]. To determine whether impaired exosome secretion compromises thermogenic function, we
used a global *Rab27a* null mouse model[42]. Beige adipocytes were differentiated from the stromal vascular
fraction of inguinal adipose tissue and then exposed to either forskolin or vehicle to assess their thermogenic
capacity. Exosomes were isolated from the conditioned media and analyzed for particle number. Both *Rab27a*
heterozygous and null beige adipocytes secreted significantly fewer exosomes than wild-type cells under both
vehicle and forskolin-stimulated conditions (Fig. 3A-B), confirming a substantial impairment in exosome
release. A loss of thermogenic function accompanied this defect. *In vitro*, *Rab27a* heterozygous and null beige
adipocytes showed markedly reduced forskolin-induced *Ucp1* mRNA expression and UCP1 protein levels
compared to wild-type controls (Fig. 3C-D), along with significantly diminished forskolin-stimulated heat
production (Fig. 3E-F). To assess whether these deficits extended *in vivo*, wild-type and *Rab27a* null mice
were housed at thermoneutrality (29°C) for 7 days to normalize basal energy expenditure, followed by a 7-day
cold exposure challenge. Inguinal adipose tissue from cold-exposed wild-type mice exhibited robust induction
of *Ucp1* transcription and UCP1 protein expression, whereas *Rab27a* null mice showed reduced responses,
with *Ucp1* mRNA and protein levels decreased by approximately 60% and 40%, respectively, compared to
wild-type controls (Fig. 3G–H; Fig. S3). Microcalorimetry revealed inguinal fat pads from cold-treated wild-type
mice generated significantly more heat than those from *Rab27a* null mice (Fig. 3I–J). These experiments
support the conclusion that RAB27A-dependent exosome trafficking is a critical component of the thermogenic
signaling cascade and that disruption of this pathway compromises the ability of adipose tissue to properly
respond to thermogenic cues. The observation that *Rab27a* heterozygous mice consistently phenocopied nulls

across all assays suggests that beige adipocyte thermogenesis may require a threshold level of RAB27A
function, raising the possibility of haploinsufficiency or other dosage-sensitive mechanisms related to exosome
secretion.
224
RAB27A deficiency reduces energy expenditure in response to cold and β3-adrenergic stimulation. To
determine whether *Rab27a* deficiency affects whole-body metabolism, wild-type and *Rab27a* heterozygous
mice were housed at room temperature (~22°C) or exposed to cold (6°C) for 7 days, then transferred to
metabolic cages at room temperature. After a 24-hour acclimation period, energy expenditure, respiratory
exchange ratio, physical activity, food intake, and water consumption were measured over the next 48 hours
(see Supplementary Table 1 for detailed measurements and corresponding p-values). Body weight was similar
between wild-type and *Rab27a* heterozygous mice under both room temperature (RT) and cold conditions;
however, heterozygous mice displayed significantly higher fat mass at RT (p = 0.0005), as shown in Figure 4A.
This excess fat was lost during cold exposure, resulting in comparable fat mass across genotypes by day 7.
Indirect calorimetry revealed no difference in total energy expenditure between genotypes at RT over the 48-
hour recording period (Fig. 4B). However, after 7 days of cold exposure, *Rab27a* heterozygous mice exhibited
reduced energy expenditure compared to wild-type controls (Fig. 4C). Quantification of total 24-hour energy
expenditure showed a significant reduction in heterozygous mice under cold conditions (p = 0.02; Fig. 4D), with
the most pronounced difference observed during the daytime phase (p = 0.0037; Fig. 4E). A similar trend was
observed during the nighttime phase, although the difference did not reach statistical significance (p = 0.0993;
Fig. 4F). Further analysis of resting energy expenditure (REE) revealed significantly lower values in
heterozygous mice during both daytime (p = 0.0362; Fig. 4G) and nighttime (p = 0.0149; Fig. 4H) cold
exposure periods. The combined 24-hour REE was also significantly reduced in *Rab27a* heterozygous mice
compared to wild-type (p = 0.0114; Fig. 4I), indicating a deficit in basal thermogenic output. Ambulatory
activity, assessed by total distance walked in the cage, was significantly reduced in *Rab27a* heterozygous mice
under RT conditions (p = 0.0462; Fig. 4J), though this difference did not persist under cold exposure. No
significant differences were observed in respiratory exchange ratio (RER), food or water intake, or sleep
duration between genotypes under either temperature condition (see Supp. Table 1). In combination, these
data indicate that *Rab27a* heterozygosity impairs cold-induced increases in resting and total energy
expenditure, independent of behavioral or nutritional factors, supporting a role for RAB27A in the regulation of
basal thermogenesis.
To assess the role of Rab27a in β3-adrenergic-driven thermogenesis, wild-type and *Rab27a* null mice
were treated with CL316,243 (x mg/kg) daily for 9 days during metabolic cage analysis (see Supp. Table 2 for
complete data). Untreated controls were excluded, as wild-type baselines were established in the cold
exposure study, and *Rab27a* heterozygous mice displayed similar phenotypes to nulls (Fig. 3A–F). As with the
cold study, at baseline, *Rab27a* null mice had significantly greater total fat mass (in grams) than wild-type
controls (p = 0.0270; Fig. 4K; n = 8 mice per group). Although both groups lost similar proportions of fat mass
(12–15%) after treatment, wild-type mice were ~4% heavier by the end of the study (p = 0.0323; Supp. Table
2). This increased body weight in wild-type mice appears to result from a significant increase in lean mass
(+6%, p = 0.0087), which the null mice failed to achieve (−0.1%; p = 0.0051 compared to WT); a known
outcome of CL316,243-induced anabolic remodeling of skeletal muscle in rodents[43,44]. Total energy
expenditure over the 9-day period was reduced in null mice (Fig. 4L), and average 24-hour energy expenditure
across this period was significantly lower in nulls (p = 0.0020; Fig. 4M). Statistical comparisons for individual
day and nighttime intervals are shown in Fig. S3B. Similar to the effects observed under cold exposure (Fig.
3G), both *Rab27a* heterozygous and null mice exhibited significantly lower *Ucp1* transcript levels following
CL316,243 treatment compared to wild-type controls (Fig. 4N), consistent with a blunted thermogenic
response. Both average daytime (p = 0.0062) and nighttime (p = 0.0014) energy expenditure were reduced in
Rab27a null mice (Fig. 4O and 4P). Respiratory exchange ratio (RER) was significantly elevated during the
daytime in *Rab27a* null mice (p = 0.0056; Fig. 4Q), suggesting a shift toward carbohydrate metabolism and
reduced fatty acid oxidation, possibly due to lower UCP1 expression in the nulls (Fig. 3G and H). Resting
energy expenditure (REE) was also broadly impaired. Both nighttime REE (p = 0.0326; Fig. 4R) and 24-hour
REE (p = 0.0012, Supp. Table 2) were significantly reduced in null mice, indicating a persistent basal
thermogenic defect. Nighttime active energy expenditure (AEE) was also lower in *Rab27a* null mice (p =
0.0003; Fig. 4S), compounding the reduction in total energy output. No significant differences were observed in
food intake, water consumption, ambulatory activity (beam breaks, walking distance), or sleep duration
(Supplemental Table 2), excluding behavioral confounders.

Taken as a whole, these experiments reveal that RAB27A is required for the full thermogenic and
anabolic response to both cold and β_3-adrenergic stimulation. In its absence, mice exhibit reduced energy
expenditure, impaired lean mass gain, and altered substrate utilization, highlighting the critical role of RAB27A
and exosome secretion in adaptive thermogenesis.

Discussion

This study uncovers a previously unappreciated role for exosome secretion in supporting the thermogenic
capacity of beige adipocytes. We show that activation of beige adipocytes is associated with increased
exosome release and that these exosomes are enriched in microRNAs typically linked to the suppression of
thermogenic programming. Impairing exosome secretion, either through *Rab27a* deficiency or pharmacologic
inhibition with the small molecules manumycin A and GW4869, leads to a blunted thermogenic response. Mice
lacking *Rab27a* exhibit significantly reduced energy expenditure in response to cold exposure and β_3-
adrenergic stimulation, underscoring the functional importance of this pathway *in vivo*. Conversely, genetic
enhancement of exosome secretion via a multi-gene booster cassette (ExoBooster) increases UCP1
expression and heat output from beige adipocytes. In sum, these results establish that exosome secretion is
not simply a byproduct of adipocyte browning but rather a necessary component of the thermogenic process.
Our findings align with and extend the established concept that exosomes are critical mediators of
intercellular communication, particularly in metabolic regulation. We propose a model in which beige
adipocytes actively secrete exosomes enriched in inhibitory microRNAs as a potential self-regulatory
mechanism to sustain thermogenic activation. Regulation of these intracellular microRNAs may be necessary
to de-repress thermogenic transcription factors and enable full activation of mitochondrial pathways. This
model is supported by the observation that inhibition of exosome secretion blunts forskolin-induced UCP1
expression and thermogenesis, and that *Rab27a*-deficient adipocytes exhibit impaired UCP1 induction,
reduced heat production, and altered substrate utilization. Furthermore, the elevation in RER in *Rab27a*-null
mice during β_3-adrenergic stimulation (Fig. 4Q) suggests a shift toward carbohydrate metabolism and impaired
fatty acid oxidation, possibly due to reduced UCP1 expression (Fig. 3).
While our study primarily focuses on beige adipocytes, exosome-mediated regulation may also occur in
classical brown fat and other adipose depots, including visceral fat. Additional studies are needed to determine
whether this pathway functions similarly across these distinct adipose tissues. However, exosome secretion
may play a more prominent role in beige adipocytes, which must transition from a relatively quiescent or
"whitened" state back to a thermogenically active phenotype. This process likely requires more extensive
cellular remodeling and may depend on additional regulatory layers, including mechanisms for clearing
inhibitory factors during browning. Moreover, although our findings center on UCP1-dependent thermogenesis,
it remains possible that exosome secretion also influences UCP1-independent pathways, such as futile cycling
of calcium, creatine, or lipids. Given the emerging heterogeneity among beige adipocytes, it is also plausible
that only specific subsets rely on exosome secretion for full thermogenic activation, an area warranting further
investigation.
Importantly, our findings have clinical relevance for obesity and type 2 diabetes. Beige and brown
adipocytes are well-established targets for increasing energy expenditure and improving glucose metabolism.
Strategies to enhance their activity have traditionally relied on cold exposure or β_3-adrenergic agonists, both of
which have translational limitations. Our work identifies exosome secretion as a novel and targetable
mechanism to augment thermogenesis in beige adipocytes, potentially offering a new approach for increasing
energy expenditure in obese individuals. Furthermore, because *Rab27a*-deficient mice fail to increase lean
mass in response to CL316,243, this work raises the possibility that exosome pathways also intersect with
anabolic remodeling and muscle-adipose crosstalk[43,44].
There are, however, several limitations to consider. First, while our data support a functional role for
exosome secretion in promoting thermogenesis, it remains unclear whether this effect is mediated by specific
exosomal cargo or by broader aspects of the secretion process itself. Our candidate screen identified several
microRNAs (miR-27a/b, miR-133a, miR-99b) that are typically linked to repression of thermogenic gene
programs, but further studies are needed to determine whether their export plays an active role in adipocyte
browning or simply correlates with it. Second, the global *Rab27a* knockout model has known effects beyond
adipose tissue, which may confound some systemic phenotypes. Future work using adipocyte-specific *Rab27a*
deletion models would help isolate the cell-autonomous effects. Lastly, while we demonstrate effects in mouse
and iPSC-derived human models, further work is required to confirm the relevance of this pathway in human
obesity.

In summary, this study reveals a novel regulatory axis linking exosome secretion to beige adipocyte
thermogenesis. By demonstrating that both enhancement and disruption of exosome pathways directly impact
energy expenditure, this work provides new insight into adipocyte biology and identifies exosome secretion as
a potential target for anti-obesity therapies.
334

Materials and Methods

Statistical Analysis. Unless otherwise stated, all data are presented as mean ± standard deviation (SD) for *in*
vitro experiments and mean ± standard error of the mean (SEM) for *in vivo* studies. The number of biological
replicates (n) is indicated in the figure legends. Comparisons between two groups were analyzed using
unpaired two-tailed Student's t-tests. For multiple group comparisons, one-way or two-way analysis of variance
(ANOVA) was performed as appropriate, followed by Tukey's multiple comparisons test. MicroRNA profiling
statistics (Fig. 1D) were performed using Global Pattern Recognition software[45]. A p-value < 0.05 was
considered statistically significant. Significance levels are indicated as follows: $P < 0.05$ (*), $P < 0.01$ (**), $P <$
0.001 (***), $P < 0.0001$ (****).

Differentiation and Activation of Human and Murine Beige Adipocytes. Human iPSC-derived beige
adipocytes were differentiated for 12 days using an adipogenic induction cocktail in EGM2 medium (Lonza), as
previously described[25]. To induce whitening, mature beige adipocytes were cultured for 4 days in non-
adipogenic basal medium consisting of high-glucose DMEM (ATCC, 30-2002) supplemented with 10%
exosome-depleted fetal bovine serum (System Biosciences, EXO-FBS-250A-1). Where indicated, cells were
treated with 2 µM manumycin A (Sigma, M6418) or 10 µM GW4869 (Sigma, D1692) for 2 hours at the
whitening phase to inhibit exosome secretion. Murine beige adipocytes were derived from the stromal vascular
fraction (SVF) of inguinal white adipose tissue harvested from female 8-week-old C57BL/6J mice. Tissue was
finely minced with a sterile razor blade and digested in isolation buffer containing 0.1% collagenase P (Roche)
at 37°C for 45–60 minutes on a nutator, with brief vortexing every 10 minutes. The digest was filtered through
a 70 µm cell strainer and centrifuged at 300 × g for 5 minutes. The SVF pellet was washed with autoMACS
running buffer (Miltenyi), centrifuged again, and resuspended in primary culture medium (high-glucose DMEM
+ 20% FBS + 1% penicillin–streptomycin). Passage 2–3 cells were differentiated for 6 days using the same
human adipogenic induction cocktail in DMEM + 10% FBS (ATCC, 30-2020), followed by 6 days of whitening
in DMEM + 10% exosome-depleted FBS. Thermogenic activation of both human and mouse whitened beige
adipocytes was induced with 10 µM forskolin for the indicated durations. For all exosome-boosting
experiments, Tg-ExoBoost and WT cells were treated with 2 µg/mL doxycycline hyclate (Sigma, D5207) for 72
hours.

Murine Mouse Models and *In Vivo* Treatment Protocols. All experiments were performed using female mice
harvested at 8-10 weeks of age. Transgenic mice expressing the doxycycline-inducible ExoBooster cassette
were generated by the Mouse Genome Modification Core at MaineHealth Institute for Research (MHIR) via
pronuclear microinjection of the linearized construct into fertilized C57BL/6J zygotes. The ExoBooster
construct includes a Tet-On 3G system and a bidirectional TRE3G promoter, enabling doxycycline-dependent
expression of a tricistronic cassette encoding STEAP3, SDC4, and NadB. Beige adipocytes used for *in vitro*
studies were derived from the stromal vascular fraction of inguinal white adipose tissue, as described above.
For in vivo activation of Tg-ExoBooster, mice were housed at thermoneutrality (29°C) for 10 days to normalize
basal energy expenditure. Doxycycline was administered in chow (625 mg/kg; Inotiv) from days 7–10. On day
10, mice were transferred to a cold environment (6°C) for an additional 3 days to stimulate thermogenesis. For
Rab27a studies, wild-type, heterozygous, and Rab27a null mice were generated as previously described[42].
Mice were housed at either room temperature (~22°C) or cold conditions (6°C) for 7 days prior to metabolic
cage analysis and phenotyping. For CL316,243 studies, mice were housed in metabolic cages and received
daily intraperitoneal injections of the β3-adrenergic receptor agonist CL316,243 (1 mg/kg body weight) for 9
consecutive days to stimulate adipose thermogenesis.

Metabolic Cage Analysis and Body Composition. Whole-body energy expenditure, respiratory exchange
ratio (RER), resting energy expenditure (REE), active energy expenditure (AEE), physical activity (distance
walked), food and water intake, and sleep behavior were monitored using the Promethion Core metabolic cage
system (Sable Systems International) after a 24-hour acclimation. Body composition was measured using dual-
energy X-ray absorptiometry (DXA; Faxitron Bioptics, LLC).

Exosome Isolation, Quantification and Inhibition. Exosomes were isolated from conditioned medium using
the miRCURY Exosome Cell/Urine/CSF Kit (Qiagen) according to the manufacturer's instructions. Particle size

distribution and concentration were measured by nanoparticle tracking analysis (NTA) using the ZetaView
instrument (Particle Metrix GmbH). Media was collected from wells seeded with equal numbers of cells and
equal volumes, allowing direct comparison of absolute particle counts across conditions. Particle
concentrations are reported as particles per mL and shown as absolute values relative to untreated controls.
Where indicated, exosome secretion was inhibited by pretreating cells with manumycin A (1 µM, Sigma) or
GW4869 (10 µM, Sigma) 2 hours before forskolin treatment, unless otherwise specified.

qPCR and Western Blotting. Total RNA, including microRNAs, was extracted using the miRNeasy Micro Kit
(Qiagen). cDNA was synthesized with qScript cDNA SuperMix (Quanta Biosciences, 95048-100), and
quantitative PCR was performed using AzuraView GreenFast qPCR Blue Mix LR (Azura Genomics, AZ-2320)
on a Bio-Rad CFX384 real-time PCR system. Primers used for amplification are listed in Table X. Gene
expression was analyzed using the ∆∆Ct method, normalized to *Actb* for mRNA or U6/18S for microRNAs, and
expressed relative to vehicle-treated or day 0 controls unless otherwise specified. Melt curve analysis
confirmed primer specificity. Protein expression was assessed by Western blot as previously described[46],
using β-actin or β-tubulin as loading controls.

Heat Production Assay via Microcalorimetry. Heat production from beige adipocytes or freshly isolated
adipose tissue was measured using high-resolution isothermal microcalorimetry on the calScreener™ system
(Symcel). Equivalent numbers of beige preadipocytes were differentiated in 200 µL of medium (as described
above) in Geltrex-coated calWell™ insert plates (Symcel, 1900402). Inserts were transferred to screw-capped
titanium vials. The calPlate™ (48-well format) was pre-equilibrated in two thermal zones for 30 minutes,
followed by a 15–30 minute stabilization period after final positioning. Continuous heat output was recorded
using calView™ software. For *ex vivo* analysis, adipose tissue biopsies were taken from identical locations
within the fat pad, placed in calWells containing 200 µL DMEM, and normalized by weight. All measurements
were normalized to internal reference wells containing media alone.

Animal Ethics. All animal studies were conducted in compliance with the National Institutes of Health Guide
for the Care and Use of Laboratory Animals and were approved by the Institutional Animal Care and Use
Committee (IACUC; protocol #2203) at the MaineHealth Institute for Research. Mice were housed in pathogen-
free, AAALAC-accredited facilities with *ad libitum* access to food and water. Every effort was made to minimize
animal suffering and reduce the number of animals used.

Data Availability. All data supporting the findings of this study are included in the article and its
Supplementary Information. Raw and processed data from RNA and microRNA qPCR screens,
microcalorimetry, and metabolic cage studies are available from the corresponding author upon reasonable
request.

Acknowledgements

This work was supported by NIH COBRE award P20GM121301 (A. Brown, L. Liaw, and C.J. Rosen) and
NIDDK award 1R01DK124261 (A. Brown). The project utilized the services of Core Facilities funded by NIH
award U54GM115516 (C.J Rosen, PI). We kindly thank Drs. Daniel Bojar and Martin Fussenegger for
providing the exosome booster plasmids.

References:
422

1. Lumeng CN, Saltiel AR. Inflammatory links between obesity and metabolic disease. *J Clin Invest*.
2011;121:2111-2117. doi: 10.1172/JCI57132
2. Wang L, Jia Q, He J, Li Y. Adipose tissue-targeting nanomedicines for obesity pharmacotherapy.
Trends Endocrinol Metab. 2025. doi: 10.1016/j.tem.2025.03.010
3. Lauti M, Kularatna M, Hill AG, MacCormick AD. Weight Regain Following Sleeve Gastrectomy-a
Systematic Review. *Obes Surg*. 2016;26:1326-1334. doi: 10.1007/s11695-016-2152-x
4. Harms M, Seale P. Brown and beige fat: development, function and therapeutic potential. *Nat Med*.
2013;19:1252-1263. doi: 10.1038/nm.3361
5. Cypess AM, Lehman S, Williams G, Tal I, Rodman D, Goldfine AB, Kuo FC, Palmer EL, Tseng YH,
Doria A, et al. Identification and importance of brown adipose tissue in adult humans. *N Engl J Med*.
2009;360:1509-1517. doi: 10.1056/NEJMoa0810780
6. Wu J, Cohen P, Spiegelman BM. Adaptive thermogenesis in adipocytes: is beige the new brown?
Genes Dev. 2013;27:234-250. doi: 10.1101/gad.211649.112

7.   Becher T, Palanisamy S, Kramer DJ, Eljalby M, Marx SJ, Wibmer AG, Butler SD, Jiang CS, Vaughan
R, Schoder H, et al. Brown adipose tissue is associated with cardiometabolic health. *Nat Med*.
2021;27:58-65. doi: 10.1038/s41591-020-1126-7
8.   Cannon B, Nedergaard J. Brown adipose tissue: function and physiological significance. *Physiol Rev*.
2004;84:277-359. doi: 10.1152/physrev.00015.2003
9.   Cero C, Lea HJ, Zhu KY, Shamsi F, Tseng YH, Cypess AM. beta3-Adrenergic receptors regulate
human brown/beige adipocyte lipolysis and thermogenesis. *JCI Insight*. 2021;6. doi:
10.1172/jci.insight.139160
10.   Rahbani JF, Scholtes C, Lagarde DM, Hussain MF, Roesler A, Dykstra CB, Bunk J, Samborska B,
O'Brien SL, Tripp E, et al. ADRA1A-Galpha(q) signalling potentiates adipocyte thermogenesis through
CKB and TNAP. *Nat Metab*. 2022;4:1459-1473. doi: 10.1038/s42255-022-00667-w
11.   Cohen P, Kajimura S. The cellular and functional complexity of thermogenic fat. *Nat Rev Mol Cell Biol*.
2021;22:393-409. doi: 10.1038/s41580-021-00350-0
12.   Wang T, Sharma AK, Wu C, Maushart CI, Ghosh A, Yang W, Stefanicka P, Kovanicova Z, Ukropec J,
Zhang J, et al. Single-nucleus transcriptomics identifies separate classes of UCP1 and futile cycle
adipocytes. *Cell Metab*. 2024;36:2130-2145 e2137. doi: 10.1016/j.cmet.2024.07.005
13.   Vargas-Castillo A, Sun Y, Smythers AL, Grauvogel L, Dumesic PA, Emont MP, Tsai LT, Rosen ED,
Zammit NW, Shaffer SM, et al. Development of a functional beige fat cell line uncovers independent
subclasses of cells expressing UCP1 and the futile creatine cycle. *Cell Metab*. 2024;36:2146-2155
e2145. doi: 10.1016/j.cmet.2024.07.002
14.   Carpentier AC, Blondin DP, Haman F, Richard D. Brown Adipose Tissue-A Translational Perspective.
Endocr Rev. 2023;44:143-192. doi: 10.1210/endrev/bnac015
15.   Malicka A, Ali A, MacCannell ADV, Roberts LD. Brown and beige adipose tissue-derived metabokine
and lipokine inter-organ signalling in health and disease. *Exp Physiol*. 2024. doi: 10.1113/EP092008
16.   Wang GX, Zhao XY, Lin JD. The brown fat secretome: metabolic functions beyond thermogenesis.
Trends Endocrinol Metab. 2015;26:231-237. doi: 10.1016/j.tem.2015.03.002
17.   Mei R, Qin W, Zheng Y, Wan Z, Liu L. Role of Adipose Tissue Derived Exosomes in Metabolic Disease.
Front Endocrinol (Lausanne). 2022;13:873865. doi: 10.3389/fendo.2022.873865
18.   Kulaj K, Harger A, Bauer M, Caliskan OS, Gupta TK, Chiang DM, Milbank E, Reber J, Karlas A,
Kotzbeck P, et al. Adipocyte-derived extracellular vesicles increase insulin secretion through transport
of insulinotropic protein cargo. *Nat Commun*. 2023;14:709. doi: 10.1038/s41467-023-36148-1
19.   Kariba Y, Yoshizawa T, Sato Y, Tsuyama T, Araki E, Yamagata K. Brown adipocyte-derived exosomal
miR-132-3p suppress hepatic Srebf1 expression and thereby attenuate expression of lipogenic genes.
Biochem Biophys Res Commun. 2020;530:500-507. doi: 10.1016/j.bbrc.2020.05.090
20.   Thomou T, Mori MA, Dreyfuss JM, Konishi M, Sakaguchi M, Wolfrum C, Rao TN, Winnay JN, Garcia-
Martin R, Grinspoon SK, et al. Adipose-derived circulating miRNAs regulate gene expression in other
tissues. *Nature*. 2017;542:450-455. doi: 10.1038/nature21365
21.   Chen Y, Buyel JJ, Hanssen MJ, Siegel F, Pan R, Naumann J, Schell M, van der Lans A, Schlein C,
Froehlich H, et al. Exosomal microRNA miR-92a concentration in serum reflects human brown fat
activity. *Nat Commun*. 2016;7:11420. doi: 10.1038/ncomms11420
22.   Ying W, Riopel M, Bandyopadhyay G, Dong Y, Birmingham A, Seo JB, Ofrecio JM, Wollam J,
Hernandez-Carretero A, Fu W, et al. Adipose Tissue Macrophage-Derived Exosomal miRNAs Can
Modulate In Vivo and In Vitro Insulin Sensitivity. *Cell*. 2017;171:372-384 e312. doi:
10.1016/j.cell.2017.08.035
23.   Vishvanath L, MacPherson KA, Hepler C, Wang QA, Shao M, Spurgin SB, Wang MY, Kusminski CM,
Morley TS, Gupta RK. Pdgfrbeta+ Mural Preadipocytes Contribute to Adipocyte Hyperplasia Induced by
High-Fat-Diet Feeding and Prolonged Cold Exposure in Adult Mice. *Cell Metab*. 2016;23:350-359. doi:
10.1016/j.cmet.2015.10.018
24.   Roh HC, Tsai LTY, Shao M, Tenen D, Shen Y, Kumari M, Lyubetskaya A, Jacobs C, Dawes B, Gupta
RK, et al. Warming Induces Significant Reprogramming of Beige, but Not Brown, Adipocyte Cellular
Identity. *Cell Metab*. 2018;27:1121-1137 e1125. doi: 10.1016/j.cmet.2018.03.005
25.   Su S, Guntur AR, Nguyen DC, Fakory SS, Doucette CC, Leech C, Lotana H, Kelley M, Kohli J, Martino
J, et al. A Renewable Source of Human Beige Adipocytes for Development of Therapies to Treat
Metabolic Syndrome. *Cell Rep*. 2018;25:3215-3228 e3219. doi: 10.1016/j.celrep.2018.11.037
26.   Karolina DS, Armugam A, Sepramaniam S, Jeyaseelan K. miRNAs and diabetes mellitus. *Expert Rev
Endocrinol Metab*. 2012;7:281-300. doi: 10.1586/eem.12.21

27. Guller I, McNaughton S, Crowley T, Gilsanz V, Kajimura S, Watt M, Russell AP. Comparative analysis of microRNA expression in mouse and human brown adipose tissue. *BMC Genomics*. 2015;16:820. doi: 10.1186/s12864-015-2045-8

28. He Y, Ding Y, Liang B, Lin J, Kim TK, Yu H, Hang H, Wang K. A Systematic Study of Dysregulated MicroRNA in Type 2 Diabetes Mellitus. *Int J Mol Sci*. 2017;18. doi: 10.3390/ijms18030456

29. Ji C, Guo X. The clinical potential of circulating microRNAs in obesity. *Nat Rev Endocrinol*. 2019;15:731-743. doi: 10.1038/s41574-019-0260-0

30. Gharanei S, Shabir K, Brown JE, Weickert MO, Barber TM, Kyrou I, Randeva HS. Regulatory microRNAs in Brown, Brite and White Adipose Tissue. *Cells*. 2020;9. doi: 10.3390/cells9112489

31. Cuevas-Ramos D, Mehta R, Aguilar-Salinas CA. Fibroblast Growth Factor 21 and Browning of White Adipose Tissue. *Front Physiol*. 2019;10:37. doi: 10.3389/fphys.2019.00037

32. Sun L, Trajkovski M. MiR-27 orchestrates the transcriptional regulation of brown adipogenesis. *Metabolism*. 2014;63:272-282. doi: 10.1016/j.metabol.2013.10.004

33. Yu J, Lv Y, Wang F, Kong X, Di W, Liu J, Sheng Y, Lv S, Ding G. MiR-27b-3p Inhibition Enhances Browning of Epididymal Fat in High-Fat Diet Induced Obese Mice. *Front Endocrinol (Lausanne)*. 2019;10:38. doi: 10.3389/fendo.2019.00038

34. Liu W, Bi P, Shan T, Yang X, Yin H, Wang YX, Liu N, Rudnicki MA, Kuang S. miR-133a regulates adipocyte browning in vivo. *PLoS Genet*. 2013;9:e1003626. doi: 10.1371/journal.pgen.1003626

35. Zhang MQ, Gao JL, Liao XD, Huang TH, Zhang MN, Wang MQ, Tian Y, Bai J, Zhou CH. miR-454 regulates triglyceride synthesis in bovine mammary epithelial cells by targeting PPAR-gamma. *Gene*. 2019;691:1-7. doi: 10.1016/j.gene.2018.12.048

36. Lee EK, Lee MJ, Abdelmohsen K, Kim W, Kim MM, Srikantan S, Martindale JL, Hutchison ER, Kim HH, Marasa BS, et al. miR-130 suppresses adipogenesis by inhibiting peroxisome proliferator-activated receptor gamma expression. *Mol Cell Biol*. 2011;31:626-638. doi: 10.1128/MCB.00894-10

37. Tao H, Wang MM, Zhang M, Zhang SP, Wang CH, Yuan WJ, Sun T, He LJ, Hu QK. MiR-126 Suppresses the Glucose-Stimulated Proliferation via IRS-2 in INS-1 beta Cells. *PLoS One*. 2016;11:e0149954. doi: 10.1371/journal.pone.0149954

38. Datta A, Kim H, Lal M, McGee L, Johnson A, Moustafa AA, Jones JC, Mondal D, Ferrer M, Abdel-Mageed AB. Manumycin A suppresses exosome biogenesis and secretion via targeted inhibition of Ras/Raf/ERK1/2 signaling and hnRNP H1 in castration-resistant prostate cancer cells. *Cancer Lett*. 2017;408:73-81. doi: 10.1016/j.canlet.2017.08.020

39. Trajkovic K, Hsu C, Chiantia S, Rajendran L, Wenzel D, Wieland F, Schwille P, Brugger B, Simons M. Ceramide triggers budding of exosome vesicles into multivesicular endosomes. *Science*. 2008;319:1244-1247. doi: 10.1126/science.1153124

40. Kojima R, Bojar D, Rizzi G, Hamri GC, El-Baba MD, Saxena P, Auslander S, Tan KR, Fussenegger M. Designer exosomes produced by implanted cells intracerebrally deliver therapeutic cargo for Parkinson's disease treatment. *Nat Commun*. 2018;9:1305. doi: 10.1038/s41467-018-03733-8

41. Ostrowski M, Carmo NB, Krumeich S, Fanget I, Raposo G, Savina A, Moita CF, Schauer K, Hume AN, Freitas RP, et al. Rab27a and Rab27b control different steps of the exosome secretion pathway. *Nat Cell Biol*. 2010;12:19-30; sup pp 11-13. doi: 10.1038/ncb2000

42. Soucy A, Potts C, Kaija A, Harrington A, McGilvrey M, Sutphin GL, Korstanje R, Tero B, Seeker J, Pinz I, et al. Effects of a Global Rab27a Null Mutation on Murine PVAT and Cardiovascular Function. *Arterioscler Thromb Vasc Biol*. 2024. doi: 10.1161/ATVBAHA.124.320969

43. Puzzo D, Raiteri R, Castaldo C, Capasso R, Pagano E, Tedesco M, Gulisano W, Drozd L, Lippiello P, Palmeri A, et al. CL316,243, a beta3-adrenergic receptor agonist, induces muscle hypertrophy and increased strength. *Sci Rep*. 2016;5:37504. doi: 10.1038/srep37504

44. Ding LN, Cheng Y, Xu LY, Zhou LQ, Guan L, Liu HM, Zhang YX, Li RM, Xu JW. The beta3 Adrenergic Receptor Agonist CL316243 Ameliorates the Metabolic Abnormalities of High-Fat Diet-Fed Rats by Activating AMPK/PGC-1alpha Signaling in Skeletal Muscle. *Diabetes Metab Syndr Obes*. 2021;14:1233-1241. doi: 10.2147/DMSO.S297351

45. Akilesh S, Shaffer DJ, Roopenian D. Customized molecular phenotyping by quantitative gene expression and pattern recognition analysis. *Genome Res*. 2003;13:1719-1727. doi: 10.1101/gr.533003

46. Doucette CC, Nguyen DC, Barteselli D, Blanchard S, Pelletier M, Kesharwani D, Jachimowicz E, Su S, Karolak M, Brown AC. Optogenetic activation of UCP1-dependent thermogenesis in brown adipocytes. *iScience*. 2023;26:106560. doi: 10.1016/j.isci.2023.106560

Figures and Tables

Figure 1. Exosome secretion correlates with thermogenic activation in beige adipocytes and is conserved across human and mouse models. (A) RT-qPCR analysis of *PGC1α* and *UCP1* transcript levels in human iPSC-derived beige adipocytes following 6-hour forskolin treatment (10 μm, n = 3). (B, C) Nanoparticle tracking analysis (NTA) of exosomes from beige adipocytes ± forskolin treatment for 6 hours, showing (B) size distribution and (C) >8-fold increase in concentration (n = 3). (D) Differentially expressed microRNAs in exosomes from beige adipocytes with or without forskolin treatment (n = 3), ranked by fold change. Orange indicates increased expression with forskolin, blue indicates decreased expression. (E) Time-course qPCR analysis of intracellular miR-27a/b-3p, miR-99b-5p, and miR-133a-5p in beige adipocytes following forskolin treatment, showing a marked reduction by 1 h (n = 3 per time point). Time zero normalized to 100%. (F) *UCP1* expression in forskolin-treated beige adipocytes is reduced by co-treatment with exosome biogenesis and release inhibitors manumycin A (manA) or GW4869 (GW) (n = 3). (G) Oxygen consumption rate (OCR) analysis via Seahorse XF analyzer showing reduced proton leak-linked respiration in forskolin-stimulated beige adipocytes ± exosome inhibitors (n = 3). (H, I) NTA of exosomes from mouse beige adipocytes ± forskolin and inhibitors: (H) size distribution by bin; (I) total particle concentration (n = 5). (J, K) qPCR showing forskolin-induced enrichment of miR-27a/b, miR-99b-5p, and miR-133a-5p in mouse beige-derived exosomes after 4-hour treatment (J) and their corresponding intracellular depletion (K) (n = 3). (L) *Ucp1* mRNA expression in forskolin-treated mouse beige adipocytes ± exosome inhibitors after 4-hour treatment (n = 3). (M) Western blot showing UCP1 protein reduction in a dose-dependent manner with exosome inhibitor co-treatment (n = 3 pooled replicates with β-tubulin as loading control). (N, O) Microcalorimetry of cultured beige adipocytes showing (N) average heat accumulation over time and (O) total heat output (quantified from N, with error bars) in the presence or absence of forskolin and exosome inhibitors (n = 3). Refer to Methods for statistical tests and significance thresholds.

Figure 2. Genetically boosted exosome secretion enhances thermogenesis in beige adipocytes. (A)
Schematic of the Tg-ExoBooster construct. The bidirectional TRE3G promoter drives upstream expression of
Tet-On 3G and downstream expression of a tricistronic STEAP3–IRES–SDC4–IRES–NadB cassette after
addition of doxycycline. p(A), polyadenylation signal; IRES, internal ribosome entry site. (B) RT-PCR validation
of ExoBooster transgene expression in cultured beige adipocytes after 72 hours doxycycline treatment (n = 3).
(C) Nanoparticle tracking analysis (NTA) of exosome size distribution from Tg-ExoBooster versus wild-type
(WT) beige adipocytes treated with doxycycline. (D) NTA analysis of total exosome particle counts from Tg-
ExoBooster versus WT beige adipocytes treated with doxycycline and with or without inhibitor co-treatment (n
= 3). (E) qPCR analysis of *Ucp1* expression in Tg-ExoBooster and WT beige adipocytes after 4-hour forskolin
stimulation (n = 3). (F) Western blot analysis of UCP1 protein levels from pooled replicates (n=3). (G, H)
Microcalorimetry traces showing (G) average heat accumulation over time and (H) total heat output (quantified
from G, with error bars) in forskolin-treated Tg-ExoBooster and wild-type beige adipocytes following
doxycycline treatment (n = 5). (I) RT-qPCR analysis of *Ucp1* expression in inguinal adipose tissue from wild-
type and Tg-ExoBooster mice after 3 days doxycycline feed and cold challenge (n = 5). (J, K) *Ex vivo*
microcalorimetry traces (J) and quantified heat output (K) from inguinal fat pads post-doxycycline and cold
challenge (n = 5). Heat data in J-K normalized to tissue weight in grams (g).

Figure 3. *Rab27a* deficiency impairs exosome secretion and thermogenesis in beige adipocytes. (A, B)
Nanoparticle tracking analysis showing (A) average exosome count and size distribution and (B) total exosome
particle counts (quantified from A, with error bars) from wild-type, *Rab27a* heterozygous, and *Rab27a* null
beige adipocytes ± forskolin (n = 4). (C) qPCR analysis of *Ucp1* mRNA expression in cultured beige adipocytes
± forskolin (n = 3). (D) Western blot of UCP1 protein in cultured beige adipocytes (n = 3 replicates pooled per
treatment). β-tubulin serves as loading control. (E, F) Microcalorimetry traces showing (E) average heat
accumulation over time and (F) total heat output (quantified from E, with error bars) from beige adipocytes (n =
3). (G) *Ucp1* expression in inguinal fat following cold exposure in wild-type, *Rab27a* heterozygous, and null
mice (n = 5 mice per group). (H) Immunoblot quantitation of UCP1 protein from inguinal adipose tissue (n = 6
mice per group). Data are normalized to β-tubulin (see Fig. S3A for blot). (I, J) Microcalorimetry traces showing
(I) average heat accumulation over time and (J) total heat output (quantified from I, with error bars) from
inguinal adipose tissue of RT and cold-exposed mice (n = 4 per group). Heat data in I-J normalized to tissue
weight in grams (g).

Figure 4. *Rab27a* deficiency reduces energy expenditure in response to cold and β3-adrenergic
stimulation. (A) Fat mass of wild-type and *Rab27a* heterozygous female mice housed at room temperature
(RT, ~22°C) or cold exposed (6°C) for 7 days (n = 6-8 mice per group for A-J). (B, C) Metabolic cage analysis
of energy expenditure over time at (B) room temperature and (C) following cold exposure. (D–F) 24-hour,
daytime, and nighttime energy expenditure after cold exposure. (G–I) Daytime, nighttime, and combined 24-
hour resting energy expenditure (REE). (J) Ambulatory activity at RT and cold. (K) Fat mass before and after
CL316,243 treatment in wild-type and *Rab27a* null mice. n = 6 mice per group for L-S. (L) Average total energy
expenditure during 9-day CL316,243 treatment. (M) Average 24-hour energy expenditure over 9 days. (N) RT-
qPCR analysis of *Ucp1* mRNA in inguinal adipose tissue ± CL316,243 treatment (day 9). (O, P) Average
daytime and nighttime energy expenditure (Q), daytime RER and (R) active energy expenditure (AEE) over 9
days.