Lessons from a gene knockout mouse model: no phenotypes induced by

deletion of the TPR4 domain of Ttc22

Jiaoyang Liu, Liankun Gu, Abin You, Hongfan Yuan, Jing Zhou, Wei Tian*, Dajun Deng*

Key Laboratory of Carcinogenesis and Translational Research (MOE/Beijing), Division of Etiology, Peking

University Cancer Hospital and Institute, Beijing, 100142, China

Corresponding authors: Wei Tian or Dajun Deng, Key Laboratory of Carcinogenesis and Translational Research

(MOE/Beijing), Division of Etiology, Peking University Cancer Hospital and Institute, Fu-Cheng-Lu #52, Haidian

District, Beijing, 100142, China, Phone:+8610-88196752 (E-mail: tianwei@bjmu.edu.cn or

dengdajun@bjmu.edu.cn; ORCID: 0000-0003-2853-7056 or 0000-0001-5161-5943)

Highlights

● Knockout of *Ttc22* exons 2 and 3 does not affect the development of C57BL mice.

● *Ttc22* knockout does not affect the induction of mouse colon cancer by AOM/DSS.

● Loss of the *Ttc22* exons 2 and 3 cannot disrupt the functions of TTC22.

Email address

Jiaoyang Liu, pocky1995@126.com

Liankun Gu, gulkun@163.com

Abin You, youabin115@126.com

Hongfan Yuan, yuanhongfan@scszlyy.org.cn

Jing Zhou, jane72@163.com

Wei Tian, tianwei@bjmu.edu.cn

Dajun Deng, dengdajun@bjmu.edu.cn

ABSTRACT

The *TTC22* gene encodes a protein containing seven tetratricopeptide repeats (TPRs), which mediate protein−protein interactions as chaperones. We previously reported that the level of *TTC22* transcript variant 1 (*TTC22v1*) was downregulated in human colon adenocarcinoma (COAD) and that TTC22 upregulated m6A-mediated *WTAP* and *SNAI1* expression via the TTC22−RPL4 interaction and subsequently promoted COAD metastasis. Thus, a commercially available C57BL/6N mouse model in which the *Ttc22* exon 2&3 encoding the TPR4 (equal to the human TTC22 TPR3) domain was knocked out via CRISPR-Cas9 was used to evaluate the contribution of *Ttc22* to the development of mice and COAD. Unfortunately, the long-term observation results demonstrated that *Ttc22* knockout (KO, including *Ttc22*$^{-/+}$ or *Ttc22*$^{-/-}$) did not affect the body weight, development, fertility, or spontaneous tumor incidence of male or female mice. No differences in the incidence of AOM/DSS-induced COAD were observed between these mouse groups, although *Ttc22* exon 2&3 deletion partially resulted in the upregulation of adaptive response genes in the colon mucosa. Further study revealed that TPR4 deletion did not disrupt the effect of TTC22 on WTAP upregulation. Consistently, TPR4 deletion did not affect the abundance of total RNA m6A in the colon tissues of the mice, suggesting that the TPR4-deleted Ttc22 mutant remains functional. In conclusion, the TPR4 domain is not essential for the Ttc22 protein. Loss of Ttc22 TPR4 caused no observable changes in the development of C57BL mice or their susceptibility to treatment with the chemical carcinogen AOM/DSS. Whether an essential sequence for a target gene is knocked out should be carefully evaluated before a gene knockout model is employed in formal experiments.

Keywords: Ttc22; TPR4 deletion; C57BL/6N mouse; development; colon cancer; AOM/DSS

INTRODUCTION

The human TTC22 protein contains seven tetratricopeptide repeats (TPRs), which mediate protein–protein interactions and are involved in multiple biological processes, including the movements of cilia and flagella and the migration of human cells [1-5]. A pilot study reported that the *TTC22* gene was among the top ten downregulated genes in the fetuses of obese women compared with lean women in the second trimester and that low *TTC22* expression in mothers may be related to fetal neurodevelopment and metabolism [6]. Differentially methylated regions of the *TTC22* gene are associated with hand grip strength [7]. DNA sequencing and transcriptome data analyses in 3,203 subjects suggested that low-frequency variants of *TTC22* were associated with LDL cholesterol [8]. These phenomena imply that TTC22 may be involved in physiological metabolism. However, the causality between TTC22 alterations and metabolism is not known.

The *TTC22* gene is comprehensively expressed in epithelial cells in the normal gut mucosa at the middle level and is extensively downregulated in gastrointestinal cancer tissues, according to the databases from the Human Protein Atlas and The Cancer Genome Atlas (TCGA) [9,10]. We initially reported that *TTC22* variant 1 (*TTC22v1*) mRNA is targeted by *miR663a*, an anti-inflammatory miRNA in the colon [11]. The level of *TTC22v1* mRNA is significantly lower in human colon adenocarcinoma (COAD) than in paired normal tissue samples, and high *TTC22* expression increases the risk of colon cancer metastasis. Our further study revealed that TTC22 increased m6A-mediated *WTAP* and *SNAI1* expression via TTC22-RPL4 binding and subsequently promoted COAD metastasis *in vitro* and *in vivo* [12], suggesting a potential role for TTC22 in the development and progression of COAD.

Therefore, we further studied the effects of Ttc22 on the development of mice and their susceptibility to the chemical carcinogen AOM/DSS via a genetically established and commercially available C57BL gene knockout mouse model in which *Ttc22* exons 2 and 3, which encode the TPR4 domain of the mouse Ttc22 protein, were knocked out via CRISPR–Cas9. Unexpectedly, no differences in the development, growth, fertility, or AOM/DSS susceptibility of the mice were induced by TPR4 deletion. Our further study results indicated that the TPR3 domain (equal to mouse TTC22 TPR4) is not essential for TTC22 to upregulate WTAP and related m6A modifications. Although commercially available gene knockout models are useful for scientists in molecular biology and pathogenesis, whether such models are eligible for use should be carefully evaluated before they are formally used in experiments.

MATERIALS AND METHODS

Ttc22 exon 2&3 knockout mouse model

Genetically established *Ttc22* knockout (Ttc22-KO) C57BL/6N mice (strain S-KO-06335) were purchased from Cyagen (Cyagen Bio. Co., Guangzhou, China). Ttc22 knockout mice were obtained via high-throughput electroporation of fertilized eggs. The CRISPR-Cas9 system and two gRNAs (gRNA#1: 5'-ggaaagugggcauaacgccuggg-3' and gRNA#2: 5'-uuccagagagauagcgaccgagg-3') targeted to *Ttc22* introns 1&3 were used to delete a 1610 bp fragment including *Ttc22* exons 2&3 encoding the Ttc22 TPR4 domain (Figure 1A). After sexual maturity, sperm were collected for cryopreservation. Through hybridization with female wild-type C57BL/6N mice, first-generation Ttc22$^{+/-}$ mice were obtained and used to generate male and female mice with different *Ttc22* genotypes (including Ttc22$^{+/+}$, Ttc22$^{+/-}$, and Ttc22$^{-/-}$), as determined by PCR sequencing via the tail DNA template and two forward primers (OuterF: 5'-tcacaacctactcaagatccagc-3' and InnerF: 5'-caactgtagggtatagcagtccc-3') and one common reverse primer (5'-ttattgcctccttggtgagttg-3'). A mouse tissue direct PCR kit (YEASEN, Shanghai, China) was used to amplify *Ttc22.* Both male and female *Ttc22$^{+/+}$* and *Ttc22$^{-/-}$* mice were maintained as seed mice in an animal facility under SPF conditions at the Peking University Cancer Hospital & Institute. The seed mice were hybridized to generate experimental mouse stocks with different *Ttc22* genotypes. The body weight of each mouse was measured once a week from 4–31 weeks of age. The animal experiments were approved by the Biomedical Ethical Committee of Peking University Cancer Hospital & Institute.

High-fat diet experiment

Four-week-old female C57BL/6N mice were fed a high-fat diet (4.73 kcal/g，H10045, Huafukang Bio. Co., Beijing, China). Additionally, a control group of four-week-old female mice was fed a regular diet. The mice were housed in a temperature-controlled room with a 12-hr light/dark cycle. All the mice had free access to food and water [13].

Fertility test

Mating tests were carried out with 6–8-week-old mice. One male Ttc22$^{+/+}$ mouse was mated with 1 Ttc22$^{+/+}$ female mouse as the control group, and 1 male Ttc22$^{-/-}$ mouse was mated with 1 Ttc22$^{-/-}$ female mouse as the Ttc22-KO group for at least 3 months. At least three mates were included in each group. The date and number of pups were recorded at birth in each cage.

Induction of COAD by AOM/DSS

Five male mice aged 6–8 weeks were included in each group within the same cage. The mice in the three groups, *Ttc22*$^{+/+}$, *Ttc22*$^{+/-}$, and *Ttc22*$^{-/-}$ were intraperitoneally administered 10 mg/kg b.w. azoxymethane (AOM) solution (1 mg/mL, freshly prepared with saline; Cat. #A5486, Sigma − Aldrich; Merck KGaA, Darmstadt, Germany) on the 1st experimental day. Beginning on the 2nd experimental day, the mice were administered dextran sodium sulfate (DSS; Cat. #24867; Sigma) in the drinking water containing 2.5% DSS (2.5 g/100 mL) for 7 continuous days. The mice were subsequently allowed to recover for three weeks (drinking water not containing DSS). The combined AOM/DSS treatment cycle was repeated two more times [14]. From the 24th to 27th experimental weeks, the mice were sacrificed, the colon was collected, and the length of the colon was measured. The intestine was opened, and the colon mucosa was carefully examined and photodocumented. Evidence of gross pathologic changes was recorded (e.g., ulceration). Then, the colon tissues were fixed with 10% formalin, rolled Swiss, embedded in paraffin, and sectioned for pathological analysis. Some of the colon tissues were snapped and frozen for RNA extraction or homogenized in RIPA buffer for protein extraction. The roll was further fixed in 10% buffered formalin overnight at room temperature and used to prepare tissue sections (4 μm). Pathology was performed by senior pathologists.

Quantitative RT−PCR (qRT−PCR)

Total RNA was extracted from fresh tissue samples via TRIzol Reagent (ZYMO Research, Beijing, China) according to the manufacturer's instructions within 5 min of euthanization. First-strand cDNA was generated via a TransScript First-Strand cDNA Synthesis Kit (TransGen Biotech, Beijing, China). Exon−exon qRT−PCR was performed via FastStart Universal SYBR Green Master Mix (Roche, Mannheim, Germany). Mouse *Gapdh* mRNA (forward primer: 5'-ggttgtctcctgcgacttca-3' and reverse primer: 5'-tggtccagggtttcttactcc-3') was used as an internal control for *Ttc22* mRNA (forward primer: 5'-gatatgcgcatggcttcgac-3' and reverse primer: 5'-tgccatggtgaagtaccagc-3'). The relative *Ttc22* mRNA level was calculated via the typical $2^{-\Delta\Delta CT}$ method.

RNA sequencing

Total RNA was extracted from the colon tissues of three male mice with different *Ttc22* genotypes via an RNA extraction kit. RNA sequencing was performed by Cloud-Seq Biotech (Shanghai, China). Briefly, total RNA was removed from the rRNAs with an NEBNext rRNA depletion kit (Biolabs, Inc., Massachusetts, USA),

which was subsequently used to construct RNA libraries via the NEBNext Ultra II directional RNA library

prep kit (Biolabs). Library sequencing was performed on an Illumina NovaSeq 6000 instrument with 150

bp paired-end reads. Paired-end reads were harvested. The high-quality clean reads were aligned to the

reference genome with HISAT2 software (v2.0.4), guided by the Ensembl gtf gene annotation file, and the

gene level FPKM was obtained as the expression profile of the mRNAs. Fold changes and p values were

calculated on the basis of the FPKM values, and the differentially expressed mRNAs were identified on the

basis of │log(FC)│>=1.0, p value<= 0.05). GO and pathway enrichment analyses were performed on the

differentially expressed mRNAs. The raw data were deposited under the Gene Expression Omnibus

accession number GSE275363.

m6A dot blot analysis

m6A dot blot analysis was performed as we previously reported [12].

Quantification of the m6A abundance via m6A-specific ELISA

The total RNA m6A level was quantified with an EpiQuik m6A RNA Methylation Quantification Kit (P-9005;

Epigentek Group, NY, USA) according to the manufacturer's instructions [15,16].

LC−MS/MS measurement of the m6A/rA ratio

Liquid-MS/MS quantification of m6A was performed by Cloudseq Biotech, Inc. (Shanghai, China), as

described previously [12].

Full-length TTC22, ΔTPR1, and ΔTPR4 expression vectors

The pCMV6-Entry-Myc/Flag full-length TTC22 and ΔTPR1 expression vectors were the same as those

previously reported [12]. The ΔTPR3 vector not containing 190-246 aa (equal to the mouse Ttc22 TPR4

domain encoding sequence) was derived from the full-length TTC22 vector. These vectors were used to

transiently transfect human colon cancer HCT116 cells (ATCC; Manassas, VA, USA).

Statistical analysis

Statistical analysis was carried out via SPSS 22. 0 software (SPSS Inc., Chicago IL, USA). The data are

presented as the means ± SDs of at least three independent experiments or as the medians (25th–75th

percentiles). Statistical analysis methods included Student's t test, one-way analysis of variance (ANOVA),

and two-way ANOVA. The exact statistical methods are described in detail for each experiment in the

Results section. All tests were two-sided. A p value <0.05 was considered to indicate statistical

significance.

RESULTS

Characterization of *Ttc22* genotypes

To evaluate the effects of the *Ttc22* gene on the development of mice and COAD, a commercially available

heterozygous Ttc22-KO C57BL/6N mouse model was used to generate mice with various *Ttc22* genotypes,

including homozygous and heterozygous Ttc22-KO ($Ttc22^{-/-}$ and $Ttc22^{+/-}$) and its wild-type control ($Ttc22^{+/+}$)

via serial hybridization, as described in the Methods section. The sequencing results indicated that a 1610

bp fragment, including *Ttc22* exons 2 and 3, which encode the Ttc22 TPR4 domain (58 aa), was deleted via

CRISPR-Cas9 and two gRNAs (Figure 1A and 1B). RNA sequencing confirmed the sequencing results (Figure

1C).

No morphologic alterations in heterozygous and homozygous Ttc22-KO mice

Compared with wild-type C57BL/6N mice, those with different *Ttc22* knockout states did not differ in

development, growth, appearance (Figure 2), body weight (Figure 3A and 3B), or fertility (Figure 4).

Considering that *TTC22* is mostly downregulated in the fetuses of obese women and that

low-frequency variants of TTC22 are associated with LDL cholesterol in the human body [1,3], we further

compared the body weights of female *Ttc22* mice fed a high-fat diet. However, no differences in the body

weights of C57BL/6N mice in different *Ttc22* genotype groups were observed (Figure 3C).

No spontaneous tumors were observed in these mice up to 40 weeks of age. As expected,

spontaneous tumors, including sebaceous carcinoma and basal carcinoma, occurred in some old mice.

However, no differences in the incidence of spontaneous tumors were detected between these *Ttc22*

genotype groups (Table S1).

No differences in AOM/DSS-induced COAD between $Ttc22^{+/+}$, $Ttc22^{+/-}$, and $Ttc22^{-/-}$ mice

Because *TTC22* is involved in COAD metastasis [5,6], we wondered whether Ttc22-KO might affect the

biological processes of the colon and the development of COAD. We performed RNA sequencing on total

RNA extracted from the colons of 58-week-old male mice. The GO and pathway analysis results revealed

that Ttc22-KO mainly upregulated genes related to microorganism infection (Figure 5), supporting the

hypothesis that Ttc22 may contribute to host defensive responses against infection.

We further studied whether Ttc22-KO affects the development of colorectal tumors induced by three

cycles of AOM/DSS treatment in male mice. By the end of the 24th or 27th experimental week,

well-differentiated or moderately differentiated adenocarcinomas were induced in the colons of all

AOM/DSS-treated mice but not in those of the 0.9% NaCl control mice (Figure 6). No differences in the

incidence or severity of colorectal tumors were observed between these mice with different *Ttc22*

genotypes.

The TPR4 domain is not essential for TTC22 function

We recently reported that the TPR1 domain, but not the TPR3 or TPR4 domain, is essential for the

TTC22-RPL4 interaction [12]. However, whether the TTC22 TPR3 or TPR4 domain contributes to WTAP

expression and total RNA m6A levels has not been previously reported. We studied the effect of the loss

of the human TTC22 TPR3 (equal to the mouse Ttc22 TPR4) domain on WTAP expression in human colon

cancer HCT116 cells (Figure 7A). As we previously reported [12], deletion of the TPR1 domain (ΔTPR1)

abolished the ability of TTC22 to increase WTAP and RNA m6A abundance. However, deletion of the TPR3

domain (ΔTPR3) did not abolish the ability of TTC22 to regulate WTAP and RNA m6A modification (Figure

7B and 7C). The results of the ELISA confirmed the results of the dot blot analysis (Figure 7D). These

findings suggest that deletion of the human TTC22 TPR3 (or mouse Ttc22 TPR4) domain did not disrupt

the upregulation of WTAP expression and m6A modification by TTC22. Consistently, the results of liquid

chromatography−mass/mass (LC−MS/MS) analysis revealed no difference in total RNA m6A abundance in

colon tissue samples from the three groups of mice (Figure 7E).

DISCUSSION

Considering that *TTC22* mediates human COAD metastasis [11,12], we studied the influence of *Ttc22*

knockout on the development of COAD in C57BL/6N mice. Unfortunately, no significant differences in the

development or fertility of mice and AOM/DSS-induced COAD were observed between wild-type and

Ttc22-KO C57BL/6N mice, probably because deletion of the Ttc22 TPR4 domain could not disrupt *Ttc22*

functions in the commercially available gene knockout model.

The AOM/DSS-mediated COAD model is an inflammation-related cancer model. The similar

prevalence of COAD induced by AOM/DSS in all of these mice suggests that the doses of AOM and DSS

administered were too high to display the effects of Ttc22-KO on colon tumorigenesis or that the loss of

the Ttc22 TPR4 (equal to the human TTC22 TPR3) domain does not disrupt the biological functions of the

Ttc22 protein in mice. Although we cannot exclude the possibility that biological inactivation of *Ttc22* may

not induce phenotypic alterations, our findings that deletion of human TTC22 TPR3 (ΔTPR3) does not

abolish *WTAP* upregulation via TTC22 and that there are no differences in total RNA m6A levels between

Ttc22-KO and Ttc22-WT mice strongly suggest that the commercial Ttc22-KO model is not an eligible

model that leads to the above negative observations.

Gene knockout animal models are important research tools for validating the biological functions of

target genes. The emergence of commercially available gene knockout mouse models has greatly

promoted research progress in biology, pathogenesis, and drug development. However, most commercial

gene knockout mouse models are not stringently created on the basis of biological inactivation of the

target genes by CRISPR-Cas9. Instead, they are often randomly created using many guide RNAs (gRNAs)

targeted to various genes in the whole genome. Because of the high labor and financial costs of

establishing a target DNA sequence-specific knockout mouse model, biological and medical researchers

like to use commercially available gene knockout mouse models directly. Our experience indicates that

careful evaluation of the eligibility of such a mouse model is essential before the model is used in formal

experiments.

In conclusion, we found that deletion of the Ttc22 TPR4 domain does not disrupt Ttc22 functions and

thus does not induce abnormal phenotypes in mice, including development, growth, or COAD induction

by AOM/DSS. Only eligible gene knockout mouse models should be used in formal experiments.

Abbreviations

COAD, colon adenocarcinoma

Ttc22, tetratricopeptide repeat domain 22

AOM, azoxymethane

DSS, dextran sodium sulfate

Funding This work was supported by grants from the Beijing Hospital Authority Mission Plan

(SML20191101) to DD and from the Science Foundation of Peking University Cancer Hospital (grant

#2022-8) to TW.

Author contributions

243 Conceptualization: DJD, WT, ABY; Funding acquisition: DJD; Investigation: JYL, WT, LKG, ABY, HFY, JZ;

244 Methodology: WT; Project administration: DJD and WT; Visualization: DJD and WT; Writing – original draft:

245 JYL, WT, and DJD; Writing – review & editing: all authors

Competing interest

247 The authors declare that they have no competing interests.

Acknowledgments

249 We acknowledge the support from Beijing Hospital Authority and Peking University Cancer Hospital for

250 this project. We also extend our gratitude to staff from the laboratory animal center at Peking University

251 Cancer Hospital for their care and maintenance of the animals.

REFERENCE

1. D'Andrea, L. TPR proteins: the versatile helix. *Trends in Biochemical Sciences*, 2003. 28(12): 655-662.

2. Hesketh SJ, Mukhopadhyay AG, Nakamura D, Toropova K, Roberts AJ. IFT-A structure reveals

carriages for membrane protein transport into cilia. Cell, 2022. 185(26): 4971-4985.e16.

3. Shamoto N, Narita K, Kubo T, Oda T, Takeda S. CFAP70 Is a Novel Axoneme-Binding Protein That

Localizes at the Base of the Outer Dynein Arm and Regulates Ciliary Motility. Cells, 2018. 7(9): p. 124.

4. Vomastek T, Iwanicki MP, Burack WR, Tiwari D, Kumar D, Parsons JT, Weber MJ, Nandicoori VK.

Extracellular signal-regulated kinase 2 (ERK2) phosphorylation sites and docking domain on the

nuclear pore complex protein Tpr cooperatively regulate ERK2-Tpr interaction. Mol Cell Biol. 2008

Nov;28(22):6954-66.

5. McCloskey A, Ibarra A, Hetzer MW. Tpr regulates the total number of nuclear pore complexes

per cell nucleus. Genes Dev. 2018 Oct 1;32(19-20):1321-1331.

6. Edlow AG, Vora NL, Hui L, Wick HC, Cowan JM, Bianchi DW. Maternal obesity affects fetal

neurodevelopmental and metabolic gene expression: a pilot study. PLoS One, 2014 Feb

18;9(2):e88661.

7. Soerensen, M. *et al.* Epigenome-wide exploratory study of monozygotic twins suggests differentially

methylated regions to associate with hand grip strength. Biogerontology, 2019;20, 627-647.

8. Yang T, Wu C, Wei P, Pan W. Integrating DNA sequencing and transcriptomic data for association

analyses of low-frequency variants and lipid traits. Hum Mol Genet. 2020 Feb 1;29(3):515-526.

9. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C,

Sjöstedt E, Asplund A, et al: Proteomics. Tissue-based map of the human proteome. Science, 2015,

347:1260419.

10. Cerami E, et al. The cBio cancer genomics portal: an open platform for exploring multidimensional

cancer genomics data. Cancer Discov. 2012, 2:401-404.

11. Tian W, Du Y, Ma Y, Zhang B, Gu L, Zhou J, Deng D. miR663a‑TTC22V1 axis inhibits colon cancer

metastasis. Oncol Rep. 2019 Mar;41(3):1718-1728.

12. You A, Tian W, Yuan H, Gu L, Zhou J, Deng D. TTC22 promotes m6A-mediated WTAP expression and

colon   cancer   metastasis   in   an   RPL4   binding-dependent   pattern.   Oncogene,   2022

Aug;41(32):3925-3938.

13. Nakamura Y, Natsume M, Yasuda A, Ishizaka M, Kawahat K, Koga J. Fructooligosaccharides suppress

high-fat diet-induced fat accumulation in C57BL/6J mice. Biofactors, 2017. 43(2): p. 145-151.

14. Angelou A, Andreatos N, Antoniou E, Zacharioudaki A, Theodoropoulos G, Damaskos C, Garmpis N,

Yuan C, Xiao W, Theocharis S, Zografos G, Papalois A, Margonis GA. A Novel Modification of the

AOM/DSS   Model   for   Inducing   Intestinal   Adenomas   in   Mice.   Anticancer   Res.   2018

Jun;38(6):3467-3470.

15. Xu Y, Zhou J, Li L, et al. FTO-mediated autophagy promotes progression of clear cell renal cell

carcinoma via regulating SIK2 mRNA stability. Int J Biol Sci. 2022; 18(15):5943-5962.

16. Ren QW, Xiang MY, Qiao JL, Liu ZJ, Zhang G, Gu LK, Zhou J, Tian W, Deng DJ. TTC7B triggers the

PI4KA-AKT1-RXRA-FTO axis and inhibits colon cancer cell proliferation by increasing RNA methylation.

Int J Biol Sci. 2025; 21(3):1127-1143.

FIGURES AND LEGENDS

Figure 1. Characterization of the genotypes of the *Ttc22* gene in C57BL/6N mice via CRISPR-Cas9. (**A**) Locations of

the deleted fragment (1610 bp), amplicons, gRNAs, and primers matched to the *Ttc22* gene. The conservation status

of *Ttc22* in placental mammals, including rats, rabbits, and humans, is displayed below. (**B**) Characterization of *Ttc22*

genotypes, as determined by PCR and sequencing. The wild-type *Ttc22* fragment is too long (2438 bp) to be

amplified via regular PCR via the OuterF and OuterR primer sets. (**C**) IGV view of *Ttc22* transcripts in RNA samples

from the colons of three male mice with different *Ttc22* genotypes, including *Ttc22* wild-type (*Ttc22*$^{+/+}$),

heterozygously deleted (*Ttc22*$^{-/+}$), and homozygously deleted (*Ttc22*$^{-/-}$) strains, as determined by RNA sequencing.

The exon 2&3 region, which is shared by both *Ttc22* variants 1 and 2 and is knocked out by CRISPR-Cas9, is

highlighted by a red rectangle.

307   **Figure 2**. The appearance of C57BL/6N male and female C57BL/6N mice with different *Ttc22* genotypes at

308   approximately 66 and 77 weeks of age

Figure 3. Body weight curves of C57BL/6N mice in different *Ttc22* genotype groups. (**A** and **B**) Body weight curves

for male and female mice fed a regular diet. (**C**) Body weight curves for female mice fed a high-fat diet. The dots and

bars represent the average and standard deviation (SD) of the body weights of the mice; *p<0.05 according to

Student's t test.

317 **Figure 4**. Fertility comparison between *Ttc22*[+/+] and *Ttc22*[-/-] mice.

Figure 5. Enrichment scores for genes upregulated and downregulated by homozygous *Ttc22* knockout in the colon

321   of 57-week-old male mice

Figure 6. Induction of colon tumors by AOM/DSS in male C57BL mice with different *Ttc22* genotypes. (**A**) Overview

of tumors in the prefixed colon; (**B**) Histological images of representative tumors induced by AOM and DSS.

Figure 7. Effect of TTC22 TPR3 deletion on WTAP expression. (**A**) Illustration of various TTC22 variants, including the

full-length TTC22 (FL) and its TPR1- or TPR3-deleted truncates (ΔTPR1 or ΔTPR3). (**B** and **C**) Effect of TTC22 TPR3

deletion on WTAP expression and total RNA m6A modification in HCT116 cells, as determined by Western blotting

and m6A-specific dot blotting. (**D**) Effect of TTC22 TPR3 deletion on total RNA m6A modification in HCT116 cells, as

determined by m6A-specific ELISA. (**E**) The ratio of total m6A to total adenosine in RNA (m6A/rA) in the colon

tissues of mice with various *Ttc22* genotypes, as determined by mass.

 TABLE LIST

    Table 1. Incidence and latency period of spontaneous lymphoma in C57BL/6N mice with different *Ttc22* genotypes

Genotype	Male			Female		
	n	Mice with tumor (%)	Range of tumor latent weeks	n	Mice with tumor (%)	Range of tumor latent weeks
Ttc22$^{+/+}$	17	3 (17.6)	45, 55, 77	12	0	-
Ttc22$^{+/-}$	14	2 (14.3)	57, 60	19	4 (21.1)	57, 60, 65, 75
Ttc22$^{-/-}$	27	2 (7.4)	54, 54	39	6 (15.4)	41, 55, 55,55, 59, 60
(sum)	58	7 (12.1)		60	10 (16.7)	