DYN-1 regulates SPD-2 and PLK-1 localization and mitotic spindle pole organization

Carter Dierlam[1], Stephanie Held[2], Jiya Hastings[2], Livinus Anyanwu[3], Robert H. Newman[2], and

Jyoti Iyer[2]*

[1]Department of Cell Biology and Molecular Genetics, University of Maryland College Park,

College Park, MD 20742, [2]Department of Biology, North Carolina Agricultural and Technical

State University, Greensboro, NC 27411, and [3]Department of Chemistry and Biochemistry,

University of Tulsa, Tulsa, OK 74104.

Number of characters = 19,747

Running title: *DYN-1 regulates SPD-2 and PLK-1*

*To whom correspondence should be addressed:

Jyoti Iyer

Email: jgiyer@ncat.edu; Tel. (336) 285-4573

Keywords: dynamin, PLK-1, SPD-2, mitosis, midbody, centrosome

ABSTRACT

Accurate cytokinesis is essential for genome stability. Although dynamin is best known for its role in vesicular trafficking, its functions during mitosis remain unclear. Here, we identify a mitotic role for the *C. elegans* dynamin homolog DYN-1 in regulating spindle assembly and the localization of the key mitotic regulators SPD-2 and PLK-1. Depletion of DYN-1 results in enlarged metaphase spindle poles and increased centrosomal SPD-2 and PLK-1 levels. These effects likely depend on DYN-1's vesicular trafficking activity. Additionally, a subset of DYN-1–depleted embryos exhibit altered PLK-1 localization at the midbody during late mitosis, correlating with midbody assembly defects. Together, these findings establish DYN-1 as a previously unrecognized regulator of spindle pole organization and early midbody assembly.

SIGNIFICANCE STATEMENT

Dynamin is primarily known for its function in endocytosis, but its involvement in mitosis is not well defined. This study shows that dynamin helps organize mitotic spindle poles, ensures proper centrosome localization of SPD-2 and PLK-1, and supports early steps of midbody assembly in *C. elegans* embryos. These findings reveal a previously unrecognized mitotic role for dynamin and suggest new links between membrane dynamics, spindle organization, and the cytokinetic machinery.

INTRODUCTION

Accurate cell division is essential for proper chromosome segregation, and mitotic errors can lead to aneuploidy—a distinguishing feature of many cancers (Levine and Holland, 2018). Understanding the molecular mechanisms that regulate mitosis is therefore critical. The GTPase dynamin is a large ~100 kDa protein that is best known for its role in the scission of endocytic vesicles (Hinshaw, 2000; Praefcke and McMahon, 2004). *C. elegans* possesses a single classical dynamin encoded by the *dyn-1* gene (Clark *et al*., 1997). Interestingly, some of the earliest studies identified dynamin as a microtubule-binding protein (Paschal *et al*., 1987; Shpetner and Vallee, 1989). However, dynamin's function in microtubule-associated processes such as spindle assembly and midbody organization during mitosis is not well understood. While DYN-1 was shown to localize to the midbody (Thompson *et al*., 2002), and mediate membrane remodeling during the late stages of abscission (König *et al*., 2017), its potential role in early midbody assembly remains unexplored.

The coiled-coil protein SPindle Defective 2 (SPD-2) (the *C. elegans* homolog of human CEP192) is essential for proper centrosome biogenesis and spindle assembly during mitosis (O'Connell *et al*., 2000; Kemp *et al*., 2004; Pelletier *et al*., 2004; Gomez-Ferreria *et al*., 2007, Zhu *et al*., 2008). SPD-2 levels at the centrosomes are regulated by proteins including SAS-7, SPD-5, DHC-1 and AIR-1, among others (Sugioka *et al*., 2017; Kemp *et al*., 2004; Pelletier *et al*., 2004). Proper levels of centrosome-associated SPD-2 are required for proper centriole duplication, pericentriolar material (PCM) organization and spindle assembly (O'Connell *et al*., 2000; Kemp *et al*., 2004; Pelletier *et al*., 2004). SPD-2 directly binds to the kinase Polo-like kinase 1 (PLK-1) to recruit the kinase to the centrosomes (Boxem *et al.,* 2008; Decker *et al.,* 2011).

PLK1 is a mitotic master regulator that is essential for several mitotic processes including

mitotic entry, centrosome dynamics, chromosome congression, chromosome alignment and

cytokinesis (Petronczki *et al.,* 2008; Archambault and Glover, 2009; Zitouni *et al.,* 2014). PLK1

exhibits a dynamic localization pattern at different stages of mitosis. During early mitosis, PLK1

localizes prominently to the centrosomes, and the kinetochores. At anaphase, it re-localizes to the

spindle midzone and at telophase, it becomes enriched at the midbody (Golsteyn *et al.,* 1995; Lee

et al., 1995; Arnaud *et al.,* 1998; Barr *et al.*, 2004; Petronczki *et al.,* 2008; Schmucker and

Sumara, 2014). The timely localization of PLK1 to various cellular components is essential for

its different important roles during mitosis. Importantly, removal of PLK1 from the centrosomes

is required for proper mitotic progression (Kishi *et al.,* 2009). Although the kinase activity of

PLK1 is thought to be required for PLK1 removal from the centrosomes (Kishi *et al.,* 2009), the

mechanism by which PLK1 is removed from the centrosomes remains unclear.

RESULTS AND DISCUSSION

***dyn-1* depletion results in abnormal metaphase spindle pole assembly and increased levels**

of SPD-2

To determine the effect of DYN-1 depletion in *C. elegans* embryos, we knocked down

endogenous DYN-1 using *RNAi* (Supplemental Figure S1, A and B)*.* Interestingly, DYN-1–

depleted embryos (n=20) were noticeably smaller than controls (n=23), exhibiting a reduced

average cytoplasmic area (~906 μm² vs. ~1200 μm²; $p < 0.0001$) (Supplemental Figure S1, C

and D). DYN-1 was previously shown to localize to the mitotic spindle at metaphase, suggesting

a potential role in metaphase spindle assembly (Thompson *et al.,* 2002). To determine the effect

of DYN-1 depletion on metaphase spindle assembly, we examined the effect of DYN-1 depletion

on spindle pole size using mCherry::tubulin as a proxy (Figure 1). Upon measuring spindle pole

area, we determined that DYN-1-depleted embryos (n=58 spindle poles) exhibited larger spindle

poles than control (*smd-1(RNAi)*)-depleted embryos (n=53 spindle poles; p=0.0018) (Figure 1, A

and B). Specifically, the average spindle pole area of control RNAi embryos was 14.6 μm^2 while

that of *dyn-1(RNAi)* embryos was 17.36 μm^2. Thus, our data have identified a novel role for the

endocytic protein DYN-1 in regulating metaphase spindle pole assembly. An important regulator

of spindle pole and centrosome size is the PCM protein SPD-2 (O'Connell *et al.*, 2000; Kemp *et*

al., 2004; Pelletier *et al.*; Decker *et al.,* 2011). Therefore, we asked whether SPD-2 levels at the

centrosomes were altered between control RNAi and *dyn-1(RNAi)* treated embryos. Our analysis

indicated that centrosome-associated SPD-2 levels were significantly increased ~1.7-fold upon

DYN-1 depletion (Figure 1, C and D; n=66 for *dyn-1(RNAi)*; n=60 for controls; p < 0.0001).

DYN-1 depletion increases levels of centrosome-associated PLK-1

Because SPD-2 directly binds to PLK-1 and recruits it to centrosomes (Boxem *et al.,* 2008;

Decker *et al.,* 2011), and SPD-2 levels increase upon DYN-1 depletion (Figure 1, C and D), we

next asked whether PLK-1 also becomes elevated at centrosomes under these conditions. To

determine if DYN-1 depletion affects centrosomal PLK-1, we performed live imaging on a strain

co-expressing mCherry::tubulin and CRISPR-tagged endogenous PLK-1::sfGFP (Martino *et al.,*

2017). We then knocked down endogenous DYN-1 in this strain using *RNAi* and performed live

imaging of control (*smd-1(RNAi)*) and *dyn-1(RNAi)* treated embryos using spinning-disk

confocal microscopy. Consistent with our hypothesis, our analysis revealed that PLK-1 levels at

the centrosomes were significantly elevated upon DYN-1 depletion (Figure 2, A and B).

Specifically, we found that PLK-1 intensity was ~2.5-fold higher in *dyn-1(RNAi)* embryos

(n=52) than controls (n=58; p < 0.0001) (Figure 2, A and B).

The *dyn-1(ky51)* GTPase mutant, which perturbs vesicular trafficking (Clark *et al.*, 1997;

Grant and Hirsch, 1999), also exhibited increased centrosomal PLK-1 (Figure 2). Specifically,

PLK-1 levels were on average elevated ~1.85-fold at *dyn-1(ky51)* centrosomes (n=46

centrosomes) as compared with control centrosomes (n=44 centrosomes; $p<0.0001$) (Figure 2, C

and D). These data indicate that the vesicular trafficking function of DYN-1 is required for its

regulation of centrosome-associated PLK-1 levels. Furthermore, these data also confirm the

specificity of our *dyn-1(RNAi)* phenotype.

To determine if total cellular levels of PLK-1 are altered between control and *dyn-*

1(ky51) embryos, we next measured PLK-1::sfGFP fluorescence intensity within the anterior and

posterior cytoplasm of metaphase embryos (Supplemental Figure S2). We reasoned that if total

PLK-1 levels are increased upon impairing DYN-1 function, we should see a proportional

increase in cytoplasmic PLK-1 fluorescence in the *dyn-1(ky51)* embryos. Our analysis revealed a

modest ~7% increase in PLK-1 levels in the anterior and posterior cytoplasm of *dyn-1(ky51)*

embryos as compared with controls ($p=0.0234$ (anterior), $p=0.0318$ (posterior)) (Supplemental

Figure S2). Given that PLK-1 levels at metaphase centrosomes increase by ~1.85-fold in *dyn-*

1(ky51) embryos (Figure 2, C and D), this modest ~1.07-fold increase in cytoplasmic PLK-1 is

unlikely to fully explain the elevated centrosome-associated PLK-1 levels observed in these

mutants.

PLK-1 persists at the centrosomes following DYN-1 depletion in a subset of embryos

during the later stages of mitosis

PLK-1 exhibits a spatially and temporally regulated localization pattern during different stages of

mitosis (Chase *et al.,* 2000; Budirahardja and Gönczy, 2008; Martino *et al.,* 2017). It transitions

from its location at the centrosomes and the kinetochores during early mitosis, to the spindle

midzone and the midbody during late mitosis (Chase *et al.,* 2000; Budirahardja and Gönczy,

2008; Martino *et al.,* 2017). While performing live imaging, we noticed several embryos where

PLK-1 appeared to persist at the centrosomes (specifically at the PCM) during the later stages of

mitosis (Figure 3A). To quantify this phenotype, we analyzed PLK-1::sfGFP fluorescence

intensity at anterior centrosomes at different cell cycle stages in both control RNAi and *dyn-*

1(RNAi) treated embryos (Figure 3; Supplemental Figure S3). Note that we measured PLK-

1::sfGFP at anterior centrosomes only because cytoplasmic background is more uniform there; in

contrast, the posterior cytoplasm contains punctate, uneven PLK-1 that makes measurements

unreliable. Our quantitative analysis revealed that there was no PLK-1 detectable at any of the

control centrosomes analyzed at 35 seconds post early telophase (n=29 centrosomes) (Figure 3,

A and B; Supplemental Figure S3). In contrast, in 35% of the centrosomes from *dyn-1(RNAi)*

treated embryos (9 out of 26 centrosomes), PLK-1 was still detectable at the centrosomes at this

stage (p=0.0005) (Figure 3, A and B; Supplemental Figure S3).

We also noted a statistically significant increase in centrosome-associated PLK-1::sfGFP

upon *dyn-1(RNAi)* (n=26 centrosomes) as compared with controls (n=29 centrosomes) via an

unpaired two-tailed t-test at early anaphase (p<0.0001), late anaphase (p=0.0004) as well as in

early telophase (p=0.0094) (Supplemental Figure S3). These data indicate that a pool of PLK-1

tends to persist at the centrosomes of DYN-1-depleted embryos during the later stages of mitosis.

DYN-1-depleted embryos exhibit abnormal midbody formation

Since DYN-1 has a known role in cytokinesis and localizes to the midbody (Thompson *et al.,*

2002), we analyzed midbody formation in DYN-1-depleted embryos. Intercellular bridge (IB)

assembly, assessed using mCherry::tubulin fluorescence, served as a proxy for midbody

formation (Hu *et al.*, 2012; Green *et al.,* 2013; König *et al.*, 2017). We noted that all of the

control RNAi embryos (n=19 embryos) exhibited a normal tubulin-intense IB during telophase

(Figure 4, A (white arrows) and B). In contrast, 50% of the *dyn-1(RNAi)* embryos (12 out of 24

embryos) did not exhibit any detectable IB during telophase (Figure 4, A (bottom panel) and B).

Interestingly, 46% of the *dyn-1(RNAi)* embryos exhibited a structurally abnormal IB that either

started forming outside the reference time-window for normal IB assembly (normal IB started

forming between 175 to 350 seconds post-metaphase) or exhibited defects in IB assembly

dynamics (Figure 4, A (middle panel) and B). Importantly, the *dyn-1(ky51)* allele also exhibited

mild defects in IB assembly (Figure 4B), demonstrating the specificity of this phenotype. While

previous ultrastructural analysis identified a role for DYN-1 in membrane remodeling during the

late stages of abscission (König et al., 2017), our study identifies a novel requirement for DYN-1

in regulating midbody organization during the initial stages of this process. Together, these data

suggest that DYN-1 function is required for proper midbody assembly.

PLK-1 localization to the midbody is defective in DYN-1-depleted embryos

Given that PLK-1 persists at the centrosomes through late mitosis in a subset of DYN-1-depleted

embryos (Figure 3), we hypothesized that this retention might correlate with abnormal PLK-1

dynamics at the midbody. To investigate this, we analyzed PLK-1 localization using two

complementary approaches: a time-matched analysis and an event-matched analysis. The time-

matched analysis (Figure 5, A and B) examined whether PLK-1 localized to the midbody during

the standard developmental window when IB formation begins (175–350 seconds post-

metaphase). Conversely, the event-matched analysis evaluated PLK-1 recruitment specifically

when an IB eventually formed, regardless of timing.

Our time-matched analysis revealed that PLK-1 failed to localize to the midbody in 83%

of *dyn-1(RNAi)* embryos (n = 24), compared to 100% localization in controls (n = 19; p <

0.0001; Figure 5A). To distinguish whether this reduction was simply due to a failure in

midbody formation, we performed an event-matched analysis restricted to *dyn-1(RNAi)* embryos

where an IB-like structure was present (n = 12). Importantly, PLK-1 localized to these formed

midbodies to varying degrees in all cases: 42% showed relatively normal accumulation, while

58% exhibited fragmented, weak, or near-background signals (Figure 5, A and C; p=0.0003).

Together, these data demonstrate that even when a midbody successfully forms following *dyn-1*

depletion, PLK-1 recruitment is frequently altered rather than entirely absent. Notably, due to

inconsistent background signals resulting from polarity defects in *dyn-1(RNAi)* embryos, we

were unable to reliably quantify PLK-1::sfGFP fluorescence intensity at midbodies, as these

background fluorescence fluctuations often led to inconsistent measurements.

In summary, our studies have identified a new role for the endocytic protein DYN-1 in

regulating metaphase spindle assembly, early midbody assembly and the localization of the

mitotic proteins SPD-2 and PLK-1. Specifically, we find that upon DYN-1 depletion, SPD-2 and

PLK-1 accumulate at the centrosomes in metaphase and this is associated with aberrant

metaphase spindle pole assembly. Furthermore, while PLK-1 is normally removed from the

centrosomes during late mitosis, upon DYN-1 depletion, we find that PLK-1 persists at a subset

of centrosomes during the later stages of mitosis.

Based upon our data, we propose a model whereby DYN-1 regulates the levels of SPD-2

and PLK-1 at metaphase centrosomes, potentially through a vesicular trafficking mechanism.

DYN-1 depletion leads to elevated SPD-2 levels at metaphase centrosomes, and because SPD-2

is a known upstream recruiter of PLK-1 (Boxem *et al.*, 2008; Decker *et al.*, 2011), this increase

in SPD-2 is accompanied by increased PLK-1 accumulation at centrosomes. The elevated levels

of these PCM components are associated with aberrant spindle pole assembly. In late mitosis,

DYN-1 depletion also disrupts the normal relationship between midbody formation and PLK-1

accumulation at the IB: embryos frequently fail to assemble a midbody, and even in cases where

a midbody-like structure does form, PLK-1 enrichment is often fragmented or reduced. While we

cannot yet distinguish whether altered PLK-1 midbody localization reflects defective midbody

assembly, impaired PLK-1 recruitment, or a combination of both, our findings indicate that

DYN-1 is required for coordinating the proper redistribution of PLK-1 between centrosomes and

the midbody.

The defects in midbody assembly were more pronounced in *dyn-1(RNAi)* embryos than in

dyn-1(ky51) mutants. Correspondingly, PLK-1::sfGFP enrichment at metaphase centrosomes

was higher following *dyn-1(RNAi)* (~2.5-fold) compared to *dyn-1(ky51)* (~1.85-fold). These

observations suggest that *dyn-1(ky51)* may be a hypomorphic allele. Alternatively, DYN-1 may

regulate these processes through mechanisms other than its role in vesicular trafficking. Future

studies should investigate these potential roles, including whether DYN-1 physically interacts

with PLK-1 to facilitate its recruitment or turnover. Rescue approaches using hypomorphic *plk-1*

alleles may also help clarify the contribution of PLK-1 dysregulation to causality of the observed

defects, but such experiments are beyond the scope of the present work.

Since our studies also uncovered the regulation of centrosome-associated SPD-2 levels

by DYN-1, an important next step is to determine the specific mechanism of this regulation.

Since SPD-2 is one of the upstream components of the PCM assembly pathway (Kemp *et al.*,

2004), future studies will also be directed to determine if other PCM protein levels at the

centrosomes are also altered upon DYN-1 depletion. Investigating these broader effects will

clarify whether the role of DYN-1 is specific to the SPD-2/PLK-1 axis or if it influences

centrosome composition and mitotic spindle assembly more generally.

MATERIALS AND METHODS

C. elegans **strains growth and maintenance**

All *C. elegans* strains described in this study were grown and maintained using standard

techniques on OP50 seeded Modified Youngren's Only Bacto-peptone (MYOB) agar plates at

20ºC (the strains were shifted to other assay-specific temperatures as needed).

C. elegans **strains and genetic crosses**

The wild-type N2 strain and the OD2425 strain expressing PLK-1::sfGFP (*plk-1(lt18[plk-*

1::sfGFP]::loxp) III) (Martino *et al.,* 2017) were obtained from the *Caenorhabditis Genetics*

Center (CGC). The OC908 strain (*bsSi30[pCW9: unc-119(+) pcdk-11.2::sfgfp::his-58::cdk-11.2*

3' UTR] II; bsIs20[pNP99: unc-119(+) tbb-1p::mCherry::tbb-2::tbb-2 3'-UTR]; bsIs2[pCK5.5:

Ppie-1::gfp::spd-2]) and the OC779 strain (*bsSi15[pKO109: spd-2p-spd-2::mCherry::spd-2 3'-*

utr, unc-119(+)] I; bsSi30[pCW9: unc-119(+) pcdk-11.2::sfgfp::his-58::cdk-11.2 3' utr] II; unc-

119(ed3) III) were obtained from Dr. Kevin O'Connell's laboratory at NIDDK, NIH. The

mCherry::tubulin–expressing strain IYR028 (*bsIs20[pNP99: unc-119(+) tbb-1p::mCherry::tbb-*

2::tbb-2 3'-UTR]) was generated in our laboratory by crossing OC908 hermaphrodites with N2

males and isolating progeny carrying only the mCherry::tubulin transgene. Following this,

mCherry::tubulin expressing males were generated from the IYR028 strain by heat shock at 30ºC

for 6 hours. These mCherry::tubulin males from the IYR028 strain were then crossed with PLK-

1::sfGFP expressing hermaphrodites from the OD2425 strain to generate the IYR038 strain

(*bsIs20[pNP99: unc-119(+) tbb-1p::mCherry::tbb-2::tbb-2 3'-UTR]; plk-1(lt18[plk-*

1::sGFP]::loxP) III) which was homozygous for both mCherry::tubulin and PLK-1::sfGFP. This

strain was validated for homozygosity of the two loci by confocal microscopy. The IYR035

strain (*dyn-1(luv35[ky51]) X*) was generated by CRISPR/Cas9 editing as described previously

(Paix *et al.,* 2015; Paix *et al.,* 2016; Iyer *et al.,* 2018; Smith *et al.*, 2020). This strain was

confirmed by both PCR and restriction digestion with Pst1 followed by agarose gel

electrophoresis as well as by DNA sequencing. The IYR043 strain (*bsIs20[pNP99: unc-119(+)*

tbb-1p::mCherry::tbb-2::tbb-2 3'-UTR]; plk-1(lt18[plk-1::sfGFP]::loxP) III; dyn-

1(luv35[ky51]) X) was homozygous for the *dyn-1(ky51)* mutation, mCherry::tubulin and PLK-

1::sfGFP. The IYR044 strain (*bsIs20[pNP99: unc-119(+) tbb-1p::mCherry::tbb-2::tbb-2 3'-*

UTR]; plk-1(lt18[plk-1::sfGFP]::loxP) III; dyn-1(+) X) was homozygous for mCherry::tubulin,

PLK-1::sfGFP and wild-type for *dyn-1*. The IYR043 and IYR044 strains were both generated by

crossing IYR035 hermaphrodites with IYR038 males (the males were made by heat shock at

30°C for 6 hours). Both these strains were validated by confocal microscopy for homozygosity of

the fluorescent markers (mCherry::tubulin and PLK-1::sfGFP). PCR and Pst1 digestion followed

by agarose gel electrophoresis as well as by DNA sequencing was used to confirm homozygosity

of the *dyn-1(ky51)* mutation (IYR043) and wild-type *dyn-1* (IYR044).

CRISPR/Cas9 editing

CRISPR/Cas9 genome editing utilizing assembled ribonucleoprotein complexes and a linear

DNA repair template (Paix *et al.,* 2015; Paix *et al.,* 2016) was performed to generate the IYR035

strain (*dyn-1(luv35[ky51]) X*) according to the protocols detailed in Iyer *et al.* (2018) and Smith

et al. (2020). The sequences of the editing reagents used to create this strain are as follows:

dyn-1(ky51) crRNA: 5'- TGAATAAGCTGCAAAATAAG -3'

dyn-1(ky51) repair template (silent mutations introduce a Pst1 restriction site): 5'-

AGACTTCTTGCCACGTGGATCAGGAATCGTAACACGTCGTTCATTGATCCTGCAGCT

TATTCAAGATCGCAATGAGTACGCCGAGTTCCTAC -3'

dyn-1(ky51) screening primers:

dyn-1(ky51) forward: 5'- CGTTGATCCCTGTGATCAAT – 3'

dyn-1(ky51) reverse: 5'- TTTCAGATGCTCAAATCATCC – 3'

RNAi by feeding

HT115 bacteria expressing the *dyn-1(RNAi)* construct were purchased from Horizon Discovery

Ltd. (Lafayette, CO) while HT115 bacteria expressing the *smd-1 RNAi* construct (control) were

obtained from Dr. Kevin O'Connell's laboratory (NIDDK, Bethesda, MD). HT115 *E. coli*

transformed with either control (*smd-1(RNAi)*) or *dyn-1(RNAi)* plasmids were grown on MYOB

plates supplemented with 100 µg/ml carbenicillin and 2 mM Isopropyl β- d-1-

thiogalactopyranoside (IPTG) as described previously (Kamath and Ahringer J 2003; Iyer *et al.,*

2022). Briefly, L4-stage *C. elegans* from the IYR038 strain were transferred onto *RNAi* plates

and incubated at 25 ºC for 24 to 36 hours. Gravid adults from these plates were then dissected to

release their embryos and 1-cell stage embryos were analyzed by confocal live imaging.

Conditions for wild-type and *dyn-1(ky51)* worms

Starved plates with the IYR043 (*dyn-1(ky51)*) and IYR044 (wild-type *dyn-1*) strains were

washed with M9 buffer and used to seed large 10 cm MYOB plates. These plates were placed at

26 ºC until L4s appeared. 100 to 125 L4s were picked from each plate and transferred onto new

individual 10 cm MYOB plates at 26 ºC overnight. Embryos from both strains were imaged the

next day using a spinning-disk confocal microscope.

Sequence of *dyn-1(RNAi)* construct

The commercially purchased *dyn-1(RNAi)* construct (Horizon Discovery Inc.) was subjected to

whole plasmid sequencing (Plasmidsaurus). The targeting sequence was found to be as follows:

5'-

ATGTCGTGGCAAAACCAGGGAATGCAGGCGTTGATCCCTGTGATCAATCGTGTCCAG

GACGCCTTCTCCCAGCTGGGCACATCTGTCAGCTTCGAACTTCCACAGATCGCCGTC

GTCGGAGGACAGTCCGCTGGAAAGTCGTCGGTGCTGGAGAATTTTGTCGGAAAAGA

CTTCTTGCCACGTGGATCAGGAATCGTAACACGTCGTCCACTTATTTTGCAGCTTATT

CAAGATCGCAATGAGTACGCCGAGTTCCTACACAAGAAGGGTCATCGCTTTGTGGAT

TTTGATGCAGTTCGGAAAGAGATTGAGGATGAGACTGATCGTGTCACTGGACAGAA

TAAGGGAATCAGTCCACATCCAATCAACTTGCGTGTCTTTTCTCCAAATGTTCTAAAT

CTGACACTCATCGATTTGCCCGGTCTCACAAAAGTGCCCGTCGGAGATCAACCAGCA

GATATTGAGCAACAGATCCGTGACATGATTCTCACATTCATCAACCGTGAGACTTGC

CTCATTCTTGCCGTCACTCCGGCCAACAGCGATCTCGCCACTTCGGATGCGTTGAAA

CTTGCGAAGGAAGTCGATCCACAGGGTCTTCGCACGATTGGAGCCCTCACCAAACTT

GACTTGATGGACGAGGGAACCGATGCTCGCGAGATCCTCGAGAACAAGCTGTTCAC

ACTTCGTCGTGGCTACGTCGGAGTTGTCAATCGTGGGCAGAAGGATATTGTCGGTCG

CAAGGATATTAGAGCTGCTTTGGACGCCGAGAGAAAGTTCTTCATCTCACACCCATC

CTACCGACATATGGCTGATCGGTTGGGAACAAGCTACCTTCAACACACTCTTAATCA

ACAGCTCACCAATCATATCCGTGATACACTGCCAACACTTCGTGATAGTCTTCAAAA

GAAGATGTTTGCTATGGAAAAGGATGTGGCCGAGTACAAGAACTACCAGCCAAATG

ATCCAGGCCGCAAGACCAAGGCTCTTTTGCAAATGGTTACCCAGTTCAATGCTGACA

TTGAGCGCTCCATTGAAGGTTCCTCTGCAAAGCTGGTTTCAGCCAATGAGCTCAGTG

GAGGAGCCCGTATCAATCGGCTTTTCCATGAGCGTTTCCCATTTGAGATTGTTAAAA

TGGAAATTGACGAGAAAGAAATGCGCAAAGAAATCCAGTATGCCATCAGAAACATT

CACGGTATCCGCGTCGGTCTCTTCACTCCGGATATGGCGTTTGAGGCAATTGCCAAA

AAGCAAATCACCCGTCTGAAGGAGCCATCGTTGAAATGCGTTGATCTGGTGGTCAAC

GAGTTGGCTAATGTGATCAGACAGTGCGCTGACACTATGGCTAGATATCCACGTCTT

CGTGACGAGCTGGAAAGAATCGTCGTCTCGCATATGCGTGAACGTGAGCAAATTGC

CAAGCAGCAAATTGGGCTCATTGTTGACTACGAACTCGCTTATATGAACACAAACCA

TGAGGATTTCATTGGATTCAGCAATGCTGAAGCAAAAGCCTCCCAAGGACAATCAG

CGAAGAAGAATCTTGGAAACCAGGTGATCAGAAAGGGCTGGCTCTCACTGAGCAAC

GTATCGTTTGTGCGTGGCTCCAAGGACAATTGGTTTGTGCTCATGTCGGACAGTTTG

AGTTGGTACAAAGATGATGAGGAGAAGGAGAAGAAGTACATGCTCCCATTGGATGG

TGTCAAGCTGAAGGATATTGAGGGTGGATTTATGTCTCGTAACCACAAGTTTGCTCT

GTTCTACCCCGACGGAAAGAACATCTACAAGGATTACAAGCAGCTTGAGTTGGGAT

GCACCAATTTGGACGAAATTGATGCGTGGAAGGCTTCATTCTTGCGTGCTGGTGTCT

ATCCAGAAAAGCAGAAGGCACAGGAAGATGAGTCCCAACAAGAGATGGAGGATAC

CTCGATTGATCCACAACTTGAGAGACAGGTGGAGACAATCCGTAATTTGGTTGATTC

CTACATGAGAATCATTACCAAGACAATTAAGGACCTGGTTCCAAAGGCGGTGATGC

ATTTGATTGTTAACCAGACAGGTGAGTTCATGAAAGATGAACTTTTGGCCCATCTCT

ACCAATGCGGCGACACTGATGCTCTCATGGAGGAATCTCAAATAGAAGCCCAGAAG

CGCGAGGAGATGCTCCGAATGTACCATGCTTGCAAGGAGGCGCTCCGCATTATCTCT

GAAGTCAACATGAGCACCCTTGGCGACCAGCCGCCGCCATTGCCGATGTCTGACTAC

CGCCCACACCCATCTGGACCTTCACCGGTGCCGCGTCCGGCTCCTGCTCCACCAGGC

GGACGTCAGGCCCCAATGCCACCACGCGGAGGACCCGGTGCCCCACCACCACCAGG

CATGAGACCACCACCAGGTGCGCCAGGAGGCGGCGGTGGCATGTACCCACCGTTGA

TTCCAAC- 3'

***dyn-1* RT-PCR**

The protocol by Ly *et al.,* 2015 was used to extract RNA from either 50 or 100 control and *dyn-*

1(RNAi) treated gravid adult worms. One modification to the protocol involved freezing the

worms in the TRIzol reagent (Thermo Fisher Scientific, Catalog #15596018) at –80 °C and

storing them until the day of the RNA extraction. RT-PCR was performed using the Qiagen

OneStep RT-PCR kit (Qiagen Inc., Catalog # 210210) following the manufacturer's instructions.

The primers used for the RT-PCR were as follows:

dyn-1 RT-PCR forward: 5'- TGTCGGAAAAGACTTCTTGC – 3'

dyn-1 RT-PCR reverse: 5'- TGTCAGATTTAGAACATTTGG – 3'

Primers sequences targeting the *act-1* gene (loading control) were obtained from Ly *et al.* (2015).

act1f*:* 5'- ACGCCAACACTGTTCTTTCC – 3'

act1r*:* 5'- GATGATCTTGATCTTCATGGTTGA – 3'

Band intensities for all RT-PCR samples were quantified using the open-source software Fiji

(Schindelin *et al.,* 2012).

***C. elegans* embryo live imaging using a spinning-disk confocal microscope**

For imaging embryos from the IYR038 and OC779 strains depicted in Figure 1:

C. elegans embryos obtained from dissected *smd-1(RNAi)* (control) and *dyn-1(RNAi)* worms

were imaged using an Olympus Yokagawa X1 spinning-disk confocal microscope as described

previously (Bergwell, Smith, *et al.,* 2023) with a few alterations. Timelapse imaging was

performed using the cellSens software (Olympus America Inc.) by taking 1.5 μm Z-stacks every

~75 seconds at a 60X magnification with a 1.42 numerical aperture (NA) objective using a Prime

95B CMOS camera. To enable quantitative analysis, all images were captured at the same laser

intensity [(50% laser intensity for both 561 nm and 488 nm channels for IYR038) and (60% laser

intensity for 561 channel and 35% laser intensity for 488 channel for OC779)] and exposure time

(700 msec exposure time for both 561 nm and 488 nm channels for both strains). All brightness

and contrast adjustments were applied equally for both the control RNAi and *dyn-1(RNAi)*

representative images to enable better image presentation and viewing.

For imaging embryos from the IYR038, IYR043, and IYR044 strains depicted in Figures 2-

5, Supplemental Figure S1, Supplemental Figure S2 and Supplemental Figure S3:

A Nikon Ti2-E inverted microscope (Nikon Instruments Inc.) outfitted with a CREST X-Light

V3 spinning-disk confocal module, a motorized XY stage with encoder feedback, and a 600-μm

Z-piezo drive was used to make time-lapse videos of embryos. Embryos were dissected and

mounted on agarose pads, and coverslips were sealed with Vaseline following previously

described procedures (Bergwell, Smith, *et al.*, 2023). Laser illumination was supplied through a

Celesta solid-state light engine (Lumencor Inc.). Time-lapse image acquisition was performed in

NIS-Elements (Nikon Instruments Inc.), capturing 1.25-μm optical Z-stacks approximately every

35 seconds with a 60× Plan Apo λD oil-immersion objective (NA 1.42; refractive index 1.515)

and an ORCA-Fusion BT sCMOS detector (Hamamatsu Inc.). To ensure consistency in

quantitative measurements, all datasets were collected under identical exposure conditions: 200

msec exposure and 60% laser power at 561 nm and 400 msec exposure and 35% laser power at

488 nm. Maximum-intensity projections of Z-stacks were generated with NIS-Elements (Nikon

Instruments Inc.). For display purposes, brightness and contrast adjustments were applied

uniformly across the entire dataset.

Spindle pole area measurement

To measure spindle pole area, the cellSens software (Olympus America Inc.) was used to obtain

maximum intensity Z-projections of the original .vsi files. These Z-projections were saved in a

.png format with the fixed scaling (left:80 and right:400) for all images and further analyzed

using the Nikon NIS-Elements software (Nikon Instruments Inc.). The calibration on the NIS-

Elements software was set to 1 pixel = 0.18333 µm. The thresholding tool in NIS-Elements was

used to define and compute the spindle pole area (in µm²) (intense mCherry::tubulin

fluorescence demarcated each spindle pole). Only regions with clearly defined spindle poles

were included in the analysis. Embryos in which spindle poles could not be reliably

distinguished were excluded. Brightness adjustments were equally applied to the images

representing spindle pole size in Figure 1A for better presentation and viewing.

SPD-2 metaphase centrosome intensity measurements

For measuring SPD-2 fluorescence intensity at centrosomes, the raw .vsi files acquired using

cellSens software (Olympus America Inc.) were opened in the Fiji software (Schindelin *et al.,*

2012) using the Bio-Formats importer plugin. The global scale was set to 1 pixel = 0.18333 µm.

Maximum-intensity Z-projections were generated and used for analyzing SPD-2 fluorescence

intensity at metaphase centrosomes. Metaphase was defined as the stage just prior to anaphase

onset, when the chromosomes had aligned at the cell center. A region of interest (ROI)

measuring 1033.33 pixels² was drawn around each metaphase centrosome and an adjacent area

of cytoplasm. The mean SPD-2 fluorescence intensity within each ROI (in arbitrary units, A.U.)

was quantified using the measure function in Fiji (Schindelin et al., 2012). Background SPD-2

fluorescence intensity (cytoplasmic ROI) was subtracted from the corresponding centrosomal

ROI value to obtain the final background-corrected SPD-2 centrosomal fluorescence intensity.

To calculate relative SPD-2 fluorescence levels normalized to control RNAi, all background-

corrected centrosomal fluorescence intensity values from both control RNAi and *dyn-1(RNAi)*

were divided by the mean background-corrected SPD-2 fluorescence intensity measured in

control RNAi embryos. Brightness, contrast, and sharpness adjustments were uniformly applied

to all images shown in Figure 1C to improve visualization of SPD-2 centrosome localization.

Embryos where the Z-projections did not capture the entire centrosome or where SPD-2 at the

centrosome could not be clearly identified due to surrounding debris were excluded from our

analysis.

PLK-1 metaphase centrosome intensity measurements

Metaphase was defined as the stage when the chromosomes had aligned at the center of the

metaphase plate. To quantify PLK-1::sfGFP fluorescence at the centrosomes, maximum intensity

Z-projections of metaphase embryos were generated and analyzed using Nikon NIS-Elements

software (Nikon Instruments Inc.). A region of interest (ROI) measuring 11.62 μm² was drawn

around each centrosome and within the cytoplasm of the same embryo. Cytoplasmic PLK-

1::sfGFP intensity was then subtracted from centrosomal PLK-1::sfGFP intensity to calculate the

background-corrected PLK-1::sfGFP centrosome fluorescence intensity, reported in arbitrary

units (A.U.). All background-corrected PLK-1::sfGFP centrosome intensities were divided by

the mean background-corrected PLK-1::sfGFP centrosome intensity of control embryos to

calculate the relative PLK-1 levels at centrosomes. Embryos with inaccurate PLK-1::sfGFP

fluorescence due to debris on top of the embryo affecting maximum intensity Z-projections or

instances where the Z-projections of the embryo did not cover the entire centrosome were

excluded from our analysis.

Analysis of PLK-1 persistence at centrosomes

To analyze PLK-1::sfGFP persistence at centrosomes at different cell cycle stages, maximum

intensity Z-projections of each timelapse sequence were quantified for centrosome-localized

PLK-1::sfGFP using the Nikon NIS-Elements software (Nikon Instruments Inc.). To quantify

centrosomal PLK-1::sfGFP levels across different cell-cycle stages, measurements were

restricted to the anterior centrosome. In the anterior cytoplasm, PLK-1::sfGFP background

fluorescence was relatively uniform, enabling consistent background subtraction and accurate

intensity quantification. By contrast, the posterior cytoplasm frequently exhibited irregular,

punctate PLK-1::sfGFP foci of variable sizes and intensities, which introduced inconsistency in

centrosome fluorescence measurements, especially at later cell cycle stages. For this reason,

posterior centrosomes were excluded from our quantitative analysis. The microtubule asters were

used to identify the position of the centrosomes at each cell cycle stage. A ROI of 11.62 μm^2 was

placed at the center of the asters as well as in its immediate vicinity (background). The PLK-

1::sfGFP fluorescence intensity at each anterior centrosome at every cell cycle stage was

determined by subtracting the background fluorescence intensity from the centrosomal

fluorescence intensity at that stage. Centrosomes with background-corrected PLK-1::sfGFP

fluorescence intensities greater than 5 arbitrary units (A.U.) were classified as "PLK-1 present"

whereas centrosomes with intensity values below this threshold were classified as "PLK-1

absent". The cell cycle stages were classified as follows: metaphase: a diamond-shaped spindle

formed and chromosomes aligned at the center, early anaphase: ~35 to 70 seconds after

metaphase when the chromosomes started separating, late anaphase: ~70 seconds after early

anaphase, early telophase: ~70 seconds after late anaphase, and 35 seconds after early telophase:

~35 seconds after early telophase. In the occasional cases where PCM disassembly prevented

identification of the aster center, the centrosome position from the preceding frame was used as a

reference. Brightness adjustments were applied equally to the representative images in Figure 3A

for better presentation and viewing. Only embryos where the Z-projections captured the entire

centrosome were used for our analysis.

Quantification of midbody abnormalities

Maximum-intensity Z-projections generated using Nikon NIS-Elements (Nikon Instruments Inc.)

were used to analyze midbody defects. Embryos were scored in a randomized and blinded

manner: image files were renamed with randomized identifiers and scored by an independent

individual. The scorer evaluated IB behavior without knowledge of genotype or treatment until

scoring was complete. An intercellular bridge (IB) was defined as present when a continuous,

tubulin-intense connection between the two daughter cells could be clearly visualized. IB

assembly was used as an indicator of midbody formation (Hu *et al.*, 2012; Green *et al.*, 2013;

König *et al.*, 2017).

A normal IB was defined as one that started forming approximately 175–350 seconds

after metaphase onset, remained stable for at least 105 seconds following its formation, and

exhibited the stereotypical IB assembly dynamics. Stereotypical IB dynamics consisted of a

sequence in which the spindle midzone coalesced into a single focused, microtubule-rich bridge-

like IB structure. Following coalescence, the IB typically increased in tubulin intensity, then

gradually decreased in intensity as it regressed. Once coalesced, normal IBs did not separate or

fragment and maintained a continuous tubulin-dense morphology until midbody resolution.

An abnormal IB was defined as one that failed to exhibit this stereotypical progression.

Abnormal IBs included bridges that did not start forming within the 175–350 second window,

did not coalesce into a single focused structure, showed fluctuating tubulin intensity, or came

apart after coalescence. An embryo was scored as lacking an IB when no IB could be detected at

any point during the remainder of the cell cycle and through the onset of centrosome separation

in the next cell cycle.

Embryos where the entire IB was not captured in the maximum intensity Z-projections

were excluded from our analysis. For better visualization and presentation, brightness and

contrast setting were applied uniformly and equally to all the representative images in Figure 4A.

Quantification of PLK-1 localization at the midbody

PLK-1 localization at the midbody was quantified using a blinded, randomized analysis using

maximum intensity Z-projections that were generated with the Nikon NIS-Elements software

(Nikon Instruments Inc.). All image files were renamed with random alphanumeric identifiers by

an independent individual who was not involved in the scoring. The scorer analyzed the

randomized dataset without knowledge of genotype, treatment condition, or imaging sequence.

After all scoring was completed and recorded, the mapping key was revealed to assign quantified

values back to experimental groups.

An IB/midbody was defined as present when a continuous, tubulin-intense connection

between the two daughter cells could be clearly visualized in Z-projections. An IB was

considered to have formed if it remained cohesive for at least three consecutive frames (~105

seconds) after its appearance. Metaphase was defined as the timepoint immediately after nuclear

envelope breakdown when the chromosomes are aligned centrally at the metaphase plate. For

both time-matched and event-matched analyses, a normal PLK-1 localization pattern was defined

as a continuous, dot-like or band-like PLK-1::sfGFP signal above the local background at the IB

center. Abnormal localization was defined as discontinuous, fragmented, or off-center PLK-

1::sfGFP fluorescence, or signal that was indistinguishable from the local background. In the

time-matched analysis, cases in which PLK-1::sfGFP failed to localize to the IB either because

IB formation was delayed or because an IB did not assemble were also scored as abnormal. For

all analyses, PLK-1 localization was scored at the frame in which the IB was fully formed and

just before the tubulin intensity at the IB began to decrease, ensuring that measurements were

taken while the midbody remained structurally relatively intact.

For time-matched analysis, PLK-1::sfGFP fluorescence was assessed at the

developmental window when the IB normally started forming in control embryos (during the

reference window of 175–350 seconds after metaphase onset).

For event-matched analysis, PLK-1::sfGFP fluorescence was assessed whenever a stable

IB formed, even if the IB formed outside of the normal developmental window for bridge

assembly.

Z-projections in which either the IB or midbody-associated PLK-1 were not fully

captured across the Z-stack were excluded from analysis to ensure accurate evaluation of PLK-1

fluorescence at the IB/midbody. Brightness adjustments were applied equally to all

representative images of PLK-1 midbody localization in Figure 5A.

Embryo size measurements

Nikon-NIS Elements (Nikon Instruments Inc.) was used to generate maximum intensity Z-

projections of the raw data and to measure the sizes of metaphase-stage 1-cell *C. elegans*

embryos. To measure embryo size, the Bézier tool in NIS-Elements was used to manually trace

the outline of each embryo, using cytoplasmic PLK-1::sfGFP fluorescence to define the

cytoplasmic embryo boundary. The enclosed area was calculated by the software and reported in

μm^2. Embryos whose boundaries could not be clearly defined by cytoplasmic PLK-1::sfGFP

fluorescence due to embryo orientation were excluded from the analysis.

Cytoplasmic PLK-1 fluorescence measurements

Maximum intensity Z-projections generated by Nikon NIS-Elements (Nikon Instruments Inc.)

were used to measure cytoplasmic PLK-1::sfGFP in the anterior and posterior cytoplasm of wild-

type and *dyn-1(ky51)* embryos. For these measurements, the Z-projections were opened in NIS-

Elements and a large ROI of 128.97 μm^2 was placed within the anterior and posterior cytoplasm

of metaphase-stage embryos away from the spindle apparatus. The mean PLK-1::sfGFP

fluorescence intensities within these ROIs were determined using the NIS-Elements software and

reported in arbitrary units (A.U.). Measurements in which the ROI could not fit within the

embryo boundaries due to smaller embryo size were excluded from our analysis.

Statistical analysis of *C. elegans* studies

Graphpad prism 10.5 and 10.6.1 (GraphPad Software, Inc., San Diego, CA) were used to

perform statistical analyses of all the data. For all the data, error bars represent the standard

deviation (s.d.) above and below the mean and the middle lines represent the mean. *p*-values

were rounded to two decimal places and considered significant if *p*<0.05. A two-tailed unpaired

t-test was used to analyze: spindle pole area (Figure 1B), SPD-2 fluorescence intensity (Figure

1D), PLK-1::sfGFP centrosome fluorescence intensities at different cell cycle stages

(Supplemental Figure S3), *dyn-1* knockdown by RT-PCR (Supplemental Figure S1B), embryo

size (cytoplasmic area) (Supplemental Figure S1D), and cytoplasmic PLK-sfGFP intensity

(Supplemental Figure S2). A Fisher's exact test was used to analyze the PLK-1::sfGFP

persistence at the centrosomes (Figure 3B), and defects in PLK-1::sfGFP localization to the

midbody (Figure 5, B and C).

ACKNOWLEDGMENTS

This work was supported by NIH/NIGMS grants (5P20GM103636-09 and 7R15GM152965-02)

and start-up funding from North Carolina A&T State University to J.I. Additional support was

provided by NIH/NIGMS grant 1R35GM153737 and NCI grant 1R01CA273095 to R.N. We

would like to acknowledge Jasmyn Grant, Journey Reddick-Umoja, Erin Haastrup, Peter Nunan,

and Mariah Moreland for technical assistance. OC908 and OC779 strains were provided by Dr.

Kevin O'Connell (NIDDK, NIH). Some of the strains used in this study were provided by the

CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

The authors declare no competing interests.

REFERENCES

Archambault V, Glover DM (2009). Polo-like kinases: conservation and divergence in their

functions and regulation. **Nat Rev Mol Cell Biol** *10*, 265–275.

Arnaud L, Pines J, Nigg EA (1998). GFP tagging reveals human Polo-like kinase 1 at the

kinetochore/centromere region of mitotic chromosomes. **Chromosoma** *107*, 424–429.

Barr FA, Silljé HHW, Nigg EA (2004). Polo-like kinases and the orchestration of cell division.

Nat Rev Mol Cell Biol *5*, 429–441.

Bergwell M, Smith A, Smith E, Dierlam C, Duran R, Haastrup E, Napier-Jameson R, Seidel R,

Potter W, Norris A, et al. (2023). A primary microcephaly-associated *sas-6* mutation perturbs

centrosome duplication, dendrite morphogenesis, and ciliogenesis in *Caenorhabditis elegans*.

Genetics *224*, iyad105.

Boxem M, Maliga Z, Klitgord N, Li N, Lemmens I, Mana M, De Lichtervelde L, Mul JD, Van

De Peut D, Devos M, et al. (2008). A protein domain-based interactome network for *C. elegans*

early embryogenesis. **Cell** *134*, 534–545.

Budirahardja Y, Gönczy P (2008). PLK-1 asymmetry contributes to asynchronous cell division

of *Caenorhabditis elegans* embryos. **Development** *135*, 1303–1313.

Chase D, Serafinas C, Ashcroft N, Kosinski M, Longo D, Ferris DK, Golden A (2000). The

polo-like kinase PLK-1 is required for nuclear envelope breakdown and the completion of

meiosis in *Caenorhabditis elegans*. **Genesis** *26*, 26–41.

Clark SG, Shurland DL, Meyerowitz EM, Bargmann CI, Van der Bliek AM (1997). A dynamin

GTPase mutation causes a rapid and reversible temperature-inducible locomotion defect in

Caenorhabditis elegans. **Proc Natl Acad Sci USA** *94*, 10438–10443.

Decker M, Jaensch S, Pozniakovsky A, Zinke A, O'Connell KF, Zachariae W, Myers E, and

Hyman AA (2011). Limiting amounts of centrosome material set centrosome size in *C. elegans*

embryos. **Curr Biol** *21*, 1259–1267.

Golsteyn RM, Mundt KE, Fry AM, Nigg EA (1995). Cell cycle regulation of the activity and

subcellular localization of Plk1, a human protein kinase implicated in mitotic spindle function. **J**

Cell Biol *129*, 1617–1628.

Gomez-Ferreria MA, Rath U, Buster DW, Chanda SK, Caldwell JS, Rines DR, Sharp DJ (2007).

Human Cep192 is required for mitotic centrosome and spindle assembly. **Curr Biol** *17*, 1960–

1966.

Grant B, Hirsh D (1999). Receptor-mediated endocytosis in the *Caenorhabditis elegans* oocyte.

Mol Biol Cell *10*, 4311–4326.

Green, R.A., J.R. Mayers, S. Wang, L. Lewellyn, A. Desai, A. Audhya, and K. Oegema (2013).

The midbody ring scaffolds the abscission machinery in the absence of midbody microtubules. **J**

Cell Biol. *203*, 505–520.

Hinshaw JE (2000). Dynamin and its role in membrane fission. **Annu Rev Cell Dev Biol** *16*,

483–519.

Hu CK, Coughlin M, Mitchison TJ (2012). Midbody assembly and its regulation during

cytokinesis. **Mol Biol Cell** *23*, 1024–1034.

Iyer J, Gentry LK, Bergwell M, Smith A, Guagliardo S, Kropp PA, Sankaralingam P, Liu Y,

Spooner E, Bowerman B, et al. (2022). The chromatin remodeling protein CHD-1 and the EFL-

1/DPL-1 transcription factor cooperatively downregulate CDK-2 to control SAS-6 levels and centriole number. **PLoS Genet** *18*, e1009799.

Iyer J, DeVaul N, Hansen T, Nebenfuehr B (2019). Using microinjection to generate genetically modified *Caenorhabditis elegans* by CRISPR/Cas9 editing. In: *Microinjection: Methods and Protocols*, ed. C Liu and Y Du, New York, NY: Humana Press, 431–457.

Kamath RS, Ahringer J (2003). Genome-wide RNAi screening in *Caenorhabditis elegans*. **Methods** *30*, 313–321.

Kemp CA, Kopish KR, Zipperlen P, Ahringer J, O'Connell KF (2004). Centrosome maturation and duplication in *C. elegans* require the coiled-coil protein SPD-2. **Dev Cell** *6*, 511–523.

Kishi K, Van Vugt MA, Okamoto KI, Hayashi Y, Yaffe MB (2009). Functional dynamics of Polo-like kinase 1 at the centrosome. **Mol Cell Biol** *29*, 3134–3150.

König J, Frankel EB, Audhya A, Müller-Reichert T (2017). Membrane remodeling during embryonic abscission in *Caenorhabditis elegans*. **J Cell Biol** *216*, 1277–1286.

Lee KS, Yuan YL, Kuriyama R, Erikson RL (1995). Plk is an M-phase-specific protein kinase and interacts with a kinesin-like protein, CHO1/MKLP1. **Mol Cell Biol** *15*, 7143–7151.

Levine MS, Holland AJ (2018). Impact of mitotic errors on cell proliferation and tumorigenesis. **Genes Dev** *32*, 620–638.

Ly K, Reid SJ, Snell RG (2015). Rapid RNA analysis of individual *Caenorhabditis elegans*. **MethodsX** *2*, 59–63.

Martino L, Morchoisne Bolhy S, Cheerambathur DK, Van Hove L, Dumont J, Joly N, Desai A, Doye V, Pintard L (2017). Channel nucleoporins recruit PLK-1 to nuclear pore complexes to direct nuclear envelope breakdown in *C. elegans*. **Dev Cell** *43*, 157–171.

O'Connell KF, Maxwell KN, White JG (2000). The *spd-2* gene is required for axis polarization

in the *C. elegans* zygote. **Dev Biol** *222*, 55–70.

Paix A, Folkmann A, Rasoloson D, Seydoux G (2015). High efficiency, homology-directed

genome editing in Caenorhabditis elegans using CRISPR-Cas9 ribonucleoprotein complexes.

Genetics *201*, 47–54.

Paix A, Schmidt H, Seydoux G (2016). Cas9-assisted recombineering in *C. elegans*: genome

editing using in vivo assembly of linear DNAs. **Nucleic Acids Res** *44*, e128.

Paschal BM, Shpetner HS, Vallee RB (1987). MAP 1C is a microtubule-activated ATPase which

translocates microtubules in vitro and has dynein-like properties. **J Cell Biol** *105*, 1273–1282.

Pelletier L, Özlü N, Hannak E, Cowan C, Habermann B, Ruer M, Müller-Reichert T, Hyman AA

(2004). The *Caenorhabditis elegans* centrosomal protein SPD-2 is required for both

pericentriolar material recruitment and centriole duplication. **Curr Biol** *14*, 863–873.

Petronczki M, Lénárt P, Peters JM (2008). Polo on the rise—from mitotic entry to cytokinesis

with Plk1. **Dev Cell** *14*, 646–659.

Praefcke GJA, McMahon HT (2004). The dynamin superfamily: universal membrane tubulation

and fission molecules?. **Nat Rev Mol Cell Biol** *5*, 133–147.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, et al. (2012). Fiji: an open-source platform for

biological-image analysis. **Nat Methods** *9*, 676–682.

Schmucker S, Sumara I (2014). Molecular dynamics of PLK1 during mitosis. **Mol Cell Oncol** 1,

e954507.

Shpetner HS, Vallee RB (1989). Identification of dynamin, a novel mechanochemical enzyme

that mediates interactions between microtubules. **Cell** *59*, 421–432.

Smith A, Bergwell M, Smith E, Mathew D, Iyer J (2020). CRISPR/Cas9 editing of the *C.*

elegans rbm-3.2 gene using the *dpy-10* Co-CRISPR screening marker and assembled

ribonucleoprotein complexes. **J Vis Exp** *166*, e62001.

Sugioka K, Hamill DR, Lowry JB, McNeely ME, Enrick M, Richter AC, Kiebler LE, Priess JR,

Bowerman B (2017). Centriolar SAS-7 acts upstream of SPD-2 to regulate centriole assembly

and pericentriolar material formation. **eLife** *6*, e20353.

Thompson HM, Skop AR, Euteneuer U, Meyer BJ, McNiven MA (2002). The large GTPase

dynamin associates with the spindle midzone and is required for cytokinesis. **Curr Biol** *12*,

2111–2117.

Zhu F, Lawo S, Bird A, Pinchev D, Ralph A, Richter C, Müller-Reichert T, Kittler R, Hyman

AA, Pelletier L (2008). The mammalian SPD-2 ortholog Cep192 regulates centrosome

biogenesis. **Curr Biol** *18*, 136–141.

Zitouni S, Nabais C, Jana SC, Guerrero A, Bettencourt-Dias M (2014). Polo-like kinases:

structural variations lead to multiple functions. **Nat Rev Mol Cell Biol** *15*, 433–452.

FIGURE LEGENDS

FIGURE 1. DYN-1 depletion increases spindle pole size and elevates SPD-2 at centrosomes. (A)

Spindle pole morphology in embryos expressing mCherry::tubulin, showing enlarged spindle

poles upon *dyn-1(RNAi)*. White arrows: spindle poles. (B) Quantification of spindle pole area

using mCherry::tubulin fluorescence. Each point = single pole; n = spindle poles analyzed. Error

bars = s.d.; middle bar = mean. (C) SPD-2 localization at metaphase centrosomes in control and

dyn-1(RNAi) embryos. Dotted outline demarcates the embryo boundary; white arrows:

centrosome-localized SPD-2 (D) Relative centrosomal SPD-2 intensity normalized to control

RNAi. n = centrosomes. Error bars = s.d.; middle bar = mean. (B, D): unpaired two-tailed t-test;

Experimental replicates: 4 for all assays. All fluorescence images include a 10 μm scale bar.

FIGURE 2. DYN-1 depletion increases centrosomal PLK-1 at metaphase. (A) PLK-1::sfGFP

localization at metaphase centrosomes in control RNAi vs *dyn-1(RNAi)* embryos. White arrows:

centrosome-localized PLK-1::sfGFP. (B) Quantification of PLK-1::sfGFP centrosomal

fluorescence at metaphase. n = centrosomes. Error bars = s.d.; middle bar = mean. (C) PLK-

1::sfGFP localization in wild-type (wt) *dyn-1* and *dyn-1(ky51)*. White arrows: centrosome-

localized PLK-1::sfGFP. (D) Relative centrosomal PLK-1 intensity in wild-type (wt) *dyn-1* and

dyn-1(ky51). n = centrosomes. Error bars = s.d.; middle bar = mean. (B, D): unpaired two-tailed

t-test; Experimental replicates: 3 for all assays. All fluorescence images include a 10 μm scale

bar.

FIGURE 3. PLK-1 persists abnormally at centrosomes during mitotic progression in *dyn-*

1(RNAi) embryos. (A) Time-lapse analysis of different mitotic stages in control RNAi and *dyn-*

1(RNAi) embryos. White arrows: anterior centrosome-localized PLK-1::sfGFP. (B) Percentage

of centrosomes retaining PLK-1 at each stage. Fisher's Exact Test; n = centrosomes analyzed.

Experimental replicates: 3 for all assays. All fluorescence images include a 10 μm scale bar.

FIGURE 4. DYN-1 depletion disrupts Intercellular Bridge (IB) assembly. (A) Time-lapse images

of mCherry::tubulin showing normal IB formation in control RNAi embryos (top panel) and

abnormal (middle panel) or missing (bottom panel) IBs in *dyn-1(RNAi)* embryos. White arrows:

IB (B) Classification of IB morphology (normal, abnormal, absent). n = embryos analyzed.

Experimental replicates: 3 for all assays. All fluorescence images include a 10 µm scale bar.

FIGURE 5. PLK-1 localization at the midbody is defective in *dyn-1(RNAi)* embryos. (A) Time-

matched and event-matched PLK-1::sfGFP localization at the midbody in control RNAi and *dyn-*

1(RNAi) embryos. White arrows: IB; yellow arrows: PLK-1::sfGFP at IB (B) Time-matched

PLK-1 scoring at the midbody. n = embryos; Fisher's Exact Test. (C) Event-matched PLK-1

scoring at the midbody. n = embryos; Fisher's Exact Test. Experimental replicates: 3 for all

assays. All fluorescence images include a 10 µm scale bar.

SUPPLEMENTAL FIGURES

FIGURE S1. *dyn-1(RNAi)* reduces *dyn-1* mRNA levels and decreases embryo size. (A) RT-PCR

validation of *dyn-1* depletion. (B) Relative *dyn-1* mRNA quantification. Error bars = s.d.; middle

bar = mean. (C) Comparison of average embryo sizes in control RNAi and *dyn-1(RNAi)*. (D)

Cytoplasmic area quantification using PLK-1::sfGFP fluorescence as a marker for the

cytoplasmic embryo border. n = embryos. Error bars = s.d. (B, D): unpaired two-tailed t-test;

Experimental replicates: 3 for all assays. All fluorescence images include a 10 µm scale bar.

FIGURE S2. Cytoplasmic PLK-1::sfGFP levels are modestly elevated in *dyn-1(ky51)* embryos as

compared to wild-type. (A) Quantification of anterior and posterior cytoplasmic PLK-1::sfGFP

in wild-type (wt) *dyn-1* and *dyn-1(ky51)* embryos. A.U.= Arbitrary Units; unpaired two-tailed t-

test; n = measurements. Error bars = s.d.; middle bar = mean. Experimental replicates: 3. All

fluorescence images include a 10 µm scale bar.

FIGURE S3. Comparison of centrosomal PLK-1 intensity across cell cycle stages in control

RNAi and *dyn-1(RNAi)* embryos. (A) Centrosomal PLK-1::sfGFP intensity at metaphase, early

and late anaphase, early telophase, and 35 seconds post early telophase. A.U.= Arbitrary Units;

unpaired two-tailed t-test; n = 29 centrosomes (control RNAi); n = 26 centrosomes (*dyn-*

1(RNAi)). Error bars = s.d.; middle bar = mean. Experimental replicates: 3. All fluorescence

images include a 10 μm scale bar.

Figure 1

A
control RNAi   *dyn-1(RNAi)*

Red: mCherry::tubulin

C
control RNAi   *dyn-1(RNAi)*

Red: mCherry::SPD-2
Green: GFP::histone

B

p=0.0018
**

Spindle pole area (μm²) using tubulin fluorescence as a proxy

control RNAi
n=53

dyn-1(RNAi)
n=58

D

Centrosomal SPD-2 fluorescence at metaphase

****

p<0.0001

Relative SPD-2 levels at centrosomes (Normalized to control RNAi)

control RNAi
n=60

dyn-1(RNAi)
n=66

Figure 2

A

| MERGE | PLK-1::sfGFP |

control RNAi

dyn-1 (RNAi)

Red: mCherry::tubulin
Green: PLK-1::sfGFP

C

| MERGE | PLK-1::sfGFP |

wt dyn-1

dyn-1(ky51)

Red: mCherry::tubulin
Green: PLK-1::sfGFP

B

Centrosomal PLK-1 fluorescence at metaphase

p<0.0001
****

Relative PLK-1 levels at centrosomes
(Normalized to control RNAi)

control RNAi
n=58

dyn-1(RNAi)
n=52

D

Centrosomal PLK-1 fluorescence at metaphase

p<0.0001
****

Relative PLK-1 levels at centrosomes
(Normalized to wt dyn-1)

wt dyn-1
n=44

dyn-1(ky51)
n=46

Figure 3

A

| | Metaphase | Early Anaphase | Late Anaphase | Early Telophase | 35 seconds after Early Telophase |

Green: PLK-1::sfGFP

Figure 4

A

	175 seconds after Metaphase	245 seconds after Metaphase	285 seconds after Metaphase	315 seconds after Metaphase	

control RNAi — Normal intercellular bridge

dyn-1 (RNAi) — Abnormal intercellular bridge

No intercellular bridge

Red: mCherry::tubulin

B

Number of embryos

- No intercellular bridge
- Abnormal intercellular bridge
- Normal intercellular bridge

control RNAi: 100%
dyn-1(RNAi): 50%, 46%
wt *dyn-1*
dyn-1(ky51): 14.3%

n=19   n=24   n=21   n=21

Figure 5

B
Time-matched PLK-1 localization at the midbody

p<0.0001
****

0%

83%

Number of embryos

control RNAi   *dyn-1(RNAi)*

n=19   n=24

□ Localized
■ Did not localize

C
Event-matched PLK-1 localization at the midbody

p=0.0003
***

0%

58%

42%

Number of embryos

control RNAi   *dyn-1(RNAi)*

n=19   n=12

□ Normal PLK-1
■ Abnormal PLK-1