Dynamin regulates PLK-1 localization and spindle pole assembly during mitosis

Carter Dierlam[1], Livinus Anyanwu[2§], Stephanie Held[3§], Robert H. Newman[3], and Jyoti Iyer[3]*

[1]Department of Cell Biology and Molecular Genetics, University of Maryland College Park,

College Park, MD 20742,[2]Department of Chemistry and Biochemistry, University of Tulsa,

Tulsa, OK 74104 and [3]Department of Biology, North Carolina Agricultural and Technical State

University, Greensboro, NC 27411.

[§]Equal Contribution

Number of characters = 19,800

11   Running title: *DYN-1 regulates PLK-1 localization*

13   To whom correspondence should be addressed:

14   Jyoti Iyer

15   Email: jgiyer@ncat.edu; Tel. (336) 285-4573

17   **Keywords**: dynamin, Plk1, mitosis, midbody, cytokinesis

ABSTRACT

Accurate cytokinesis is essential for maintaining genomic integrity. Although the GTPase

dynamin has been well studied for its role in vesicular trafficking, its function during mitosis

remains poorly understood. In this study, we uncover a novel role for the *C. elegans* dynamin

homolog, DYN-1, in regulating mitotic spindle pole assembly and the spatiotemporal

localization of the key mitotic kinase Polo-like kinase 1 (PLK-1). Our studies demonstrate that

the depletion of DYN-1 leads to enlarged metaphase spindle poles and elevated levels of

centrosome-associated PLK-1. Strikingly, PLK-1 fails to re-localize from centrosomes to the

midbody during late mitosis in a subset of DYN-1–depleted embryos, correlating with abnormal

PLK-1 localization at the midbody and defective midbody formation. Importantly, this

phenotype is likely not due to increased total PLK-1 protein levels, as DNM2 (human homolog

of DYN-1) depletion in HeLa cells did not alter total Plk1 abundance. Together, our findings

identify DYN-1 as a new regulator of PLK-1 localization during mitosis and suggest that failure

to remove PLK-1 from centrosomes may underlie cytokinesis defects that are observed upon

DYN-1 depletion.

SIGNIFICANCE STATEMENT

Dynamin is primarily known for its function in endocytosis, but its involvement in mitosis is not

well defined. This study shows that dynamin helps organize the mitotic spindle and ensures

proper localization of the master mitotic kinase PLK-1 in *C. elegans* embryos. These findings

reveal a previously unrecognized role for dynamin in mitosis and suggest new links between

membrane dynamics and the cell division machinery.

INTRODUCTION

Proper cytokinesis is a prerequisite for the accurate partitioning of the chromosomes to yield two genetically identical daughter cells. Cytokinesis defects can lead to aneuploidy, which is a characteristic feature of cancer cells. Therefore, it is essential to understand the molecular mechanisms governing cytokinesis. The GTPase dynamin is a large ~100 kDa protein that is best known for its role in the scission of endocytic vesicles (Hinshaw JE 2000; Menon *et al.,* 2013; Khurana *et al.,* 2023). The dynamin protein is strongly conserved between humans and the multicellular eukaryotic nematode *Caenorhabditis elegans* (*C. elegans*). *C. elegans* have a single classical dynamin isoform called DYN-1. There is over 60% identity at the amino acid level between human DNM2 and *C. elegans* DYN-1 (Clustal Omega). Interestingly, some of the earliest studies identified dynamin as a microtubule-binding protein (Shpetner and Vallee, 1989). However, dynamin's function in microtubule-associated processes is less well understood. Studies in various organisms including *C. elegans* and human cells have revealed that dynamin is required for proper cytokinesis (Thompson *et al.,* 2002; Wienke *et al.,* 1999; Pelissier *et al.,* 2003). However, mechanistic gaps still exist in our understanding of how DNM2 regulates cytokinesis.

Polo-like kinase 1 (Plk1) is a mitotic master regulator that is essential for several mitotic processes including mitotic entry, centrosome dynamics, polarity establishment, chromosome congression, chromosome alignment and cytokinesis (Petronczki *et al.,* 2008; Archambault and Glover, 2009; Zitouni *et al.,* 2014, Nishi *et al.*, 2008). Importantly, Plk1 exhibits a dynamic localization pattern at different stages of mitosis. During early mitosis, Plk1 localizes to the centrosomes and the kinetochores. At anaphase, it re-localizes to the spindle midzone and at telophase, it localizes to the midbody (Golsteyn *et al.,* 1995; Lee *et al.,* 1995; Arnaud *et al.,*

1998; Petronczki *et al.,* 2008; Kishi *et al.,* 2009; Schmucker and Sumara, 2014). The timely

localization of Plk1 to various cellular components is essential for its different important roles

during mitosis. Importantly, like DNM2, Plk1 is also required for proper cytokinesis (Petronczki

et al., 2008; Archambault and Glover, 2009; Zitouni *et al.,* 2014). Plk1 activity is required for

recruitment of the RhoA guanine nucleotide exchange factor (GEF), Ect2, to the spindle midzone

and equatorial cortex through phosphorylation of the centralspindlin component CYK4. Ect2, in

turn, mediates RhoA activation to facilitate contractile ring assembly and cleavage furrow

ingression, thereby ensuring proper cytokinesis (Brennan *et al.,* 2007; Burkard *et al.,* 2007;

Petronczki *et al.,* 2007; Santamaria *et al.,* 2007; Petronczki *et al.,* 2008; Burkard *et al.,* 2009;

Wolfe *et al.,* 2009). Importantly, removal of Plk1 from the centrosomes is required for proper

mitotic progression (Kishi *et al.,* 2009). Although the kinase activity of Plk1 is thought to be

required for Plk1 removal from the centrosomes (Kishi *et al.,* 2009), the mechanism by which

Plk1 is removed from the centrosomes remains unclear.

RESULTS AND DISCUSSION

***dyn-1 RNAi* embryos exhibit cytokinesis defects**

To study the role of DYN-1 in cytokinesis, we depleted endogenous DYN-1 from *C. elegans*

embryos expressing mCherry-SPD-2 (marks centrosomes) and GFP-histone (marks DNA) using

RNAi. Importantly, our *RNAi* procedure successfully reproduced the cytokinesis failure

phenotype that was previously reported by Thompson *et al*. upon *dyn-1 RNAi* in *C. elegans*

(Thompson *et al.,* 2002)*.* While all of the control (S-adenosyl-methionine decarboxylase (*smd-*

1)) *RNAi* treated embryos proceeded normally through cell division (n=28 embryos), 45% of the

dyn-1 RNAi embryos failed cytokinesis (n=31 embryos) (Fisher's exact test, p<0.0001) **(Figure**

S1). Since our *RNAi* depletion was able to replicate the cytokinesis failure phenotype that was

previously reported upon *dyn-1 RNAi* in *C. elegans* by another group (Thompson *et al.,* 2002),

we concluded that our depletion was successful. Therefore, we proceeded to investigate the

downstream effects of DYN-1 depletion.

***dyn-1* depletion results in abnormal metaphase spindle pole assembly**

Interestingly, DYN-1 was shown to localize to the mitotic spindle at metaphase, suggesting a

potential role in metaphase spindle assembly (Thompson *et al.,* 2002). To determine the effect of

DYN-1 depletion on metaphase spindle assembly, we depleted DYN-1 using *RNAi* and

examined the effect of DYN-1 depletion on metaphase spindle assembly using mCherry-tubulin

as a proxy **(Figure 1)**. Upon measuring spindle pole area (as assessed by mCherry-tubulin

fluorescence intensity), we determined that DYN-1-depleted embryos (n=58 spindle poles)

exhibited a larger spindle pole area than control (*smd-1*) *RNAi* depleted embryos (n=53 spindle

poles; unpaired two-tailed t-test, p=0.0018) **(Figure 1)**. Specifically, the average spindle pole

area of control *RNAi* treated embryos was 14.6 μm^2 while that of *dyn-1 RNAi* treated embryos

was 17.36 μm^2. Thus, our data have identified a novel role for the endocytic protein DYN-1 in

regulating metaphase spindle pole assembly. To explore this role further, we examined the

mechanism by which DYN-1 might regulate spindle assembly. One protein that is an important

regulator of mitotic spindle pole assembly is the kinase PLK-1. Furthermore, PLK-1 exhibits a

partially overlapping localization pattern with DYN-1 during mitosis (Chase *et al.,* 2000;

Budirahardja and Gönczy, 2008; Martino *et al.,* 2017; Thompson *et al.,* 2002). Therefore, we

questioned whether PLK-1 levels at the centrosomes were altered between control and *dyn-1*

RNAi treated embryos.

DYN-1 depletion increases the levels of centrosome-associated PLK-1

To determine whether DYN-1 depletion affects the levels of centrosome-associated PLK-1, we

created a new *C. elegans* strain expressing mCherry-Tubulin and CRISPR-tagged endogenous

PLK-1-sfGFP (Martino *et al.,* 2017) via genetic crossing. We then knocked down endogenous

DYN-1 in this strain using *RNAi* and performed live imaging of control (*smd-1) RNAi* and *dyn-1*

RNAi treated embryos using spinning-disk confocal microscopy. Consistent with our hypothesis,

our analysis revealed that PLK-1 levels at the centrosomes were elevated upon DYN-1 depletion

(Figure 2). Specifically, we found that the average PLK-1 fluorescence intensity in DYN-1-

depleted embryos was 51.02 arbitrary units (n=40 centrosomes). This was about 1.5 times higher

than that of control (*smd-1) RNAi* treated embryos (average PLK-1 fluorescence intensity =

33.13 arbitrary units; n=46 centrosomes; unpaired two-tailed t-test, $p<0.0001$). These data

indicate that DYN-1 normally acts to negatively regulate PLK-1 localization to the centrosomes

through an as-yet unidentified mechanism. Therefore, in the absence of DYN-1, PLK-1 is no

longer restricted from localizing to the centrosomes and consequently, its levels at the

centrosomes increase. These increased levels of centrosome-associated PLK-1 upon DYN-1

depletion **(Figure 2)** are consistent with the enlarged spindle poles that we observed upon DYN-

1 depletion **(Figure 1).** However, it is currently unclear whether DYN-1 regulates PLK-1

localization via vesicular trafficking or through an endocytosis-independent mechanism. Future

studies will be directed towards further dissecting the molecular mechanism by which DYN-1

regulates PLK-1 localization.

PLK-1 persists at the centrosomes following DYN-1 depletion in a subset of embryos

during the later stages of mitosis

PLK-1 exhibits a spatially and temporally regulated localization during different stages of

mitosis (Chase *et al.,* 2000; Budirahardja and Gönczy, 2008; Martino *et al.,* 2017). PLK-1

primarily localizes to the centrosomes and to the kinetochores from prophase to metaphase.

Upon anaphase onset, PLK-1 starts re-localizing to the spindle midzone. Finally, during

telophase, PLK-1 completely disappears from the centrosomes and is recruited to the midbody to

facilitate abscission (Chase *et al.,* 2000; Budirahardja and Gönczy, 2008; Martino *et al.,* 2017).

During the course of our imaging experiments, we noticed several embryos where PLK-1

appeared to persist at the centrosomes (specifically at the pericentriolar material) during

telophase **(Figure 3A)**. To quantify this phenotype, we analyzed PLK-1 localization to the

centrosomes at different cell cycle stages using maximum-intensity Z-projections of the time-

lapse images from both control (*smd-1*) and *dyn-1 RNAi* treated embryos **(Figure 3B)**.

Centrosomes where PLK-1 fluorescence intensity was visually determined to be above that of

the background cytoplasmic PLK-1-sfGFP fluorescence were scored as being positive for PLK-

1. Our analysis revealed that in control embryos at late telophase, there was no PLK-1 detectable

at any of the centrosomes analyzed (n=48 centrosomes) **(Figure 3A and Figure 3B)**. In contrast,

in 15% of the centrosomes from *dyn-1 RNAi* treated embryos (in 7 out of 48 centrosomes), PLK-

1 was still detectable at the centrosomes at late telophase (Fisher's exact test, p=0.0124) **(Figure**

3A and Figure 3B). We also noted a marked increase in the percentage of centrosomes with

PLK-1 during early telophase upon DYN-1 depletion **(Figure 3B)**. Specifically, while only 13%

of the centrosomes (6 out of 48 centrosomes) from control embryos had PLK-1 associated with

them during early telophase, 40% of the centrosomes (19 out of 48 centrosomes) from DYN-1-

depleted embryos still had PLK-1 associated with them at this stage (Fisher's exact test,

p=0.0047) **(Figure 3B)**. These data suggest that a pool of PLK-1 tends to persist at the

centrosomes of DYN-1-depleted embryos during the later stages of mitosis. Since DYN-1

depletion results in an elevation of PLK-1 levels at the centrosomes during early mitosis, it is

likely that these increased levels of PLK-1 make it difficult for PLK-1 to be effectively removed

from the centrosomes during the later stages of mitosis in a subset of *dyn-1 RNAi* embryos.

157 DYN-1-depleted embryos exhibit abnormal midbody formation

Since DYN-1 has a known role in cytokinesis and localizes to the midbody (Thompson *et al.,*

2002), we analyzed midbody formation in DYN-1-depleted embryos using mCherry-tubulin

fluorescence as a proxy for midbody assembly. We noted that all of the control *RNAi* treated

embryos (n=26 embryos) exhibited a normal detectable bridge-like midbody during telophase as

determined by intense tubulin fluorescence **(Figure 4A** (top panel, white arrow) **and Figure 4B)**.

In contrast, 72% of the *dyn-1 RNAi* treated embryos (18 out of 25 embryos) either exhibited a

defective midbody or did not exhibit any detectable midbody during telophase **(Figure 4A**

(bottom panel) **and Figure 4B)** (Fisher's exact test, p<0.0001). In some of these instances, we

observed that in DYN-1-depleted embryos, the midbody formed abnormally after telophase had

already been completed and the embryo had entered the subsequent cell cycle. Together, these

data suggest that DYN-1 function is required for proper midbody assembly.

169 PLK-1 localization to the midbody is defective in embryos depleted of DYN-1

Since our previous data demonstrated that PLK-1 continued to localize to the centrosomes at the

later mitotic stages in a subset of DYN-1-depleted embryos **(Figure 3)**, we hypothesized that this

may correlate with an abnormal localization of PLK-1 to the midbody during late mitosis.

Indeed, time-lapse live imaging revealed that PLK-1-sfGFP localization to the midbody was

defective in DYN-1 depleted embryos **(Figure 5)**. Specifically, we found that while all of the

control embryos (n=26 embryos) exhibited a stereotypical localization of PLK-1-sfGFP at the

center of the midbody, in about 72% of the DYN-1-depleted embryos (18 out of 25 embryos),

PLK-1-sfGFP failed to localize correctly to the midbody **(Figure 5)** (Fisher's exact test,

p<0.0001). As reported previously, in some severe cases, the cleavage furrow failed to form

completely upon DYN-1 depletion (Thompson *et al.,* 2002). In these cases, we observed that

PLK-1-sfGFP localization was severely defective during both anaphase and telophase

(Supplementary Movie S1). Notably, PLK-1-sfGFP failed to localize correctly to the midbody

in 17 out of the 18 embryos that were quantified as having midbody defects in **Figure 4**. These

data indicate that proper PLK-1 localization to the midbody is likely a prerequisite for normal

midbody assembly.

Total protein levels of Plk1 are not increased upon DNM2 depletion in HeLa cells

Since we observed increased PLK-1 levels at the centrosomes upon DYN-1 depletion **(Figure 2)**,

we asked whether this was occurring due to an overall increase in total PLK-1 levels or due to a

specific enhancement of PLK-1-sfGFP localization to centrosomes. To answer this question, we

performed western blot analysis of *C. elegans* whole worm lysates from control and *dyn-1 RNAi*

worms and probed them for PLK-1-sfGFP. Unfortunately, we could not detect any PLK-1-sfGFP

expression in our lysates, possibly due to its low abundance in whole worm lysates (data not

shown). Since both PLK-1 and DYN-1 proteins are conserved between humans and *C. elegans*

(Clustal Omega)*,* we turned to HeLa cells to answer this question. We were able to successfully

knockdown DNM2 (human DYN-1 homolog) in HeLa cells **(Figure S2A).** The DNM2

knockdown efficiency in our experiments ranged from ~81% to 95.4% **(Figure S2B)**.

Normalized Plk1 expression levels measured from four independent experiments did not reveal a

statistically significant increase in Plk1 levels upon DNM2 depletion **(Figure S2C)** (unpaired

two-tailed t-test with Welch's correction, p=0.4516). These data support the notion that the total

cellular levels of Plk1 are likely not increased upon DNM2 depletion.

In summary, our studies have identified a new role for the endocytic protein DYN-1 in

regulating metaphase spindle assembly and PLK-1 localization during mitosis. Specifically, we

find that upon DYN-1 depletion, PLK-1 accumulates at the centrosomes in metaphase and this is

associated with aberrant metaphase spindle pole assembly. Furthermore, while PLK-1 is

normally removed from the centrosomes during late mitosis, upon DYN-1 depletion, we find that

PLK-1 persists at the centrosomes during the later stages of mitosis. We hypothesize that a

failure to remove PLK-1 from the centrosomes in a timely manner likely causes midbody defects

and results in cytokinesis failure in DYN-1-depleted embryos.

Moving forward, a detailed mechanistic analysis will be carried out to further investigate

the cross-talk between DYN-1 and PLK-1. Specifically, we will utilize mutants such as *dyn-

1(ky51ts)* to determine if the vesicular trafficking function of DYN-1 is required for this

phenotype (Clark *et al.,* 1997; Grant and Hirsh 1999). Future experiments will be also designed

to test if DYN-1 and PLK-1 physically interact or if perturbing the functions of other endocytic

proteins phenocopies the DYN-1 depletion phenotype. We will also determine whether the

localization of key cytokinesis regulators (e.g. CYK-4, NOP-1, NMY-2, ANI-1, RHO-1, ECT-2)

is altered upon DYN-1 depletion. Rescue experiments involving partial knockdown of PLK-1 in

dyn-1 RNAi embryos could potentially clarify whether elevated centrosomal PLK-1 levels

contribute to the observed mitotic and cytokinetic defects of DYN-1 depletion. However, in our

experience, achieving consistent and titratable knockdown of PLK-1 using *RNAi* by feeding has

proven challenging. As a result, this experiment was not pursued in the present study. As an

alternative, future studies will utilize the temperature-sensitive *plk-1(or683ts)* mutant to achieve

partial loss of PLK-1 function under semi-permissive conditions (O'Rourke *et al.,* 2011). This

allele was shown to remove PLK-1 from centrosomes in five hours at the restrictive temperature

of 25 °C (Martino *et al.,* 2017). To test whether reducing PLK-1 levels can rescue the cytokinesis

defects of *dyn-1*–depleted embryos, we will need to carefully titrate the inactivation window to

achieve centrosomal PLK-1 levels compatible with normal cell division. While promising, this

strategy is technically challenging, as it requires precise timing of embryo imaging within a

narrow window to achieve the optimal level of PLK-1 reduction. Despite these constraints, this

approach offers a powerful means to directly test whether elevated PLK-1 at centrosomes

underlies the mitotic and cytokinetic phenotypes observed upon DYN-1 depletion.

MATERIALS AND METHODS

C. elegans **strains growth and maintenance**

All *C. elegans* strains described in this study were grown and maintained using standard

techniques on OP50 seeded Modified Youngren's Only Bacto-peptone (MYOB) agar plates at

20ºC (the strains were shifted to other assay-specific temperatures as needed).

C. elegans **strains and genetic crosses**

The wild-type N2 strain and the OD2425 strain expressing PLK-1-sfGFP (*plk-1(lt18[plk-*

1::sfGFP]::loxp) III) (Martino *et al.,* 2017) were obtained from the *Caenorhabditis Genetics*

Center (CGC). The OC908 strain (*bsSi30[pCW9: unc-119(+) pcdk-11.2::sfgfp::his-58::cdk-11.2*

3' UTR] II; bsIs20[pNP99: unc-119(+) tbb-1p::mCherry::tbb-2::tbb-2 3'-UTR]; bsIs2[pCK5.5:

Ppie-1::gfp::spd-2]) and the OC779 strain (*bsSi15[pKO109: spd-2p-spd-2::mCherry::spd-2 3'-*

utr, unc-119(+)] I; bsSi30[pCW9: unc-119(+) pcdk-11.2::sfgfp::his-58::cdk-11.2 3' utr] II; unc-

119(ed3) III) were obtained from Dr. Kevin O'Connell's laboratory at NIDDK, NIH. The

mCherry–tubulin–expressing strain IYR028 (*bsIs20[pNP99: unc-119(+) tbb-1p::mCherry::tbb-*

2::tbb-2 3'-UTR]) was generated in our laboratory by crossing OC908 hermaphrodites with N2

males and isolating progeny carrying only the mCherry-tubulin transgene. Following this,

mCherry-tubulin expressing males were generated from the IYR028 strain by heat shock at 30°C

for 6 hours. These mCherry-tubulin males from the IYR028 strain were then crossed with PLK-

1-sfGFP expressing hermaphrodites from the OD2425 strain to generate the IYR038 strain

(*bsIs20[pNP99: unc-119(+) tbb-1p::mCherry::tbb-2::tbb-2 3'-UTR]; plk-1(lt18[plk-*

1::sGFP]::loxP) III) which was homozygous for both mCherry-Tubulin and PLK-1-sfGFP. This

IYR038 strain was used for all but one of the imaging experiments described in this study. The

experiment involving the analysis of cytokinesis failure utilized the OC779 strain expressing

mCherry SPD-2 and GFP-histone.

RNAi **by feeding**

HT115 bacteria expressing the *dyn-1 RNAi* construct were purchased from Horizon Discovery

Ltd. (Lafayette, CO) while HT115 bacteria expressing the *smd-1 RNAi* construct (control) were

obtained from Dr. Kevin O'Connell's laboratory (NIDDK, Bethesda, MD). HT115 *E. coli*

transformed with either control (*smd-1*) or *dyn-1 RNAi* plasmids were grown on MYOB plates

supplemented with 50 µg/ml carbenicillin and 2 mM Isopropyl β- d-1-thiogalactopyranoside

(IPTG) as described previously (Kamath and Ahringer J 2003; Iyer *et al.,* 2022). Briefly, L4-

stage *C. elegans* from either the IYR038 or the OC779 strains were transferred onto the

respective *RNAi* plates and incubated at 25 ºC for 24 to 36 hours. Gravid adults from these plates

were then dissected to release their embryos and 1-cell stage embryos were analyzed by confocal

live imaging.

C. elegans **embryo live imaging using a spinning-disk confocal microscope**

C. elegans embryos obtained from dissected *smd-1* and *dyn-1 RNAi* treated worms were imaged

using an Olympus Yokagawa X1 spinning-disk confocal microscope as described previously

(Bergwell *et al.,* 2023) with a few alterations. Timelapse imaging was performed using the

cellSens software (Olympus America Inc.) by taking 1.5 µm Z-stacks every ~75 seconds at a

60X magnification with a 1.42 numerical aperture (NA) objective using a Prime 95B CMOS

camera. To enable quantitative analysis, all images were captured at the same laser intensity

(50% laser intensity for both 561 nm and 488 nm channels) and exposure time (700 msec

exposure time for both 561 nm and 488 nm channels). Maximum intensity Z-projections were

obtained using the cellSens software and converted to a .png format for all representative images

and analyses. All brightness and contrast adjustments were applied equally for both the control

and *dyn-1 RNAi* representative images to enable better image presentation and viewing.

Cytokinesis failure quantification

To score cytokinesis failure, the development of each embryo was monitored from the 1-cell

stage to the 4-cell stage. If the embryonic nuclei coalesced back to set up a multipolar spindle at

the 2-cell stage, the embryo was scored as having failed cytokinesis.

Spindle pole area measurement

To measure spindle pole area, the cellSens software (Olympus America Inc.) was used to obtain

maximum intensity Z-projections of the original .vsi files. These Z-projections were saved in a

.png format with the fixed scaling (left:80 and right:400) for all images and further analyzed

using the Nikon NIS-Elements software (Nikon Instruments Inc.). The calibration on the NIS-

Elements software was set to 1 pixel = 0.183 µm. The thresholding tool in NIS-Elements was

used to define and compute the spindle pole area (in µm²) (intense mCherry-tubulin fluorescence

demarcated each spindle pole). Only regions with clearly defined spindle poles were included in

the analysis. Embryos in which spindle poles could not be reliably distinguished were excluded.

PLK-1 metaphase centrosome intensity measurements

In our time-lapse imaging experiments, the stage immediately prior to anaphase onset (~75

seconds before anaphase) was designated as metaphase. Notably, there was some embryo-to-

embryo variability in chromosome alignment at this stage; in some embryos, the chromosomes

were centrally aligned, while in others, they were still in the process of aligning. To quantify

PLK-1–sfGFP fluorescence at the centrosomes, .avi files of maximum intensity Z-projections of

metaphase embryos were analyzed using Nikon NIS-Elements software (Nikon Instruments,

Inc.), with image calibration set to 1 pixel = 0.183333 μm. A region of interest (ROI) measuring

13.81 μm² was drawn around each centrosome and within the cytoplasm of the same embryo.

Cytoplasmic PLK-1–sfGFP intensity was then subtracted from centrosomal PLK-1-sfGFP

intensity to calculate the mean PLK-1-sfGFP fluorescence intensity, reported in arbitrary units.

Embryos with inaccurate PLK-1-sfGFP fluorescence due to debris on top of the embryo

affecting maximum intensity Z-projections were excluded from our analysis. Embryos where the

centrosomal PLK-1–sfGFP signal could not be clearly distinguished from the chromosomal

PLK-1-sfGFP signal were also excluded from our analysis.

Analysis of PLK-1 persistence at centrosomes

To analyze PLK-1-sfGFP persistence at centrosomes, maximum intensity Z-projections of each

timelapse sequence were visually analyzed for the presence or the absence of PLK-1-sfGFP at

the centrosomes. Centrosomes where the fluorescence intensity of PLK-1-sfGFP was visually

determined to be greater than the cytoplasmic background PLK-1-sfGFP fluorescence were

scored as being positive for PLK-1-sfGFP. The cell cycle stages were classified as follows:

metaphase: the chromosomes were aligning at the center of the cell and a diamond-shaped

spindle was formed, early anaphase: ~75 seconds after metaphase when the chromosomes start to

separate, late anaphase: ~75 seconds after early anaphase, early telophase: ~75 seconds after late

anaphase and late telophase: ~75 seconds after early telophase.

Quantification of midbody abnormalities

Maximum intensity Z-projections of the 561 (red) channel from the imaged IYR038 strain were

analyzed for midbody defects. Midbody assembly was analyzed at either 300 or 375 seconds

after metaphase onset. A normal midbody was classified as a structure where an intense tubulin

bridge was clearly visible at the abscission area close to the center of the embryo. An abnormal

midbody was defined as one where either no intense tubulin bridge connecting the two daughter

cells was detected or where a structure resembling the midbody formed after the cell had already

entered the next cell cycle.

Quantification of PLK-1 midbody localization defects

Maximum intensity Z-projections were used to quantify PLK-1-sfGFP midbody localization

defects. PLK-1-sfGFP localization to the midbody was analyzed at either 300 or 375 seconds

after metaphase onset. Embryos where a bright PLK-1-sfGFP spot could be detected at the

abscission point during early to late telophase were classified as exhibiting a normal PLK-1

midbody localization. Embryos where PLK-1-sfGFP exhibited a scattered localization at the

cleavage furrow during early to late telophase were classified as exhibiting an abnormal PLK-1

localization.

Statistical analysis of *C. elegans* studies

Graphpad prism 10.5 (GraphPad Software, Inc., San Diego, CA) was used to perform statistical

analyses of all the data. For all the data, error bars represent the standard deviation (s.d.) above

and below the mean and the middle lines represent the mean. *p*-values were rounded to two

decimal places and considered significant if $p<0.05$. A two-tailed unpaired t-test was used to

analyze the spindle pole area and the PLK-1-sfGFP fluorescence intensity data. A two-tailed

unpaired t-test with Welch's correction was used to analyze band intensities from western

blotting. A Fisher's exact test was used to analyze the cytokinesis defects, PLK-1-sfGFP

persistence at the centrosomes, defects in midbody formation and defects in PLK-1-sfGFP

localization to the midbody.

Cell culture and siRNA transfections

HeLa cells were maintained by culturing them in Dulbecco's Modified Eagle Medium (DMEM)

(Gibco, Catalog #11-995-040) with 10% fetal bovine serum (FBS) (Corning, Catalog

#MT35010CV) and 1% penicillin/streptomycin (1% P/S) (Gibco, Catalog #15-140-122) in a

37°C incubator with 5% CO_2. For DNM2 siRNA experiments, HeLa cells were seeded at a

density of up to $2.4X10^5$ cells per well (20 to 30% confluence) into each well of a 6-well plate

and allowed to adhere to the plate for 24 hours prior to transfection. The cells were rinsed twice

with DPBS (Gibco, Catalog # 14040-117) prior to transfection. Control and DNM2 siRNA

constructs were obtained from Santa Cruz Biotechnology Inc. (Control siRNA Catalog # sc-

37007; DNM2 siRNA Catalog # sc-35236). The siRNA constructs were transfected into HeLa

cells using Lipofectamine LTX reagent (Thermo Fisher Scientific, Inc., Catalog # 15-338-030) in

1 ml of reduced serum OPTI-MEM medium (Gibco, Catalog # 31985070) according to the

manufacturer's instructions. After six hours, 1 ml of DMEM containing 20% FBS and 2% P/S

was added to the transfected cells. After 24 hours, the media containing the transfection reagent

was removed and replaced with fresh DMEM with 10%FBS and 1% P/S. The cells were allowed

to grow for 48 hours and then lysed with RIPA buffer (Santa Cruz Biotechnology Inc, Catalog #

sc-24948A) containing 1 mM phenylmethylsulfonyl fluoride (PMSF) (Thermo Fisher Scientific,

Inc., Catalog # 36978) and a protease inhibitor cocktail (Pierce Protease Inhibitor Tablets,

EDTA-free, Catalog #A32965). Lysate protein concentrations were measured using a Bradford

assay (Thermo Scientific Inc., Catalog # 23200).

Western blotting

The HeLa cell lysates from control and DNM2 siRNA treated cells were subjected to western

blot analysis using the Bio-Rad wet transfer method. Briefly, approximately 20 micrograms of

protein from each sample were loaded onto a well of an SDS-PAGE gel (Genscript Corporation,

Catalog # M00656). The gel was run until adequate protein separation was achieved. Following

electrophoresis, the proteins were subjected to an overnight transfer onto a 0.2 μm nitrocellulose

membrane (Thermo Fisher Scientific Inc., Catalog # LC2000) for 16 hours at 30V using the

Towbin buffer (25 mM Tris base, 192 mM Glycine, 20% (v/v) Methanol) at 4ºC. The membrane

was then stained with the Ponceau S solution (Fisher Scientific Inc. Catalog # A40000279) to

ensure good protein transfer. Following this, the membranes were blocked with blocking buffer

(1X Tris-buffered saline (TBS) (0.05 M Tris, 0.15 M sodium chloride, pH=7.4), 0.2% gelatin

from cold water fish skin (Sigma-Aldrich Inc., Catalog # G7041), 0.1% Tween 20 (Thermo

Scientific Inc., Catalog # J20605-AP) and 0.02% sodium azide (RPI Research Products, Catalog

S24080-250)) at room temperature for 30 minutes. Once blocked, the membranes were probed

with a rabbit polyclonal anti-DNM2 antibody at a 1:1000 dilution (Abcam Inc., Catalog #

ab3457), a mouse monoclonal anti-tubulin antibody at a 1:100 dilution (Santa Cruz

Biotechnology Inc., Catalog # sc-32293), a mouse monoclonal anti-Plk1 antibody at a 1:100

dilution (Santa Cruz Biotechnology Inc., Catalog # sc-17783) for 2 hours at room temperature.

Subsequently, the membrane was washed 3 times with 1X TBS supplemented with 0.1% Tween-

20 (1XTBST) and incubated with the corresponding secondary antibodies (IRDye 680RD goat

anti-rabbit IgG secondary antibody (LI-COR Biotech, LLC, Catalog # 926-68071) and IRDye

800CW donkey anti-mouse IgG secondary antibody (LI-COR Biotech, LLC, Catalog # 926-

32212)) at a 1:10,000 dilution for 1 hour at room temperature. The membrane was washed again

3 times using 1XTBST and imaged using the LI-COR Odyssey Fc imaging system (LI-COR

Biotech, LLC).

Western blot intensity analysis

The band intensities for the different proteins were quantified using the LI-COR Image Studio

software (LI-COR Biotech LLC) and normalized with respect to the loading control (tubulin).

Band intensities from four independent western blot experiments were analyzed and plotted.

ACKNOWLEDGMENTS

This work was supported by NIH/NIGMS grant 5P20GM103636-09 and start-up funding from

North Carolina A&T State University to J.I. Additional support was provided by NIH/NIGMS

grant 1R35GM153737 and NCI grant 1R01CA273095 (to R.N.). We thank the University of

Tulsa for supporting Livinus Anyanwu and for providing infrastructure used in several

experiments. OC908 and OC779 strains were generously provided by Dr. Kevin O'Connell

(NIDDK, NIH). Some of the strains used in this study were provided by the CGC, which is

funded by NIH Office of Research Infrastructure Programs (P40 OD010440). The authors

declare no competing interests.

REFERENCES

Archambault V, Glover DM (2009). Polo-like kinases: conservation and divergence in their

functions and regulation. **Nat Rev Mol Cell Biol** *10*(4), 265-275.

Arnaud L, Pines J, Nigg EA (1998). GFP tagging reveals human Polo-like kinase 1 at the

kinetochore/centromere region of mitotic chromosomes. **Chromosoma** *107*(6), 424-429.

Bergwell M, Smith A, Smith E, Dierlam C, Duran R, Haastrup E, Napier-Jameson R, Seidel R, Potter W, Norris A, *et al.* (2023). A primary microcephaly-associated sas-6 mutation perturbs centrosome duplication, dendrite morphogenesis, and ciliogenesis in Caenorhabditis elegans. **Genetics** *224*(4), p.iyad105.

Brennan IM, Peters U, Kapoor TM, Straight AF (2007). Polo-like kinase controls vertebrate spindle elongation and cytokinesis. **PloS one** *2*(5), e409.

Budirahardja Y, Gönczy P (2008). PLK-1 asymmetry contributes to asynchronous cell division of *C. elegans* embryos. **Development** *135*(7), 1303–1313.

Burkard ME, Randall CL, Larochelle S, Zhang C, Shokat KM, Fisher RP, Jallepalli PV (2007). Chemical genetics reveals the requirement for Polo-like kinase 1 activity in positioning RhoA and triggering cytokinesis in human cells. **Proc Natl Acad Sci USA** *104*(11), 4383-4388.

Burkard ME, Maciejowski J, Rodriguez-Bravo V, Repka M, Lowery DM, Clauser KR, Zhang C, Shokat KM, Carr SA, Yaffe MB, *et al.* (2009). Plk1 self-organization and priming phosphorylation of HsCYK-4 at the spindle midzone regulate the onset of division in human cells. **PLoS Biol** *7*(5), p.e1000111.

Chase D, Serafinas C, Ashcroft N, Kosinski M, Longo D, Ferris DK, Golden, A (2000). The polo-like kinase PLK-1 is required for nuclear envelope breakdown and the completion of meiosis in Caenorhabditis elegans. **Genesis**, *26*(1), 26-41.

Clark SG, Shurland DL, Meyerowitz EM, Bargmann CI, Van Der Bliek AM (1997). A dynamin GTPase mutation causes a rapid and reversible temperature-inducible locomotion defect in C. elegans. **Proc Natl Acad Sci USA** *94*(19), 10438-10443.

Golsteyn RM, Mundt KE, Fry AM, Nigg EA (1995). Cell cycle regulation of the activity and

subcellular localization of Plk1, a human protein kinase implicated in mitotic spindle function. **J

Cell Biol** *129*(6), 1617-1628.

Grant B, Hirsh D (1999). Receptor-mediated endocytosis in the Caenorhabditis elegans

oocyte. **Mol Biol Cell** *10*(12), 4311-4326.

Hinshaw JE (2000). Dynamin and its role in membrane fission. **Annu Rev Cell Dev** *16*(1), 483-

519.

Iyer J, Gentry LK, Bergwell M, Smith A, Guagliardo S, Kropp PA, Sankaralingam P, Liu Y,

Spooner E, Bowerman B, *et al.* (2022). The chromatin remodeling protein CHD-1 and the EFL-

1/DPL-1 transcription factor cooperatively down regulate CDK-2 to control SAS-6 levels and

centriole number. **PLoS Genet** *18*(4), p.e1009799.

Kamath RS, Ahringer J (2003). Genome-wide RNAi screening in Caenorhabditis elegans.

Methods *30*(4), 313-321.

Khurana H, Pucadyil TJ (2023). "Gearing" up for dynamin-catalyzed membrane fission. **Curr

Opin Cell Biol** *83*, 102204.

Kishi K, Van Vugt MA, Okamoto KI, Hayashi Y, Yaffe MB (2009). Functional dynamics of

Polo-like kinase 1 at the centrosome. **Mol Cell Biol** *29*(11), 3134-3150.

Lee KS, Yuan YLO, Kuriyama R, Erikson RL (1995). Plk is an M-phase-specific protein kinase

and interacts with a kinesin-like protein, CHO1/MKLP-1. **Mol Cell Biol** *15*(12), 7143-7151.

Martino L, Morchoisne-Bolhy S, Cheerambathur DK, Van Hove L, Dumont J, Joly N, Desai A,

Doye V, Pintard L (2017). Channel nucleoporins recruit PLK-1 to nuclear pore complexes to

direct nuclear envelope breakdown in C. elegans. **Dev Cell** *43*(2), 157-171.

Menon M, Schafer DA (2013). Dynamin: expanding its scope to the cytoskeleton. **Int Rev Cell**

Mol Biol *302*, 187-219.

Nishi Y, Rogers E, Robertson SM, Lin R (2008). Polo kinases regulate C. elegans embryonic

polarity via binding to DYRK2-primed MEX-5 and MEX-6. **Development** *135*(4), 687–697.

O'Rourke SM, Carter C, Carter L, Christensen SN, Jones MP, Nash B, Price MH, Turnbull DW,

Garner AR, Hamill DR, *et al.* (2011). A survey of new temperature-sensitive, embryonic-lethal

mutations in C. elegans: 24 alleles of thirteen genes. **PLoS One** 6, e16644.

Pelissier A, Chauvin JP, Lecuit T (2003). Trafficking through Rab11 endosomes is required for

cellularization during Drosophila embryogenesis. **Curr Biol** *13*(21), 1848-1857.

Petronczki M, Glotzer M, Kraut N, Peters JM (2007). Polo-like kinase 1 triggers the initiation of

cytokinesis in human cells by promoting recruitment of the RhoGEF Ect2 to the central

spindle. **Dev Cell** *12*(5), 713-725.

Petronczki M, Lénárt P, Peters JM (2008). Polo on the rise—from mitotic entry to cytokinesis

with Plk1. **Dev Cell** *14*(5), 646-659.

Santamaria A, Neef R, Eberspächer U, Eis K, Husemann M, Mumberg D, Prechtl S, Schulze V,

Siemeister G, Wortmann L, *et al.* (2007). Use of the novel Plk1 inhibitor ZK-thiazolidinone to

elucidate functions of Plk1 in early and late stages of mitosis. **Mol Biol Cell** *18*(10), 4024-4036.

Schmucker S, Sumara I (2014). Molecular dynamics of PLK1 during mitosis. **Mol Cell Oncol**

1(2), e954507.

Shpetner HS, Vallee RB (1989). Identification of dynamin, a novel mechanochemical enzyme

that mediates interactions between microtubules. **Cell** *59*(3), 421-432.

Thompson HM, Skop AR, Euteneuer U, Meyer BJ, McNiven MA (2002). The large GTPase

dynamin associates with the spindle midzone and is required for cytokinesis. **Curr Biol**, *12*(24),

2111-2117.

Wienke DC, Knetsch ML, Neuhaus EM, Reedy MC, Manstein DJ (1999). Disruption of a

dynamin homologue affects endocytosis, organelle morphology, and cytokinesis in

Dictyostelium discoideum. **Mol Biol Cell** *10*(1), 225-243.

Wolfe BA, Takaki T, Petronczki M, Glotzer M (2009). Polo-like kinase 1 directs assembly of the

HsCyk-4 RhoGAP/Ect2 RhoGEF complex to initiate cleavage furrow formation. **PLoS**

Biol *7*(5), e1000110.

Zitouni S, Nabais C, Jana SC, Guerrero A, Bettencourt-Dias M (2014). Polo-like kinases:

structural variations lead to multiple functions. **Nat Rev Mol Cell Biol** *15*(7), 433-452.

Figure 1

A

B

Red: mCherry::Tubulin

Figure 1: DYN-1 depletion increases spindle pole size. (A) Stills from live imaging of *C. elegans* embryos demonstrating that spindle pole size is increased upon *dyn-1 RNAi*. White arrows highlight spindle poles. Scale bar = 10 µm. (B) Quantification of spindle pole area in control and DYN-1-depleted embryos using mCherry-tubulin fluorescence as a proxy for the spindle poles. Each circle represents the area of a single spindle pole. n = number of spindle poles analyzed. Error bars represent the s.d; middle lines represent the mean. Unpaired two-tailed t-test, p=0.0018.

Figure 2

Figure 2: DYN-1 depletion increases centrosomal PLK-1-sfGFP localization. (A) Stills from live imaging assays of *C. elegans* embryos expressing PLK-1-sfGFP (green) and mCherry-tubulin (red) demonstrating that centrosomal localization of PLK-1-sfGFP is increased upon *dyn-1* depletion. White arrows represent centrosome-localized PLK-1. Scale bar = 10 µm. (B) Quantification of PLK-1-sfGFP fluorescence intensity upon control and *dyn-1 RNAi*. Each circle represents the PLK-1-sfGFP fluorescence intensity at a single centrosome. n = number of centrosomes analyzed. Error bars represent the s.d; middle lines represent the mean. Unpaired two-tailed t-test, p<0.0001.

Figure 3

A

B

Figure 3: PLK-1-sfGFP persists at the centrosomes upon *dyn-1* knockdown (A) Stills from live imaging experiments of *C. elegans* embryos expressing PLK-1-sfGFP (green) demonstrating that PLK-1-sfGFP persists at the centrosomes upon *dyn-1* depletion. The metaphase timepoint was annotated as the zero timepoint. White arrows represent PLK-1-sfGFP at centrosomes. Scale bar = 10 µm. (B) Quantification of PLK-1-sfGFP localization at the centrosomes at different cell cycle stages. n = number of centrosomes analyzed.

Figure 4

A

B

Figure 4: DYN-1 knockdown causes midbody defects. (A) Stills from live imaging of *C. elegans* embryos demonstrating that midbody assembly is defective upon DYN-1 depletion. Red: mCherry-tubulin, the white arrow represents a normal midbody. Scale bar = 10 µm. (B) Quantification of midbody defects observed upon *dyn-1* depletion. n= number of embryos analyzed. Fisher's exact test, $p < 0.0001$.

Figure 5

A

B

Figure 5: PLK-1-sfGFP fails to localize correctly to the midbody upon *dyn-1 RNAi*. (A)
Stills from live imaging of *C. elegans* embryos expressing PLK-1-sfGFP (green) and mCherry-
tubulin (red) demonstrating that the midbody localization of PLK-1 is perturbed upon *dyn-1*
depletion. The white arrow represents a normal PLK-1 localization at the midbody. Scale bar =
10 μm. (B) Quantification of PLK-1-sfGFP midbody localization upon *dyn-1 RNAi*. n = number
of embryos analyzed. Fisher's exact test, p<0.0001.