Differential sensitivity to LINE 1-induced damage contributes to the expansion of Tet2-deficient HSCs upon chronic inflammatory stress

Anne Stolz[1,2,3], Cheng Xuan[1,2,3], Christelle Freitas[4], Rabie Chelbi[1,2,3,5], Hassan Ait-Ougouram[1,2,3], David Fandrei[1,2,3], Mengliang Ye[6], Elisabeth Nelson[4], Claire Maillard[4], Mathieu Bohm[1,2,3,5], Salomeh El-Choufani[1,2,3], Audrey Porquet[1,2,3], Weiwei Zhang[1,2,3], Kahina Issoulaiene[1,2,3], Christian Muchardt[7], Elina Zueva[6], Michele Goodhardt[4], David Garrick[4*], Françoise Porteu[1,2,3*] and Emilie Elvira-Matelot[1,2,3*§]

[1]INSERM UMR1287,

[2]Université Paris Saclay,

[3]Gustave Roussy Cancer Center, 94805 Villejuif, France.

[4]INSERM UMR1342, Institut de Recherche Saint Louis, Université Paris Cité, 75010 Paris, France.

[5]Inovarion, 75005 Paris, France

[6]INSERM U932, Institute Curie, Paris Sciences-et-Lettres Research University, 75005 Paris

[7]UMR 7238, Institut de Biologie Paris Seine, 75252 Paris

[*]Equal contribution

[§]to whom the correspondence should be sent:

Emilie Elvira-Matelot (emilie.elvira-matelot@inserm.fr)

INSERM U1287
Cancer Campus Gustave Roussy, PR1
114 rue Édouard-Vaillant
94805 Villejuif Cedex-France
Tel: (33) 1 42 11 23 48

Running title: LINE-1 as a new player in expansion of Tet2[-/-] HSC upon inflammation.

SUMMARY

Chronic inflammation that accompanies aging impairs the function of hematopoietic stem cells (HSCs) and promotes expansion of *TET2*-mutated cells, leading to clonal hematopoiesis of indeterminate potential (CHIP). The molecular mechanisms underlying these processes are still unclear. We recently showed that age and stresses like irradiation reduce the heterochromatin mark H3K9me3, leading to the epigenetic derepression of transposable elements such as LINE-1/L1, and L1-induced HSC functional changes through DNA damage and transcriptomic alterations. However, the impact of TEs on the clonal expansion of *Tet2*$^{-/-}$ HSCs upon chronic inflammation is unknown. Here we show that in wild type (WT) HSC, chronic inflammation induced by repeated low doses of lipopolysaccharide (LPS) triggers H3K9me3 loss at L1. We further show that L1s are involved in LPS-induced DNA damage and reduction of HSC clonogenicity. In contrast, *Tet2*$^{-/-}$ HSCs resist LPS-induced L1 epigenetic derepression and associated DNA damage. As a result, WT HSCs show decreased competitiveness against *Tet2*$^{-/-}$ HSCs upon chronic inflammation. These findings identify epigenetic control of L1s as a key mediator of inflammation-induced HSC dysfunction and clonal expansion of *Tet2*-mutant cells.

Keywords: hematopoietic stem cells, clonal hematopoiesis of indeterminate potential, inflammation, heterochromatin, transposable elements.

Non-standard abbreviations: HSC : hematopoietic stem cell; CHIP: clonal hematopoiesis of indeterminate potential, MDS: myelodysplastic syndromes, AML: acute myeloid leukemia, CMML: chronic myelomonocytic leukemia, TE: transposable element; LTR: long terminal repeat; ERV: endogenous retrovirus; LINE/SINE: long/short interspersed elements

INTRODUCTION

As cells age, they acquire somatic mutations due to environmental exposure and replication errors. Hematopoietic stem cells (HSCs), that give rise to all blood cells throughout life, are not spared by this phenomenon. While most of these mutations will have no functional consequence, some may lead to a clonal advantage and subsequent expansion. In the hematopoietic tissue, this phenomenon is referred to as clonal hematopoiesis of indeterminate potential (CHIP). CHIP is defined as the presence of somatic mutations with a variant allele frequency ≥2% in the peripheral blood of individuals without evidence of hematologic disease (Busque et al., 2012; Jaiswal et al., 2014; Genovese et al., 2014). The frequency of allelic variants at a given time depends both on the age at which the mutation appears (and the time the clone has had to expand) and on the clone's ability to gain an advantage over non-mutated clones. The prevalence of CHIP increases with age and in a context of low-grade inflammation, and is present in more than 20% of people over the age of 70 (Jakobsen et al., 2024).

Most CHIP mutations are found in genes commonly mutated in myelodysplastic syndrome (MDS) and acute myeloid leukemia (AML), with mutations in epigenetic factors such as TET2, DNMT3A and ASXL1 being the most frequent (Jaiswal et al., 2014; Genovese et al., 2014; Kar et al., 2022; Bick et al., 2020). CHIP is associated with an increased risk of atherosclerosis and cardiovascular diseases (Fuster et al., 2017; Sano et al., 2018b; a) and with a 10-fold increased risk of leukemia (Jaiswal et al., 2014; Genovese et al., 2014; Bowman et al., 2018; Mitchell et al., 2021). It is thus considered as a preleukemic state. Around 75% of AML patients have a CHIP mutation years before diagnosis (Desai et al., 2018). Understanding the mechanisms by which CHIP mutations confer a selective advantage to HSCs and drive the emergence of preleukemic clones is therefore of critical importance.

Several studies have investigated the mechanisms by which specific mutant clones gain an advantage over non-mutated HSCs and have led to a better understanding of the selective environments that drive their expansion (Florez et al., 2022). For example, clones with mutations in DNA repair pathways are selected and expand preferentially in response to genotoxic stresses, such as ionizing radiations, in the context of space exploration (Brojakowska et al., 2022; Mencia-Trinchant et al., 2020), or in radiotherapy and chemotherapy (Takahashi et al., 2017; Coombs et al., 2017; Bolton et al., 2020; Gillis et al., 2017; Desai and Roboz, 2019). In contrast, expansion of clones mutated in epigenetic factors, such as TET2, appears to be favored by the emergence of an inflammatory microenvironment in the bone marrow which is a characteristic feature of aging (inflammaging) (Cook et al., 2020; Cai et al., 2018; Caiado et al., 2023; McClatchy et al., 2023). Further, since TET2 mutant HSCs display a myeloid-biased potential (Quivoron et al., 2011; Li et al., 2011; Moran-Crusio et al., 2011; Ko et al., 2011; Jakobsen et al., 2024) and give rise to differentiated blood cells with enhanced inflammatory properties (Yeaton et al., 2022; Pronier et al., 2022; Zhang et al., 2015; Sano et al., 2018b; Fuster et al., 2017), the expansion of these mutant HSCs is also likely to further promote the inflammatory microenvironment of the bone marrow in a feed-forward manner. This proinflammatory role has been linked to the higher risk of developing cardiovascular or autoimmune diseases (Cobo et al., 2022). *TET2* mutated HSCs are resistant to inflammatory stimuli and cytokines that can be produced by their myeloid progeny (Cai et al., 2018; Caiado et al., 2023; Abegunde et al., 2018; Jakobsen et al., 2024), and thus present a selective advantage which eventually leads to their expansion (Cook et al., 2020; Avagyan et al., 2021; Speck, 2022; Jakobsen et al., 2024). However, the molecular mechanisms involved in the clonal expansion of *TET2* mutated HSCs upon inflammation are still unclear.

Reorganization of heterochromatin is one of the most commonly reported changes in aging and senescence. Maintenance of HSC identity is dependent on the trimethylation of lysine 9 of histone H3 (H3K9me3) (Koide et al., 2016), and loss of H3K9me3 leads to an accelerated hematopoietic aging (Keenan et al., 2020). We have shown that decreased H3K9me3 occurs in both human and mouse HSCs with age and is associated with a decrease of B cell differentiation potential (Djeghloul et al., 2016). One very important role of heterochromatin, through DNA methylation and H3K9me3, is the repression of transposable elements (TEs). Retrotransposable elements (RTEs), which are further classified as either long terminal repeat (LTR) sequences (including endogenous retroviruses (ERV)), or non-LTR elements (including long and short interspersed elements (LINE-1/L1; SINE)), represent almost half of the human and mouse genomes. Originally thought of as junk DNA, they are gaining increasing interest, especially in aging and cancer which are strongly associated with chromatin disorganization. RTEs constitute an important source of genomic instability since they can spread in the genome through a copy/paste mechanism and induce DNA damage (Gasior et al., 2006; Belancio et al., 2010; Barbieri et al., 2018). TEs also serve as a source of endogenous double-stranded (ds) RNA or cytoplasmic cDNA that are sensed as endogenous genomic parasites by members of the RIG-I-like (RLR) receptors or cGAS/STING pathways, eventually triggering the production of interferon type-I (IFN) and other inflammatory cytokines through activation of IRF3/NF-κB pathways (Gazquez-Gutierrez et al., 2021).

Erosion of heterochromatin upon aging is accompanied by overexpression of TEs and increased low-grade inflammation (De Cecco et al., 2013, 2019). We have previously shown that decreased H3K9me3 in HSCs upon aging and irradiation stress is associated with an increased expression of the evolutionarily recent and active L1s and LTRs (L1Md and IAP respectively in mouse) (Barbieri et al., 2018; Djeghloul et al., 2016; Pelinski et al., 2022). TE overexpression contributes to irradiation-induced long-term DNA damage in HSCs and loss

of HSC function which can be reversed by reverse transcriptase inhibitors (RTIs) (Barbieri et al., 2018). TE expression is also increased following chromatin decondensation in HSCs after chemotherapies, leading to intrinsic sterile inflammation and HSC exit from quiescence (Clapes et al., 2021). These data suggest that age or stress-induced alterations in heterochromatin and TE derepression may be a common and central molecular mechanism involved in HSC functional changes. However, the contribution of these mechanisms to the expansion of TET2 mutated cells upon inflammation has never been addressed.

In this study, we demonstrate that chronic inflammation, mediated by repeated low doses of lipopolysaccharide (LPS) mimicking inflammaging, triggers heterochromatin erosion, derepression of L1 elements and L1-induced DNA damage in wild-type HSCs. In contrast, *Tet2*[-/-] HSCs resist the deleterious effects of inflammation by epigenetically repressing L1, ultimately leading to their clonal expansion.

RESULTS

Chronic inflammation leads to decreased H3K9me3 heterochromatin mark in HSCs

To investigate the differences between wild-type (WT) and *Tet2* mutated HSCs that may explain the latter's clonal advantage upon inflammation, we first characterized the effect of chronic inflammation on the heterochromatin of WT cells. For this, we used a mouse model of chronic inflammation mediated by serial injections of low doses of LPS, which has been shown to mimic the low-level chronic inflammation observed with age and leads to impaired HSC function and aging-like changes in hematopoiesis (Esplin et al., 2011; Lescale et al., 2010; Dykstra et al., 2011; Rossi et al., 2005).

In these experiments, WT mice were injected with 6 µg LPS every 2 days for 4 weeks (**Figure 1A),** and the effect of this treatment on the repressive histone modification H3K9me3 in HSCs was analyzed by immunofluorescence. Interestingly, as with aging (Djeghloul et al.,

2016), we observed a significant decrease in global levels of H3K9me3 after LPS treatment in Lin- Sca+ Kit+ CD34- Flk2- HSCs (**Figure 1B**). This change was not due to a decrease in the total level of histone H3 which is rather increased in the presence of LPS (**Figure 1C**). The reduction in H3K9me3 induced by LPS was still detectable in HSCs isolated from mice 8 weeks after the end of the LPS treatment, indicating that LPS induces a stable epigenetic change **(Figure 1D)**. This parallels the long-term damage to HSC function, which is also maintained after removal of the inflammatory stimulus (Esplin et al., 2011; Takizawa et al., 2017). To determine if decreased H3K9me3 in LPS-treated HSCs could be a general feature of chronic inflammation, we analyzed HSC heterochromatin in a second model of chronic inflammation, the Db/Db obese mice (Chua et al., 1996). These mice carry a mutation in the *LepR* gene encoding the Leptin receptor, leading to rapid weight gain associated with systemic inflammation and a myeloid-biased hematopoietic compartment (Nagareddy et al., 2014). Immunofluorescence staining revealed a significant decrease in global levels of H3K9me3 in HSCs of Db/Db mice relative to that of control (Db/+) littermates **(Figure 1E)**. A similar, although less pronounced, reduction in H3K9me3 was also observed in Lin- Sca+ Kit+ CD34+ Flk2- short-term reconstituting stem cells (ST-HSCs) of Db/Db mice. These results support our findings in LPS-treated mice and strongly suggest that chronic inflammation, like aging, leads to impairment of H3K9me3 heterochromatin in HSCs. To investigate this further, we generated mixed bone marrow chimeras by reconstitution of lethally irradiated mice with a 1:1 ratio of bone marrow cells from wild-type mice and from mice deleted for *Tlr4*, encoding the Toll-like receptor 4, the pattern-recognition receptor responsible for the detection of LPS stimulation in HSCs (Takizawa et al., 2017). The mice were treated with LPS as described above, and the effect on H3K9me3 was analyzed by immunofluorescence in WT and *Tlr4*[-/-] HSCs, which could be distinguished using CD45.1/CD45.2 isotypic markers (**Figure 1F**). Strikingly, while a decrease in H3K9me3 was

observed in CD45.1[+] WT HSCs, no decrease was detected in CD45.2[+] *Tlr4[-/-]* HSCs from the same LPS-treated mice, indicating that LPS-mediated changes in HSC heterochromatin occur via a direct interaction with Tlr4 receptors expressed by the stem cells. Altogether, these findings indicate that chronic inflammation directly impacts the chromatin state of HSCs, leading to a stable reduction of H3K9me3.

Chronic inflammation induces a loss of H3K9me3 at L1Md and IAP, the most recent subfamilies of L1s and LTRs

To further characterize the loss of H3K9me3 that occurs upon chronic inflammation, we performed H3K9me3 CUT&Tag experiments in HSCs sorted from LPS-treated mice (**Figure 1A).** We have previously shown that in HSCs, H3K9me3 is mainly enriched at TEs, and more particularly at L1Md elements, the most recent subfamilies of LINEs (Pelinski et al., 2022). In order to capture the information at these evolutionary youngest TE subfamilies, we performed a combined analysis of unique and multiple-mapping reads, as previously described (Pelinski et al., 2022). Focused differential enrichment analysis at TE genomic loci revealed 598 loci with significantly altered enrichment of H3K9me3, with 403 (67%) and 195 (33%) TEs showing decreased or increased H3K9me3 respectively upon LPS treatment (**Figure 2A).** Altogether across these 598 differentially altered TEs, there was a significant ($p<0.0001$) overall decrease in H3K9me3 concentration (**Figure 2B).** Among the TEs with decreased H3K9me3 following LPS treatment, there was a significant ($p<0.0001$-hypergeometric test) enrichment of LINEs (62.5%) and LTR (36.0%) as compared to the repartition of these TEs in the mouse genome (24.7% and 22.1% respectively) (**Figure 2C**). LINEs were also significantly enriched in TEs gaining H3K9me3 (43.8%), but to a lesser extent than TEs which lose H3K9me3 (**Figure 2C**). Consistent with these results, analysis across all 598 differentially affected TE loci confirmed a decrease in H3K9me3 at LINEs and at LTRs,

while SINE and DNA transposons displayed an increase in H3K9me3 (although DNA transposons represent relatively few loci) (**Figure 2D**). These data were confirmed in an independent H3K9me3 CUT&Tag experiment (**Figure S1 A-D**).

Since we and others have reported that evolutionarily recent TE subfamilies, particularly L1Md, are subject to tighter epigenetic regulation as compared to older TEs (Barau et al., 2016; Pezic et al., 2014; Pelinski et al., 2022), we further analyzed changes in H3K9me3 enrichment at LINEs and LTRs according to their evolutionary age, as calculated in (Sookdeo et al., 2013). As previously reported, we observed a negative correlation between the age of the LINE elements and H3K9me3 enrichment in WT HSCs, with the youngest repeats (typically L1Md_A, L1Md_Tf, and L1Md_Gf), showing the highest enrichment. Loss of H3K9me3 was mainly observed at these youngest subfamilies (**Figure 2E and S2A**). With respect to LTRs, while the anti-correlation between H3K9me3 enrichment and the age of LTRs is less pronounced, again the strongest enrichment of H3K9me3 was also observed at the youngest elements, especially IAPEz and MERVL (**Figure 2F and S2B**) and H3K9me3 loss in response to LPS treatment is mainly observed at these youngest subfamilies.

Finally, plot profiles of H3K9me3 enrichment across all L1Md and IAP sequences confirmed significant decrease of H3K9me3 at these elements in response to LPS treatment (**Figure 2G-H**). This decrease was particularly evident at elements which show the highest enrichment of H3K9me3 in the basal state: L1Md_A (**Figure S2C)** and IAPEz **(Figure S2D)** subfamilies. As previously described in embryonic stem cells (Bulut-Karslioglu et al., 2014), intact L1Md and IAP (conservation score > 97%), present a stronger H3K9me3 enrichment than degenerate elements (conservation score < 97%), and are the most affected by H3K9me3 loss upon LPS treatment (**Figure S2E-H).** Finally, H3K9me3 loss was specific for long L1Md (>5kb), that have more chance to contain the sequences encoding for the reverse transcription machinery (endonuclease and reverse transcriptase), compared to truncated L1Md (<5kb)

(Figure S2I-J). Altogether, these data suggest that chronic LPS treatment induces a decrease of H3K9me3 at evolutionarily recent and intact subfamilies of L1s and LTRs that are the most likely to be damaging to the cell.

Chronic inflammation induces a gain of H3K9me3 at L1Md in *Tet2*[-/-] HSCs

We next investigated the impact of chronic inflammation on H3K9me3 enrichment at TEs in *Tet2*[-/-] HSCs. We found 241 TE loci with significantly differential enrichment of H3K9me3 following LPS treatment. Surprisingly, by contrast with what was observed in WT HSCs, more loci (133, 55%) were observed to gain H3K9me3 following LPS treatment than those that lost it (108, 45%) (**Figure 3A).** Accordingly, the distribution of the H3K9me3 concentration at these differential peaks showed a slight but significant increase following LPS treatment (**Figure 3B).** Compared to the genomic distribution of TEs, there was a significant (p<0.001) enrichment of LTR (32.4%) among the TEs with decreased H3K9me3, while there was a significant (p<0.01) enrichment of LINEs (38.2%) among the TEs gaining H3K9me3 (**Figure 3C**). Consistent with this, analysis of H3K9me3 concentration at the 241 loci with differential H3K9me3 enrichment between *Tet2*[-/-]-LPS and *Tet2*[-/-] conditions confirmed a significant increase in H3K9me3 at LINEs, but no significant changes at other TE classes (**Figure 3D**). Finally, plot profile of H3K9me3 concentration along all L1Md sequences showed a significant increase of H3K9me3 at L1Mds upon LPS treatment in *Tet2*[-/-] HSCs (**Figure 3E**). This increase was notably observed at intact rather than degenerate L1Mds (**Figure S3A-B**) and long (>5kb) vs truncated (<5kb) L1Mds (**Figure S3C-D).** Altogether, these data show that LPS had inverse effects on H3K9me3 enrichment at L1 in WT and *Tet2*[-/-] HSCs.

Chronic LPS induces distinct transcriptomic changes in WT and *Tet2*[-/-] HSCs

To further investigate these different responses to LPS, we next compared the effect of chronic LPS treatment on the transcriptome in WT and $Tet2^{-/-}$ HSCs. Gene analysis revealed 332 differentially expressed genes (DEGs; p<0.05) upon LPS treatment in WT HSCs, of which 198 (60%) are down- and 134 (40%) are up-regulated (**Figure S4A**), and 1260 DEGs found after LPS treatment of $Tet2^{-/-}$ HSCs, of which 746 (59%) are down-regulated and 514 (41%) are up-regulated (**Figure S4B**) (**supplementary table S1**). Of the 1541 genes whose expression was altered by LPS in either WT and $Tet2^{-/-}$ HSCs, only 51 genes (3.3%) were altered in both genotypes (**Figure S4C**), of which only 19 (1.2%) changed in the same direction (shown in green in **Figure S4D**).

We then performed Gene Set Enrichment Analysis (GSEA) on Hallmark gene sets (FDR<0.05, NES <-1.5 or >1.5) and showed that chronic LPS treatment led to inverse effects on gene pathways in WT and $Tet2^{-/-}$ HSCs. Indeed, while signatures of proliferative pathways (E2F and MYC targets) are downregulated following LPS treatment in WT HSCs (**Figure 4A-B**), they are positively enriched in $Tet2^{-/-}$ HSCs (**Figure 4A,C**). We also observed inverse responses of the TNFA signaling via NFKB pathway upon LPS-treatment in WT (enriched) and $Tet2^{-/-}$ (depleted) HSCs (**Figure 4A,D-E**).

Altogether, these data indicate that in comparison to WT HSCs, $Tet2^{-/-}$ HSCs show a distinct transcriptomic response to inflammation, associated with increased enrichment of proliferative pathways and decreased enrichment of inflammatory pathways.

As H3K9me3 is mainly found at TEs (Pelinski et al., 2022), we analyzed the consequences of its deregulation upon LPS in WT and $Tet2^{-/-}$ HSCs on TE expression using TEtranscripts (Jin et al., 2015) (multiple mapping reads). LPS significantly (p<0.05) increased expression of 94 (97%) and decreased expression of 3 (3%) TE subfamilies (**Figure 4F**) (**supplementary table S2**). On the contrary, among the 57 significantly deregulated TE subfamilies upon LPS treatment in $Tet2^{-/-}$ HSCs, only 7 (12%) were up-regulated and 50 (88%) were downregulated

(**Figure 4G**). Focusing on L1s that are inversely epigenetically regulated in WT and $Tet2^{-/-}$ HSCs by LPS, all of the significantly deregulated L1 in WT and $Tet2^{-/-}$ HSCs were up- and down-regulated respectively upon LPS treatment (**Figure S4E-F**). Altogether, these data indicate that chronic LPS treatment leads to opposite effects on TE and L1 expression in WT and $Tet2^{-/-}$ HSCs.

Derepression of TEs upon LPS treatment in WT HSC is associated with the accumulation of DNA damage and HSC functional changes

We and others have shown that L1 derepression and mobilization in HSCs are associated with double strand breaks (DSBs), as measured by the presence of γH2AX foci (Gasior et al., 2006; Belancio et al., 2010; Barbieri et al., 2018). Inflammation has also been shown to lead to genomic instability in HSPCs/HSCs (Takizawa et al., 2017; Rodriguez-Meira et al., 2023). L1s of greater than 5kb in length could encode an intact ORF2 protein with endonuclease and reverse transcriptase activities which can potentially induce DNA damage. In order to characterize the consequences of the epigenetic derepression of these L1s in WT HSCs upon LPS treatment, we assessed γH2AX foci by immunofluorescence in HSCs sorted from mice at the end of chronic LPS treatment in vivo (**Figure 1A**), or in HSCs from untreated mice following culture in the presence or absence of 10μg/ml LPS for 48h in vitro. In both cases, LPS treatment induced a significant increase in the number of γH2AX foci in HSCs (**Figure 5A-D**). In the in vitro conditions, the induction of γH2AX foci by LPS was inhibited in the presence of lamivudine (3TC), a reverse transcriptase inhibitor known to act on L1 reverse transcriptase activity (**Figure 5C-D**), indicating that L1 is responsible for DNA damage induced by LPS in HSCs.

To further assess whether L1-induced DNA damage affects HSC function upon LPS

treatment, we transduced Lin- cells with either an sh-control or an sh-L1 fused to mCherry reporter gene, that was previously shown to reduce L1Md expression in HSPCs (Clapes et al., 2021). We then plated mCherry positive HSCs in methylcellulose, in the presence or absence of 10μg/ml LPS, and assessed their clonogenicity (**Figure 5E).** While LPS induced a significant decrease in the number of colonies formed by HSCs transduced with an sh-control, it did not affect the clonogenicity of HSCs transduced with an sh-L1 (**Figure 5F).**

Altogether, these data suggest that loss of H3K9me3 at TEs, notably at L1s, contributes to the DNA damage and decrease in clonogenicity observed in HSC in response to chronic inflammation.

Tet2[−/−] **HSCs are protected from the L1-mediated deleterious effects of LPS, leading to their expansion.**

We then assessed whether epigenetic changes observed at L1 elements in *Tet2*[−/−] HSCs may prevent the accumulation of DNA damage and functional alterations. Indeed, in contrast with findings in WT HSCs, LPS did not increase γH2AX foci in *Tet2*[−/−] HSCs both *in vitro* in culture and *in vivo* after chronic injections (**Figure 6A-C**). As reported previously (Esplin et al., 2011), treatment with LPS decreased the reconstitution capacity of WT HSCs (**Figure S5A-D**). To assess how TET2 loss affects the response of HSC to LPS in vivo, hematopoietic chimeras were created by transplanting CD45.1 recipient mice with 0.3×10^6 bone marrow (BM) cells from CD45.2 WT or *Tet2*[−/−] mice (10%) in competition with 2.7×10^6 BM (90%) cells from CD45.1 WT mice. At 1 month post-engraftment, recipient mice were treated with PBS (control) or LPS every two days for 30 days and reconstitution was assessed in BM and peripheral blood (PB) at the end of 5 months (**Figure 6D**). LPS treatment had no significant impact on peripheral blood engraftment of either WT or *Tet2*[−/−] cells after 5 months (**Figure S5E**). However, the proportion of WT CD45.2[+] cells in Lin- Sca+ Kit+ (LSK) and HSC

populations (**Figure 6 E-F**), as well as their absolute numbers (**Figure 6 G-H)** were significantly reduced following LPS treatment as compared to PBS-treated mice. As previously described (Cai et al., 2018; Caiado et al., 2023), $Tet2^{-/-}$ HSCs present a significantly higher reconstitution capacity than WT HSCs (**Figure 6 E, Figure S5E**). In contrast to WT HSCs, the proportion of CD45.2$^+$ $Tet2^{-/-}$ in LSK and HSC populations (**Figure 6 E-F**) as well as their absolute numbers (**Figure 6 G-H)** were not affected by LPS treatment. To determine the functional consequences of LPS exposure on long-term HSC function, 5.0×10^6 BM cells from primary transplants were engrafted to secondary CD45.1 recipients. The effect of LPS on WT HSCs was still observed in these secondary transplantations, indicating a long-term influence on HSC function, while the reconstitution capacity of $Tet2^{-/-}$ HSCs was again unaffected (**Figure 6 I**).

To assess the role played by L1s in the resistance of $Tet2^{-/-}$ HSCs towards LPS treatment, we performed in vitro competition assays between WT HSCs transduced with either an sh-control or an sh-LINE fused to mCherry reporter gene, and $Tet2^{-/-}$ HSCs transduced with an sh-control fused to GFP reporter gene and followed the relative expansion of $Tet2^{-/-}$ HSCs (GFP+) after 3 days in culture (**Figure 7A**). The results showed that LPS had a significantly (p<0.01) stronger effect on the expansion of $Tet2^{-/-}$ HSCs when in competition with WT HSCs transduced with an sh-control than when in competition with WT HSCs transduced with an sh-L1 (**Figure 7B and S6A)**.

Altogether, these results suggest that the relative expansion of $Tet2^{-/-}$ HSCs induced by chronic inflammation is dependent on the presence of L1 transcripts and their deleterious effect on WT HSCs (**Figure 7C**).

DISCUSSION

Through their ability to modify the transcriptome and to induce genomic instability, TEs can exert an oncogenic activity (Chuong et al., 2017; Liang et al., 2024). However, when expressed above a certain threshold, TEs can induce DNA damage, senescence, and a state of "sterile inflammation" and thus play the role of tumor suppressors (Zhao et al., 2021; De Cecco et al., 2019). We previously showed that derepression of young active L1Md, which occurs upon H3K9me3 loss, is partly responsible for the sustained DNA damage which is observed in HSCs even a long-time after irradiation. Preventing this damage with RTI is associated with a significant improvement of HSC reconstitution capacities (Barbieri et al., 2018; Pelinski et al., 2022). We show here that L1 epigenetic derepression following H3K9me3 loss contributes to DNA damage in HSCs and their decreased clonogenicity upon chronic inflammation. We further show that $Tet2^{-/-}$ HSCs are refractory to LPS-induced epigenetic derepression of L1 elements and L1-induced DNA damage, giving them an advantage compared to WT HSCs, and eventually leading to their clonal expansion upon inflammation.

Global heterochromatin loss was proposed as a theory of aging (Tsurumi and Li, 2012). The current study adds to our previous findings and those of others, which demonstrate that heterochromatin loss and overexpression of TEs is observed in HSC upon chronological aging (Djeghloul et al., 2016; Hong et al., 2023), as well as in premature aging phenotypes induced by irradiation- (Barbieri et al., 2018; Pelinski et al., 2022) or inflammatory stress (this study). In all cases, RTI or depletion of L1 elements can enhance the clonogenic potential of HSPCs upon aging. These data suggest that heterochromatin loss at TEs and TE derepression may be a common and central molecular mechanism involved in the alteration of HSC function upon aging and stress-induced premature aging. Rather than a global loss of heterochromatin, it has

been proposed that aging may be the result of a redistribution of heterochromatin (Tsurumi and Li, 2012). We show that chronic inflammation leads to a relocalization of H3K9me3 at the different TE classes in HSCs, with a decreased concentration at some LINEs and LTR but an increased enrichment at some SINEs and DNA. These data are in agreement with those of Hong et al. who reported a spatial relocalization of H3K9me3-marked heterochromatin upon aging in HSPCs, which is accompanied by overexpression of TEs, notably LTRs, leading to TE-induced inflammatory response (Hong et al., 2023). Interestingly it has been shown that compared to HSCs, myeloid progenitors exhibit a relocalization of H3K9me3 from the periphery to the center of the nucleus (Ugarte et al., 2015), mimicking the redistribution of H3K9me3 observed in HSCs upon aging by Hong et al. (Ugarte et al., 2015; Hong et al., 2023). As HSCs present a biased differentiation towards the myeloid lineage upon aging, it would be of interest to study the role played by H3K9me3 relocalization and TE derepression on the induction of a myeloid program in HSCs.

Selective pressures on HSCs modulate their functions and responses to stress. Hematopoietic aging can be seen as the net result of accumulated DNA damage and functional alterations in combination with selective pressures arising from various age-associated stresses which eventually favor the emergence of competitive clones capable of self-renewal in the aged bone marrow. These emerging clones also display a predisposition to oncogenic transformation. We show here that the differential epigenetic regulation of L1 upon LPS treatment in $Tet2^{-/-}$ vs WT HSCs and the resistance of $Tet2^{-/-}$ HSCs towards TE-induced DNA damage may be a new player in the resistance of $Tet2^{-/-}$ HSCs towards inflammation, which contributes to their clonal expansion. Our findings add to previous studies showing that $Tet2^{-/-}$ HSCs are resistant towards epigenetic changes such as heterochromatin relocalization and loss of DNA methylation, which are observed upon aging and chronic IL1β treatment, respectively (Hong et al., 2023; McClatchy et al., 2023). Heterochromatin relocalization was

specifically associated with upregulation of LTRs in WT HSPCs upon aging, and the induction of an inflammatory response, which was not observed in $Tet2^{-/-}$ HSPCs (Hong et al., 2023). In our study of a more restricted HSC population, we did not observe differential regulation of LTRs, with H3K9me3 enrichment at LTRs decreasing in both WT and $Tet2^{-/-}$ HSCs upon LPS treatment. We observed that LPS induced opposing effects at intact L1s, which displayed decreased and increased H3K9me3 enrichment in WT and $Tet2^{-/-}$ HSCs respectively. Using sh-L1, we further demonstrate that L1s are necessary for the decreased competitivity of WT HSCs relative to $Tet2^{-/-}$ HSCs.

It has previously been shown in mouse embryonic stem cells that members of the HUSH complex are required for specific repression of L1 elements, and particularly the most recent L1Mds by recruiting the SETDB1 methyltransferase and H3K9me3 (Robbez-Masson et al., 2018; Liu et al., 2018). Inactivation of members of the HUSH complex leads to impaired proliferation of an AML cell line through L1-induced DNA damage (Gu et al., 2021). However, our RNA-seq analysis did not reveal a significant upregulation of members of the HUSH complex, such as Tasor, Mphosph8 (mpp8) or Pphln1 (periphilin 1), that could account for the specific increase in H3K9me3 enrichment at L1s in $Tet2^{-/-}$ HSCs upon LPS treatment (**supplementary table1**).

Previous studies have shown that the survival advantage of TET2-deficient HSCs under inflammatory conditions is linked to changes in anti- vs pro-apoptotic gene expression, increased expression of DNA repair pathways, and an increase in proliferative and HSC self-renewal transcriptomic signatures (Cai et al., 2018; Abegunde et al., 2018; Caiado et al., 2023; McClatchy et al., 2023). As compared to WT HSCs, $Tet2^{-/-}$ HSCs also show resistance to DNA demethylation of transcription factor binding sites involved in differentiation, which could contribute to their enhanced self-renewal capacity (McClatchy et al., 2023). We did not observe a maintenance of HSC signatures in $Tet2^{-/-}$ HSCs upon LPS treatment as compared to

WT in our RNA-seq data (data not shown). However, we confirm that LPS induced an increase in DNA repair pathways in *Tet2*$^{-/-}$ HSCs, which was not observed in WT cells (**Figure 4A**). In addition to the epigenetic repression of L1, the upregulation of DNA repair genes may thus also play a role in the resolution of DNA damage accumulation in *Tet2*$^{-/-}$ HSCs and their resistance towards LPS treatment. We further confirm the increase in proliferative pathways in *Tet2*$^{-/-}$ HSCs upon inflammation. However, in our hands, this was not associated with the expansion of *Tet2*$^{-/-}$ HSCs upon LPS treatment as defined by the total number of cells (in wells with at least one cell at a given time) and colony size (defined by the total number of cells per well) (**Figure S6 B-C**). Finally, as previously observed in TET2-mutated vs WT HSCs in individuals with CHIP (Jakobsen et al., 2024), our data show an attenuated transcriptional response of *Tet2*$^{-/-}$ HSCs towards inflammation, with a significant decrease of the TNFA_signalling_via_NFKB pathway. Hence, in addition to L1 epigenetic repression, this attenuated response may further increase the clonal advantage of *Tet2*$^{-/-}$ HSCs, as previously shown in Tet2 mutant HSC in zebrafish (Avagyan et al., 2021). Interestingly, we also observed a significant decrease in interferon gamma response pathway in *Tet2*$^{-/-}$ HSCs (**Figure 4A**), that could be the consequence of TE repression.

Following from our study, it will be of interest to understand how repression of TEs contributes to the predisposition to oncogenic transformation which is characteristic of Tet2-mutant HSC. Interestingly, expression of TEs is repressed in leukemic stem cells from patients with AML (Colombo et al., 2017). Further, in patients with chronic myelomonocytic leukemia (CMML), 60% of whom present with biallelic mutations of TET2, we recently showed an increase of the repressive H3K9me2 mark at TEs and a repression of immune associated transcripts. This could allow the resistance and further expansion of mutated cells as compared to their WT counterparts upon aging (Hidaoui et al., 2024). Epigenetic silencing of TEs by the HUSH complex and SETDB1 H3K9 methyltransferase is required for AML

cell line proliferation, by preventing TE-induced DNA damage and interferon response (Gu et al., 2021; Cuellar et al., 2017). Together, these findings suggest that TE repression could be an important early event contributing to clonal evolution and the establishment of a pre-leukemic state.

METHODS

Mice strain and treatment

$Tet2^{-/-}$ and WT littermates were provided by Olivier Bernard (INSERM U1170 – Gustave roussy)(Quivoron et al., 2011). Db/Db mice (BKS(D)-Leprdb/db/JOrlRj) and Db/+ controls (BKS(D)-Leprdb/+/JOrlRj) (14 weeks old) were obtained from Janvier Lab. TLR4$^{-/-}$ mice were provided by Christian Muchardt (Institut de Biologie Paris Seine). C57BL/6 CD45.2 or CD45.1 mice (6-8 weeks old) were from Envigo or Charles River laboratories respectively. All the mice were housed in a specific pathogen-free environment. All procedures were reviewed and approved by the Animal Care Committee N°26 (APAFIS #23286-2019121214392327 and APAFIS #35147-2022020315419972). $Tet2^{-/-}$ and WT littermates received intraperitoneal injections of 6 µg LPS-B5 Ultrapure (Invivogen, tlrl-pb5lps) every 2 days for 28 days and control mice were injected with 1x PBS (adaptation of the protocol described by Esplin et al. (2011)). Mice were irradiated with 9.5 Gy with an X-ray irradiator (RX irradiator X- RAD 320) or with a cesium irradiator (GSR D1®)-) for reconstitution experiments with total BM, or shRNA-transduced Lin- cells, respectively. Engraftments were performed by retro-orbital injections.

Cell harvest and culture

BM was harvested from femur, tibia and hip bones. Total BM was depleted of differentiated hematopoietic cells (lineage-positive; lin+ cells) using Mouse Hematopoietic Progenitor Stem Cell Enrichment Set (BD Biosciences) and LT-HSCs (Lin-, Sca+, c-Kit+,CD34-, Flk2-) or ST-HSCs (Lin-, Sca+, c-Kit+,CD34+, Flk2-) were sorted using ARIA3, ARIA Fusion or Influx cell sorters (BD Franklin Lakes, NJ, USA), using the following antibodies: CD3ε (Lin) – APC clone 145-2C11 (553066, BD), TER-119 (Lin) – APC clone Ter-119 (557909), CD45R/B220 (Lin) – APC clone RA3-6B2 (553092), Ly6G-6C (Lin)-APC clone RB6-8C5 (553129), Ly-6A/E (Sca-1) - PeCy7 clone d7 (558162), CD117 (c-Kit) – PE or PerCP-Cy5.5, clone 2B8 (553355 or 560557 respectively), CD34– FITC or AF700 clone RAM34 (560238 or 560518), CD135 (Flk2) – BV421 or PE clone A2F10.1 (562898 or 553842 respectively), all from BD Biosciences, and collected in StemSpan (StemCell Technologies).

Freshly isolated murine HSCs were cultured in StemSpan medium supplemented with 1% penicillin/streptomycin (P/S), containing the following cytokines (all from Miltenyi Biotech): FTL3-Ligand (FTL3-L, 10ng/ml), stem cell factor (SCF, 100ng/mL) interleukin 3 (IL3, 10ng/mL) and interleukin 6 (IL6, 10ng/mL), at 37°C, 5% CO_2. For treatments, 10μg/mL LPS-B5 Ultrapure (Invivogen, tlrl-pb5lps) or 10μM lamivudine (Sigma, 3TC- L1295-10MG) were added at the start of the cultures (day 0) for 48h.

Immunofluorescence

For H3K9me3 analysis: sorted HSCs were pipetted onto poly-lysine-coated slides (2.000-5.000 cells per spot), and allowed to fix to the slide for 1 h at room temperature in a humid chamber. Cells were then fixed with 2% formaldehyde / PBS for 20 min and permeabilized in 0.5% Triton X-100 / PBS for 15 min at room temperature. After washing three times with

PBS, cells were blocked for 1 h in 5% BSA / 5% goat serum / 0.05% Tween / PBS and were incubated with anti-H3K9me3 (07-523, Millipore) or anti-H3 (ab18521, Abcam) primary antibodies diluted in blocking solution) overnight at 4°C in a humid chamber. Slides were washed three times in PBS and incubated for 1 h at room temperature in 1% BSA / PBS containing anti-rabbit-Alexa488 (A11008, Invitrogen). After three washings in PBS, slides were mounted using VectaShield containing DAPI (H1200, Vector Laboratories). Stained cells were analyzed using a 63× objective lens Zeiss Lsm 800 Confocal microscope and Zeiss Zen blue software. Mean fluorescence intensity (Integrated density) of primary antibodies was measured using ImageJ software.

For analysis of DNA damage, 3000-5000 HSCs were cytospun onto glass slides and immunofluorescence was performed as previously described (de Laval et al., 2013b). γH2AX antibody was purchased from Millipore (05-636-I) and used at 1/2000. Detection was performed using Alexa Fluor 555-coupled anti-mouse secondary antibody (1/600) (Invitrogen, A-21425). slides were visualized using an SPE confocal microscope (Leica).

Cell transduction with sh-L1

Lentiviral vectors expressing control sh-RNA (scramble sequence: ATCTCGCTTGGGCGAGAGTAAGTAGTGAAGCCACAGATGTACTTACTCTCGCCCA AGCGAGAG) or shRNA targeting L1 (tgctgttgacagtgagcgCTAAACAGGTGATGGAAATGATAGTGAAGCCACAGATGTATCA TTTCCATCACCTGTTTAGtgcctactgcctcgga), together with GFP or mCherry reporter genes, were kindly provided by Dr. Antonio Morales-Hernandez (University of Michigan). Lentivirus was prepared using a three-plasmid system: (1) shRNA transfer vector-, (2) Gag/Pol- + Rev/Tat- (pCMV), and (3) envelope plasmid (pMD2.G) by cotransfection of HEK293T cells using jetPRIME® (OZYME, POL101000046). Supernatant was collected 48

h later, cleared, titred onto HEK293T cells and stored at −80 °C. 50μL of virus were used to transduce 2 million Lin- cells and incubated for 72h before sorting of GFP+ or mCherry+ HSCs.

CFU assays

350 HSCs (Lin-, Sca+, c-Kit+,CD34-, Flk2-) transduced with sh-L1 or sh-control were treated or not with 10μg/mL LPS-B5 Ultrapure (Invivogen, tlrl-pb5lps)LPS for 48h in liquid culture before seeding in 24-well plates containing 500μl of methylcellulose (M3434, StemCell), supplemented with 1% P/S (ThermoFischer Scientific). Colonies were counted after 7 days.

CUT&Tag

CUT&Tag-IT assay kit (Active Motif) was used on 3,000 HSCs according to manufacturer's instructions. Cells were incubated O/N with 0.5μg of H3K9me3 (C15410193-Diagenode). Transposed DNA fragments were amplified 18-fold by PCR using adapters supplied with the kit. PCR purification was carried out using the same Kit to remove remaining primers and large fragments. The quantity and quality of the libraries were assessed on Agilent 2100 Bioanalyzer (Agilent Technologies 50567-4626). Sequencing of the libraries was performed on the NovaSeq-6000 at Gustave Roussy (Illumina; 50 bp paired-end reads).

CUT&Tag Analysis

Quality control was performed using FastQC (v0.11.9) and MultiQC (v1.11). Reads were aligned to the mm10 (UCSC genome) reference using Bowtie2 (v2.4.1) with the following parameters: --end-to-end --very-sensitive --no-mixed --no-discordant --phred33 -I 10 -X 700. Duplicate reads were removed with Picard (v2.26.9) using the parameters --REMOVE_DUPLICATES true --VALIDATION_STRINGENCY LENIENT. Aligned read

quality scores were set to 0, retaining all reads using Samtools (v1.13) with the parameter -q 0. Aligned reads were sorted and indexed with Samtools (v1.13). A coverage track (bigWig) was generated using deeptools (v3.5.0) with parameters -bs 5 --normalizeUsing BPM. All bam and bigwig files were generated by downsampling of each sample to that of the sample with the lowest number of reads

The Bioconductor package Diffbind (Ross-Innes et al., 2012) (v3.8.4) was employed in R v4.2.3 to quantify H3K9me3 levels at TEs, using a reconstructed version of the RepeatMasker annotation which allows the assembly of different fragments of an element (Bailly-Bechet et al., 2014; Walter et al., 2016). Paired-end mode was enabled during the read counting step using the SummarizeOverlaps method. The default mapping quality threshold (mapQCth) was adjusted to 0 for multimapping analysis. Differential binding affinity was assessed using the DBA_DESEQ2 method. The normalized H3K9me3 concentration across all TE loci within the same family or subfamily was aggregated to calculate the total H3K9me3 concentration per TE family. Differential binding at peaks was determined using a P-value threshold of 0.05.

Correlation plots of TE age and H3K9me3 concentration was plotted with ggplot2 (v3.5.1) in R (v4.2.3), assuming a genomic substitution rate of 1.1% per million years (Sookdeo et al., 2013) and summing the mean H3K9me3 concentration at TEs from the same subfamily.

A coverage track (bigWig) was generated using deeptools' bamCoverage with the following parameters: -bs 5 -normalizeUsing BPM. Scores over different genomic regions were calculated using deeptools' computeMatrix with the following parameters --beforeRegionStartLength 1000 --afterRegionStartLength 1000 --regionBodyLength 2500 missingDataAsZero. The mean density profile for 2-3 samples was then plotted with the SEM using the R package collection tidyverse (Wickham et al., 2019). Statistics of the signal were calculated using the R package Rseb (Gregoricchio et al., 2022) and the Wilcoxon test.

RNA-seq

HSCs from individual mice were lysed in Tri-Reagent (Zymo Research) and stored at −80°C until used. Total RNA was extracted using the Direct-Zol RNA microprep kit (Zymo research). All samples were subjected to SMARTer® Stranded Total RNA-Seq Kit v3 - Pico Input Mammalian (Takara Bio USA, Inc., 634485) following the manufacturer's instructions. The fragmentation time was adjusted according to RNA quality, after evaluation on the Agilent Bioanalyzer. Libraries were pooled and sequenced (2 x 100bp) on the NovaSeq6000 (Illumina).

RNA-seq analysis

Quality control was performed using FastQC (v0.11.9) and MultiQC (v1.11). UMI was extracted using umi_tools (v1.1.2). Adapters were removed with Trimgalor (v0.6.5). The FASTQ file was filtered for rRNA sequences using SortMeRNA (v3.4.6). Reads were aligned to the GRCm38 mouse genome reference using STAR (v2.7.5a) and gene quantification was performed using salmon (v1.6.0). Differential expression analysis was performed using DESeq2 R package with P-value <0.05.

For pathway enrichment analysis and data visualization, we performed Gene Set Enrichment Analysis (GSEA) using the GSEA command line (v 4.3.3, Broad Institute) (Subramanian et al., 2005). Normalized counts from the DESeq2 differential expression analysis were used as input. Enrichment analysis was performed against the mm10 Hallmark gene sets obtained from the Molecular Signatures Database (MSigDB) (Castanza et al., 2023). Default parameters were used, except for the permutation type which was set to "gene_set" and metric set to Diff_of_Classes.

Enrichment scores and statistical significance were calculated using 1000 gene set permutations. Results were considered significant at a false discovery rate (FDR) of less than 0.05. GSEA results were plotted and visualized using the R package tidyverse.

TETranscripts

To allow sufficient number of multireads during alignment for TE analysis, --outFilterMultimapNmax and --winAnchorMultimapNmax were set to 100. TE quantification was performed using TETranscripts (v2.2.3) with the parameter mode set to multi. Differential expression analysis was performed using the DESeq2 R package with P-value <0.05.

CUT&Tag and RNA-seq analysis were performed on the Core Cluster of the Institut Français de Bioinformatique (IFB) (ANR-11-INBS-0013).

Statistical analysis

Results were statistically evaluated using GraphPad Prism version 9.0 software (GraphPad Software Inc.). The results are displayed as the means and SEM. The value of *, $P < 0.05$ was considered as significant, and **, $P < 0.01$ or ***, $P < 0.001$ as highly significant.

FIGURE LEGENDS

Figure 1. Chronic inflammation leads to decreased H3K9me3 heterochromatin in HSCs.

(A) Schematic of the chronic inflammation model used in this study. Young mice were injected with either PBS (control-CTRL) or a low dose of LPS (6 μg) every 48 h over 4 weeks followed by immunofluorescence (IF), CUT&Tag or RNA-seq analysis of HSC. **(B-C)**

Immunofluorescence analysis of H3K9me3 (B) and total histone H3 (C) in HSCs isolated from control or LPS-treated mice (minimum 9 animals per group). Representative images (left), and quantitation of fluorescent signal (right, minimum of 250 cells). **(D)** Mice were treated with PBS (control, n=3) or LPS (n=4) as in (A) and HSC were isolated at 8 weeks after completion of treatment for analysis of H3K9me3 (*left panel*) and total histone H3 (*right panel*) by IF. **(E)** IF analysis of H3K9me3 in HSC (left panel), and short-term (ST)-HSC (right panel) isolated from mice heterozygous (Db/+) or homozygous (Db/Db) for mutation of the *LepR* gene (3-6 animals per group). **(F)** (*Top*) Schematic of the experiment: Equal numbers of WT (CD45.1[+]) and Tlr4[-/-] (CD45.2[+]) bone marrow cells were engrafted into lethally irradiated hosts. At 12 weeks post engraftment, mice were treated with PBS (CTRL, 3 mice) or LPS (4 mice) for 4 weeks as in (A), and HSCs were isolated for analysis of H3K9me3 by IF (*bottom*). For all quantitation of immunofluorescence, signal has been normalized to the mean value observed in the control group. Internal box plots indicate Q1, median and Q3 of each distribution. (**** $p<0.0001$; *** $p<0.001$; ** $p<0.01$; * $p<0.05$; ns indicates not significant; Mann Whitney).

Figure 2. Chronic LPS treatment decreases H3K9me3 at the most recent subfamilies of LINEs and LTR in HSC. (A) Volcano plot showing changes in H3K9me3 enrichment at TE loci in HSC between WT-LPS and WT conditions. TEs presenting a significant (*p<0.05)* increase (red) or decrease (green) of H3K9me3 enrichment which are analysed further in (B-D) are indicated. **(B)** Violin plot representing the distribution of H3K9me3 concentration at each TE locus retrieved in (A). **(C)** Repartition of the % of each family of TEs among the total TE loci in the mouse mm10 genome (*top*) and among the TEs retrieved in (A) presenting a significant decrease (*middle* - loss) or increase (*bottom* – gain) of H3K9me3. Hypergeometric test ***p<0.001, ****p<0.0001. **(D)** Violin plot representing the distribution

of H3K9me3 concentration at each locus retrieved in (A) for LINE, LTR, SINE and DNA families of TEs in WT and WT-LPS conditions. Wilcoxon test. **p<0.01; ****p<0.0001. **(E-F)** Correlation plot representing H3K9me3 concentration quantified at all LINE (E) or LTR (F) elements *vs* their age in million years (My) in WT and WT-LPS conditions. R, Pearson correlation coefficient; p, pvalue. **(G-H)** Plot profile of the mean +/- SEM H3K9me3 enrichment along the L1Md (G) or IAP (H) sequences +/- 1kb flanking regions in WT-LPS (orange) *vs* WT (blue) conditions (2-3 samples per condition).

Figure 3. LPS induces a gain of H3K9me3 at L1Mds in *Tet2*^{-/-} HSCs. (A) Volcano plot showing changes in H3K9me3 enrichment at TE loci in HSC between *Tet2*^{-/-} mice +/- LPS. TEs presenting a significant (*p<0.05)* increase (red) or decrease (green) enrichment of H3K9me3 following LPS treatment (analysed further in (B-D)) are indicated. **(B)** Violin plot representing the distribution of H3K9me3 concentration at each TE locus retrieved in (A). **(C)** Repartition of the % of each family of TEs among the total TE loci in the mouse mm10 genome (*top*) and among the TE retrieved in (A) presenting a significant decrease (*middle -* loss) or increase (*bottom –* gain) of H3K9me3. Hypergeometric test **p<0.01; ***p<0.001 **(D)** Violin plots representing the distribution of H3K9me3 concentration at each locus retrieved in (A) for LINE, LTR, SINE and DNA families of TEs in *Tet2*^{-/-}-LPS and *Tet2*^{-/-} conditions. Wilcoxon test. **p<0.01; **(E)** Plot profile of the mean +/- SEM H3K9me3 enrichment along L1Md sequences +/- 1kb flanking regions in *Tet2*^{-/-}-LPS (red) *vs* *Tet2*^{-/-} (green) conditions (2 samples per condition).

Figure 4. LPS induces differential transcriptomic responses in WT and *Tet2*^{-/-} HSCs. (A) GSEA analysis showing Hallmark gene sets significantly differentially enriched (padj<0.05 and -1.5<Normalized enrichment score (NES) or NES>1.5) in WT-LPS vs WT (left) and/or in

Tet2⁻ᐟ⁻-LPS vs *Tet2⁻ᐟ⁻* (right) comparisons. **(B-E)** Enrichment plots for E2F TARGETS (B-C) and TNF-α signalling via NF-κB (D-E) hallmark genesets in WT-LPS vs WT (B,D) and *Tet2⁻ᐟ⁻*-LPS vs *Tet2⁻ᐟ⁻* (C,E) conditions. NES: normalized enrichment score. **(F-G)** violin plot representing TE subfamilies expression for each TE subfamily significantly (p<0.05) differentially expressed in WT-LPS vs WT (F) and *Tet2⁻ᐟ⁻*-LPS vs *Tet2⁻ᐟ⁻* (G) conditions.

Figure 5. TE derepression upon LPS treatment in WT HSC is associated with the accumulation of DNA damage and loss of HSC function. (A-B) Representative images (A) and quantification (B) of γH2AX foci in HSCs sorted from WT mice treated or not with LPS in vivo. n=2 independent experiments, pools of 2 mice for each experiment, 38 to 55 cells counted per slide. 2 way ANOVA with Sidak's multiple comparison test *p<0.05 for 0-1-2 foci per cell. **(C-D)** Representative images (C) and quantification (D) of γH2AX foci in HSCs +/- 10μg/ml LPS and +/- 10μM lamivudine after 48h of culture in vitro. n=2 independent experiments, 3 pools of 2 mice for each experiment, 40-42 cells counted per slide. 1-way ANOVA with Tukey's multiple comparison test. **** p<0.0001 **,p<0.01, ns: non-significant **(E)** Experimental design. Lineage – (Lin-) cells were transduced with either an sh-control or an sh-L1 and mcherry+ HSCs were sorted and plated in methylcellulose in the presence or absence of LPS 3 days after transduction. Figure created with biorender **(F)** Number of colonies formed from 5000 HSCs normalized to untreated cells. N=3 independent experiments. *p<0.05 1-way ANOVA with Tukey's multiple comparison test.

Figure 6. *Tet2⁻ᐟ⁻* HSCs are resistant to the deleterious effects of LPS. (A-C) representative images (A) and quantification (B-C) of γH2AX foci in WT or *Tet2⁻ᐟ⁻* HSCs +/- 10μg/ml LPS in vitro (B) and after PBS or LPS treatment in vivo (C). n=3 independent experiments, 1 pool of 2 mice in each, 38 to 52 cells counted per slide. 2-way ANOVA with Tukey's multiple

comparisons test. *p<0.05; **p<0.01, ***p<0.001 for 0-1-2 foci comparisons; #p<0.05, ##p<0.01 for 3-4-5-6 foci comparison. **(D)** Experimental design for testing the response of WT or *Tet2*$^{-/-}$ cells to LPS treatment in chimeric mice in vivo. BMMC : Bone marrow mononuclear cells. **(E-F)** % CD45.2 in LSK (E) and HSCs (F) and **(G-H)** absolute numbers of LSK (G) and HSCs (H). **(I)** % CD45.2 in blood upon secondary reconstitution. One dot represents a single mouse. n=2 independent experiments *p<0.05; **p<0.01, ***p<0.001, ****p<0.0001 1-way ANOVA with Tukey's multiple comparison test; #p<0.05 unpaired two-tailed t-test. ns = non significant.

Figure 7. Relative expansion of *Tet2*$^{-/-}$ HSCs in response to LPS is dependent on L1 expression in wild-type cells **(A)** Experimental design to analyze *Tet2*$^{-/-}$ vs WT HSCs competition in vitro in the presence or absence of LPS. 70% of WT HSCs transduced with either an sh-control or sh-LINE fused to mCherry reporter gene were cultured in vitro with 30% of *Tet2*$^{-/-}$ HSCs transduced with an sh-control fused to GFP reporter gene, in the presence or absence of LPS. **(B)** Ratio of *Tet2*$^{-/-}$ HSCs (GFP+) between LPS and non-LPS conditions when in competition with WT HSCs (mcherry+) transduced with an sh-control or with an sh-L1. **p<0.01 Unpaired t-test. **(C) Model.** Upon chronic inflammatory stress, the function of WT HSCs is altered through loss of H3K9me3 at L1, L1 overexpression and L1-induced accumulation of DNA damage. *Tet2*$^{-/-}$ are resistant to L1-induced deleterious effects upon LPS treatment through epigenetic repression, eventually leading to their clonal expansion. Figure created with biorender.

SUPPLEMENTAL MATERIAL

Figure S1 (related to figure 2). (A-D) Analysis of an independent H3K9me3 CUT&Tag experiment in HSC of WT-LPS and WT. **(A)** Volcano plot of the TE differentially enriched in H3K9me3 between WT-LPS and WT conditions. The horizontal axis represents the − log10(pValue) and the vertical axis the log2 fold change (FC). TE presenting a significant (*p<0.05)* increase (red) or decrease (green) enrichment in H3K9me3 are shown. **(B)** Violin plot representing the distribution of H3K9me3 concentration at each TE locus retrieved in (A). **(C)** Repartition of the % of each family of TE among the total TE loci in the mouse mm10 genome (*top*) and among the TE retrieved in (A) presenting a significant decrease (*middle* - loss) or increase (*bottom* − gain) of H3K9me3. Hypergeometric test. ***p<0.01, ****p<0.0001 **(D)** Violin plot representing the distribution of H3K9me3 concentration at each locus retrieved in (A) for LINE, LTR, SINE and DNA families of TEs in WT and WT-LPS conditions. Wilcoxon test. ****p<0.0001.

Figure S2 (related to figure 2). (A-B) Correlation plot representing H3K9me3 concentration quantified at differentially enriched LINE (A) or LTR (B) elements retrieved in Figure 2A *vs* their age in million years (My) in WT and WT-LPS conditions. R, Pearson correlation coefficient; p, pvalue. **(C-J)** Plot profiles of the mean +/- SEM H3K9me3 enrichment along L1MdA (C) and IAPEz (D) sequences, intact (E) vs degenerate (F) L1Md, intact (G) vs degenerate (H) IAP, (intact meaning >97% conservation as previously defined (Bulut-Karslioglu et al., 2014)), and >5kb (I) and <5kb (J) L1Md sequences +/- 1kb flanking regions in WT-LPS (orange) *vs* WT (blue) conditions. (2-3 samples per condition)

Figure S3 (related to figure 3). **(A-D)** Plot profile of the mean +/- SEM H3K9me3

enrichment along intact (A) or degenerate (B) and >5kb (C) or <5kb (D) L1Md sequences +/-1kb flanking regions in *Tet2*[-/-]-LPS (red) and *Tet2*[-/-] (green) conditions (2 samples per condition).

Figure S4 (related to Figure 4): (A-B) Volcano plot of gene expression changes between WT-LPS and WT (A) and between *Tet2*[-/-]-LPS and *Tet2*[-/-] (B) conditions. The vertical axis represents the −log10(pValue) and the horizontal axis the log2 fold change (FC). Genes presenting a significant (*p<0.05, 0.9<FC<1.1)* increase (red) or decrease (green) expression are shown. **(C-D)** Venn diagram of DEGs (*p<0.05, 0.9<FC<1.1*) in observed between WT-LPS vs WT and *Tet2*[-/-]-LPS vs *Tet2*[-/-]comparisons. In (C), all DEGs are shown, regardless of the direction of change and in (D), the genes have been separated according to the direction of change observed upon LPS treatment, with colours indicating the number of genes inversely (red) or commonly (green) deregulated in WT and *Tet2*[-/-] HSCs. **(E-F)** violin plot representing L1 subfamilies expression for each L1 subfamily significantly (p<0.05) differentially expressed in WT-LPS vs WT (E) and *Tet2*[-/-]-LPS vs *Tet2*[-/-] (F) conditions.

Figure S5 (related to Figure 6). (A) Experimental design for reconstitution experiments using BM from mice treated with PBS (WT) or LPS (WT-LPS) every two days for 30 days; BMMC, bone marrow mononuclear cells. **(B)** Percentage of CD45.2 donor contribution in blood in mice transplanted with WT or WT-LPS cells 5 months after reconstitution. **(C-D)** Percentage of LSK CD45.2+ (C) or LSK CD34- Flk2- CD45.2+ (HSC) (D) donor contribution in BM 5 months after reconstitution. 5 mice per group. Mann Whitney test *p<0.05. **(E)** Percentage of CD45.2 cells in blood after LPS treatment in mice engrafted with 10% WT or *Tet2*[-/-] BM cells 3 months after the end of the treatment. 4-5 mice per group in 2 independent experiments. One-way ANOVA with Tukey's multiple comparison test.

Figure S6 (related to Figure 7).

(A) Gating strategy for studying sh-control or sh-LINE transduced HSCs. **(B-C)** Single cell liquid cultures. Cell number distribution per well (B), and proliferation index (C) of WT or *Tet2^-/-* HSCs treated or not with LPS at day 7. n=4 independent experiments. Two-way ANOVA with tukey's multiple comparison test *p<0.05, **p<0.01 for the >31 cells group.

Acknowledgments

We thank animal facility, the Genomic and the Imaging and Cytometry Platforms of Gustave Roussy for CUT&Tag and RNA sequencing and cell sorting and confocal analysis, respectively. We also thank S. Duchez and members of the Platforme Technologique of the Institut de Recherche Saint Louis. We thank O. Bernard (INSERM U1170, Gustave Roussy) for providing *Tet2^-/-* mice, C. Muchardt (Institut de biologie paris Seine) for providing *Tlr4^-/-* mice, A. Morales-Hernandez (University of Michigan, US) for providing sh-L1, C. Marty (INSERM U1287, Gustave Roussy) for her advice on cell transduction and S. Lotersztajn for advice and fruitful discussion concerning the Db/Db mouse model. We also thank A. Teissandier and D. Bourc'his for providing us the reconstructed repeatMasker database. This work was supported by INSERM and grants from Ligue National Contre le Cancer (LNCC, Equipe labellisée EL2020) and ARC Foundation (n° ARCPGA2023110007352_7985) to F. P. and Agence National de la Recherche (ANR-15-CE14-0003 to M.G., ANR-23-CE14-0017 to F.P., and ANRJCJC20-CE14-0018-01 to E.E.M), and ARC Foundation (ARCPJA2022070005354), Fondation de France (n° 00135490/ PR-151961) and EHA kick off grant (EHA KOG 2022) to E.E.M.

A.S. is recipient of a fellowship from LNCC, C.X. is recipient of a fellowship from the Chinese Scientific Council, C.F. is supported by ANR (ANR-15-CE14-0003 to M.G. and C.M.) and A.P is a recipient from the Ministère de l'Enseignement Supérieur de la Recherche

et de l'Innovation. M.B is supported by INCA (PLBIO N°2020-095 to F.P.) and ANR (ANR-23-CE14-0017 to F.P.), R.C is supported by ANR (ANRJCJC20-CE14-0018-01 to E.E.M) and ANR (ANR-23-CE14-0017 to F.P.), H.A.O is supported by ARC foundation (n° ARCPGA2023110007352_7985 to F. P.), and M.Y is supported by ANR (ANR-23-CE14-0017 to F.P.).

The authors declare no conflict of interest.

Author contributions

A.S., C.X., C.F., W.Z., K.I , C. Ma and A.P. performed cytometry, IF experiments and analysed the results. A.S. performed CUT&Tag and RNA-seq experiments. A.S., C.F., E.N. S.EC and E.E.M performed reconstitution experiments and analyzed the results. A.S. performed in vitro competition assays and analyzed the results. M.B. and C.X. performed CFU experiments and analyzed the results. R.C., D.F., H.A.O, M.Y, E.Z. and E.E.M. performed bioinformatic analyses. M.G. designed and supervised the initial studies in WT HSCs, and analyzed the results, D.G., F.P. and E.E.M. designed and supervised the study, analyzed the results and wrote the manuscript. M.G., C.M., F.P., and E.E.M. contributed to funding acquisition.

REFERENCES

Abegunde, S.O., R. Buckstein, R.A. Wells, and M.J. Rauh. 2018. An inflammatory environment containing TNFα favors Tet2-mutant clonal hematopoiesis. *Exp Hematol*. 59:60–65. doi:10.1016/j.exphem.2017.11.002.

Avagyan, S., J.E. Henninger, W.P. Mannherz, M. Mistry, J. Yoon, S. Yang, M.C. Weber, J.L. Moore, and L.I. Zon. 2021. Resistance to inflammation underlies enhanced fitness in clonal hematopoiesis. *Science*. 374:768–772. doi:10.1126/science.aba9304.

Bailly-Bechet, M., A. Haudry, and E. Lerat. 2014. "One code to find them all": a perl tool to

conveniently parse RepeatMasker output files. *Mobile DNA*. 5:13. doi:10.1186/1759-8753-5-13.

Barau, J., A. Teissandier, N. Zamudio, S. Roy, V. Nalesso, Y. Hérault, F. Guillou, and D. Bourc'his. 2016. The DNA methyltransferase DNMT3C protects male germ cells from transposon activity. *Science*. 354:909–912. doi:10.1126/science.aah5143.

Barbieri, D., E. Elvira-Matelot, Y. Pelinski, L. Genève, B. de Laval, G. Yogarajah, C. Pecquet, S.N. Constantinescu, and F. Porteu. 2018. Thrombopoietin protects hematopoietic stem cells from retrotransposon-mediated damage by promoting an antiviral response. *Journal of Experimental Medicine*. 215:1463–1480. doi:10.1084/jem.20170997.

Belancio, V.P., A.M. Roy-Engel, R.R. Pochampally, and P. Deininger. 2010. Somatic expression of LINE-1 elements in human tissues. *Nucleic Acids Research*. 38:3909–3922. doi:10.1093/nar/gkq132.

Bick, A.G., J.S. Weinstock, S.K. Nandakumar, C.P. Fulco, E.L. Bao, S.M. Zekavat, M.D. Szeto, X. Liao, M.J. Leventhal, J. Nasser, K. Chang, C. Laurie, B.B. Burugula, C.J. Gibson, A. Niroula, A.E. Lin, M.A. Taub, F. Aguet, K. Ardlie, B.D. Mitchell, K.C. Barnes, A. Moscati, M. Fornage, S. Redline, B.M. Psaty, E.K. Silverman, S.T. Weiss, N.D. Palmer, R.S. Vasan, E.G. Burchard, S.L.R. Kardia, J. He, R.C. Kaplan, N.L. Smith, D.K. Arnett, D.A. Schwartz, A. Correa, M. de Andrade, X. Guo, B.A. Konkle, B. Custer, J.M. Peralta, H. Gui, D.A. Meyers, S.T. McGarvey, I.Y.-D. Chen, M.B. Shoemaker, P.A. Peyser, J.G. Broome, S.M. Gogarten, F.F. Wang, Q. Wong, M.E. Montasser, M. Daya, E.E. Kenny, K.E. North, L.J. Launer, B.E. Cade, J.C. Bis, M.H. Cho, J. Lasky-Su, D.W. Bowden, L.A. Cupples, A.C.Y. Mak, L.C. Becker, J.A. Smith, T.N. Kelly, S. Aslibekyan, S.R. Heckbert, H.K. Tiwari, I.V. Yang, J.A. Heit, S.A. Lubitz, J.M. Johnsen, J.E. Curran, S.E. Wenzel, D.E. Weeks, D.C. Rao, D. Darbar, J.-Y. Moon, R.P. Tracy, E.J. Buth, N. Rafaels, R.J.F. Loos, P. Durda, Y. Liu, L. Hou, J. Lee, P. Kachroo, B.I. Freedman, D. Levy, L.F. Bielak, J.E. Hixson, J.S. Floyd, E.A. Whitsel, P.T. Ellinor, M.R. Irvin, T.E. Fingerlin, et al. 2020. Inherited causes of clonal haematopoiesis in 97,691 whole genomes. *Nature*. 586:763–768. doi:10.1038/s41586-020-2819-2.

Bolton, K.L., R.N. Ptashkin, T. Gao, L. Braunstein, S.M. Devlin, D. Kelly, M. Patel, A. Berthon, A. Syed, M. Yabe, C.C. Coombs, N.M. Caltabellotta, M. Walsh, K. Offit, Z. Stadler, D. Mandelker, J. Schulman, A. Patel, J. Philip, E. Bernard, G. Gundem, J.E.A. Ossa, M. Levine, J.S.M. Martinez, N. Farnoud, D. Glodzik, S. Li, M.E. Robson, C. Lee, P.D.P. Pharoah, K.H. Stopsack, B. Spitzer, S. Mantha, J. Fagin, L. Boucai, C.J. Gibson, B.L. Ebert, A.L. Young, T. Druley, K. Takahashi, N. Gillis, M. Ball, E. Padron, D.M. Hyman, J. Baselga, L. Norton, S. Gardos, V.M. Klimek, H. Scher, D. Bajorin, E. Paraiso, R. Benayed, M.E. Arcila, M. Ladanyi, D.B. Solit, M.F. Berger, M. Tallman, M. Garcia-Closas, N. Chatterjee, L.A. Diaz, R.L. Levine, L.M. Morton, A. Zehir, and E. Papaemmanuil. 2020. Cancer therapy shapes the fitness landscape of clonal hematopoiesis. *Nat Genet*. 52:1219–1226. doi:10.1038/s41588-020-00710-0.

Bowman, R.L., L. Busque, and R.L. Levine. 2018. Clonal Hematopoiesis and Evolution to Hematopoietic Malignancies. *Cell Stem Cell*. 22:157–170. doi:10.1016/j.stem.2018.01.011.

Brojakowska, A., A. Kour, M.C. Thel, E. Park, M. Bisserier, V.N.S. Garikipati, L. Hadri, P.J. Mills, K. Walsh, and D.A. Goukassian. 2022. Retrospective analysis of somatic mutations and

clonal hematopoiesis in astronauts. *Commun Biol*. 5:828. doi:10.1038/s42003-022-03777-z.

Bulut-Karslioglu, A., I.A. De La Rosa-Velázquez, F. Ramirez, M. Barenboim, M. Onishi-Seebacher, J. Arand, C. Galán, G.E. Winter, B. Engist, B. Gerle, R.J. O'Sullivan, J.H.A. Martens, J. Walter, T. Manke, M. Lachner, and T. Jenuwein. 2014. Suv39h-dependent H3K9me3 marks intact retrotransposons and silences LINE elements in mouse embryonic stem cells. *Mol Cell*. 55:277–290. doi:10.1016/j.molcel.2014.05.029.

Busque, L., J.P. Patel, M.E. Figueroa, A. Vasanthakumar, S. Provost, Z. Hamilou, L. Mollica, J. Li, A. Viale, A. Heguy, M. Hassimi, N. Socci, P.K. Bhatt, M. Gonen, C.E. Mason, A. Melnick, L.A. Godley, C.W. Brennan, O. Abdel-Wahab, and R.L. Levine. 2012. Recurrent somatic TET2 mutations in normal elderly individuals with clonal hematopoiesis. *Nat Genet*. 44:1179–1181. doi:10.1038/ng.2413.

Cai, Z., J.J. Kotzin, B. Ramdas, S. Chen, S. Nelanuthala, L.R. Palam, R. Pandey, R.S. Mali, Y. Liu, M.R. Kelley, G. Sandusky, M. Mohseni, A. Williams, J. Henao-Mejia, and R. Kapur. 2018. Inhibition of Inflammatory Signaling in Tet2 Mutant Preleukemic Cells Mitigates Stress-Induced Abnormalities and Clonal Hematopoiesis. *Cell Stem Cell*. 23:833-849.e5. doi:10.1016/j.stem.2018.10.013.

Caiado, F., L.V. Kovtonyuk, N.G. Gonullu, J. Fullin, S. Boettcher, and M.G. Manz. 2023. Aging drives *Tet2* +/− clonal hematopoiesis via IL-1 signaling. *Blood*. 141:886–903. doi:10.1182/blood.2022016835.

Castanza, A.S., J.M. Recla, D. Eby, H. Thorvaldsdóttir, C.J. Bult, and J.P. Mesirov. 2023. Extending support for mouse data in the Molecular Signatures Database (MSigDB). *Nat Methods*. 20:1619–1620. doi:10.1038/s41592-023-02014-7.

Chua, S.C., W.K. Chung, X.S. Wu-Peng, Y. Zhang, S.M. Liu, L. Tartaglia, and R.L. Leibel. 1996. Phenotypes of mouse diabetes and rat fatty due to mutations in the OB (leptin) receptor. *Science*. 271:994–996. doi:10.1126/science.271.5251.994.

Chuong, E.B., N.C. Elde, and C. Feschotte. 2017. Regulatory activities of transposable elements: from conflicts to benefits. *Nat Rev Genet*. 18:71–86. doi:10.1038/nrg.2016.139.

Clapes, T., A. Polyzou, P. Prater, null Sagar, A. Morales-Hernández, M.G. Ferrarini, N. Kehrer, S. Lefkopoulos, V. Bergo, B. Hummel, N. Obier, D. Maticzka, A. Bridgeman, J.S. Herman, I. Ilik, L. Klaeylé, J. Rehwinkel, S. McKinney-Freeman, R. Backofen, A. Akhtar, N. Cabezas-Wallscheid, R. Sawarkar, R. Rebollo, D. Grün, and E. Trompouki. 2021. Chemotherapy-induced transposable elements activate MDA5 to enhance haematopoietic regeneration. *Nat Cell Biol*. 23:704–717. doi:10.1038/s41556-021-00707-9.

Cobo, I., T. Tanaka, C.K. Glass, and C. Yeang. 2022. Clonal hematopoiesis driven by DNMT3A and TET2 mutations: role in monocyte and macrophage biology and atherosclerotic cardiovascular disease. *Current Opinion in Hematology*. 29:1–7. doi:10.1097/MOH.0000000000000688.

Colombo, A.R., A. Zubair, D. Thiagarajan, S. Nuzhdin, T.J. Triche, and G. Ramsingh. 2017. Suppression of Transposable Elements in Leukemic Stem Cells. *Sci Rep*. 7:7029. doi:10.1038/s41598-017-07356-9.

Cook, E.K., M. Luo, and M.J. Rauh. 2020. Clonal hematopoiesis and inflammation: Partners in leukemogenesis and comorbidity. *Experimental Hematology*. 83:85–94. doi:10.1016/j.exphem.2020.01.011.

Coombs, C.C., A. Zehir, S.M. Devlin, A. Kishtagari, A. Syed, P. Jonsson, D.M. Hyman, D.B. Solit, M.E. Robson, J. Baselga, M.E. Arcila, M. Ladanyi, M.S. Tallman, R.L. Levine, and M.F. Berger. 2017. Therapy-Related Clonal Hematopoiesis in Patients with Non-hematologic Cancers Is Common and Associated with Adverse Clinical Outcomes. *Cell Stem Cell*. 21:374-382.e4. doi:10.1016/j.stem.2017.07.010.

Cuellar, T.L., A.-M. Herzner, X. Zhang, Y. Goyal, C. Watanabe, B.A. Friedman, V. Janakiraman, S. Durinck, J. Stinson, D. Arnott, T.K. Cheung, S. Chaudhuri, Z. Modrusan, J.M. Doerr, M. Classon, and B. Haley. 2017. ---Silencing of retrotransposons by SETDB1 inhibits the interferon response in acute myeloid leukemia--. *Journal of Cell Biology*. 216:3535–3549. doi:10.1083/jcb.201612160.

De Cecco, M., S.W. Criscione, E.J. Peckham, S. Hillenmeyer, E.A. Hamm, J. Manivannan, A.L. Peterson, J.A. Kreiling, N. Neretti, and J.M. Sedivy. 2013. Genomes of replicatively senescent cells undergo global epigenetic changes leading to gene silencing and activation of transposable elements. *Aging Cell*. 12:247–256. doi:10.1111/acel.12047.

De Cecco, M., T. Ito, A.P. Petrashen, A.E. Elias, N.J. Skvir, S.W. Criscione, A. Caligiana, G. Brocculi, E.M. Adney, J.D. Boeke, O. Le, C. Beauséjour, J. Ambati, K. Ambati, M. Simon, A. Seluanov, V. Gorbunova, P.E. Slagboom, S.L. Helfand, N. Neretti, and J.M. Sedivy. 2019. L1 drives IFN in senescent cells and promotes age-associated inflammation. *Nature*. 566:73–78. doi:10.1038/s41586-018-0784-9.

Desai, P., N. Mencia-Trinchant, O. Savenkov, M.S. Simon, G. Cheang, S. Lee, M. Samuel, E.K. Ritchie, M.L. Guzman, K.V. Ballman, G.J. Roboz, and D.C. Hassane. 2018. Somatic mutations precede acute myeloid leukemia years before diagnosis. *Nat Med*. 24:1015–1023. doi:10.1038/s41591-018-0081-z.

Desai, P., and G.J. Roboz. 2019. Clonal Hematopoiesis and therapy related MDS/AML. *Best Practice & Research Clinical Haematology*. 32:13–23. doi:10.1016/j.beha.2019.02.006.

Djeghloul, D., K. Kuranda, I. Kuzniak, D. Barbieri, I. Naguibneva, C. Choisy, J.-C. Bories, C. Dosquet, M. Pla, V. Vanneaux, G. Socié, F. Porteu, D. Garrick, and M. Goodhardt. 2016. Age-Associated Decrease of the Histone Methyltransferase SUV39H1 in HSC Perturbs Heterochromatin and B Lymphoid Differentiation. *Stem Cell Reports*. 6:970–984. doi:10.1016/j.stemcr.2016.05.007.

Dykstra, B., S. Olthof, J. Schreuder, M. Ritsema, and G. de Haan. 2011. Clonal analysis reveals multiple functional defects of aged murine hematopoietic stem cells. *J Exp Med*. 208:2691–2703. doi:10.1084/jem.20111490.

Esplin, B.L., T. Shimazu, R.S. Welner, K.P. Garrett, L. Nie, Q. Zhang, M.B. Humphrey, Q. Yang, L.A. Borghesi, and P.W. Kincade. 2011. Chronic Exposure to a TLR Ligand Injures Hematopoietic Stem Cells. *J.I.* 186:5367–5375. doi:10.4049/jimmunol.1003438.

Florez, M.A., B.T. Tran, T.K. Wathan, J. DeGregori, E.M. Pietras, and K.Y. King. 2022. Clonal hematopoiesis: Mutation-specific adaptation to environmental change. *Cell Stem Cell*. 29:882–904. doi:10.1016/j.stem.2022.05.006.

Fuster, J.J., S. MacLauchlan, M.A. Zuriaga, M.N. Polackal, A.C. Ostriker, R. Chakraborty, C.-L. Wu, S. Sano, S. Muralidharan, C. Rius, J. Vuong, S. Jacob, V. Muralidhar, A.A.B. Robertson, M.A. Cooper, V. Andrés, K.K. Hirschi, K.A. Martin, and K. Walsh. 2017. Clonal hematopoiesis associated with TET2 deficiency accelerates atherosclerosis development in mice. *Science*. 355:842–847. doi:10.1126/science.aag1381.

Gasior, S.L., T.P. Wakeman, B. Xu, and P.L. Deininger. 2006. The human LINE-1 retrotransposon creates DNA double-strand breaks. *J Mol Biol*. 357:1383–1393. doi:10.1016/j.jmb.2006.01.089.

Gazquez-Gutierrez, A., J. Witteveldt, S. R Heras, and S. Macias. 2021. Sensing of transposable elements by the antiviral innate immune system. *RNA*. rna.078721.121. doi:10.1261/rna.078721.121.

Genovese, G., A.K. Kähler, R.E. Handsaker, J. Lindberg, S.A. Rose, S.F. Bakhoum, K. Chambert, E. Mick, B.M. Neale, M. Fromer, S.M. Purcell, O. Svantesson, M. Landén, M. Höglund, S. Lehmann, S.B. Gabriel, J.L. Moran, E.S. Lander, P.F. Sullivan, P. Sklar, H. Grönberg, C.M. Hultman, and S.A. McCarroll. 2014. Clonal Hematopoiesis and Blood-Cancer Risk Inferred from Blood DNA Sequence. *N Engl J Med*. 371:2477–2487. doi:10.1056/NEJMoa1409405.

Gillis, N.K., M. Ball, Q. Zhang, Z. Ma, Y. Zhao, S.J. Yoder, M.E. Balasis, T.E. Mesa, D.A. Sallman, J.E. Lancet, R.S. Komrokji, A.F. List, H.L. McLeod, M. Alsina, R. Baz, K.H. Shain, D.E. Rollison, and E. Padron. 2017. Clonal haemopoiesis and therapy-related myeloid malignancies in elderly patients: a proof-of-concept, case-control study. *The Lancet Oncology*. 18:112–121. doi:10.1016/S1470-2045(16)30627-1.

Gregoricchio, S., L. Polit, M. Esposito, J. Berthelet, L. Delestré, E. Evanno, M. Diop, I. Gallais, H. Aleth, M. Poplineau, W. Zwart, F. Rosenbauer, F. Rodrigues-Lima, E. Duprez, V. Boeva, and C. Guillouf. 2022. HDAC1 and PRC2 mediate combinatorial control in SPI1/PU.1-dependent gene repression in murine erythroleukaemia. *Nucleic Acids Res*. 50:7938–7958. doi:10.1093/nar/gkac613.

Gu, Z., Y. Liu, Y. Zhang, H. Cao, J. Lyu, X. Wang, A. Wylie, S.J. Newkirk, A.E. Jones, M. Lee, G.A. Botten, M. Deng, K.E. Dickerson, C.C. Zhang, W. An, J.M. Abrams, and J. Xu. 2021. Silencing of LINE-1 retrotransposons is a selective dependency of myeloid leukemia. *Nat Genet*. 53:672–682. doi:10.1038/s41588-021-00829-8.

Hidaoui, D., R. Chelbie, M. Bohm, A. Porquet, A. Polyzou, V. Alcazer, S. Depil, A. Imanci, M. Morabito, A. Renneville, D. selimoglu-Buet, S. Thépot, R. Itzykson, L. Laplane, N. Droin, E. Trompouki, E. Elvira-Matelot, E. Solary, and F. Porteu. 2024. Targeting Heterochromatin Eliminates Malignant Hematopoietic Stem and Progenitor Cells in Chronic Myelomonocytic Leukemia Through Reactivation of Retroelements and Innate Immune pathways. Cancer Biology.

Hong, T., J. Li, L. Guo, M. Cavalier, T. Wang, Y. Dou, A. DeLaFuente, S. Fang, A. Guzman,

K. Wohlan, C. Kapadia, C. Rosas, Y. Yang, C.C. Yin, S. Li, M.J. You, X. Cheng, M.A. Goodell, Y. Zhou, and Y. Huang. 2023. TET2 modulates spatial relocalization of heterochromatin in aged hematopoietic stem and progenitor cells. *Nat Aging*. 3:1387–1400. doi:10.1038/s43587-023-00505-y.

Jaiswal, S., P. Fontanillas, J. Flannick, A. Manning, P.V. Grauman, B.G. Mar, R.C. Lindsley, C.H. Mermel, N. Burtt, A. Chavez, J.M. Higgins, V. Moltchanov, F.C. Kuo, M.J. Kluk, B. Henderson, L. Kinnunen, H.A. Koistinen, C. Ladenvall, G. Getz, A. Correa, B.F. Banahan, S. Gabriel, S. Kathiresan, H.M. Stringham, M.I. McCarthy, M. Boehnke, J. Tuomilehto, C. Haiman, L. Groop, G. Atzmon, J.G. Wilson, D. Neuberg, D. Altshuler, and B.L. Ebert. 2014. Age-Related Clonal Hematopoiesis Associated with Adverse Outcomes. *N Engl J Med*. 371:2488–2498. doi:10.1056/NEJMoa1408617.

Jakobsen, N.A., S. Turkalj, A.G.X. Zeng, B. Stoilova, M. Metzner, S. Rahmig, M.S. Nagree, S. Shah, R. Moore, B. Usukhbayar, M. Angulo Salazar, G.-A. Gafencu, A. Kennedy, S. Newman, B.J.L. Kendrick, A.H. Taylor, R. Afinowi-Luitz, R. Gundle, B. Watkins, K. Wheway, D. Beazley, A. Murison, A.G. Aguilar-Navarro, E. Flores-Figueroa, S.G. Dakin, A.J. Carr, C. Nerlov, J.E. Dick, S.Z. Xie, and P. Vyas. 2024. Selective advantage of mutant stem cells in human clonal hematopoiesis is associated with attenuated response to inflammation and aging. *Cell Stem Cell*. S1934590924002078. doi:10.1016/j.stem.2024.05.010.

Jin, Y., O.H. Tam, E. Paniagua, and M. Hammell. 2015. TEtranscripts: a package for including transposable elements in differential expression analysis of RNA-seq datasets. *Bioinformatics*. 31:3593–3599. doi:10.1093/bioinformatics/btv422.

Kar, S.P., P.M. Quiros, M. Gu, T. Jiang, J. Mitchell, R. Langdon, V. Iyer, C. Barcena, M.S. Vijayabaskar, M.A. Fabre, P. Carter, S. Petrovski, S. Burgess, and G.S. Vassiliou. 2022. Genome-wide analyses of 200,453 individuals yield new insights into the causes and consequences of clonal hematopoiesis. *Nat Genet*. 54:1155–1166. doi:10.1038/s41588-022-01121-z.

Keenan, C.R., N. Iannarella, G. Naselli, N.G. Bediaga, T.M. Johanson, L.C. Harrison, and R.S. Allan. 2020. Extreme disruption of heterochromatin is required for accelerated hematopoietic aging. *Blood*. 135:2049–2058. doi:10.1182/blood.2019002990.

Ko, M., H.S. Bandukwala, J. An, E.D. Lamperti, E.C. Thompson, R. Hastie, A. Tsangaratou, K. Rajewsky, S.B. Koralov, and A. Rao. 2011. Ten-Eleven-Translocation 2 (TET2) negatively regulates homeostasis and differentiation of hematopoietic stem cells in mice. *Proc. Natl. Acad. Sci. U.S.A.* 108:14566–14571. doi:10.1073/pnas.1112317108.

Koide, S., M. Oshima, K. Takubo, S. Yamazaki, E. Nitta, A. Saraya, K. Aoyama, Y. Kato, S. Miyagi, Y. Nakajima-Takagi, T. Chiba, H. Matsui, F. Arai, Y. Suzuki, H. Kimura, H. Nakauchi, T. Suda, Y. Shinkai, and A. Iwama. 2016. Setdb1 maintains hematopoietic stem and progenitor cells by restricting the ectopic activation of nonhematopoietic genes. *Blood*. 128:638–649. doi:10.1182/blood-2016-01-694810.

Lescale, C., S. Dias, J. Maës, A. Cumano, P. Szabo, D. Charron, M.E. Weksler, C. Dosquet, P. Vieira, and M. Goodhardt. 2010. Reduced EBF expression underlies loss of B-cell potential of hematopoietic progenitors with age. *Aging Cell*. 9:410–419. doi:10.1111/j.1474-

9726.2010.00566.x.

Li, Z., X. Cai, C.-L. Cai, J. Wang, W. Zhang, B.E. Petersen, F.-C. Yang, and M. Xu. 2011. Deletion of Tet2 in mice leads to dysregulated hematopoietic stem cells and subsequent development of myeloid malignancies. *Blood*. 118:4509–4518. doi:10.1182/blood-2010-12-325241.

Liang, Y., X. Qu, N.M. Shah, and T. Wang. 2024. Towards targeting transposable elements for cancer therapy. *Nat Rev Cancer*. 24:123–140. doi:10.1038/s41568-023-00653-8.

Liu, N., C.H. Lee, T. Swigut, E. Grow, B. Gu, M.C. Bassik, and J. Wysocka. 2018. Selective silencing of euchromatic L1s revealed by genome-wide screens for L1 regulators. *Nature*. 553:228–232. doi:10.1038/nature25179.

McClatchy, J., R. Strogantsev, E. Wolfe, H.Y. Lin, M. Mohammadhosseini, B.A. Davis, C. Eden, D. Goldman, W.H. Fleming, P. Conley, G. Wu, L. Cimmino, H. Mohammed, and A. Agarwal. 2023. Clonal hematopoiesis related TET2 loss-of-function impedes IL1β-mediated epigenetic reprogramming in hematopoietic stem and progenitor cells. *Nat Commun*. 14:8102. doi:10.1038/s41467-023-43697-y.

Mencia-Trinchant, N., M.J. MacKay, C. Chin, E. Afshinnekoo, J. Foox, C. Meydan, D. Butler, C. Mozsary, N.A. Vernice, C. Darby, M.C. Schatz, S.M. Bailey, A.M. Melnick, M.L. Guzman, K. Bolton, L.Z. Braunstein, F. Garrett-Bakelman, R.L. Levine, D.C. Hassane, and C.E. Mason. 2020. Clonal Hematopoiesis Before, During, and After Human Spaceflight. *Cell Reports*. 33:108458. doi:10.1016/j.celrep.2020.108458.

Mitchell, S.R., J. Gopakumar, and S. Jaiswal. 2021. Insights into clonal hematopoiesis and its relation to cancer risk. *Current Opinion in Genetics & Development*. 66:63–69. doi:10.1016/j.gde.2020.12.004.

Moran-Crusio, K., L. Reavie, A. Shih, O. Abdel-Wahab, D. Ndiaye-Lobry, C. Lobry, M.E. Figueroa, A. Vasanthakumar, J. Patel, X. Zhao, F. Perna, S. Pandey, J. Madzo, C. Song, Q. Dai, C. He, S. Ibrahim, M. Beran, J. Zavadil, S.D. Nimer, A. Melnick, L.A. Godley, I. Aifantis, and R.L. Levine. 2011. Tet2 loss leads to increased hematopoietic stem cell self-renewal and myeloid transformation. *Cancer Cell*. 20:11–24. doi:10.1016/j.ccr.2011.06.001.

Nagareddy, P.R., M. Kraakman, S.L. Masters, R.A. Stirzaker, D.J. Gorman, R.W. Grant, D. Dragoljevic, E.S. Hong, A. Abdel-Latif, S.S. Smyth, S.H. Choi, J. Korner, K.E. Bornfeldt, E.A. Fisher, V.D. Dixit, A.R. Tall, I.J. Goldberg, and A.J. Murphy. 2014. Adipose tissue macrophages promote myelopoiesis and monocytosis in obesity. *Cell Metab*. 19:821–835. doi:10.1016/j.cmet.2014.03.029.

Pelinski, Y., D. Hidaoui, A. Stolz, F. Hermetet, R. Chelbi, M.K. Diop, A.M. Chioukh, F. Porteu, and E. Elvira-Matelot. 2022. NF-κB signaling controls H3K9me3 levels at intronic LINE-1 and hematopoietic stem cell genes in cis. *Journal of Experimental Medicine*. 219:e20211356. doi:c.

Pezic, D., S.A. Manakov, R. Sachidanandam, and A.A. Aravin. 2014. piRNA pathway targets active LINE1 elements to establish the repressive H3K9me3 mark in germ cells. *Genes Dev*. 28:1410–1428. doi:10.1101/gad.240895.114.

Pronier, E., A. Imanci, D. Selimoglu-Buet, B. Badaoui, R. Itzykson, T. Roger, C. Jego, A. Naimo, M. Francillette, M. Breckler, O. Wagner-Ballon, M.E. Figueroa, M. Aglave, D. Gautheret, F. Porteu, O.A. Bernard, W. Vainchenker, F. Delhommeau, E. Solary, and N.M. Droin. 2022. Macrophage migration inhibitory factor is overproduced through EGR1 in TET2low resting monocytes. *Commun Biol*. 5:110. doi:10.1038/s42003-022-03057-w.

Quivoron, C., L. Couronné, V. Della Valle, C.K. Lopez, I. Plo, O. Wagner-Ballon, M. Do Cruzeiro, F. Delhommeau, B. Arnulf, M.-H. Stern, L. Godley, P. Opolon, H. Tilly, E. Solary, Y. Duffourd, P. Dessen, H. Merle-Beral, F. Nguyen-Khac, M. Fontenay, W. Vainchenker, C. Bastard, T. Mercher, and O.A. Bernard. 2011. TET2 Inactivation Results in Pleiotropic Hematopoietic Abnormalities in Mouse and Is a Recurrent Event during Human Lymphomagenesis. *Cancer Cell*. 20:25–38. doi:10.1016/j.ccr.2011.06.003.

Robbez-Masson, L., C.H.C. Tie, L. Conde, H. Tunbak, C. Husovsky, I.A. Tchasovnikarova, R.T. Timms, J. Herrero, P.J. Lehner, and H.M. Rowe. 2018. The HUSH complex cooperates with TRIM28 to repress young retrotransposons and new genes. *Genome Res*. 28:836–845. doi:10.1101/gr.228171.117.

Rodriguez-Meira, A., R. Norfo, S. Wen, A.L. Chédeville, H. Rahman, J. O'Sullivan, G. Wang, E. Louka, W.W. Kretzschmar, A. Paterson, C. Brierley, J.-E. Martin, C. Demeule, M. Bashton, N. Sousos, D. Moralli, L. Subha Meem, J. Carrelha, B. Wu, A. Hamblin, H. Guermouche, F. Pasquier, C. Marzac, F. Girodon, W. Vainchenker, M. Drummond, C. Harrison, J.R. Chapman, I. Plo, S.E.W. Jacobsen, B. Psaila, S. Thongjuea, I. Antony-Debré, and A.J. Mead. 2023. Single-cell multi-omics identifies chronic inflammation as a driver of TP53-mutant leukemic evolution. *Nat Genet*. 55:1531–1541. doi:10.1038/s41588-023-01480-1.

Rossi, D.J., D. Bryder, J.M. Zahn, H. Ahlenius, R. Sonu, A.J. Wagers, and I.L. Weissman. 2005. Cell intrinsic alterations underlie hematopoietic stem cell aging. *Proc Natl Acad Sci U S A*. 102:9194–9199. doi:10.1073/pnas.0503280102.

Ross-Innes, C.S., R. Stark, A.E. Teschendorff, K.A. Holmes, H.R. Ali, M.J. Dunning, G.D. Brown, O. Gojis, I.O. Ellis, A.R. Green, S. Ali, S.-F. Chin, C. Palmieri, C. Caldas, and J.S. Carroll. 2012. Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. *Nature*. 481:389–393. doi:10.1038/nature10730.

Sano, S., K. Oshima, Y. Wang, Y. Katanasaka, M. Sano, and K. Walsh. 2018a. CRISPR-Mediated Gene Editing to Assess the Roles of Tet2 and Dnmt3a in Clonal Hematopoiesis and Cardiovascular Disease. *Circ Res*. 123:335–341. doi:10.1161/CIRCRESAHA.118.313225.

Sano, S., K. Oshima, Y. Wang, S. MacLauchlan, Y. Katanasaka, M. Sano, M.A. Zuriaga, M. Yoshiyama, D. Goukassian, M.A. Cooper, J.J. Fuster, and K. Walsh. 2018b. Tet2-Mediated Clonal Hematopoiesis Accelerates Heart Failure Through a Mechanism Involving the IL-1β/NLRP3 Inflammasome. *J Am Coll Cardiol*. 71:875–886. doi:10.1016/j.jacc.2017.12.037.

Sookdeo, A., C.M. Hepp, M.A. McClure, and S. Boissinot. 2013. Revisiting the evolution of mouse LINE-1 in the genomic era. *Mob DNA*. 4:3. doi:10.1186/1759-8753-4-3.

Speck, N.A. 2022. A Pernicious Cycle Affecting Premalignant Stem Cells. *N Engl J Med*.

386:596–598. doi:10.1056/NEJMcibr2117528.

Subramanian, A., P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette, A. Paulovich, S.L. Pomeroy, T.R. Golub, E.S. Lander, and J.P. Mesirov. 2005. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U.S.A*. 102:15545–15550. doi:10.1073/pnas.0506580102.

Takahashi, K., F. Wang, H. Kantarjian, D. Doss, K. Khanna, E. Thompson, L. Zhao, K. Patel, S. Neelapu, C. Gumbs, C. Bueso-Ramos, C.D. DiNardo, S. Colla, F. Ravandi, J. Zhang, X. Huang, X. Wu, F. Samaniego, G. Garcia-Manero, and P.A. Futreal. 2017. Preleukaemic clonal haemopoiesis and risk of therapy-related myeloid neoplasms: a case-control study. *The Lancet Oncology*. 18:100–111. doi:10.1016/S1470-2045(16)30626-X.

Takizawa, H., K. Fritsch, L.V. Kovtonyuk, Y. Saito, C. Yakkala, K. Jacobs, A.K. Ahuja, M. Lopes, A. Hausmann, W.-D. Hardt, Á. Gomariz, C. Nombela-Arrieta, and M.G. Manz. 2017. Pathogen-Induced TLR4-TRIF Innate Immune Signaling in Hematopoietic Stem Cells Promotes Proliferation but Reduces Competitive Fitness. *Cell Stem Cell*. 21:225-240.e5. doi:10.1016/j.stem.2017.06.013.

Tsurumi, A., and W.X. Li. 2012. Global heterochromatin loss: a unifying theory of aging? *Epigenetics*. 7:680–688. doi:10.4161/epi.20540.

Ugarte, F., R. Sousae, B. Cinquin, E.W. Martin, J. Krietsch, G. Sanchez, M. Inman, H. Tsang, M. Warr, E. Passegué, C.A. Larabell, and E.C. Forsberg. 2015. Progressive Chromatin Condensation and H3K9 Methylation Regulate the Differentiation of Embryonic and Hematopoietic Stem Cells. *Stem Cell Reports*. 5:728–740. doi:10.1016/j.stemcr.2015.09.009.

Walter, M., A. Teissandier, R. Pérez-Palacios, and D. Bourc'his. 2016. An epigenetic switch ensures transposon repression upon dynamic loss of DNA methylation in embryonic stem cells. *Elife*. 5:e11418. doi:10.7554/eLife.11418.

Wickham, H., M. Averick, J. Bryan, W. Chang, L. McGowan, R. François, G. Grolemund, A. Hayes, L. Henry, J. Hester, M. Kuhn, T. Pedersen, E. Miller, S. Bache, K. Müller, J. Ooms, D. Robinson, D. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K. Woo, and H. Yutani. 2019. Welcome to the Tidyverse. *JOSS*. 4:1686. doi:10.21105/joss.01686.

Yeaton, A., G. Cayanan, S. Loghavi, I. Dolgalev, E.M. Leddin, C.E. Loo, H. Torabifard, D. Nicolet, J. Wang, K. Corrigan, V. Paraskevopoulou, D.T. Starczynowski, E. Wang, O. Abdel-Wahab, A.D. Viny, R.M. Stone, J.C. Byrd, O.A. Guryanova, R.M. Kohli, G.A. Cisneros, A. Tsirigos, A.-K. Eisfeld, I. Aifantis, and M. Guillamot. 2022. The Impact of Inflammation-Induced Tumor Plasticity during Myeloid Transformation. *Cancer Discovery*. 12:2392–2413. doi:10.1158/2159-8290.CD-21-1146.

Zhang, Q., K. Zhao, Q. Shen, Y. Han, Y. Gu, X. Li, D. Zhao, Y. Liu, C. Wang, X. Zhang, X. Su, J. Liu, W. Ge, R.L. Levine, N. Li, and X. Cao. 2015. Tet2 is required to resolve inflammation by recruiting Hdac2 to specifically repress IL-6. *Nature*. 525:389–393. doi:10.1038/nature15252.

Zhao, Y., E. Oreskovic, Q. Zhang, Q. Lu, A. Gilman, Y.S. Lin, J. He, Z. Zheng, J.Y. Lu, J. Lee, Z. Ke, J. Ablaeva, M.J. Sweet, S. Horvath, Z. Zhang, E. Nevo, A. Seluanov, and V.

Gorbunova. 2021. Transposon-triggered innate immune response confers cancer resistance to the blind mole rat. *Nat Immunol*. 22:1219–1230. doi:10.1038/s41590-021-01027-8.

Fig 1. Chronic inflammation leads to decreased H3K9me3 heterochromatin mark in HSCS

Figure 1. Chronic inflammation leads to decreased H3K9me3 heterochromatin in HSCs. (A) Schematic of the chronic inflammation model used in this study. Young mice were injected with either PBS (control-CTRL) or a low dose of LPS (6 μg) every 48 h over 4 weeks followed by immunofluorescence (IF), CUT&Tag or RNA-seq analysis of HSC. **(B-C)** Immunofluorescence analysis of H3K9me3 (B) and total histone H3 (C) in HSCs isolated from control or LPS-treated mice (minimum 9 animals per group). Representative images (left), and quantitation of fluorescent signal (right, minimum of 250 cells). **(D)** Mice were treated with PBS (control, n=3) or LPS (n=4) as in (A) and HSC were isolated at 8 weeks after completion of treatment for analysis of H3K9me3 (*left panel*) and total histone H3 (*right panel*) by IF. **(E)** IF analysis of H3K9me3 in HSC (left panel), and short-term (ST)-HSC (right panel) isolated from mice heterozygous (Db/+) or homozygous (Db/Db) for mutation of the *LepR* gene (3-6 animals per group). **(F)** (*Top*) Schematic of the experiment: Equal numbers of WT (CD45.1$^+$) and Tlr4$^{-/-}$ (CD45.2$^+$) bone marrow cells were engrafted into lethally irradiated hosts. At 12 weeks post engraftment, mice were treated with PBS (CTRL, 3 mice) or LPS (4 mice) for 4 weeks as in (A), and HSCs were isolated for analysis of H3K9me3 by IF (*bottom*). For all quantitation of immunofluorescence, signal has been normalized to the mean value observed in the control group. Internal box plots indicate Q1, median and Q3 of each distribution. (**** p<0.0001; *** p<0.001; ** p<0.01; * p<0.05; ns indicates not significant; Mann Whitney).

FIGURE 2

Stolz et al.

Fig 2. Chronic LPS treatment induces a loss of H3K9me3 at the most recent subfamilies of L1s and LTRs

Figure 2. Chronic LPS treatment decreases H3K9me3 at the most recent subfamilies of LINEs and LTR in HSC. (A) Volcano plot showing changes in H3K9me3 enrichment at TE loci in HSC between WT-LPS and WT conditions. TEs presenting a significant (*p<0.05)* increase (red) or decrease (green) of H3K9me3 enrichment which are analysed further in (B-D) are indicated. **(B)** Violin plot representing the distribution of H3K9me3 concentration at each TE locus retrieved in (A). **(C)** Repartition of the % of each family of TEs among the total TE loci in the mouse mm10 genome (*top*) and among the TEs retrieved in (A) presenting a significant decrease (*middle* - loss) or increase (*bottom* – gain) of H3K9me3. Hypergeometric test ***p<0.001, ****p<0.0001. **(D)** Violin plot representing the distribution of H3K9me3 concentration at each locus retrieved in (A) for LINE, LTR, SINE and DNA families of TEs in WT and WT-LPS conditions. Wilcoxon test. **p<0.01; ****p<0.0001. **(E-F)** Correlation plot representing H3K9me3 concentration quantified at all LINE (E) or LTR (F) elements *vs* their age in million years (My) in WT and WT-LPS conditions. R, Pearson correlation coefficient; p, pvalue. **(G-H)** Plot profile of the mean +/- SEM H3K9me3 enrichment along the L1Md (G) or IAP (H) sequences +/- 1kb flanking regions in WT-LPS (orange) *vs* WT (blue) conditions (2-3 samples per condition).

Figure 3. LPS induces a gain of H3K9me3 at L1Mds in *Tet2⁻/⁻* HSCs .

Figure 3. LPS induces a gain of H3K9me3 at L1Mds in *Tet2^{-/-}* HSCs. (A) Volcano plot showing changes in H3K9me3 enrichment at TE loci in HSC between *Tet2^{-/-}* mice +/- LPS. TEs presenting a significant (*p<0.05)* increase (red) or decrease (green) enrichment of H3K9me3 following LPS treatment (analysed further in (B-D)) are indicated. **(B)** Violin plot representing the distribution of H3K9me3 concentration at each TE locus retrieved in (A). **(C)** Repartition of the % of each family of TEs among the total TE loci in the mouse mm10 genome (*top*) and among the TE retrieved in (A) presenting a significant decrease (*middle* - loss) or increase (*bottom* – gain) of H3K9me3. Hypergeometric test **p<0.01; ***p<0.001 **(D)** Violin plots representing the distribution of H3K9me3 concentration at each locus retrieved in (A) for LINE, LTR, SINE and DNA families of TEs in *Tet2^{-/-}*-LPS and *Tet2^{-/-}* conditions. Wilcoxon test. **p<0.01; **(E)** Plot profile of the mean +/- SEM H3K9me3 enrichment along L1Md sequences +/- 1kb flanking regions in *Tet2^{-/-}*-LPS (red) *vs Tet2^{-/-}* (green) conditions (2 samples per condition).

A.

Fig 4. LPS induces differential transcriptomic responses in WT and *Tet2*^-/- HSCs.

Figure 4. LPS induces differential transcriptomic responses in WT and *Tet2*^{-/-} HSCs. (A) GSEA analysis showing Hallmark gene sets significantly differentially enriched (padj<0.05 and -1.5<Normalized enrichment score (NES) or NES>1.5) in WT-LPS vs WT (left) and/or in *Tet2*^{-/-}-LPS vs *Tet2*^{-/-} (right) comparisons. **(B-E)** Enrichment plots for E2F TARGETS (B-C) and TNF-a signalling via NF-kB (D-E) hallmark genesets in WT-LPS vs WT (B,D) and *Tet2*^{-/-}-LPS vs *Tet2*^{-/-} (C,E) conditions. NES: normalized enrichment score. **(F-G)** violin plot representing TE subfamilies expression for each TE subfamily significantly (p<0.05) differentially expressed in WT-LPS vs WT (F) and *Tet2*^{-/-}-LPS vs *Tet2*^{-/-} (G) conditions.

Fig 5. TE derepression upon LPS treatment is associated with the accumulation of DNA damage and loss of HSC function

Figure 5. TE derepression upon LPS treatment in WT HSC is associated with the accumulation of DNA damage and loss of HSC function. (A-B) Representative images (A) and quantification (B) of gH2AX foci in HSCs sorted from WT mice treated or not with LPS in vivo. n=2 independent experiments, pools of 2 mice for each experiment, 38 to 55 cells counted per slide. 2 way ANOVA with Sidak's multiple comparison test *p<0.05 for 0-1-2 foci per cell. **(C-D)** Representative images (C) and quantification (D) of gH2AX foci in HSCs +/- 10µg/ml LPS and +/- 10µM lamivudine after 48h of culture in vitro. n=2 independent experiments, 3 pools of 2 mice for each experiment, 40-42 cells counted per slide. 1-way ANOVA with Tukey's multiple comparison test. **** p<0.0001 **,p<0.01, ns: non-significant **(E)** Experimental design. Lineage – (Lin-) cells were transduced with either an sh-control or an sh-L1 and mcherry+ HSCs were sorted and plated in methylcellulose in the presence or absence of LPS 3 days after transduction. Figure created with biorender **(F)** Number of colonies formed from 5000 HSCs normalized to untreated cells. N=3 independent experiments. *p<0.05 1-way ANOVA with Tukey's multiple comparison test.

Figure 6. *Tet2⁻/⁻* HSCs are resistant to the deleterious effects of LPS

Figure 6. *Tet2*$^{-/-}$ **HSCs are resistant to the deleterious effects of LPS. (A-C)** representative images (A) and quantification (B-C) of gH2AX foci in WT or *Tet2*$^{-/-}$ HSCs +/- 10µg/ml LPS in vitro (B) and after PBS or LPS treatment in vivo (C). n=3 independent experiments, 1 pool of 2 mice in each, 38 to 52 cells counted per slide. 2-way ANOVA with Tukey's multiple comparisons test. *p<0.05; **p<0.01, ***p<0.001 for 0-1-2 foci comparisons; #p<0.05, ##p<0.01 for 3-4-5-6 foci comparison. **(D)** Experimental design for testing the response of WT or *Tet2*$^{-/-}$ cells to LPS treatment in chimeric mice in vivo. BMMC : Bone marrow mononuclear cells. **(E-F)** % CD45.2 in LSK (E) and HSCs (F) and **(G-H)** absolute numbers of LSK (G) and HSCs (H). **(I)** % CD45.2 in blood upon secondary reconstitution. One dot represents a single mouse. n=2 independent experiments *p<0.05; **p<0.01, ***p<0.001, ****p<0.0001 1-way ANOVA with Tukey's multiple comparison test; #p<0.05 unpaired two-tailed t-test. ns = non significant.

A.

70 % WT HSCs
Sh-control-mCherry or
Sh-LINE-mCherry

3 days

+/- LPS

30 % Tet2$^{-/-}$ HSCs
Sh-control-GFP

B.

C.

Fig 7. Relative expansion of *Tet2*$^{-/-}$ HSCs in response to LPS is dependent on L1 expression in wild-type cells

Figure 7. Relative expansion of *Tet2*^{-/-} HSCs in response to LPS is dependent on L1 expression in wild-type cells **(A)** Experimental design to analyze *Tet2*^{-/-} vs WT HSCs competition in vitro in the presence or absence of LPS. 70% of WT HSCs transduced with either an sh-control or sh-LINE fused to mCherry reporter gene were cultured in vitro with 30% of *Tet2*^{-/-} HSCs transduced with an sh-control fused to GFP reporter gene, in the presence or absence of LPS. **(B)** Ratio of *Tet2*^{-/-} HSCs (GFP+) between LPS and non-LPS conditions when in competition with WT HSCs (mcherry+) transduced with an sh-control or with an sh-L1. **$p<0.01$ Unpaired t-test. **(C) Model.** Upon chronic inflammatory stress, the function of WT HSCs is altered through loss of H3K9me3 at L1, L1 overexpression and L1-induced accumulation of DNA damage. *Tet2*^{-/-} are resistant to L1-induced deleterious effects upon LPS treatment through epigenetic repression, eventually leading to their clonal expansion. Figure created with biorender.