Title:

Mothers respond to biological pup calls with heart rate changes in Japanese house bats, *Pipistrellus*

abramus

Author names and affiliations:

Kazuki YOSHINO-HASHIZAWA[1,2*], Midori HIRAGOCHI[1*], Kohta I KOBAYASI[1], Shizuko

HIRYU[1]

[1] Graduate School of Life and Medical Sciences, Doshisha University, 1-3 Tatara-Miyakodani,

Kyotanabe, Kyoto, 610-0394 Japan.

[2] School of Life Science and Technology, Institute of Science Tokyo, 4259 Nagatsuta, Midori,

Yokohama, Kanagawa, 226-8501 Japan.

* This symbol indicates equally contributed authors

ORCID iDs: 0000-0002-3569-4220 (KYH), 0000-0001-6303-4767 (KIK), 0000-0002-7106-1244

(SH)

To whom correspondence should be addressed.

kyoshino@ultrasonics.jp (KYH)

Keywords (least 3 keywords):

isolation call, electrocardiogram, autonomic nervous system, maternal care, motivation

Statements and Declarations

Competing Interests

The authors declare no conflict of interest.

Abstract

Maternal care is essential for offspring survival in mammals, especially in colonial species

where mothers must recognize their own young among many. In the Japanese house bat, *Pipistrellus*

abramus, mothers identify their pups using acoustic cues, particularly isolation calls (ICs) produced

by newborns. However, little is known about the physiological responses underlying maternal

recognition. Here, we investigated emotional responses of mother bats to pup calls by measuring

heart rate (HR) changes during a controlled playback experiment. We recorded ICs from 2-day-old

pups and echolocation calls (ECs) from 30-day-old pups, then presented these sounds to mothers

after pup independence. HR significantly increased in response to their own pups' calls, both ICs and

ECs, while calls from non-family pups did not elicit significant changes. Among these, ECs from

their own older pups evoked the largest average HR increase, suggesting sustained maternal

responsiveness despite developmental changes in call structure. Additionally, unfamiliar adult calls

induced HR elevation, possibly reflecting general social arousal. These findings indicate that *P.*

abramus mothers exhibit selective physiological arousal to their own pups' calls, with HR serving as

a sensitive marker of maternal recognition and motivation based on dynamic acoustic information.

Introduction

Maternal lactation is a fundamental behavior in mammals, providing offspring with essential

nutrients required for survival and proper development during the early stages of life. If a female were

to nurse the offspring of another species or unrelated individuals, her ability to provide sufficient milk

to her own offspring would be compromised, ultimately reducing her reproductive fitness.

Consequently, through long-term adaptive evolution, adult female mammals have developed the

ability to accurately recognize their own offspring [1]. Mammalian females utilize multiple sensory

modalities for offspring recognition, including visual [2], olfactory [3], and auditory [4] cues, or a

combination of these [5]. Among these, acoustic cues are particularly important in low-visibility

environments and are widely used by many animals, especially nocturnal species. Acoustic signals

emitted from pups, including isolation calls (ICs), are recognized by their mothers [6–9]. ICs are often

perceived as distress calls, reflecting the negative emotions states of offspring [10], and play critical

role in mother-pup communication and individual recognition, particularly when pups are separated

from their mothers [11]. Most research on mother-offspring communication via ICs has focused on

ultrasonic vocalizations (USVs) in rodents, commonly used as experimental models due to their

versatility [12]. Maternal behavior tests in rodents have demonstrated that mothers exhibit nurturing

behaviors – such as pup retrieval-, immediately after locating their pups via USVs. However, mice

and certain other rodents are known to care not only for their own pups but also for unrelated pups,

primarily relying on olfactory cues for pup recognition [13–15].

In contrast, echolocating bats present a unique model for studying mother-offspring

communication based on acoustic cures. It has been reported that bat pups produce ICs, and mother

bats can discriminate own pups using these calls [16–21]. ICs in bats are typically characterized by

long duration, low frequency, and a high repetition rate, allowing mothers to identify their own pups

even within large colonies [17,18,20]. Moreover, despite developmental changes in call structure,

mothers may maintain the ability to recognize their pups throughout the nursery period [22].

Collectively, these findings suggest that mother bats are well-suited model animals for studying

maternal responses to ICs, particularly in the context of individual recognition.

While behavioral responses, such as pup retrieval, have been widely used to infer maternal

recognition, such behaviors do not fully capture the underlying emotional and motivational states of

the mothers. Previous studies have suggested that ICs elicit nurturing behaviors via emotional arousal

[23]; however, arousal is not always apparent through behavior alone. To assess internal emotional

states directly, physiological measures offer a valuable complementary approach. Heart rate (HR) is

widely used as a minimally invasive physiological indicator of emotional arousal across various

species. Monitoring HR can reveal subtle internal state changes that may not be observable through

behavior, providing insight into the affective component of maternal responses. HR measurements in

bats may help clarify whether mothers exhibit specific physiological responses to their own pups' calls,

or whether all ICs are perceived as signals of negative emotional valence. In this study, we measured

HR responses in mother bats during playback of pup vocalizations to investigate their emotional

reactions. Building on these approaches, our study aims to explore maternal emotions and motivation

in bats from a psychophysiological perspective by examining HR responses to acoustic stimuli, with

a particular focus on the discrimination of calls from their own versus non-family pups.

Materials and Methods

Study species and captive status

The target species was the Japanese house bat *Pipistrellus abramus* (Temminck, 1840), an

insectivorous, aerial-hunting species belonging to the bat family Vespertilionidae. For echolocation,

adults of this species use frequency-modulated (FM) pulses that sweep downward from initial

frequencies around 80 kHz to terminal frequencies (TF) of around 40 kHz [24,25]. Their auditory

sensitivity exhibits a broad U-shaped curve in the 4-80 kHz frequency range, with low thresholds

between 20 and 50 kHz and the highest sensitivity in the 35-50 kHz range [26]. Vocalizations in *P.*

abramus pups differ from adult echolocation calls and appear to serve as precursors of echolocation

calls as well as ICs [22]. They raises their pups in colonies of 50–200 individuals, where multiple

families of females and their pups coexist [27].

In May 2022, four late-pregnancy female bats were captured from a colony roosting in

bridge girders near the campus of Doshisha University, in Kyotanabe, Kyoto, Japan. Each female was

individually banded with a marked ring on her forearms (2.0 mm internal diameter, 4.0 mm height,

Porzana Ltd, Winchelsea, U.K.). All four females gave birth to litters of three pups in the laboratory

over three days, from 20 to 22 June 2022. We checked the condition of the pups daily, and they were

well cared for by their mothers. The pups' body mass ranged from 1.1 to 1.3 g, whereas adult body

mass ranged from 5 to 8 g. Since two pups died shortly after birth (pup-Ba from mother-B and pup-

Db from mother-D), we recorded the vocalizations of the ten surviving captive-born pups (three female

and seven male) from the four mothers during their first postnatal month. The pups were identified as

follows: pup-Aa, -Ab, and -Ac from mother-A; pup-Bb and -Bc from mother-B, pup-Ca, -Cb, and -

Cc from mother-C; pup-Da and -Dc from mother-D. Each pup was marked using a harmless animal

marker (FG2200G, EXEC, Japan) applied to a small area of skin on its back. All mothers and pups

were kept in a rearing cage (25 cm × 17 cm × 17 cm) under controlled conditions, including a

regulated temperature (25 ± 2°C), humidity (55 ± 5%), and light–dark cycles following the natural

photoperiod in Kyotanabe, Kyoto, Japan. The female bats were fed mealworms, provided with water,

and given an adequate amount of vitamins to ensure proper nutrition.

Experiment I: Recording vocalization from pups

Pup vocalizations were recorded daily during their first post-natal month. To record each

pup vocalizations, one pup was gently separated from its mother and put on an echo-soundproof

sponge (30 cm × 30 cm) in the experimental room. After the recording, the pups were returned to their

mother, and the same procedure was repeated for its siblings. Vocalizations were recorded using an

ultrasonic microphone (Ultrasoundgate CM16/CMPA, Avisoft Bioacoustics, Germany) placed at

approximately 20 cm from the pup for 3 minutes with a sampling frequency of 250 kHz and a

resolution of 16 bits. As a control, we also recorded vocalizations from one adult male bat using same

recording setup. A custom MATLAB program was used to analyze the acoustic characteristics of the

calls, including duration, terminal frequency (TF), and initial frequency (IF), obtained from the

spectrogram at -20 dB relative to the peak intensity of the call. These characteristics were then used

to classify the calls into the ICs and echolocation calls (ECs), based on previous study [22].

Experiment II: ECG recording from mother bats

To avoid interference with maternal care, electrocardiograms of four mother bats were

recorded from 38 to 45 days after delivery, when the pups had matured. During these recordings, sound

stimuli were presented. All sound stimuli were taken from pup vocalizations recorded in Experiment

I, specifically from males at 2 days after birth (DAB) for ICs and at 30 DAB for ECs, and were

normalized to a length of 3 seconds, containing between 15 to 33 calls per stimulus.

Five types of sound stimuli were presented to each mother bat: (I) ICs from their own pups

at 2 DAB, (II) ECs from their own pups at 30 DAB, (III) ICs from non-family pups at 2 DAB, (IV)

ECs from non-family pups at 30 DAB, and (V) ECs from an unfamiliar adult bat. For stimuli III and

IV, two or three non-family pup calls were uses, consisting of calls from pups of different mothers

(e.g., mother A was exposed pup Bc's and Cc's calls for both stimuli III and IV). All stimuli were

normalized the stimuli based on peak-to-peak amplitude, as it was difficult to control the loudness due

to differences in call duration and frequency range of ECs and ICs.

This experiment was carried out in a soundproof chamber (1.4 m×1.4 m×2.1 m). Each

subject was positioned in the center of a circular table using soft sponges for stabilization. Sound

stimuli were presented at 100 dB peak-to-peak SPL via a tweeter (PT-R7III, Pioneer, Japan) located

approximately 15 cm from the subject. The stimuli were played through an audio interface (Rubix44,

Roland, Japan) at a sampling frequency of 192 kHz and a resolution of 16 bits, amplified by an

amplifier (STR-DH190, Sony, Japan) connected to the tweeter.

A total 39 sound stimuli were presented to each mother over two days, with each of the five

stimulus types presented at least six times. One of the four mothers died before the second day of the

experiment and was therefore exposed to only 18 stimuli. The order of stimulus presentation was

randomized, except for the first stimulus of each day. To avoid potential startle responses from the

first stimulus, the adult EC was always presented first and was excluded from subsequent analysis.

ECG electrodes were attached subcutaneously in a Lead II configuration at three positions:

left leg (positive), right thumb (negative), and right leg (reference). Lead II is known to provide a

highest signal-to-noise ratio for ECG recording in echolocating bats [28]. Subjects were anesthetized

with 2–4% isoflurane (Pfizer, USA) during surgery. Fur was shaved at three dorsal sites (base of the

right arm, left leg, and right leg), and 2% xylocaine jelly (Sandoz, Switzerland) was applied as a local

anesthetic to facilitate the subcutaneous insertion of small safety-pin electrodes. Electrodes were

connected to the ECG recording system (C3410 / C3314 / C3216 / C3100, Intan technology, USA)

and, signals were recorded using vendor software (RHX software, Intan technology, USA) at a

sampling rate of 30 kHz. ECG recordings were obtained by measuring the potential differences

between the positive and the negative electrodes. Before presenting sound stimuli, subjects were

allowed to acclimate until their HR stabilized at approximately 200-300 beats per minute (bpm). ECGs

were recorded from 50 seconds before to 100 seconds after the onset of each sound stimulus (one trial).

After each presentation, we ensured that the HR returned to baseline before presenting the next

stimulus.

ECG processing and statistical analysis

All digital signal processing was performed using a custom MATLAB program (R2021a,

MathWorks, USA). To reduce noise, a 1–650 Hz band-pass filter was applied to the raw ECG.

Instantaneous HR was calculated at the reciprocal of the R-R interval, which was automatically

detected from the filtered ECG signal. HR values were then binned into one-second intervals, and

relative HR changes were calculated by subtracting the average HR in -10 to 0 second period as the

baseline. HR increases were quantified by averaging HR values over the 0–30 second period following

the onset of each sound stimulus type [29].

All statistical analyzes were performed using a custom MATLAB program (R2024a,

MathWorks, USA). To test for significant increases in HR in response to each acoustic stimulus, we

applied a one-sample Wilcoxon signed-rank test with Bonferroni correction to average HR values over

the 0–30 second period after the stimulus onset in Experiment II.

Results

Experiment I: Developmental changes in ultrasonic vocalizations by pups

All pups vocalized ICs until approximately 10 to 12 DAB (Fig. 1 top). Although ICs were already present at 0 DAB, their amplitude was unstable, making it difficult to evaluate their acoustic components. At 2 DAB, the fundamental frequency component of ICs consisted of FM sounds that descended from the initial frequency (IF) of the first harmonic, approximately 35 kHz, to the terminal frequency (TF), around 20 kHz, across all pups (see Table S1). As the pups developed, both IF and TF with development, accompanied by a reduction in duration. By 10 DAB, their ICs reached approximately 40 kHz (IF), 25 kHz (TF) and a duration of 10 ms.

In contrast, ECs first appeared at approximately 8 DAB (Fig. 1 bottom). At this stage, the IF of the first harmonic was around 50 kHz, and the TF was around 30 kHz. Both frequencies increased with postnatal development, and the duration gradually shortened, reaching levels comparable to those of adult ECs by 30 DAB (see Table S1). Figure 2A shows an example of postnatal vocalization development in a representative pup (Ab). During this developmental progress, their vocalizations gradually shifted toward the acoustic characteristics of adult ECs, and this pattern was consistently observed in other pups as well (Fig. 2B).

Experiment II: Mother bat's ECG changes for sound stimuli

As illustrated in Fig. 3, the ECG and HR responses of the mother bats were recorded before and after the presentation of sound stimuli. ECG signals were recorded with a sufficient signal-to-noise ratio to allow clear identification of R waves for HR calculation (Fig. 3A). In all five types of stimuli, the average HR showed an immediate increase following stimulus onset (Fig. 3B).

The largest increase in average HR was observed in response to (II) ECs from their own

pups at 30 DAB, reaching approximately 130 bpm at its peak, which was significantly higher than

baseline levels ($P = 0.001$; Fig. 3C). Additionally, (V) ECs from an unfamiliar adult bat also induced

significantly higher HR increases ($P = 0.001$). In response to (I) ICs from their own pups at 2 DAB,

HR increased by approximately 70 bpm (Fig. 3B), which was also significantly higher than baseline

($P = 0.040$). As shown in Fig. 3C, although (III) ICs from non-family pups at 2 DAB and (IV) ECs

from non-family pups at 30 DAB appeared to induce slight increases in average HR, these increases

were not statistically significant (III: $P = 0.137$, IV: $P = 0.171$).

Discussion

In Experiment II, the HR of mother bats significantly increased in response to their own

pups' vocalization at both 2 DAB and 30 DAB (stimuli I and II), whereas call from non-family pups

(stimuli III and IV) did not induce significant changes (Fig. 3B, 3C). These results suggested that *P.*

abramus mothers recognize their own pups based on acoustic features of their calls, and that this

recognition is reflected in physiological changes, specifically increased HR. Previous studies in

rodents have shown that ICs are associated with distress signals, eliciting arousal or stress responses

in mother [10]. It is therefore plausible that mother bats also experience arousal upon hearing their

own pups' calls—both ICs and ECs—which may promote maternal behaviors such as nursing or

retrieval. In other species, ECs have been shown to encode various types of social information,

including sex, individual identity, group membership, and behavioral states [30–34]. Similarly, *P.*

abramus may also extract social information from ECs, which could explain the strong HR response

to their own pups' ECs at 30 DAB. Notably, in *P. pipistrellus*, mothers do not exhibit care toward

non-family pups [35]. The absence of significant HR increase in response to non-family pup calls

(stimuli III and IV) in *P. abramus* may reflect a similarly low level of maternal motivation toward

unrelated offspring. Interestingly, the mothers exhibited significant HR increases in response to (V)

ECs from an unfamiliar adult bat, even though this stimulus was intended as social control. While we

could not directly assess the cause of this response, one possibility is that, around 40 days postpartum

when the experiment was conducted, females were not yet in the mating season. Consequently, hearing

adult male calls might have triggered a rejection-related physiological response, potentially mediated

by ventromedial hypothalamus and sympathetic nervous system [36,37]. Further investigation is

needed to clarify the neural mechanisms underlying reproductive status, social cue processing, and

vocal communication in mother bats.

When we compared the mothers' HR in response to their own pups' calls at two different

postnatal days, both responses were significantly higher than baseline (stimulus I and II in Fig. 3B,

3C). ICs at 2 DAB were assumed to play an extremely important role in the pup's survival and were

expected to elicit a stronger maternal response than ECs at 30 DAB. However, our HR results did not

support this assumption (I: $P = 0.040$, II: $P = 0.001$; Wilcoxon signed-rank test with Bonferroni

correction). One possible explanation for this discrepancy is that mother bats, goats, and sea lions

[7,38], may continuously update their memory of pup calls on a daily basis. In this study, ECG

measurements of mother bats began approximately 40 days after delivery (38 to 45 days), when the

pups had already become independent. This timing was chosen to avoid undue stress on lactating

mothers and to prevent the risk of harming immature pups, which cannot maintain their body

temperature for more than an hour. It is therefore possible that the calls at 30 DAB more closely

resembled the most recently experienced pup vocalizations, whereas the sounds at 2 DAB may have

appeared unfamiliar or unnatural during the experiment. As shown in Figure 1, the acoustic

characteristics of pup vocalizations change daily. We therefore hypothesize that mothers' recognition

abilities undergo continuous adaptation. To address memory-related effects linked to vocal changes

during development, future studies should incorporate experimental designs such as daily auditory

playback to evaluate how dynamic memory updating influences maternal motivation.

The findings of this study demonstrate that changes in the internal state of mother bats in

response to pup calls can be assessed by monitoring increases in HR. Specifically, we found that HR

increases were selectively induced by calls from their own pups but not by calls from non-family pups,

suggesting that HR reliably reflects maternal recognition and motivation in *P. abramus.* Previous

studies have reported HR increases in echolocating bats in response to negative social calls [29,39–

41], indicating that HR is a useful physiological indicator of emotional arousal elicited by acoustic

stimuli. However, a limitation of using HR alone is its inability to distinguish between positive and

negative emotional valences, as elevated HR can accompany both distress and affiliative responses.

In other animal species, including humans, heart rate variability (HRV) is commonly used to infer the

balance between sympathetic and parasympathetic nervous system activity and to provide a more

nuanced assessment of emotional state [42–44]. To evaluate more complex emotional dimensions,

such as the strength of mother-offspring bonds, feelings of security, empathy-like responses, and

developmental maturity, it will be important to combine HR measurements with HRV analyzes.

HRV has proven particularly effective in detecting subtle emotional responses to social

acoustic stimuli, such as vocal communication. Nevertheless, in bats, standard frequency band

definitions for low-frequency (LF) and high-frequency (HF) components, essential for HRV power

spectrum analysis, have not been established, likely due to their highly dynamic HR patterns. Further

investigation is needed to define species-specific HRV parameters and to establish reliable

physiological markers of autonomic function and emotional states in bats.

References

1. Klug H, Bonsall MB, Alonzo SH. 2013 The origin of parental care in relation to male and female

life history. *Ecol. Evol.* **3**, 779–791. (doi:10.1002/ece3.493)

2. Dasser V. 1988 A social concept in Java monkeys. *Anim. Behav.* **36**, 225–230. (doi:10.1016/S0003-

3472(88)80265-3)

3. Porter RH. 1998 Olfaction and human kin recognition. *Genetica* **104**, 259–63.

(doi:10.1023/a:1026404319384)

4. Briefer E, McElligott AG. 2011 Mutual mother-offspring vocal recognition in an ungulate hider

species (Capra hircus). *Anim. Cogn.* **14**, 585–98. (doi:10.1007/s10071-011-0396-3)

5. Wierucka K, Charrier I, Harcourt R, Pitcher BJ. 2018 Visual cues do not enhance sea lion pups'

response to multimodal maternal cues. *Sci. Rep.* **8**, 1–7. (doi:10.1038/s41598-018-28171-w)

6. Nagasawa M, Okabe S, Mogi K, Kikusui T. 2012 Oxytocin and mutual communication in mother-

infant bonding. *Front. Hum. Neurosci.* **6**, 31. (doi:10.3389/fnhum.2012.00031)

7. Pitcher BJ, Harcourt RG, Charrier I. 2010 Rapid onset of maternal vocal recognition in a colonially

breeding mammal, the Australian sea lion. *PLoS One* **5**, e12195.

(doi:10.1371/journal.pone.0012195)

8. Sèbe F, Aubin T, Boué A, Poindron P. 2008 Mother-young vocal communication and acoustic

recognition promote preferential nursing in sheep. *J. Exp. Biol.* **211**, 3554–3562.

(doi:10.1242/jeb.016055)

9.   Geissler DB, Ehret G. 2002 Time-critical integration of formants for perception of communication

calls in mice. *Proc. Natl. Acad. Sci.* **99**, 9021–9025. (doi:10.1073/pnas.122606499)

10.   Okabe S, Kanno K. 2023 Acoustic Properties and Biological Significance of Ultrasonic

Vocalizations in Rodents: Emotional Expressions. In *Acoustic Communication in Animals*, pp.

153–173. Singapore: Springer Nature Singapore. (doi:10.1007/978-981-99-0831-8_10)

11.   Camaclang AE, Hollis L, Barclay RMR. 2006 Variation in body temperature and isolation calls of

juvenile big brown bats, Eptesicus fuscus. *Anim. Behav.* **71**, 657–662.

(doi:10.1016/j.anbehav.2005.07.009)

12.   Ehret G. 2005 Infant rodent ultrasounds - A gate to the understanding of sound communication.

Behav. Genet. **35**, 19–29. (doi:10.1007/s10519-004-0853-8)

13.   Ebensperger LA, Hurtado MJ, Valdivia I. 2006 Lactating Females Do Not Discriminate Between

Their Own Young and Unrelated Pups in the Communally Breeding Rodent, Octodon degus.

Ethology **112**, 921–929. (doi:10.1111/j.1439-0310.2006.01251.x)

14.   Wang Z, Storm DR. 2011 Maternal behavior is impaired in female mice lacking type 3 adenylyl

cyclase. *Neuropsychopharmacology* **36**, 772–81. (doi:10.1038/npp.2010.211)

15.   Okabe S, Nagasawa M, Kihara T, Kato M, Harada T, Koshida N, Mogi K, Kikusui T. 2010 The

Effects of Social Experience and Gonadal Hormones on Retrieving Behavior of Mice and their

Responses to Pup Ultrasonic Vocalizations. *Zoolog. Sci.* **27**, 790–795. (doi:10.2108/zsj.27.790)

16.   Scherrer JA, Wilkinson GS. 1993 Evening bat isolation calls provide evidence for heritable

signatures. *Anim. Behav.* **46**, 847–860. (doi:10.1006/anbe.1993.1270)

17.   Matsumura S. 1981 Mother-Infant Communication in a Horseshoe Bat (Rhinolophus

304   ferrumequinum nippon): Vocal Communication in Three-Week-Old Infants. *J. Mammal.* **62**, 20–

305   28. (doi:10.2307/1380474)

306   18.   Matsumura S. 1979 Mother-Infant Communication in a Horseshoe Bat (Rhinolophus

307   ferrumequinum nippon): Development of Vocalization. *J. Mammal.* **60**, 76–84.

308   (doi:10.2307/1379760)

309   19.   Tan X, Li Y, Sun K, Jin L, Feng J. 2022 Mutual mother-pup acoustic identification in Asian

310   particolored bats. *Ecol. Evol.* **12**. (doi:10.1002/ece3.9554)

311   20.   Pfalzer G, Kusch J. 2003 Structure and variability of bat social calls: Implications for specificity

312   and individual recognition. *J. Zool.* **261**, 21–33. (doi:10.1017/S0952836903003935)

313   21.   Li Y, Sun D, Tan X, Chi T, Liu Y, Jin L, Feng J. 2020 Isolation calls from twin siblings are similar

314   in the Asian parti-coloured bat, Vespertilio sinensis. *Bioacoustics* **29**, 697–708.

315   (doi:10.1080/09524622.2019.1673819)

316   22.   Hiryu S, Riquimaroux H. 2011 Developmental changes in ultrasonic vocalizations by infant

317   Japanese echolocating bats, Pipistrellus abramus. *J. Acoust. Soc. Am.* **130**, EL147-53.

318   (doi:10.1121/1.3632044)

319   23.   Bell RW. 1974 Ultrasounds in small rodents: Arousal-produced and arousal-producing. *Dev.*

320   *Psychobiol.* **7**, 39–42. (doi:10.1002/dev.420070107)

24. Hiryu S, Hagino T, Fujioka E, Riquimaroux H, Watanabe Y. 2008 Adaptive echolocation sounds

of insectivorous bats, Pipistrellus abramus, during foraging flights in the field. *J. Acoust. Soc. Am.*

124, EL51-6. (doi:10.1121/1.2947629)

25. Fujioka E, Mantani S, Hiryu S, Riquimaroux H, Watanabe Y. 2011 Echolocation and flight strategy

of Japanese house bats during natural foraging, revealed by a microphone array system. *J. Acoust.*

Soc. Am. **129**, 1081–1088. (doi:10.1121/1.3523300)

26. Boku S, Riquimaroux H, Simmons AM, Simmons JA. 2015 Auditory brainstem response of the

Japanese house bat (Pipistrellus abramus). *J. Acoust. Soc. Am.* **137**, 1063–1068.

(doi:10.1121/1.4908212)

27. Funakoshi K, Katahira R, Ikeda H. 2009 Night-roost usage and nocturnal behavior in the Japanese

house-dwelling bat, Pipistrellus abramus. *Mammal Study* **34**, 131–139. (doi:10.3106/041.034.0302)

28. Mihova D, Hechavarría JC. 2016 The electrocardiogram signal of Seba's short-tailed bat, Carollia

perspicillata. *J. Comp. Physiol. A* **202**, 517–526. (doi:10.1007/s00359-016-1098-8)

29. Yoshino-Hashizawa K, Nishiuchi Y, Hiragochi M, Kihara M, Kobayasi KI, Hiryu S. 2023 The

distress context of social calls evokes a fear response in the bat Pipistrellus abramus. *J. Exp. Biol.*

226, jeb246271. (doi:10.1242/jeb.246271)

30. Knörnschild M, Jung K, Nagy M, Metz M, Kalko E. 2012 Bat echolocation calls facilitate social

communication. *Proc. R. Soc. B Biol. Sci.* **279**, 4827–4835. (doi:10.1098/rspb.2012.1995)

31. Yovel Y, Melcon ML, Franz MO, Denzinger A, Schnitzler HU. 2009 The voice of bats: How

340   greater mouse-eared bats recognize individuals based on their echolocation calls. *PLoS Comput.*

341   *Biol.* **5**. (doi:10.1371/journal.pcbi.1000400)

342 32. Tan X, Lin A, Sun K, Jin L, Feng J. 2022 Greater Horseshoe Bats Recognize the Sex and Individual

343   Identity of Conspecifics from Their Echolocation Calls. *Animals* **12**. (doi:10.3390/ani12243490)

344 33. Masters WM, Raver KAS, Kazial KA. 1995 Sonar signals of big brown bats, Eptesicus fuscus,

345   contain information about individual identity, age and family affiliation. *Anim. Behav.* **50**, 1243–

346   1260. (doi:10.1016/0003-3472(95)80041-7)

347 34. Barclay RMR, Jacobs DS. 2023 Interindividual communication by bats via echolocation. *Can. J.*

348   *Zool.* **101**, 128–143. (doi:10.1139/cjz-2022-0121)

349 35. Hughes PM, Speakman JR, Jones G, Racey PA. 1989 Suckling behaviour in the pipistrelle bat

350   (Pipistrellus pipistrellus). *J. Zool.* **219**, 665–670. (doi:10.1111/j.1469-7998.1989.tb02607.x)

351 36. Gutierrez-Castellanos N, Husain BFA, Dias IC, Nomoto K, Duarte MA, Ferreira L, Lacoste B,

352   Lima SQ. 2025 A hypothalamic node for the cyclical control of female sexual rejection. *Neuron*

353   **113**, 277-290.e8. (doi:10.1016/j.neuron.2024.10.026)

354 37. Shao J, Chen Y, Gao D, Liu Y, Hu N, Yin L, Zhang X, Yang F. 2023 Ventromedial hypothalamus

355   relays chronic stress inputs and exerts bidirectional regulation on anxiety state and related

356   sympathetic activity. *Front. Cell. Neurosci.* **17**, 1–14. (doi:10.3389/fncel.2023.1281919)

357 38. Briefer EF, de la Torre MP, McElligott AG. 2012 Mother goats do not forget their kids' calls. *Proc.*

358   *R. Soc. B Biol. Sci.* **279**, 3749–3755. (doi:10.1098/rspb.2012.0986)

39. Ma J, Naumann RT, Kanwal JS. 2010 Fear conditioned discrimination of frequency modulated

sweeps within species-specific calls of mustached bats. *PLoS One* **5**.

(doi:10.1371/journal.pone.0010579)

40. Gadziola MA, Shanbhag SJ, Wenstrup JJ. 2015 Two distinct representations of social vocalizations

in the basolateral amygdala. *J. Neurophysiol.* **115**, 868–886. (doi:10.1152/jn.00953.2015)

41. Gadziola MA, Grimsley JMS, Faure PA, Wenstrup JJ. 2012 Social Vocalizations of Big Brown

Bats Vary with Behavioral Context. *PLoS One* **7**, e44550. (doi:10.1371/journal.pone.0044550)

42. Liu J, Wei W, Kuang H, Tsien JZ, Zhao F. 2014 Heart rate and heart rate variability assessment

identifies individual differences in fear response magnitudes to earthquake, free fall, and air puff in

mice. *PLoS One* **9**. (doi:10.1371/journal.pone.0093270)

43. Friedman BH. 2007 An autonomic flexibility-neurovisceral integration model of anxiety and

cardiac vagal tone. *Biol. Psychol.* **74**, 185–199. (doi:10.1016/j.biopsycho.2005.08.009)

44. Carravieri A, Müller MS, Yoda K, Hayama SI, Yamamoto M. 2016 Dominant parasympathetic

modulation of heart rate and heart rate variability in a wild-caught seabird. *Physiol. Biochem. Zool.*

89, 263–276. (doi:10.1086/686894)

Acknowledgments

We are grateful to Motoki Kihara, Mayuko Mogi and Daisuke Namazue for their assistance with the

recordings. We also thank Yu Teshima and Andreas Mayer for their helpful comments on the

additional analysis.

381
382 **Fig. 1. Developmental changes in ultrasonic vocalizations by pups of *P. abramus*. Spectrograms of ultrasound**
383 **vocalizations produced by a pup. Upper row: isolation calls (ICs) recorded when the pup was separated from its**
384 **mother, from 0 to 12 DAB. Bottom row: echolocation calls (ECs) recorded when the pup was actively exploring**
385 **from 8 to 30 DAB, along with a representative EC from an adult bat.**
386

Fig. 2. Developmental changes in ultrasonic vocalizations by pups of *P. abramus*. (A) Example of developmental

changes in duration, initial frequency (IF) and terminal frequency (TF) from 0 to 30 DAB in a representative

pup (Ab). Red to blue points indicate changes over time, and black points represent ECs from an adult bat. (B)

Average developmental changes in vocalization parameters across pups from different mothers, along with the

adult bat EC (black star marker). The large open circle marker indicates vocalizations at 0 DAB, and the large

open square marker indicates those at 30 DAB.

Fig. 3. Heart rate responses to acoustic presentation in mother bats. (A) Example of a raw electrocardiogram (ECG) recording from a mother bat. The gray area indicates the period of acoustic stimulus presentation with echolocation calls (ECs) from the mother's own pup. (B) Average heart rate (HR) responses of all mothers to acoustic stimuli. The baseline was calculated as the average HR during the period from -10 to 0 seconds before stimulus onset. The five stimulus types were: (I) isolation calls (ICs) from their own pups at 2 DAB, (II) ECs from their own pups at 30 DAB, (III) ICs from non-family pups at 2 DAB, (IV) ECs from non-family pups at 30 DAB, and (V) ECs from an unfamiliar adult bat. (C) Distribution of HR increases from 0 to 30 seconds after stimulus onset. Each box plot shows the median (red line), interquartile range (box edges), and the minimum and maximum values (whiskers). Statistical significance is indicated by *$P<0.05$, **$P<0.01$ (Wilcoxon sign rank test with Bonferroni correction).

Funding information

This work was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grant [Numbers JP 21H05295, 22H01503 to SH, 23KJ2080, and 25KJ0298 to KYH], the Japan Science and Technology Agency (JST) SPRING Grant [Number JPMJST2129 to KYH], the JST establishment of university fellowships towards the creation of science technology innovation Grant [Number JPMJFS2145 to KYH] and the Japan Science Society (JSS) Sasakawa Scientific Research Grant [Number 2021-5028 to KYH].

Ethics statement

All experiments complied with the Principles of Animal Care (publication no.86-23 [revised 1985] of the National Institutes of Health) and all Japanese laws. All procedures were approved by the Animal Experiment Committee at Doshisha University.

Competing interests statement

The authors declare there are no conflicts of interest.

Data accessibility statement

Data are available at figshare (doi: 10.6084/m9.figshare.29614925). Other raw data supporting the findings of this study are available from the corresponding author on a reasonable request.

Artificial Intelligence (AI) declaration

We have not used AI-assisted technologies in creating this article.