1

Geographic Divergence in Population Genomics and Shell Morphology

Reveal History of Glacial Refugia in a Coastal Dogwhelk

Emily K. Longman[1], Joaquin C.B. Nunez[1], Eric Sanford[2,3], & Melissa Pespeni[1]

[1] Department of Biology, University of Vermont, Burlington, VT, USA

[2] Bodega Marine Laboratory, University of California Davis, Bodega Bay, CA, USA

[3] Department of Evolution and Ecology, University of California Davis, Davis, CA, USA

Corresponding Author: Emily K. Longman, Emily.Longman@uvm.edu

Keywords: *Nucella canaliculata*, Population genomics, Geometric morphometrics, Phylogeography, Ecological genomics

Abstract:

Studying contemporaneous spatial patterns of genomic diversity can yield important insights into the evolutionary processes that structure populations and shape patterns of adaptation. In contrast to the large number of marine species with planktonic larvae, populations of marine taxa with low dispersal and deep evolutionary divergences offer an opportunity to reveal the phylogeographic histories of marine ecosystems. Here, we constructed a draft genome assembly for the low-dispersing marine dogwhelk, *Nucella canaliculata*, and studied patterns of genomic diversity and shell morphometrics in 19 populations distributed along ~1,500 km of the west coast of North America. We observed significant population structure with a strong phylogeographic break at Monterey Bay, which was matched with divergence in shell morphology. Genomic patterns, concomitant with computer simulations, suggest that there were at least two refugial populations during the last glacial maximum that subsequently experienced post-glacial expansion and admixture. Furthermore, by linking genotype to phenotype, we identified candidate loci associated with the molecular bases of variation in shell morphology. These findings highlight that historical events can play a persistent role in structuring the genomic diversity of modern coastal populations, with important consequences for spatial variation in the adaptive capacity and vulnerability of populations to future climatic changes.

Introduction:

Introduction:

Understanding the population structure and demography of a species is essential for

interpreting patterns of adaptation and linking phenotypic variation to its genetic bases [1].

Spatially and temporally varying selection pressures [2], historical events [3,4], biogeographic

barriers to dispersal [5], and life history patterns [6], can all shape the present-day geographic

distribution of populations and the genetic diversity they harbor. However, our understanding of

population structure and subsequently the molecular bases of adaptation in many species is

significantly hindered by the lack of genomic resources for non-model organisms [7,8]. This

demographic information is of pressing concern because an understanding of extant levels of

genomic diversity is needed to predict a species' adaptive capacity and vulnerability to a rapidly

changing climate [9-11].

In marine ecosystems, understanding population connectivity and genetic diversity is

further complicated by extrinsic factors such as oceanographic processes and the open and three-

dimensional nature of marine environments [12–15], as well as intrinsic factors including the

wide range of life history strategies and developmental modes present in marine organisms

[6,16]. The location of phylogeographic breaks and the level of population structure often reflect

a species' reproductive strategy; species with direct development and low dispersal potential

typically have much more defined genetic structure than species with planktonic larvae

[13,14,17,18].

Using marine taxa with low dispersal to understand historical responses to climate change

could be a useful tool to predict how marine species will respond during the modern era of rapid

global change [19]. This is because the genomes of species with low dispersal often reflect

deeper evolutionary divergence over longer timescales [19–21]. The low levels of gene flow in

taxa with direct development allow for the historical separation among populations to persist for

many generations [13,20]. Additionally, advances in genomics and single nucleotide

polymorphism (SNP) based approaches can provide much greater resolution of these

demographic patterns [12,22,23]. Thus, studying the phylogeographic structure of low-dispersing

species can lend insight into the processes shaping the evolutionary history of marine ecosystems

and the impact of past and possibly future climatic changes.

Fossil assemblages of calcifying marine species along the northeast Pacific from the late

Pleistocene show that the geographic distribution of many species shifted equatorially in

response to Pleistocene climatic fluctuations [24–26]. Classic genetic models of Pleistocene

glaciation suggest that these climatic changes should have impacts on present-day genetic

diversity, with greater genetic diversity in areas of glacial refugia than areas of population

expansion, and high dissimilarity between refugial populations due to the deeper divergence time

[3,4]. However, to date, relatively few phylogeographic studies along the west coast of North

America show evidence that historical events, such as Pleistocene glaciation, left a signature on

contemporary patterns of genetic diversity [13,19,20,27]. Instead, because most species studied

to date have planktonic larvae and display high levels of connectivity, their population structure

has been primarily attributed to oceanographic forces and vicariance, with Point Conception and

Cape Mendocino serving as common phylogeographic breaks [13,15,28]. In contrast, the few

studies on low-dispersing taxa suggest phylogeographical breaks at Monterey Bay and the long

stretches of sandy beaches around Los Angeles, California [13,19,28].

One such marine invertebrate with low dispersal potential is the channeled dogwhelk,

Nucella canaliculata, which is a species of predatory gastropod (Family Muricidae) that lives on

rocky shores from central California to Alaska and consumes mussel and barnacle prey via a

chemo-mechanical mode of feeding [29–30]. Females lay benthic egg capsules, and crawl-away

young establish in the surrounding area [29]. Additionally, individuals reside only in mussel beds

on wave-exposed headlands, thus populations are isolated due to intervening sandy beaches.

Previous research has shown that this species displays a pattern of local adaptation with striking

spatial variation in predatory drilling ability [30–33], foraging behavior [34], and heat tolerance

[35]. Despite the wealth of ecological knowledge about *N. canaliculata*, there is a lack of

genomic resources for the species, limiting our understanding of the molecular mechanisms

generating these adaptive patterns. Genetic studies of the mitochondrial locus cytochrome

oxidase I (*mtCOI*) suggest that populations display a pattern of isolation by distance and that the

southern-most sampled population in central California was the most genetically diverged [30].

The biogeographic patterns of intraspecific trait variation and the species habitat specificity

suggest that the population genomics of this low dispersing taxa may reveal spatially complex

phylogeographic patterns.

Here, we present a high-quality draft genome assembly generated from a single *Nucella*

canaliculata individual, a genome-wide panel of genetic variation from 19 populations spanning

~1,500km of the west coast of North America using pooled-sequencing [36], and analyses of

shell morphometrics from dogwhelks collected across the 19 field sites. There is a long history of

malacologists using geographic variation in shell morphology as an indicator of potential

isolation and genetic divergence among gastropod populations [37,38]. Thus, our analyses

explore whether spatial variation in morphological traits is concordant with the phylogeographic

patterns revealed by the genomic data. Additionally, we integrate the genomic and

morphological data to begin to link genetic and phenotypic variation. Specifically with these

genomic and morphological datasets, we aimed to address the following questions: 1) How is

genetic diversity in *N. canaliculata* spatially structured along ~1,500km of the west coast of

North America? 2) Do patterns of genetic diversity appear to be structured more strongly by

contemporary or historical processes? 3) Does variation in shell morphology suggest a similar

biogeographic pattern of divergence as the genomic data? 4) Lastly, what are the molecular bases

for intraspecific variation in shell morphology? Overall, our analyses revealed strong population

structure with congruence between the genomic and phenotypic datasets, as well as evidence of

glacial refugia.

Methods:

Genome Sequencing and Assembly

Genome assembly was based on source DNA extracted from one adult female dogwhelk

collected from Bodega Marine Reserve, California (38°19'8", -123°4'28") in June 2022. High

molecular weight DNA extraction from flash frozen foot tissue, library preparation and

sequencing (Oxford Nanopore PromethION and Element Bio AVITI 600 flow cell) occurred at

the DNA Technologies and Expression Analysis Core at the UC Davis Genome Center (see

Appendix S1: Section S1 for more details).

We assembled the genome using a hybrid assembly approach using `DBG2OLC` [39]. This

approach combines the continuity of the Oxford Nanopore long reads and the quality of the

AVITI short reads (see Appendix S1: Section S1 for more details). Genome quality and

completeness was quantified using `QUAST` [40] and `BUSCO` [v5.7.1] [41]. Repeats and low

complexity DNA sequences were masked using `RepeatMasker` [v4.1.6] with the model

trained on the Pacific oyster, *Crassostrea gigas* [42]. Genome annotation of the softmasked

genome occurred using `Augustus` [v3.5.0] trained on *Drosophila melanogaster* [43].

Field Collections and DNA Extractions

We collected 20 adult (>18 mm in length) *N. canaliculata* from 19 populations ($N = 380$

total individuals) along the Pacific coast of North America (Table S1). Collections occurred

during low tide at the 19 rocky headlands from October 2023 to July 2024. Directly after field

collections, we sampled and preserved in 95% EtOH foot muscle tissue for each dogwhelk. We

extracted the DNA from each individual dogwhelk using an E.Z.N.A Mollusc & Insect DNA Kit

(Omega Bio-Tek, Norcross, Georgia) with minor modifications. We determined DNA quality

and quantity using gel electrophoresis, a Nanodrop, and a Qubit. For each population, DNA from

the 20 individuals was pooled in equimolar amount. Subsequent library preparation

(Watchmaker DNA library preparation kit [Watchmaker Genomics, Boulder, Colorado] with 6

PCR cycles) and sequencing (150 base pair paired-end sequencing on two lanes of NovaSeqX

25B) was performed at the DNA Technologies and Expression Analysis Core at the UC Davis

Genome Center.

Bioinformatic Processing and SNP Calling

We assessed the quality of the raw pooled sequences using `FastQC` [44]. Adapters were

detected and low-quality reads were trimmed on a sliding window using a minimum phred score

of 20 with `fastp` [v0.23.4] [45]. Reads were mapped to the softmasked *N. canaliculata*

reference genome using the Burrows-Wheeler algorithm, `bwa-mem2` [2.2.1] [46]. We filtered

(quality threshold of 40), sorted and removed duplicate reads from the binary alignment map

(bam) files using `SAMtools` [v1.9] [47] and `PicardTools` [v1.9] [48]. The bam files from

the two sequencing lanes were merged and the final bam files were indexed using `SAMtools`

[v1.9] [47]. Mean coverage of the bam files was 33.13, with an average mapping rate of 57.25

(Table S2). We used `freebayes` [v1.3.6] to call variants from the 19 bam files [49] and

subsequently performed stringent filtering of the VCF file with `vcftools` [v0.1.16] and the

`poolfstat` R package [50] (Appendix S1: Section S2).

Population Structure and Genomic Diversity

Population structure was analyzed using principal component analysis (PCA) and F_{ST}.

The VCF file was converted to a pooldata object using the *vcf2pooldata* function in the

`poolfstat` R package [50]. A PCA was performed using the *randomallele* function. We

calculated a global F_{ST} metric, a block-jackknife estimation of F_{ST}, and a multi-locus F_{ST} over a

sliding window of 1kb using the *computeFST* function. Additionally, we estimated pairwise-

population F_{ST} using the *compute.pairwiseFST* function using the ANOVA method.

Population demography was further characterized by analyzing the Ω matrix in

`BayPass` [v2.41] [51]. An Ω matrix was generated using 100 pilot runs, then converted into a

covariance matrix then into dissimilarity matrix, which was subsequently visualized as a

hierarchical clustering tree using the ape R package [52].

We tested for isolation by distance (IBD) by evaluating whether there was a relationship

between pairwise genetic distance, $F_{ST}/(1- F_{ST})$, and geographic distance. Given that *N.*

canaliculata lacks a planktonic larval stage, distance between sites was measured as the distance

along the coastline using the measure tool on Google Maps. We used Mantel tests (1,000

permutations) to determine the significance of the relationships using the `ecodist` R package.

We calculated nucleotide diversity (π), Watterson's estimator (θ), and Tajima's D for

each population using `npstat` [v1] [53]. To do so, the bam file for each population was

converted to pileup format using `SAMtools` [v1.9] [47], then calculations for each statistic were

performed on a sliding window with a length of 25kb with a minimum and maximum coverage

of 20 and 120, respectively, a minimum base quality of 30 and a minimum allele count of 5.

Demographic Inference

Demographic Inference

We compared whether two post-glacial demography models fit our data using the

coalescent framework implemented in `moments` [v1.2.2] [54]. The models tested were: 1)

expansion from a single glacial refugium, and 2) a secondary encounter between two glacial

refugia. We used a pipeline similar to that deployed in [55] and [56] to fit `moments` to Pool-Seq

data (see Appendix S1: Section S3 for more details). For each model we report the Akaike

information criterion (AIC) for 10 optimization runs. To translate coalescent units into

chronological time, we scaled the output of the model (i.e., θ) by the factor $4L\mu$. Where L is the

length of sequence and μ is the mutation rate of *Drosophila melanogaster* 2.8×10^{-9} [57].

In addition to model inference, we also conducted forward genetic simulations using

`SLiM` [58]. The goal of these forward simulations was to contextualize the patterns of genetic

variation expected under two proposed scenarios. Our SLiM simulations consisted of seven

demes arranged in a stepping-stone fashion (see Appendix S1: Section S3 for more details). In

the one-refugium model, the first deme recolonizes the entire habitat after an extinction event. In

contrast, in the two refugia model, the habitat is recolonized by demes 1 and 7 post an extinction

event.

Dogwhelk Shell Geometric Morphometrics

Dogwhelk Shell Geometric Morphometrics

Shell morphology was characterized using landmark analysis. We took photographs of

the ventral surfaces of 380 field-collected *N. canaliculata* dogwhelks from the 19 populations

using a Canon DSLR camera with a Canon EF 28-136mm f/3.5-5.6 IS lens on a tripod. We

utilized `tpsUtil` (v1.83) and `tpsDig` (v2.31) to place 15 landmarks on predefined locations

on the ventral surface of the shell [59]. Dogwhelks with damaged shells were removed from the

analyses (n=10). We used the `geomorph` package in R [60] and the program `MorphoJ` [61] to

perform statistical analyses of the landmark coordinates. We performed a Procrustes

transformation to remove variation in dogwhelk orientation using the *gpagen* R function then

then analyzed the Procrustes coordinates using PCA. We also tested for intraspecific variation in

shell shape as a function of size (quantified as the log of the centroid size for each dogwhelk),

site or latitude, and their interaction with a Procrustes ANOVA using the R function *procD.lm*

with 1,000 permutations.

Detection of the Genomic Bases of Morphological Variation

We identified outliers associated with shell morphology using `BayPass` [v2.41] [51].

The coefficient of variation for PC1 and PC2 of the morphological data were used as phenotypic

input data for `BayPass`. Outlier SNPs were identified using a Bonferroni threshold (alpha =

$0.05/8{,}277{,}206 = 6.04\text{x}10^{-9}$), then subsequently annotated with `SnpEff` [62]. Putative functional

annotation of proteins was done using `blast` against Molluscan taxa within the UniProtKB

reference proteomes – Swiss-Prot database.

<u>Results:</u>

***The First Draft Genome of* Nucella canaliculata**

We utilized a hybrid assembly approach using a combination of Oxford Nanopore

Technologies PromethION long reads and Element Bio AVITI short reads to generate the first

genome of *Nucella canaliculata*. The *de novo* assembly was very large (a total assembly length

of 1.57 Gb with an N50 of 146,860 bp) and consisted of numerous scaffolds (18,944 contigs)

(Table 1). BUSCO scores indicate that the draft genome was 91.0% and 80.7% complete for

Eukaryota and Mollusca, respectively (Table 2).

Table 1. *Nucella canaliculata* genome assembly metrics.

Assembly Length (bp)	1,571,681,309
Number of Contigs/Scaffolds (bp)	18,944
Largest Contig/Scaffold (bp)	2,196,136
GC %	42.52%
N50 (bp)	146,860
L50 (bp)	3,117

218 **Table 2.** Quality of *Nucella canaliculata* genome assembly.

	Complete BUSCOs	Complete and Single Copy BUSCOs	Complete and Duplicated BUSCOs	Fragmented BUSCOs	Missing BUSCOs
Eukaryota (n = 255)	91.0%	85.1%	5.9%	4.7%	4.3%
Mollusca (n = 5295)	80.7%	70.9%	9.8%	3.2%	16.1%

Patterns of Population Structure

The Pool-Seq data were used to analyze patterns of population structure and spatial

patterns of genomic diversity. After stringent filtering, our final SNP list for the 19 pools

contained 8,277,206 SNPs. The principal component analysis (PCA) showed strong clustering of

populations based on geography (Figure 1; Figure S1). PC1 captured 30.54% of the variation and

strongly partitioned the populations into two clusters, one north and one south of Monterey Bay.

PC2 (8.83% variation), PC3 (7.49% variation), PC4 (5.4% variation), and PC5 (4.93% variation)

all further subdivide the seven populations in the southern cluster. It was not until PC6 (4.79%

variation) that there is notable separation among the 12 populations in the cluster north of

Monterey Bay.

Figure 1. Population structure and genetic variation among the 19 *Nucella canaliculata*
populations along the coast of California and Oregon, USA. Projections of the first two principal
component analysis (PCA) loadings on the map of the collection sites: A) PC1 and B) PC2. The
first map states the site codes and other biogeographically relevant landmarks. See Table S1 for
an explanation of the population codes and coordinates.

The genome wide mean F_{ST} was 0.582, a high value for a marine species (Table S3).

Multi-locus F_{ST} indicated high levels of F_{ST} distributed evenly across the genome (Figure S2).

Pair-wise F_{ST} among the 19 populations ranged from 0.002 to 0.8 (Table S4). The greatest

differentiation occurred between populations south versus north of Monterey Bay (top right

corner of Figure 2A). There was little genetic differentiation among the populations north of

Monterey Bay. In contrast, there was highly variable amounts of genetic differentiation among

the populations south of Monterey Bay (bottom right corner of Figure 2A).

244
Figure 2. A) Heatmap of pairwise F_{ST} for the 19 *Nucella canaliculata* populations. Blue and
yellow indicate high and low levels of genetic differentiation, respectively. B) The log of the
geographic distance in km between sites and pairwise F_{ST}. Color indicates which sites are being
compared — both in the southern cluster (red), both in the northern cluster (blue) and a comparison
between a northern and southern site (yellow). C) Hierarchical clustering tree based on the omega
matrix from `BayPass`. See Table S1 for definitions of population codes.

Across the 19 populations, *Nucella canaliculata* displayed a pattern of isolation by

distance (Mantel tests, R^2=0.480, *P*-value=0.001; Figure 2B). However, this pattern is only

present when contrasting populations across the southern versus northern genetic cluster, and

within the southern cluster (Figure 2B). Despite their greater geographic distance, there was no

pattern of isolation by distance among the populations north of Monterey Bay (blue dots in

Figure 2B). Hierarchical clustering based on the `BayPass` population covariance matrix

recapitulated the clustering of the populations into two genetically distinct groups (Figure 2C).

All 19 populations had very low levels of genetic variation, heterozygosity and Tajima's

D (Figure 3; Figure S3; Table S4). The two populations just south of Monterey Bay, Soberanes

Point and Point Lobos, had the highest nucleotide diversity and heterozygosity (Figure 3; Figure

S3). Additionally, despite mean nucleotide diversity being similar across most of the populations

(Table S4), the distribution and median nucleotide diversity strongly differed by geography

(Figure 3A). The majority of populations south of Monterey Bay had a broader range of

nucleotide diversity and low median π, while the populations north of Monterey Bay have a very

narrow range of nucleotide diversity and higher π. Almost all populations, except the three

populations closest to Monterey Bay (SBR, PL and PGP) had negative median Tajima's D

values indicating an excess of rare alleles in each population (Figure 3B; Table S4). The 11 most

northern populations had mean Tajima's D values that were more negative than the five most

southern populations. This pattern is indicative of an excess of rare alleles in the northern

populations and is consistent with recent population expansion.

272
Figure 3. A) Nucleotide diversity ($Log_{10}(\pi)$), and B) Tajima's D across the 19 *Nucella*
canaliculata populations. Calculations were performed on a sliding window with a window size
of 25kb. The dashed vertical line for Tajima's D indicates 0, the expected value under a neutral
model. The colors match the 19 populations indicated in Table S1 and are ordered from north to
south. C) Within-population block-jackknife estimate ± standard error of heterozygosity from
`poolfstat`. Populations are ordered north to south from top to bottom. D) Distribution of AIC
values for the two demographic models tested using `moments`. The mean ± standard deviation of
AIC among all model replicates is shown (the best model is reported in the text). E) Hypothetical
levels of genetic variation derived from simulating the two proposed demographic scenarios using
`SLiM`.

Genomic Evidence of Post-Glacial Demography

We performed coalescent demographic inference with `moments` [54] to compare

whether our data matched either of two phylogeographic models: (1) a single southern refugium

expanding northward, and (2) a secondary contact (admixture) between two glacial refugia.

These models were tested on three central California populations (PGP, PL, and SBR) that

showed potential admixture signals in the genomic diversity analyses (Figure 3A). Our results

indicate that, for all three populations, the admixture model provides a better fit based on AIC

values (Figure 3D). The ΔAIC (Log_{10} transformed) values for the admixture model, relative to

the expansion model, were: PGP = 5.40; PL = 6.62; and SBR = 6.83.

For PGP, the best-fitting model estimated an admixture event around 10,500 years ago,

with a very low southern ancestry contribution (1%), indicating that most of its genetic ancestry

derives from northern populations. This aligns with its position as the southern boundary of the

hypothetical northern refugia. In contrast, PL and SBR exhibit strong signatures of admixture

between northern and southern populations. For PL, the best model estimated an admixture event

approximately 10,231 years ago, with 57% of its ancestry derived from southern populations.

Similarly, for SBR, admixture time was estimated at 7,451 years ago, with 59% southern

ancestry. These results suggest that PL and SBR are situated at the "core" of the admixture zone

between the hypothetical northern and southern glacial refugia.

To further investigate these patterns, we conducted forward genetic simulations using

`SLiM` [58] to compare the expected levels of genetic variation under our proposed demographic

scenarios. Among the two simulated models, the "two refugia" scenario produced patterns of

genetic variation most consistent with our empirical data. In this model, the core admixture

populations, i.e., SBR and PL, exhibited the highest levels of genetic variation (Figure 3E),

mirroring the observed data (Figure 3A). Additionally, in the simulations, genetic variation

decreased monotonically away from the admixture zone at a rate proportional to the simulated

population size of each refugium (in our simulations, refugium 2 had a smaller population size

relative to refugium 1). In contrast, the one refugium model produced a different pattern, in

which genetic variation steadily declined with range expansion, lacking the pronounced peaks in

diversity seen in the admixed populations.

Shell Geometric Morphometrics Show Patterns of Spatial Differentiation

Given the strong signature of genetic divergence among populations, we explored

whether shell morphology displayed a similar pattern of divergence. We performed landmark

analysis on images of the ventral surface of the dogwhelks (Figure 4A). PC1 (21.60% of the

variance) and PC2 (19.65%) explained most of the shape variation and geographically separated

the populations based on shell shape (Figure 4B; Figure S4). PC1 described a shell shape with a

lack of elongation of the spire and a rotundness of the main body whorl and aperture (wireframe

image for PC1 on Figure 4B). PC2 described shape changes in the widest point of the outer lip

and the angle of the siphonal canal. Dogwhelks from populations south of Monterey Bay occupy

a greater amount of PC space than the northern dogwhelks despite there being fewer populations

analyzed (Figure 4B). When comparing pairwise Procrustes distances, the greatest divergence in

shape was between the populations south of Monterey Bay (except HZD) with those from

northern California and southern Oregon (dark blue/teal section in the top right of Figure 4C).

Size (measured as centroid size) and population explained 5.5% and 21.6% of the dogwhelk

shape variation (Figure 4D; Table S7). Similarly, size, latitude and their interaction all explained

a significant amount of the variance in dogwhelk shape (Figure 4D; Table S7).

Figure 4. A) Diagram showing the position of the 15 landmarks used in the geometric morphometric analysis of *Nucella canaliculata* shells from 19 populations. B) Principal component 1 and 2 of the Procrustes coordinates. Wireframes display the shape variation for each PC (purple) in relation to the mean shape (black). Dogwhelks are colored by population. See Table S1 for an explanation of population codes. C) Heatmap of Procrustes distances among the 19 populations. D) The centroid size (log transformed) as a function of latitude. The dotted line indicates the location of Monterey Bay. Colors indicate the 19 populations as specified in B.

A Shell Matrix Protein Emerges as a Genomic Outlier of Local Adaptation

We used `BayPass` to identify outlier SNPs associated with morphological variation. Seven Bonferroni corrected outlier SNPs were identified associated with the coefficient of variation for PC1 (Figure 5A; Table S8), and six outlier SNPs were identified associated with the coefficient of variation for PC2 (Figure 5B; Table S8). Four candidate loci were in the same protein (gene g26813), with one SNP (position 41601) identified in both analyses (Table S8).

Most of the populations south of Monterey Bay were fixed at these four loci, while the

populations north of Monterey Bay were polymorphic with variable allele frequencies (Figure

5C). Three of the loci show strong associations with morphology — when the allele frequency of

reference allele approaches 0 the morphometric space becomes negative (Figure S5). To identify

the function of this protein of interest, we ran `blastp`. This protein has been identified as a

shell matrix protein in multiple gastropod species including the zebra mussel, *Dreissena*

polymorpha (UniProt entry: A0A9D4NEM1; percent identity 26.2%, positive matches = 52.4%,

E-value = 0.0002) [63], and the lettuce slug, *Elysia crispata* (UniProt entry: A0AAE1DT89;

percent identity 24.3%, positive matches = 45.0%, E-value = 0.00044) [64].

353
Figure 5. Signatures of selection from `BayPass` for the coefficient of variation A) for PC1 and
B) for PC2 of the *Nucella canaliculata* shell morphology. The solid blue line indicates Bonferroni
corrected outliers and the dotted blue line indicates significance of SNPs at a value of 0.001.
Bonferroni corrected outliers are identified in black, with the four candidate loci on gene g26813
in red. C) Allele frequencies of the four candidate loci on gene g26813 as a function of latitude.
The solid black line indicates the location of Monterey Bay.

<u>Discussion:</u>

The present-day genomic diversity and population structure of a species reflects a

combination of contemporary and historical processes [5]. In contrast to most marine species

which have high population connectivity due to planktonic larvae, the genomics of marine taxa

with direct development and low dispersal provide an especially powerful opportunity to analyze

deeper evolutionary histories and the impacts of historical events [19–21]. Our pooled

sequencing of 19 populations of the low-dispersing channeled dogwhelk, *Nucella canaliculata*,

across ~1,500 km of the west coast of North America indicated a high level of population

structure, with a phylogeographic split at Monterey Bay, and a pattern of genomic diversity that

revealed a history of being structured by past geological events. The contemporary genomic

patterns and accompanying computer simulations suggest that during the last glacial maximum

there were at least two refugial populations, north and south of Monterey Bay, that experienced

post-glacial expansions and admixture. This biogeographic pattern of divergence was further

supported by spatial variation in shell morphology. Additionally, we identified putatively

adaptive outlier loci that may be the molecular drivers of variation in shell morphology. Our

findings have important management implications; if the range of this species shifts poleward, as

predicted with climate change, there is risk of losing the unique genetic diversity that is primarily

maintained in the populations in the southern portion of the species' range limit [20,27,65].

Phylogeography and Glacial Refugia

Pleistocene glaciation drove changes in species distributions, with most temperate marine

species in the northern hemisphere retreating southward to escape glacial ice sheets [24,25].

Geological assemblages from Pleistocene terraces along the west coast of North America

indicate that many species, including *N. canaliculata*, shifted their ranges due to climatic

fluctuations in accord with shifting habitat conditions [24–26]. This historical event is assumed

to have left a signature on the current genetic composition of populations. Genetic models

suggest high genetic diversity within areas of glacial refugia and high dissimilarity between

refugial populations, and low genetic diversity in the area of subsequent population expansion

[3]. In marine environments, particularly the Northeastern Pacific, there has been mixed

evidence that this historical event has left imprints on the genetics of extant marine populations

[13,15,19,27]. The high dispersal potential present in most marine species [6] and the importance

of oceanographic processes and vicariant barriers in structuring genetic diversity typically

overwhelms any remnant signature of Pleistocene glaciation [15,66]. Additionally, genomic

advancements in SNP based approaches are increasing our phylogeographic resolution and

power to detect the persistent influence of glaciation [12,27].

The present-day phylogeographic structure of *N. canaliculata* showed clear evidence of

glacial refugia, followed by population expansion and admixture (Figure 2; Figure 3). Our results

revealed two genetically distinct clusters with Monterey Bay being the phylogeographic

breakpoint (Figure 1). All populations north of Monterey Bay were genetically similar and have

a negative Tajima's D value, indicating population expansion (Figure 3B). In contrast, the

populations between Monterey Bay and the species southern range limit of Point Conception

contained variable levels of nucleotide diversity and were highly diverged from each other

despite their spatial proximity (Figure 2B; Figure 3A). These patterns closely match those

predicted from genetic models of Pleistocene glaciation [3,7]. Additionally, the populations

nearest Monterey Bay show a unique signature of higher nucleotide diversity and heterozygosity

compared to the populations to the north and south, suggesting potential admixture in this region

(Figure 3).

Our `moments` analyses and `SLiM` simulation results provided greater clarity to this

glacial pattern as well as unique insights about the fine scale demography of the species,

particularly about the evolutionary histories of the populations nearest Monterey Bay. These

analyses suggest that at least one northern population refugium persisted through the cycles of

glaciation, rather than the entire contemporary species being derived from southerly populations

(Figure 3D; Figure 3E). These refugial populations expanded post-glacially and experienced a

secondary contact near Monterey Bay. Further, the admixture proportions show that the two

populations just south of Monterey Bay (Point Lobos [PL] and Soberanes Point [SBR]) are the

'true' admixed zone, while the region just north of Monterey Bay (i.e., Pigeon Point [PGP]) is

the likely southern edge of the northern refugia. The drastic drop in admixture proportions

between Point Lobos and Pigeon Point, in the context of the F_{ST} findings (Figure 2), suggests

that admixture was relatively recent, and that migration is low. An important caveat of our

analysis is that several key genetic parameters for *Nucella* remain uncharacterized; thus, we

relied on published rates from the model system *Drosophila*. Consequently, parameters related to

population divergence and admixture timing should be considered approximations and

interpreted with caution.

Genetic variation in the northern populations indicates that the phylogeographic history

of *N. canaliculata* may be more complex than suggested by our models of glacial refugia. Under

a simple scenario of a "large" southern refugia and a "small" (or remanent) northern refugia, we

should expect to see a sharp decline in the levels of standing genetic variation towards the

northern edge (Figure 3D). Yet, in our real data, we see a peak in variation at Soberanes Point

(SBR) and Point Lobos (PL), with variable variation in the southern populations and stable levels

of genetic variation in the northern populations that are, on average, higher than the southern

sites (Figure 3A). One possible hypothesis to explain this departure from expectations is that the

southern refugia was smaller than the northern refugia; this would be unusual, but not impossible

(see [4]). For example, strong competition with other snails in the refugia or limitations in prey

availability could have kept populations small [67]. An alternative hypothesis could be that there

were multiple northern refugia that experienced sequential mixing post-glaciation, giving rise to

the elevated variation. To clarify these patterns and determine the exact location and size of the

northern refugia, we would need thorough sampling of populations near the species northern

range limit, as well as ancient samples.

Understanding the present-day phylogeographic structure of a species is particularly

important as species are being forced to either rapidly adapt or shift their ranges in response to

anthropogenic climate change [9–11]. Most of the unique genetic diversity of *N. canaliculata*

(i.e., suggested by high F_{ST}) was in the populations south of Monterey Bay. Given that

populations at the warmer, trailing edge of a range are more likely to go extinct in response to

climate change, this could mean losing large amounts of endemic diversity [20,27], particularly

for species with low dispersal. In accord, a recent study that surveyed several coastal marine

species, including *N. canaliculata*, across their ranges in the northeast Pacific found that

populations near their equatorward edge are demographically vulnerable to climate change and

environmental stress [68]. Population density of *N. canaliculata* declined latitudinally from

Oregon to central California in association with increasing aerial and water temperature stress,

and a reduction in rocky habitat availability [68]. Small populations are highly vulnerable to

local extirpation due to Allee effects and stochastic events [69]. There is accumulating evidence

that marine heatwaves and other oceanic extreme events are becoming increasingly common

[70]. Thus, there is substantial risk that the genetically unique, but demographically small,

populations of *N. canaliculata* near their trailing edge could face local extirpation.

Adaptive consequences of low dispersal potential

The phylogeographic pattern of defined population structure and low levels of genetic

diversity that we observed has likely impacted the adaptive potential and evolutionary trajectory

of the *N. canaliculata* populations. Previous research studying intraspecific variation in

predatory drilling ability in *N. canaliculata* showed that the species displays a pattern of local

adaptation [30–33]. However, rather than a smooth latitudinal gradient, the adaptive landscape

appears to be a spatial mosaic, with specific sites deviating from regional patterns [32,33].

Despite their geographic separation, the two populations with the greatest drilling ability are

Soberanes Point (SBR) and Van Damme (VD). Our genomic results provide important

demographic context to this phenotypic pattern and suggest that these two populations have

strikingly different genetic backgrounds, with Soberanes Point (SBR) being an admixed

population and Van Damme (VD) being in the northern genetic cluster. Thus, this phenotypic

pattern of parallel adaptation could be due to the populations evolving strong drilling using

independent molecular pathways either via the same or different morphological mechanism. This

hypothesis is further supported by the fact that these two populations display very different

behavioral patterns when exposed to predator cues [34]. Further research that links genotype to

phenotype would be needed to elucidate the molecular mechanisms behind this adaptive

landscape.

For many species, linking phenotype to genotype and understanding patterns of

adaptation are significantly hindered by the lack of genomic resources for non-model organisms

[7,8]. The genome assembly presented here is the first for the species, and the second assembly

for the ecologically important genus *Nucella* [71]. Extraction of high-quality molluscan DNA is

a challenge due to the presence of mucopolysaccharides and other secondary metabolites [72].

Additionally, molluscan genomes are large, have a high proportion of repetitive regions, and

typically harbor high heterozygosity, thus genome assembly is computationally difficult and

relatively few genomes have been assembled, despite there being more than 100,000 living

molluscs and incredibly high morphological diversity [73,74]. Like previously published

molluscan genomes [71,73], the *N. canaliculata* assembly presented here was very large and

consisted of numerous scaffolds. Despite this, the assembly was smaller than the estimated

genome size for the species of 2.54pg±0.057 [75], a pattern not uncommon in molluscan

genomes [74].

Shell Morphology and Species Taxonomy

The study of gastropod shell morphology predates Linnaean taxonomy [37,38]. Given

their ubiquity and longevity, the calcareous shells of gastropods have long been used to infer

taxonomy and evolutionary patterns [76]. Despite this, few modern phylogeographic studies of

marine gastropods incorporate morphological data alongside genetic data (but see [19,77]). We

found evidence that *N. canaliculata* dogwhelks show spatial variation in shell morphology, with

the greatest difference between populations south versus north of Monterey Bay (Figure 4C),

consistent with patterns of geographic divergence suggested by the genomic data. Dogwhelks

from southern populations were typically more rotund and had a shorter spire, compared to

dogwhelks from northern populations (Figure 4B). Additionally, the size of the dogwhelks varied

latitudinally with the largest dogwhelks present at Soberanes Point (SBR) and Point Lobos (PL)

(Figure 4D). Shell morphology in *Nucella* is a result of both phenotypic plasticity [34] and

adaptive evolution [78]. Consequently, the morphometric data (Figure 4) showed a less stark

pattern of divergence among the populations than the genomic data (Figure 2). Notwithstanding,

the concordance of our datasets increases our confidence in the biogeographic pattern.

Our genomic and morphometric results both indicate a strong spatial signal of divergence

with a breakpoint at Monterey Bay. This biogeographic pattern in *N. canaliculata* matches an

overlooked taxonomic hypothesis, based on shell morphology, that there may be a northern and

southern subspecies divided at Monterey Bay [79]. In this context, what had previously been

hypothesized as two morphological subspecies, are according to our data, two

phylogeographically distinct genetic clusters whose morphological differences reflect divergent

histories of demography and selection. Future research with more intensive sampling,

particularly around Monterey Bay and at the species northern range limit, as well as individual

sequencing is necessary to resolve the taxonomy of the species.

To test for signatures of selection and begin to link genotype to phenotype, we ran

`BayPass` on the coefficient of variation for PC1 and PC2 of the shell morphology dataset. This

framework allowed us to identify outlier loci while controlling for population structure. We

identified a handful of SNPs associated with both coefficients of variation (Figure 5). Notably,

one protein stood out in both analyses. The four candidate loci on this protein display a spatial

pattern such that they are fixed in the populations south of Monterey Bay and polymorphic in the

populations north of Monterey Bay (Figure 5C). This protein has been identified as a shell matrix

protein in other gastropod species, including the zebra mussel, *Dreissena polymorpha* [63], and

the lettuce slug, *Elysia crispata* [64]. We hypothesize that this protein is likely a candidate for

morphological variation and local adaptation and should be studied further.

Overall, this research highlights the power of genomic approaches to advance our

understanding of how historical events have shaped contemporary demography in coastal

species. The geographic distribution of genomic diversity in *Nucella canaliculata* reveals a

persistent signature of glacial events in the northeast Pacific, with important implications for

adaptive capacity and conservation in an era of rapid global change.

Data Accessibility Statement: The *Nucella canaliculata* genome assembly can be found at

BioProject accession number PRJNA1282036 and the raw reads for the assembly can be found

under the BioProject accession number PRJNA1282048. Raw pooled-sequencing reads can be

found under the SRA BioProject accession number PRJNA1276871, and the individual

BioSample accession numbers can be found in Table S1. A GitHub repository with the code for

analyzing the data can be found at https://github.com/emily-

longman/Nucella_can_Pop_Genomics.

Ethics: Collections in California were performed under CA Fish & Wildlife Scientific Collecting

Permits S-191200004-19122-001 and S-190980002-23226-001, and specific use amendments S-

190980003-23226-001-01 and S-190980003-23226-001-02, and CA State Parks permit 24-820-

015 and amendment 24-820-015A. Collections in Oregon were performed under the taking

permit 27976.

Author's contributions: E.K.L.: conceptualization, data curation, formal analysis, funding

acquisition, investigation, methodology, project administration, resources, software, supervision,

validation, visualization, writing – original draft, writing – review & editing; J.C.B.N.: formal

analysis, software, visualization, writing – review & editing; E.S.: funding acquisition,

investigation, writing – review & editing; M.P.: conceptualization, methodology, resources,

supervision, writing – review & editing. All authors gave final approval for publication and

agreed to be held accountable for the work performed therein.

Funding: This research was supported by an NSF postdoctoral fellowship (OCE-2307933)

awarded to E.K.L. and the National Science Foundation grant OCE-1851462 awarded to E.S.

Acknowledgements: We thank the Vandenberg Space Force Base, the UC Reserve System and

the Abbott family for access to field sites. We thank N. Longman and J. Fosnight for assistance

with the field collections, and R. A. Bay, B. B. Cameron, and E. S. Nielsen for help with

optimizing the extraction protocol and discussions about sequencing approaches for the genome.

References:

[1] Hughes, A.R., Inouye, B.D., Johnson, M.T., Underwood, N., & Vellend, M. (2008).

Ecological consequences of genetic diversity. *Ecology Letters*, *11*(6), 609–623.

(https://doi.org/10.1111/j.1461-0248.2008.01179.x)

[2] Saccheri, I., & Hanski, I. (2006). Natural selection and population dynamics. *Trends in*

Ecology & Evolution, *21*(6), 341–347. (https://doi.org/10.1016/j.tree.2006.03.018)

[3] Hewitt, G.M. (1996). Some genetic consequences of ice ages, and their role in divergence

and speciation. *Biological journal of the Linnean Society*, *58*(3), 247–276.

(https://doi.org/10.1111/j.1095-8312.1996.tb01434.x)

[4] Maggs, C.A., Castilho, R., Foltz, D., Henzler, C., Jolly, M.T., Kelly, J., Olsen, J., Perez, K.E.,

Stam, W., Väinölä, R., et al. (2008). Evaluating signatures of glacial refugia for North

Atlantic benthic marine taxa. *Ecology, 89*(sp11), S108–S122. (https://doi.org/10.1890/08-

0257.1)

[5] Avise, J.C. (2000). *Phylogeography: the history and formation of species*. Harvard

University Press. (https://doi.org/10.2307/j.ctv1nzfgj7)

[6] Thorson, G. (1950). Reproductive and larval ecology of marine bottom invertebrates.

Biological Reviews of the Cambridge Philosophical Society, 25, 1–45.

(https://doi.org/10.1111/j.1469-185X.1950.tb00585.x)

[7] Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local

adaptation. *Nature Reviews Genetics*, *14*(11), 807–820. (https://doi.org/10.1038/nrg3522)

[8] Da Fonesca, R.R. Albrechtsen, A., Themudo, G.E., Ramos-Madrigal, J., Sibbesen, J.A.,

Maretty, L., Zepeda-Mendoza, M.L., Campos, P.F., Heller, R., & Pereira, R.J. (2016).

Next generation biology: sequencing and data analysis approaches for non-model

organisms. *Marine genomics, 30*, 3–13. (https://doi.org/10.1016/j.margen.2016.04.012)

[9] Waldvogel, A.M., Feldmeyer, B., Rolshausen, G., Exposito-Alonso, M., Rellstab, C., Kofler,

R., Mock, T., Schmid, K., Schmitt, I., Bataillon, T., et al. (2020). Evolutionary genomics

can improve prediction of species' responses to climate change. *Evolution Letters*, *4*(1),

4–18. (https://doi.org/10.1002/evl3.154)

[10] Urban, M.C., Swaegers, J., Stoks, R., Snook, R.R., Otto, S.P., Noble, D.W., Moiron, M.,

Hällfors, M.H., Gómez-Llano, M., Fior, S., et al. (2024). When and how can we predict

adaptive responses to climate change? *Evolution Letters*, *8*(1), 172–187.

(https://doi.org/10.1093/evlett/qrad038)

[11] Bernatchez, L., Ferchaud, A.L., Berger, C.S., Venney, C.J., & Xuereb, A. (2024). Genomics

for monitoring and understanding species responses to global climate change. *Nature*

Reviews Genetics, *25*(3), 165–183. (https://doi.org/10.1038/s41576-023-00657-y)

[12] Crawford, D.L., & Oleksiak, M.F. (2016). Ecological population genomics in the marine

environment. *Briefings in Gunctional Genomics*, *15*(5), 342–351.

(https://doi.org/10.1093/bfgp/elw008)

[13] Pelc, R.A., Warner, R.R., & Gaines, S.D. (2009). Geographical patterns of genetic structure

in marine species with contrasting life histories. *Journal of Biogeography*, 36(10), 1881–

1890. (https://doi.org/10.1111/j.1365-2699.2009.02138.x)

[14] Pascual, M., Rives, B., Schunter, C., & Macpherson, E. (2017). Impact of life history traits

on gene flow: A multispecies systematic review across oceanographic barriers in the

Mediterranean Sea. *PLoS One*, *12*(5), e0176419.

(https://doi.org/10.1371/journal.pone.0176419)

[15] Kelly, R.P., & Palumbi, S.R. (2010). Genetic structure among 50 species of the northeastern

Pacific rocky intertidal community. *PloS one*, *5*(1), e8594.

(https://doi.org/10.1371/journal.pone.0008594)

[16] Grosberg R.K. & C.W. Cunningham. (2001). Genetic structure in the sea: from populations

to communities. In Marine Community Ecology, ed. M.D. Bertness, S.D. Gaines, M.E.

Hay, pp. 61–84. Sunderland, MA: Sinauer

[17] Riginos, C., Douglas, K.E., Jin, Y., Shanahan, D.F., & Treml, E.A. (2011). Effects of

geography and life history traits on genetic differentiation in benthic marine

fishes. *Ecography*, *34*(4), 566–575. (https://doi.org/10.1111/j.1600-0587.2010.06511.x)

[18] Arranz, V., Fewster, R.M., & Lavery, S.D. (2024). Concordance and drivers of marine

spatial structure determined using genogeographic clustering. *Journal of*

Biogeography, *51*(10), 2034–2046. (https://doi.org/10.1111/jbi.14967)

[19] Hellberg, M.E., Balch, D.P., & Roy, K. (2001). Climate-driven range expansion and

morphological evolution in a marine gastropod. *Science, 292*(5522), 1707–1710.

(https://doi.org/10.1126/science.1060102)

[20] Fenberg, P.B., Posbic, K., & Hellberg, M.E. (2014). Historical and recent processes shaping

the geographic range of a rocky intertidal gastropod: phylogeography, ecology, and

habitat availability. *Ecology and Evolution*, *4*(16), 3244–3255.

(https://doi.org/10.1002/ece3.1181)

[21] Jacobs, D.K., Haney, T.A., & Louie, K.D. (2004). Genes, diversity, and geologic process on

the Pacific coast. *Annual Review of Earth and Planetary Sciences*, *32*(1), 601–652.

(https://doi.org/10.1146/annurev.earth.32.092203.122436)

[22] McGaughran, A., Liggins, L., Marske, K.A., Dawson, M.N., Schiebelhut, L.M., Lavery,

S.D., Knowles, LL., Moritz, C. & Riginos, C. (2022). Comparative phylogeography in

the genomic age: Opportunities and challenges. *Journal of Biogeography*, *49*(12), 2130–

2144. (https://doi.org/10.1111/jbi.14481)

[23] Bringloe, T.T., Fort, A., Inaba, M., Sulpice, R., Ghriofa, C.N., Mols-Mortensen, A., Filbee-

Dexter, K., Vieira, C., Kawai, H., Hanyuda, T., et al. (2022). Whole genome population

structure of North Atlantic kelp confirms high-latitude glacial refugia. *Molecular*

Ecology, *31*(24), 6473–6488. (https://doi.org/10.1111/mec.16714)

[24] Addicott, W.O. (1966). *Late Pleistocene marine paleoecology and zoogeography in central*

California. US Government Printing Office. (https://doi.org/10.3133/pp523C)

[25] Fields, P.A., Graham, J.B., Rosenblatt, R.H. & Somero, G.N. (1993). Effects of expected

global climate change on marine faunas. *Trends in Ecology & Evolution, 8*(10), 361–367.

(https://doi.org/10.1016/0169-5347(93)90220-J)

[26] Roy, K., Valentine, J.W., Jablonski, D., & Kidwell, S.M. (1996). Scales of climatic

variability and time averaging in Pleistocene biotas: implications for ecology and

evolution. *Trends in ecology & evolution*, *11*(11), 458–463.

(https://doi.org/10.1016/0169-5347(96)10054-9)

[27] Nielsen, E.S., Walkes, S., Sones, J.L., Fenberg, P.B., Paz-García, D.A., Cameron, B.B.,

Grosberg, R.K., Sanford, E. & Bay, R.A. (2024). Pushed waves, trailing edges, and

extreme events: Eco-evolutionary dynamics of a geographic range shift in the owl limpet,

Lottia gigantea. Global Change Biology, *30*(7), e17414.

(https://doi.org/10.1111/gcb.17414)

[28] Dawson, M.N. (2001). Phylogeography in coastal marine animals: a solution from

California? *Journal of Biogeography, 28*(6), 723–736. (https://doi.org/10.1046/j.1365-

2699.2001.00572.x)

[29] Marko, P.B., Moran, A.L., Kolotuchina, N.K., & Zaslavskaya, N.I. (2014). Phylogenetics of

the gastropod genus Nucella (Neogastropoda: Muricidae): species identities, timing of

diversification and correlated patterns of life-history evolution. *Journal of Molluscan*

Studies, 80(4), 341–353. (https://doi.org/10.1093/mollus/eyu024)

[30] Sanford, E., Roth, M.S., Johns, G.C., Wares, J.P., & Somero, G.N. (2003). Local selection

and latitudinal variation in a marine predator-prey interactions. *Science, 300*(5622),

1135–1137. (https://doi.org/10.1126/science.1083437)

[31] Sanford, E. & Worth, D.J. (2010). Local adaptation along a continuous coastline: prey

recruitment drives differentiation of a predatory snail. *Ecology, 91*(3), 891–901.

(https://doi.org/10.1890/09-0536.1)

[32] Sanford, E. & Worth, D.J. (2009). Genetic differences among populations of a marine snail

drive geographic variation in predation. *Ecology, 90*(11), 3108–3118.

(https://doi.org/10.1890/08-2055.1)

[33] Longman, E.K. & Sanford, E. (2025). Biogeographic variation in mussel shell thickness and

657   drilling predation on rocky shores. *Oecologia, 207*, 126. (https://doi.org/10.1007/s00442-

658   025-05760-x)

659   [34] Neylan, I.P., Longman, E.K., Sanford, E., Stachowicz, J.J., & Sih, A. (2024). Long-term

660   anti-predator learning and memory differ across populations and sexes in an intertidal

661   snail. *Proceedings B, 291*(2032), 20240944. (https://doi.org/10.1098/rspb.2024.0944)

662   [35] Kuo, E.S., & Sanford, E. (2009). Geographic variation in the upper thermal limits of an

663   intertidal snail: implications for climate envelope models. *Marine Ecology Progress*

664   *Series*, *388*, 137–146. (https://doi.org/10.3354/meps08102)

665   [36] Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of

666   individuals—mining genome-wide polymorphism data without big funding. *Nature*

667   *Reviews Genetics*, *15*(11), 749–763. (https://doi.org/10.1038/nrg3803)

668   [37] Fretter, V. & Graham, A. (1962). British Prosobranch Molluscs; Their functional anatomy

669   and ecology. London: Ray Soc. 755 pp.

670   [38] Bieler, R. (1992). Gastropod phylogeny and systematics. *Annual Review of Ecology and*

671   *Systematics*, 311–338.

672   [39] Ye, C., Hill, C. M., Wu, S., Ruan, J., & Ma, Z. (2016). DBG2OLC: efficient assembly of

673   large genomes using long erroneous reads of the third generation sequencing

674   technologies. *Scientific Reports*, 6(1), 31900. (https://doi.org/10.1038/srep31900)

675   [40] Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: quality assessment

676   tool for genome assemblies. *Bioinformatics, 29*(8), 1072–1075.

677   (https://doi.org/10.1093/bioinformatics/btt086)

[41] Manni, M., Berkeley, M.R., Seppey, M., Simão, F.A., Zdobnov, E.M. (2021). BUSCO

update: novel and streamlined workflows along with broader and deeper phylogenetic

coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology*

and Evolution, 38(10), 4647–4654. (https://doi.org/10.1093/molbev/msab199)

[42] Smit, A.F.A., Hubley, R., Green P. (2013-2015). RepeatMasker Open-4.0.

http://www.repeatmasker.org

[43] Stanke, M., Keller, O., Gunduz, I., Hayes, A., Waack, S., & Morgenstern, B. (2006).

AUGUSTUS: ab initio prediction of alternative transcripts. *Nucleic Acids Research, 34*,

W435–W439. (https://doi.org/10.1093/nar/gkl200)

[44] Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data.*

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

[45] Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). Fastp: An ultra-fast all-in-one FASTQ

processor. *Bioinformatics, 34*(17), i884–i890.

(https://doi.org/10.1093/bioinformatics/bty560)

[46] Vasimuddin, M. Sanchit, M., Henge, L., Srinivas A. (2019). Efficient architecture-aware

acceleration of BWA-MEM for multicore systems. *IEEE Parallel and Distributed*

Processing Symposiums (IPDPS). (https://doi:10.1109/IPDPS.2019.00041)

[47] Li, J., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,

G., Durbin, R, 1000 Genome Project Data Processing Subgroup. (2009) The sequence

alignment/map format and SAMtools. *Bioinformatics, 25*(16), 2078–2079.

(https://doi.org/10.1093/bioinformatics/btp352)

[48] Broad Institute. (2019). Picard Toolkit. GitHub Repository.

https://broadinstitute.github.io/picard/

[49] Garrison, E., & Marth, G. (2012). Haplotype-based variant detection from short-read

sequencing. *arXiv*, 1207.3907. (https://doi.org/10.48550/arXiv.1207.3907)

[50] Gautier, M., Vitalis, R., Flori, L., & Estoup, A. (2022). f-Statistic estimation and admixture

graph construction with Pool-seq or allele count data using the R package poolfstat.

Molecular Ecology Resources, 22, 1394–1416. (https://doi.org/10.1111/1755-

0998.13557)

[51] Gautier, M. (2015). Genome-wide scan for adaptive divergence and association with

population-specific covariates. *Genetics, 201*(4), 1555–1579.

(https://doi.org/10.1534/genetics.115.181453)

[52] Paradis, E., & Schliep, K. (2019). Ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics, 35*, 526–528.

(https://doi.org/10.1093/bioinformatics/bty633)

[53] Ferretti, L., Ramos-Onsins, S.E., & Pérez-Enciso, M. (2013). Population genomics from

pool sequencing. *Molecular ecology*, *22*(22), 5561–5576.

(https://doi.org/10.1111/mec.12522)

[54] Jouganous, J., Long, W., Ragsdale, A.P., & Gravel, S. (2017). Inferring the joint

demographic history of multiple populations: beyond the diffusion

approximation. *Genetics*, *206*(3), 1549–1567.

(https://doi.org/10.1534/genetics.117.200493)

[54] Kapun, M., Nunez, J. C., Bogaerts-Márquez, M., Murga-Moreno, J., Paris, M., Outten, J.,

Coronado-Zamora, M., Tern, C., Rota-Stabelli, O., García Guerreiro, M.P., et al. (2021).

Drosophila evolution over space and time (DEST): a new population genomics

723 resource. *Molecular biology and evolution*, *38*(12), 5782–5805.

724 (https://doi.org/10.1093/molbev/msab259)

725 [56] Nunez, J.C.B., Coronado-Zamora, M., Gautier, M., Kapun, M., Steindl, S., Ometto, L.,

726 Hoedjes, K.M., Beets, J., Wiberg, R.A.W., Mazzeo, G.R., et al. (*in press*) Footprints of

727 worldwide adaptation in structured populations of *D. melanogaster* through the expanded

728 DEST 2.0 genomic resource. *Molecular biology and evolution.*

729 [57] Keightley, P.D., Ness, R.W., Halligan, D.L., & Haddrill, P.R. (2014). Estimation of the

730 spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib

731 family. *Genetics, 196*(1), 313–320. (https://doi.org/10.1534/genetics.113.158758)

732 [58] Haller, B.C., & Messer, P.W. (2023). SLiM 4: multispecies eco-evolutionary modeling. *The*

733 *American Naturalist*, *201*(5), E127–E139. (https://doi.org/10.1086/723601)

734 [59] Rohlf, F.J. (2015). The tps series of software. *Hystrix, 26*(1), 9–12.

735 (https://doi.org/10.4404/hystrix-26.1-11264)

736 [60] Adams, D.C., Collyer, M., Kaliontzopoulou, A., & Baken, E. (2025). *Geomorph: Geometric*

737 *morphometric analyses of 2D and 3D landmark data*. R package version 4.0.10.

738 (https://doi.org/10.32614/CRAN.package.geomorph)

739 [61] Klingenberg, C.P. (2011). MorphoJ: an integrated software package for geometric

740 morphometrics. *Molecular ecology resources*, *11*(2), 353-357.

741 (https://doi.org/10.1111/j.1755-0998.2010.02924.x)

742 [62] Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X., &

743 Ruden, D.M. (2012). A program for annotating and predicting the effects of single

744 nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster*

745 strain w1118; iso-2; iso-3. *fly*, *6*(2), 80–92. (https://doi.org/10.4161/fly.19695)

[63] McCartney, M.A., Auch, B., Kono, T., Mallez, S., Zhang, Y., Obille, A., Becker, A.,

Abrahante, J.E., Garbe, J., Badalamenti, J.P. et al. (2022). The genome of the zebra

mussel, *Dreissena polymorpha*: a resource for comparative genomics, invasion genetics,

and biocontrol. *G3*, *12*(2), jkab423. (https://doi.org/10.1093/g3journal/jkab423)

[64] Eastman, K.E., Pendleton, A.L., Shaikh, M.A., Suttiyut, T., Ogas, R., Tomko, P., Gavelis,

G., Widhalm, J.R. & Wisecaver, J.H. (2023). A reference genome for the long-term

kleptoplast-retaining sea slug *Elysia crispata* morphotype clarki. *G3: Genes, Genomes,*

Genetics, *13*(12), jkad234. (https://doi.org/10.1093/g3journal/jkad234)

[65] Zarzyczny, K.M., Hellberg, M.E., Lugli, E.B., MacLean, M., Paz-García, D.A., Rius, M.,

Ross, E.G., Treviño Balandra, E.X., Vanstone, J., Williams, S.T. & Fenberg, P.B. (2024).

Opposing genetic patterns of range shifting temperate and tropical gastropods in an area

undergoing tropicalisation. *Journal of Biogeography*, *51*(2), 246–262.

(https://doi.org/10.1111/jbi.14744)

[66] Wares, J.P., & Cunningham, C.W. (2001). Phylogeography and historical ecology of the

north Atlantic intertidal. *Evolution*, *55*(12), 2455–2469. (https://doi.org/10.1111/j.0014-

3820.2001.tb00760.x)

[67] Spight, T.M., & Emlen, J. (1976). Clutch sizes of two marine snails with a changing food

supply. *Ecology*, *57*(6), 1162–1178. (https://doi.org/10.2307/1935042)

[68] Gravem, S.A., Bachhuber, S., Bignami, S., Chiachi, A.E., Field, L.C., Gaddam, R.N.,

Raimondi, P.T., & Menge, B.A. (2025). Biogeographic Patterns in Density, Recruitment,

Body Size and Zonation of Rocky Intertidal Predators Suggest Increased Population

Vulnerability Near Southern Range Limits. *Journal of Biogeography*, *52*(2), 257–273.

(https://doi.org/10.1111/jbi.15029)

[69] Stephens, P.A., & Sutherland, W.J. (1999). Consequences of the Allee effect for behaviour,

ecology and conservation. *Trends in ecology & evolution*, *14*(10), 401–405.

(https://doi.org/10.1016/S0169-5347(99)01684-5)

[70] Smith, K.E., Aubin, M., Burrows, M.T., Filbee-Dexter, K., Hobday, A. J., Holbrook, N.J.,

King, N.G., Moore, P.J., Sen Gupta, A., Thomsen, M. et al. (2024). Global impacts of

marine heatwaves on coastal foundation species. *Nature Communications*, *15*(1), 5052.

(https://doi.org/10.1038/s41467-024-49307-9)

[71] Kadiwala, J., Hesketh, A., De Weerd, H., Ritch, H., & Shakur, R. (2025). The first genome

Assembly of the Dogwhelk *Nucella lapillus*, a bioindicator species for the marine

environment. *Scientific Data*, *12*(1), 1–8. (https://doi.org/10.1038/s41597-025-04764-9)

[72] Adema, C. M. (2021). Sticky problems: extraction of nucleic acids from

molluscs. *Philosophical Transactions of the Royal Society B*, *376*(1825), 20200162.

(https://doi.org/10.1098/rstb.2020.0162)

[73] Sun, J., Li, R., Chen, C., Sigwart, J.D., & Kocot, K.M. (2021). Benchmarking Oxford

Nanopore read assemblers for high-quality molluscan genomes. *Philosophical*

Transactions of the Royal Society B, *376*(1825), 20200160.

(https://doi.org/10.1098/rstb.2020.0160)

[74] Chen, Z., Baeza, J.A., Chen, C., Gonzalez, M.T., González, V.L., Greve, C., Kocot, K.M.,

Arbizu, P.M., Moles, J., Schell, T., et al. (2025). A genome-based phylogeny for

Mollusca is concordant with fossils and morphology. *Science, 387*(6737), 1001–1007.

(https://doi.org/10.1126/science.ads0215)

[75] Pascoe, P.L. Jha, A.N., & Dixon, D.R. (2004). Variation of karyotype composition and

genome size in some muricid gastropods from the northern hemisphere. *Journal of*

Molluscan Studies, 70(4), 389–398. (https://doi.org/10.1093/mollus/70.4.389)

[76] Vermeij, G. J. (1993). *Evolution and escalation: an ecological history of life*. Princeton

University Press. (https://doi.org/10.2307/j.ctv18zhf8b)

[77] Tirado, T., Saura, M., Rolán-Alvarez, E., & Quesada, H. (2016). Historical biogeography of

the marine snail Littorina saxatilis inferred from haplotype and shell morphology

evolution in NW Spain. *PLoS One*, *11*(8), e0161287.

(https://doi.org/10.1371/journal.pone.0161287)

[78] McIntire, L.C., Kota, T., Flagor, T.E., & Bourdeau, P.E. (2021). Weak directional selection

on inducible defensive shell traits in a marine gastropod by the inducing shell-breaking

predator. *Marine Biology, 168*, 1–12. (https://doi.org/10.1007/s00227-021-03860-9)

[79] McLean, J.H. (2006). Hypothesis for the recognition of *Nucella analoga* (Forbes, 18520 in

the northeastern Pacific. *Festivus, 38*, 17–21.