A thalamic hub integrates brainstem and stress signals to dynamically regulate REM

sleep

Aletta Magyar[1,2,3,4], Péter Berki[1,3], Sándor Borbély[1], Boglárka Barsy[1,2], Anna Virág

Bakacsi[1,2,3], Magor L Lőrincz[5], Danqian Liu[6], Ferenc Mátyás[1,2,7]

1. Institute of Experimental Medicine, Budapest, Hungary;

2. Institute of Cognitive Neuroscience and Psychology, Research Centre for Natural Sciences,

Budapest, Hungary;

3. János Szentágothai Doctoral School of Neurosciences, Semmelweis University, Budapest,

Hungary;

4. Institute of Biophysics and Radiation Biology, Semmelweis University, Budapest,

Hungary;

5. Department of Physiology, University of Szeged, Szeged, Hungary;

6. Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of

Sciences, China-Hungary Belt-and-Road Joint laboratory on Brain Science, Shanghai, China;

7. Department of Anatomy and Histology, University of Veterinary Medicine, Budapest,

Hungary

Corresponding author: Ferenc Mátyás (matyasf@koki.hu).

Abstract

Rapid eye movement (REM) sleep, a brain state critical for sleep quality and cortical

cognition, is tightly controlled by brainstem circuits and highly sensitive to stress. Yet, how

these systems interact to regulate REM sleep and its associated forebrain rhythms remains

elusive. Here, we identify a thalamic hub that conveys medullary REM-promoting signals,

and integrates stress inputs to regulate REM sleep. A subpopulation of paraventricular

thalamic neurons that collaterally project to the cortex and nucleus accumbens (PVT$_{\rightarrow NAc}$)

selectively responds to activation of medulla REM-promoting neurons, and bidirectionally

modulates REM-associated theta oscillations in an activity-dependent manner. Their low-

frequency activation promotes theta rhythms during REM sleep, while high-frequency

activation suppresses them, mirroring the neuronal signatures of acute and chronic stress,

respectively. Distinct patterns of PVT$_{\rightarrow NAc}$ neurons underlie the bidirectional stress

modulation of REM sleep, partly by differentially engaging prefrontal microcircuits via their

collateral projections. Together, our findings uncover a thalamic integrative hub that couples

sleep-regulating and stress pathways to adaptively control REM sleep expression.

Introduction

Neuronal oscillations of cortical networks are fundamental for regulating brain states and influencing cognitive functions. In particular, Rapid Eye Movement (REM) sleep has been linked to key cognitive processes such as memory formation and problem solving, in both human and mice[1–5]. Furthermore, disruption of REM sleep and the associated theta oscillation (4-8 Hz) has been implicated in various psychiatric disorders, including post-traumatic stress disorder, anxiety disorders, cognitive impairments in Parkinson's disease as well as Alzheimer's disease[1,6–9]. These findings suggest that cortical REM rhythm could serve as valuable biomarker for the development of such pathologies.

REM sleep has been shown to be regulated by brain-wide neuronal networks [10–13]. Brainstem circuits, including the medullar GABAergic signals, have been identified as crucial executive hubs for generating REM sleep and cortical theta oscillation[13–16]. Recent evidence has highlighted that cortical regions, whose excitation–inhibition balance is altered during REM sleep, is directly involved in the regulation of REM sleep[17,18]. However, in the absence of direct cortical innervation, it remains unclear how medullar ascending signals are transmitted to forebrain circuits to modulate REM sleep theta oscillations, and consequently, regulate REM states.

The stress-sensitive nature of REM sleep implies the existence of a neural switching system capable of integrating sleep-, salience- and stress-related information from brainstem and midbrain structures[19,20]. Cortical functions related to sleep and cognition are strongly influenced by its thalamic inputs[21–23], particularly those originating from the midline thalamic paraventricular nucleus (PVT)[24–26]. Neuronal activity of PVT plays a key role in encoding salience, arousal and homeostatic signals within the forebrain[25–29], as well as in regulating behavioral responses to stress[30–34]. Notably, the PVT receives broad ascending inputs from the brainstem and sends extensive projections to the prefrontal cortex (PFC). This positions the

PVT as a candidate link between medullary and cortical mechanisms of REM sleep that can

integrate arousal and stress information.

Thus, in this work, by combining circuit tracing, optogenetic manipulation, and cell-

type-specific electrophysiological recordings, we identified the engagement of the PVT in the

regulation of REM sleep and the associated theta oscillation at the level of PFC operation.

Our data show that a subpopulation within the PVT that send collateral projections to the PFC

and nucleus accumbens ($PVT_{\rightarrow NAc}$) was selectively activated by direct REM-promoting

VLM_{GABA} inputs, and display REM sleep-associated activation. The activity of $PVT_{\rightarrow NAc}$ cells

not only regulates theta oscillations and the associated excitation/inhibition balance in the

PFC during physiological sleep, but also recruits adaptive mechanisms in stress-dependent

manner. Taken together, our findings uncover a brainstem–thalamus–cortex circuit through

which the paraventricular thalamus dynamically regulates REM sleep and cortical theta

oscillations, providing a framework for understanding how sleep and stress interact to shape

higher-order brain functions.

Results

PVT transmits VLM_{GABA} signal to the PFC

We first examined whether activation of the REM-promoting ventrolateral

GABAergic neurons (VLM_{GABA}) could affect cortical theta oscillations, as it was earlier

reported in drug-free condition[14], by combining cell-type-specific optogenetics and

electrophysiological recordings in the PFC. Specifically, VLM_{GABA} cells were virally

transduced with Cre-dependent recombinant adeno-associated virus (AAV) expressing

Channelrhodopsin 2 (ChR2) in vGAT-Cre mice (Fig. 1a). After 3-4 weeks of recovery,

VLM_{GABA} neurons were optogenetically stimulated, which – indeed – evoked an increase in

cortical theta oscillatory activity in all tested mice, measured in the 4-8 Hz frequency band

(Fig. 1b and Extended Data Fig. 1, also see *Methods*).

To search for a neuronal pathway through which the VLM$_{GABA}$ signal can reach PFC,

we mapped the brain-wide efferentation of VLM$_{GABA}$ neurons. First, we confirmed that

cortical regions in general, including the PFC, receive no or negligible direct innervation from

the VLM$_{GABA}$ neurons (Extended Data Fig. 2). Conversely, widespread VLM$_{GABA}$ inputs were

observed subcortically, in the brainstem and hypothalamus, with dense axonal arborization

also observed in the PVT. A quantitative analysis revealed that VLM$_{GABA}$ neurons

topographically projected to the PVT with the strongest input localized to the middle-

posterior region (Fig. 1c,d), where PFC-projecting PVT neurons were identified by retrograde

labeling (Fig. 1e,f). These suggest that these PVT neurons may represent a key relay linking

medullary REM-sleep promoting activity to cortical networks.

To demonstrate functional involvement of the PVT in REM sleep, we next examined

the activity of PVT neurons across sleep-wake cycles. To target PVT neurons, we used the

calretinin (CR, '*Calb2*')-Cre mice, as the majority of PFC-projecting PVT cells express

calretinin[25]. We injected the PVT of CR-Cre mice with a Cre-dependent recombinant AAV

expressing ChR2. Three weeks later, we implanted optrode-tetrode configured wire bundles

surrounding an optical fiber to record the activity of optogenetically identified PVT cells

during natural home-cage sleep. Simultaneously, EEG and EMG electrodes were placed

above the frontal cortex and in the neck muscle to classify brain states (awake, non-REM,

REM) based on theta (4-8 Hz for REM) and delta (1-4 Hz for NREM) oscillations, together

with EMG signals (Extended Data Fig. 1a and 3a).

Analysis of identified PVT cells (N = 34 cells, n = 3 mice; Extended Data Fig. 3b)

revealed diverse state-dependent activity modulations (Extended Data Fig. 3c-e). At

population level, PVT neurons fired at significantly higher rates during REM sleep compared

to NREM sleep (Extended Data Fig. 3c).

The most pronounced increase in the PVT firing occurred at the NREM-to-REM transitions

(Extended Data Fig. 3e,f) highlighting their REM-dependence activity pattern. Moreover, a

negative correlation was also observed between baseline firing rate and REM-induced firing

rate changes, indicating that neurons with lower baseline activity exhibited greater REM-

associated activation (Extended Data Fig. 3d).

Next, we examined whether VLM_{GABA} signaling contributes to shaping the sleep

stage-dependent activity of PVT neurons and, in turn, cortical oscillations. To do so, we

virally expressed NpHR3.0 in VLM_{GABA} cells of vGAT-Cre mice (n = 7) and performed

simultaneous in vivo extracellular recordings from PVT and PFC while optogenetically

inhibiting VLM_{GABA}-to-PVT terminals under anesthesia (Fig. 1g). This configuration allowed

us to perform cell-type-selective thalamic recordings over extended periods. It also enabled us

to differentiate local field potential signal (LFP) from superficial (layers 1-3) and deep (layers

5-6) PFC regions, given that sleep oscillations exhibit laminar profiles, with higher REM

associated cortical activity in superficial layers[18,35–37]. Consistent with the heterogeneous sleep

state-dependent activity of PVT cells (Extended Data Fig. 3c,d,g,h), inhibition of VLM_{GABA}

input produced diverse firing rate changes in individual PVT cells, particularly during REM-

like high-theta periods (N=95 cells, n = 7 mice; Fig. 1h). Notably, VLM_{GABA} modulated cells

(PVT_{MOD}; N = 68) exhibited lower baseline firing rates than non-modulated ones (PVT_{MOD}, N

= 68: 0.74±0.85 Hz; PVT_{NON}, N = 27: 1.85±1.97 Hz), and only PVT_{MOD} neurons showed

increased activity during the NREM-to-REM transitions (Fig. 1i,j and Extended Data Fig.

2a,b). Moreover, REM-associated firing rate changes correlated with baseline activity only in

PVT_{MOD} but not in PVT_{NON} neurons (Fig. 1k) and only firing of PVT_{MOD} neurons was phase-

locked to the ascending theta cycle, suggesting their involvement in promoting theta

oscillations (Fig. 1l).

We also found that optogenetic silencing of VLM_{GABA} input, both non-selectively and

specifically during high-theta states, disrupted ongoing theta oscillations in superficial but not

in deep PFC layers (Fig. 1m and Extended Data Fig. 4c-j). Notably, REM-associated silencing

exerted a stronger theta-suppressive effect in 6 out of 7 mice (Extended Data Fig. 4k).

Altogether, these findings indicate that a subpopulation of PVT neurons is under REM-related

VLM_{GABA} control and influences theta oscillations in a VLM_{GABA}-dependent manner.

$PVT_{\rightarrow NAc}$ neurons selectively mediate the REM sleep-promoting and theta-modulating

effect of VLM_{GABA} neurons

A subset of low firing PVT neurons mediates VLM-driven theta oscillatory changes.

The PVT-to-nucleus accumbens ($PVT_{\rightarrow NAc}$) and PVT-to-amygdala ($PVT_{\rightarrow AMY}$) pathways

represent the two largest PVT subpopulations, each with distinct connectivity and

functions[25,27,38,39]. To determine which subgroup is under VLM_{GABA} control and exerts a

laminar-selective influence on cortical oscillation and REM sleep, we employed multiple

approaches. First, we retrogradely labeled $PVT_{\rightarrow NAc}$ or $PVT_{\rightarrow AMY}$ neurons using rgAAV-

eYFP while VLM_{GABA} cells were transduced with AAV carrying ChR2 in vGAT-Cre mice.

After 8–10 weeks, we conducted acute slice electrophysiology on coronal slices from the

middle-posterior PVT (Fig. 2a,b). Optogenetic activation of VLM_{GABA}-to-PVT fibers evoked

$GABA_A$-mediated inhibitory postsynaptic potentials (eIPSCs) in both $PVT_{\rightarrow NAc}$ and

$PVT_{\rightarrow AMY}$ neurons (Fig. 2c). However, eIPSCs amplitudes were significantly larger than

spontaneous IPSCs in $PVT_{\rightarrow NAc}$ but not $PVT_{\rightarrow AMY}$ neurons (Fig. 2d). Additionally, $PVT_{\rightarrow NAc}$

neurons exhibited stronger and more reliable inhibitory responses than $PVT_{\rightarrow AMY}$ neurons

across all stimulation frequencies (Fig. 2e,f). While VLM_{GABA}-to-$PVT_{\rightarrow AMY}$ synapses

displayed short-term depression at 5/10/20 Hz, VLM_{GABA}-to-$PVT_{\rightarrow NAc}$ synapses maintained

stable transmission (Fig. 2e,f), suggesting that $PVT_{\rightarrow NAc}$ neurons more efficiently relay

VLM_{GABA} inputs.

To confirm the selective involvement of PVT$_{\rightarrow NAc}$ neurons in REM sleep, we used

retrograde labeling[25] in CR-Cre mice to express ChR2 in either PVT$_{\rightarrow NAc}$ or PVT$_{\rightarrow AMY}$

neurons (Extended Data Fig. 5a-g), and recorded their firing rates across natural sleep stages

(Fig. 2g and Extended Data Fig. 6a,b). PVT$_{\rightarrow NAc}$ neurons had lower baseline activity and

stronger REM modulation compared to PVT$_{\rightarrow AMY}$ neurons (Fig. 2h and Extended Data Fig.

6b,c). Only PVT$_{\rightarrow NAc}$ neurons increased their firing at NREM-to-REM transitions (Fig. 2i and

Extended Data Fig. 6d) and showed a negative correlation between baseline activity and

REM-induced firing changes (Fig. 2j). Furthermore, PVT$_{\rightarrow NAc}$ firing was phase-locked to the

ascending theta cycle, whereas PVT$_{\rightarrow AMY}$ firing aligned with the descending phase,

suggesting a PVT$_{\rightarrow NAc}$-mediated role in theta oscillations (Fig. 2k) resembling VLM$_{GABA}$-

modulated PVT neurons (Fig. 1k,l). These findings establish PVT$_{\rightarrow NAc}$ as the key PVT

To directly asses the causal role of PVT$_{\rightarrow NAc}$ and PVT$_{\rightarrow AMY}$ neurons in REM sleep, we

optogenetically activated these neurons at two different frequencies during natural sleep.

Since PVT$_{\rightarrow NAc}$ activity naturally rises to ∼1 Hz before NREM-to-REM transitions (Fig. 2i)

and maintains this rate throughout REM bouts, we applied a 1 Hz stimulation protocol. In

addition, a 10 Hz protocol was also carried out as activity of PVT cells were shown to reach

this frequency[25]. Optogenetic activation of PVT$_{\rightarrow NAc}$ activation at 1 Hz (n = 7 mice) enhanced

REM sleep, increasing theta band power and REM episode duration (Fig. 2l,n and Extended

Data Fig. 6e,g). In contrast, 10 Hz PVT$_{\rightarrow NAc}$ activation suppressed REM-associated theta

oscillations (Fig. 2m,n, and Extended Data Fig. 6f,h). Furthermore, PVT$_{\rightarrow NAc}$ activations

triggered sleep-stage transitions with greater probability and efficacy and produced more

pronounced REM disruptions, compared to PVT$_{\rightarrow AMY}$ (n = 8) or control PVT$_{YFP}$ stimulations

(n =3; Fig. 2l-o and Extended Data Fig. 6e-j). Together, these results indicate that PVT$_{\rightarrow NAc}$

neurons exert bidirectional control over REM sleep in an activity-dependent manner.

PVT$_{\rightarrow NAc}$ bidirectionally modulate the VLM$_{GABA}$ effects on theta oscillation

The previous results collectively suggest that PVT$_{\rightarrow NAc}$ is able to maintain or suppress

the VLM$_{GABA}$ input-mediated effects on theta oscillation. To directly verify this, PVT$_{CR+}$ and

VLM$_{GABA}$ were transduced with AAVs carrying ChR2 in CR-Cre/vGAT-FLPO mice. After

recovery, antidromic activation of PVT PVT$_{\rightarrow NAc}$ and orthodromic activation of VLM$_{GABA}$

neurons were simultaneously applied while thalamic neuronal and cortical LFP activities were

monitored (Fig. 3a). Optogenetic activation of VLM$_{GABA}$ selectively altered the firing rate of

the photo-identified PVT$_{\rightarrow NAc}$ but not of the un-identified PVT neurons (PVT$_{UN}$; Fig. 3b).

Notably, VLM$_{GABA}$ activation increased PVT$_{\rightarrow NAc}$ firing to ~1 Hz. This elevation was further

enchanced when VLM$_{GABA}$ stimulation was paired with 1 Hz PVT$_{\rightarrow NAc}$ activation, and

increased even more when it was combined with 10 Hz thalamic stimulation (Fig. 3c and

Extended Data Fig. 7). In addition to changes in mean firing rate, PVT$_{\rightarrow NAc}$ neurons shifted

from phasic to tonic firing modes when 1 Hz thalamic stimulation was replaced by 10 Hz

stimulation together with VLM$_{GABA}$ activation (Fig. 3d), as reflected by the interspike interval

distribution (Fig. 3e).

Meanwhile, this progressively elevated PVT$_{\rightarrow NAc}$ activity bidirectionally modulated the theta

rhythm-promoting effects of VLM$_{GABA}$ input: while 1 Hz PVT$_{\rightarrow NAc}$ activation maintained

theta band power, 10 Hz thalamic stimulation suppressed it (Fig. 3f,g). These results confirm

that, despite the ongoing VLM$_{GABA}$ signaling, PVT$_{\rightarrow NAc}$ neurons can either enhance or

suppress REM-associated PFC theta oscillations in an activity- and firing mode-dependent

manner.

Stress-induced REM sleep modulation by PVT$_{\rightarrow NAc}$ neurons

Having identified PVT$_{\rightarrow NAc}$ neurons as a key mediator of VLM$_{GABA}$-to-PFC signaling

in physiological regulation of REM sleep and the associated cortical theta oscillation, we next

investigated whether this population also contributes to stress-induced modulation of REM

sleep. Stress exerts bidirectional effects on REM sleep: while acute stress transiently enhances

REM sleep as an adaptive response, chronic or intense stress suppresses it[40–43]. Thus, we

employed a 4-day restraint stress paradigm, allowing us to monitor the activity of the same,

optogenetically identified PVT cells during both acute and chronic stress as well as

characteristics of REM sleep.

Following a habituation day (R0), mice underwent 3 days of restraint stress (R1-3), with day 1

(R1) representing acute phase and days 2-3 (R2-3) modelling chronic phase of the restraint

stress (Fig. 4a). Throughout the experiment, frontal cortical EEG, EMG, and longitudinal

thalamic single unit recordings were conducted during habituation/restraint and the

subsequent 6-hour post restraint sleep periods (PRS0-3). Mice (n=6 mice) exhibited biphasic

REM sleep modulation due to the restraint stress: REM sleep and the associated theta band

power increased following acute phase, but were progressively suppressed upon repeated

restraint, chronic stress. By the third day, both of them were significantly reduced compared

to baseline (Fig. 4b and Extended Data Fig. 8a,b). Concurrently, $PVT_{\rightarrow NAc}$ neurons (N=11

cells; n=3 mice) showed increased firing during both the restraint and post-restraint sleep

periods, with some cells reaching mean firing rates of 10 Hz, which response was absent in

$PVT_{\rightarrow AMY}$ neurons (N=9 cells, n=3 mice; Fig. 4c,d and Extended Data Fig. 8c-e).

To establish a causal link between the progressive increase in $PVT_{\rightarrow NAc}$ activity and

bidirectional REM sleep modulation, we optogenetically activated $PVT_{\rightarrow NAc}$ neurons (10 Hz,

10 s) during sleep across multiple days while monitoring frontal cortical EEG activity (n = 7

mice; Fig. 4e). Control groups included mice expressing ChR2 in $PVT_{\rightarrow AMY}$ neurons (n = 8)

and YFP in PVT (n = 3).

Despite individual variability, $PVT_{\rightarrow NAc}$ stimulation produced bidirectional REM modulation,

mirroring the effects of restraint stress: stimulation on the first two days (S1-2) increased

REM sleep and theta band power, resembling the acute stress responses, whereas continued

stimulation (S3-6) progressively suppressed both, similar to chronic stress (Fig. 6f and

Extended Data Fig. 8h). PVT$_{\rightarrow NAc}$ firing also exhibited a progressive increase in non-

stimulated periods, paralleling the impact of restraint stress (Fig. 4h and Extended Data Fig.

8k). In contrast, stimulation of PVT$_{\rightarrow AMY}$ and YFP expressing mice showed no REM sleep

alterations (Fig. 4g and Extended Data Fig. 8i,l) meanwhile the activity of PVT$_{\rightarrow AMY}$ cells did

not increase but decreased (Fig. 4h). The comparable REM sleep dynamics induced by

restraint stress and PVT$_{\rightarrow NAc}$ activation (Fig. 6, b,f and Extended Data Fig. 8j) indicate that

progressive PVT$_{\rightarrow NAc}$ recruitment mediates stress-dependent REM sleep plasticity.

We also aimed to determine whether the PVT$_{\rightarrow NAc}$ is embedded into a brain-wide

network collecting the necessary stress and other theta-rhythm signals to shape sleep

architecture in a stress-dependent manner. Rabies-mediated monosynaptic retrograde tracing

revealed that PVT$_{\rightarrow NAc}$ neurons receive a broader array of inputs than PVT$_{\rightarrow AMY}$, including

projections from stress-sensitive and REM-regulating hypothalamic, mesencephalic and

brainstem nuclei (Extended Data Fig. 9a-h). These findings position PVT$_{\rightarrow NAc}$ as a central hub

for integrating stress and REM rhythm signals and modulating theta oscillation during sleep,

accordingly.

Activity-dependent effects of PVT$_{\rightarrow NAc}$ on PFC circuit

Next, we examined the impacts of the presently described medullo-thalamic

connection on cortical microcircuit dynamics responsible for REM-associated theta

oscillation[18,44]. During acute *in vivo* extracellular LFP recordings, we also monitored single-

unit activity of principal neurons (PN) as well as fast-spiking (FS) and non-fast-spiking (NFS)

interneurons in PFC, classified based on their electrophysiological properties (Extended Data

Fig. 10a-d). In parallel with PVT activity and cortical theta oscillation changes (Fig. 1),

prefrontal cortical interneurons (IN) and PN were also modulated by the inhibition of

VLM$_{GABA}$-to-PVT terminals. A larger proportion of IN and PN exhibited activity changes

when optogenetic silencing occurred during REM-like high theta states compared to all trials

Our further analysis revealed that FS neurons decreased while PN and NFS neurons increased

their activity upon inhibition of VLM$_{GABA}$-to-PVT inputs, coinciding with the reduction of

theta oscillation (Extended Data Fig. 10f-i). These findings suggest that the VLM$_{GABA}$ inputs

exert an influence on PVT, which in turn, shape the REM-associated theta rhythm via the PFC

circuit.

To find out whether selective activation of PVT$_{\rightarrow NAc}$ – which bidirectionally

modulates REM sleep and the associated theta oscillations (Fig. 2) – can drive similar changes

in prefrontal microcircuit[18,44], we further dissected the PVT-PFC network. First, we compared

the thalamic axonal distribution of PVT$_{\rightarrow NAc}$ and PVT$_{\rightarrow AMY}$ cells in the PFC. Analyzing the

retro-anterogradely labeled thalamic axonal patterns (see *Methods*), PVT$_{\rightarrow NAc}$ neurons were

found to preferentially target the superficial layers, while PVT$_{\rightarrow AMY}$ innervated the deep

layers (Fig. 5a,b and Extended Data Fig. 11). Next, we investigated the existence of laminar-

specific impact of PVT$_{\rightarrow NAc}$ and PVT$_{\rightarrow AMY}$ activation on cellular cortical activity. Antidromic

activation of PVT$_{\rightarrow NAc}$ and PVT$_{\rightarrow AMY}$ neurons was robust (Extended Data Fig. 12) and

induced distinct, layer-specific effects in prefrontal cortical single-unit activity, consistent

with their projection patterns. PVT$_{\rightarrow NAc}$ activation engaged a larger fraction of PFC neurons

and resulted in a better signal-to noise ratio in superficial layers compared to PVT$_{\rightarrow AMY}$ (Fig.

5c-h).

Notably, in anesthetized condition, 1 Hz and 10 Hz stimulation of PVT$_{\rightarrow NAc}$ (Extended Data

Fig. 13a,b) elicited opposing changes in the superficial PFC theta power (increased and

decreased; respectively) mirroring the effects observed in natural sleep conditions (Fig. 2).

Similar LFP dynamics were detected in the primary somatosensory cortex (Extended Data

Fig. 13c,d), indicating that PVT$_{\rightarrow NAc}$ — but not PVT$_{\rightarrow AMY}$ — can orchestrate global changes

in cortical theta oscillation.

When 1 Hz PVT$_{\rightarrow NAc}$ activation enhanced theta power in PFC, FS neuron activity increased,

while NFS and PN activity decreased. The opposite occurred with 10 Hz stimulation, which

suppressed theta power (Fig. 5i and Extended Data Fig. 13e). Similar shifts in PFC

microcircuit were observed during spontaneous sleep state transitions: from low-to-high theta

(as in NREM-to-REM) and high-theta-to-desynchronized states (as in REM-to wake

transitions; Fig. 5j and Extended Data Fig. 13f). Altogether, these findings indicate that

PVT$_{\rightarrow NAc}$, but not PVT$_{\rightarrow AMY}$ (Fig. 5k and Extended Data Fig. 13g) selectively shapes the

superficial PFC microcircuit dynamics underlying theta oscillation in an activity-dependent

and stress-dependent manner (Fig. 6).

Discussion

Establishing a neuronal network linking REM generator brainstem to PFC has broad

implications for understanding how REM sleep architecture adapts to homeostatic needs or

stress experiences. Our findings reveal a convergence of inputs that regulate REM sleep

rhythm, such as from the REM rhythm-generating VLM$_{GABA}$ neurons[14,15] (Figs. 1-3), and

from stress-signal carrying brain-wide pathways[45,46] onto the PVT$_{\rightarrow NAc}$ population (Extended

Data Fig. 9). In an activity- and firing mode-dependent manner, PVT$_{\rightarrow NAc}$ maintains or

suppresses the effect of VLM$_{GABA}$ signaling and drives shifts in the excitatory/inhibitory (E/I)

balance within the PFC, which, in turn, are manifested as bidirectional changes in REM-

associated theta oscillation (Figs. 3,5 and 6). We further demonstrate that PVT$_{\rightarrow NAc}$ activity

increases progressively with prolonged stress exposure, mediating dual roles of stress in

regulating REM sleep: acute stress results in a REM rebound, whereas repeated stress elevates

PVT$_{\rightarrow NAc}$ activity and and disrupts REM sleep (Fig. 4).

VLM_{GABA} inputs to the $PVT_{\rightarrow NAc}$ were shown to play an important role in fine-tuning of

REM sleep-associated theta oscillation. Short (5 ms) activation of VLM_{GABA} neurons exerts

an inhibitory effect on individual $PVT_{\rightarrow NAc}$ cell by eliciting IPSCs (Fig. 2). However, longer

(10 s) medullary inhibition caused facilitation at population level (Fig. 3). Prolonged

inhibition can gradually raise intracellular chloride levels shifting the chloride equilibrium

potential toward more depolarized values and thereby biasing the membrane potential closer

to the thalamic firing threshold. Under such conditions, the same inhibitory input can become

functionally excitatory in hippocampal and hypothalamic cells[47–49], a scenario that may also

be present in $PVT_{\rightarrow NAc}$ cells explaining their increased firing upon longer medullary

inhibition (Fig. 2). This moderate enhancement in $PVT_{\rightarrow NAc}$ activity biases the system toward

REM sleep transition.

In addition to the VLM_{GABA} input, the $PVT_{\rightarrow NAc}$ integrates a range of subcortical inputs

essential for fine-tuning REM rhythm [such as the locus coeruleus[34,50], lateral hypothalamus[51],

tegmental/pontine area[52] and preoptic area[53]], regulating homeostatic needs[45] as well as

providing adequate behavioral responses to stress[30–33] (Extended Data Fig. 9). The

convergence of all of these inputs can further modulate $PVT_{\rightarrow NAc}$ firing mode and activity,

which, in turn, re-evaluates REM rhythm transition to the PFC and modulates REM sleep

theta oscillation, accordingly (Fig. 3).

The emergence of REM sleep-coupled theta oscillation in the cortex highly depends on

the cortical E/I balance[18,37,44]. Our data highlight that $PVT_{\rightarrow NAc}$-PFC input acts as a critical

modulator capable of altering PFC inhibitory circuits to adaptively promote or suppress

cortical theta oscillation, in an activity-dependent manner (Fig 5). During acute (mild) stress

or when homeostatic REM sleep pressure increases, $PVT_{\rightarrow NAc}$ neurons likely receive both

moderately enhanced stress- or homeostasis-related excitatory input and prolonged VLM-

mediated inhibition. This combination increases $PVT_{\rightarrow NAc}$ activity engaging fast-spiking

(FS) interneuron–mediated feedforward inhibition and shifting the cortical E/I balance toward

REM sleep promotion[18,54,55]. Conversely, stronger excitatory inputs due to chronic stress can

cause over-activation and tonic firing of PVT$_{\rightarrow NAc}$ neurons, shifting the E/I balance toward

non-FS-based inhibition and REM sleep suppression. Consistently, we showed that acute 1

Hz and 10 Hz optogenetic stimulation of PVT$_{\rightarrow NAc}$ recapitulated homoeostatic changes in

theta oscillation, whereas chronic stimulations mimicked stress-induced alterations of REM

sleep. Repeated 10 Hz activation of PVT$_{\rightarrow NAc}$ neurons for 1-2 days resulted in facilitation of

REM sleep theta oscillation similar to acute stress, while prolonged stimulation (up to 6 days)

strongly reduced it similar to chronic stress. These findings provide causal evidence that

establishes PVT$_{\rightarrow NAc}$ activity as an important mechanism in fine-tuning of REM sleep and its

associated theta oscillation in accordance with physiological and behavioral states.

Notably, stressed-triggered thalamic hyperactivity is persistent, causing REM sleep

facilitation or reduction for hours or even days, in contrast to the transient thalamic activity

increase observed at natural NREM-to-REM transitions. Beyond fast glutamatergic

transmissions, these long-lasting changes in sleep likely depend on slower neuromodulatory

signaling mechanisms. One candidate is nitric oxide (NO), whose levels has been shown to be

suppressed by acute stress while elevated by repeated stress[56]. As cortical NO level can

influence E/I balance[57] and that the PVT$_{\rightarrow NAc}$ cells express NO synthetizing enzymes

(Extended Data Fig. 14), thalamic NO release may contribute to activity-dependent shifts in

PFC inhibitory circuits underlying REM sleep modulation. Similarly, acute stress has been

shown to elevate corticotrophin-releasing hormone (CRH) contributing to REM sleep

rebound[58–61]. Given that PVT$_{\rightarrow NAc}$ cells also express CRH[62], their stress-induced activation

and CRH release may also influence cortical activity over longer timescales. Together, NO

and CRH signaling in PVT➔NAc neurons may mediate the long-term impact of stress on

REM sleep regulation.

Encoding stress signals and shaping adaptive stress responses are widely distributed

across forebrain circuits, including the NAc and PFC[63–67]. Recently, PVT neurons,

particularly those with accumbal or cortical collaterals, have been shown to modulate stress-

induced and motivational behaviors[27,31,68–70]. By identifying the sleep rhythm regulatory role

of PVT$_{\rightarrow NAc}$ cells, which co-project to both forebrain regions, this thalamic cell group may

serve as a shared mechanism linking REM sleep expression with stress-coping forebrain

networks, thereby coordinating REM sleep regulation with its associated functions.

The laminar-specific cortical connectivity of PVT$_{\rightarrow NAc}$ neurons (Fig. 5) is also highly

dedicated for REM sleep theta modulation, as these preferentially target the superficial layers.

These cortical layers possess a unique role in REM sleep, supported by strong feed-forward

and feed-back inhibitory mechanisms[37,71–74]. Cortical FS neurons form dense perisomatic-

targeting and axo-axonic connections with neighbouring pyramidal cells, mediating fast local

feed-forward inhibition, that is particularly critical for REM initiation and maintenance within

superficial layers[18,36,37,44,75,76]. In contrast, NFS neurons provide a feed-back inhibition,

reducing REM-associated theta activity and promoting slow wave sleep[77]. In addition, layer

2/3, but not layer 5, pyramidal neurons in the retrosplenial cortex are strongly activated during

REM stage, indicating that superficial principal cells are indispensable for REM sleep

regulations[37]. As shifts in cortical inhibition underlying REM sleep kinetics are primarily

affected by PVT$_{\rightarrow NAc}$ neurons, distinguishing them from neighbouring PVT$_{\rightarrow AMY}$ cells,

PVT$_{\rightarrow NAc}$ neurons are ideal to fulfil the role of 'conductor' in state-dependent regulation of

REM-related cortical dynamics.

Taken together, their activity dynamics and input-output organization uniquely position

PVT$_{\rightarrow NAc}$ neurons to modulate cortical REM sleep in a homeostatic and stress-dependent

manner, exerting opposite effects on theta oscillations via superficial cortical inhibitory

circuits. As homeostatic E/I balance-mediated REM sleep is essential for PFC-dependent

functions[78–80], stress-driven neuronal reorganization – modulated by PVT$_{\rightarrow \text{NAc}}$ activity – may

lead to persistent changes within frontal cortical structure and in REM-associated cognitive

performance[3,72,81], and ultimately, contribute to the development of psychiatric disorders[82–85].

Methods

Animals

Adult (2–5 months old female and male) CR (*Calb2*)-Cre (RRID: IMSR_JAX:010774), vGAT (*Slc32a1*)-Cre and -FLPO (from The Jackson Laboratory) mice were used. We also crossbred mice to have CR-Cre/vGAT-Flpo mice for the experiments. Female mice were used only for the anatomical investigations. Animals from the same gender were group-housed in a humidity- and temperature-controlled environment. Animals were kept under standard laboratory conditions on a 12 h light/12 h dark cycle with food and water available ad libitum. The Regional and Institutional Committee of the Research Centre for Natural Sciences, the Institute of Experimental Medicine and the National Animal Research Authorities of Hungary (PE/EA/00080-4/2023) approved all procedures. After all surgeries, animals received subcutaneous Rimadyl (Carprofen, 1.4 mg/kg).

Stereotactic surgeries

Viral injection

Mice were anesthetized with ketamine–xylazine (5:1, 3× dilution, ketamine: 100 mg/kg; xylazine: 4 mg/kg) and injected with one of the following AAV vectors with Nanoliter Injector (WPI) at a rate of 1 nl s^{-1} (50-200 nL): AAV-Ef1a-DIO-ChR2-eYFP (Penn Vector Core; titer: 5×10^{12} to 1×10^{13} genome copies (GC)/ml), AAV5-EF1a-DIO-eYFP-WPRE-hGH (Penn Vector Core; titer: 5×1012 GC/ml), AAV-Ef1a-DIO-ChR2-mCherry (Penn Vector Core; titer: 5×10^{12} to 1×10^{13} GC/ml), AAV-DFO-ChR2-eYFP virus (titer: 1×10^{13} GC/ml), AAV5-EF1a-DIO-mCherry (UNC Vector Core; titer: 5.3×10^{12} GC/ml), AAV-Ef1a-DIO-eNpHR3.0-eYFP (Penn Vector Core; titer: 3×10^{13} GC/ml), AAV-CAG-FLEXFRT-ChR2-mCherry (Addgene; titer: $\geq 7 \times 10^{12}$ GC/ml) and AAVrg-CAG-GFP (Addgene; titer: 7×10^{12} GC/ml). The target regions are (with anterior–posterior, lateral and dorsal–ventral

coordinates): dorsal midline thalamus (paraventricular thalamus, PVT; −1-1.2/±0/3.2–2.8 mm

(n=35 animals); amygdala (AMY; −1.8/±3.4/-3.7 mm) (n=18 animals); nucleus accumbens

(NAc; +1.4/±1.1/–3.9–4.2 mm) (n=16 animals) and ventrolateral medulla (VLM; −7.3/±1.4/–

5.3 mm) (n= 27 animals).

Rabies tracing

The tracing method has been used and validated earlier[86]. CR-Cre mice were injected with

100 nL of AAV2/8-hSynFLEX-TVA-p2A-eGFP-p2A-oG (4.5×10^{12} GC/mL) into the PVT.

After 4 weeks, mice were injected with rabies(ΔG)-EnvA-mCherry (100 nL; 3.5×10^{7} GC/mL)

into the NAc or AMY (N = 3-3). Mice were perfused after 9-10 days of survival. Both the

rabies (mCherry) and the AAV (eGFP) signal were amplified by immunofluorescent staining

procedure (see below).

Immunohistochemistry

All anatomical data were obtained from immunohistochemically stained mouse brain slices.

Firstly, the tissue blocks were cut with a VT1200S Vibratome (Leica) into 50 μm coronal

sections. Free-floating brain sections were intensively washed (5 times for 10 min) with 0.1 M

PB. During immunohistochemical processing, all antibodies were diluted in 0.1 M PB. For

high-quality fluorescent labeling, sections were treated with a blocking solution containing

10% normal donkey serum (NDS, Sigma-Aldrich, S30-M) or normal goat serum (NGS,

Vector, S-1000, RRID: AB_2336615) and 0.5% Triton-X (Sigma-Aldrich) for 30 min at room

temperature (RT). In the case of electrode track analysis, sections were cryoprotected in 30%

sucrose solution overnight, and freeze thawed over liquid nitrogen and then, treated with 10%

normal donkey serum solution to keep the visibility of the DiI (1,1′-Dioctadecyl-3,3,3′,3′-

tetramethylindocarbocyanine perchlorate; Santa Cruz Biotechnology).

Then, the sections were incubated in primary antibody solution at RT overnight or for 2–3

days (at 4 °C). The following primary antibodies were used: green fluorescent protein (GFP,

chicken, Life Technology: A10262, RRID: AB_2534023; 1:2000), mCherry (rabbit,

BioVision: 5993-100, RRID: AB_1975001; 1:2000), FoxP2 (mouse, Merck Millipore:

MABE415, RRID: AB_2721039; 1:500; Invitrogen: MA5-31419, RRID: AB_2787055;

1:2000; rabbit, Abcam: ab16046, RRID: AB_2107107; 1:500), Calb1(rabbit, SWANT: CB38,

RRID: AB_10000340; 1:2000; mouse, SWANT: 300, RRID: AB_10000347; 1:2000), PV

(mouse, SWANT: PV235, RRID: AB_10000343; 1:2000), CR (mouse, SWANT: 6B3, RRID:

AB_10000320; 1:2000), Ctip2 (rat, Abcam: ab18465, RRID: AB_2064130; 1:500), WFS1

(rabbit, Proteintech: 26995-1-AP; RRID: AB_2880717; 1:500) and NOS (mouse, Sigma:

N2280; RRID: AB_260754; 1:300). After the primary antibody incubation, the sections were

treated with the following secondary IgG antibodies for fluorescent staining (1:500; 2 h at

RT): Alexa 488-conjugated goat anti-chicken (GACh-A488; Molecular Probes: A11039,

RRID: AB_142924), Alexa 488-conjugated donkey anti-mouse (DAM-A488; Jackson: 715-

545-150, RRID: AB_2340846), Alexa 555-conjugated donkey anti-rabbit (DAR-A555;

Invitrogen: A31572, RRID: AB_162543), Cy3-conjugated donkey anti-mouse (DAM-CY3;

Jackson: 715-165-151, RRID: AB_2340813), Cy3-conjugated donkey anti-rabbit (DAR-CY3;

Jackson: 711-165-152, RRID:AB_2307443), Alexa 594-conjugated donkey anti-rabbit (DAR-

A594; Molecular Probes: A21207, RRID: AB_141637), Alexa 647-conjugated donkey anti-

rabbit (DAR-A647; Jackson: 711-605-152, RRID: AB_2492288). In some cases, staining was

enhanced after primary antibody incubation with biotinylated secondary antibodies (1:300; 2

h at RT), which are the followings: biotinylated goat anti-rabbit IgG (bHAR; Vector

Laboratories: BA-1100, RRID: AB_2336201), biotinylated goat anti- mouse IgG (bHAM;

Vector Laboratories: BA-2000, RRID: AB_2313581), Elite Avidin-Biotin Complex (eABC;

Vector Laboratories: PK-6100, RRID: AB_2336819), streptavidin-conjugated green

fluorescent antibodies (1:1000-2000; 2 hr. at RT) (SA-A488; Jackson: 016-540-084, RRID:

AB_2337249), streptavidin-conjugated red fluorescent antibodies (SA-CY3; Jackson: 016-

160-084, RRID: AB_2337244), streptavidin-conjugated far red fluorescent antibodies (SA-

A647; Jackson: 016-600-084, RRID: AB_2341101). All the fluorescent slices were mounted

in Vectashield Mounting Medium (Vector Laboratories: H-1000, RRID: AB_2336789) or in

Vectashield Mounting Medium with DAPI (Vector Laboratories: H-1200, RRID:

AB_2336790).

Alternatively, we used DAB-Ni as a chromogen for Rabies. Sections were cryoprotected in

30% sucrose solution overnight, and freeze thawed over liquid nitrogen and then were treated

with 1% H_2O_2 solution (10 min) for endogenous peroxide blocking. Next, the slices were

incubated in 10% normal donkey serum (30 min, RT). After primary antibody incubation,

slices were treated with biotinylated secondary antibodies (goat anti-rabbit or horse anti-goat

IgG) and eABC (see above). To visualize the labeled cells, sections were developed with

DAB-Ni. Sections were then dehydrated in xylol and mounted in DePex (Serva).

Identification of cortical layers

To differentiate the cortical layers in PrL and the primary somatosensory cortex (S1), we used

combination of neurochemical markers[87]. The cortical layer 2/3 (L2/3) was identified by using

Calb1 staining, while layer 6 (L6) was visualized with fork head box protein P2 (FoxP2)

staining. Furthermore, for dissemination of the layer 5 were used Chicken ovalbumin

upstream promoter transcription factor-interacting proteins 2 (Ctip2) and anti-Wolfram

syndrome 1 (WFS1). The dorsoventral borders of the PrL cortex is determined by the

parvalbumin (PV) immunostaining[88].

Axon density analysis

We performed retro-anterograde viral labeling approach in CR-Cre mice to visualize PFC

projections of PVT$_{\rightarrow NAc}$ and PVT$_{\rightarrow AMY}$ neurons as described earlier[25]. Shortly, AAV injection

into NAc or AMY retrogadely-labeled CR+ neurons in PVT. If these PVT cells had cortical

collaterals, the virus propagated in an anterograde fashion and visualized axons in PFC as

well. These retro-anterograde viral labelings were also amplified and the slices were

counterstained for the above mentioned markers to identify the PrL cortex layers and borders.

Confocal Z-stack (slice number: 50; total thickness: 23.34-40.3 μm, PVT$_{\rightarrow NAc}$: 36.31±2.66

μm; PVT$_{\rightarrow AMY}$: 31.83±4.32 μm) images were taken with confocal microscope (Zeiss LSM

710; Zeiss ZEN 2010B SP1 Release version 6.0; Carl Zeiss Microimaging GmbH) a ×63

objective (10+x Plan Apochromat 10x/0.45 M27; 20x Plan Apochromat 20x/0.8 M27; 63x

Plan Apochromat 63x/1.4 Oil DIC M27) at four different AP levels between Bregma +2.22

and +1.54 mm. The length of the retro-anterograde labelled CR+ axons were counted with a

use of a custom-written Fiji macro[25] in a given region of interest (available at

https://github.com/baabek/Axon-density-analyzer-ImageJ-script.git). The macro calculated

axon length for each stack and the total axon length was summed up for each layers and slices

in each animal. Then, the relative axon density was counted for each layers and slices (total

axon length / slice volume (ROI area * interval)), and finally the relative axon density was

calculated for the animal (RAD = total axon length/total volume). Furthermore, we

determined the total axon distribution by layers for each animal (layer RAD/all RAD).

To analyze the distribution of VLM$_{GABA}$ axons in PVT, we used the same methods. The area

of the PVT was determined by calretinin (CR) immunostaining[88]. The distribution of the

anterogradely labelled VLM$_{GABA}$ axons were counted between Bregma -0.3 and -2.1 mm in

PVT.

Chronic optrode implantations

Four weeks after AAV-Ef1a-DIO-ChR2-eYFP and AAV5-EF1a-DIO-eYFP virus injections

into AMY or NAC in CR-Cre mice, four custom-fabricated tungsten tetrodes (12.5 μm in

diameter, California Fine Wire) were implanted into the PVT (AP/ML/DV (−1/±0.6/–2.8 mm

at 10°). The tungsten electrodes were attached with a multimode optic fiber (105-μm core

diameter, NA = 0.22; Thorlabs) (200–300 μm left between the optic fiber and tetrode tips),

with the help of a polyimide tube (0.008 ID, Neuralynx). The electrode wires were joined to

an electrode interface board (EIB-16, Neuralynx) by gold electrode contact pins (Neuralynx).

The ground, reference, EMG, EEG wires (Phoenix Wire Inc.) were soldered to the appropriate

electrode interface board point. The tetrode impedances measured before the implantation at 1

kHz were kept between 240 and 750 kΩ (Intan Technologies). Ground and reference screws

were implanted in parietal bone, respectively; an EMG wire was inserted into the neck

muscle; an EEG screw was placed in frontal bone over PFC cortex. Finally, all pieces were

secured onto the skull by multiple layers of dental acrylic (Paladur, Heraeus Kulzer) and

shielded by a copper web. Mice were left for at least 4 days to recover and then, handled for

several days.

In vivo electrophysiology in anesthetized preparations

In vivo recordings were performed 4–12 weeks after the viral injections (AAV-Ef1a-DIO-

ChR2-eYFP or AAV-CAG-FLEXFRT-ChR2-mCherry into the PVT or AAV-Ef1a-DIO-

eNpHR3.0-eYFP into the VLM) from the prelimbic cortex, primer somatosensory cortex and

the paraventricular thalamus in urethane anesthetized (20 m/m % dilution; 0.005ml/1g) CR-

Cre or vGAT-Cre transgenic mice. The head of the animal was fixed in a stereotaxic frame; a

screw was driven into the occipital or into the left parietal bone, which served as a reference

electrode. LFP activity as well as the extracellular unit activities from PrL, PVT and S1 were

monitored with silicon probes (A1x16-5mm-50-703, L16; A1x32-10mm-50-177, L32;

A2x16-10mm-100-500-177, BL32; A1x32-5mm-50-177, L32; Neuronexus), stained with

fluorescent DiI. The wideband neural data (0.1–7500 Hz) were amplified (gain: 192×) and

digitalized at 20 kHz (Intan Technologies). The electrodes were lowered into PrL

(AP/ML/DV +2/–2.5/–3.5 mm/60°), into S1 cortex (AP/ML/DV –1.2/+3.2/–1, –0.8 mm/20°)

and into PVT (AP/ML/DV –1.1/+1/–3.2 mm/20°).

For the 'antidromic–orthodromic' activation[25], the optic fibers were placed into the NAc and

AMY, where the PVT_{CR+} fibers were activated, and the evoked LFP and unit responses were

detected in PrL, S1 and PVT. Under these conditions, action potentials first traveled

antidromically and, at a putative branching, could turn to orthodromic direction as well. The

NAc and AMY did not contain CR+ cells, which projected back to the PVT, so the thalamic

stimulation was due to direct antidromic activation. In the cases of PVT somatic and

VLM_{GABA}→PVT terminal manipulations, the optic fiber was placed above the PVT. For the

VLM_{GABA} somatic stimulation, the optic fiber was placed above the VLM (AP/ML/DV –

7.2/+3.3/–5.6/20°). Optogenetic activation consisted of 473-nm blue-light pulses (1s or 10s, 1

or 10 Hz, ~10-15 mW, Laserglow Technologies). In the case of the 1 s long activations, the

stimulus was repeated 30 times and the interstimulus intervals were 60 s, while the 10 s long

stimulations were repeated 10 times with 250 or 300 s interstimulus intervals. The optogenetic

inhibition with NphR3.0 consisted of 532-nm green-light pulses (10s, ~15 mW). The lasers

were triggered by analog signals driven by a National Instruments acquisition board (USB-

6353) and Matlab. Analog trigger pulses were also delivered to the recording system and

registered parallel with neural data with zero latency. After recordings, animals were

transcardially perfused, and coronal sections were cut from the paraformaldehyde-fixed

brains. The position of the silicon probes were verified based on DiI labeling of the electrode

track. For determination of the layer specific mapping, we used different neurochemical

markers (see above). Mice with incorrect virus injections or electrode positions were excluded

from the analysis (N=4 mice).

In vivo electrophysiology of freely behaving mice

In vivo recordings from freely behaving CR-Cre mice were performed 4-5 days after the optrode implantation (into the PVT). EEG, EMG and extracellular PVT unit activities were monitored with optrodes and wire implantation (see previously). Of note, chronic EEG recording with frontal cortical screw electrodes and acute LFP recordings with multisite silicon probes resulted in comparable sleep oscillation fluctuations, similar duration of single theta band (~REM) episodes as well as similar theta band power values (Fig. S1). These findings indicate that both approaches are equally suitable for investigating the neuronal processes underlying REM-associated theta oscillations.

The wideband neural data (0.1–7500 Hz) were amplified (gain: 192×) and digitized at 20 kHz. Optogenetic tagging and activation consisted of 473-nm blue-light pulses (5 ms 0.5–1 Hz, for tagging; 10s 1 Hz; 10s 10 Hz; ~15 mW). The evoked unit responses were detected in PVT_{CR+}, $PVT_{→NAc}$ or $PVT_{→AMY}$ neurons simultaneously with the EEG and EMG signal. The $PVT_{→NAc}$ or $PVT_{→AMY}$ neurons were transduced with ChR2 carrying AAV vectors using retro-anterograde viral injections. For optical identification, 5 ms long pulses were repeated 30 times, while the 10 s long 1 Hz and 10 Hz stimulations were repeated several times with random interstimulus intervals, which was minimum 300 s long. The behavior of mice was captured on video at 30 frames/s.

We measured the spontaneous sleep-wake transitions and sleep architecture of the $PVT_{→NAc}$ (N= 6 mice) and the $PVT_{→AMY}$ mice (N= 7 mice) for several days. Each day 10s 1 Hz and 10s 10 Hz stimulation were randomly applied.

Restraint stress

The spontaneous sleep-wake transitions and sleep architecture were measured in both light and dark phases (EEG, supplemented with EMG signal) and the activity of the $PVT_{→NAc}$ and

PVT→AMY (N= 6 mice) cells. During the four-day experiment 19 units of PVT→NAc and 20

units of PVT→AMY neurons were followed. Only cells that could be followed through the

entire stress paradigm were included in the further analysis (PVT→NAc, N = 11 and PVT→AMY,

N = 9). Daily measurements were merged, re-clustered, allowing reliable identification of

corresponding data points. The first day served as a pre-restraint day (light-dark phase)

followed by three consecutive days of restraint stress. Mice were exposed to 90 min of

restraint stress in a falcon tube at the end of the light phase. Due to the tetrode implantation, a

longitudinal gap was made on the top of the falcon tube. Immediately after the restraint stress,

sleep architecture was measured during the first 6 h in the dark phase.

At the end of the experiments, animals were transcardially perfused, and coronal sections

were cut from the paraformaldehyde-fixed brains. The position of the optrodes and EEG

screw were verified with help of the electrode track.

Acute slice electrophysiology

For acute slice preparation, mice (vGAT-Cre, 3-month-old, N_{AMY}=5, N_{NAc}=4) were

decapitated under deep isoflurane anesthesia. The brain was removed and placed into an ice-

cold cutting solution, which had been bubbled with 95% O_2–5% CO_2 (carbogen gas) for at

least 30 min before use. The cutting solution contained the following (in mM): *205 sucrose,*

2.5 KCl, 26 $NaHCO_3$, 0.5 $CaCl_2$, 5 $MgCl_2$, 1.25 NaH_2PO_4, 10 glucose, saturated with 95%

O_2–5% CO_2. Coronal slices of 300 μm thickness were cut using a Vibratome (Leica

VT1000S) and placed into an interface-type incubation chamber for recovery submerged in

standard artificial cerebrospinal fluid (ACSF) containing the following (in mM): *126 NaCl,*

2.5 KCl, 26 $NaHCO_3$, 2 $CaCl_2$, 2 $MgCl_2$, 1.25 NaH_2PO_4, 10 glucose, saturated with 95% O_2–

5% CO_2. During measurements, slices were transferred individually into a submerged-type

recording chamber with a superfusion system allowing constantly bubbled (95% O_2–5% CO_2)

ACSF to flow at a rate of 3-3.5 ml/min. The ACSF was adjusted to 300-305 mOsm. All

measurements were carried out at 33 –34°C, temperature was maintained by a dual flow

heater (Supertech Instruments). The pipette solution contained (in mM): 110 D-gluconic acid

potassium salt, 4 NaCl, 20 HEPES, 0,1 EGTA, 10 phosphocreatine di(tris) salt, 2 ATP

magnesium salt, 0.3 GTP sodium salt, pH: 7.3, 280-300 mOsm and biocytin (Sigma). Pipette

resistances were 3-6 MΩ when filled with pipette solution. Visualization of slices and

selection of retrogradely labelled AMY or NAc projecting PVT cells was guided by AAVrg-

CAG-GFP signal (see: animals/Viral injection) and was imaged under an upright microscope

(BX61WI; Olympus, Tokyo, Japan equipped with infrared-differential interference contrast

optics and a UV lamp). Only cells located deeper than ~50 µm (measured from the slice

surface) were targeted. All cells were recorded in voltage-clamp mode and held at -65 mV

holding potential during the formation of the gigaseal. Series resistance was constantly

monitored after the whole-cell configuration was established, and individual recordings taken

for analysis showed stability in series resistance between a 5% margin during the whole

recording. After the whole-cell configuration was established, cells were clamped at -10 mV

holding potential in order to record inhibitory postsynaptic currents (IPSCs). In order to evoke

IPSCs, trains of 5*5ms laser light pulses (473 nm, calibrated to ~10 mW at slice surface,

Sanctity Laser) were delivered at 1, 5, 10, and 20 Hz frequency by an optic fiber positioned

above the slices. Recordings were performed with a Multiclamp 700B amplifier (Molecular

Devices). Data was digitized at 20 kHz with a DAQ board (National Instruments, USB-6353).

All data were processed and analyzed off-line using standard built-in functions of Python

2.7.0. environment. A threshold for IPSC peak detection was calculated for each neuron from

a 2-minute-long baseline before stimulation trains. The threshold value was set to be $3*SD_{mean}$

of the signal. IPSPs were considered to be evoked (eIPSCs) if the peak was detected within

the 2-10 ms range after a given stimulus on-set. Spontaneous IPSCs were detected for

quantification from the 2-minute-long baseline region before stimulation trains.

Neural data processing

LFP analyses

The natural sleep and anesthetized measurements were analyzed in a similar way. The

LFP/EEG/EMG signals were downsampled at 3000 Hz. The power of delta (1–4 Hz), theta

(4-8 Hz), sigma/spindle activity (9–14 Hz), alpha (8–13 Hz), beta (13-30 Hz) frequency bands

were differentiated[89]. For the further analysis the signals were filtered on a given frequency,

thereafter were the absolute value of the filtered signals calculated, on which we performed

smoothing in 100 ms window (RMS; root-means square) (**Fig. 2o,** Extended Data Fig. 1a,3f).

Sleep–wake states were determined by using RMS. In the natural sleep conditions, awake

state was characterized by high alpha, beta activity (alpha, beta RMS) and low delta power

(delta RMS) with high EMG (threshold >0.8), while sleep was characterized by low alpha and

high delta activity with low EMG, and it was further subdivided into NREM and REM state.

For the further classification the EMG signal was normalized. REM was determined as a high

sigma/delta (sigma/delta RMS) ratio associated, with low delta and high theta power and low

EMG (threshold <0.5). NREM was determined as a low sigma/delta ratio associated with high

delta power or spindle activity and low EMG (threshold <0.8). NREM sleep was not divided

into further subsections. In the anaesthetized conditions, desynchronized state was

characterized by high alpha, beta activity (alpha, beta RMS) and low delta power (delta

RMS), while sleep was characterized by low alpha and high delta activity, and it was further

subdivided into NREM-like and REM-like state. REM-like states were determined as a high

sigma/delta (sigma/delta RMS) ratio associated with low delta and high theta power. NREM

was determined as a low sigma/delta ratio associated with high delta power or spindle

activity. All animals and measurements had a different sigma/delta RMS ratio, thus, the ratio

was calculated for each trial and for each animal; the pre-stimulus ratio was compared to the

measurement specific median. An 18 s time window, the baseline (sleep) period (between -20

s and -2 s), was used to determine pre-stimulus sleep stages. During spontaneous sleep

architecture monitoring, the oscillation fluctuation was monitored in 5 s time windows.

The delta and theta RMS changes were calculated for the normalized signal after the optical

stimulation or for the spontaneous state. The starting point of the elevated delta and theta

RMS segment was determined by the time point, when the normalized signal exceeded the

baseline for the first time after the stimulus, while the end point of the changes was the return

of the RMS signal to baseline. The lengths of the delta and theta RMS changes (recovery-

time) were given by the time period between these two points. The REM onset was given by

the time when all the criteria of REM state appeared at the same time. 10s 1/10 Hz 10 mW

optogenetic activations were applied to analyze changes in LFP activity (delta and theta RMS,

FFT, modulation index (MI)). When the sleep was 'light' (the EEG was desynchronized) for

longer periods, the stimulation protocol was paused. Sham stimuli were used as the control.

The starting point of a sham stimulus was the half point of the ISI. Sleep–wake states were

determined in the case of the sham stimuli, too.

Under natural sleep conditions, the FFT and RMS changes were calculated for the frontal

EEG signal. In anesthetized measurements, we could distinguish superficial and deep layers

due to the insertion path of the extracellular electrode. Based on the distribution of the

$PVT_{\rightarrow NAc}$ and $PVT_{\rightarrow AMY}$ fibers, as well as the location of the intertelenchephalic and

corticothalamic/pyramidal track neurons in the PFC[87], layers 1, 2-3 and 5a were taken as the

superficial, while layers 5b and 6 were considered as the deep territories. From each animal,

one channel per superficial vs. deep layer was further analyzed. The channels were selected

based on the best signal-to-noise ratio and were located at least 300 μm away from each other.

For identifying the state-dependent firing rate for PVT cells, EMG onset as an indicator for

sleep/wake transition was set as described above. Awake periods were only accepted when

these were preceded by a 30 s sleeping phase and were longer than 500 ms, defined as the

lower limit for minimal arousal[25]. Peri-event time histograms (PSTH) were defined for each

cell around the detected EMG onsets. Z-score values of the firing rates were given upon

PSTH calculation for each cell to a 20 s baseline (sleep) period (between –30 s and –10 s,

calculated from the onset of the EMG signal). Significant changes of the firing rates were

defined upon at least two significant ($z > 1.92$, $P > 0.05$) neighboring z-score (1 s) bins in the

[–10, 10]-s interval around EMG onset. Mean z-scores for $PVT_{\rightarrow NAc}$ or $PVT_{\rightarrow AMY}$ neurons are

presented[25]. To determine the sleep state specific EMG changes, distinct calculation methods

were used. Each animal and measurement had different EMG levels, thus for the EMG onset

latency and duration were calculated for every trial, and for each animal; the pre-stimulus

ratio was compared to the measurement specific median. If the median pre-stimulus EMG

level was lower than the post-stimulus EMG level for each stimulus up to 18 s baseline

(sleep) period (between –20 s and –2 s), then it was determined as an EMG onset.

To validate the probability of stimulation-induced state transition, we analyzed the RMS

changes. The probability was assessed when the animal transitioned from one state (X) to

other (Y) within a fixed time window (t = 10 s). First, all stimulation events (p) were

identified during which the animal was in state X, then the number of events (q) were

determined in which the animal entered state Y. State Y was considered valid if it was

maintained for the entire time window or exceeded it (d). The cumulative probability of

transitions were calculated according to the following equation ($P(X \rightarrow Y \,|\, t \leq d) = q \,/\, p$), in

order to quantify the likelihood of stimulation-induced state transition (**Fig. 2o**). To validate

the efficiency of stimulation-induced theta modulation, an efficiency index was calculated.

This index was defined as the product of the probability of transition (see above) and the

duration of achived state (in secundum): efficiency index = transition probability * state

duration (**Fig. 2o**).

Single cell analyses

The natural sleep and anesthetized measurements were analyzed in a similar way. Noise

filtering by average subtraction and raw electrophysiological recordings were filtered (>500

Hz) for spike detection. Spike detection and principal-component-analysis-based automatic

clustering were performed using SpikeDetekt and KlustaKwik (2019 edition) respectively[90].

Cell grouping was manually refined using KlustaViewa (2019 edition). A group of spikes was

considered to be generated by a single neuron if they formed a discrete, well-isolated cluster

and had an autocorrelogram with a refractory period (if average bin values of the first 1.5 ms

did not reach the autocorrelogram's asymptote line). Cells were merged in those cases where

units had the same autocorrelogram and a similarly shaped action potential on the same

channel points to avoid enumerating the same cell more than once. Further data analyses were

carried out using custom-built Matlab routines. All the presented electrophysiological data

was derived from individually identified single units (clustered cells).

Putative principal neurons (glutamatergic cortical neurons) and GABAergic cortical

interneurons were separated by the spike half-width, the autocorrelogram[91] and the baseline

firing rate [92], which we considered as the most reliable parameters for this purpose. Based on

previous data on cortical GABAergic interneurons, we distinguished fast spiking (FS) and

non-fast spiking (NFS) interneurons based on their spike half-width, baseline firing rate and

autocorrelogram (Extended Data Fig. 10a-d).

We calculated the signal-to-noise ratio to determine the strength of the evoked feed-back

inhibition on the cortical signal (**Fig. 5e,h**). To define it, we calculated the variance of the

evoked response between two time points: T_0 and $T_{evoked\ MUA\ peak}$. The $T_{evoked\ MUA\ peak}$ is the

positive peak of the MUA for the 1s 10 Hz optical stimulation. The signal-to-noise ratio was

then estimated as the mean at T_0 and $T_{evoked\ MUA\ peak}$ divided by standard deviation at T_0 and

$T_{evoked\ MUA\ peak}$. The evoked signal-to-noise ratio was calculated for the first three evoked

MUA peaks, which were compared to the signal-to-noise ratio of the stage before the stimulus

(-100 ms-0 ms).

Significant changes in evoked responses of PVT cells were validated with MI. MI was

calculated for the firing rate (FR) from the mean PSTH of the pre period (between -10 s and 0

s) which was compared to the mean PSTH of the post period (between 0 s and +10 s). We

expressed the modulation index as the ratio of firing rate: $(FR_{post} - FR_{pre})/(FR_{post} + FR_{pre})$[93].

Neurons with MI between 0.2 and 1 were considered as activated and with MI between -0.2

and -1, as inhibited. Both the activated and inhibited neurons were assigned to the modulated

PVT (PVT_{MOD}) population. Based on these, neurons with MI between 0.2 and -0.2 were

assumed as non-modulated cells (PVT_{NON}; **Fig. 1i,j**).

Furthermore, in the chronic and anaesthetized data, thalamic unit activity was correlated with

theta waves (phase-locking analysis in **Fig. 1l and 2k**). Based on the dominance of the

distinct frequency band (delta, theta, sigma, gamma, EMG) the sleep stages were

automatically scored. The different oscillations were normalized to the maximum power.

REM was assigned to suprathreshold theta and subthreshold delta power, with low EMG. The

sorted REM stages are cut out from the full measurement. After low-pass filtering the peaks

of theta waves were defined. We have defined theta waves within a 0.18 s width value and

with maximum 100-microvolt amplitude. Based on the width of waves the start and the end

point of the theta waves are calculated. Using this data, sinusoidal waves were fitted and these

hypothetical sinusoidal waves were phase-locked to the spiking activity of thalamic neurons.

Statistical analysis

The number of analyzed slices and number of cells are shown with N, while n represents the

number of mice. Data from independent experiments or animals were pooled when it was

possible. Statistical significance was assessed using Kruskal-Wallis ANOVA with Mann–

Whitney U-tests, Friedman ANOVA with Wilcoxon Matched Pairs Test, Paired t-test or

Student's t-test with normality test, as well as analysis of variance (one-way ANOVA,

repeated measures ANOVA) to analyze the axon density data or the electrophysiological data.

Normality was tested using the Shapiro-Wilk test. Pearson's χ^2 tests were used to compare the

distribution of the responsive cortical cells. Post hoc test is used for ANOVA analysis (Fisher

LSD or Bonferroni Correction). For correlation analysis Kendall's τ coefficient was used to

measure the ordinal association between two measured quantities. The Kolmogorov–Smirnov

test was used to validate the distribution of a sample and to compare the distributions of two

independent sample groups. An Equivalence Independent Samples T-test (TOST) was

performed to assess the correspondence between data obtained under anesthesia and in freely

moving conditions [94,95]. Statistical analysis was performed using JASP, MATLAB or

Microsoft Office Excel. Significance levels are indicated as follows: $0.1 > \#P > 0.05$; $*P <$

0.05; $**P < 0.01$; $***P < 0.001$; $****P < 0.0001$. No statistical methods were used to

predetermine sample size, but it is comparable to previously published work[25,96,97].

References

1.   della Monica, C., Johnsen, S., Atzori, G., Groeger, J. A. & Dijk, D.-J. Rapid Eye

Movement Sleep, Sleep Continuity and Slow Wave Sleep as Predictors of Cognition,

Mood, and Subjective Sleep Quality in Healthy Men and Women, Aged 20–84 Years.

Front. Psychiatry **9,** (2018).

2.   Lewis, P. A., Knoblich, G. & Poe, G. How Memory Replay in Sleep Boosts Creative

Problem-Solving. *Trends Cogn. Sci.* **22,** 491–503 (2018).

3.   Li, W., Ma, L., Yang, G. & Gan, W.-B. REM sleep selectively prunes and maintains

new synapses in development and learning. *Nat. Neurosci.* **20,** 427–437 (2017).

4.   Walker, M. P., Liston, C., Hobson, J. A. & Stickgold, R. Cognitive flexibility across

the sleep–wake cycle: REM-sleep enhancement of anagram problem solving. *Cogn.*

Brain Res. **14,** 317–324 (2002).

5.   Nishida, M., Pearsall, J., Buckner, R. L. & Walker, M. P. REM Sleep, Prefrontal Theta,

and the Consolidation of Human Emotional Memory. *Cereb. Cortex* **19,** 1158–1166

(2009).

6.   Nir, Y. *et al.* Selective neuronal lapses precede human cognitive lapses following sleep

deprivation. *Nat. Med.* **23,** 1474–1480 (2017).

7.   Wulff, K., Gatti, S., Wettstein, J. G. & Foster, R. G. Sleep and circadian rhythm

disruption in psychiatric and neurodegenerative disease. *Nat. Rev. Neurosci.* **11,** 589–

599 (2010).

8.   Jozwiak, N. *et al.* REM Sleep Behavior Disorder and Cognitive Impairment in

Parkinson's Disease. *Sleep* **40,** (2017).

9.   Katsuki, F., Gerashchenko, D. & Brown, R. E. Alterations of sleep oscillations in

Alzheimer's disease: A potential role for GABAergic neurons in the cortex,

hippocampus, and thalamus. *Brain Res. Bull.* **187,** 181–198 (2022).

10. Brown, R. E., Basheer, R., McKenna, J. T., Strecker, R. E. & McCarley, R. W. Control

of sleep and wakefulness. *Physiological Reviews* (2012).

doi:10.1152/physrev.00032.2011

11. Peever, J. & Fuller, P. M. The Biology of REM Sleep. *Current Biology* (2017).

doi:10.1016/j.cub.2017.10.026

12. Scammell, T. E., Arrigoni, E. & Lipton, J. O. Neural Circuitry of Wakefulness and

Sleep. *Neuron* (2017). doi:10.1016/j.neuron.2017.01.014

13. Peever, J. & Fuller, P. M. Neuroscience: A Distributed Neural Network Controls REM

Sleep. *Curr. Biol.* **26,** R34–R35 (2016).

14. Weber, F. *et al.* Control of REM sleep by ventral medulla GABAergic neurons. *Nature*

526, 435–438 (2015).

15. Valencia Garcia, S. *et al.* Ventromedial medulla inhibitory neuron inactivation induces

REM sleep without atonia and REM sleep behavior disorder. *Nat. Commun.* **9,** 504

(2018).

16. Dong, Y. & Liu, D. Cracking the complexity of REM sleep. *Trends Neurosci.* (2025).

doi:10.1016/j.tins.2025.10.008

17. Hong, J., Lozano, D. E., Beier, K. T., Chung, S. & Weber, F. Prefrontal cortical

regulation of REM sleep. *Nat. Neurosci.* **26,** 1820–1832 (2023).

18. Aime, M. *et al.* Paradoxical somatodendritic decoupling supports cortical plasticity

817  during REM sleep. *Science (80-.).* **376,** 724–730 (2022).

818 19. Benington, J. H. & Heller, H. C. REM-sleep timing is controlled homeostatically by

819  accumulation of REM-sleep propensity in non-REM sleep. *Am. J. Physiol. Integr.*

820  *Comp. Physiol.* **266,** R1992–R2000 (1994).

821 20. Vivaldi, E. A., Ocampo, A., Wyneken, U., Roncagliolo, M. & Zapata, A. M. Short-

822  term homeostasis of active sleep and the architecture of sleep in the rat. *J.*

823  *Neurophysiol.* (1994). doi:10.1152/jn.1994.72.4.1745

824 21. Steriade, M., McCormick, D. A. & Sejnowski, T. J. Thalamocortical oscillations in the

825  sleeping and aroused brain. *Science* **262,** 679–685 (1993).

826 22. Sherman SM, G. R. *Exploring the thalamus and its role in cortical function. 2nd ed.*

827  (MIT Press, 2006).

828 23. Shine, J. M., Lewis, L. D., Garrett, D. D. & Hwang, K. The impact of the human

829  thalamus on brain-wide information processing. *Nat. Rev. Neurosci.* **24,** 416–430

830  (2023).

831 24. Vertes, R. P., Linley, S. B. & Hoover, W. B. Limbic circuitry of the midline thalamus.

832  *Neuroscience and Biobehavioral Reviews* **54,** 89–107 (2015).

833 25. Mátyás, F. *et al.* A highly collateralized thalamic cell type with arousal-predicting

834  activity serves as a key hub for graded state transitions in the forebrain. *Nat. Neurosci.*

835  **21,** 1551–1562 (2018).

836 26. Ren, S. *et al.* The paraventricular thalamus is a critical thalamic area for wakefulness.

837  *Science (80-.).* **362,** 429–434 (2018).

838 27. Ma, J. *et al.* Divergent projections of the paraventricular nucleus of the thalamus

mediate the selection of passive and active defensive behaviors. *Nat. Neurosci.* **24,**

1429–1440 (2021).

28. Zhu, Y. *et al.* Dynamic salience processing in paraventricular thalamus gates

associative learning. *Science (80-.).* **362,** 423–429 (2018).

29. Colavito, V., Tesoriero, C., Wirtu, A. T., Grassi-Zucconi, G. & Bentivoglio, M. Limbic

thalamus and state-dependent behavior: The paraventricular nucleus of the thalamic

midline as a node in circadian timing and sleep/wake-regulatory networks. *Neurosci.*

Biobehav. Rev. **54,** 3–17 (2015).

30. Herman, J. P. Neural control of chronic stress adaptation. *Front. Behav. Neurosci.* **7,**

(2013).

31. Hsu, D. T., Kirouac, G. J., Zubieta, J.-K. & Bhatnagar, S. Contributions of the

paraventricular thalamic nucleus in the regulation of stress, motivation, and mood.

Front. Behav. Neurosci. **8,** (2014).

32. Pliota, P. *et al.* Stress peptides sensitize fear circuitry to promote passive coping. *Mol.*

Psychiatry **25,** 428–441 (2020).

33. Jász, A. *et al.* Persistently increased post-stress activity of paraventricular thalamic

neurons is essential for the emergence of stress-induced alterations in behaviour. *PLOS*

Biol. **23,** e3002962 (2025).

34. Beas, B. S. *et al.* The locus coeruleus drives disinhibition in the midline thalamus via a

dopaminergic mechanism. *Nature Neuroscience* (2018). doi:10.1038/s41593-018-

0167-4

35. Csercsa, R. *et al.* Laminar analysis of slow wave activity in humans. *Brain* **133,** 2814–

2829 (2010).

36. Funk, C. M., Honjoh, S., Rodriguez, A. V., Cirelli, C. & Tononi, G. Local Slow Waves

in Superficial Layers of Primary Cortical Areas during REM Sleep. *Curr. Biol.* **26,**

396–403 (2016).

37. Dong, Y., Li, J., Zhou, M., Du, Y. & Liu, D. Cortical regulation of two-stage rapid eye

movement sleep. *Nat. Neurosci. 2022 2512* **25,** 1675–1682 (2022).

38. Dong, X., Li, S. & Kirouac, G. J. Collateralization of projections from the

paraventricular nucleus of the thalamus to the nucleus accumbens, bed nucleus of the

stria terminalis, and central nucleus of the amygdala. *Brain Structure and Function* 1–

17 (2017). doi:10.1007/s00429-017-1445-8

39. Aquino-Miranda, G. *et al.* Functional properties of corticothalamic circuits targeting

paraventricular thalamic neurons. *Neuron* **112,** 4060–4080.e7 (2024).

40. Suchecki, D., Tiba, P. A. & Machado, R. B. REM Sleep Rebound as an Adaptive

Response to Stressful Situations. *Front. Neurol.* **3,** (2012).

41. Kim, E. J. & Dimsdale, J. E. The effect of psychosocial stress on sleep: A review of

polysomnographic evidence. *Behav. Sleep Med.* **5,** 256–278 (2007).

42. Yu, X. *et al.* A specific circuit in the midbrain detects stress and induces restorative

sleep. *Science (80-.).* **377,** 63–72 (2022).

43. Yu, X., Nollet, M., Franks, N. P. & Wisden, W. Sleep and the recovery from stress.

Neuron **113,** p2910-2926 (2025).

44. Niethard, N. *et al.* Sleep-Stage-Specific Regulation of Cortical Excitation and

Inhibition. *Curr. Biol.* (2016). doi:10.1016/j.cub.2016.08.035

45.   Penzo, M. A. & Gao, C. The paraventricular nucleus of the thalamus: an integrative

node underlying homeostatic behavior. *Trends Neurosci.* **44,** 538–549 (2021).

46.   Kirouac, G. J. Placing the paraventricular nucleus of the thalamus within the brain

circuits that control behavior. *Neuroscience and Biobehavioral Reviews* **56,** 315–329

(2015).

47.   Misgeld, U., Deisz, R. A., Dodt, H. U. & Lux, H. D. The role of chloride transport in

postsynaptic inhibition of hippocampal neurons. *Science (80-.).* **232,** 1413–1415

(1986).

48.   Hee, J. C. *et al.* Excitatory actions of GABA in the suprachiasmatic nucleus. *J.*

Neurosci. **28,** 5450–5459 (2008).

49.   Sørensen, A. T. *et al.* Altered Chloride Homeostasis Decreases the Action Potential

Threshold and Increases Hyperexcitability in Hippocampal Neurons. *eneuro* **4,**

ENEURO.0172-17.2017 (2017).

50.   Osorio-Forero, A. *et al.* Infraslow noradrenergic locus coeruleus activity fluctuations

are gatekeepers of the NREM–REM sleep cycle. *Nat. Neurosci.* **28,** 84–96 (2025).

51.   Jego, S. *et al.* Optogenetic identification of a rapid eye movement sleep modulatory

circuit in the hypothalamus. *Nat. Neurosci.* **16,** 1637–1643 (2013).

52.   Van Dort, C. J. *et al.* Optogenetic activation of cholinergic neurons in the PPT or LDT

induces REM sleep. *Proc. Natl. Acad. Sci.* **112,** 584–589 (2015).

53.   Kroeger, D. *et al.* Galanin neurons in the ventrolateral preoptic area promote sleep and

heat loss in mice. *Nat. Commun.* **9,** 4129 (2018).

54.   Perlman, G., Tanti, A. & Mechawar, N. Parvalbumin interneuron alterations in stress-

related mood disorders: A systematic review. *Neurobiol. Stress* **15,** 100380 (2021).

55.   Chen, C.-C., Lu, J., Yang, R., Ding, J. B. & Zuo, Y. Selective activation of

parvalbumin interneurons prevents stress-induced synapse loss and perceptual defects.

Mol. Psychiatry **23,** 1614–1625 (2018).

56.   Gądek-Michalska, A., Tadeusz, J., Rachwalska, P., Spyrka, J. & Bugajski, J. Effect of

repeated restraint on homotypic stress-induced nitric oxide synthases expression in

brain structures regulating HPA axis. *Pharmacol. Reports* **64,** 1381–1390 (2012).

57.   Le Roux, N., Amar, M., Moreau, A. W. & Fossier, P. Roles of nitric oxide in the

homeostatic control of the excitation–inhibition balance in rat visual cortical networks.

Neuroscience **163,** 942–951 (2009).

58.   Chrousos, G. P. & Gold, P. W. The concepts of stress and stress system disorders.

Overview of physical and behavioral homeostasis. *JAMA* **267,** 1244–52 (1992).

59.   Kimura, M. *et al.* Conditional corticotropin-releasing hormone overexpression in the

mouse forebrain enhances rapid eye movement sleep. *Mol. Psychiatry* **15,** 154–165

(2010).

60.   Marrosu, F., Gessa, G. L., Giagheddu, M. & Fratta, W. Corticotropin-releasing factor

(CRF) increases paradoxical sleep (PS) rebound in PS-deprived rats. *Brain Res.* **515,**

315–318 (1990).

61.   Steiger, A. Sleep and the hypothalamo–pituitary–adrenocortical system. *Sleep Med.*

Rev. **6,** 125–138 (2002).

62.   Itoga, C. A. *et al.* New viral-genetic mapping uncovers an enrichment of corticotropin-

releasing hormone-expressing neuronal inputs to the nucleus accumbens from stress-

related brain regions. *J. Comp. Neurol.* **527,** 2474–2487 (2019).

63. Arnsten, A. F. T. Stress signalling pathways that impair prefrontal cortex structure and

function. *Nat. Rev. Neurosci.* **10,** 410–422 (2009).

64. Cerqueira, J. J., Mailliet, F., Almeida, O. F. X., Jay, T. M. & Sousa, N. The Prefrontal

Cortex as a Key Target of the Maladaptive Response to Stress. *J. Neurosci.* **27,** 2781–

2787 (2007).

65. Maier, S. F. & Watkins, L. R. Role of the medial prefrontal cortex in coping and

resilience. *Brain Res.* **1355,** 52–60 (2010).

66. Cabib, S. & Puglisi-Allegra, S. The mesoaccumbens dopamine in coping with stress.

Neurosci. Biobehav. Rev. **36,** 79–89 (2012).

67. Campioni, M. R., Xu, M. & McGehee, D. S. Stress-induced changes in nucleus

accumbens glutamate synaptic plasticity. *J. Neurophysiol.* (2009).

doi:10.1152/jn.91111.2008

68. Dong, X., Li, S. & Kirouac, G. J. A projection from the paraventricular nucleus of the

thalamus to the shell of the nucleus accumbens contributes to footshock stress-induced

social avoidance. *Neurobiol. Stress* (2020). doi:10.1016/j.ynstr.2020.100266

69. De Groote, A. & de Kerchove d'Exaerde, A. Thalamo-Nucleus Accumbens Projections

in Motivated Behaviors and Addiction. *Front. Syst. Neurosci.* **15,** (2021).

70. Wang, T. *et al.* Paraventricular thalamus dynamically modulates aversive memory via

tuning prefrontal inhibitory circuitry. *J. Neurosci.* **43,** 3630–3646. (2023).

71. Lai, C. S. W., Franke, T. F. & Gan, W. B. Opposite effects of fear conditioning and

extinction on dendritic spine remodelling. *Nat. 2012 4837387* **483,** 87–91 (2012).

72.   Zhou, Y. *et al.* REM sleep promotes experience-dependent dendritic spine elimination

in the mouse cortex. *Nat. Commun.* **11,** 4819 (2020).

73.   Pi, H. J. *et al.* Cortical interneurons that specialize in disinhibitory control. *Nat. 2013*

5037477 **503,** 521–524 (2013).

74.   Ferguson, B. R. & Gao, W. J. Pv interneurons: critical regulators of E/I balance for

prefrontal cortex-dependent behavior and psychiatric disorders. *Front. Neural Circuits*

12, 37 (2018).

75.   Fino, E. & Yuste, R. Dense Inhibitory Connectivity in Neocortex. *Neuron* **69,** 1188–

1203 (2011).

76.   Packer, A. M. & Yuste, R. Dense, Unspecific Connectivity of Neocortical

Parvalbumin-Positive Interneurons: A Canonical Microcircuit for Inhibition? *J.*

Neurosci. **31,** 13260–13271 (2011).

77.   Funk, C. M. *et al.* Role of somatostatin-positive cortical interneurons in the generation

of sleep slow waves. *J. Neurosci.* **37,** 9132–9148 (2017).

78.   Yizhar, O. *et al.* Neocortical excitation/inhibition balance in information processing

and social dysfunction. *Nature* **477,** 171–178 (2011).

79.   Sohal, V. S. & Rubenstein, J. L. R. Excitation-inhibition balance as a framework for

investigating mechanisms in neuropsychiatric disorders. *Mol. Psychiatry* **24,** 1248–

1257 (2019).

80.   Selten, M., van Bokhoven, H. & Nadif Kasri, N. Inhibitory control of the

excitatory/inhibitory balance in psychiatric disorders. *F1000Research* **7,** 23 (2018).

81.   Radley, J. J. *et al.* Repeated Stress Induces Dendritic Spine Loss in the Rat Medial

Prefrontal Cortex. *Cereb. Cortex* **16,** 313–320 (2006).

82.   Canter, R. G., Penney, J. & Tsai, L.-H. The road to restoring neural circuits for the

treatment of Alzheimer's disease. *Nature* **539,** 187–196 (2016).

83.   Busche, M. A. & Konnerth, A. Impairments of neural circuit function in Alzheimer's

disease. *Philos. Trans. R. Soc. B Biol. Sci.* **371,** 20150429 (2016).

84.   Shu, Y., Hasenstaub, A. & McCormick, D. A. Turning on and off recurrent balanced

cortical activity. *Nature* **423,** 288–293 (2003).

85.   Ferguson, B. R. & Gao, W. J. Thalamic Control of Cognition and Social Behavior Via

Regulation of Gamma-Aminobutyric Acidergic Signaling and Excitation/Inhibition

Balance in the Medial Prefrontal Cortex. *Biol. Psychiatry* **83,** 657–669 (2018).

86.   Barsy, B. *et al.* Associative and plastic thalamic signaling to the lateral amygdala

controls fear behavior. *Nat. Neurosci.* **23,** 625–637 (2020).

87.   Babiczky, Á. & Matyas, F. Molecular characteristics and laminar distribution of

prefrontal neurons projecting to the mesolimbic system. *Elife* **11,** (2022).

88.   Mátyás, F., Lee, J., Shin, H. S. & Acsády, L. The fear circuit of the mouse forebrain:

Connections between the mediodorsal thalamus, frontal cortices and basolateral

amygdala. *Eur. J. Neurosci.* **39,** 1810–1823 (2014).

89.   Steriade, M. Corticothalamic resonance, states of vigilance and mentation.

Neuroscience **101,** 243–276 (2000).

90.   Rossant, C. *et al.* Spike sorting for large, dense electrode arrays. *Nat. Neurosci.* **19,**

634–641 (2016).

91.   Barthó, P. *et al.* Characterization of Neocortical Principal Cells and Interneurons by

Network Interactions and Extracellular Features. *J. Neurophysiol.* **92,** 600–608 (2004).

92.   Gentet, L. J., Avermann, M., Matyas, F., Staiger, J. F. & Petersen, C. C. H. Membrane

Potential Dynamics of GABAergic Neurons in the Barrel Cortex of Behaving Mice.

Neuron **65,** 422–435 (2010).

93.   Michaiel, A. M., Parker, P. R. L. & Niell, C. M. A Hallucinogenic Serotonin-2A

Receptor Agonist Reduces Visual Response Gain and Alters Temporal Dynamics in

Mouse V1. *Cell Rep.* **26,** 3475–3483.e4 (2019).

94.   Walker, E. & Nowacki, A. S. Understanding equivalence and noninferiority testing.

Journal of General Internal Medicine **26,** 192–196 (2011).

95.   Pardoe, H. R. *et al.* Pooling Morphometric Estimates: A Statistical Equivalence

Approach. *J. Neuroimaging* **26,** 109–115 (2016).

96.   Do-Monte, F. H., Quinõnes-Laracuente, K. & Quirk, G. J. A temporal shift in the

circuits mediating retrieval of fear memory. *Nat. 2015 5197544* **519,** 460–463 (2015).

97.   Do-Monte, F. H., Minier-Toribio, A., Quiñones-Laracuente, K., Medina-Colón, E. M.

& Quirk, G. J. Thalamic Regulation of Sucrose Seeking during Unexpected Reward

Omission. *Neuron* **94,** 388–400.e4 (2017).

1009

1010

Acknowledgements

We thank Z.J. Huang and László Acsády for providing us transgenic mice, Tamás Herczeg, Réka Erdős, Anna Fehér, and Katalin Varga for laboratory assistance, Csaba Dávid and Ákos Babiczky for advising us on axonal analysis and Gábor Nyíri for sharing primary antibodies. We thank Tallie Z. Baram, Antoine Adamantidis, Nadia Urbain and Iván Soltész for comments and discussions about the manuscript. We also thank the Institute of Molecular Life Sciences of the HUN-REN Research Centre for Natural Sciences for the use of Zeiss microscope and we would like to thank the Light Microscopy Center of HUN-REN Institute of Experimental Medicine. We also thank the Virus Technology Unit of HUN-REN Institute of Experimental Medicine. This work was supported by the following funding agencies: Ministry of Innovation and Technology of Hungary from the National Research, Development and Innovation Fund (K138836, and KKP126998 to FM; FK135285, FK144583 to SB; PD124034 to BB; HU-RIZONT-2024-00003 to AM); Hungarian Brain Research Program (2017-1.2.1-NKP-2017-00002 to FM; NAP2022-I-7/2022 to MLL); New National Excellence Program of the Ministry for Innovation and Technology (ÚNKP-21-5-ÁTE-2 to FM; ÚNKP-22-2-III-ELTE-539 to AVB, ELKH SA-48/2021); National Natural Science Foundation of China (8234101126 to DL); Cooperative Doctoral Program (KDP-2020-1015461 to AM); New national excellence program of the ministry for culture and innovation from the source of the national research, development and innovation fund (EKÖP-2024-87 to PB, EKÖP-2024-276 to AVB, 2025-2.1.1-EKÖP-2025-00014 to AM). FM was a János Bolyai Research Fellow.

Contributions

1034 A.M., P.B., B.B. and F.M. conceived and designed the project. A.M., P.B., B.B., A.V.B. and

1035 F.M. performed experiments. A.M., P.B., S.B. and F.M. analysed data. S.B., M.L.L and F.M.

1036 supervised electrophysiology experiments. F.M. supervised the entire project. A.M., P.B.,

1037 M.L.L., D.L. and F.M. prepared figures and wrote the original draft. The other authors made

1038 valuable comments on the manuscript and figures.

1039

1040 **Data availability**

1041 All data that support the findings of this study are available from the corresponding author

1042 upon request. Source data are provided with this paper.

1043

1044 **Code availability**

1045 Custom-written codes used to analyze data from this study are available from the

1046 corresponding author upon request.

1047

1048 **Declaration of interests**

1049 The authors declare no competing interests.

Fig. 1: VLM_GABA shape REM-related cortical theta oscillation via PVT

a, L*eft,* experimental design for prefrontal cortical LFP recordings with optogenetic activation of VLM_GABA neurons. *Right*, injection sites targeting VLM_GABA cells with AAV-DIO-ChR2 in vGAT-Cre mice. **b,** *Left*, effect of VLM_GABA stimulation on the LFP spectrogram. *Right*, population data for theta (θ) band power changes induced by VLM_GABA stimulation (n = 8 animals; Wilcoxon matched-pairs test). **c,** Quantification of topographic distribution of VLM_GABA-to-PVT inputs (n = 3 animals; Shapiro-Wilk W test with paired t-test). **d,** Representative confocal images of eYFP-labeled VLM_GABA-to-PVT (green) at three anteroposterior levels. Cyan color shows the CR counterstaining indicating territory of PVT. **e,** Schematic (*left*) of and injection site (*right*) targeting the PFC with retrograde (CTB) tracer. **f,** Representative confocal images of CTB-labeled PFC-projecting PVT cells (red) at three anteroposterior levels corresponding to **d**. Cyan color shows the CR counterstaining indicating territory of PVT. Insets, high magnification images of CTB labeled cells. Asterisks indicate

cortico-thalamic fibers. **g,** Experimental design for simultaneous PVT and PFC recordings with optogenetic silencing (NpHr3.0) of VLM_{GABA}-to-PVT inputs. **h,** Pie charts showing state-dependent responsiveness of PVT cells (N = 95; n = 7 mice) to inhibition of VLM_{GABA}-to-PVT inputs (Pearson's χ2 test). *ALL STIM,* all trials; *High Theta STIM,* inhibition during high theta power (REM-like) periods. **i,** Baseline (BL) and evoked PVT activity by inhibition of VLM_{GABA}-to-PVT inputs during high theta trials ($NpHR_{HT}$) clustered by responsiveness to the inhibition [PVT_{MOD} (purple) vs. PVT_{NON} (orange)] [Wilcoxon matched-pairs test for BL vs. $NpHR_{HT}$; Mann-Whitney U test for PVT_{MOD} (N=68 cells) vs. $PVT_{NON-MOD}$ (N=27 cells)]. **j,** Firing rate changes upon NREM-like (N)-to-REM-like (R) state transition were observed only in PVT_{MOD} cells (Wilcoxon matched-pairs test for NREM vs. REM; Mann-Whitney U test for PVT_{MOD} vs. PVT_{NON}). **k,** Negative correlation between mean firing rate (mFR) and high theta-induced firing rate changes (R/N) observed only in PVT_{MOD} cells (Kendall's TAU Correlations). **l,** Vector diagrams represent distinct, phase-locked firing of PVT_{MOD} and PVT_{NON} cells during theta waves (Pearson's χ2 test). **m,** *Left,* Effect of optogenetic silencing of VLM_{GABA}-to-PVT inputs on the LFP spectrogram. *Right,* Population data for theta (θ) band power changes provoked by inhibition of VLM_{GABA}-to-PVT inputs (green) relative to pre-stim (black) in *Hight Theta* STIM trials (n = 7 animals; Wilcoxon matched-pairs test). *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$. (Statistics in **Extended Data Table 1**).

Fig. 2: PVT→NAc neurons mediate the impact of VLM_GABA on cortical theta oscillation and REM sleep

a, Experimental design for slice electrophysiology in PVT of vGAT-Cre mice examining VLM_GABA inputs onto projection-specific PVT cells. **b,** Representative high-magnification Z-stack confocal images (3 μm total in depth) showing biocytin-filled (white), PVT→NAc (green; *left*) and PVT→AMY (green; *right*) cells contacted by VLM_GABA inputs (red). **c,** *Left,* optogenetically evoked GABAergic IPSCs (eIPSC; Ctrl) in PVT cells (N = 4) were completely abolished by gabazine (GBZ; orange). *Right,* example traces of a PVT→NAc cell in Ctrl (black) and GBZ (orange) conditions (Student's t-test). **d,** Population data for spontaneous and evoked IPSCs of PVT→NAc (green) and PVT→AMY cells (red; N = 7-7; Student's t-test). **e,** Example traces of a PVT→NAc and PVT→AMY cell for frequency-dependent eIPSCs. **f,** Population data for eIPSC amplitude and success rate (One-Way ANOVA, Bonferroni's post hoc comparisons). **g,** Experimental design for in vivo recordings of PVT→NAc or PVT→AMY cells during natural sleep. **h,** Differences in mean firing rate (mFR) of PVT→NAc (N = 25 cells; n = 7 animals) or PVT→AMY cells (N = 36 cells; n = 8 animals; Mann-Whitney U test). **i,** Elevation in mean firing rate at NREM-to-REM transition for PVT→NAc (*left*) but not PVT→AMY cells (*right*; Wilcoxon matched-pairs test). Pink color, 1 s bins with significant elevation (s.b.) compared to baseline (bl, grey). **j,** Negative correlation between mean firing rate (mFR) and REM induced firing rate changes (R/N) restricted to PVT→NAc cells Kendall's TAU Correlations). **k,** Distinct phase-locked firing of PVT→NAc (green) and PVT→AMY cells (red) during REM-associated theta oscillation (Pearson's χ2 test). *Top,* example unit activity of a PVT→NAc (green) and PVT→AMY cell (red). *Bottom,* vector diagram of population data. **l,m,** Frequency-dependent bidirectional effects of optogenetic stimulation of PVT→NAc and PVT→AMY cells on REM-associated theta oscillation indicated by wavelets (**l**, **m**, *left*) and quantification of theta band power changes (**l,m**, *right*). **n,** Summary data for the impact of frequency-dependent optogenetic stimulation on theta band power (n = 7-8 mice; Wilcoxon matched-pairs test). **o,** Population data for optogenetically evoked state transitions (*top*), duration of REM episodes (*middle*) and the

calculated effectiveness to change REM sleep (*bottom*). [#]$P < 0.1$; *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.0001$. (Statistics in **Extended Data Table 1**).

Fig. 3: PVT$_{\rightarrow NAc}$ bidirectionally modulate the VLM$_{GABA}$ effect on theta oscillation in activity- and firing mode-dependent manner

a, Experimental design for investigating the impact of VLM$_{GABA}$ and PVT$_{\rightarrow NAc}$ optogenetic co-activation. **b,** Baseline (BL) and evoked PVT activity induced by stimulation of VLM$_{GABA}$ neurons clustered by responsiveness to antidromic phototagging from NAc (PVT$_{\rightarrow NAc}$; N=41 cells) vs. untagged cells (PVT$_{UN}$,N= 39 cells; n=8 mice; Wilcoxon matched-pairs test for BL vs. VLM$_{ChR2}$; Mann-Whitney U test for PVT$_{\rightarrow NAc}$ vs. PVT$_{UN}$). **c,** Baseline and evoked PVT$_{\rightarrow NAc}$ activity during paired stimulation of VLM$_{GABA}$ and PVT$_{\rightarrow NAc}$ at 1 and 10 Hz (Wilcoxon matched-pairs test). **d,** Averaged activity changes of PVT$_{\rightarrow NAc}$ neurons evoked by VLM$_{GABA}$ orthodromic and PVT$_{\rightarrow NAc}$ antidromic optogenetic co-activation indicated as z-scores. **e,** Inter-spike interval (ISI) distribution for PVT$_{\rightarrow NAc}$ neurons during pre-STIM and STIM periods (10 Hz VLM$_{GABA}$ + 1 Hz PVT$_{\rightarrow NAc}$ and 10 Hz VLM$_{GABA}$ + 10 Hz PVT$_{\rightarrow NAc}$). Note the multimodal ISI distribution (arrows) observed during 10 Hz VLM$_{GABA}$ + 1 Hz PVT$_{\rightarrow NAc}$ stimulation (Kolmogorov-Smirnov test). **f,** Effect of the three stimulation protocols on the LFP spectrogram. **g,** Frequency-dependent, bidirectional effects of paired stimulation of VLM$_{GABA}$ and PVT$_{\rightarrow NAc}$ on theta oscillation (Wilcoxon matched-pairs test). #P < 0.1; *P < 0.05; **P < 0.01; ****P < 0.0001. (Statistics in **Extended Data Table 1**).

Fig. 4: Activity changes of PVT→NAc neurons are also responsible for stress-induced bidirectional modulation of REM sleep

a, Experimental design for investigating stress-dependent REM sleep alterations. **b,** Acute (Ac.) and chronic (Chr.) restraint induced changes in REM state during post-restraint sleep periods (PRS1-3) compared to the habituation day (Hab.; PRS0) indicated as proportion of REM sleep (*left*) and theta-band power (*right*) (n = 6 mice; Wilcoxon matched-pairs test). **c,d,** Population data for restraint-induced activity changes of PVT→NAc (*left*; N = 11 cells) and PVT→AMY cells (*right*; N = 9 cells) during restraint [R0-R3; (**c**)] and post-restraint REM sleep periods [PRS0-PRS3; (**d**)] (Repeated Measures ANOVA with post hoc LSD test). **e,** Experimental design for investigating the impacts of optogenetic activation of PVT→NAc (n = 7 mice) and PVT→AMY neurons (n = 8 mice) on REM sleep. **f,g,** Optogenetic stimulation (blue) of PVT→NAc (**f**) but not PVT→AMY (**g**) triggered REM state changes similar to those induced by acute and chronic restraint stress (Wilcoxon matched-pairs test). **h,** Population data indicating REM-sleep-related activity changes of PVT→NAc (N = 25) and PVT→AMY (N= 36) neurons during non-stim periods (Repeated Measures ANOVA with post hoc LSD test). *P < 0.05; **P < 0.01; ****P < 0.0001. (Statistics in **Extended Data Table 1**).

Fig. 5: Frequency-dependent and layer-specific effects of PVT→NAc on PFC circuits underlying modulation of cortical theta oscillation

a, Experimental design for investigating the effects of PVT→NAc or PVT→AMY activation on PFC circuits. **b,** Distinct, layer-specific distribution of PVT→NAc and PVT→AMY inputs in PFC (One-way ANOVA with post hoc LSD test). **c-h,** Distinct, layer-specific effects of PVT→NAc and PVT→AMY activation in PFC. Representative cases for evoked multiunit activity (eMUA) in the superficial (**c**) and deep layers of PFC (**f**); population data for eMUA (**a,g**; n= 4 mice; Wilcoxon matched-pairs test); population data for signal-to-noise ratio of eMUA (**e,h**; n=4 mice; Repeated Measures ANOVA with post hoc LSD test). **i,** Opposite effects of PFC FS (N=13 cells), NFS (N= 33 cells) and PN (N= 37 cells) triggered by 1 Hz and 10 Hz activation of PVT→NAc (n=7 mice; Wilcoxon matched-pairs test for baseline vs 1 Hz vs. 10 Hz; Mann-Whitney U test for FS vs. NFS vs PN). **j,** Similar shifts in PFC FS, NFS and PN activation at spontaneous state changes: from high delta periods to high theta periods [similar to NREM (N) → REM (R)] and from high theta periods to desynchronized state [similar to REM (R) → post-REM (R)] (n=7 mice; Wilcoxon matched-pairs test for delta→theta vs. theta→desyn; Mann-Whitney U test for FS vs. NFS vs. PN). **k,** PVT→AMY stimulation did not elicit state-dependent activation of PFC FS, NFS and PN (n=7 mice; Wilcoxon matched-pairs test for baseline vs 1 Hz vs 10 Hz; Kruskal-Wallis ANOVA for FS vs. NFS vs. PN). *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001. (Statistics in **Extended Data Table 1**).

Fig. 6: Circuit diagram for PVT→NAc activity-mediated REM sleep alterations in normal and stressful conditions.