Novel Gain of Function Mouse Model of KCNT1-Related Epilepsy

Sarah L. Olguin[1,2], Daniel Davis[3,4]; Ammara Rehman[1,5], Sean W. Berg[1,2], Elizabeth Sahagun[1,2]; Payton S. Oswalt[3], Mandar Patil[1,5], Paula D.M. Sullivan[1,5], Bing Zhang[6], **Elizabeth C. Bryda[3,4*], Roy Ben-Shalom[1,5*,] Jill L. Silverman[1,2*]**

Affiliations:
[1]MIND Institute, University of California Davis School of Medicine, Sacramento, CA 95817, USA. [2]Department of Psychiatry and Behavioral Sciences, University of California Davis School of Medicine, Sacramento, CA 95817, USA. [3]Veterinary Pathobiology, College of Veterinary Medicine, University of Missouri, Columbia, MO, 65201, USA. [4] MU Animal Modeling Core University of Missouri, Columbia, MO, 65201, USA. [5]Department of Neurology, University of California Davis School of Medicine, Sacramento, CA 95817, USA [6]Division of Biological Sciences, University of Missouri, Columbia, MO 65201, USA.

*Co-Corresponding Authors:
Jill L. Silverman
MIND Institute and Department of Psychiatry and Behavioral Sciences
4625 2[nd] Ave Suite 1001A
Sacramento, CA 95817
Phone: 916-734-8531
jsilverman@ucdavis.edu

*Co-Corresponding Authors:
Elizabeth Bryda, Ph.D.
Department of Veterinary Pathobiology, University of Missouri
4011 Discovery Drive
Columbia, MO 65201
Phone: (573) 882-5504
brydae@missouri.edu

*Co-Corresponding Authors:
Roy Ben-Shalom
MIND Institute and Department of Neurology
rbenshalom@ucdavis.edu

Keywords: *KCNT1, Epilepsy, epileptic encephalopathy, behavior, seizures, excitability, mouse model, hyperexcitability, epilepsy, channel, potassium, DEE*

Conflict of Interest: The authors declare no conflicts of interest.

Acknowledgements: This work was supported by the MIND Institute's Intellectual and Developmental Disabilities Resource Center NIH P50HD103526 (LA), the NIMH ARTP R01MH117130 (JAG), and Research Council Grant URC-21-027 from the Office of Research,

University of Missouri (ECB). The mouse model was generated by the University of Missouri
Animal Modeling Core.
Funding Sources: This work was supported by the MIND Institute's Intellectual and
Developmental Disabilities Resource Center P50HD103526 and an MDBR award from the
KCNT1 Foundation and the University of Pennsylvania Orphan Disease Center (JLS, SLO),
Hartwell Foundation Biomedical Fellowship (JLS; SLO), and Research Council Grant URC-21-
027 from the Office of Research, University of Missouri (ECB).
52
53

Abstract

KCNT1-related epilepsy is an autosomal dominant neurodevelopmental disorder with at

least 64 known human variants, each with unique electrophysiological and epileptic

characteristics. A multi-disciplinary collaboration generated a novel mouse model (C57BL/6-

Kcnt1^{em1Bryd}) carrying the G269S variant, corresponding to human G288S, located within the

coding region of the channel pore. Network excitability of cultured cortical neurons from

Kcnt1^{+/G269S} exhibited sustained hyperexcitability and hypersynchronous bursting while

Kcnt1^{G269S/G269S} neurons showed early excessive bursting followed by network collapse,

suggesting excitotoxicity. *Kcnt1^{+/G269S}* displayed poor motor coordination, erratic breathing, and

increased apneas. Critically, *Kcnt1^{+/G269S}* were more susceptible to thermal-induced seizures in

early life. In summary, these data: (i) provide a novel mouse model of KCNT1-related epilepsy,

(ii) provide strong *in vitro* evidence of neuronal hyperexcitability, (iii) illustrate early-life seizures

as a functional outcome measure, and (iv) lay the groundwork for future analysis of neural activity

in vivo and modeling circuit level dynamics *in vitro* and *in silico*.

Significance Statement

Gain-of-function mutations in the sodium-gated potassium channel KCNT1 have been

linked to pediatric epilepsy of varying severity. The human *KCNT1* variant G288S (G269S in

mice) is linked to Autosomal Dominant Nocturnal Frontal Lobe Epilepsy (ADNFLE), Epilepsy

of Infancy with Migrating Focal Seizures (EIMFS), and other severe developmental epileptic

encephalopathies. There are currently no therapeutics to prevent the progression of *KCNT1*-

related epilepsy, therefore, the scientific community requires a novel mouse model that is well

characterized, *in vitro* and *in vivo,* to screen and assess targeted therapeutics. Herein, we

engineered a novel mouse to assess developmental and adult phenotypes resulting from the

G288S/G269S variant, *in vitro* and *in vivo*, to advance translation toward therapeutic testing for

individuals with *KCNT1*-related epilepsy.

Introduction

Genetic studies have shown that approximately one-third of neurodevelopmental disorders (NDDs) are associated with gene mutations affecting neuronal ion channels, emphasizing their critical role in brain pathology[1-3]. To understand how these mutations alter channel kinetics, *in vivo* experimental models are essential[4-7]. The *KCNT1* gene encodes a sodium activated potassium (K_{Na^+}) channel that regulates neuronal excitability by influencing the resting membrane potential and hyperpolarization following action potentials. Mutations in *KCNT1* result in diverse electrophysiological profiles, contributing to distinct epileptic syndromes such as Epilepsy of Infancy with Migrating Focal Seizures (EIMFS), Autosomal Dominant Nocturnal Frontal Lobe Epilepsy (ADNFLE), West syndrome, and Ohtahara syndrome[8]. To date, 64 *KCNT1* mutations have been identified, across 248 individuals[9,10], however, few *in vivo* mouse models exist, which has curbed therapeutic development.

Homozygous *Kcnt1*[-/-] mice exhibit poor motor ability and impaired motor learning, however, seizure related metrics were similar to wild-type (WT) control mice[11]. A mouse model carrying the R455H variant (human R474H) resulted in increased potassium currents in glutamatergic neurons and suppression of inhibitory neuron firing[12,13]. This occurs when dysfunctional channels activate rapidly in response to sodium influx during action potential upstroke, which can limit sodium channel inactivation and promote hyperexcitability. Video-EEG of *Kcnt1*[+/R455H] mice revealed persistent interictal spikes, spontaneous seizures, and a reduced threshold for pentylenetetrazol (PTZ)-induced seizures[11].

The homozygous P924L variant causes frequent debilitating seizures and developmental compromise[14]. Another experimental model, Y777H (human Y796H) exhibited sleep disturbances due to elevated seizure bouts and length[15]. Both *Kcnt1*[+/Y777H] and *Kcnt1*[Y777H/Y777H]

were susceptible to kainic acid (KA) and PTZ-induced seizures (Shore et al., 2020). Conditional

expression of Y777H had no effect on glutamatergic neurons but, increased subthreshold K_{Na^+}

currents in inhibitory neurons[16]. This cell type-specific modulation creates network imbalance,

functionally silencing certain inhibitory neurons while excitatory transmission remains intact or

enhanced[10].

Finally, the loss-of-function (LOF) variant, P932I, is associated with severe early infantile

epileptic encephalopathy (EE) and abnormal myelination[17]. Each *KCNT1* variant affects channel

kinetics differently—altering activation thresholds, open probability, and current amplitude[10].

These differences impact neuronal excitability and seizure susceptibility, varying considerably

across variants, neuronal subtypes, and background strain of the rodent model[18-20] adding

complexity to understanding circuit-level dysfunction in *KCNT1*-related epilepsies.

D*rosophila melanogaster* lack a *KCNT1* ortholog, allowing for the introduction of the

human *KCNT1* (*hKCNT1)* gene. While the WT *hKCNT1* gene showed no phenotype, expression

of GOF human variants exhibited seizure activity rates of 41% (G288S), 48% (R398Q), and 38%

(R928C), respectively[21,22]. Zhang's laboratory illustrates that *hKCNT1* G288S and R928C

mutations increased potassium conductance, impaired larval locomotion, hyperpolarized the

resting membrane potential, and reduced muscle resistance[21,22]. Consistent with these findings,

expression of the G288S variant in HEK cells exhibits: (i) a 21-fold increase in K_{Na^+} current

amplitude; (ii) an approximate 20% increase in channel open probability; (iii) a hyperpolarizing

shift in activation threshold, reducing voltage sensitivity; and (iv) minimal changes in channel

kinetics[10]. These data encouraged the generation of the *Kcnt1* GOF G288S (G269S in mice)

experimental model, termed B6-*Kcnt1*[em1Bryd]. This model enabled targeted characterization of

electrophysiological properties *in vitro* and functional outcomes *in vivo* which provide a valuable

platform for studying *KCNT1*-related epilepsy.

Materials and Methods.

Generation of the G288S Variant (G269S Mouse Model): All experimental procedures were

performed in accordance with the National Institutes of Health (NIH) Guide for the Care and Use

of Laboratory Animals and approved by the Institutional Animal Care and Use Committees

(IACUC) at the University of Missouri and University of California, Davis School of Medicine.

The novel *Kcnt1* mutant mouse model was generated by the MU Animal Modeling Core

(Columbia, MO) using a standard knock in method involving CRISPR/Cas9 genome editing. The

strategy for generating the *Kcnt1* exon 11 codon change was to replace the wild-type mouse *Kcnt1*

gene codon GGG with AGC creating a missense mutation. Glycine (G) is replaced by serine (S)

at amino acid position 269 in the mouse *Kcnt1* protein which mimics the G288S human variant.

C57BL/6J (stock #000664) animals were purchased from the Jackson Laboratory (Bar Harbor,

ME). Super ovulated females (4-5 weeks of age) were mated to stud males (~10 weeks of age)

overnight and zygotes were collected the following morning. sgRNAs were ordered as chemically

modified synthetic sgRNAs through Synthego (Redwood City, CA). The chemical modifications

on each sgRNA were 2′-O-methyl analogs and 3′ phosphonothioate internucleotide linkages at the

first three 5′ and 3′ terminal RNA residues. The repair template was chemically synthesized by

IDT (Coralville, IA) as a 200bp single-stranded DNA oligo (ssODN). The ssODN was

complementary to the non-target strand and contained symmetrical homology arms. An additional

silent mutation was introduced for genotyping purposes.

sgRNA sequence: 5′-TCCAGGTGCTGGATCCCAC-3′. Repair template sequence (sgRNA

sequence   disrupted   by   desired   mutation):   5′-

CTGCTGCAGCCTCTCCCCCACTGCTGCAGCCCCAGCCCCACTGCTGCAGCCCCAGCC

CCACTGCTGCAGCCCTGATCTTCCTGTTTCTAGGACCTGT**AG**CATCCAGCACCTGGA

GCGAGCAGGTGGCAACTTGAACCTGCTGACCTCCTTCTACTTCTGCATCGTGACTTTC

TCAACTGTGGGCTTCGGTGATGTGAC -3′. A mix containing a final concentration of 3.0μM

sgRNA, 2.0μM enhanced specificity Cas9 protein (Sigma) and 1.6μM ssODN was made

immediately prior to electroporation. CRISPR sgRNA/Cas9 RNP complexes were formed by

gently mixing the sgRNA and Cas9 protein together and incubating at room temperature for 10min.

After RNP formation, the ssODN repair template was added to the mixture. Using a NepaGene21

electroporator along with a 1.5-mm gap glass slide electrode, zygotes were electroporated under

the following conditions: Poring pulse: 40V, 3.5ms length, 50ms interval, 10% decay rate, positive

polarity (x4 pulses) Transfer pulse: 5V, 50ms length, 50ms interval, 40% decay rate, alternating

polarity (x5 pulses). Manipulated zygotes were then surgically transferred to pseudo-pregnant

surrogate CD1 females (8 weeks of age) purchased from Charles River Laboratories (Wilmington,

MA). Multiple animals carrying the desired single base pair change were identified by genotype

analysis and confirmed by nucleotide sequence analysis of the region around the base pair change.

The founders confirmed to carry the mutation were backcrossed to C57BL6/J for 2 generations

and then to C57BL6/NJ (Jackson laboratory #005304) for 2 generations to establish the strain

C57BL/6NJ *Kcnt1[em1Bryd]* which is available at the Mutant Mouse Resource and Research Center

at the university of Missouri (MMRRC#075727) and is available upon request at www.nmrrc.org.

In spring 2021, animals were transferred to UC Davis, School of Medicine (Sacramento, CA) to

establish a colony. This colony was maintained by backcrossing to C57BL/6J for >10 generations.

Genotyping: Genomic DNA was extracted by placing a small amount of tissue (ear or tail) in

100μl alkaline lysis reagent (25mM NaOH and 0. mM EDTA at a pH of 12) and heating at 95 ˚C

for 1hr. This was followed by addition of 100µl neutralization buffer (40mM Tris-HCl) with

vortexing for 5secs to mix. The <u>PCR primers were Kcnt1.F1</u>: 5'-cattccacactgcagccctgtctct-3'and

<u>Kcnt1.R1</u>: 5'-caagggtgacacagatcaggatgaccac- 3'. The 25µl reaction contained the following:

<u>12.5µl</u> EmeraldAmp® Supermix (Takara Bio USA Inc., San Jose, CA); <u>10.0µl</u> H$_2$O; <u>0.75µl</u>

Forward primer (10µM); <u>0.75µl</u> Reverse primer (10µM); <u>1µl</u> genomic DNA (~5-50ng). Thermal

cycler conditions were 1) 95 °C, 3mins, 2) 95 °C, 30secs, 3) 69 °C, 30secs, 4) 72 °C, 1min. Steps

2-4 were repeated 40 times, 5) 72°C, 7mins, 6) 4 °C, hold. The expected PCR amplicon = 403 bp**.**

Following PCR amplification, 2.8µl CutSmart buffer and 1.0µl *BtsC*I (20 units) (New England

Biolabs, Ipswich, MA) were added directly to the PCR reaction and incubated at 50°C for 2hrs.

Gel electrophoresis was performed with a 3% agarose gel in 1X TAE buffer at 80V for 1hr.

Expected restriction fragment sizes: wild type allele = 19bp, 23bp, 42bp, and 319bp; mutant allele

= 19bp, 23bp, 42bp, 113bp, 206bp. The 319bp fragment in the wild type allele is cut into 113bp

and 206bp fragments in the mutant allele.

<u>Cell Culture:</u> Test mice were generated by crossing male and female *Kcnt1$^{+/G269S}$* to yield pups of

three genotypes (*Kcnt1$^{+/+}$, Kcnt1$^{+/G269S}$*, and *Kcnt1$^{G269S/G269S}$*) and neonatal cortices (PND 0-2) were

collected. Primary cortical neuron-glia cultures were prepared as previously described[23,24]. Briefly,

cortices were minced with a razor blade prior to incubation in Hibernate A containing papain and

DNase. The tissue was then triturated with glass pipettes to dissociate cells from the extracellular

matrix (Bellco Glass, Vineland, NJ). Dissociated cells were counted and plated as described below.

<u>High Density Microelectrode Array Plating:</u> MaxWell Biosystems high density microelectrode

arrays (HD-MEAs) were used to assess electrophysiological activity of primary neuronal cultures,

as previously described[25,26]. MaxTwo 6-well HD-MEA plates were sterilized, filled with 2mL of

plating medium (Neurobasal Plus medium (Gibco, ThermoFisher Scientific), 2% B27 Plus

supplement (×50) (Gibco, ThermoFisher Scientific), 1x GlutaMAX supplement (Invitrogen), 10%

heat-inactivated horse serum (Invitrogen), and HEPES buffer (1M, pH 7.5) (Fisher Scientific)),

then incubated for 2 days at 37°C with 5% CO_2. On the day of cell plating, the plating media was

removed, and chips washed with sterile water, coated with primary coating (50μl of 0.07%

polyethyleneimine (PEI, Sigma-Aldrich)), and incubated for 1hr. The primary coating was

removed, the chips were washed 3 times with sterile water, and vacuum dried. Secary coating (50ul

Laminin (0.02 mg/mL; Sigma-Aldrich)) was applied and incubated (~ 1hr) until ready to plate

cells. The laminin coating was removed, 50uL of primary cortical cells were added to the HD-

MEAs, yielding approximately 125,000 cells per chip, and incubated for 1hr before the addition

of 2mL plating medium. 24 hours later, 50% of media was removed and replaced with cell culture

maintenance media (Dulbecco's Modified Eagle Medium (DMEM) supplemented with GlutaMax,

D-glucose and sodium pyruvate (Gibco, ThermoFisher Scientific) and 10% horse serum). Media

changes occurred twice weekly by removing 50% of the media and replacing it with an equal

volume of cell culture media. Neural activity was recorded twice a week as described below.

**HD-MEA Electrophysiological Recordings and Data Processing**: HD-MEA recordings were

performed between days _in vitro_ (DIV) 3 to 30 to capture the developmental trajectories of

dissociated neuronal cultures. Due to differences in recording schedules across experiments, some

DIVs varied slightly between litters (e.g., DIV 13 vs 14, DIV 27 vs. 28). Each session began with

the Activity Scan Assay, which recorded 30-sec segments across the entire chip under 'Sparse 7x'

configuration to assess overall activity of the chip. Based on spike amplitude and spatial

distribution, 1,020 electrodes were selected from regions exhibiting high activity (spike amplitude)

to maximize neuron coverage in the Network Scan Assay. Spontaneous activity was recorded for

300s at a 10 kHz sampling rate using the MaxTwo device (MaxWell Biosystems).

Data processing focused on two aspects: (1) The activity of the culture over time, and (2)

network-level dynamics reflecting population synchrony and communication. Raw extracellular

signals were bandpass filtered between 300Hz and 3,000Hz to isolate multi-unit activity (MUA).

Spike detection was performed using a threshold set at five times the standard deviation of baseline

noise. A recording channel was considered active if it exceeded a predefined minimum firing rate

and spike amplitude threshold. The mean firing rate and mean spike amplitude were computed

from all active channels to assess the overall culture health and neuronal proliferation. To

characterize network-level activity, detected spikes were transformed into spike raster plots and

binned to generate population firing rates. A Gaussian kernel was applied to these binned spike

trains to estimate a continuous measure of population activity, capturing burst dynamics and

synchronous events. Statistical comparisons between genotypes were conducted using repeated

measures ANOVA over time and one-way ANOVAs followed by tukey's post hoc tests for

selected group comparisons. To ensure analytical consistency, adjacent DIVs with ±1 day variation

were grouped during analysis. All analyses were conducted using custom MATLAB R2024 scripts

in combination with the MaxWell Biosystems Toolbox (v23.2).

Subjects for Developmental Milestones, Behavioral Assays, and Seizure Susceptibility: All mice

were housed in a temperature- and humidity-controlled vivarium maintained on a 12:12 light-dark

cycle and tested during the light cycle. Animals were fed a standard diet of Teklad global 18%

protein rodent diets 2918 (Envigo, Hayward, CA, USA). Wild-type (WT) and heterozygous

Kcnt1$^{+/G269S}$ offspring were labeled by paw tattoo over postnatal days (PND) 2-4 using non-toxic

animal tattoo ink (Ketchum Manufacturing Inc., Brockville, ON, Canada). Tail biopsies (1–2 mm)

from pups were taken for genotyping, following the UC Davis IACUC policy regarding tissue

collection. Mice were weaned at PND21, and group housed by sex with 2-4 animals per cage.

Assays were performed in order of least to most stressful and between the hours of 9:00 AM PST

and 4:00 PM PST during the light phase. All behavior testing was conducted by an experimenter

blinded to genotype and included both sexes. Mice were allowed to habituate in their home cages

in a dimly lit room adjacent to the testing room for 1hr prior to the start of testing to limit any

effect of transporting between the vivarium and testing suite. Between subjects, testing apparatus

surfaces were cleaned using 70% ethanol and allowed to dry.

Behavioral Assays: Four Cohorts of mice were used for a combination of developmental

milestones, seizure-related outcomes, behavioral phenotypes, and respiratory physiology, with full

details below. Cohort 1 was used to assess developmental milestones (PND 3, 5, 9, and 11,

Supplemental Table 1) with behavioral testing order: open field, rotarod, novel object

recognition, contextual and cued fear conditioning and, a two hit seizure assessment using low

stimulus electroconvulsive threshold (ECT, Ugo Basile, Gemonio (VA), Italy) and high dose

(lethal) chemoconvulsant seizure induction with pentylenetetrazol (PTZ) intraperitoneally (i.p.).

These were not counterbalanced for order of administration because the PTZ dose is lethal,

therefore it had to be the "sec hit." To assess epileptogenesis or epileptic progression, while

simultaneously characterizing changes in seizure susceptibility with respect to prior epileptic

events, animals were initially exposed to an electroconvulsive induced seizure to eliminate

duration and severity variability at lower chemo convulsant doses[27]. Cohort 2 comprised 4 litters

and Cohort 3 comprised 5 litters (a replicate Cohort of Cohort 2) were tested during a critical time

window (PND 21-23) for hyperthermia-induced seizures. Finally, to further investigate *KCNT1*'s

potential influence on respiratory physiology, Cohort 4 mice comprised of littermates from 8 litters

that underwent whole body plethysmography (WBP). Given that age-matched males are larger

than females in body weight and size, both contributing parameters that influence the volume of

air respirated during the plethysmograph diagnosis, Cohort 4 subjects were sex-separated to

eliminate sex-originating variables for WBP presentation and analysis.

(i) Open Field Locomotion Exploratory locomotion in a novel open field environment was assayed

in a room illuminated at ~40lux in a 40 cm×40 cm×30.5 cm arena as previously described[28-31].

Exploratory locomotion is an essential assay for gross motor abilities and physical activity over

time, identifying key motoric phenotypes, that utilize a conserved neural circuit from flies to

humans.

(ii) Rotarod Motor coordination, balance, and motor learning were assessed using an Ugo-Basile

accelerating rotarod (Stoelting Co, Wood Dale, IL) as previously published[32-34]. Mice are placed

on a rotating cylinder that slowly increases from 4 to 40 revolutions per min (rpm) over a 5min

trial, with 3 trials per day using a 60min inter-trial interval, and subsequently repeated over 3

consecutive days for a total of 9 trials. Performance was scored as latency to fall (sec) from the

cylinder with a maximum latency of 300sec.

(iii) Whole Body Plethysmography (WBP) Whole-body plethysmography (WBP) chambers

(Buxco, Data Sciences International, St. Paul, MN) were utilized to assess respiratory parameters,

such as inhalation/exhalation volumes, as well as the occurrence and time spent in apneas (an

absence of inhaling or exhaling).

(iv) Electroconvulsive Seizure Threshold (ECT) Low intensity (5mA with a 0.9 ms pulse width,

299 Hz, for 200 msec), electroconvulsive shock was administered via ear clamps to assess seizure

threshold via an ECT Racine Scale consisting of: (0) walks away, (1) brief stun and walks away,

(2) very brief jaw, forelimb clonus, lasting less than 1sec, (3) prolonged jaw, forelimb clonus,

ventral or dorsal neck flexion, loss of posture, lasting several secs, (4) tonic extension of forelimbs

(90 degree angle to torso), (5) tonic extension of hind limbs (90 degree angle to torso). The ECT

Racine Scale is unique in that animals typically experience a single seizure event resulting in a

single score and then recover over time.

(v) Seizure susceptibility by a Lethal Dose of the Chemoconvulsant Pentylenetetrazol (PTZ)

Dosing was conducted between 9am-2pm in a well-lit room and followed our validated dosing

regimens for C57BL/6J. Subjects were weighed and administered PTZ (70 mg/kg, i.p.), then

placed in a clean, empty cage with subsequent seizure stages live scored using a modified Racine

scale for a maximum of 30mins. Stages scored were: (1) first jerk/Straub's tail, (2) loss of righting

reflex, (3) generalized clonic-tonic seizure, (4) full tonic extension, (5) death. The PTZ Racine

Scale is unique in that animals typically progress through the stages over time beginning with 1

and ending at 5.

(vi) Hyperthermia-induced Seizure Susceptibility Postnatal day (PND) 21-23 subjects were

utilized to reflect the same approximate vulnerable developmental window as the human

population[35,36]. Subjects were habituated in an adjacent room for at least 30mins in low light (~30

lux). Subjects were given a brief anesthetic exposure (0.1ml 4% isoflurane for 30sec to 1min until

loss of consciousness) for insertion of rectal temperature probe (Physitemp RET-4, Clifton, NJ,

USA), prior to placement in the testing arena. A heat lamp placed above the arena was positioned

at a distance to maintain an internal body temperature of 37.5°C for a 10min habituation period.

Following arena habituation, the distance of the heat lamp was dynamically adjusted to avoid rapid

overheating and gradually raise the internal body temperature evenly by approximately 0.5°C

every 2mins. Testing continues until a behavioral seizure score of 3 or above is observed or until

the body's internal temperature reaches 42.5°C, in which case the heat lamp is immediately turned

off and distanced from the arena to allow the subject to cool to 40°C proceeded by removal. An

observer blind to genotype records the type of seizure using a modified Racine Scale (1. Staring,

2. Head nodding, 3. Unilateral forelimb clonus, 4. Bilateral forelimb clonus, 5. Rearing and falling,

6. Clonic seizure) previously described by[37]. The febrile Racine Scale is unique in that animals

typically progress through the stages over time and then recover once they are removed from the

heat source. Animals are returned to their home cage and provided hydrogel once they are active

and exhibit normal ambulatory behavior (~30mins).

Statistical Analysis Data were analyzed with GraphPad Prism by two-tailed Student's t-test for

single parameter comparison between the two genotypes or a One-Way Repeated Measures

ANOVA for comparison of genotype over time (e.g. development, open field, pre-and post-fear

conditioning). Significance levels were set at $p < 0.05$. Significant ANOVAs were followed by

Bonferroni-Dunn posthoc testing.

Results.

Mouse model generation. The *KCNT1* gene sequences differ slightly between human and mouse: the altered amino acid in the G288S human KCNT1 variant is at amino acid position 269 in the mouse protein (**Fig. 1**). Genome editing with CRISPR/Cas9 was used to make the mouse model: the WT mouse *Kcnt1* exon 11 codon (GGG) was replaced with AGC to create the missense mutation. This leads to glycine (G) being replaced by serine (S) at amino acid position 269 (G269S) in the mouse Kcnt1 protein this new allele (*Kcnt1^{em1Bryd}*) was nucleotide sequence verified. The mutation was generated on a C57BL/6J genetic background and was maintained by backcrossing heterozygous animals to B6/J and B6/NJ mice.

High-Density microelectrode array analysis reveals distinct network activity patterns in Kcnt1$^{+/G269S}$ and Kcnt1$^{G269S/G269S}$. To investigate the functional consequences of the *Kcnt1* G269S mutation at the network level, we employed high-density microelectrode array (HD-MEA) recordings of primary cortical neurons from wild-type (*Kcnt1$^{+/+}$*), heterozygous (*Kcnt1$^{+/G269S}$*), and homozygous (*Kcnt1$^{G269S/G269S}$*) mice. The Maxwell Biosystems HD-MEA platform featuring 26,400 electrodes with 1,024 simultaneous recording channels enabled detailed spatial and temporal mapping of neuronal network activity patterns (**Fig. 2A**). Extracellular signals were recorded from multiple channels allowing for precise detection of action potentials across the neuronal population (**Fig. 2B, C**) from DIV 13-21, spanning the critical network formation and early maturation phases. To ensure comparable recording conditions across genotypes, we quantified the percentage of active electrodes: all three genotypes showed similar electrode activation rates (approximately 70-75%, **Fig. 2D**), indicating that the G269S mutation did not significantly affect neuronal viability or density during initial plating and attachment. However, despite similar electrode coverage, mean firing rates (DIV13-21) differed significantly between

genotypes (**Fig. 2E;** $F_{(2,168)}$=13.21, p<0.0001) in a gene-dosage dependent manner: *Kcnt1*[G269S/G269S]

neurons had the highest mean firing rate (4.50±0.28 Hz), followed by *Kcnt1*[+/G269S] (3.56 ± 0.14Hz)

and *Kcnt1*[+/+] neurons (2.84±0.15 Hz). Tukey's post hoc tests revealed that *Kcnt1*[G269S/G269S] neurons

fired significantly more than *Kcnt1*[+/G269S] (p=0.001) and *Kcnt1*[+/+] neurons (p<0.0001); while

Kcnt1[+/G269S] neuronal firing rate did not reach significance from *Kcnt1*[+/+] (p=0.051).

Network development and bursting dynamics. Early network development corresponds to DIV

13 when synchronous activity begins to emerge and spread across the neuronal population[38,39],

while DIV 27 represents the mature network phase characterized by established bursting patterns

and stabilized connectivity[40-42]. At DIV 13, we observed distinctive firing patterns across the three

genotypes (**Fig. 3A-C** $F_{(2,52)}$=6.515, p=0.003). During this network formation phase,

Kcnt1[G269S/G269S] cultures displayed more frequent bursting activity compared to *Kcnt1*[+/+] (tukey's

post hoc p=0.002), with *Kcnt1*[+/G269S] networks showing intermediate burst frequency

(tukey$_{Kcnt1+/+vsKcnt1+/G269S}$ p=0.039; tukey $_{Kcnt1+/G269SvsKcnt1G269S/G269S}$ p=0.199). By DIV 27, the

differences in activity patterns became more pronounced but showed a different trajectory than at

DIV 13 (**Fig. 3D-F** $F_{(2,40)}$=14.60, p<0.0001). While *Kcnt1*[+/+] and *Kcnt1*[+/G269S] cultures maintained

consistent bursting patterns with increased synchronization characteristic of mature networks,

Kcnt1[G269S/G269S] cultures exhibited markedly diminished burst amplitude and altered burst patterns

(tukey$_{Kcnt1+/+vsKcnt1G269S/G269S}$ p=0.149; tukey$_{Kcnt1+/G269SvsKcnt1G269S/G269S}$ p<0.0001). This unexpected

reversal in the homozygous phenotype suggests potential network dysfunction emerging with

prolonged exposure to hyperexcitability, possibly through excitotoxic mechanisms.

Temporal evolution of network hyperexcitability. Quantitative analysis of burst frequency across

developmental timepoints (**Fig. 3G** $F_{TIME(2,109)}$=7.516, p=0.0003; $F_{GENOTYPE(2,52)}$=8.027, p=0.0009)

revealed a striking biphasic pattern in *Kcnt1*[G269S/G269S] cultures. Between DIV 13-17,

Kcnt1$^{G269S/G269S}$ networks exhibited significantly higher burst counts (107.9± 7.21) per recording,

compared to *Kcnt1$^{+/+}$* (62.00 ± 7.41, tukey's p=0.002) reflecting initial hyperexcitability during

network formation. However, bursting activity declined in *Kcnt1$^{G269S/G269S}$* networks (87.19± 7.31)

with frequency falling below *Kcnt1$^{+/+}$* levels (108.4± 5.34) by DIV 24-27 ($F_{(2,40)}$=9.77, p=0.0004;

tukey's p=0.162). In contrast, *Kcnt1$^{+/G269S}$* cultures increased burst frequency over development,

showing significantly elevated activity (95.15±4.74; tukey's p=0.0003) at DIV13-17 compared to

Kcnt1$^{+/+}$ and rising to *Kcnt1$^{+/+}$* levels (129.40±6.91 tukey's p=0.121) by DIV 24-27. Meanwhile,

Kcnt1$^{+/G269S}$ cultures burst frequency was below *Kcnt1$^{G269S/G269S}$* networks at DIV 13-17 (tukey's

p=0.275) and rose above their levels by DIV 24-27 (tukey's p=0.0002). This indicates that the

Kcnt1$^{+/G269S}$ confers a persistent and sustainable hyperexcitability phenotype that remains

compatible with ongoing network function, unlike *Kcnt1$^{G269S/G269S}$*.

Further analysis of the firing pattern variability was performed, using the Fano factor, the

variance-to-mean ratio of spike counts, where a higher Fano factor reflects the healthy stochastic

variability essential for proper circuit formation and refinement[43]. *Kcnt1$^{+/+}$* networks maintained

high Fano factor values (**Fig. 3H,** 656.31 ± 71.94; DIV 13 $F_{(2,51)}$=7.057, p=0.002; DIV 27

$F_{(2,39)}$=12.36, p<0.0001). In contrast, both *Kcnt1$^{G269S/G269S}$* (303.11 ± 32.09; tukey's p=0.004;) and

Kcnt1$^{+/G269S}$ networks (326.67 ± 28.08; tukey's p=0.003) showed significantly lower values during

early network formation DIV 13. While the Fano factor of *Kcnt1$^{+/G269S}$* networks gradually

increased toward *Kcnt1$^{+/+}$* levels by DIV 27 (tukey's p=0.253), *Kcnt1$^{G269S/G269S}$* networks remained

locked in an abnormally regularized firing pattern (tukey's p<0.001). This developmental

trajectory—initial hyperexcitability followed by progressive reduction in network activity in

Kcnt1$^{G269S/G269S}$ cultures, contrasted with sustained hyperexcitability in *Kcnt1$^{+/G269S}$* cultures—

suggests that the G269S GOF mutation affects network development in a gene-dosage dependent

manner. Based on these electrophysiological findings, we decided to pursue only the $Kcnt1^{+/G269S}$

model for subsequent *in vivo* behavioral characterization.

Exploratory locomotion and motor coordination in the G288S $Kcnt1^{+/G269S}$ mouse model. Novel

exploration in the open field was assessed between age, sex matched littermate controls, for effects

resulting from $Kcnt1^{+/G269S}$. Total activity across a 30min session illustrated an effect over time in

both genotypes (**Fig. 4A**; $F_{TIME(3,148)}=7.706$, p<0.0001), showing acclimation to the arena, as

expected. A trend toward hyperactivity in the $Kcnt1^{+/G269S}$ mice was observed (**Fig. 4A**;

$F_{(1,42)}=0.6074$, p=0.4404) when the 30min session was summed (**Fig. 4B**; $t_{(1,40)}=2.156$, p=0.0372).

Motor coordination and motor learning were measured using the accelerating rotarod. We

observed a significant reduction in latencies to fall in $Kcnt1^{+/G269S}$ compared to $Kcnt1^{+/+}$ age

matched mice (**Fig. 4C**; $F_{GENOTYPE(1,42)}=6.79$, p=0.0126) indicating poor motor coordination.

However, over the three training days we observed improved performance over time, (**Fig. 4C**;

$F_{TIME(2,84)}=36.05$, p<0.0001), in both genotypes illustrating intact motor learning.

Cognitive ability assessed by two standardized assays in the $Kcnt1^{+/G288S}$ mouse model is intact.

Cognitive ability was assessed in two different tasks novel object recognition (NOR) memory and

fear conditioning. During the familiarization phase of the NOR task, $Kcnt1^{+/G269S}$ mice explored

the objects substantially less than the $Kcnt1^{+/+}$ control mice (**Fig. S1A**: $F_{FAMILIARIZATION}$

$_{GENOTYPE(1,29)}=0.00306$, p<0.0001), with a mean investigation time in $Kcnt1^{+/+}$ subjects of

52.285sec, while in $Kcnt1^{+/G269S}$ the mean investigation time was 26.865sec. During the novel

investigation phase, both genotypes spent greater time investigating the novel object, versus the

familiar object (**Fig. S1B**: $F_{Kcnt1+/+ (1,41)}=4.813$, p<0.05 and $F_{Kcnt1+/G269S(1,36)}=18.50$, p<0.001). The

discrimination index was calculated by dividing the difference in investigation time between the

novel and familiar object by the total amount of exploration of both objects (**Fig. S1C**:

t$_{(1,40)}$=0.7566), clearly illustrating a lack of cognitive impairments in this assay. Cued and contextual fear conditioning illustrated a strong effect of training (learning from the tone-shock pairings (UCS-CS) (**Fig. S1D**; F$_{TRAINING(1,40)}$=107.8, p<0.0001). While a genotype by training interaction was observed, (**Fig. S1D**; F$_{TRAININGvsGENOTYPE(1,40)}$=4.951, p<0.04), no genotype effect, during training, was observed, (**Fig. S1D**; F$_{GENOTYPE(1, 40)}$=0.7672, p=0.3863), which is a key control metric, since no genotype effect across training allows for the investigation of differences in memory and recall. Contextual recall 24 hrs. after UCS-CS training did not differ between genotypes (**Fig. S1E**; t$_{(1,50)}$=0.4610, p=0.6468). However, cued recall, performed 48hrs after initial training, illustrated recall of the cued-shock pairing by high levels of freezing illustrated post cue but not pre cue in *Kcnt1$^{+/G269S}$* compared to *Kcnt1$^{+/+}$* age matched mice (**Fig. S1F**; F$_{CUE(1,40)}$=11.05, p<0.0001).

Behavioral seizure quantification, in the Kcnt1$^{+/G269S}$ mouse model by multiple methods. Firstly, susceptibility to seizures early in life (post-natal day 21-23) was induced by hyperthermia, analogous to fever induced (febrile) seizures in Cohort 2 animals. Susceptibility was measured by probability to develop behavioral seizures, temperature to first seizure, and severity of seizure by a modified Febrile Racine score. *Kcnt1$^{+/G269S}$* animals had a higher average Febrile Racine score than *Kcnt1$^{+/+}$* controls (**Fig. 5A**: t$_{(1,31)}$=2.146, p<0.05) and a lower temperature threshold (**Fig. 5B**: t$_{(1,16)}$=2.849, df=16, p<0.01). *Kcnt1$^{+/G269S}$* animals also had an increased probability of developing a behavioral seizure (**Fig. 5C**: χ2=7.506, p<0.0005), over their *Kcnt1$^{+/+}$* counterparts. Importantly, a replicate Cohort by a separate investigator produced similar results (**Fig. S2A-C**). Following behavioral assay completion, Cohort 1 mice were measured for epileptic progression after induction with electro- and chemoconvulsant seizures. Seizures in Cohort 1 animals were first induced by ear-clip electroconvulsive threshold (ECT) wherein animals reached similar seizure

levels (**Fig. 5D**: $t_{(1, 40)}$=0.17, p=0.86) and recovered post-seizure induction in less than 5 minutes

(**Fig. 5E**: $t_{(1,37)}$=1.28, p=2.08). Both *Kcnt1$^{+/G269S}$* and *Kcnt1$^{+/+}$* animals had similar probability of

recovery post-seizure experience (**Fig. 5F**: $\chi2$=7.506, p=0.65). Following PTZ administration at

70 mg/kg, a lethal dose, *Kcnt1$^{+/G269S}$* and *Kcnt1$^{+/+}$* mice reached each seizure stage at similar rates

(**Fig. 5G**: $F_{(1,37)}$=0.01, p=0.92), including latency to first jerk, latency to loss of righting, latency

to generalized clonic-tonic seizure, latency to full extension, and latency to death. Additionally,

the probability of survival was similar between *Kcnt1$^{+/G269S}$* and *Kcnt1$^{+/+}$* animals (**Fig. 3H**:

$F_{(1,12)}$=0.499, p=0.49). Taken together, the *Kcnt1$^{+/G269S}$* animals were highly sensitive to early life

heat-induced seizures, as opposed to the adult exposure to either electroconvulsive or

chemoconvulsive induced seizures. Since the ECT and PTZ seizure inductions were conducted in

the same animals we correlated the seizure severity between the two induction methods, in order

to determine if one would be predictive of the other, i.e., correlation (**Fig. S3**). Latency to first jerk

after PTZ administration was negatively correlated with seizure severity in ECT in *Kcnt1$^{+/+}$*

animals (**Fig. S3A** r=-0.46, p=0.05). While latency to generalized tonic-clonic PTZ seizures were

negatively correlated with seizure severity after ECT in *Kcnt1$^{+/G269S}$* (**Fig S3G** r=-0.57, p=0.02).

Together these correlations indicate that mice that experienced the most intense responses to ECT,

also moved more quickly through the Racine scale stages during PTZ induced seizures.

Respiratory parameters identified erratic breathing and apnea in the Kcnt1$^{+/G288S}$ mouse model.

Our laboratory was interested in defining respiratory metrics at baseline, given KCNT1-related

epilepsy's increased incidence of Sudden Unexpected Death in Epilepsy (SUDEP) and elevated

apneas[44-46]. Animals were assessed for 2hrs during the light cycle for respiratory parameters in the

whole-body plethysmography (WBP) chambers. *Kcnt1$^{+/G269S}$* and *Kcnt1$^{+/+}$* breathing

characteristics were quantified, including indices such as inspiration and expiration volumes, as

well as the presence of apneas. Animals were separated by sex as aged matched female mice are

smaller than their male counterparts resulting in inherent sex differences for these measures.

Percentage of time spent breathing normal was similar between $Kcnt1^{+/G269S}$ and $Kcnt1^{+/+}$ females

(**Fig. 6A**: $t_{(1,18)}=0.4896$, p=0.6297), whereas males exhibited decreased normal breathing

($t_{(1,17)}=2.391$, p=0.0258). Additionally, males and females of both genotypes had similar levels of

erratic breathing (**Fig. 6B**: females $t_{(1,18)}=1.675$, p=0.1094; males $t_{(1,17)}=0.7847$, p=0.4410). Lastly,

the number of apnea events was elevated in $Kcnt1^{+/G269}$ mice in males (**Fig. 6C**: $t_{(1,17)}$,t=2.391,

p=0.0258) and not in females ($t_{(1,18)}$,t=0.4896, p=0.6297), compared to their respective sex matched

$Kcnt1^{+/+}$ controls. Taken together, this data may indicate that male $Kcnt1^{+/G269S}$ mice experience

more respiratory difficulties than age matched females.

Discussion.

Herein, we generated a novel experimental model for *KCNT1*-related epilepsy and channelopathies, a subgroup of neurodevelopmental disorders. This model mouse carries the mouse equivalent (G269S) of the human GOF variant G288S. CRISPR/Cas9 was used in C57BL/6J mice to replace the WT gene codon GGG with AGC, creating the missense mutation that substitutes glycine (G) to serine (S) at amino acid position 269 in the mouse Kcnt1 protein. The new allele (*Kcnt1em1Bryd*) was nucleotide sequence verified, and the mice were maintained on C57BL6/NJ and were donated to the Mutant Mouse Resource and Research Center (MMRRC:075727-MU). Although the initial model was maintained on a C57BL/6NJ background, we backcrossed it onto C57BL/6J to align with existing models of developmental and epileptic encephalopathy (DEE) and rare genetic neurodevelopmental disorders (NDDs). However, we acknowledge the limitations of the C57BL/6J strain in seizure studies, as documented in prior literature[18-20,47-49]. Given the reported differences in seizure susceptibility and responsiveness between sub strains, future experiments may return to the C57BL/6NJ background to better understand how genetic background influences phenotype, particularly of spontaneous recurrent seizures (SRS), seizure induction and potential of therapeutic testing[18,50]. Such comparisons are essential for evaluating the generalizability of findings and optimizing translational relevance[51].

We performed functional characterization of primary cortical neurons from *Kcnt1*[+/+], *Kcnt1*[+/G269S], and *Kcnt1*[G269S/G269S] mice *in vitro* revealing gene dose-dependent differences in excitability. Homozygous *Kcnt1*[G269S/G269S] cultures showed an initial phase of hyperexcitability followed by network collapse, while heterozygous *Kcnt1*[+/G269S] cultures maintained sustained hyperexcitability. Strikingly, both mutant genotypes exhibited a ~50% reduction in firing pattern variability (Fano factor: *Kcnt1*[+/+] ≈ 656 vs. mutants ≈ 300–327), indicating abnormally regularized

activity. This loss of stochastic variability likely disrupts normal circuit development [43]. These

findings are consistent with known cell type-specific effects of KCNT1 gain-of-function (GOF)

variants, which disproportionately impair inhibitory neurons—particularly non-fast-spiking

GABAergic interneurons[13,52]. The G269S mutation, located in the channel pore region, appears

especially severe: homozygous cultures lost bursting activity by DIV 28, suggesting excitotoxicity

due to excessive firing. In contrast, the heterozygous model retains viable hyperexcitability, more

accurately reflecting the human condition and offering a robust platform for therapeutic

development. These findings suggest that approaches targeting improved inhibitory tone and

excitation-inhibition balance, rather than KCNT1 channel blockade alone, may prove more

effective for treating *KCNT1*-related epilepsies[53-55].

Next, we discovered that *Kcnt1*$^{+/G269S}$ mice exhibited seizures following early life

hyperthermia, a preclinical assay utilized as an analog for febrile seizures or seizures resulting

from high temperature fevers[37]. Compared to age-matched control littermates, *Kcnt1*$^{+/G269S}$ mice

experienced seizure onset at lower temperatures, had a higher incidence of seizures, and scored

higher on a modified febrile Racine scale, indicating increased severity across two Cohorts. In

contrast, adult *Kcnt1*$^{+/G269S}$ mice that were subjected to a 2-hit seizure paradigm—induction by

electroconvulsive threshold (ECT) followed by a high dose of pentylenetetrazol (PTZ, 70

mg/kg)—were indistinguishable from *Kcnt1*$^{+/+}$ controls, on every metric. Finally, a tailored

behavioral phenotyping battery was performed on a seizure naïve Cohort of *Kcnt1*$^{+/G269S}$ and

Kcnt1$^{+/+}$ littermates that included: developmental milestones, exploratory locomotion, motor

coordination and motor learning, novel object learning and memory, and contextual and cued

learning and memory. Additionally, we evaluated respiratory physiology using whole-body

plethysmography (WBP), which was included given the clinical reports of asthma and pulmonary

hemorrhage in *KCNT1*-related epilepsy[56-59], as well as the presence of apneas leading to disrupted sleep as a common occurrence in genetic NDDs, particularly when facial dysmorphic traits are evident[60-62]. While *Kcnt1*$^{+/G269S}$ and *Kcnt1*$^{+/+}$ littermates were developmentally indistinguishable, *Kcnt1*$^{+/G269S}$ mice were hyperactive in the 30min open field arena and lacked motor coordination on days 1 and 2 on the rotarod. Finally, *Kcnt1*$^{+/G269S}$ mice exhibited increased apneas and erratic breathing which align with the clinical presentation.

In summary, we report for the first time a novel Kcnt1 G269S knock-in mouse model, backcrossed onto the C57BL/6J background. This model represents a significant advancement in the study of *KCNT1*-related epilepsy, developmental and epileptic encephalopathies (DEEs), and neurodevelopmental channelopathies. We hypothesize that early-life seizures, as observed in this model, are driving phenotypes that closely align with clinical features of *KCNT1*-related epilepsy, including motor hyperactivity, learning and memory impairments, apneas, spontaneous seizures, and sleep disturbances. Notably, homozygous *Kcnt1*$^{G269S/G269S}$ mice are not construct valid, thus reducing the relevance, required for translational research, and making this heterozygous model ideal for both *in vitro* and *in vivo* therapeutic testing.

Limited clinical success of KCNT1 channel blockers such as quinidine has been well documented[53,54,63-65]. This work illustrates that future therapeutic strategies may be more effective if they focus on: (i) restoring inhibitory tone; (ii) reducing potassium efflux; and (iii) correcting excitation-inhibition (E/I) imbalance. Future studies will incorporate long-term wireless EEG to assess cortical activity, including spiking, spike trains, sleep architecture, and spectral power density over multiple days. Cortical EEG is a critical translational biomarker and is routinely used by our lab and others[26,37,66-68], especially given the nuance of the C57BL6/J background strain[69]. Moreover, we will report sensory evoked potentials, another common translational biomarker.

Additionally, our adult ECT experiments were conducted using ear-clip stimulation at low

intensity. Based on recommendations from the Frankel lab (personal communication), we have

since modified our ECT apparatus to allow for transcorneal stimulation at higher intensities. This

implementation will be reported in future studies, as data is currently being acquired. These

analyses have the ability to illustrate the severity of this GOF variant and enable deeper *in vivo*

phenotyping. These enhancements will contribute to a more comprehensive understanding of

KCNT1 variant effects and support improved patient stratification.

The G288S mutation was originally identified via whole-exome sequencing in two

unrelated patients with early-onset DEE[65]. As previously noted, this variant does not correspond

to a single epilepsy subtype[70], highlighting the importance of understanding how G288S/G269S

modulates network excitability, insights will inform therapeutic development, and might yield

information relevant to the most affected individuals and inform stratification. Groundwork in

KCNT1-related epilepsy was outlined by many as summarized in **Table 1**, but we must highlight

generous contributions and personal communication with the Wayne Frankel laboratory

(Columbia University) who generated the first ever mouse models, the knock-in (KI), human

variants Y777H and Y796H and illustrated motor cortex hyperexcitability, early-onset seizures,

and hyperactivity in KI/KI mice[16,52]. In 2024, Wu et al. discovered the R455H mutation which

exhibited increased sodium conductance, increased excitatory neuron activity, and suppression of

excitability in inhibitory interneurons in a mouse model, which altered sodium channel expression,

with effect sizes of 4-6-fold increases in KI/KI mice. While previous findings in homozygous

variant mice are significant, with substantial effect sizes to test for therapeutic reversal, there

remains an unmet need for a more construct valid single copy KI variant model, since affected

individuals with *KCNT1*-related epilepsy are heterozygous. The model presented herein has filled

561 this unmet need by its generation and clear phenotypic excitability in cortical neurons and early

562 life heat seizure sensitivity as an *in vivo* outcome measure, malleable for future therapeutic testing,

563 making it a powerful tool for translation and platforms of future therapeutic discovery.

564 **Conclusions**

565 Overall, characterization of $Kcnt1^{+/G269S}$ mice indicates a sensitivity to hyperthermia-induced

566 seizures and a motor learning delay. Future studies are required to investigate the impact of early

567 life seizures on adult motor, cognitive, and sleep measures as well as EEG before developing novel

568 therapeutic interventions for individuals with *KCNT1*-related epilepsy.

570
Figure 1. Generation of the *Kcnt1*^{G269S} mutant mouse model.

A) The KCNT1 protein consists of six hydrophobic transmembrane segments with a pore loop
domain between the last two transmembrane segments. It also has a large intracellular carboxy-
terminal region containing tandem RCK domains and an NAD+ binding domain. The mouse
G269S (corresponding to human G288S) mutation, indicated by a red star, occurs within the
channel pore. **B)** The mouse model was engineered using CRISPR-Cas9 genome editing to replace
the endogenous GGG (Gly) with AGC (Ser). **C)** Confirmation of the engineered DNA base pair
changes using Sanger sequencing. The heterozygous sample in the chromatogram shows
overlapping G (black) and A (green) peaks for the G to A change and overlapping G (black) and
C (blue) peaks for the G to C change.

582
Fig. 2. High-density MEA recordings reveal genotype-dependent differences in neuronal
network activity.
(A) Image of the high-density microelectrode array (HD-MEA) chip used in this study (26,000
electrodes; MaxOne platform, Maxwell Biosystems). (B) Example of raw extracellular action
potential (eAP) traces recorded from active electrodes. (C) Representative firing rate map from a
cortical culture, showing spatial distribution of neuronal activity across the MEA surface.
Warmer colors indicate higher firing rates (color scale: 2–12 Hz). (D) Percentage of active
electrodes per well for wild-type *Kcnt1*[+/+], heterozygous *Kcnt1*[+/G269S], and homozygous
Kcnt1[G269S/G269S,] neuronal networks. (E) Quantification of mean firing rate (Hz) across genotypes.
Kcnt1[G269S/G269S] neurons fired significantly more than both *Kcnt1*[+/G269S] (t = −3.00, p = 0.0036,
) and *Kcnt1*[+/+] (t = 5.27, p <0.0001, *), while *Kcnt1*[+/G269S] neurons also differed significantly
from *Kcnt1*[+/+] (t = 3.51, p = 0.0007, ***). Bars represent mean ± SEM; individual points reflect
biological replicates.

596
Fig. 3. Developmental progression of burst dynamics in *Kcnt1^+/+*, *Kcnt1^+/G269S*, and
***Kcnt1^G269S/G269S* in primary neuron culture.**
(A–F) Representative raster plots (top) and population firing rate traces (bottom) from wild-type
(*Kcnt1^+/+*, A, D), heterozygous (*Kcnt1^+/G269S*, B, E), and homozygous (*Kcnt1^G269S/G269S*, C,F)
cortical networks recorded at DIV13 (A–C) and DIV27 (D–F). Each raster shows 120secs of
activity across 1024 electrodes. Firing rate traces display synchronized burst events detected using
a Gaussian kernel convolution; red dots indicate burst peaks. *Kcnt1^+/+* and *Kcnt1^+/G269S* networks
display sparse, periodic bursts at DIV13 that become more frequent and structured by DIV27. In

contrast, $Kcnt1^{G269S/G269S}$ networks show highly frequent and irregular bursting at DIV13, followed
by diminished activity and synchrony at DIV27. (G) Quantification of burst count across
developmental timepoints (DIV13–27). $Kcnt1^{G269S/G269S}$ networks display a biphasic pattern with
early hyperexcitability (DIV13–17) and later hypoactivity, whereas $Kcnt1^{+/G269S}$ cultures maintain
and increase burst frequency over the developmental time. (H) Fano factor, a measure of firing
variability, is significantly reduced in both mutant genotypes compared to $Kcnt1^{+/+}$ at all
timepoints, indicating abnormally regularized activity. Bars represent mean ± SEM; individual
data points represent biological replicates. Error bars indicate mean +/- SEM. $p < 0.05$ (*), $p <$
0.01 (**), $p < 0.001$ (***).
614

615
Fig. 4. Elevated motor activity in a novel open field arena using the total activity parameter
over a 30min period in *Kcnt1*^{+/G269S} vs *Kcnt1*^{+/+}. Gross motor abilities were assessed using
activity in a novel open field arena by: A, B) total activity across a 30min session. In Panel A, data
are shown in 5min bins. C) *Kcnt1*^{+/G269S} have poor motor coordination compared to *Kcnt1*^{+/+}
controls using the standard accelerating rotarod paradigm. Error bars indicate mean +/- SEM. *
p<0.05
622

Figure 5: Seizure responsiveness to Hyperthermia-, Electroconvulsive-, and Chemoconvulsive-induction.

A) Seizure severity using modified Febrile Racine scale to Hyperthermia-induced seizure. B) Average temperature to behavioral seizure. C) Probability of behavioral seizure as temperature is increased. D) Seizure severity using a modified ECT Racine scale in response to electroconvulsive shock threshold (ECT; 5mA with a 0.9ms pulse width, 299 Hz, for 200 msec). E) Time to recovery (sec) post ECT induction. F) Probability of recovery post ECT. G) Latency to each stage of a modified Racine scale in response to chemoconvulsive treatment with pentylenetetrazole (PTZ, i.p., 70 mg/kg). H) Probability of survival post PTZ. Error bars indicate mean +/- SEM. * p<0.05.

634
635 **Fig. 6: Respiratory apneas and erratic breathing assessed by whole body plethysmography**
636 **(WBP) in *Kcnt1*[+/G269S] and *Kcnt1* [+/+] mice.** Whole body plethysmography evaluation for A) % of
637 time breathing normal, B) % of Time breathing erratic, C) number of apneas. Error bars indicate
638 mean +/- SEM. * $p < 0.05$
639

References

640
641
1   Satterstrom, F. K. *et al.* Large-Scale Exome Sequencing Study Implicates Both Developmental
and Functional Changes in the Neurobiology of Autism. *Cell* **180**, 568-584 e523 (2020).
https://doi.org/10.1016/j.cell.2019.12.036
2   Noebels, J. Precision physiology and rescue of brain ion channel disorders. *J Gen Physiol* **149**,
533-546 (2017). https://doi.org/10.1085/jgp.201711759
3   Daghsni, M. *et al.* Autism throughout genetics: Perusal of the implication of ion channels. *Brain*
Behav **8**, e00978 (2018). https://doi.org/10.1002/brb3.978
4   Sanders, S. J. *et al.* Progress in Understanding and Treating SCN2A-Mediated Disorders. *Trends*
Neurosci **41**, 442-456 (2018). https://doi.org/10.1016/j.tins.2018.03.011
5   Smith, R. S. & Walsh, C. A. Ion Channel Functions in Early Brain Development. *Trends Neurosci*
43, 103-114 (2020). https://doi.org/10.1016/j.tins.2019.12.004
6   Niemi, M. E. K. *et al.* Common genetic variants contribute to risk of rare severe
neurodevelopmental disorders. *Nature* **562**, 268-271 (2018). https://doi.org/10.1038/s41586-
018-0566-4
7   Almog, Y. *et al.* Developmental alterations in firing properties of hippocampal CA1 inhibitory and
excitatory neurons in a mouse model of Dravet syndrome. *Neurobiol Dis* **148**, 105209 (2021).
https://doi.org/10.1016/j.nbd.2020.105209
8   Hinckley, C. A. *et al.* Functional evaluation of epilepsy-associated KCNT1 variants in multiple
cellular systems reveals a predominant gain of function impact on channel properties. *Epilepsia*
64, 2126-2136 (2023). https://doi.org/10.1111/epi.17648
9   Bonardi, C. M. *et al.* KCNT1-related epilepsies and epileptic encephalopathies: phenotypic and
mutational spectrum. *Brain* **144**, 3635-3650 (2021). https://doi.org/10.1093/brain/awab219
10   Rychkov, G. Y. *et al.* Functional Effects of Epilepsy Associated KCNT1 Mutations Suggest
Pathogenesis via Aberrant Inhibitory Neuronal Activity. *Int J Mol Sci* **23** (2022).
https://doi.org/10.3390/ijms232315133
11   Quraishi, I. H. *et al.* An Epilepsy-Associated KCNT1 Mutation Enhances Excitability of Human
iPSC-Derived Neurons by Increasing Slack K(Na) Currents. *J Neurosci* **39**, 7438-7449 (2019).
https://doi.org/10.1523/JNEUROSCI.1628-18.2019
12   Lim, C. X., Ricos, M. G., Dibbens, L. M. & Heron, S. E. KCNT1 mutations in seizure disorders: the
phenotypic spectrum and functional effects. *J Med Genet* **53**, 217-225 (2016).
https://doi.org/10.1136/jmedgenet-2015-103508
13   Wu, J., Quraishi, I. H., Zhang, Y., Bromwich, M. & Kaczmarek, L. K. Disease-causing Slack
potassium channel mutations produce opposite effects on excitability of excitatory and
inhibitory neurons. *Cell Rep* **43**, 113904 (2024). https://doi.org/10.1016/j.celrep.2024.113904
14   Burbano, L. E. *et al.* Antisense oligonucleotide therapy for KCNT1 encephalopathy. *JCI Insight* **7**
(2022). https://doi.org/10.1172/jci.insight.146090
15   Wang, R., Teng, S., Turanchik, M., Zhen, F. & Peng, Y. Tonic-clonic seizures induce hypersomnia
and suppress rapid eye movement sleep in mouse models of epilepsy. *Sleep Adv* **6**, zpaf009
(2025). https://doi.org/10.1093/sleepadvances/zpaf009
16   Shore, A. N. *et al.* Heterozygous expression of a Kcnt1 gain-of-function variant has differential
effects on SST- and PV-expressing cortical GABAergic neurons. *bioRxiv* (2024).
https://doi.org/10.1101/2023.10.11.561953
17   Vanderver, A. *et al.* Identification of a novel de novo p.Phe932Ile KCNT1 mutation in a patient
with leukoencephalopathy and severe epilepsy. *Pediatr Neurol* **50**, 112-114 (2014).
https://doi.org/10.1016/j.pediatrneurol.2013.06.024

| 687 | 18 | Copping, N. A., Adhikari, A., Petkova, S. P. & Silverman, J. L. Genetic backgrounds have unique seizure response profiles and behavioral outcomes following convulsant administration. *Epilepsy Behav* **101**, 106547 (2019). https://doi.org/10.1016/j.yebeh.2019.106547 |

18   Copping, N. A., Adhikari, A., Petkova, S. P. & Silverman, J. L. Genetic backgrounds have unique
seizure response profiles and behavioral outcomes following convulsant administration. *Epilepsy*
Behav **101**, 106547 (2019). https://doi.org/10.1016/j.yebeh.2019.106547

19   Miller, A. R., Hawkins, N. A., McCollom, C. E. & Kearney, J. A. Mapping genetic modifiers of
survival in a mouse model of Dravet syndrome. *Genes Brain Behav* **13**, 163-172 (2014).
https://doi.org/10.1111/gbb.12099

20   Yang, Y. & Frankel, W. N. Genetic approaches to studying mouse models of human seizure
disorders. *Adv Exp Med Biol* **548**, 1-11 (2004). https://doi.org/10.1007/978-1-4757-6376-8_1

21   Zhang, Q., Liu, Y., Xu, J., Teng, Y. & Zhang, Z. The Functional Properties, Physiological Roles,
Channelopathy and Pharmacological Characteristics of the Slack (KCNT1) Channel. *Adv Exp Med*
Biol **1349**, 387-400 (2021). https://doi.org/10.1007/978-981-16-4254-8_18

22   Ehaideb, S. *et al.* Allostatic Compensatory Mechanisms Enable the Mutant KCNT1 Channels to
Induce Seizures and Hyperexcitability. *iScience* (2018). https://doi.org/
http://dx.doi.org/10.2139/ssrn.3188397

23   Seiler, S. *et al.* Astrocyte-derived factors regulate CNS myelination. *Glia* **72**, 2038-2060 (2024).
https://doi.org/10.1002/glia.24596

24   Hanes, A. L. *et al.* Divergent Synaptic Scaling of Miniature EPSCs following Activity Blockade in
Dissociated Neuronal Cultures. *J Neurosci* **40**, 4090-4102 (2020).
https://doi.org/10.1523/JNEUROSCI.1393-19.2020

25   Ben-Shalom, R., Aviv, A., Razon, B. & Korngreen, A. Optimizing ion channel models using a
parallel genetic algorithm on graphical processors. *J Neurosci Methods* **206**, 183-194 (2012).
https://doi.org/10.1016/j.jneumeth.2012.02.024

26   Fenton, T. A. *et al.* Hyperexcitability and translational phenotypes in a preclinical mouse model
of SYNGAP1-related intellectual disability. *Transl Psychiatry* **14**, 405 (2024).
https://doi.org/10.1038/s41398-024-03077-6

27   Sharma, S., Puttachary, S., Thippeswamy, A., Kanthasamy, A. G. & Thippeswamy, T. Status
Epilepticus: Behavioral and Electroencephalography Seizure Correlates in Kainate Experimental
Models. *Front Neurol* **9**, 7 (2018). https://doi.org/10.3389/fneur.2018.00007

28   Adhikari, A. *et al.* Touchscreen cognitive deficits, hyperexcitability and hyperactivity in males and
females using two models of Cdkl5 deficiency. *Hum Mol Genet* **31**, 3032-3050 (2022).
https://doi.org/10.1093/hmg/ddac091

29   Copping, N. A. *et al.* Neuronal overexpression of Ube3a isoform 2 causes behavioral
impairments and neuroanatomical pathology relevant to 15q11.2-q13.3 duplication syndrome.
Hum Mol Genet **26**, 3995-4010 (2017). https://doi.org/10.1093/hmg/ddx289

30   Gompers, A. L. *et al.* Germline Chd8 haploinsufficiency alters brain development in mouse. *Nat*
Neurosci **20**, 1062-1073 (2017). https://doi.org/10.1038/nn.4592

31   Silverman, J. L., Babineau, B. A., Oliver, C. F., Karras, M. N. & Crawley, J. N. Influence of
stimulant-induced hyperactivity on social approach in the BTBR mouse model of autism.
Neuropharmacology **68**, 210-222 (2013). https://doi.org/10.1016/j.neuropharm.2012.07.042

32   Berg, E. L. *et al.* Insulin-like growth factor-2 does not improve behavioral deficits in mouse and
rat models of Angelman Syndrome. *Mol Autism* **12**, 59 (2021). https://doi.org/10.1186/s13229-
021-00467-1

33   Huie, E. Z. *et al.* Peptidomimetic inhibitors targeting TrkB/PSD-95 signaling improves cognition
and seizure outcomes in an Angelman Syndrome mouse model. *Neuropsychopharmacology* **50**,
772-782 (2025). https://doi.org/10.1038/s41386-024-02020-z

34   Petkova, S. P. *et al.* Gait as a quantitative translational outcome measure in Angelman
syndrome. *Autism Res* **15**, 821-833 (2022). https://doi.org/10.1002/aur.2697

35   Symonds, J. D. *et al.* Incidence and phenotypes of childhood-onset genetic epilepsies: a
prospective population-based national Cohort. *Brain* **142**, 2303-2318 (2019).
https://doi.org/10.1093/brain/awz195

36   Cottam, N. C. *et al.* From Circuits to Lifespan: Translating Mouse and Human Timelines with
Neuroimaging-Based Tractography. *J Neurosci* **45** (2025).
https://doi.org/10.1523/JNEUROSCI.1429-24.2025

37   Born, H. A. *et al.* Early Developmental EEG and Seizure Phenotypes in a Full Gene Deletion of
Ubiquitin Protein Ligase E3A Rat Model of Angelman Syndrome. *eNeuro* **8** (2021).
https://doi.org/10.1523/ENEURO.0345-20.2020

38   Bikbaev, A., Frischknecht, R. & Heine, M. Brain extracellular matrix retains connectivity in
neuronal networks. *Sci Rep* **5**, 14527 (2015). https://doi.org/10.1038/srep14527

39   Cabrera-Garcia, D. *et al.* Early prediction of developing spontaneous activity in cultured neuronal
networks. *Sci Rep* **11**, 20407 (2021). https://doi.org/10.1038/s41598-021-99538-9

40   Vertes, P. E., Alexander-Bloch, A. & Bullmore, E. T. Generative models of rich clubs in Hebbian
neuronal networks and large-scale human brain networks. *Philos Trans R Soc Lond B Biol Sci* **369**
(2014). https://doi.org/10.1098/rstb.2013.0531

41   Spencer, M. C. *et al.* Multiscale evolving complex network model of functional connectivity in
neuronal cultures. *IEEE Trans Biomed Eng* **59**, 30-34 (2012).
https://doi.org/10.1109/TBME.2011.2171340

42   Downes, J. H. *et al.* Emergence of a small-world functional network in cultured neurons. *PLoS*
Comput Biol **8**, e1002522 (2012). https://doi.org/10.1371/journal.pcbi.1002522

43   Wagenaar, D. A., Pine, J. & Potter, S. M. Searching for plasticity in dissociated cortical cultures
on multi-electrode arrays. *J Negat Results Biomed* **5**, 16 (2006). https://doi.org/10.1186/1477-
5751-5-16

44   Moller, R. S. *et al.* Mutations in KCNT1 cause a spectrum of focal epilepsies. *Epilepsia* **56**, e114-
120 (2015). https://doi.org/10.1111/epi.13071

45   Kuchenbuch, M. *et al.* KCNT1 epilepsy with migrating focal seizures shows a temporal sequence
with poor outcome, high mortality and SUDEP. *Brain* **142**, 2996-3008 (2019).
https://doi.org/10.1093/brain/awz240

46   Chorny, L. E., Nordli, D. R., 3rd & Galan, F. Breath-holding spells: A typical case with EEG and
video correlate. *Epileptic Disord* (2024). https://doi.org/10.1002/epd2.20324

47   Frankel, W. N., Taylor, L., Beyer, B., Tempel, B. L. & White, H. S. Electroconvulsive thresholds of
inbred mouse strains. *Genomics* **74**, 306-312 (2001). https://doi.org/10.1006/geno.2001.6564

48   Mistry, A. M. *et al.* Strain- and age-dependent hippocampal neuron sodium currents correlate
with epilepsy severity in Dravet syndrome mice. *Neurobiol Dis* **65**, 1-11 (2014).
https://doi.org/10.1016/j.nbd.2014.01.006

49   Sukoff Rizzo, S. J., McTighe, S. & McKinzie, D. L. Genetic Background and Sex: Impact on
Generalizability of Research Findings in Pharmacology Studies. *Handb Exp Pharmacol* **257**, 147-
162 (2020). https://doi.org/10.1007/164_2019_282

50   Calsbeek, J. J. *et al.* Persistent neuropathology and behavioral deficits in a mouse model of
status epilepticus induced by acute intoxication with diisopropylfluorophosphate.
Neurotoxicology **87**, 106-119 (2021). https://doi.org/10.1016/j.neuro.2021.09.001

51   Sittig, L. J. *et al.* Genetic Background Limits Generalizability of Genotype-Phenotype
Relationships. *Neuron* **91**, 1253-1259 (2016). https://doi.org/10.1016/j.neuron.2016.08.013

52   Shore, A. N. *et al.* Reduced GABAergic Neuron Excitability, Altered Synaptic Connectivity, and
Seizures in a KCNT1 Gain-of-Function Mouse Model of Childhood Epilepsy. *Cell Rep* **33**, 108303
(2020). https://doi.org/10.1016/j.celrep.2020.108303

53   Numis, A. L. *et al.* Lack of response to quinidine in KCNT1-related neonatal epilepsy. *Epilepsia* **59**,
1889-1898 (2018). https://doi.org/10.1111/epi.14551

54   Xu, D. *et al.* Precision therapy with quinidine of KCNT1-related epileptic disorders: A systematic
review. *Br J Clin Pharmacol* **88**, 5096-5112 (2022). https://doi.org/10.1111/bcp.15479

55   Liu, C. Q. *et al.* Efficient strategies based on behavioral and electrophysiological methods for
epilepsy-related gene screening in the Drosophila model. *Front Mol Neurosci* **16**, 1121877
(2023). https://doi.org/10.3389/fnmol.2023.1121877

56   Cassibba, J. *et al.* Pulmonary Hemorrhage Revealing Multiple Vascular Malformations in a Child
with KCNT1 Developmental Epileptic Encephalopathy. *J Pediatr* **237**, 311-312 (2021).
https://doi.org/10.1016/j.jpeds.2021.06.019

57   Chauhan, D., Verhoeven, P. A., Kohli, U., Udassi, J. P. & Mascio, C. E. Life-threatening pulmonary
haemorrhage treated with coil embolisation followed by lobectomy in a patient with KCNT1
mutation. *Cardiol Young* **34**, 701-703 (2024). https://doi.org/10.1017/S104795112300447X

58   Garancini, N. *et al.* Pulmonary vascular malformations in genetic epileptic encephalopathy: A
rare, fatal case report. *Respir Med Case Rep* **53**, 102155 (2025).
https://doi.org/10.1016/j.rmcr.2024.102155

59   Kawasaki, Y. *et al.* Three Cases of KCNT1 Mutations: Malignant Migrating Partial Seizures in
Infancy with Massive Systemic to Pulmonary Collateral Arteries. *J Pediatr* **191**, 270-274 (2017).
https://doi.org/10.1016/j.jpeds.2017.08.057

60   Fusco, A. F. *et al.* Respiratory dysfunction in a mouse model of spinocerebellar ataxia type 7. *Dis*
Model Mech **14** (2021). https://doi.org/10.1242/dmm.048893

61   Gertler, T., Bearden, D., Bhattacharjee, A. & Carvill, G. in *GeneReviews((R))* (eds M. P. Adam *et*
al.) (1993).

62   Hagebeuk, E. E., van den Bossche, R. A. & de Weerd, A. W. Respiratory and sleep disorders in
female children with atypical Rett syndrome caused by mutations in the CDKL5 gene. *Dev Med*
Child Neurol **55**, 480-484 (2013). https://doi.org/10.1111/j.1469-8749.2012.04432.x

63   Jia, Y. *et al.* Quinidine Therapy for Lennox-Gastaut Syndrome With KCNT1 Mutation. A Case
Report and Literature Review. *Front Neurol* **10**, 64 (2019).
https://doi.org/10.3389/fneur.2019.00064

64   Lin, Z. *et al.* Efficacy of Anti-seizure Medications, Quinidine, and Ketogenic Diet Therapy for
KCNT1-Related Epilepsy and Genotype-Efficacy Correlation Analysis. *Front Neurol* **12**, 834971
(2021). https://doi.org/10.3389/fneur.2021.834971

65   Rizzo, F. *et al.* Characterization of two de novoKCNT1 mutations in children with malignant
migrating partial seizures in infancy. *Mol Cell Neurosci* **72**, 54-63 (2016).
https://doi.org/10.1016/j.mcn.2016.01.004

66   Copping, N. A. & Silverman, J. L. Abnormal electrophysiological phenotypes and sleep deficits in
a mouse model of Angelman Syndrome. *Mol Autism* **12**, 9 (2021).
https://doi.org/10.1186/s13229-021-00416-y

67   Gonzalez, E. A. *et al.* Sex-specific acute and chronic neurotoxicity of acute
diisopropylfluorophosphate (DFP)-intoxication in juvenile Sprague-Dawley rats. *Curr Res Toxicol*
2, 341-356 (2021). https://doi.org/10.1016/j.crtox.2021.09.002

68   Haigh, J. L. *et al.* Deletion of a non-canonical regulatory sequence causes loss of Scn1a
expression and epileptic phenotypes in mice. *Genome Med* **13**, 69 (2021).
https://doi.org/10.1186/s13073-021-00884-0

69   Purtell, H. *et al.* Electrographic spikes are common in wildtype mice. *Epilepsy Behav* **89**, 94-98
(2018). https://doi.org/10.1016/j.yebeh.2018.09.003

70   Andreone, B. J. *et al.* Durable suppression of seizures in a preclinical model of KCNT1 genetic
epilepsy with divalent small interfering RNA. *Epilepsia* (2025).
https://doi.org/10.1111/epi.18278

830
831

Translational Stage	Subjects/Model System	Genotypes Tested	Phenotypes found	
Clinical	5 female and 7 male MMPSI patients	A934T; R428Q; R474H; I760M	EEG: migrating seizures. Brain MRI: delayed myelination, thin corpus callosum. Hypotonia	
Clinical	248 patients	152 EIMFS; 37 DEE; 53 ADNFLE and 6 with other phenotypes	Literature review: clinical seizures, sleep disturbances, hypotonia, and other features	
Clinical	43 patients	n=1 unless otherwise stated: A477T; A934T; R398Q; R398Leu; R428Q (n=2); R474H (n=3); R929Q; R950Q (n=3); R961H; R961S; Q651R; G288S (n=3); K947E; F932S	Quinidine treatment resulted in a >50% seizure reduction in patients with G288S; F932S; R961S; R950Q; A477T while seizures worsened for 3 patients with R474H variant.	
Clinical	30 patients: 17 boys and 13 girls	not specified	Ketogenic diet therapy was effective regardless of the variant, especially effective in patients with functional domain variant-related epilepsy. Quinidine was effective in patients with non-functional domain variant-related epilepsy.	
Preclinical (human sequence)	Human iPSC-derived neurons	Homozygous P924L	Sodium-dependent potassium currents are increased; duration of action potentials is decreased in current clamp recordings with an increase in the total number of action potentials	
Preclinical (human DNA sequence; non neuronal specific)	HEK293T cells	T314A	decreased current amplitude abolished channel activation	
		G288S, R398Q, N449S, L781V, E893K, M896V, F932L, R928C, S937G, L942F, R961H, A965T	increased current amplitude	
		L781V, E893K, M896V, R928C, F932L	increased channel open probabilities	
Preclinical (human DNA sequence; non neuronal specific)	*Xenopus laevis* oocytes and HEK cell lines	36 variants	significant increases in current magnitude for G288S, A934T, R389Q, R428Q, and R474H variants. G288S and R398Q increased channel kinetics. R474H reduced channel kinetics. 28 of 33 variants tested had GOF impacts on current magnitude, V1/2, or both	

Preclinical (species specific DNA sequence; non neuronal specific)	*Xenopus laevis* oocyte	hKCNT1 mutations M896I, R398Q, Y796H, R928C, R428Q, A934T, P924L and the rat rKcnt1 mutant R907C	KCNT1 mutations increase current magnitude and hasten (M896I, R928C and A934T) or slow (R398Q) activation, increase current amplitude, and modulate channel kinetics. Quinidine at 300uM was effective in blocking current amplitude with varying sensitivity.	
In vivo preclinical	*Drosophila*	hKCNT1, G288S, R928C	Increases in K+ conductance, hyper excitability and seizures	Eha
In vivo and *In vitro* preclinical	*Drosophila* and HEK293T cells	hKCNT1, G288S, R398Q or R928C	Expression in GABAergic neurons resulted in seizure phenotypes resistant to carbamazepine, valproic acid, vigabatrin, quinidine, and cannabidiol.	
In vivo and *In vitro* preclinical	CHO cells and mice	Cells: L456F of human KNa1.1 Mouse: Kcnt1 L457F Background strain: C57BL/6J	No significant differences between genotypes on Rotarod, Zero maze, Open Field, home cage. No differences by GT to seizure susceptibility by flurothyl (20ul/min). While KI/+ and KI/KI mice were susceptible to both Kainate (30mg/kg) and Pentylenetrazole (60mg/kg) induction.	
In vitro preclinical (neuronal specific)	Mouse primary neurons	Y796H (mouse Y777H) Background strain: C57BL/6NJ	No effects on glutamatergic or VIP neuron function. SST neurons became hypoexcitable with a higher rheobase current and lower action potential firing frequency, whereas PV neurons became hyperexcitable with a lower rheobase current and higher AP firing frequency	
In vivo preclinical	Mice	Y796H (mouse Y777H) Background strain: C57BL/6NJ	Alterations in GABAergic neurons including hyperexcitability and hyper-synchronicity. Additional Epileptiform activity was found in HOM mice only via Video-EEG.	

In vivo preclinical	Mice	Y796H (mouse Y777H) crossed with Snap25-GCaMP6s Background strain: C57BL/6NJ	Widefield Ca2+ imaging in HOM and WT mice showed no differences in mesoscale interictal activity	T
In vivo preclinical	Mice and *Xenopus laevis* oocytes	Y796H (mouse Y777H) Background strain: C57BL/6 13 hKCNT1B variants	hKCNT1B and variants were characterized in *Xenopus laevis* oocytes for activity and responsiveness to 13 potassium channel blockers, of which tipepidine and hydroquinine were the most effective. Additionally, these two drugs were effective at blocking spontaneous seizures in Y777H HOM mice.	
In vivo preclinical	Mice	Y796H (mouse Y777H) generated on background strain C57BL/6NJ, but backcrossed to C57BL/6J in this study (number of crossings not specified)	Spontaneous seizures in KCNT1 HOM mice occur during NREM sleep and suppress REM sleep. (NREM: synchronized EEG with high-amplitude, delta frequency [0.5–4 Hz] activity and low EMG activity; REM: high power at theta frequencies [6–9 Hz] and low EMG activity)	
In vivo preclinical	Mice	R474H (mouse R455H) developed on C57BL/6J and backcrossed for 4-5 generations Knock-out mice (Kcnt1$^{-/-}$) developed on Sv/129 and backcrossed with C57BL/6NJ for 2 generations. WTs were from a parallel colony of Sv/129 backcrossed with C57BL/6NJ	Knock out mice were hypoactive in open field and on the rotarod, with no differences in seizure susceptibility from WT. Homozygous R455H was embryonic lethal. R455H/+ mice had increased spontaneous seizures in EEG and a decreased threshold for PTZ (60mg/kg) induced seizures compared to WT with no motor differences in the open field or rotarod	
In vitro and *Ex Vivo* preclinical	Mouse primary neurons and slice	R474H (mouse R455H) Background strain: C57BL/6J	Na+-dependent K+ (KNa) and voltage-dependent sodium (NaV) currents are increased in both excitatory and inhibitory cortical neurons leading to increases in firing of excitatory neurons and suppressing firing of inhibitory neurons.	

In vivo preclinical	Mice	Knock-out mice (Kcnt1^{-/-}) developed on C57BL/6J and backcrossed to C57BL/6NJ for >6 generations	Complete loss of hippocampal LTP and LTD, Reduced NMDA receptor signaling, suppresses the expression of multiple proteins that localize to axons, dendrites and myelin sheath, but increases levels of inner mitochondrial membrane proteins
		R455H/+ Background strain: C57BL/6J	Some protein changes, but much milder compared to KO. ASO treatment did reverse some of the protein changes seen in this model

832 **Table 1. Summary of clinical and preclinical data using model systems or clinical studies**
833 **related to the novel translational model herein, *Kcnt1*^{+/G288S}.**