Title Pages

Article Type:

Research Paper

Title:

Aphid infestation induces plant-sex-specific changes in floral chemistry and pollinator

behaviour in *Silene latifolia*

Runing Title:

Sex-specific floral responses to aphid infestation

Authors:

Kaya B. Zill[1], Thomas Stegemann[2], Elisabeth Kaltenegger[3], Wolfgang Bilger[5], Tobias J.

Demetrowitsch[4], Henry Berndt[6], Alexandra Erfmeier[1], Sybille B. Unsicker[5], Karin Schrieber[1]

Affiliations:

[1] Kiel University, Faculty of Mathematics and Natural Sciences, Institute for Ecosystem

Research, Geobotany

[2] Kiel University, Faculty of Mathematics and Natural Sciences, Pharmaceutical Institute,

Pharmaceutical Biology

[3] Kiel University, Faculty of Mathematics and Natural Sciences, Botanical Institute and

Botanical Garden, Biochemical Ecology and Molecular Evolution

[4] Kiel University, Faculty of Agricultural and Nutritional Sciences, Institute for Human Nutrition

and Food Science, Division of Food Technology

[5] Kiel University, Faculty of Mathematics and Natural Sciences, Botanical Institute and

Botanical Garden, Plant-Environment-Interactions

[6] Kiel University, Mathematics and Natural Sciences, Zoological Institute, Marine Symbiosis

E-Mail addresses and Orchid IDs:

Karin Schrieber: kschrieber@ecology.uni-kiel.de, 0000-0001-7181-2741

Thomas Stegemann: tstegemann@pharmazie.uni-kiel.de, 0000-0002-0391-3337

Title Pages

Sybille Unsicker: sunsicker@bot.uni-kiel.de, 0000-0002-9738-0075

Tobias Demetrowitsch: tdemetrowitsch@foodtech.uni-kiel.de, 0000-0002-3528-0255

Elisabeth Kaltenegger: ekaltenegger@bot.uni-kiel.de, 0000-0001-6542-1509

Wolfgang Bilger: wbilger@bot.uni-kiel.de, 0000-0001-7800-4210

Alexandra Erfmeier: aerfmeier@ecology.uni-kiel.de, 0000-0002-1002-9216

Karin Schwarz: info@foodtech.uni-kiel.de, 0000-0002-3943-0564

Henry Berndt: hberndt@zoologie.uni-kiel.de, 0000-0002-0735-5539

Kaya Zill: NA, stu218014@mail.uni-kiel.de

Corresponding Author:

Dr. Karin Schrieber

Kiel University

Institute for Ecosystem Research

Olshausenstr. 75

24118 Kiel

Phone: +49 (0) 431 880-4082

E-mail: kschrieber@ecology.uni-kiel.de

Counts:

Words main text excl. material and methods: 5148

Figures: 6

Tables: 3

Supporting information: 2

Highlight, Abstract, Keywords

Highlight:

Aphid infestation alters multiple visual and chemical floral traits in a plant sex-specific manner,

leading to reduced attractiveness to moth pollinators in female plants, but not in males.

Abstract:

Pollinators consistently share the complex information and resource landscape of their host

plants with herbivores. Yet, how sap feeders affect floral attractiveness to pollinators remains

poorly understood, despite the critical role of this tripartite interaction in natural and agricultural

ecosystems. These intricate interactions may be further enriched by pronounced sexual

dimorphisms in dioecious species. In this study, we examined how infestation by the

oligophagous aphid *Brachycaudus lychnidis* affects sex-specific interactions among the

dioecious plant *Silene latifolia* and its specialist moth pollinator *Hadena bicruris*. We exposed

male and female plants to aphid herbivory and evaluated its effects on floral traits (visual cues,

floral scent, and nectar chemistry) and pollinator behaviour. Contrary to sexual selection theory,

female plants were not less attractive than males under control conditions. Instead, our results

suggest that females exhibit plasticity in floral trait expression depending on pollination

success. Aphid infestation altered floral traits in a sex-specific way, with stronger declines in

female attractiveness linked to nectar compounds potentially acting as feeding cues or

behavioural modulators. This indicates sex-specific defence strategies and possible

involvement of indirect defences in stabilizing this specialized plant–pollinator–herbivore

system.

Keywords: *Brachycaudus*, dioecy, *Hadena bicruris*, nectar chemistry, nocturnal pollination,

metabolomics, *Silene latifolia*, sexual dimorphism, plant-pollinator-herbivore interaction

Graphical Abstract:

Plant-sex specific effect of aphid infestation on floral traits (number, size, colour, scent composition, nectar quantity and composition) and pollinator behaviour.

Abbreviations used in the main text:

- ANOVA: analysis of variance
- FC, FA, MC, MA: female control, female aphid infested, male control, male aphid infested
- GC-MS: gas chromatography-mass spectrometry
- (G)L(M)M: (generalized) linear (mixed effects) model
- NMDS: nonmetric multi-dimensional scaling
- UHPLC-ESI-qTof-MS/MS: quadrupole time of flight - electro spray ionization - tandem mass spectrometry
- PERMANOVA: permutational analysis of variance
- VOCs: Volatile organic compounds

1. Introduction

Understanding plant × pollinator × herbivore interactions and their component species' co-evolutionary dynamics has become a central goal in chemical ecology (Kessler and Chautá, 2020; De-la-Cruz *et al.*, 2022; Muola *et al.*, 2022). The complex plant traits mediating interactions with pollinators and herbivores have been shown to overlap significantly. While the ecological consequences of this shared information space are now well studied in systems involving chewing-biting herbivores, the effect of sap feeders on communication and resource exchange with pollinators is poorly understood, despite their critical role in the function of both, natural and agricultural ecosystems (Morales-Hojas, 2017; Noriega *et al.*, 2018).

Plants evolve diverse traits to attract pollinators and modulate their behaviour thereby maximizing their own fitness with minimal resource investment (Santamaría and Rodríguez-Gironés, 2015). Next to flowering phenology and visual signals (flower number, spatial arrangement, size, shape, hue, and colour luminance), floral chemistry plays a central role in ensuring these complex functions (Borghi *et al.*, 2017; Trunschke *et al.*, 2021; Wong *et al.*, 2023). The emission of floral volatile organic compounds (VOCs) can trigger oriented movement of the receiver towards the emitter, and elicit landing, feeding as well as reproductive behaviour (Muhlemann *et al.*, 2014). Such scent signals can be honest and guide visitors to a high-quality reward, or deceive them about the presence of food or mates, thereby exploiting their pollination service (Ito *et al.*, 2021; Peakall, 2023). Nectar - the primary nutritional reward provided to pollinators - is a complex secretion composed of sugars, amino acids, lipids, proteins and diverse secondary metabolites (Nicolson, 2022), and thus far more than a ready-to-use energy source for feeding animals. Plants may use nectar as a toolkit to manipulate the behaviour of their floral visitors. Secondary nectar metabolites can cause phagostimulation and satiety; improve learning capacity and memory; promote muscle performance, motivation and arousal; but also initiate sluggish and imprecise movements (Nepi *et al.*, 2018; Mustard, 2020; Barberis *et al.*, 2023). These very functions facilitate the exploitation of highly efficient pollination services while reducing energy investment in nectar sugars and amino acids, thus not specifically benefiting the floral visitors. Although we are becoming increasingly aware of this beautiful complexity, there are still few studies that take a holistic view of the pollination syndrome, considering the combined roles of visual traits, floral VOCs, and nectar metabolites.

Pollinators universally share the complex information and resource landscape of their host plants with herbivores. There is now evidence for interactions among traits and genes controlling plant reproduction and defence strategies, leading to complex reciprocal co-evolutionary feedbacks (Kessler and Chautá, 2020; De-la-Cruz *et al.*, 2022). Herbivores may

directly reduce plant attractiveness to pollinators through aggression or the consumption of

floral tissues and vegetative structures necessary for their production (McCall and Irwin, 2006).

Multiple floral traits have thus evolved to protect floral tissues from herbivory (Paul *et al.*, 2022).

Constitutive and induced plant defences may not only alter a plant's suitability to herbivores,

but also deter pollinators (Adler *et al.*, 2012) or reduce resource allocation to pollinator

attraction (Stamp, 2003). Specifically chemical plant traits are considered to mediate multiple

other antagonistic and synergistic pleiotropic effects on pollinator attraction and herbivore

defence (Muola *et al.*, 2022). So far, insight into such complex tripartite interactions comes

exclusively from chewing-biting herbivores. Aphids, which are among the most damaging plant

pests (Dedryver *et al.*, 2010), however, have been largely overlooked. These exceptionally fast

reproducing insects feed passively on the phloem sap of their hosts, thereby rapidly depleting

them of essential nutrients, altering their source-sink relationships, and excreting phytotoxins

as well as phytopathogenic viruses with their saliva (Nalam *et al.*, 2019; Twayana *et al.*, 2022).

Plants in turn, actively defend against aphids by changing the composition of sugars and amino

acids in the phloem sap or loading it with deterrents, toxins or anti-digestive proteins (Nalam

et al., 2019; Twayana *et al.*, 2022). Floral traits attracting pollinators are specifically prone to

the direct and indirect effects of aphid infestation, as aphids often feed close to reproductive

structures on flowering stems. This proximity suggests a high potential for interference with

pollinator-mediated reproductive processes, warranting closer empirical attention.

The eco-evolutionary link between pollinator attraction and herbivore defence becomes

particularly apparent in dioecious species (Johnson *et al.*, 2015). Plants that allocate male and

female reproductive function to different individuals often exhibit complementary sexual

divergence in both traits based on dramatic differences in resource demands for successful

reproduction (Barrett and Hough, 2013; Johnson *et al.*, 2015). Male success is predicted to

largely depend on efficient pollen transfer and thus pollinator attraction, while female success

relies on pollination, as well as on the production and defence of seeds, and their dispersal

(Moore and Pannell, 2011). Consequently, females are predicted to allocate fewer resources

to floral attractiveness but exhibit more efficient herbivory defence and tolerance as compared

to males (Avila-Sakar and Romanow, 2012; Barrett and Hough, 2013). As recent research has

produced contradictory evidence for these theories (Cornelissen and Stiling, 2005; Tang-

Martínez, 2012; Sargent and McKeough, 2022), a central question remains: What are the sex-

specific ecological consequences of parallel interactions with pollinators and herbivores in

dioecious plant species? Addressing this question is key to understanding how co-selection by

pollinators and herbivores contributes to the evolution of divergent reproductive strategies in

dioecious plants.

A system that is ideal to study these complex interplays consists of the dioecious plant *Silene*

latifolia Poir. (Caryophyllaceae), its nocturnal pollinator *Hadena bicruris* Hufn*.* (Noctuidae)*,*

and the aphid *Brachycaudus lychnidis* L. (Aphididae). The species involved in this system

engage in highly specialized relationships. Researchers have only begun to uncover the

sophisticated functional traits that have co-evolved to stabilize the cost-benefit ratio of their

individual bipartite interactions (Blair and Wolfe, 2004; Dötterl *et al.*, 2006; Labouche and

Bernasconi, 2010; Schrieber *et al.*, 2021). Insight into the ecology of the tripartite interaction,

however, is lacking. Here, we exposed female and male *S. latifolia* plants to control and *B.*

lychnidis infestation treatments to assess flowering phenology, spatial flower arrangement,

flower colour, the chemical composition of floral scent and nectar as well as nectar quantity,

and attractiveness to *H. bicruris*. We specifically aimed at characterizing i) sexual dimorphisms

in floral attractiveness; ii) the effect of aphid infestation on floral attractiveness to pollinators;

and iii) sex-specific differences in the susceptibility to aphid infestation and the quality and

magnitude of responses to aphid infestation.

2. Methods

2.1 Study organisms

Silene latifolia (white campion, *Figure 1A*) is a short-lived perennial plant within the
Caryophyllaceae family, which is native to ruderal habitats across Eurasia and North Africa.
The species belongs to the few angiosperms featuring a dioecious reproductive system and
being simultaneously pollinated by specialized animals (Charlesworth, 1993; Käfer *et al.*,
2017). *Silene latifolia* plants exhibit a distinct moth pollination syndrome. Their white, large,
and funnel-shaped flowers open at dusk to release a scent bouquet composed of more than
60 VOCs (Dötterl and Jürgens, 2005; Dötterl *et al.*, 2005) to advertise large amounts of nectar
rewards (Tew *et al.*, 2021) with a mostly unknown chemical composition. Male and female *S.*
latifolia are known to exhibit pronounced sexual dimorphisms in a broad set of floral traits
(Gehring *et al.*, 2004; Dötterl *et al.*, 2005; Delph *et al.*, 2010; Schrieber *et al.*, 2021), which are
largely in line with theory of sexual selection (Bateman, 1948).

Hadena bicruris (campion owl, *Figure 1B*) - a crepuscular (active during twilight) moth within
the Noctuidae family - is the most relevant pollinator species of *S. latifolia* in Eurasia (Jürgens
et al., 1996; van Putten *et al.*, 2007). Together both species form one of only ~13 worldwide
known nursery pollination systems (Dufaÿ and Anstett, 2003). Female moths pollinate female
plants while ovipositing on flower ovaries to provide their hatching larvae with nutritious seeds
and shelter by developing fruits. The pollination services provided by male moths, together
with the high egg and larval mortality caused by parasitoids and predators of *H. bicruris*,
compensate for the costs of seed consumption (Labouche and Bernasconi, 2010; Villacañas
de Castro and Hoffmeister, 2020). Together these factors shift the outcome of the interaction
for the plant towards the latter end of a seed predation-pollination continuum. A substantial
fraction of VOCs in the floral scent of *S. latifolia* triggers antennal and behavioural responses
in *H. bicruris* (Dötterl *et al.*, 2006). While the function of floral scents in this intricate co-
evolutionary relationship is well studied (Dötterl *et al.*, 2006), the chemistry of nectar - the only
reward and main food source provided to adult moths - is largely unexplored.

A third species participating in this highly specialized system within Eurasia is the holocyclic
aphid *Brachycaudus lychnidis (Figure 1C)*, which spends both the parthenogenetic and sexual
part of its life cycle on *Silene*. The aphid feeds mainly on the phloem sap of flowering stems.

2.2 Plant and insect cultivation

Seeds from five *S. latifolia* mother plants were collected in a population in Wiesengrund,
Germany (51°41'17.1"N, 14°35'45.8"E) and experimentally outcrossed for two generations
under greenhouse conditions (see Schrieber *et al.*, 2019 for details). We cultivated 30 plants

per maternal family individually in 360 mL pots filled with organic potting soil (Floragard TKS®

2 Instant Plus, DE) in a greenhouse at 25 °C / 15 °C and 60 % RH in a 16 h day / 8 h night

cycle. When plants started flowering after 6 weeks, we selected two males and two females

from each of the five maternal families (N = 20). The selected plants were transferred into 8 L

pots filled with the above-mentioned substrate and pruned to a height of 15 cm to induce over-

compensatory growth. Please note that *S. latifolia* is adapted to grazing/mowing and thus

cutting is near-natural (Jessen *et al.*, 2022). Moreover, cutting was necessary to induce the

emergence of multiple shoots and sufficient flower production for the experiment. Plants were

fertilized with 0.3 g Universol® Yellow (Everris-Headquarters, NL) dissolved in 200 mL of water

every other week and prophylactically treated with sulphur vapor to prevent fungal infections

as well as with biological pest control agents to prevent thrips and uncontrolled aphid

infestation (agents: *Amblyseius cucumeris against thrips; Aphidius ervi, Aphidius colemani and*

Lysiphlebus testaceipes against aphids Katz Biotech GmbH, DE). The preventive treatment

with parasitoid wasps was stopped three weeks before controlled aphid infestation.

Adult *H. bicruris* moths were collected in different wild *S. latifolia* populations around Kiel,

Germany (54°19′ N, 10°8′ O) in 2022 and kept together in ventilated, transparent 20 x 30 cm

boxes equipped with egg cartons and tissue papers within a climate chamber (MLR-352H,

PHC Corporation, Ōizumi-machi, JP) at 25 °C / 15 °C and 60 % RH in a 16 h day / 8 h night

cycle. They were fed a honey-water solution (6 g organic flower honey, 14 mL distilled water)

from artificial PET *S. latifolia* flowers. In addition, moths received one fresh male and female

S. latifolia flower per day, as they did not reproduce without them. Individual flowers were

harvested from a pool of food plants and presented in Eppendorf tubes filled with water.

Hadena bicruris eggs, which were deposited all across the breeding boxes, were collected and

placed on moist filter paper in individual ventilated plastic containers (50 mL). Upon hatching,

larvae were provided with an artificial diet according to [43], which was additionally

supplemented with 20 mL of *S. latifolia* seeds per 500 mL to increase acceptance. Pupated *H.*

bicruris were moved to ventilated 50 mL containers filled with sterile sand and newly hatched

moths were transferred into the above-mentioned boxes for adults.

Brachycaudus lychnidis aphids were collected and identified according to

(influentialpoints.com, 2024) in spring 2024 in different wild *S. latifolia* populations around Kiel,

Germany (54°19′ N, 10°8′ O). They were cultivated in net cages on a balanced mixture of male

and female *S. latifolia* plants from our study population in a climate chamber (HGC 1514

Module 7, Weiss Umwelttechnik GmbH, DE) at 23 °C and 55 % RH, in a 16 h day / 8 h night

cycle. Food plants were watered and exchanged on demand.

2.3 Experimental setup and data acquisition

In mid-July 2024 at the age of 13 weeks, all selected plants were enclosed in 150 × 50 cm organza mesh bags and one individual per sex (male, female) × maternal family (I-V) combination was inoculated with 50 adult aphids *(Figure 2A, Supplementary Figure S1A-C)*. All aphid colonies showed an initial decline in population size of max. 10 % and switched to positive growth rates max. three days after the inoculation. Thus, our approach does not account for potential plant sex-specific host choice, early phase plant defences preventing colonization success, and indirect defence; but focuses on the effects of established aphid colonies having optimal growth conditions. Data on plant stress indicators, pollinator choice, visual flower traits, floral scent, and nectar *(Figure 2B-F)* were collected over a 40-day period following aphid inoculation *(Figure 2G)*.

2.3.1 Plant stress indicators

To quantify the disruption of physiological homeostasis under aphid infestation, we optically assessed the chlorophyll concentration, and the absorbance by epidermally located UV-absorbing compounds, mainly flavonoids using the LEAF-STATE-ANALYZER LSA-2050 (Heinz Walz GmbH, DE) *(Figure 2B&G)* for all plants. As reference, data provided with the instrument for the lower leaf side of *Sedum telephium* was used. Data were acquired directly before and 40 days after aphid inoculation between 11:00 am and 12:00 noon in five leaves per plant individual. The selected leaves were evenly distributed along the vertical axis of the plants and well exposed to light (i.e., not significantly shaded by other leaves) and the measured area, located in the upper middle part of the lamina, was free of honeydew and aphids, which typically feed on stems. Immediately after these measurements, we counted the number of aphid individuals per size category (small: < 1.5 mm, medium: < 2 mm, large: > 2 mm) directly on the plant. Subsequently, the aphids were carefully washed off the plants with warm tap water, collected on filter paper and dried for 24 h at 25 °C, then for 24 h at 65 °C, and finally for 24 h at 105 °C to determine total aphid dry biomass per plant.

2.3.2 Pollinator behaviour

Behavioural assays with *H. bicruris* were conducted under laboratory conditions 30-39 days after aphid inoculation *(Figure 2C&G)* between 8:00 pm and 0:30 am - the peak activity time of our captive moths. For the time of data acquisition in the choice assays, adult moths were kept individually in ventilated, transparent boxes (15 × 12 × 11 cm) equipped with shelters and a feeding station featuring a honey-water solution as described in *chapter 2.2*. For each assay, the artificial diet was removed in the morning and replaced in the evening with two fully

developed (male: fully dehisced anthers, female: fully expanded and papillous stigma), freshly

picked flowers from the experimental plants, which were carefully fixed in lidless Eppendorf

tubes filled with tap water. The two flowers were positioned on opposite sides of the boxes for

maximum spatial separation *(Supplementary Figure S1D)* to provide the moths with the choice

among the following treatment groups: control females *versus* aphid infested females, control

males *versus* aphid infested males, control females *versus* control males, aphid infested

females *versus* aphid infested males. The moth individuals used in these behavioural tests (N

= 11) differed in age and sex (5 females, 6 males; moth sex was accounted for in the statistical

analysis) and were tested in multiple-choice trials using a fully randomized design with respect

to plant treatment groups and individuals, completing a total of 66 trials. Each night, up to eight

moths were used, with two observers monitoring four moths each. The boxes were evenly lit

with dim red LED lamps to allow observation without confounding the trials (the visual system

of moths cannot capture red light). Three behavioural parameters were quantified for each of

the two flowers: (i) the number of visits, with a visit starting and ending with physical contact to

the flower; (ii) the number of feeding attempts per flower, with an attempt starting/ending by

unrolling/rolling up and inserting/removing the proboscis into/from the corolla tube; and (iii) the

total time spent with any physical contact on the flower.

2.3.3   Visual flower traits

We measured a broad set of visual flower traits at the plant individual and single flower level

throughout the experiment *(Figure 2D&G)*.

Flower colour was quantified using a digital image transformation method that accounts for

natural light conditions and the visual system of the pollinator based on (Troscianko and

Stevens, 2015). Images were acquired 21-25 days after aphid inoculation for three to six

flowers (depending on availability) from all plant individuals. They were taken outdoors in an

open area, free from shadows cast by trees or buildings, during the last hour of daylight on

rain-free days to match the natural light conditions perceived by *Hadena* (Bopp and

Gottsberger, 2004). A fully developed but fresh flower was plugged into a custom-made black

ethylene vinyl acetate platform, which featured two reflectance standards (PTFB 10% and

Spectralon 99%, Labsphere, UK) along with a size reference *(Supplementary Figure 1E)*.

Images were captured using a modified Samsung NX1000 camera (Suwon, KOR) with its

sensor filter removed, to achieve full-spectrum sensitivity (300 – 1000 nm). The camera was

equipped with a UV-sensitive lens (Nikon EL 80 mm, JP). To capture both visible and ultraviolet

(UV) light, we took one photo with a UV/IR blocking filter (UV/IR Cut, transmitting 400-700 nm,

Baader Planetarium, DE) and another one with a UV pass plus IR blocking filter (U-filter,

transmitting 300-400 nm, Baader Planetarium, DE) using a custom-made filter slide mount. All

photos were taken with an aperture of 5.6, and ISO of 800, whereas the shutter speed was

adjusted to the available light conditions to receive well-exposed photos. Post-processing was

carried out using the Multispectral Image Calibration and Analysis (MICA) Toolbox plugin

(Troscianko and Stevens, 2015) within ImageJ 1.47 t (Rueden *et al.*, 2017). We selected the

flower silhouette for colour analysis. The images were linearized to correct the camera's non-

linear light response and standardized based on the two reflectance standards to compensate

for variations in natural lighting. We fitted the multispectral images to a mathematical model

incorporating (i) the spectral sensitivity of our Samsung NX1000-Nikkor EL 80 mm 300-700 nm

camera (data derived from Troscianko and Stevens, 2015); (ii) the spectral sensitivities of the

three photoreceptors in the moth compound eye (data derived from Johnsen *et al.,* 2006); and

(iii) the spectral composition of sun light during dusk (data derived from Johnsen *et al.,* 2006).

For further details see Schrieber *et al.* (2021). The outcomes of these analyses are the

percentage of green, blue and UV-A dusk sun radiation reflected by the flower, which are

perceivable by a moth's visual system.

Flower area was measured from the same images for three to six flowers per plant individual.

We assessed the exact area covered by a single, well developed petal limb, the area covered

by the exact flower silhouette and the area covered by the circular perimeter *(Figure 2D, 1 =*

flower perimeter, 2 = flower silhouette, 3 = petal silhouette). The circular perimeter reflects the

flower's maximum extension, but not its visual impact on pollinators due to natural variation in

the shape of single petals, while the area covered by the exact flower silhouette may represent

what pollinators see but not the plant's investment into flowers (petal overlap), which can be

captured by individual petal area.

The number of opened flowers and flower buds per plant individual were recorded at three

time points during the experiment (10, 20 and 40 days after aphid inoculation, *Figure 2D&G*)

and corrected for the flowers removed for other analyses. Directly after the removal of aphids

(40 days after aphid inoculation), plant aboveground parts were cut off directly on top of the

soil and separated into shoots (without leaves), and flowers. Plant parts were dried individually

in paper bags for 24 h at 65 °C followed by an additional 24 h at 105 °C to determine dry

biomass. During data acquisition for pollinator behaviour, flower colour, and nectar rewards,

we collected the harvested flowers in plant individual-specific bags to assess their dry biomass

as described above and added it to the total flower dry biomass.

2.3.4   Flower scent

At day 27 after the aphid inoculation, floral VOC emission was collected for all experimental

plants during the peak emission in *S. latifolia* between 10:00 pm (sunset) and 1:00 am (Dötterl

et al., 2005) using a mobile push-pull system as described in Berkum *et al.*, (2024) . Up to

three fresh and fully developed flowers per plant from different shoots were enclosed in 35 cm

× 31 cm PET bags (RUBIN® Bratschlauch, DE), which were closed at both ends with cable

ties *(Figure 2E)*. Compressed air entered the system after passing through an activated

charcoal filter on the lower side of the bag at a flow rate of 1 L min^{-1} and was pulled out at the

top through an adsorbent Poropak filter (Alltech, USA), using a vacuum at the rate of 0.6 L/min.

After VOC collection, the filters were eluted with 200 µL dichloromethane (Sigma Aldrich, DE)

containing 10 ng/µL of nonyl acetate (Sigma Aldrich, DE) as an internal standard. As the

headspace of treated plants also included aphids, we additionally determined which

compounds were exclusively released by *Brachycaudus*. We enclosed 60 aphids (20 small: <

1.5 mm, 20 medium: > 1.5 mm, 20 large: > 2 mm) from our laboratory population in the above-

described PET bags to collect and analyse their volatile emissions as described for the plants

with three replicates.

Samples were analysed on a gas chromatograph (TRACE GC1310, ThermoFisher Scientific,

DE) equipped with a TraceGOLD TG-5MS column (30 m × 0.25 mm × 0.25 µm + 5 m;

ThermoFisher Scientific, DE) and coupled with a mass spectrometer (TSQ Duo MS,

ThermoFisher Scientific, DE). A sample volume of 1 µL was injected with a split ratio of 1:10 at

250 °C and migrated with a helium-flow of 1 mL min^{-1}. The transfer line was set to 270 °C and

the ion source of the mass spectrometer was operated at 230 °C and 70 eV for ionization. The

temperature program started at 40 °C for 2 min, ramped to 180 °C at a rate of 6 °C min^{-1},

ramped to 300 °C at a rate of 60 °C min^{-1}, and finished with a 5 min hold time at 300 °C. Mass

spectra of separated compounds were acquired in full scan mode between 33 and 350 m/z.

An alkane standard mix (C_8-C_{20}, Sigma Aldrich, DE) was analysed under the same conditions

to calculate Kovats retention indices (RI) for the targeted compounds. The data processing

was performed with Chromeleon^{TM} 7.3 (ThermoFisher Scientific, DE). The putative identity of

compounds was determined via comparisons of mass spectra and RI's with databases (NIST

Mass Spectral Search Program Version 3.0, (Smith *et al.*, 2004)) and (Dötterl *et al.*, 2005). The

identities were assigned according to the following criteria: match factor to the NIST databank

>750 and RI difference < 10. The base peak in the mass spectrum was used as quantitation

ion. Additionally, two secondary ions (confirmation ions) were selected per compound to

ensure correct peak assignment *(Supplementary Table S1A)*. The relative compound

abundance was quantified by dividing the peak area (based on individual quantitation ion) of

each volatile by the peak area of the internal standard nonyl acetate (quantitation ion 43,

confirmation ions 56, 70). To account for variation in flower number in the headspace, scent

abundance was normalized per flower. We selected all VOCs for the statistical analyses that

could not be clearly identified as contaminations.

370   2.3.5   Nectar quantity and chemistry

We measured the nectar volume for the flowers that have been harvested for the colour and

size analyses, directly after they have been photographed. All nectar was extracted from the

base of the ovary / the base of the stamens and anthopophor into 1 µL microcapillary tubes

(Minicaps NA-HEP, Hirschmann Laborgeräte, DE). The length of the nectar column was

measured with a calliper to determine the exact volume.

Nectar chemistry was addressed based on a standardized nectar volume via Ultra High

Performance Liquid Chromatography - Quadrupole Time of Flight - Electro Spray Ionization -

Tandem Mass Spectrometry (UHPLC-ESI-qTof-MS/MS). For these analyses, we additionally

collected all nectar from three fresh, fully developed flowers per plant individual using 1 µL

microcapillary tubes on day 28 after aphid infestation. After determining the exact nectar

volume as described above, each microcapillary was rinsed with 30 µL of HPLC grade water

(Agilent, USA) into a 0.5 mL Eppendorf tube, in which all nectar from one plant individual was

pooled. The samples were then vortexed, snap-frozen in liquid N_2 and stored at -80 °C.

Samples underwent a modified simultaneous metabolite, protein, and lipid extraction

(SIMPLEX) as described in (Schrieber *et al.*, 2023) to obtain both, hydrophilic and lipophilic

nectar metabolites. After defrosting and vortexing, 0.1 mL of nectar solution was mixed with

0.5 mL of ultra-pure methanol (Carl Roth, GE) and 4 mL of methyl-tert-butylether (LC-grade,

Carl Roth, GE). Samples were incubated for 30 min at 25 rpm in an end-over-end shaker

before phase separation was induced by adding 0.5 mL of ultra-pure water (Carl Roth, GE).

After 10 min of incubation, samples were centrifuged at 4000 *g* and 4 °C for 10 min and the

upper lipophilic phase was collected. The lower hydrophilic phase was used as a template to

repeat the above-described protocol for the extraction of hydrophilic metabolites. Extracts were

re-suspended in 0.5 mL of either ultra-pure water and ultra-pure methanol (hydrophilic extracts:

50 / 50, v / v, Carl Roth, GE) or ultra-pure isopropanol and LC-grade chloroform (lipophilic

extracts: 3:1, v / v, Carl Roth, GE). The dilution of the resulting nectar stock solutions for

analyses via UPLC-qTof-MS/MS was then adjusted to differences in the volume of extracted

nectar among plant individuals. Hence, the volume of analysed nectar was standardized for

these analyses.

For UHPLC-ESI-qTof-MS/MS analyses, 2 µL of extract were injected at 30 °C into a UHPLC

(Nexara, Shimadzu, DE) containing a C_{18} Column (100 × 2.1 mm, 1.7 µm, Luna Omega Polar,

Phenomenx, DE). The solvent system was composed of (A) water with 0.1% formic acid, and

(B) acetonitrile with 0,1% formic acid (all LC-grade, Carl Roth, DE). The gradient program

started at a flow rate of 0.3 mL min^{-1} with 5 % B, was then raised over 20 min to 95 % B, and

finally kept at 95 % B for an additional 5 min. Afterwards, the column was re-equilibrated at 5%

B for 6 min. Chromatographically separated analytes entered a qTOF-MS/MS (x500R, Sciex,

DE) with the following parameters: ion source gas 45 psi (nitrogen), curtain gas 35 psi

(nitrogen), CAD gas 7 psi (nitrogen), electrospray ion source temperature 500 °C, ion spray

voltage 5500 V. The MS acquisitions parameters were: MS1 scans at 100-1000 m/z with a

data-dependent MS2 trigger having a collision energy (CID) of 40 eV and a scan range of 50-

1000 m/z. The acquired data were then converted to .mzML format and further processed in

mzMine (Schmid *et al.*, 2023). Metabolic features with a maximum m/z difference of 0.001 amu

and maximum retention time difference of 0.1 min were bucketed and treated as identical

features in the following analysis. For feature annotation, the acquired MS/MS spectra were

compared against public mass spectral libraries from Global Natural Products Social Network

(GNPS) (Wang *et al.*, 2016) and the MassBank (MassBank consortium and its contributors,

2024) using the NIST search algorithm (cosine similarity of 0.6) and the MassBank formula

(cosine similarity of 0.5). Features that remained unidentified were further annotated using

COSMIC and CSI:FingerID according to the SIRIUS software package (Hoffmann *et al.*, 2022).

2.3.6   Statistical analyses

All statistical analyses were implemented in R 4.4.1 (R Development Core Team, 2024). Plant

stress indicators, spatial flower traits, flower colour traits, and floral rewards were analysed as

responses in (generalized) linear mixed effects models - (G)LMMs, which included the aphid

infestation treatment, plant sex and the interaction among both variables as predictors. Highly

correlated traits were combined into a single response and analysed within one model. These

models included the different types of summarized response traits as a factorial co-predictor,

for which all possible interactions with sex and aphid infestation treatment were tested

according to the scheme *response at plant individual level (level of summary) ~ cofactor:* leaf

chlorophyll content (on day 0 and 40 of the experiment) ~ time point; leaf epidermal light

absorption by flavonoids (on day 0 and 40 of the experiment) ~ time point; number aphids

(belonging to the size category small, medium and large) ~ aphid size category; flower number

(on day 10, 20 and 40 of the experiment) ~ time point; dry biomass (of a single flower, all

flowers, and flowering stems) ~ biomass component; display area (of a single petal limb, all

petal limbs, and the full corolla expansion) ~ flower component; flower light reflection (in the

green, blue and UV-A range) ~ colour quality. The remaining traits (nectar volume) were

acquired across several flowers per plant individual and analysed in this high resolution (i.e.,

not based on means). Consequently, all models had repeated measurements on the plant

individual level and thus included a random effect for the plant individual. All models were

checked for violations of their underlying assumptions using tests and plots from the R-

package "DHARMa" (Hartig, 2022) and were adjusted with regard to error family, dispersion

or zero-inflation, if necessary. An overview of all model parameters is provided in *Table 1*.

Based on these models, we calculated type III ANOVA tables (based on Wald- χ^2 tests) while

setting all contrasts to sum-zero coding using the R-package "car" (Fox and Weisberg, 2019).

If factorial predictors were involved in significant interactions, we additionally calculated post-

hoc contrasts among the estimated means of their levels within levels of other factors involved

in the respective interaction (R-package "emmeans"; Lenth, 2024)).

The binary choice tests were analysed with two models testing (i) preferences for a specific

aphid infestation treatment within sexes (assays: control female *versus* aphid-infested female,

control male *versus* aphid-infested male); and (ii) preferences for a specific plant sex within

aphid infestation treatments (assays: control female *versus* control male, aphid infested female

versus aphid-infested male). Model i / ii included the behavioural parameter in control relative

to aphid infested (i) or female relative to male (ii) flowers as binomial response and the flowers

sexes (i) or the flowers aphid infestation treatments (ii) as predictors. Further predictors

included in both models were the behavioural parameter type (visits, feeding attempts, time;

all acquired in the same moth and trial), and its interaction with the previously mentioned

model-specific predictor. In addition, both models included the sex of the moth as a cofactor

for which only additive effects were tested, as sample sizes were not large enough to reliably

test for all possible interactions. Moth individual and trial were set as random effects *(Table 1)*.

Model checking, adjustment and downstream analyses with type III ANOVA were implemented

as described above. Post-hoc contrast (R-package "emmeans"; Lenth, 2024) were used to test

for a significant deviation of the binary outcome from the intercept of 0.5 (i.e., preference or

avoidance) within the different plant sex × behavioural type combinations (i) or aphid infestation

treatment × behavioural type combinations (ii).

The statistical approaches used for the analyses of the metabolome are identical for floral

nectar and scent. In a first step, we performed a permutational analyses of variance

(PERMANOVA) using the Bray-Curtis dissimilarities of the full metabolome datasets (n = 41

features in scent, n = 4,385 features in nectar assessed via UPLC-qTof-MS/MS) as the

response distance matrix. We included aphid infestation treatment, sex, and their interactions

as predictors and ran the models with 100,000 permutations (R-package "vegan" (Oksanen *et*

al., 2024)). As previously implemented tests for multivariate homogeneity of group dispersions

in the distance matrices among all treatment levels yielded no deviation for all datasets, the

predictor effects assessed via PERMANOVA are based on differences in group centroids

rather than differences in group dispersions (Anderson and Walsh, 2013). The multivariate

group separations among plant infestation treatments and sexes were illustrated with Non-

metric Multidimensional Scaling (NMDS) plots (R-package "vegan" (Oksanen *et al.*, 2024)). To

identify the compounds responsible for this separation, we used an approach that is

comparable to a volcano plot (Cui and Churchill, 2003) and simultaneously allows the statistical

assessment of interaction effects among plant sex and aphid infestation treatment. Instead of

applying two-sided significance tests (t or u-tests), we entered all metabolites as responses

into a linear model loop to test for the main and interactive effects of plant sex and aphid

infestation treatment. Each metabolite ran with three alternative models as untransformed,

squareroot-transfromed and log-transformed response. To check for violations of model

assumptions and select the best of the three alternative models for each metabolite

accordingly, we extracted the results from the Kolmogorov-Smirnov test for uniformity

(normality of residuals) and Levene's test for homogeneity of variance, which are provided for

targeted model optimization in the R-package "DHARMa" (Hartig, 2022). If model assumptions

were violated in all three alternative models, the respective metabolite was not further

analysed. The best fit models for the remaining metabolites were further analysed with type III

ANOVAs based on F-tests (R-package "car" (Fox and Weisberg, 2019)) and the calculation of

post-hoc contrasts (R-package "emmeans" (Lenth, 2024)) as described in the previous section.

P-values were corrected for inflation of alpha errors by multiple testing (false discovery rate:

FDR) according to (Benjamini and Hochberg, 1995). If a metabolites p-value in a given group

comparison ($F_{C\&A}/M_{C\&A}$, $T_{F\&M}/C_{F\&M}$, FC/FT, MC/MT, FC/MC, FT/MT) was significant and the

corresponding $\log_2$-fold change in the intensity of this metabolite was either ≤ -1 (halving of

intensity in the first relative to the second listed group) or ≥ 1 (doubling of intensity in the first

relative to the second listed group) it was considered to be modulated.

3. Results

4.1 Plant stress indicators

After 40 days of inoculation with *B. lychnidis*, aphid load in terms of weight and number was equally high in female and male *S. latifolia* plants *(Table 1; weight: females = 0.60 +/- 0.12, males = 0.53 +/- 0.12, p>0.05, F_{2DF}= 0.42; number: females = 566 +/- 169, males = 484 +/- 146, p>0.05, F_{2DF}= 0.95)*. Leaf dry mass per plant individual, leaf chlorophyll content and absorption by epidermal flavonoids were not affected by the aphid infestation treatment but significantly higher in females than males in both, control and aphid infested plants *(Table 1, leaf number: females = 5.33 +/- 0.56, males = 3.50 +/- 0.56, p<0.05, F_{2DF}= 5.28; chlorophyll content: females = 27.5 +/- 1.37, males = 22.9 +/- 0.91, p<0.001, χ^2_{2DF}= 14.37; absorption by epidermal flavonoids: females = 0.70 +/- 0.03, males = 0.62 +/- 0.03, p< 0.05, χ^2_{2DF}= 4.80)*. Moreover, leaf chlorophyll contents decreased from 35.9 +/- 1.01 to 17.6 +/- 0.80 nmol cm^{-2} *(p<0.001, χ^2_{2DF}= 60.34)*, whereas absorption by epidermal flavonoids increased from 0.61 to 0.71 *(p<0.001, χ^2_{2DF}= 49.47)* from the beginning towards the end of the experiment, regardless of sex and aphid infestation treatment *(Table 1*, column cofactor*)*.

3.1 Effects of plant sex and aphid infestation on pollinator behaviour

A subset of our binary choice assays tested behavioural moth responses to single flowers from control versus aphid-infested plants, and whether these responses varied between male and female plants. We detected significant preferences for flowers from control plants over aphid-infested ones across all behavioural parameters in females, but not in males *(Figure 3A; z-ratio female plants C-A: 2.11* [visits], 2.29* [feeding attempts], 3.54*** [time spent]; z-ratio male plants C-A: 0.23^{ns} [visits], 0.05^{ns} [feeding attempts], 0.93^{ns} [time spent])*. Consequently, the effect of plant sex on preference for control plants was highly significant *(Table 1, p< 0.001, χ^2_{2DF}= 101.20)*. In addition, we detected a marginally significant main effect of pollinator sex *(Table 1, p<0.06, χ^2_{2DF}= 3.72)*, with only female but not male *H. bicruris* moths showing a significant preference for control over aphid-infested *S. latifolia* plants *(z-ratio female moth C-A: 2.32*, z-ratio male moth C-A: -0.03^{ns})*.

Another subset of our binary choice assays tested whether moths prefer single flowers from one plant sex over the other and whether this preference is modified by the aphid infestation treatment. The statistical analysis yielded no significant preferences of the moths for either plant sex *(Figure 3B; z-ratio control plants F-M: -0.78^{ns} [visits], -0.76^{ns} [feeding attempts], -0.59^{ns} [time spent]; z-ratio aphid infested plants F-M: -0.45^{ns} [visits], -0.66^{ns} [feeding attempts],*

528   *-0.55^{ns} [time spent])*. Neither the aphid infestation treatment of the plants nor moth sex shaped

529   the choice of plant sexes *(Table 1)*.

3.2 Effects of plant sex and aphid infestation on visual flower traits

Flower number was significantly shaped by the three-way interaction sex × infestation

treatment × and infestation time *(Table 1, Figure 4A, p< 0.05, χ^2_{2DF}= 8.88)*. Post-hoc

comparisons revealed that female plants suffered both earlier and stronger declines in flower

number upon aphid infestation as compared to males *(female t-ratio C-A: 2.05^{ns} [10 days],*

4.25*** [20 days], 9.26*** [40 days]; male t-ratio C-A: 0.40^{ns} [10 days], 1.36^{ns} [20 days], 4.74***

[40 days]). Correspondingly, females produced significantly fewer flowers than males only in

aphid infested plants and the magnitude of this effect increased over time *(control t-ratio F-M:*

-2.14^{ns} [10 days], -1.41^{ns} [20 days], -1.26^{ns} [40 days]; infestation t-ratio F-M: -3.84** [10 days],

-4.31*** [20 days], -5.31*** [40 days]). Reproductive biomass depended on the three-way

interaction sex × infestation treatment × component *(Table 1, Figure 4B, p< 0.001, χ^2_{2DF}=*

16.86). Post-hoc comparisons revealed that the component single flower mass was not

affected by the infestation treatment *(t-ratio C-A: 1.17^{ns} [female], 1.14^{ns} [male])* but generally

higher in females than males *(t-ratio F-M: 6.22*** [control], 5.76*** [infested])*. In contrast, the

component total flower mass declined severely under aphid infestation, specifically in females

(t-ratio C-A: 12.89*** [female], 7.26*** [male]) and was accordingly higher in females only in

controls but not under aphid infestation *(t-ratio F-M: 5.22*** [control], -0.41^{ns} [infested])*. The

mass component of flowering stems was not affected by aphid infestation *(t-ratio C-A: 2.73$^{n.s}$*

[female], 0.86$^{n.s}$ [male]), and plant sex *(t-ratio F-M: 2.56^{ns} [control], 0.68^{ns} [infested])*. Flower

light reflectance *(Table 1, Figure 4C)* was significantly shaped by the two two-way interactions

sex × aphid infestation treatment *(p< 0.05, χ^2_{1DF}= 4.91)* and aphid infestation treatment × colour

type *(p< 0.05, χ^2_{2DF}= 8.27)*. Post-hoc comparisons showed that the aphid-infestation treatment

reduced floral light reflectance in the green and blue spectrum only in male, but not in female

plants *(female t-ratio C-A: 1.39^{ns} [green], 1.69$^{ns.}$ [blue], 0.11^{ns} [UV]; male t-ratio C-A: 3.00**

[green], 3.22* [blue], 2.09^{ns} [UV]). Flower area was neither affected by plant sex, aphid

infestation treatment or their interaction *(Table 1)*.

3.1 Effects of plant sex and aphid infestation on the composition of floral scent

We detected a total of 41 VOCs in the floral headspace of *S. latifolia*, with values normalized

per flower. Of these 35 could be putatively identified via matches of RIs and mass spectra with

databases *(Supplementary Table S1A)*. A large fraction of these VOCs have been previously

described in *S. latifolia* (Dötterl *et al.*, 2005; Mamadalieva *et al.*, 2014) and of these, 16 have

been shown to elicit antennal and/or behavioural responses in *H. bicruris* (Dötterl *et al.*, 2006)*.

The PERMANOVA detected a marginally significant main effect of aphid infestation treatment

on the composition of these floral VOCs *(Table 1, p < 0.09, F_{1DF} = 2.41, R^2 = 0.13)* but no

divergence among sexes or interactive effects. This is in accordance with both, the group

dispersion visualized in the NMDS ordination plot *(Figure 5A)* and the outcomes of the model

loop *(Figure 5B, Supplementary Table S1A)*. Without FDR correction, the loop identified eight

compounds shaped by the main effect of treatment with reduced abundance in aphid infested

as compared to control plants. These included five compounds known to elicit behavioural

responses in *H. bicruris* according to Dötterl *et al.* (2006) -benzaldehyde, trans-β-ocimene, lilac

aldehyde A, lilac aldehyde B+C, and lilac alcohol D- as well as two of their degradation

products, lilac alcohol formate A and lilac alcohol formate B. One compound which was not

tested for behavioural effects on *H. bicruris* so far - phenethyl benzoate - was significantly

modulated by the interaction plant sex × aphid infestation treatment and decreased under

aphid infestation in female plants but not in males. When accounting for inflation of alpha

errors, all these effects were non-significant.

3.2 Effects of plant sex and aphid infestation on nectar traits

Nectar volume was unaffected by the aphid infestation treatment but considerably shaped by

plant sex with female flowers featuring higher nectar amounts *(Table 1, Figure 6A, p< 0.001,*

χ^2_{1DF}= 46.10). The untargeted metabolome analyses of a standardized nectar volume per plant

via UPLC-qTof-MS/MS yielded abundance data for 4,385 features. Of these, 761 features were

tentatively identified via matches of their mass spectra with databases or via cheminformatics

according (Hoffmann *et al.*, 2022), and an additional 71 could be assigned to a compound

class *(Supplementary Table 1B)*. This is the first detailed description of nectar chemistry in the

ecological model system *S. latifolia.* The species' nectar contained various nutritious

metabolites, including sugars (D-glucose, D-fructose, sucrose, melezitose, stachyose,

nystose, mannobiose, palatinose), proteinogenic and non-proteinogenic amino acids (e.g.,

proline, glutamic acid, tyrosine ornithine) and numerous oligopeptides, nucleosides

(adenosine, adenosine monophosphate, a guanosine derivate), and multiple lipids (e.g., fatty

alkyls; fatty acids, amides and alcohols; ceramides; glycerophosphoethanolamines,

sphingolipids; prostaglandins). Its diverse secondary metabolites comprised vitamins (e.g.,

calcitriol, riboflavin), flavonoids (e.g., saponarin, vitexin-2"-O-rhamnoside), terpenoids (e.g.,

myrtenol, perillyl alcohol, raddeanin D, pisumoside A), as well as fatty amides and amines

(oleamide -a neuroactive compound [Mendelson and Basile, 2001] which has so far not been

described in nectar, tyramine, homospermidine). Most of these secondary compounds are

likely derived from plant metabolism, but several have a known fungal or bacterial origin and

thus likely derive from the nectar microbiome (e.g., oleandomycin, enniatin B, asperiamide B,

N-acetylglucosamine [chitin monomer]).

The PERMANOVA detected a highly significant effect of plant sex *(Table 1, p < 0.001, F_{1DF} =*

10.65, R² = 0.39), but no effects of infestation treatment or the interaction sex × infestation

treatment on the composition of the metabolome in a standardized volume of nectar*.* The group

dispersions visualized in ordination plots via NMDS illustrate a clear differentiation of the nectar

metabolome among female and male plants and additionally indicate different responses to

the aphid infestation treatment in males and females in some features *(Figure 6B)*. The results

from the linear model loop further support this observation *(Figure 6C-F, Table 2,*

Supplementary Table S1B). When accounting (not accounting) for multiple testing, a total of

219 (3) features had significantly and remarkably ($\log_2$ fold change >1) higher abundance in

females than males, including six features that were absent in the latter. These did not include

primary nutritional compounds such as sugars or free amino acids, but seven oligopeptides,

nine fatty amides, six terpenoids (including three triterpenoid steroids), a glycerophosphate, a

flavonoid and the microbial compound N-acetylglucosamine [chitin monomer]. The remaining

discriminatory features, including those unique to female nectar, could not be identified *(Figure*

6C, Table 2, Supplementary Table 1B). On the other hand, 45 (15) features had significant and

remarkably ($\log_2$ fold change <-1) higher abundances in males than females with 20 of them

being absent in the latter. These included the free proteinogenic amino acid tyrosine (unique),

N-(3-Methylbutanoyl)valine (unique), two oligopeptides (one unique), a macrocyclic

despipeptide of probable microbial origin (unique), two flavonoid glycosides (one unique), a

glycerosphingolipid, and a guanosine derivate (unique, highly significant). The aphid

infestation treatment had clearly weaker effects on the composition of plant nectar. Plants

showed increased abundances of 0 (10) features including the tetrasaccharide stachyose; and

decreased abundances of 0 (18) features including the amino acid phenylalanine, a

disaccharidester, and the isoflavonoid genistin *(Figure 6D, Table 2, Supplementary Table S1B)*.

The interaction plant sex × aphid infestation treatment shaped the abundance of 21 (7) features

(Figure 6E&F; please note that both plots display the same metabolic features with different

group comparisons; Table 2; Supplementary Table S1B). These interactions had seven major

effect directions, which are listed as labels at the top of both plots. The most prominent

interaction pattern was significantly and markedly higher metabolite abundance in females

compared to males under control conditions, but not under aphid infestation in 16 (0) features,

which included a nucleoside analogue. The second most prominent interaction pattern

correlated with moth behaviour. In total 1(5) features had significantly and markedly reduced

abundances under aphid infestation in females, but not in males. We analysed the MS² spectra

for these six features in detail. Most of them did not yield high confidence matches or similarity

scores to known compounds, suggesting these metabolites are not previously reported natural

products. Our detailed analyses provided chemical class annotations and further details on

structural features allowing to discuss their putative function in the context of plant-pollinator

interactions (*Supplementary Table S1C*). These compounds include a pyrrolidine derivative

($C_{17}H_{18}N$, m/z 237.021) with aromatic substituents (m/z = 70.0648 → pyrrolidinegroup, m/z =

77.0382 → phenylmethylgroup, m/z = 91.0539 → phenylgroup), a xanthine derivative with

additional substituents ($C_{14}H_{14}N_4O_5S$, m/z 351.07), a benzenesulfonyl compound with a

sugar-like substituent ($C_{14}H_{21}NO_7S$, m/z 351.07), two ceramides of which one bared multiple

amino substituents ($C_{32}H_{65}NO_4$, m/z 528.3914; $C_{23}H_{51}N_5O_4$, m/z 484.3645) and a flavonoid

($C_{31}H_{24}O_9$, m/z 541.1492).

4. Discussion

Building on the pronounced sexual differentiation in floral traits of the dioecious plant *Silene latifolia*, we examined how controlled infestation by the oligophagous aphid *Brachycaudus lychnidis* affects interactions with the specialist moth pollinator *Hadena bicruris*. Contrary to classical expectations from sexual selection theory, female plants were not outcompeted by males in floral attractiveness under control conditions. Instead, they exhibited high plasticity in floral trait expression and produced diverse, potentially bioactive nectar metabolites, some of which may specifically target female pollinator behaviour. Under aphid infestation, however, female floral attractiveness declined more strongly than in males, suggesting sex-specific defence strategies and highlighting the potential role of indirect defences in stabilizing plant–pollinator–herbivore interactions.

4.1 Female *Silene latifolia* plants challenge expectations: no evidence of reduced floral attractiveness compared to males

Bateman's theory of sexual selection predicts that, relative to males, female reproductive success is more limited by available resources and therefore does not increase with the number of mates, leading to less attractive flowers that serve as a resource exploited and competed for by males (Bateman, 1948). In the present study, we found no evidence for these predictions.

Hadena bicruris moths did not generally prefer single flowers of either sex (*Figure 4B*, *Table 1* - pollinator choice F-M). Female and male plants exhibited no general differences in flower numbers (*Figure 4A* - control conditions), flower sizes (*Table 1* - petal area), and flower colours (*Figure 4C*). Female flowers even featured higher nectar volumes as compared to males (*Figure 6A*) and major nectar sugar constituents showed no sex-specific differences (*Figure 6C*, *Table 3*, *Supplementary Table S1B*). Instead, we observed higher abundances of over 200 other primary and secondary metabolites in females than males – some of which likely shape the nutritional and pharmacological value of nectar for moth pollinators. These included seven oligopeptides, partially enriched in branched-chain amino acids, which may serve as a rapid source of NADH, $FADH_2$, and acetyl-CoA for ATP production, supporting energy demands during flight or allocation to egg production in female moths (Levin *et al.*, 2017; Cavigliasso *et al.*, 2023); nine fatty amides potentially serving as energy stores, metabolic fuel or behavioural modulators (Ranger *et al.*, 2005; Yoshinaga, 2016; Levin *et al.*, 2017); six terpenoids (of which two were steroidal glycosides) that may provide selective feeding or oviposition cues (González-Coloma *et al.*, 2011; Boncan *et al.*, 2020); a glycerophospholipid that may support the maintenance and generation of somatic tissue due to its central role in rapid membrane

synthesis (Toprak *et al.*, 2020); and a flavonoid which may act as health promotor in moth
pollinators (Palmer-Young *et al.*, 2019). The function and further identification of the numerous
female biased metabolites will be the subject of future studies. As only female plants bear the
cost of the interaction with *H. bicruris* (fruit predation), and only female moths impose this cost,
there is great potential for the discovery of nectar metabolites interfering with reproduction in
female moths. Male *S. latifolia* plants, on the other hand, exhibited higher abundances of over
40 compounds, including several from the same chemical classes and with similar putative
functions as those dominant in female nectar (BCAA-rich oligopeptides, flavonoid glycosides,
membrane building lipids such as ceramides and glycosphingolipids). Most strikingly, tyrosine
and a derivate of guanosine were unique to male nectar. Both compounds can be considered
essential for the nutrition of moth pollinators that rely exclusively on nectar as a nitrogen source
and may enhance its taste and memorability accordingly (Nepi, 2014). In addition, tyrosine is
a precursor for the synthesis of the neurotransmitters tyramine and dopamine, which shape
numerous insect behavioural traits like e.g., motivation and learning (Barberis *et al.*, 2023). To
the best of our knowledge, our study reports one of the most extensive cases of sexual
differentiation in the composition of nectar for a dioecious plant to date (see also (Nepi *et al.*,
2001; Dötterl *et al.*, 2014)). Since *H. bicruris* showed no preference for either plant sex at the
single-flower level (*Figure 4B*, *Table 1* - pollinator choice F-M), we conclude that these striking
differences complement each other functionally to stabilize a dioecious system relying on
frequent visits to both sexes, with plants promoting this by offering essential nectar resources
in only one sex.

Overall, our study offers a more nuanced view of female floral traits than predicted by
Bateman's principle. A potential criticism is that females in our greenhouse setting did not
exhibit lower floral attractiveness as they were not pollinated and hence did not invest in seeds,
which aligns with long-term personal observations of visual flower traits in *S. latifolia*. However,
this very aspect underscores the limitations of Bateman's theory as it supports that females
exhibit adaptive plasticity in floral traits, with attractiveness decreasing only as pollination
success increases. High attractiveness of unpollinated female plants may promote female
mate competition, post-pollination mate choice in females [84], multiple paternity in fruits, and
ultimately greater offspring diversity and fitness [85]. This would be in line with the theory of
flexible sex role behaviour [83], which -similar to Bateman's principle- originates from zoology
and may be applicable to understanding reproductive strategies in dioecious plants. The
substantial among-population variation in sexual dimorphism observed in *S. latifolia* in previous
studies (Schrieber *et al.*, 2017, 2019, 2021) further contradicts the assumption of a fixed
evolutionary trajectory, instead pointing toward locally driven and context-dependent trait
evolution.

4.2 Aphid infestation reduces female floral attractiveness via sex-specific trait changes

In accordance with our hypothesis, infestation with *B. lychnidis* aphids affected a wide range

of traits in *S. latifolia* flowers, as well as the behavioural responses of *H. bicruris* moth

pollinators. Although most of the observed effects clearly depended on plant sex, we begin

with a general discussion of aphid-induced changes, as these have, to our knowledge, not

been previously reported.

Aphid infestation reduced flower numbers (more strongly in females than males, *Figure 4A*),

and reduced flower light reflection in the blue and green range (only in males, *Figure 4C*).

These adverse effects likely result from both resource depletion and the reallocation of

resources toward plant defence (Twayana *et al.*, 2022), and may substantially limit plant

visibility to pollinators (Barbot *et al.*, 2022). Aphid infestation also moderately reduced the

emission of five key volatile organic compounds (VOCs) known to elicit antennal and feeding

responses in *H. bicruris* (Dötterl *et al.*, 2006), at the level of individual flowers (*Figure 5B*, *Table*

2). Nectar volume was not lower in aphid-infested plants (*Figure 6A*), likely because they

produced fewer flowers and could consequently invest more nectar in each one. While the

major sugar constituents remained largely unaffected (*Figure 6D*, *Table 3*), stachyose levels

increased in both sexes, suggesting that plants load their phloem with this poorly digestible

tetrasaccharide to impair aphid performance (Twayana *et al.*, 2022), while simultaneously

ensuring pollination by enzymatically breaking it down into simpler sugars such as fructose,

glucose, or sucrose in nectar. Conversely, levels of phenylalanine, a nucleoside analogue, a

phenolic acid glycoside, and an isoflavonoid were reduced. These changes may result either

directly from phloem-feeding-induced resource depletion or reflect a passive carry-over of

defensive phloem sap modifications targeting aphids (Twayana *et al.*, 2022). Although most of

these alterations affected both sexes and would be expected to generally reduce pollinator

attraction, this was observed only in one sex. It is important to note that our choice assays did

not account for the markedly reduced flower numbers at the whole-plant level, which likely

amplify the effects of floral chemical changes and thus have a stronger negative impact on

pollinator attraction.

At the single-flower level, *H. bicruris* preferred flowers from control plants over aphid-infested

ones only in *S. latifolia* females, but not males (*Figure 3*A). These preferences were also

dependent on moth sex, being primarily exhibited by female *H. bicruris* (*Table 1*). This

behaviour coincided with a reduction of the floral VOC phenylethyl benzoate - a compound not

previously described in *S. latifolia* but found in other moth-pollinated species [84] - which was

observed only in aphid-infested female plants, but not in males (*Figure 5B*, *Table 2*).

Additionally, six metabolic features in the nectar of *S. latifolia* were reduced by aphid infestation

only in female plants (*Figure 6F*, *Table 3*). Four of them simultaneously had higher abundances

in female than male plants only under control but not under aphid infestation conditions

(*Supplementary table S1B*). Although these effects were only observed before FDR correction,

the corresponding effect direction in pollinator behaviour suggests biological relevance. None

of these metabolites belonged to the major nutritional nectar constituents (sugars, amino acids)

or could be clearly identified as a previously described natural product. Our detailed

chemoinformatic analysis of MS^2 spectra elucidated compound classes and further structural

components indicating a potential role of these metabolites as selective feeding cues or

behavioural modulators for female *H. bicruris*. One candidate compound contains both

pyrrolidine and phenyl substituents, a structural motif found in several synthetic psychoactive

compounds that inhibit monoamine reuptake transporters, particularly dopamine, and are

associated with stimulant activity (Meyer *et al.*, 2013). Another notable candidate is a xanthine

derivative, a class of compounds widely documented to enhance learning, memory, and

arousal in pollinators (Mustard, 2020). The benzenesulfonyl compound containing amino and

methoxy sugar-like moieties, the two ceramides and the flavonoid may possibly contribute to

specific recognition by pollinators via contact cues. A detailed structural characterization of

these novel natural products via NMR analysis of larger nectar volumes as well as individual

and combinatorial bioassays are necessary to validate these putative ecological functions.

In summary, our findings support that male plants have more efficient plastic responses to

maintain their floral attractiveness under severe herbivore pressure. This is rather difficult to

explain with higher resource demands for reproduction in females, as our setup prevented

pollination and fruit set. Instead, it points to different defence strategies in males and females,

which cannot be fully disentangled with our setup. Our approach bypassed aphid host

preferences; direct plant defences preventing the early establishment of aphids, the attraction

of aphid parasitoids and predators, and it did not account for resource limitation and

competition among sexes that all occur in the field. Indeed, female plants in some dioecious

species have been shown to be less frequently attacked by herbivores (Sargent and

McKeough, 2022), to investigate more into direct defence (Sargent and McKeough, 2022) and

potentially also into indirect defences (Aranda-Rickert *et al.*, 2021). In fact it has been shown

that *S. latifolia* females in Europe are clearly less frequently attacked by aphids than males

(Wolfe *et al.*, 2004). Hence, they are unlikely to experience aphid infestation as strong as in

our experiment under more natural conditions, which may be one explanation for their higher

susceptibility. Yet, the cause for lower aphid abundances on female plants remains to be

investigated in the future. Sex-specific indirect defences against aphids appear to be a

particularly promising research route here. The seed-predating larvae of *H. bicruris* are

attacked by a broad range of parasitoids and predators (Villacañas de Castro and Hoffmeister,
2020) that -if the plant is involved at all- should be attracted by females only. There is ample
evidence that the evolution of indirect defences against different enemy species and guilds is
tightly coupled via shared plant signalling and VOC synthesis pathways or the attraction of
generalist predators (Kessler and Baldwin, 2001; Heil and McKey, 2003; Rasmann *et al.*,
2005). Finally, since female plants are often reported to exploit resources more rapidly than
males (Jiang *et al.*, 2016; Bürli *et al.*, 2022), which is indicated for *S. latifolia* in our study
through higher leaf biomass and greater chlorophyll content in females (*Table 1*), competition
among sexes in resource limited environments may substantially shift the sex-specific
susceptibility patterns.

5. Conclusion and outlook

Our study provides evidence for multidirectional sexual differentiation of visual and chemical
flower traits and their responses to aphid infestation in a dioecious plant species. It supports
that the expression of sexual dimorphisms in floral attractiveness can be highly plastic
depending on the presence of both, pollinators and herbivores, thereby opening exciting
avenues for future research. Controlled experiments manipulating the presence of both,
pollinators and aphids, ideally under field conditions reflecting natural conditions, while
integrating data collection on plant, pollinator and aphid reproductive success with quantitative
leaf and flower chemistry are a next step in disentangling the sex-specific co-evolution in this
tripartite system. In such future studies, it might be more purposeful to focus on the
complementarity of female and male defence and floral attractiveness traits in stabilizing a
dioecious reproductive system that is dependent on a nursery pollinator and involves several
other parties (aphids, aphid and caterpillar parasitoids and predators), instead of seeking for
the better defended or more attractive sex.

6. Acknowledgements

Tobias Möckel from the FabLab in Kiel developed a customised camera attachment for light
filter switches that was used for the assessment of flower colour composition. Reza Esmaeilnia
Talemi supported the collection of plant nectar for metabolome analyses and Meike Pfeiler
prepared nectar extracts. Stefan Dötterl advised on the establishment of our *Hadena*
laboratory population, and Eric Folz was involved in the maintenance of the aphid and

pollinator laboratory populations. Roman Adler designed vector graphs for the study design
illustrations; and Verena Zajonc and Katharina Grosser provided valuable technical support.

7. Supplementary data

815 Supplementary Table S1: Summary of floral metabolites in *Silene latifolia*: functional annotation
816 and responses to sex and aphid infestation
817 Supplementary Figure S1: Representative images from the experimental procedure

8. Author contributions:

819 "Conceptualization, K.S.; Methodology, K.S., T.S., E.K, T.D., S.U., and W.B., Validation, K.S.,
820 T.S., E.K, and W.B.; Formal Analysis, K.S.; Investigation, K.Z., E.K., T.D., and T.S.; Resources,
821 A.E.; S.U., T.S., E.K, T.D. and W.B.; Data Curation, K.S.; Writing – Original Draft Preparation,
822 K.S. and K.Z.; Writing – Review & Editing, K.S., E.K, W.B., S.U., H.B., T.D., T.S; Visualization,
823 K.S.; Supervision, K.S., S.U.; Project Administration, K.S.; Funding Acquisition, K.S.

9. Conflict of interest:

825 No conflict of interest is declared

10. Funding

827 This study has been funded by the program for the promotion of young female scientists from
828 the Faculty of Mathematics and Natural Sciences of Kiel University

11. Data availability

830 All data / code supporting this article will be deposited in dryad / zenedo and made publicly
831 available upon editorial acceptance of the manuscript

12. References

Adler LS, Seifert MG, Wink M, Morse GE. 2012. Reliance on pollinators predicts defensive chemistry across tobacco species. Ecology Letters **15**, 1140–1148.

Anderson MJ, Walsh DCI. 2013. PERMANOVA, ANOSIM, and the Mantel test in the face of heterogeneous dispersions: What null hypothesis are you testing? Ecological Monographs **83**, 557–574.

Aranda-Rickert A, Torréns J, Yela NI, Brizuela MM, Di Stilio VS. 2021. Distance Dependent Contribution of Ants to Pollination but Not Defense in a Dioecious, Ambophilous Gymnosperm. Frontiers in Plant Science **12**, 722405.

Avila-Sakar G, Romanow CA. 2012. Divergence in Defence against Herbivores between Males and Females of Dioecious Plant Species. International Journal of Evolutionary Biology **2012**, 897157.

Barberis M, Calabrese D, Galloni M, Nepi M. 2023. Secondary Metabolites in Nectar-Mediated Plant-Pollinator Relationships. Plants **12**, 550.

Barbot E, Dufaÿ M, Tonnabel J, Godé C, De Cauwer I. 2022. On the function of flower number: disentangling fertility from pollinator-mediated selection. Proceedings of the Royal Society B: Biological Sciences **289**, 20221987.

Barrett SCH, Hough J. 2013. Sexual dimorphism in flowering plants. Journal of Experimental Botany **64**, 67–82.

Bateman AJ. 1948. Intra-sexual selection in Drosophila. Heredity **2**, 349–368.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B (Methodological) **57**, 289–300.

Berkum PM, Giorgi FD, Rothe B, Durka W, Gershenzon J, Roscher C, Unsicker SB. 2024. Selection strengthens the relationship between plant diversity and the metabolic profile of Plantago lanceolata. bioRxiv.

Blair AC, Wolfe LM. 2004. The evolution of an invasive plant: an experimental study with Silene latifolia. Ecology **85**, 3035–3042.

Boncan DAT, Tsang SSK, Li C, Lee IHT, Lam H-M, Chan T-F, Hui JHL. 2020. Terpenes and Terpenoids in Plants: Interactions with Environment and Insects. International Journal of Molecular Sciences **21**, 7382.

Bopp S, Gottsberger G. 2004. Importance of Silene latifolia ssp. alba and S. dioica (Caryophyllaceae) as host plants of the parasitic pollinator Hadena bicruris (Lepidoptera, Noctuidae). Oikos **105**, 221–228.

Borghi M, Fernie AR, Schiestl FP, Bouwmeester HJ. 2017. The Sexual Advantage of Looking, Smelling, and Tasting Good: The Metabolic Network that Produces Signals for Pollinators. Trends in Plant Science **22**, 338–350.

Bürli S, Pannell JR, Tonnabel J. 2022. Environmental variation in sex ratios and sexual
dimorphism in three wind-pollinated dioecious plant species. Oikos **2022**, e08651.

Cavigliasso F, Savary L, Spangenberg JE, Gallart-Ayala H, Ivanisevic J, Kawecki TJ.
2023. Experimental evolution of metabolism under nutrient restriction: enhanced amino acid
catabolism and a key role of branched-chain amino acids. Evolution Letters **7**, 273–284.

Charlesworth D. 1993. Why are Unisexual Flowers Associated with Wind Pollination and
Unspecialized Pollinators? The American Naturalist **141**, 481–490.

Cornelissen T, Stiling P. 2005. Sex-biased herbivory: a meta-analysis of the effects of gender
on plant-herbivore interactions. Oikos **111**, 488–500.

Cui X, Churchill GA. 2003. Statistical tests for differential expression in cDNA microarray
experiments. Genome Biology **4**, 210.

Dedryver C-A, Le Ralec A, Fabre F. 2010. The conflicting relationships between aphids and
men: A review of aphid damage and control strategies. Comptes Rendus Biologies **333**, 539–
553.

**De-la-Cruz IM, Batsleer F, Bonte D, Diller C, Hytönen T, Muola A, Osorio S, Posé D,
Vandegehuchte ML, Stenberg JA**. 2022. Evolutionary Ecology of Plant-Arthropod
Interactions in Light of the "Omics" Sciences: A Broad Guide. Frontiers in Plant Science **13**.

Delph LF, Arntz AM, Scotti-Saintagne C, Scotti I. 2010. The Genomic Architecture of Sexual
Dimorphism in the Dioecious Plant Silene Latifolia. Evolution **64**, 2873–2886.

Dötterl S, Glück U, Jürgens A, Woodring J, Aas G. 2014. Floral Reward, Advertisement and
Attractiveness to Honey Bees in Dioecious Salix caprea. PLOS ONE **9**, e93421.

Dötterl S, Jürgens A. 2005. Spatial fragrance patterns in flowers of Silene latifolia: Lilac
compounds as olfactory nectar guides? Plant Systematics and Evolution **255**, 99–109.

Dötterl S, Jurgens A, Seifert K, Laube T, Weissbecker B, Schutz S. 2006. Nursery
pollination by a moth in Silene latifolia: the role of odours in eliciting antennal and behavioural
responses. New Phytologist **169**, 707–718.

Dötterl S, Wolfe LM, Jürgens A. 2005. Qualitative and quantitative analyses of flower scent
in Silene latifolia. Phytochemistry **66**, 203–213.

Dufaÿ M, Anstett M-C. 2003. Conflicts between plants and pollinators that reproduce within
inflorescences: evolutionary variations on a theme. Oikos **100**, 3–14.

Elzinga JA, Biere A, Harvey A. 2002. The rearing of the gregarious koinobiont endoparasitoid
Microplitis tristis (Hymenoptera: Braconidae) on its natural host Hadena bicruris (Lepidoptera:
Noctuidae). Proceedings of the Section Experimental and Applied Entomology of the
Netherlands Entomological Society (2002), 109–115.

Fox J, Weisberg S. 2019. *An R Companion to Applied Regression*. SAGE Publications.

Gehring JL, Scoby * Jennifer, Parsons M, Delph LF. 2004. Whole-plant investment in nectar
is greater for males than pollinated females in the dioecious plant Silene latifolia. Evolutionary
Ecology Research **6**, 1237–1252.

González-Coloma A, López-Balboa C, Santana O, Reina M, Fraga BM. 2011. Triterpene-
based plant defenses. Phytochemistry Reviews **10**, 245–260.

Hartig F. 2022. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. R package version 0.4.6 **https://CRAN.R-project.org/package=DHARMa**.

Heil M, McKey D. 2003. Protective Ant-Plant Interactions as Model Systems in Ecological and Evolutionary Research. Annual Review of Ecology, Evolution, and Systematics **34**, 425–553.

Hoffmann MA, Nothias L-F, Ludwig M, Fleischauer M, Gentry EC, Witting M, Dorrestein PC, Dührkop K, Böcker S. 2022. High-confidence structural annotation of metabolites absent from spectral libraries. Nature Biotechnology **40**, 411–421.

https://influentialpoints.com/Gallery/Brachycaudus_lychnidis_Campion_aphids.htm. 2024. *in press.*

Ito K, Suzuki MF, Mochizuki K. 2021. Evolution of honest reward signal in flowers. Proceedings of the Royal Society B: Biological Sciences **288**, 20202848.

Jessen M-T, Auge H, Harpole WS, Hautier Y, Eskelinen A. 2022. Grazing and light modify Silene latifolia responses to nutrients and future climate. PLOS ONE **17**, e0276789.

Jiang H, Zhang S, Lei Y, Xu G, Zhang D. 2016. Alternative Growth and Defensive Strategies Reveal Potential and Gender Specific Trade-Offs in Dioecious Plants Salix paraplesia to Nutrient Availability. Frontiers in Plant Science **7**, 1064.

Johnsen S, Kelber A, Warrant E, Sweeney AM, Widder EA, Lee RL, Hernández-Andrés J. 2006. Crepuscular and nocturnal illumination and its effects on color perception by the nocturnal hawkmoth Deilephila elpenor. Journal of Experimental Biology **209**, 789–800.

Johnson MTJ, Campbell SA, Barrett SCH. 2015. Evolutionary Interactions Between Plant Reproduction and Defense Against Herbivores. Annual Review of Ecology, Evolution, and Systematics **46**, 191–213.

Jürgens A, Witt T, Gottsberger G. 1996. Reproduction and Pollination in Central European Populations of Silene and Saponaria Species. Botanica Acta **109**, 316–324.

Käfer J, Marais GAB, Pannell JR. 2017. On the rarity of dioecy in flowering plants. Molecular Ecology **26**, 1225–1241.

Kessler A, Baldwin IT. 2001. Defensive Function of Herbivore-Induced Plant Volatile Emissions in Nature. Science **291**, 2141–2144.

Kessler A, Chautá A. 2020. The ecological consequences of herbivore-induced plant responses on plant–pollinator interactions. Emerging Topics in Life Sciences **4**, 33–43.

Labouche A-M, Bernasconi G. 2010. Male moths provide pollination benefits in the Silene latifolia–Hadena bicruris nursery pollination system. Functional Ecology **24**, 534–544.

Lenth R. 2024. emmeans: Estimated Marginal Means, aka Least-Squares Means. https://CRAN.R-project.org/package=emmeans **R package version 1.10.2.**

Levin E, McCue MD, Davidowitz G. 2017. More than just sugar: allocation of nectar amino acids and fatty acids in a Lepidopteran. Proceedings of the Royal Society B: Biological Sciences **284**, 20162126.

Mamadalieva NZ, Lafont R, Wink M. 2014. Diversity of Secondary Metabolites in the Genus
Silene L. (Caryophyllaceae)—Structures, Distribution, and Biological Properties. Diversity **6**,
415–499.

MassBank consortium and its contributors. 2024. MassBank/MassBank-data: Release
version 2024.11. Zenedo **https://doi.org/10.5281/zenodo.14221628**.

McCall AC, Irwin RE. 2006. Florivory: the intersection of pollination and herbivory. Ecology
Letters **9**, 1351–1365.

Mendelson WB, Basile AS. 2001. The hypnotic actions of the fatty acid amide, oleamide.
Neuropsychopharmacology: Official Publication of the American College of
Neuropsychopharmacology **25**, S36-39.

Meyer MR, Schmitt S, Maurer HH. 2013. Studies on the metabolism and detectability of the
emerging drug of abuse diphenyl-2-pyrrolidinemethanol (D2PM) in rat urine using GC-MS and
LC-HR-MS/MS. Journal of Mass Spectrometry **48**, 243–249.

Moore JC, Pannell JR. 2011. Sexual selection in plants. Current Biology **21**, R176–R182.

Morales-Hojas R. 2017. Molecular ecology of insect pests of agricultural importance: the case
of aphids. Ecological Entomology **42**, 18–27.

Muhlemann JK, Klempien A, Dudareva N. 2014. Floral volatiles: from biosynthesis to
function. Plant, Cell & Environment **37**, 1936–1949.

Muola A, Lucas-Barbosa D, Kessler A. 2022. Editorial: Mechanisms underlying plant-
pollinator-herbivore interactions. Frontiers in Plant Science **13**.

Mustard JA. 2020. Neuroactive nectar: compounds in nectar that interact with neurons.
Arthropod-Plant Interactions **14**, 151–159.

Nalam V, Louis J, Shah J. 2019. Plant defense against aphids, the pest extraordinaire. Plant
Science **279**, 96–107.

Nepi M. 2014. Beyond nectar sweetness: the hidden ecological role of non-protein amino acids
in nectar. Journal of Ecology **102**, 108–115.

Nepi M, Grasso DA, Mancuso S. 2018. Nectar in Plant–Insect Mutualistic Relationships:
From Food Reward to Partner Manipulation. Frontiers in Plant Science **9**, 1063.

Nepi M, Guarnieri M, Pacini E. 2001. Nectar Secretion, Reabsorption, and Sugar
Composition in Male and Female Flowers of Cucurbita pepo. International Journal of Plant
Sciences **162**, 353–358.

Nicolson SW. 2022. Sweet solutions: nectar chemistry and quality. Philosophical Transactions
of the Royal Society B: Biological Sciences **377**, 20210163.

Noriega JA, Hortal J, Azcárate FM, *et al.* 2018. Research trends in ecosystem services
provided by insects. Basic and Applied Ecology **26**, 8–23.

Oksanen J, Blanchet FG, Friendly M, *et al.* 2024. Vegan: Community Ecology Package.
https://CRAN.R-project.org/package=vegan **R package version 2.6-8**.

Palmer-Young EC, Farrell IW, Adler LS, Milano NJ, Egan PA, Junker RR, Irwin RE,
Stevenson PC. 2019. Chemistry of floral rewards: intra- and interspecific variability of nectar
and pollen secondary metabolites across taxa. Ecological Monographs **89**, e01335.

Paul I, Poddar Sarkar M, Bhadoria PBS. 2022. Floral secondary metabolites in context of
biotic and abiotic stress factors. Chemoecology **32**, 49–68.

Peakall R. 2023. Pollination by sexual deception. Current Biology **33**, R489–R496.

R Development Core Team. 2024. R: A Language and Environment for Statistical Computing.
R Foundation for Statistical Computing.

Ranger CM, Winter REK, Rottinghaus GE, Backus EA, Johnson DW. 2005. Mass spectral
characterization of fatty acid amides from alfalfa trichomes and their deterrence against the
potato leafhopper. Phytochemistry **66**, 529–541.

Rasmann S, Köllner TG, Degenhardt J, Hiltpold I, Toepfer S, Kuhlmann U, Gershenzon
J, Turlings TCJ. 2005. Recruitment of entomopathogenic nematodes by insect-damaged
maize roots. Nature **434**, 732–737.

Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, Eliceiri KW. 2017.
ImageJ2: ImageJ for the next generation of scientific image data. BMC Bioinformatics **18**, 529.

Santamaría L, Rodríguez-Gironés MA. 2015. Are flowers red in teeth and claw? Exploitation
barriers and the antagonist nature of mutualisms. Evolutionary Ecology **29**, 311–322.

Sargent RD, McKeough AD. 2022. New Evidence Suggests No Sex Bias in Herbivory or Plant
Defense. The American Naturalist **200**, 435–447.

Schmid R, Heuckeroth S, Korf A, *et al.* 2023. Integrative analysis of multimodal mass
spectrometry data in MZmine 3. Nature Biotechnology **41**, 447–449.

Schrieber K, Glüsing S, Peters L, Eichert B, Althoff M, Schwarz K, Erfmeier A,
Demetrowitsch T. 2023. Population divergence in heat and drought responses of a coastal
plant: from metabolic phenotypes to plant morphology and growth. (T Obata, Ed.). Journal of
Experimental Botany, erad147.

Schrieber K, Paul SC, Höche LV, Salas AC, Didszun R, Mößnang J, Müller C, Erfmeier A,
Eilers EJ. 2021. Inbreeding in a dioecious plant has sex- and population origin-specific effects
on its interactions with pollinators. (MC Aldrich and Y Joo, Eds.). eLife **10**, e65610.

Schrieber K, Wolf S, Wypior C, Höhlig D, Hensen I, Lachmuth S. 2017. Adaptive and non-
adaptive evolution of trait means and genetic trait correlations for herbivory resistance and
performance in an invasive plant. Oikos **126**, 572–582.

Schrieber K, Wolf S, Wypior C, Höhlig D, Keller SR, Hensen I, Lachmuth S. 2019. Release
from natural enemies mitigates inbreeding depression in native and invasive *Silene latifolia*
populations. Ecology and Evolution **9**, 3564–3576.

Smith RN, Martell AE, Motekaitis RJ. 2004. NIST standard reference database. NIST
Critically Selected Stability Constants of Metal Complexes Database **Ver 2**.

Stamp N. 2003. Out Of The Quagmire Of Plant Defense Hypotheses. The Quarterly Review
of Biology **78**, 23–55.

Tang-Martínez Z. 2012. Repetition of Bateman challenges the paradigm. Proceedings of the
National Academy of Sciences of the United States of America **109**, 11476–11477.

Tew NE, Memmott J, Vaughan IP, Bird S, Stone GN, Potts SG, Baldock KCR. 2021.
Quantifying nectar production by flowering plants in urban and rural landscapes. (I Bartomeus,
Ed.). Journal of Ecology **109**, 1747–1757.

Toprak U, Hegedus D, Doğan C, Güney G. 2020. A journey into the world of insect lipid
metabolism. Archives of Insect Biochemistry and Physiology **104**, e21682.

Troscianko J, Stevens M. 2015. Image calibration and analysis toolbox – a free software suite
for objectively measuring reflectance, colour and pattern. Methods in Ecology and Evolution **6**,
1320–1331.

Trunschke J, Lunau K, Pyke GH, Ren Z-X, Wang H. 2021. Flower Color Evolution and the
Evidence of Pollinator-Mediated Selection. Frontiers in Plant Science **12**.

Twayana M, Girija AM, Mohan V, Shah J. 2022. Phloem: At the center of action in plant
defense against aphids. Journal of Plant Physiology **273**, 153695.

van Putten WF, Elzinga JA, Biere A. 2007. Host Fidelity of the Pollinator Guilds of Silene
dioica and Silene latifolia: Possible Consequences for Sympatric Host Race Differentiation of
a Vectored Plant Disease. International Journal of Plant Sciences **168**, 421–434.

Villacañas de Castro C, Hoffmeister TS. 2020*a*. Friend or foe? A parasitic wasp shifts the
cost/benefit ratio in a nursery pollination system impacting plant fitness. Ecology and Evolution
10, 4220–4232.

Villacañas de Castro C, Hoffmeister TS. 2020*b*. Friend or foe? A parasitic wasp shifts the
cost/benefit ratio in a nursery pollination system impacting plant fitness. Ecology and Evolution
10, 4220–4232.

Wang M, Carver JJ, Phelan VV, *et al.* 2016. Sharing and community curation of mass
spectrometry data with Global Natural Products Social Molecular Networking. Nature
Biotechnology **34**, 828–837.

Wolfe LM, Elzinga JA, Biere A. 2004. Increased susceptibility to enemies following
introduction in the invasive plant Silene latifolia. Ecology Letters **7**, 813–820.

Wong DCJ, Pichersky E, Peakall R. 2023. Many different flowers make a bouquet: Lessons
from specialized metabolite diversity in plant–pollinator interactions. Current Opinion in Plant
Biology **73**, 102332.

Yoshinaga N. 2016. Physiological function and ecological aspects of fatty acid-amino acid
conjugates in insects†. Bioscience, Biotechnology, and Biochemistry **80**, 1274–1282.

1056

Tables

Table 1: *Overview of statistical analyses investigating the effects of aphid infestation treatment (atm) with Brachycaudus lychnidis and Silene latifolia plant sex on plant stress indicators, behaviour of the pollinator Hadena bicruris and plant traits involved in communication and resource exchange with pollinators.*

	(1) Sex	(2) Atm	(3) Cofactor†	(4) Sex × Atm	(5) Sex × Cofactor†	(6) Atm × Cofactor†	(7) Sex × Amt. × Cofactor†	(8) Sex moth	Random effects
Aphid number None, GLMM, NB, none	0.328	-	**0.000**	-	-	0.207	-	-	Plant individual
Aphid weight None, LM, G, none	0.535	-	-	-	-	-	-	-	-
Leaf biomass None, LM, G, none	**0.035**	0.516	-	0.397	-	-	-	-	-
Chlorophyll Log, LMM, G, D~(1-7)	**0.000**	0.774	**0.000**	0.206	0.530	0.495	0.938	-	Plant individual
Absorption by Flavonoids None, LMM, G, none	**0.028**	0.637	**0.000**	0.167	0.421-	0.539	0.448	-	Plant individual
Pollinator choice C-A None, GLMM, B, none	**0.000**	††	**0.000**	††	0.266	††	††	0.052	Moth individual, trial
Pollinator choice F-M None, GLMM, BB, none	††	0.909	0.954	††	††	0.962	††	0.862	Moth individual, trial
Flower number None, GLMM, P, Z~ I D~(1)	**0.000**[ii]	0.082[ii]	**0.000**[ii]	0.238[ii]	0.757[ii]	**0.000**[ii]	**0.012**	-	Plant individual
Flower biomass None, LMM, G, D~(3)	**0.000**[ii]	**0.001**[ii]	**0.000**[ii]	0.123[ii]	**0.000**[ii]	**0.000**[ii]	**0.000**	-	Plant individual
Flower colour None, LMM, G, D~(1,3,5)	0.394[ii]	**0.000**[ii]	**0.000**[ii]	**0.027**	0.977	**0.016**	0.6908	-	Plant individual
Flower area None, GLMM, NB, D~(1-7)	0.348	0.092	**0.000**	0.645	0.124	0.942	0.53	-	Plant individual
Scent metabolome PERMANOVA (Bray-Curtis)	0.550	0.081	-	0.477	-	-	-	-	-
Nectar volume None, LMM, G, none	**0.000**	0.554	-	0.225	-	-	-	-	Plant individual
Nectar metabolome (LC) PERMANOVA (Bray-Curtis)	**0.000**	0.204	-	0.308	-	-	-	-	-

The table provides an overview of the model structures with regard to the included responses (column 1), predictors and random effects (row 1); as well as further model characteristics such as response transformations (log = natural logarithm, none = none), model types (GLMM, LMM, LM, PERMANOVA), the assumed error families (B = binomial, BB = betabinomial, G = Gaussian, NB = overdispersed Poisson with quadratic parametrization, P = Poisson), and additional zero-inflation / dispersion formulas (Z / D~ = account for access of zeroes in Poisson data / account for predictor-specific variance inhomogeneity, none = none). Moreover, the table provides an overview of p-values from type III ANOVAs (based on Wald χ²-test for (G)LMMs,

Tables

F-tests for LMs, and permutation statistics with pseudo-F estimation for PERMANOVA). Significant predictor terms are highlighted with bold lettering. Main effects and lower order interaction terms involved in significant (higher order) interactions are marked with "ii". Effects that were not tested are marked with "-". Please note that (†) the identity of cofactors for each response is described in detail in chapter 3.3.6; and (††) that in pollinator behaviour models, the actual test whether pollinators prefer control (C) over aphid infested (A) plants or female (F) over male (M) plants is a post-hoc on the intercept of the binomial model, see chapter 3.2 for details.

Tables

Table 2: Overview of metabolites highlighted in Figure 5B.

ID Figure 5	Effect(s)	Effect direction(s)	Molecular formula	Compund class	Compound identity	ID S1
1	None	-	C7H8O2	Benzenoids	2-Methoxy phenol	C15
2	None	-	C8H10O	Benzenoids	2-Phenylethanol	C19
3	Atm	↓	C7H6O	Benzenoids	Benzaldehyde	C07
4	None	-	C8H8O2	Benzenoids	Methyl benzoate	C16
5	None	-	C8H8O3	Benzenoids	Methyl salicylate	C24
6	None	-	C8H8O	Benzenoids	Phenylacetaldehyde	C11
7	None	-	C8H14O2	Fatty acid derivatives	cis-3-Hexenyl acetate	C09
8	None	-	C9H16O2	Mono-terpenoids	Lilac alcohol A	C25
9	None	-	C9H16O2	Mono-terpenoids	Lilac alcohol B + C	C26
10	Atm	↓	C9H16O2	Mono-terpenoids	Lilac alcohol D	C27
11	Atm	↓	C10H16O3	Mono-terpenoids	Lilac alcohol formate A	C34_35
12	Atm	↓	C10H16O3	Mono-terpenoids	Lilac alcohol formate B	C36
13	Atm	↓	C9H14O2	Mono-terpenoids	Lilac aldehyde A	C21
14	Atm	↓	C9H14O2	Mono-terpenoids	Lilac aldehyde B+C	C22
15	None	-	C9H14O2	Mono-terpenoids	Lilac aldehyde D	C23
16	Atm	↓	C10H16	Mono-terpenoids	trans-ß-Ocimene	C12
17	None	-	C5H11NO	N-bearing compounds	anti-3-Methyl-butyl-aldoxime	C06
18	None	-	C5H11NO	N-bearing compounds	syn-3-Methyl-butyl aldoxime	C04
19	Sex*Atm	F↓M-	C15H14O2	Phenyl-propanoids	Benzoic acid, 2-phenylethyl ester	C40
20	Atm	↓	-	-	-	C20

Metabolites highlighted with IDs in Figure 5 were significantly (uncorrected p-value < 0.05) and remarkably (log$_2$ fold change <-1 or >1) modulated by aphid infestation treatment (atm; ↓ = reduced, ↑ = increased, - = unaffected by aphid infestation) and sex (F=female, M=male) and/or have been shown to elicit behavioural or antennal responses in the pollinator Hadena bicruris according to (Dötterl *et al.*, 2006) *(all except Figure ID 19 and 20). Metabolites are listed with their molecular formula, compound class, putative identity, and ID in the full dataset Supplementary Table S1A.*

Tables

 Table 3: Overview of metabolites highlighted in *Figure 6C-F*.

ID Fig 6	Effect direction sex	Effect direction atm	Compund class	Compound identity	ID S1
1	-	-	Carbohydrates & conjugates	D-Glucose	ID0000
2	-	-	Carbohydrates & conjugates	D-Fructose	ID110_125_129
3	-	-	Carbohydrates & conjugates	Sucrose	ID60_138
4	-	-	Carbohydrates & conjugates	Nystose	ID0080
5	-	-	Carbohydrates & conjugates	Melezitose	ID71_142
6	-	-	Carbohydrates & conjugates	Palatinose	ID0117
7	-	↑	Carbohydrates & conjugates	Stachyose	ID0140
8	-	-	Carbohydrates & conjugates	2α-Mannobiose	ID0136
9	F>M	-	Carbohydrates & conjugates	N-Acetylglucosamine	ID0162
10	-	↓	Carbohydrates & conjugates	†Disaccharide glycosylglycerol ester	ID0115
11	**C:F<M, A:F=M**	**F↔, M↓**	**Carbohydrates & conjugates**	**† Nucleoside analogue**	**ID0054**
12	**C:F>M, A:F=M**	**F↔, M↔**	**Carbohydrates & conjugates**	**†Nucleoside analogue**	**ID0111**
13	-	↓	Carbohydrates & conjugates	†Phenolic acid glycoside	ID0033
14	-	↓	Carbohydrates & conjugates	Geinistin	ID0455
15	-	↓	Carbohydrates & conjugates	-	ID0050
16	-	↓	Amino acids, peptides, analogues	DL-Phenylalanine	ID0194
17	**F<M**	**-**	**Amino acids, peptides, analogues**	**L-Tyrosine**	**ID0179**
18	**F<M**	**-**	**Amino acids, peptides, analogues**	**N-(3-Methylbutanoyl)valine**	**ID0164**
19	**F>M**	**-**	**Amino acids, peptides, analogues**	**H-Ser-Leu$_5$-OH**	**ID1690**
20	**F<M**	**-**	**Amino acids, peptides, analogues**	**L-PheNH$_2$,Lys-Leu-Val-Phe**	**ID2889**
21	**F>M**	**-**	**Amino acids, peptides, analogues**	**Unknown peptide**	**ID1655**
22	**F>M**	**-**	**Amino acids, peptides, analogues**	**Unknown peptide**	**ID1700**
23	**F<M**	**-**	**Amino acids, peptides, analogues**	**Unknown peptide**	**ID0047**
24	**F<M**	**-**	**Amino acids, peptides, analogues**	**†Macrocyclic despipeptide**	**ID1489**
25	**F>M**	**-**	**Amino acids, peptides, analogues**	**† Amino alcohol diester**	**ID1870**
26	*C:F>M, A:F=M*	*F↓, M↔*	*Amines*	*Pyrrolidine derivate*	*ID0040*
27	**F<M**	**-**	**Purines and purine derivatives**	**† Guanosine derivate**	**ID0165**
28	*C:F=M, A:F=M*	*F↓, M↔*	*Purines and purine derivatives*	*Xanthine derivate*	*ID0103*
29	**F>M**	**-**	**Glycerophosphates**	**† PG(O-20:1/17:1)**	**ID2141**
30	**C:F>>M, A:F>M**	**F↔, M↔**	**Glycerophosphoethanolamines**	**† Alkyl-PE(C15:0)**	**ID1683**
31	**F<M**	**-**	**Glycosphingolipids**	**† β-GalCer (C20mu:0)**	**ID2724**
32	**C:F=M, A:F<M**	**F↔, M↔**	**Phosphatidylethanolamine**	**† PE(16:0/18:sie2)**	**ID3072**
33	F<M	-	Ceramides	† Bis(bishexyloxybiphenyl)amine	ID4068
34	**C:F<M, A:F=M**	**F↔, M↔**	**Ceramides**	**Palmitoleoyl ethanolamide**	**ID0990**
35	*C:F>M, A:F=M*	*F↓, M↔*	*Ceramides*	*Hydroxylated ceramide*	*ID1697*
36	***C:F=M, A:F=M***	***F↓, M↔***	***Ceramides***	***-***	***ID1771***
37	**F<M**	**-**	**Fatty amides**	**† GABA derivate**	**ID2723**
38	**F>M**	**-**	**Fatty amides**	**† Aminoglycerol-Derivate**	**ID1842**
39	**F>M**	**-**	**Fatty amides**	Fumaric acid diester	**ID1952**
40	**F>M**	**-**	**Fatty amides**	†Glyco-hydroxy fatty amide	**ID1787**
41	**F>M**	**-**	**Fatty amides**	**-**	**ID1633**
42	**F>M**	**-**	**Polyol**	**† Trimethyltrihydroxyundecanyl**	**ID1664**
43	F<M	-	Flavonoid glycosides	† Spiroflavonoid glycoside	ID0293
44	**F<M**	**-**	**Flavonoid glycosides**	**Vitexin-7-Rutinoside**	**ID0294**
45	**F>M**	**-**	**Flavonoid glycosides**	**-**	**ID1611**
46	*C:F>M, A:F=M*	*F↓, M↔*	*Flavonoids*	*-*	*ID0131*
47	**F>M**	**-**	**Terpene glycosides**	**-**	**ID2302**
48	**F>M**	**-**	**Triterpenoids**	**† Triterpenoid steroidal glycosides**	**ID2436**
49	**F>M**	**-**	**Triterpenoids**	**† Triterpenoid steroidal glycosides**	**ID2372**
50	*C:F=M, A:F=M*	*F↓, M↔*	*Benzenesulfonyl compounds*	*-*	*ID0091*

Tables

Metabolites highlighted with IDs in Figure 6C-F *were saccharides (primary nutritional compounds) or putatively identified compounds that were significantly (FDR corrected [bold lettering] / uncorrected p-value <0.05) and remarkably (log$_2$ fold change <-1 or >1) modulated by plant sex (F=female, M=male, entries only in column 2), aphid infestation treatment (atm; ↓ = reduced, ↑ = increased under aphid infestation, entries only in column 3) or the interaction plant sex × aphid infestation treatment (entries in column 2 and 3). The abundance of features printed in italic correlates with moth behaviour (*Figure 3*). Metabolites are listed with their compound class, putative identity, and ID in the full dataset* Supplementary Table S1B*. †: IUPAC names for the regarding metabolites are replaced with simplified names for compact presentation and fully listed* S1B*.*

Figures

Figure 1: *Plant-pollinator-herbivore study system. (A) Silene latifolia (female), (B) Hadena bicruris (male), (C) adult Brachycaudus lychnidis individual and colony feeding on flowering stems of S. latifolia.*

Figures

Figure 2: *Overview of (A) the experimental setup (abbreviations: FC = control female, FA = aphid infested female, MC = control male, MA = aphid infested male); the data acquisition for (B) the plant stress indicators (i) absorbance by chlorophyll and epidermally located flavonoids and (ii) aphid load in terms of weight and number, (C) pollinator behaviour, (D) visual flower traits related to (i) color, (ii) quantity and (iii) display area, (E) floral scent, and (F) flower nectar; as well as (G) the time line for data acquisition after inoculation of plants with aphids.*

Figures

Figure 3: *Effects of Silene latifolia plant sex and infestation by Brachycaudus lychnidis on the behaviour of the moth pollinator Hadena bicruris in terms of flower visits, feeding attempts and the total time spent on a flower. (A) Illustrates choice tests for control (C, deep coloured) versus aphid infested (A, light coloured) female (F, red) and male (M, turquoise) plants (FC/FA: n=11, MC/MA: n=17). (B) Illustrates choice tests for female (red) versus male (turquoise) control (C, dark coloured) and aphid infested (A, light coloured) plants (FC/MC: n=19, FA/MA: n=9). Graphs combine Box-Whisker plots of raw data with the marginal means +/- standard error (black squares) estimated for all groups by the (G)LMMs. Asterisks indicate significant deviations of the binary outcome from 0.5 (i.e., significant preference > 0.5 > aversion; significance levels: ***p<0.001, **p<0.01, *p<0.05).*

Figures

Figure 4: *Combined effects of Silene latifolia plant sex and Brachycaudus lychnidis aphid infestation treatment on (A) flower numbers, (B) reproductive biomasses, and (C) flower colours. The graphs combine Box-Whisker plots of raw data for female control (FC, dark red, n = $5_{(A,B)}/14_{(C)}$), female infested (FA, light red, n = $5_{(A,B)}/7_{(C)}$), male control (MC, dark turquoise, n = $5_{(A,B)}/13_{(C)}$), and male infested (MA, light turquoise, n = $5_{(A,B)}/13_{(C)}$) plants with the marginal means +/- standard error (black squares) for these groups estimated by the (G)LMMs. Significant effects detected via type III ANOVAs based on Wald χ^2 tests are denoted at the top of the plot, while the results of relevant post-hoc comparisons are denoted within the plots (significance levels: ***p<0.001, **p<0.01, *p<0.05).*

Figure 5: *Combined effects of Brachycaudus lychnidis aphid infestation treatment and plant sex on floral headspace VOC composition in Silene latifolia. (A) The NMDS ordination for female control (FC, dark red, n = 4), female infested (FA, light red, n = 4), male control (MC, dark turquoise, n = 5), and male infested (MA, light turquoise, n = 5) plants shows plant individual scores (points), distances to group centroids (spiders), and their standard error (ellipses). (B) Volcano plot summarizing the results of model loops, i.e., the main and interactive effects of plant sex and aphid infestation treatment on the abundance of individual metabolites. Metabolic features modulated by aphid infestation are highlighted (log$_2$ fold change <-1 or >1 and uncorrected [after FDR correction all effects non-significant] p-value <0.05 = coloured) with their effect direction (increased | decreased under aphid infestation (↑ | ↓) = brown | beige). A single metabolic feature (ID 18) was shaped by the interaction aphid infestation treatment × plant sex and is illustrated with one point for female (F) and male (M) plants. Numbered metabolic features correspond to Table 2. For further details see Supplementary Table S1A.*

Figures

Figure 6: *Combined effects of Silene latifolia plant sex and Brachycaudus lychnidis aphid infestation treatment on* (A) *nectar volume, and* (B-F) *the metabolic composition of a standardized nectar volume.* (A) *Box-Whisker plots of raw data for female control (FC, dark red, n = 23), female infested (FA, light red, n = 13), male control (MC, dark turquoise, n = 25), and male infested (MA, light turquoise, n = 25) plants with the marginal means +/- standard error (black squares) for these groups estimated by the (G)LMMs. Significant effects detected via type III ANOVAs based on Wald χ^2 tests are denoted at the top of the plot, while the results of relevant post-hoc comparisons are denoted within the plots (significance levels: ***p<0.001, **p<0.01, *p<0.05).* (B) *The NMDS ordination for FC (n = 5), FA (n = 3), MC (n = 5), and MA (n = 5) plants shows plant individual scores (points), distances to group centroids (spiders), and their standard error (ellipses).* (C-F) *Volcano plots illustrate the results of model loops, i.e., the main and interactive effects of plant sex and aphid infestation treatment on the abundance of individual metabolic features. Modulated features are highlighted (FDR corrected [uncorrected] p-value <0.05 and $\log_2$ fold change <-1 or >1 = coloured with black margin [coloured without margin]) with their effect direction (plant sex: F>M = red, M>F = turquoise, aphid infestation treatment: increased under aphid infestation (↑) = brown, decreased (↓) = beige). Interaction effect plots adopt the same modulation colour coding for* (E) *the sex effect within control and aphid infested plants and* (F) *the aphid infestation effect within females and males. Identical features within these group comparisons are connected by lines. Numbered metabolic features correspond to* Table 3*. A maximum of 15 identified features was highlighted for each modulation direction. For further details see* Supplementary Table S1B*.*