Title Pages

Article Type:

Research Paper

Title:

Aphid infestation induces plant-sex-specific changes in floral chemistry and pollinator

behaviour in *Silene latifolia*

Runing Title:

Sex-specific floral responses to aphid infestation

Authors:

Kaya B. Zill[1], Thomas Stegemann[2], Elisabeth Kaltenegger[3], Wolfgang Bilger[5], Tobias J.

Demetrowitsch[4], Henry Berndt[6], Alexandra Erfmeier[1], Sybille B. Unsicker[5], Karin Schrieber[1]

Affiliations:

[1] Kiel University, Faculty of Mathematics and Natural Sciences, Institute for Ecosystem

Research, Geobotany

[2] Kiel University, Faculty of Mathematics and Natural Sciences, Pharmaceutical Institute,

Pharmaceutical Biology

[3] Kiel University, Faculty of Mathematics and Natural Sciences, Botanical Institute and

Botanical Garden, Biochemical Ecology and Molecular Evolution

[4] Kiel University, Faculty of Agricultural and Nutritional Sciences, Institute for Human Nutrition

and Food Science, Division of Food Technology

[5] Kiel University, Faculty of Mathematics and Natural Sciences, Botanical Institute and

Botanical Garden, Plant-Environment-Interactions

[6] Kiel University, Mathematics and Natural Sciences, Zoological Institute, Marine Symbiosis

E-Mail addresses and Orchid IDs:

Karin Schrieber: kschrieber@ecology.uni-kiel.de, 0000-0001-7181-2741

Thomas Stegemann: tstegemann@pharmazie.uni-kiel.de, 0000-0002-0391-3337

Sybille Unsicker: sunsicker@bot.uni-kiel.de, 0000-0002-9738-0075

Tobias Demetrowitsch: tdemetrowitsch@foodtech.uni-kiel.de, 0000-0002-3528-0255

Elisabeth Kaltenegger: ekaltenegger@bot.uni-kiel.de, 0000-0001-6542-1509

Wolfgang Bilger: wbilger@bot.uni-kiel.de, 0000-0001-7800-4210

Alexandra Erfmeier: aerfmeier@ecology.uni-kiel.de, 0000-0002-1002-9216Henry Berndt:

hberndt@zoologie.uni-kiel.de, 0000-0002-0735-5539

Kaya Zill: NA, stu218014@mail.uni-kiel.de

Corresponding Author:

Dr. Karin Schrieber

Kiel University

Institute for Ecosystem Research

Olshausenstr. 75

24118 Kiel

Phone: +49 (0) 431 880-4082

E-mail: kschrieber@ecology.uni-kiel.de

Counts:

Words main text excl. material and methods: 5457

Figures: 6

Tables: 3

Supporting   information: 3

Highlight:

Aphid infestation alters multiple visual and chemical floral traits in a plant sex-specific

manner, leading to reduced attractiveness to moth pollinators in female plants, but not in

males.

Abstract:

Pollinators share the complex information and resource landscape of their host plants with

herbivores. Yet, how sap feeders affect floral attractiveness to pollinators remains poorly

understood, despite the critical role of this tripartite interaction in natural and agricultural

ecosystems. In dioecious plant species, which display pronounced sexual dimorphism, these

intricate interactions may vary in magnitude and direction between females and males, with

significant implications for plant population dynamics and species co-evolution. In this study,

we examined how infestation by the oligophagous aphid *Brachycaudus lychnidis* affects sex-

specific interactions among the dioecious plant *Silene latifolia* and its specialist moth

pollinator *Hadena bicruris*. We exposed male and female plants to aphid herbivory and

evaluated its effects on floral traits (visual cues, floral scent, and nectar chemistry) and

pollinator behaviour. While aphid infestation affected some floral traits equally in both sexes

and others more strongly in males or in females, we observed stronger declines in female

attractiveness to pollinators, which were mainly linked to nectar compounds potentially acting

as feeding cues or behavioural modulators. We discuss our results in the light of sexual

selection and plant defence theory while emphasizing the complementarity of female and

male traits in stabilizing this specialized plant-pollinator-herbivore system.

Keywords: *Brachycaudus*, dioecy, *Hadena bicruris*, nectar chemistry, nocturnal pollination,

metabolomics, *Silene latifolia*, sexual dimorphism, plant-pollinator-herbivore interaction

Highlight, Abstract, Keywords

Abbreviations used in the main text:

- ANOVA: analysis of variance
- FC, FA, MC, MA: female control, female aphid infested, male control, male aphid infested
- GC-MS: gas chromatography-mass spectrometry
- (G)L(M)M: (generalized) linear (mixed effects) model
- IQR: Interquartile range
- NMDS: nonmetric multi-dimensional scaling
- UHPLC-ESI-qTof-MS/MS: quadrupole time of flight - electro spray ionization - tandem mass spectrometry
- PERMANOVA: permutational analysis of variance
- Q1/3: Quartile 1/3
- VOCs: Volatile organic compounds

1. Introduction

Understanding plant × pollinator × herbivore interactions and their component species' co-evolutionary dynamics has recently become a central goal in chemical ecology (Moreira *et al.*, 2019; Kessler and Chautá, 2020; De-la-Cruz *et al.*, 2022; Muola *et al.*, 2022). The complex plant traits mediating interactions with pollinators and herbivores show considerable overlap, and we are beginning to understand the ecology of this shared trait space in systems involving chewing-biting herbivores. However, the impact of sap feeders on communication and resource exchange with pollinators remains largely unexplored, despite their critical role in the function of both natural and agricultural ecosystems (Morales-Hojas, 2017; Noriega *et al.*, 2018).

Plants evolve diverse traits to attract pollinators and modulate their behaviour thereby maximizing their own fitness with minimal resource investment (Santamaría and Rodríguez-Gironés, 2015). Next to flowering phenology and visual signals (flower number, spatial arrangement, size, shape, hue, and colour luminance), floral chemistry plays a central role in ensuring these complex functions (Borghi *et al.*, 2017; Trunschke *et al.*, 2021; Wong *et al.*, 2023). The emission of floral volatile organic compounds (VOCs) can trigger oriented movement of the receiver towards the emitter, and elicit landing, feeding as well as reproductive behaviour (Muhlemann *et al.*, 2014). Such scent signals can be honest and guide visitors to a high-quality reward, or deceive them about the presence of food or mates, thereby exploiting their pollination service (Ito *et al.*, 2021; Peakall, 2023). Nectar - the primary nutritional reward provided to pollinators - is a complex secretion composed of sugars, amino acids, lipids, proteins and diverse secondary metabolites (Nicolson, 2022), and thus not only a ready-to-use energy source for feeding animals. Plants may use nectar as a toolkit to manipulate the behaviour of their floral visitors. Secondary nectar metabolites can cause phagostimulation and satiety; improve learning capacity and memory; promote muscle performance, motivation and arousal; but also initiate sluggish and imprecise movements (Nepi *et al.*, 2018; Mustard, 2020; Barberis *et al.*, 2023). These various functions facilitate the exploitation of highly efficient pollination services while reducing energy investment in nectar sugars and amino acids. Although we are becoming increasingly aware of this beautiful complexity, there are still few studies that take a holistic view of the pollination syndrome, considering the combined roles of visual traits, floral VOCs, and nectar metabolites.

Pollinators universally share the complex information and resource landscape of their host plants with herbivores. There is now evidence for interactions among traits and genes controlling plant reproduction and defence strategies, leading to complex reciprocal co-

evolutionary feedbacks (Kessler and Chautá, 2020; De-la-Cruz *et al.*, 2022). Herbivores may
directly reduce plant attractiveness to pollinators through aggression or the consumption of
floral tissues and vegetative structures necessary for flower production (McCall and Irwin,
2006). Multiple floral traits have thus evolved to protect floral tissues from herbivory (Paul *et*
al., 2022). Constitutive and induced plant defences may not only alter a plant's suitability to
herbivores, but also deter pollinators (Adler *et al.*, 2012) or reduce resource allocation to
pollinator attraction (Stamp, 2003). Specifically chemical plant traits are considered to
mediate multiple other antagonistic and synergistic pleiotropic effects on pollinator attraction
and herbivore defence (Kessler and Chautá, 2020). Defence compounds induced by
herbivores can for example be allocated to nectar or pollen and deter or harm pollinators
(Rivest and Forrest, 2020), while specific floral VOC attracting pollinators can be used as a
feeding cue for florivores (Sasidharan *et al.*, 2023). So far, insight into such complex tripartite
interactions comes almost exclusively from chewing-biting herbivores. Aphids, which are
among the most damaging plant pests (Dedryver *et al.*, 2010), however, have been largely
overlooked (but see Pareja et al., 2012). These exceptionally fast reproducing insects cause
considerable stress in plants. They passively feed on phloem sap, thereby rapidly depleting
their hosts of essential nutrients, altering their source-sink relationships, and excreting
phytotoxins as well as phytopathogenic viruses with their saliva (Nalam *et al.*, 2019;
Twayana *et al.*, 2022). Plants in turn, actively defend against aphids by changing the
composition of sugars and amino acids in the phloem sap or loading it with deterrents, toxins
or anti-digestive proteins (Nalam *et al.*, 2019; Twayana *et al.*, 2022). Floral traits attracting
pollinators are specifically prone to the direct and indirect effects of aphid infestation, as
aphids often feed close to reproductive structures on flowering stems. This proximity
suggests a high potential for interference with pollinator-mediated reproductive processes,
warranting closer empirical attention.

The eco-evolutionary link between pollinator attraction and herbivore defence becomes
particularly apparent in dioecious species (Johnson *et al.*, 2015). Plants that allocate male
and female reproductive function to different individuals often exhibit complementary sexual
divergence in both traits based on dramatic differences in resource demands for successful
reproduction (Barrett and Hough, 2013; Johnson *et al.*, 2015). Male success is predicted to
largely depend on efficient pollen transfer and thus pollinator attraction, while female success
relies on pollination, as well as on the production and defence of seeds, and their dispersal
(Moore and Pannell, 2011). Consequently, females are predicted to allocate fewer resources
to floral attractiveness but exhibit more efficient herbivory defence and tolerance as
compared to males (Avila-Sakar and Romanow, 2012; Barrett and Hough, 2013). As recent
research has produced contradictory evidence for these theories (Cornelissen and Stiling,

2005; Tang-Martínez, 2012; Sargent and McKeough, 2022), a central question remains:
What are the sex-specific ecological consequences of parallel interactions with pollinators
and herbivores in dioecious plant species? Addressing this question is key to understanding
how co-selection by pollinators and herbivores contributes to the evolution of divergent
reproductive strategies in dioecious plants.

A system that is ideal to study these complex interplays consists of the dioecious plant *Silene*
latifolia POIR. (Caryophyllaceae), its nocturnal pollinator *Hadena bicruris* HUFN. (Noctuidae)*,*
and the aphid *Brachycaudus lychnidis* L. (Aphididae). The species involved in this system
engage in highly specialized relationships. Researchers have only begun to uncover the
sophisticated functional traits that have co-evolved to stabilize the cost-benefit ratio of their
individual bipartite interactions (Blair and Wolfe, 2004; Dötterl *et al.*, 2006; Labouche and
Bernasconi, 2010; Schrieber *et al.*, 2021). Insight into the ecology of the tripartite interaction,
however, is lacking. Here, we exposed female and male *S. latifolia* plants to control and *B.*
lychnidis infestation treatments to assess flowering phenology, spatial flower arrangement,
flower colour, the chemical composition of floral scent and nectar as well as nectar quantity,
and attractiveness to *H. bicruris*. We specifically aimed at characterizing i) sexual
dimorphisms in floral attractiveness; ii) the effect of aphid infestation on floral attractiveness
to pollinators; and iii) sex-specific differences in the susceptibility to aphid infestation and the
quality and magnitude of responses to aphid infestation.

2. Methods

2.1 Study organisms

Silene latifolia (white campion, *Figure 1A*) is a short-lived dioecious plant species within the Caryophyllaceae family, which is native to ruderal habitats across Eurasia and North Africa. The plant exhibits a distinct moth pollination syndrome. Its white, large, and funnel-shaped flowers open at dusk to release a scent bouquet composed of more than 60 VOCs (Dötterl and Jürgens, 2005; Dötterl *et al.*, 2005) to advertise large amounts of nectar rewards (Tew *et al.*, 2021) with a mostly unknown chemical composition. Male and female *S. latifolia* are known to exhibit pronounced sexual dimorphisms in a broad set of floral traits. Observed differences in flower number, size, colour, total floral scent emission, and nectar volume as well as sugar content (Gehring *et al.*, 2004; Waelti *et al.*, 2009*a*; Delph *et al.*, 2010; Schrieber *et al.*, 2021)are largely in line with theory of sexual selection, i.e., higher trait values in male plants (Bateman, 1948).

Silene latifolia is readily visited by diurnal generalist insects and nocturnal moths, with *Hadena bicruris* (campion owl, *Figure 1B*) - a crepuscular moth within the Noctuidae family - being the most frequent and relevant pollinator species in Eurasia (Jürgens *et al.*, 1996; van Putten *et al.*, 2007; Schrieber *et al.*, 2021). Together both species form one of only ~16 worldwide known nursery pollination systems (Hossaert-McKey *et al.*, 2010). Female moths pollinate female plants and oviposit on flower ovaries to provide their hatching larvae with nutritious seeds and shelter by developing fruits. The pollination services provided by male moths, together with the high egg and larval mortality caused by parasitoids and predators of *H. bicruris*, can compensate for the costs of seed consumption (Labouche and Bernasconi, 2010; Villacañas de Castro and Hoffmeister, 2020). Together these factors shift the outcome of the *Silene latifolia* × *Hadena bicruris* interaction for the plant towards the latter end of a seed predation-pollination continuum. A substantial fraction of VOCs in the floral scent of *S. latifolia* triggers antennal and behavioural responses in *H. bicruris* (Dötterl *et al.*, 2006). While the function of floral scents in this intricate co-evolutionary relationship is well studied (Dötterl *et al.*, 2006), the functional chemistry of nectar - the only reward and main food source provided to adult moths - remains unexplored beyond its basic sugar composition (Shykoff, 1997; Gehring *et al.*, 2004; Witt *et al.*, 2013).

A third species participating in this highly specialized system within Eurasia is the holocyclic aphid *Brachycaudus lychnidis (Figure 1C)*, which spends both the parthenogenetic and sexual part of its life cycle on *S. latifolia*. The aphid feeds mainly on the phloem sap of flowering stems.

2.2 Plant and insect cultivation

Seeds from five *S. latifolia* mother plants were collected in a population in Wiesengrund,
Germany (51°41'17.1"N, 14°35'45.8"E) and experimentally outcrossed for two generations
under greenhouse conditions (see Schrieber *et al*., 2019 for details). We cultivated 30 plants
per maternal family individually in 360 mL pots filled with organic potting soil (Floragard
TKS® 2 Instant Plus, DE) in a greenhouse at 25 °C / 15 °C and 60 % RH in a 16 h day / 8 h
night cycle. When plants started flowering after 6 weeks, we selected two males and two
females from each of the five maternal families (N = 20). The selected plants were
transferred into 8 L pots filled with the above-mentioned substrate and pruned to a height of
15 cm to induce over-compensatory growth. Please note that *S. latifolia* is adapted to
grazing/mowing and thus cutting is near-natural (Jessen *et al*., 2022). Moreover, cutting was
necessary to induce the emergence of multiple shoots and sufficient flower production for the
experiment. Plants were fertilized with 0.3 g Universol® Yellow (Everris-Headquarters, NL)
dissolved in 200 mL of water every other week and prophylactically treated with sulphur
vapor to prevent fungal infections as well as with biological pest control agents to prevent
thrips and uncontrolled aphid infestation (agents: *Amblyseius cucumeris against thrips;*
Aphidius ervi, Aphidius colemani and Lysiphlebus testaceipes against aphids Katz Biotech
GmbH, DE). The preventive treatment with parasitoid wasps was stopped three weeks
before controlled aphid infestation.

Adult *H. bicruris* moths were collected in different wild *S. latifolia* populations around Kiel,
Germany (54°19′ N, 10°8′ O) in 2022 and kept together in ventilated, transparent 20 x 30 cm
boxes equipped with egg cartons and tissue papers within a climate chamber (MLR-352H,
PHC Corporation, Ōizumi-machi, JP) at 25 °C / 15 °C and 60 % RH in a 16 h day / 8 h night
cycle. They were fed a honey-water solution (6 g organic flower honey, 14 mL distilled water)
from plastic *S. latifolia* flowers. In addition, moths received one fresh male and female *S.*
latifolia flower per day, as they did not reproduce without them. Individual flowers were
harvested from a pool of food plants and presented in Eppendorf tubes filled with water.
Hadena bicruris eggs, which were deposited all across the breeding boxes, were collected
and placed on moist filter paper in individual ventilated plastic containers (50 mL). Upon
hatching, larvae were provided with an artificial diet according to (Elzinga *et al*., 2002), which
was additionally supplemented with 20 mL of *S. latifolia* seeds per 500 mL to increase
acceptance. Pupated *H. bicruris* were moved to ventilated 50 mL containers filled with sterile
sand and newly hatched moths were transferred into the above-mentioned breeding boxes
and food supply for adults.

Brachycaudus lychnidis aphids were collected and identified according to
(influentialpoints.com, 2024) in spring 2024 in different wild *S. latifolia* populations around

Kiel, Germany (54°19′ N, 10°8′ O). They were cultivated in net cages on a balanced mixture

of male and female *S. latifolia* plants from our study population in a climate chamber (HGC

1514 Module 7, Weiss Umwelttechnik GmbH, DE) at 23 °C and 55 % RH, in a 16 h day / 8 h

night cycle. Food plants were watered and exchanged on demand.

2.3 Experimental setup and data acquisition

In mid-July 2024 at the age of 13 weeks, all selected plants were enclosed in 150 × 50 cm

organza mesh bags and one individual per sex (male, female) × maternal family (I-V)

combination was inoculated with 50 adult aphids *(Figure 2A, Supplementary Figure S1A-C)*.

All aphid colonies showed an initial decline in population size of max. 10 % and switched to

positive growth rates max. three days after the inoculation. Thus, our approach does not

account for potential plant sex-specific host choice, early phase plant defences preventing

colonization success, and indirect defence; but focuses on the effects of established aphid

colonies having optimal growth conditions. Data on plant stress indicators, pollinator choice,

visual flower traits, floral scent, and nectar *(Figure 2B-F)* were collected over a 40-day period

following aphid inoculation *(Figure 2G)*.

2.3.1   Plant stress indicators

To quantify the disruption of physiological homeostasis under aphid infestation, we optically

assessed the chlorophyll concentration, and the absorbance by epidermally located UV-

absorbing compounds, mainly flavonoids using the LEAF-STATE-ANALYZER LSA-2050

(Heinz Walz GmbH, DE) *(Figure 2B&G)* for all plants. As reference, data provided with the

instrument for the lower leaf side of *Sedum telephium* was used. Data were acquired directly

before and 40 days after aphid inoculation between 11:00 am and 12:00 noon in five leaves

per plant individual. The selected leaves were evenly distributed along the vertical axis of the

plants and well exposed to light (i.e., not significantly shaded by other leaves) and the

measured area, located in the upper middle part of the lamina, was free of honeydew and

aphids, which typically feed on stems. Immediately after these measurements, we counted

the number of aphid individuals per size category (small: < 1.5 mm, medium: < 2 mm, large:

> 2 mm) directly on the plant. Subsequently, the aphids were carefully washed off the plants

with warm tap water, collected on filter paper and dried for 24 h at 25 °C, then for 24 h at 65

°C, and finally for 24 h at 105 °C to determine total aphid dry biomass per plant.

2.3.2 Pollinator behaviour

Behavioural assays with *H. bicruris* were conducted under laboratory conditions 30-39 days

after aphid inoculation *(Figure 2C&G)* between 8:00 pm and 0:30 am - the peak activity time

of our captive moths. For the choice assays, freshly hatched adult moths were kept

individually in ventilated, transparent boxes (15 × 12 × 11 cm) equipped with shelters and a

feeding station featuring a honey-water solution as described in *chapter 2.2*. For each assay,

the artificial diet was removed in the morning and replaced in the evening with two fully

developed (male: fully dehisced anthers, female: fully expanded and papillous stigma; flower

age 2-3 days), freshly picked flowers from the experimental plants, which were carefully fixed

in lidless Eppendorf tubes filled with tap water. The two flowers were positioned on opposite

sides of the boxes for maximum spatial separation *(Supplementary Figure S1D)* to provide

the moths with the binary choice among the following four treatment groups: control females

versus aphid infested females, control males *versus* aphid infested males, control females

versus control males, aphid infested females *versus* aphid infested males. The moth

individuals used in these behavioural tests (N = 11) differed in age and sex (5 females, 6

males; moth sex was accounted for in the statistical analysis) and were tested one to two

times in each of the four treatment groups (depending on flower availability) using a fully

randomized design that resulted in a total of 61 trials. Each night, up to eight moths were

used, with two observers monitoring four moths each. The boxes were evenly lit with dim red

LED lamps to allow observation without confounding the trials (the visual system of moths

cannot capture red light). Three behavioural parameters were quantified for each of the two

flowers: (i) the number of visits, with a visit starting and ending with physical contact to the

flower; (ii) the number of feeding attempts per flower, with an attempt starting/ending by

unrolling/rolling up and inserting/removing the proboscis into/from the corolla tube; and (iii)

the total time spent with any physical contact on the flower. These three parameters were

assessed for the first four visits within each trial. We deliberately decided against first-visit

choice tests, as these give chance undue weight and moth individuals frequently revisit

individual flowers or probe several flowers before feeding extensively on one under natural

conditions (personal observation made in Schrieber *et al*., 2021). Female moths did not

oviposit on the flowers offered in the choice test because they did not mate in isolation.

2.3.3 Visual flower traits

We measured a broad set of visual flower traits at the plant individual and single flower level

throughout the experiment *(Figure 2D&G)*.

Flower colour was quantified using a digital image transformation method that accounts for
natural light conditions and the visual system of the pollinator based on (Troscianko and
Stevens, 2015). Images were acquired 21-25 days after aphid inoculation for three to six
flowers (depending on availability) from all plant individuals. They were taken outdoors in an
open area, free from shadows cast by trees or buildings, during the last hour of daylight on
rain-free days to match the natural light conditions perceived by *Hadena* (Bopp and
Gottsberger, 2004). A fully developed but fresh flower (age 2-3 days) was plugged into a
custom-made black ethylene vinyl acetate platform, which featured two reflectance standards
(PTFB 10% and Spectralon 99%, Labsphere, UK) along with a size reference
(Supplementary Figure 1E). Images were captured using a modified Samsung NX1000
camera (Suwon, KOR) with its sensor filter removed, to achieve full-spectrum sensitivity (300
– 1000 nm). The camera was equipped with a UV-sensitive lens (Nikon EL 80 mm, JP). To
capture both visible and ultraviolet (UV) light, we took one photo with a UV/IR blocking filter
(UV/IR Cut, transmitting 400-700 nm, Baader Planetarium, DE) and another one with a UV
pass plus IR blocking filter (U-filter, transmitting 300-400 nm, Baader Planetarium, DE) using
a custom-made filter slide mount. All photos were taken with an aperture of 5.6, and ISO of
800, whereas the shutter speed was adjusted to the available light conditions to receive well-
exposed photos. Post-processing was carried out using the Multispectral Image Calibration
and Analysis (MICA) Toolbox plugin (Troscianko and Stevens, 2015) within ImageJ 1.47 t
(Rueden *et al.*, 2017). We selected the flower silhouette for colour analysis. The images were
linearized to correct the camera's non-linear light response and standardized based on the
two reflectance standards to compensate for variations in natural lighting. We fitted the
multispectral images to a mathematical model incorporating (i) the spectral sensitivity of our
Samsung NX1000-Nikkor EL 80 mm 300-700 nm camera (data derived from Troscianko and
Stevens, 2015); (ii) the spectral sensitivities of the three photoreceptors in the moth
compound eye (data derived from Johnsen *et al.*, 2006); and (iii) the spectral composition of
sun light during dusk (data derived from Johnsen *et al.*, 2006). For further details see
Schrieber *et al.* (2021). The outcomes of these analyses are the percentage of green, blue
and UV-A dusk sun radiation reflected by the flower, which are perceivable by a moth's visual
system.

Flower area was measured from the same images for three to six flowers per plant individual.
We assessed the exact area covered by a single, well developed petal limb, the area
covered by the exact flower silhouette and the area covered by the circular perimeter *(Figure*
2D, 1 = flower perimeter, 2 = flower silhouette, 3 = petal silhouette). The circular perimeter
reflects the flower's maximum extension, but not its visual impact on pollinators due to
natural variation in the shape of single petals, while the area covered by the exact flower

silhouette may represent what pollinators see but not the plant's investment into flowers
(petal overlap), which can be captured by individual petal area.

The number of opened flowers and flower buds per plant individual were recorded at three
time points during the experiment (10, 20 and 40 days after aphid inoculation, *Figure 2D&G*)
and corrected for the flowers removed for other analyses. Directly after the removal of aphids
(40 days after aphid inoculation), plant aboveground parts were cut off directly on top of the
soil and separated into shoots (without leaves), and flowers. Plant parts were dried
individually in paper bags for 24 h at 65 °C followed by an additional 24 h at 105 °C to
determine dry biomass. During data acquisition for pollinator behaviour, flower colour, and
nectar rewards, we collected the harvested flowers in plant individual-specific bags to assess
their dry biomass as described above and added it to the total flower dry biomass.

2.3.4 Flower scent

At day 27 after the aphid inoculation, floral VOC emission was collected for all experimental
plants during the peak emission in *S. latifolia* between 10:00 pm (sunset) and 1:00 am
(Dötterl *et al.*, 2005) using a mobile push-pull system as described in Berkum *et al.*, (2024) .
Up to three fresh and fully developed flowers per plant (age 2-3 days) from different shoots
were enclosed in 35 cm × 31 cm PET bags (RUBIN® Bratschlauch, DE), which were closed
at both ends with cable ties *(Figure 2E)*. Compressed air entered the system after passing
through an activated charcoal filter on the lower side of the bag at a flow rate of 1 L min^{-1} and
was pulled out at the top through an adsorbent Poropak filter (Alltech, USA), using a vacuum
at the rate of 0.6 L/min. After VOC collection, the filters were eluted with 200 μL
dichloromethane (Sigma Aldrich, DE) containing 10 ng/μL of nonyl acetate (Sigma Aldrich,
DE) as an internal standard. As the headspace of treated plants also included aphids, we
additionally determined which compounds were exclusively released by *Brachycaudus*. We
enclosed 60 aphids (20 small: < 1.5 mm, 20 medium: > 1.5 mm, 20 large: > 2 mm) from our
laboratory population in the above-described PET bags to collect and analyse their volatile
emissions as described for the plants with three replicates.

Samples were analysed on a gas chromatograph (TRACE GC1310, ThermoFisher Scientific,
DE) equipped with a TraceGOLD TG-5MS column (30 m × 0.25 mm × 0.25 μm + 5 m;
ThermoFisher Scientific, DE) and coupled with a mass spectrometer (TSQ Duo MS,
ThermoFisher Scientific, DE). A sample volume of 1 μL was injected with a split ratio of 1:10
at 250 °C and migrated with a helium-flow of 1 mL min^{-1}. The transfer line was set to 270 °C
and the ion source of the mass spectrometer was operated at 230 °C and 70 eV for
ionization. The temperature program started at 40 °C for 2 min, ramped to 180 °C at a rate of

6 °C min^{-1}, ramped to 300 °C at a rate of 60 °C min^{-1}, and finished with a 5 min hold time at

300 °C. Mass spectra of separated compounds were acquired in full scan mode between 33

and 350 m/z. An alkane standard mix (C$_8$-C$_{20}$, Sigma Aldrich, DE) was analysed under the

same conditions to calculate linear retention indices (RI) for the targeted compounds. The

data processing was performed with Chromeleon^{TM} 7.3 (ThermoFisher Scientific, DE). The

putative identity of compounds was determined via comparisons of mass spectra and RI's

with databases (NIST Mass Spectral Search Program Version 3.0, Smith *et al.*, 2004) and

(Dötterl *et al.*, 2005). The identities were assigned according to the following criteria: match

factor to the NIST databank >750 and RI difference < 10. We selected all VOCs for the

statistical analyses that could not be clearly identified as contaminations. The base peak in

the mass spectrum of these VOC was used as quantitation ion. Additionally, two secondary

ions (confirmation ions) were selected per compound to ensure correct peak assignment

(Supplementary Table S1A). The relative compound abundance was quantified by dividing

the peak area (based on individual quantitation ion) of each volatile by the peak area of the

internal standard nonyl acetate (quantitation ion 43, confirmation ions 56, 70). To account for

variation in flower number in the headspace, relative scent abundance was normalized per

flower. In addition to this semi-quantitative approach, we created another dataset to focus on

qualitative differences only by dividing the peak area of each VOC by the total peak area of

all compounds.

2.3.5 Nectar quantity and chemistry

We measured the nectar volume for the flowers that have been harvested for the colour and

size analyses, directly after they have been photographed. All nectar was extracted from the

base of the ovary / the base of the stamens and anthopophor into 1 µL microcapillary tubes

(Minicaps NA-HEP, Hirschmann Laborgeräte, DE). The length of the nectar column was

measured with a calliper to determine the exact volume.

Nectar chemistry was addressed based on a standardized nectar volume via Ultra High

Performance Liquid Chromatography - Quadrupole Time of Flight - Electro Spray Ionization -

Tandem Mass Spectrometry (UHPLC-ESI-qTof-MS/MS). For these analyses, we additionally

collected all nectar from three fresh, fully developed flowers per plant individual using 1 µL

microcapillary tubes on day 28 after aphid infestation. After determining the exact nectar

volume as described above, each microcapillary was rinsed with 30 µL of HPLC grade water

(Agilent, USA) into a 0.5 mL Eppendorf tube, in which all nectar from one plant individual was

pooled. The samples were then vortexed, snap-frozen in liquid N$_2$ and stored at -80 °C.

Samples underwent a modified simultaneous metabolite, protein, and lipid extraction

(SIMPLEX) as described in (Schrieber *et al.*, 2023) to obtain both, hydrophilic and lipophilic

nectar metabolites. After defrosting and vortexing, 0.1 mL of nectar solution was mixed with

0.5 mL of ultra-pure methanol (Carl Roth, GE) and 4 mL of methyl-tert-butylether (LC-grade,

Carl Roth, GE). Samples were incubated for 30 min at 25 rpm in an end-over-end shaker

before phase separation was induced by adding 0.5 mL of ultra-pure water (Carl Roth, GE).

After 10 min of incubation, samples were centrifuged at 4000 *g* and 4 °C for 10 min and the

upper lipophilic phase was collected. The lower hydrophilic phase was used as a template to

repeat the above-described protocol for the extraction of hydrophilic metabolites. Extracts

were re-suspended in 0.5 mL of either ultra-pure water and ultra-pure methanol (hydrophilic

extracts: 50 / 50, v / v, Carl Roth, GE) or ultra-pure isopropanol and LC-grade chloroform

(lipophilic extracts: 3:1, v / v, Carl Roth, GE). The dilution of the resulting nectar stock

solutions for analyses via UPLC-qTof-MS/MS was then adjusted to differences in the volume

of extracted nectar among plant individuals. Hence, the volume of analysed nectar was

standardized for these analyses.

For UHPLC-ESI-qTof-MS/MS analyses, 2 µL of extract were injected at 30 °C into a UHPLC

(Nexara, Shimadzu, DE) containing a C_{18} Column (100 × 2.1 mm, 1.7 µm, Luna Omega

Polar, Phenomenx, DE). The solvent system was composed of (A) water with 0.1% formic

acid, and (B) acetonitrile with 0,1% formic acid (all LC-grade, Carl Roth, DE). The gradient

program started at a flow rate of 0.3 mL min^{-1} with 5 % B, was then raised over 20 min to 95

% B, and finally kept at 95 % B for an additional 5 min. Afterwards, the column was re-

equilibrated at 5% B for 6 min. Chromatographically separated analytes entered a qTOF-

MS/MS (x500R, Sciex, DE) with the following parameters: ion source gas 45 psi (nitrogen),

curtain gas 35 psi (nitrogen), CAD gas 7 psi (nitrogen), electrospray ion source temperature

500 °C, ion spray voltage 5500 V. The MS acquisitions parameters were: MS1 scans at 100-

1000 m/z with a data-dependent MS2 trigger having a collision energy (CID) of 40 eV and a

scan range of 50-1000 m/z. The acquired data were then converted to .mzML format and

further processed in mzMine (Schmid *et al.*, 2023). Metabolic features with a maximum m/z

difference of 0.001 amu and maximum retention time difference of 0.1 min were bucketed

and treated as identical features in the following analysis. For feature annotation, the

acquired MS/MS spectra were compared against public mass spectral libraries from Global

Natural Products Social Network (GNPS) (Wang *et al.*, 2016) and the MassBank (MassBank

consortium and its contributors, 2024) using the NIST search algorithm (cosine similarity of

0.6) and the MassBank formula (cosine similarity of 0.5). Features that remained unidentified

were further annotated using COSMIC and CSI:FingerID according to the SIRIUS software

package (Hoffmann *et al.*, 2022).

2.3.6   Statistical analyses

All statistical analyses were implemented in R 4.4.1 (R Development Core Team, 2024).
Plant stress indicators, spatial flower traits, flower colour traits, and floral rewards were
analysed as responses in (generalized) linear mixed effects models - (G)LMMs, which
included the aphid infestation treatment, plant sex and the interaction among both variables
as predictors. Highly correlated traits were combined into a single response and analysed
within one model. These models included the different types of summarized response traits
as a factorial co-predictor, for which all possible interactions with sex and aphid infestation
treatment were tested according to the scheme *response at plant individual level (level of*
summary) ~ cofactor: leaf chlorophyll content (on day 0 and 40 of the experiment) ~ time
point; leaf epidermal light absorption by flavonoids (on day 0 and 40 of the experiment) ~
time point; number aphids (belonging to the size category small, medium and large) ~ aphid
size category; flower number (on day 10, 20 and 40 of the experiment) ~ time point; dry
biomass (of a single flower, all flowers, and flowering stems) ~ biomass component; display
area (of a single petal limb, all petal limbs, and the full corolla expansion) ~ flower
component; flower light reflection (in the green, blue and UV-A range) ~ colour quality. The
remaining traits (nectar volume) were acquired across several flowers per plant individual
and analysed in this high resolution (i.e., not based on means). Consequently, all models had
repeated measurements on the plant individual level and thus included a random effect for
the plant individual. All models were checked for violations of their underlying assumptions
using tests and plots from the R-package "DHARMa" (Hartig, 2022) and were adjusted with
regard to error family, dispersion or zero-inflation, if necessary. An overview of all model
parameters is provided in *Table 1*. Based on these models, we calculated type III ANOVA
tables (based on Wald- χ^2 tests) while setting all contrasts to sum-zero coding using the R-
package "car" (Fox and Weisberg, 2019). If factorial predictors were involved in significant
interactions, we additionally calculated post-hoc contrasts among the estimated means of
their levels within levels of other factors involved in the respective interaction (R-package
"emmeans"; Lenth, 2024)).

The choice tests were analysed with two binomial models testing (i) preferences for a
specific aphid infestation treatment within sexes (assays: control female *versus* aphid-
infested female, control male *versus* aphid-infested male); and (ii) preferences for a specific
plant sex within aphid infestation treatments (assays: control female *versus* control male,
aphid infested female *versus* aphid-infested male). Model i / ii included the behavioural
parameter in control relative to aphid infested flowers, i.e., the proportion of visits, feeding
attempts or time spent in control relative to infested flowers (i) or female relative to male
flowers, i.e., the proportion of visits, feeding attempts or time spent in control female to male

flowers (ii) as binomial response and the flowers sexes (i) or the flowers aphid infestation

treatments (ii) as predictors. Further predictors included in both models were the behavioural

parameter type (visits, feeding attempts, time; all acquired in the same moth and trial), and

its interaction with the previously mentioned model-specific predictor. In addition, both

models included the sex of the moth as a cofactor for which only additive effects were tested,

as sample sizes were not large enough to reliably test for all possible interactions. Moth

individual and trial were set as random effects *(Table 1)*. Model checking, adjustment and

downstream analyses with type III ANOVA were implemented as described above. Post-hoc

contrast (R-package "emmeans"; Lenth, 2024) were used to test for a significant deviation of

the binary outcome from the intercept of 0.5 (i.e., preference or avoidance) within the

different plant sex × behavioural type combinations (i) or aphid infestation treatment ×

behavioural type combinations (ii).

The statistical approaches used for the analyses of the metabolome are identical for floral

nectar and scent. In a first step, we performed a permutational analyses of variance

(PERMANOVA) using the Bray-Curtis dissimilarities of the full metabolome datasets (n = 33

features in scent [for both, semi-quantitative and qualitative only data sets], n = 4,385

features in nectar assessed via UPLC-qTof-MS/MS) as the response distance matrix. We

included aphid infestation treatment, sex, and their interactions as predictors and ran the

models with 100,000 permutations (R-package "vegan" (Oksanen *et al.*, 2024)). As

previously implemented tests for multivariate homogeneity of group dispersions in the

distance matrices among all treatment levels yielded no deviation for all datasets, the

predictor effects assessed via PERMANOVA are based on differences in group centroids

rather than differences in group dispersions (Anderson and Walsh, 2013). The multivariate

group separations among plant infestation treatments and sexes were illustrated with Non-

metric Multidimensional Scaling (NMDS) plots (R-package "vegan" (Oksanen *et al.*, 2024)).

To identify the compounds responsible for this separation, we used an approach that is

comparable to a volcano plot (Cui and Churchill, 2003) and simultaneously allows the

statistical assessment of interaction effects among plant sex and aphid infestation treatment.

Instead of applying two-sided significance tests (t or u-tests), we entered all metabolites as

responses into a linear model loop to test for the main and interactive effects of plant sex and

aphid infestation treatment. Each metabolite ran with three alternative models as

untransformed, squareroot-transfromed and log-transformed response. To check for

violations of model assumptions and select the best of the three alternative models for each

metabolite accordingly, we extracted the results from the Kolmogorov-Smirnov test for

uniformity (normality of residuals) and Levene's test for homogeneity of variance, which are

provided for targeted model optimization in the R-package "DHARMa" (Hartig, 2022). If

model assumptions were violated in all three alternative models, the respective metabolite
was not further analysed. The best fit models for the remaining metabolites were further
analysed with type III ANOVAs based on F-tests (R-package "car" (Fox and Weisberg, 2019))
and the calculation of post-hoc contrasts (R-package "emmeans" (Lenth, 2024)) as described
in the previous section. *P*-values were corrected for inflation of alpha errors by multiple
testing (false discovery rate: FDR) according to (Benjamini and Hochberg, 1995). If a
metabolites p-value in a given group comparison ($F_{C\&A}/M_{C\&A}$, $T_{F\&M}/C_{F\&M}$, FC/FT, MC/MT,
FC/MC, FT/MT) was significant and the corresponding $\log_2$-fold change in the intensity of this
metabolite was either ≤ -1 (halving of intensity in the first relative to the second listed group)
or ≥ 1 (doubling of intensity in the first relative to the second listed group) it was considered
to be modulated.

3. Results

4.1 Plant stress indicators

After 40 days of inoculation with *B. lychnidis*, aphid load in terms of weight and number was

equally high in female and male *S. latifolia* plants *(Table 1)*. Females had significantly higher

trait values than males for leaf dry mass per plant individual (*females = 5.33 +/- 0.56, males*

= 3.50 +/- 0.56, p<0.05, F_{2DF}= 5.28), leaf chlorophyll content (*females = 27.5 +/- 1.37, males*

= 22.9 +/- 0.91, p<0.001, χ^2_{2DF}= 14.37) and absorption by epidermal flavonoids (*females =*

0.70 +/- 0.03, males = 0.62 +/- 0.03, p< 0.05, χ^2_{2DF}= 4.80), but aphid infestation treatment

had no effects on these traits *(Table 1)*. Moreover, leaf chlorophyll contents decreased

(p<0.001, χ^2_{2DF}= 60.34), whereas absorption by epidermal flavonoids increased *(p<0.001,*

χ^2_{2DF}= 49.47) from the beginning towards the end of the experiment, regardless of sex and

aphid infestation treatment *(Table 1*, column cofactor*)*.

3.1 Effects of plant sex and aphid infestation on pollinator behaviour

A subset of our choice assays tested behavioural moth responses to single flowers from

control versus aphid-infested plants, and whether these responses varied between male and

female plants. The effects of aphid infestation treatment and plant sex on choice behaviour

were marginally significant and highly significant, respectively *(Table 1; p= 0.072, χ^2_{2DF}= 3.23;*

p< 0.001, χ^2_{2DF}= 142.91). Post-hoc comparisons showed that only females, but not males,

exhibited a highly significant preference for flowers from control plants over aphid-infested

ones across all behavioural parameters *(Figure 3A; z-ratio female plants C-A: 3.36*** [visits],*

3.23*** [feeding attempts], 4.19*** [time spent]; z-ratio male plants C-A: -0.08^{ns} [visits], -

0.24^{ns} [feeding attempts], -0.56^{ns} [time spent]). In addition, we detected a marginally

significant main effect of pollinator sex *(Table 1, Supplementary Figure S2, p=0.097, χ^2_{2DF}=*

2.89), indicating that the overall choice behaviour differed slightly between females and

males. Post-hoc tests showed that only female but not male *H. bicruris* moths showed a

significant preference for control over aphid-infested *S. latifolia* plants *(z-ratio female moth C-*

A: 2.45*, z-ratio male moth C-A: 0.16^{ns}, Supplementary Figure S2).

Another subset of our choice assays tested whether moths prefer single flowers from one

plant sex over the other and whether this preference is modified by the aphid infestation

treatment. The statistical analysis yielded no significant preferences of the moths for either

plant sex and neither aphid infestation treatment nor their own sex shaped their choice

(Figure 3B, Table 1).

3.2 Effects of plant sex and aphid infestation on visual flower traits

Flower number was significantly shaped by the three-way interaction sex × infestation treatment × and infestation time *(Table 1, Figure 4A, p< 0.05, χ^2_{2DF}= 8.88)*. Post-hoc comparisons revealed that female plants suffered both earlier and stronger declines in flower number upon aphid infestation as compared to males *(female t-ratio C-A: 2.05^{ns} [10 days], 4.25*** [20 days], 9.26*** [40 days]; male t-ratio C-A: 0.40^{ns} [10 days], 1.36^{ns} [20 days], 4.74*** [40 days])*. Correspondingly, females produced significantly fewer flowers than males only in aphid infested plants and the magnitude of this effect increased over time *(control t-ratio F-M: -2.14^{ns} [10 days], -1.41^{ns} [20 days], -1.26^{ns} [40 days]; infestation t-ratio F-M: -3.84** [10 days], -4.31*** [20 days], -5.31*** [40 days])*. Reproductive biomass depended on the three-way interaction sex × infestation treatment × component *(Table 1, Figure 4B, p< 0.001, χ^2_{2DF}= 16.86)*. Post-hoc comparisons revealed that the component single flower mass was not affected by the infestation treatment *(t-ratio C-A: 1.17^{ns} [female], 1.14^{ns} [male])* but generally higher in females than males *(t-ratio F-M: 6.22*** [control], 5.76*** [infested])*. In contrast, the component total flower mass declined severely under aphid infestation, specifically in females *(t-ratio C-A: 12.89*** [female], 7.26*** [male])* and was accordingly higher in females only in controls but not under aphid infestation *(t-ratio F-M: 5.22*** [control], -0.41^{ns} [infested])*. The mass component of flowering stems was not affected by aphid infestation *(t-ratio C-A: 2.73$^{n.s}$ [female], 0.86$^{n.s}$ [male])*, and plant sex *(t-ratio F-M: 2.56^{ns} [control], 0.68^{ns} [infested])*. Flower light reflectance *(Table 1, Figure 4C)* was significantly shaped by the two two-way interactions sex × aphid infestation treatment *(p< 0.05, χ^2_{1DF}= 4.91)* and aphid infestation treatment × colour type *(p< 0.05, χ^2_{2DF}= 8.27)*. Post-hoc comparisons showed that the aphid-infestation treatment reduced floral light reflectance in the green and blue spectrum only in male, but not in female plants *(female t-ratio C-A: 1.39^{ns} [green], 1.69$^{ns.}$ [blue], 0.11^{ns} [UV]; male t-ratio C-A: 3.00* [green], 3.22* [blue], 2.09^{ns} [UV])*. Flower area was neither affected by plant sex, aphid infestation treatment or their interaction *(Table 1)*.

3.1 Effects of plant sex and aphid infestation on the composition of floral scent

We detected a total of 33 VOCs in the floral headspace of *S. latifolia*, with values normalized per flower. Of these 32 could be putatively identified via matches of RIs and mass spectra with databases *(Supplementary Table S1A)*. A large fraction of these VOCs have been previously described in *S. latifolia* (Dötterl *et al.*, 2005; Mamadalieva *et al.*, 2014) and of these, 16 have been shown to elicit antennal and/or behavioural responses in *H. bicruris* (Dötterl *et al.*, 2006)*.*

The PERMANOVA on the semi-quantitative VOC dataset detected a marginally significant
main effect of aphid infestation treatment on the composition of these floral VOCs *(Table 1, p*
< 0.087, F_{1DF} = 2.30, R^2 = 0.13) but no divergence among sexes or interactive effects. This is
in accordance with both, the group dispersion visualized in the NMDS ordination plot *(Figure*
5A) and the outcomes of the model loop *(Figure 5B, Supplementary Table S1A)*. Without
FDR correction, the loop identified eight compounds shaped by the main effect of treatment
with reduced abundance in aphid infested as compared to control plants. These included five
compounds known to elicit behavioural responses in *H. bicruris* according to Dötterl *et al.*
(2006) -benzaldehyde, trans-β-ocimene, lilac aldehyde A, lilac aldehyde B+C, and lilac
alcohol D- as well as two of their degradation products, lilac alcohol formate A and lilac
alcohol formate B. One compound which was not tested for behavioural effects on *H. bicruris*
so far - phenethyl benzoate - was significantly modulated by the interaction plant sex × aphid
infestation treatment and decreased under aphid infestation in female plants but not in
males. When accounting for inflation of alpha errors, all these effects were non-significant.
These effects were detected despite the limited sample size, and might become more
pronounced with additional replication. Multivariate analyses of qualitative differences only,
yielded no significant effects of aphid infestation treatment and plant sex *(Table 1)*.

3.2 Effects of plant sex and aphid infestation on nectar traits

Nectar volume was unaffected by the aphid infestation treatment but considerably shaped by
plant sex with female flowers featuring higher nectar amounts *(Table 1, Figure 6A, p< 0.001,*
χ^2_{1DF}= 46.10). The untargeted metabolome analyses of a standardized nectar volume per
plant via UPLC-qTof-MS/MS yielded abundance data for 4,385 features. Of these, 761
features were tentatively identified via matches of their mass spectra with databases or via
cheminformatics according (Hoffmann *et al.*, 2022), and an additional 71 could be assigned
to a compound class *(Supplementary Table 1B)*. This is the first detailed description of nectar
chemistry in the ecological model system *S. latifolia.* The species' nectar contained various
nutritious metabolites, including sugars (D-glucose, D-fructose, sucrose, melezitose,
stachyose, nystose, mannobiose, palatinose), proteinogenic and non-proteinogenic amino
acids (e.g., proline, glutamic acid, tyrosine ornithine) and numerous oligopeptides,
nucleosides (adenosine, adenosine monophosphate, a guanosine derivate), and multiple
lipids (e.g., fatty alkyls; fatty acids, amides and alcohols; ceramides;
glycerophosphoethanolamines, sphingolipids; prostaglandins). Its diverse secondary
metabolites comprised vitamins (e.g., calcitriol, riboflavin), flavonoids (e.g., saponarin,
vitexin-2''-O-rhamnoside), terpenoids (e.g., myrtenol, perillyl alcohol, raddeanin D,
pisumoside A), as well as fatty amides and amines (oleamide -a neuroactive compound

[Mendelson and Basile, 2001] which has so far not been described in nectar, tyramine,
homospermidine). Most of these secondary compounds are likely derived from plant
metabolism, but several have a known fungal or bacterial origin and thus likely derive from
the nectar microbiome (e.g., oleandomycin, enniatin B, asperiamide B, N-acetylglucosamine
[chitin monomer]).

The PERMANOVA detected a highly significant effect of plant sex *(Table 1, p < 0.001, F_{1DF} =*
10.65, R^2 = 0.39), but no effects of infestation treatment or the interaction sex × infestation
treatment on the composition of the metabolome in a standardized volume of nectar. The
group dispersions visualized in ordination plots via NMDS illustrate a clear differentiation of
the nectar metabolome among female and male plants and additionally indicate different
responses to the aphid infestation treatment in males and females in some features *(Figure*
6B). The results from the linear model loop further support this observation *(Figure 6C-F,*
Table 2, Supplementary Table S1B). When accounting (not accounting) for multiple testing, a
total of 219 (3) features had significantly and remarkably (log$_2$ fold change >1) higher
abundance in females than males, including six features that were absent in the latter. These
did not include primary nutritional compounds such as sugars or free amino acids, but seven
oligopeptides, nine fatty amides, six terpenoids (including three triterpenoid steroids), a
glycerophosphate, a flavonoid and the microbial compound N-acetylglucosamine [chitin
monomer]. The remaining discriminatory features, including those unique to female nectar,
could not be identified *(Figure 6C, Table 2, Supplementary Table 1B)*. On the other hand, 45
(15) features had significant and remarkably (log$_2$ fold change <-1) higher abundances in
males than females with 20 of them being absent in the latter. These included the free
proteinogenic amino acid tyrosine (unique), N-(3-Methylbutanoyl)valine (unique), two
oligopeptides (one unique), a macrocyclic despipeptide of probable microbial origin (unique),
two flavonoid glycosides (one unique), a glycerosphingolipid, and a guanosine derivate
(unique, highly significant). The aphid infestation treatment had clearly weaker effects on the
composition of plant nectar. When accounting (not accounting) for multiple testing, plants
showed increased abundances of 0 (10) features including the tetrasaccharide stachyose;
and decreased abundances of 0 (18) features including the amino acid phenylalanine, a
disaccharidester, and the isoflavonoid genistin *(Figure 6D, Table 2, Supplementary Table*
S1B). The interaction plant sex × aphid infestation treatment shaped the abundance of 21 (7)
features *(Figure 6E&F; please note that both plots display the same metabolic features with*
different group comparisons; Table 2; Supplementary Table S1B). These interactions had
seven major effect directions, which are listed as labels at the top of both plots. The most
prominent interaction pattern was significantly and markedly higher metabolite abundance in
females compared to males under control conditions, but not under aphid infestation in 16 (0)

features, which included a nucleoside analogue. The second most prominent interaction

pattern correlated with moth behaviour. In total 1(5) features had significantly and markedly

reduced abundances under aphid infestation in females, but not in males. We analysed the

MS² spectra for these six features in detail. Most of them did not yield high confidence

matches or similarity scores to known compounds, suggesting these metabolites are not

previously reported natural products. Our detailed analyses provided chemical class

annotations and further details on structural features allowing to discuss their putative

function in the context of plant-pollinator interactions (*Supplementary Table S1C*). These

compounds include a pyrrolidine derivative ($C_{17}H_{18}N$, m/z 237.021) with aromatic

substituents (m/z = 70.0648 → pyrrolidinegroup, m/z = 77.0382 → phenylmethylgroup, m/z =

91.0539 → phenylgroup), a xanthine derivative with additional substituents ($C_{14}H_{14}N_4O_5S$,

m/z 351.07), a benzenesulfonyl compound with a sugar-like substituent ($C_{14}H_{21}NO_7S$, m/z

351.07), two ceramides of which one bared multiple amino substituents ($C_{32}H_{65}NO_4$, m/z

528.3914; $C_{23}H_{51}N_5O_4$, m/z 484.3645) and a flavonoid ($C_{31}H_{24}O_9$, m/z 541.1492).

4. Discussion

Building on the pronounced sexual differentiation in floral traits of the dioecious plant *Silene*
latifolia, we examined how controlled infestation by the oligophagous aphid *Brachycaudus*
lychnidis affects interactions with the specialist moth pollinator *Hadena bicruris*. Contrary to
classical expectations from sexual selection theory, female plants were not outcompeted by
males in floral attractiveness under control conditions. Instead, they exhibited high plasticity
in floral trait expression and produced diverse, potentially bioactive nectar metabolites, some
of which may specifically target female pollinator behaviour. While aphid infestation affected
some floral traits equally in both sexes and others more strongly in males or in females, we
observed stronger declines in female attractiveness to *H. bicruris*. This points to sex-specific
plastic responses, such as increased attraction of alternative pollinators in females, as well
as sex-specific defence strategies, which we discuss here in detail.

4.1 Female *Silene latifolia* plants challenge expectations: no evidence of reduced floral
attractiveness compared to males

Bateman's theory of sexual selection predicts that, relative to males, female reproductive
success is more limited by available resources and therefore does not increase with the
number of mates, leading to less attractive flowers that serve as a resource exploited and
competed for by males (Bateman, 1948). In the present study, we found no evidence for
these predictions.

Hadena bicruris moths did not generally prefer single flowers of either sex (*Figure 4B*, *Table*
1 - pollinator choice F-M). Female and male plants exhibited no general differences in flower
numbers (*Figure 4A* - control conditions), flower sizes (*Table 1* - petal area), and flower
colours (*Figure 4C*). Female flowers even featured higher nectar volumes as compared to
males (*Figure 6A*) and major nectar sugar constituents showed no sex-specific differences
(*Figure 6C*, *Table 3*, *Supplementary Table S1B*). Instead, we observed higher abundances of
over 200 other primary and secondary metabolites in females than males, including six that
were unique to female nectar. Some of these compounds likely shape the nutritional and
pharmacological value of nectar for moth pollinators. These included seven oligopeptides,
partially enriched in branched-chain amino acids, which may serve as a rapid source of
NADH, $FADH_2$, and acetyl-CoA for ATP production, supporting energy demands during flight
or allocation to egg production in female moths (Levin *et al.*, 2017; Cavigliasso *et al.*, 2023);
nine fatty amides potentially serving as energy stores, metabolic fuel or behavioural
modulators (Ranger *et al.*, 2005; Yoshinaga, 2016; Levin *et al.*, 2017); six terpenoids (of
which two were steroidal glycosides) that may provide selective feeding or oviposition cues

(González-Coloma *et al.*, 2011; Boncan *et al.*, 2020); a glycerophospholipid that may support
the maintenance and generation of somatic tissue due to its central role in rapid membrane
synthesis (Toprak *et al.*, 2020); and a flavonoid which may act as health promotor in moth
pollinators (Palmer☐Young *et al.*, 2019). The function and further identification of the
numerous female biased metabolites, specifically the unidentified unique ones, will be the
subject of future studies. As only female plants bear the cost of the interaction with *H. bicruris*
(fruit predation), and only female moths impose this cost, there is great potential for the
discovery of nectar metabolites interfering with or stimulating reproduction in female moths.
Male *S. latifolia* plants, on the other hand, exhibited higher abundances of over 40
compounds, including several from the same chemical classes and with similar putative
functions as those dominant in female nectar (BCAA-rich oligopeptides, flavonoid glycosides,
membrane building lipids such as ceramides and glycosphingolipids). Most strikingly, tyrosine
and a derivate of guanosine were unique to male nectar. Both compounds can be considered
essential for the nutrition of moth pollinators that rely exclusively on nectar as a nitrogen
source and may enhance its taste and memorability accordingly (Nepi, 2014). In addition,
tyrosine is a precursor for the synthesis of the neurotransmitters tyramine and dopamine,
which shape numerous insect behavioural traits like e.g., motivation and learning (Barberis *et*
al., 2023). To the best of our knowledge, our study reports one of the most extensive cases of
sexual differentiation in the composition of nectar for a dioecious plant to date (see also Nepi
et al., 2001; Dötterl *et al.*, 2014). Since *H. bicruris* showed no preference for either plant sex
at the single-flower level (*Figure 4B*, *Table 1* - pollinator choice F-M), we conclude that these
striking differences complement each other functionally to stabilize a dioecious system
relying on frequent visits to both sexes, with plants promoting this by offering essential nectar
resources in only one sex.

Overall, our study offers a more nuanced view of female floral traits than predicted by
Bateman's principle. A potential criticism is that females in our greenhouse setting did not
exhibit lower floral attractiveness as they were not pollinated and hence did not invest in
seeds, which aligns with long-term personal observations of visual flower traits in *S. latifolia*.
However, this very aspect underscores the limitations of Bateman's theory as it supports that
females exhibit adaptive plasticity in floral traits, with attractiveness decreasing only as
pollination success increases. High attractiveness of unpollinated female plants may promote
female mate competition, post-pollination mate choice in females (Tonnabel *et al.*, 2021),
multiple paternity in fruits, and ultimately greater offspring diversity and fitness (Teixeira and
Bernasconi, 2007). This would be in line with the theory of flexible sex role behaviour
(Gowaty and Hubbell, 2005), which -similar to Bateman's principle- originates from zoology
and may be applicable to understanding reproductive strategies in dioecious plants. The

substantial among-population variation in sexual dimorphism observed in *S. latifolia* in

previous studies (Dötterl *et al.*, 2009; Schrieber *et al.*, 2017, 2019, 2021) further contradicts

the assumption of a fixed evolutionary trajectory, instead pointing toward locally driven and

context-dependent trait evolution.

4.2 Aphid infestation reduces female floral attractiveness via sex-specific trait

changes

In accordance with our hypothesis, infestation with *B. lychnidis* aphids affected a wide range

of traits in *S. latifolia* flowers, as well as the behavioural responses of *H. bicruris* moth

pollinators. Although most of the observed effects clearly depended on plant sex, we begin

with a general discussion of aphid-induced changes, as these have, to our knowledge, not

been previously reported.

Aphid infestation reduced flower numbers (more strongly in females than males, *Figure 4A*),

and reduced flower light reflection in the blue and green range (only in males, *Figure 4C*).

These adverse effects likely result from both resource depletion and the reallocation of

resources toward plant defence (Twayana *et al.*, 2022), and may substantially limit plant

visibility to pollinators (Barbot *et al.*, 2022). Aphid infestation also moderately reduced the

emission of five key volatile organic compounds (VOCs) known to elicit antennal and feeding

responses in *H. bicruris* (Dötterl *et al.*, 2006), at the level of individual flowers (*Figure 5B*,

Table 2). Nectar volume was not lower in aphid-infested plants (*Figure 6A*), likely because

they produced fewer flowers and could consequently invest more nectar in each one. While

the major sugar constituents remained largely unaffected (*Figure 6D*, *Table 3*), stachyose

levels increased in both sexes, suggesting that plants load their phloem with this poorly

digestible tetrasaccharide to impair aphid performance (Twayana *et al.*, 2022), while

simultaneously ensuring pollination by enzymatically breaking it down into simpler sugars

such as fructose, glucose, or sucrose in nectar. Conversely, levels of phenylalanine, a

nucleoside analogue, a phenolic acid glycoside, and an isoflavonoid were reduced. These

changes may result either directly from phloem-feeding-induced resource depletion or reflect

a passive carry-over of defensive phloem sap modifications targeting aphids (Twayana *et al.*,

2022). Although most of these alterations affected both sexes and would be expected to

generally reduce pollinator attraction, this was observed only in one sex.

At the single-flower level, *H. bicruris* preferred flowers from control plants over aphid-infested

ones only in *S. latifolia* females, but not males (*Figure 3A*). These preferences were also

dependent on moth sex, being primarily exhibited by female *H. bicruris* (*Table 1*). This

behaviour coincided with a reduction of the floral VOC phenylethyl benzoate - a compound

not previously described in *S. latifolia* but found in other moth-pollinated species (El-Sayed,
2011) - which was observed only in aphid-infested female plants, but not in males (*Figure*
5B, *Table 2*). Additionally, six metabolic features in the nectar of *S. latifolia* were reduced by
aphid infestation only in female plants (*Figure 6F*, *Table 3*). Four of them simultaneously had
higher abundances in female than male plants only under control but not under aphid
infestation conditions (*Supplementary table S1B*). Although these effects were only observed
before FDR correction, the corresponding effect direction in pollinator behaviour suggests
biological relevance. None of these metabolites belonged to the major nutritional nectar
constituents (sugars, amino acids) or could be clearly identified as a previously described
natural product. Our detailed chemoinformatic analysis of MS^2 spectra elucidated compound
classes and further structural components indicating a potential role of these metabolites as
selective feeding cues or behavioural modulators for female *H. bicruris*. One candidate
compound contains both pyrrolidine and phenyl substituents, a structural motif found in
several synthetic psychoactive compounds that inhibit monoamine reuptake transporters,
particularly dopamine, and are associated with stimulant activity (Meyer *et al.*, 2013). Another
notable candidate is a xanthine derivative, a class of compounds widely documented to
enhance learning, memory, and arousal in pollinators (Mustard, 2020). The benzenesulfonyl
compound containing amino and methoxy sugar-like moieties, the two ceramides and the
flavonoid may possibly contribute to specific recognition by pollinators via contact cues. A
detailed structural characterization of these novel natural products via NMR analysis of larger
nectar volumes as well as individual and combinatorial bioassays are necessary to validate
these putative ecological functions.

Even if more than just a small subset of the many measured floral traits had declined more
severely upon aphid infestation in females than in males (as was in fact not the case), it
would remain an overly simplistic conclusion that females are more vulnerable to stress. It
remains to be clarified which of these responses represent a direct negative consequence of
stress and which reflect a functional plastic adjustment under natural conditions. The finding
that the nursery pollinator *H. bicruris* avoided flowers of aphid-infested *S. latifolia* females
only *(Figure 3)*, combined with the greater choosiness of female moths *(Supplementary*
Figure S2), suggests that, under field conditions, infested female plants could experience a
shift in the outcome of the interaction towards the latter end of the seed predation-pollination
continuum. Additionally, reduced female flower attractiveness to *H. bicruris*, potentially
mediated by shifts in nectar metabolite composition, could be associated with increased
reliance on or even attraction of other pollinators that are less efficient but also less
damaging. Moreover, the role of sex-specific defence strategies against aphids deserves
particular attention in future studies. Our approach bypassed aphid host preferences; direct

plant defences preventing the early establishment of aphids, the attraction of aphid
parasitoids and predators that all occur in the field. Indeed, female plants in some dioecious
species have been shown to be less frequently attacked by herbivores (Sargent and
McKeough, 2022), to investigate more into direct defence (Sargent and McKeough, 2022)
and potentially also into indirect defences (Aranda-Rickert *et al.*, 2021). In fact it has been
shown that *S. latifolia* females in Europe are clearly less frequently attacked by aphids than
males (Wolfe *et al.*, 2004). Hence, they are unlikely to experience aphid infestation as strong
as in our experiment under more natural conditions. Yet, the cause for lower aphid
abundances on female plants remains to be investigated in the future. Sex-specific indirect
defences against aphids appear to be a particularly promising research route here. The
seed-predating larvae of *H. bicruris* are attacked by a broad range of parasitoids and
predators (Villacañas de Castro and Hoffmeister, 2020) that -if the plant is involved at all-
should be attracted by females only. There is ample evidence that the evolution of indirect
defences against different enemy species and guilds is tightly coupled via shared plant
signalling and VOC synthesis pathways or the attraction of generalist predators (Kessler and
Baldwin, 2001; Heil and McKey, 2003; Rasmann *et al.*, 2005).

5. Conclusion and outlook

Our study provides evidence for multidirectional sexual differentiation of visual and chemical
flower traits and their responses to aphid infestation in a dioecious plant species. It supports
that the expression of sexual dimorphisms in floral attractiveness can be highly plastic
depending on the presence of both, pollinators and herbivores, thereby opening exciting
avenues for future research. Controlled experiments manipulating the presence of both,
pollinators and aphids, ideally under field conditions reflecting natural conditions, while
integrating data collection on plant, pollinator and aphid reproductive success with
quantitative leaf and flower chemistry are a next step in disentangling the sex-specific co-
evolution in this tripartite system. A particular focus will be on the many newly discovered
sex-specific nectar metabolites that potentially serve as oviposition cues, physiologically
reduce fecundity or offspring survival, or force visits to both plant sexes. In such future
studies, it might be more purposeful to focus on the complementarity of female and male
defence and floral attractiveness traits in stabilizing a dioecious reproductive system
involving a nursery pollinator and several other parties (aphids, aphid and caterpillar
parasitoids and predators), instead of seeking for the better defended or more attractive sex.

6. Acknowledgements

Tobias Möckel from the FabLab in Kiel developed a customised camera attachment for light
filter switches that was used for the assessment of flower colour composition. Reza
Esmaeilnia Talemi supported the collection of plant nectar for metabolome analyses and
Meike Pfeiler prepared nectar extracts. Stefan Dötterl advised on the establishment of our
Hadena bicruris laboratory population, and Eric Folz was involved in the maintenance of the
aphid and pollinator laboratory populations. Roman Adler designed vector graphs for the
study design illustrations; and Verena Zajonc and Katharina Grosser provided valuable
technical support.

7. Supplementary data

Supplementary Table S1: Summary of floral metabolites in *Silene latifolia*: functional
annotation and responses to sex and aphid infestation
Supplementary Figure S1: Representative images from the experimental procedure
Supplementary Figure S2: Effects of moth sex on choice behaviour

8. Author contributions:

Conceptualization: K.S.; Methodology: K.S., T.S., E.K., S.U., W.B., and T.D.; Investigation:
K.Z., T.S., and E.K.; Validation: K.S., T.S., E.K., and W.B.; Formal Analysis: K.S.; Resources:
T.S., E.K., A.E., S.U., T.D., and W.B.; Data Curation: K.S.; Writing – Original Draft
Preparation: K.S.; Writing – Review & Editing: K.S., E.K., W.B., S.U., T.S, T.D., and H.B.;
Writing – Manuscript Revision: K.S., and E.K.; Visualization: K.S.; Supervision: K.S., S.U.;
Project Administration: K.S.; Funding Acquisition: K.S.

9. Conflict of interest:

No conflict of interest is declared

10. Funding

This study has been funded by the program for the promotion of young female scientists from

the Faculty of Mathematics and Natural Sciences of Kiel University

11. Data availability

All data supporting this article are be deposited in dryad under the following link:

https://doi.org/10.5061/dryad.83bk3jb6q. The code used for statistical analyses and

generation of graphs is deposited in zenedo under the following link:

https://doi.org/10.5281/zenodo.17700170.

12. References

Adler LS, Seifert MG, Wink M, Morse GE. 2012. Reliance on pollinators predicts defensive

chemistry across tobacco species. Ecology Letters **15**, 1140–1148.

Anderson MJ, Walsh DCI. 2013. PERMANOVA, ANOSIM, and the Mantel test in the face of
heterogeneous dispersions: What null hypothesis are you testing? Ecological Monographs
83, 557–574.

Aranda-Rickert A, Torréns J, Yela NI, Brizuela MM, Di Stilio VS. 2021. Distance
Dependent Contribution of Ants to Pollination but Not Defense in a Dioecious, Ambophilous
Gymnosperm. Frontiers in Plant Science **12**, 722405.

Avila-Sakar G, Romanow CA. 2012. Divergence in Defence against Herbivores between
Males and Females of Dioecious Plant Species. International Journal of Evolutionary Biology
2012, 897157.

Barberis M, Calabrese D, Galloni M, Nepi M. 2023. Secondary Metabolites in Nectar-
Mediated Plant-Pollinator Relationships. Plants **12**, 550.

Barbot E, Dufaÿ M, Tonnabel J, Godé C, De Cauwer I. 2022. On the function of flower
number: disentangling fertility from pollinator-mediated selection. Proceedings of the Royal
Society B: Biological Sciences **289**, 20221987.

Barrett SCH, Hough J. 2013. Sexual dimorphism in flowering plants. Journal of
Experimental Botany **64**, 67–82.

Bateman AJ. 1948. Intra-sexual selection in Drosophila. Heredity **2**, 349–368.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B
(Methodological) **57**, 289–300.

Berkum PM, Giorgi FD, Rothe B, Durka W, Gershenzon J, Roscher C, Unsicker SB.
2024. Selection strengthens the relationship between plant diversity and the metabolic profile
of Plantago lanceolata. bioRxiv.

Blair AC, Wolfe LM. 2004. The evolution of an invasive plant: an experimental study with
Silene latifolia. Ecology **85**, 3035–3042.

Boncan DAT, Tsang SSK, Li C, Lee IHT, Lam H-M, Chan T-F, Hui JHL. 2020. Terpenes
and Terpenoids in Plants: Interactions with Environment and Insects. International Journal of
Molecular Sciences **21**, 7382.

Bopp S, Gottsberger G. 2004. Importance of *Silene latifolia* ssp. *alba* and *S. dioica*
(Caryophyllaceae) as host plants of the parasitic pollinator *Hadena bicruris* (Lepidoptera,
Noctuidae). Oikos **105**, 221–228.

Borghi M, Fernie AR, Schiestl FP, Bouwmeester HJ. 2017. The Sexual Advantage of
Looking, Smelling, and Tasting Good: The Metabolic Network that Produces Signals for
Pollinators. Trends in Plant Science **22**, 338–350.

Cavigliasso F, Savary L, Spangenberg JE, Gallart-Ayala H, Ivanisevic J, Kawecki TJ.
2023. Experimental evolution of metabolism under nutrient restriction: enhanced amino acid
catabolism and a key role of branched-chain amino acids. Evolution Letters **7**, 273–284.

Cornelissen T, Stiling P. 2005. Sex-biased herbivory: a meta-analysis of the effects of
gender on plant-herbivore interactions. Oikos **111**, 488–500.

Cui X, Churchill GA. 2003. Statistical tests for differential expression in cDNA microarray
experiments. Genome Biology **4**, 210.

Dedryver C-A, Le Ralec A, Fabre F. 2010. The conflicting relationships between aphids and
men: A review of aphid damage and control strategies. Comptes Rendus Biologies **333**, 539–
553.

De-la-Cruz IM, Batsleer F, Bonte D, Diller C, Hytönen T, Muola A, Osorio S, Posé D,
Vandegehuchte ML, Stenberg JA. 2022. Evolutionary Ecology of Plant-Arthropod
Interactions in Light of the "Omics" Sciences: A Broad Guide. Frontiers in Plant Science **13**.

Delph LF, Arntz AM, Scotti-Saintagne C, Scotti I. 2010. The Genomic Architecture of
Sexual Dimorphism in the Dioecious Plant Silene Latifolia. Evolution **64**, 2873–2886.

Dötterl S, Glück U, Jürgens A, Woodring J, Aas G. 2014. Floral Reward, Advertisement
and Attractiveness to Honey Bees in Dioecious *Salix caprea*. PLOS ONE **9**, e93421.

Dötterl S, Jürgens A. 2005. Spatial fragrance patterns in flowers of *Silene latifolia*: Lilac
compounds as olfactory nectar guides? Plant Systematics and Evolution **255**, 99–109.

Dötterl S, Jurgens A, Seifert K, Laube T, Weissbecker B, Schutz S. 2006. Nursery
pollination by a moth in *Silene latifolia*: the role of odours in eliciting antennal and behavioural
responses. New Phytologist **169**, 707–718.

Dötterl S, Jürgens A, Wolfe L, Biere A. 2009. Disease Status and Population Origin Effects
on Floral Scent: Potential Consequences for Oviposition and Fruit Predation in A Complex
Interaction Between A Plant, Fungus, and Noctuid Moth. Journal of Chemical Ecology **35**,
307–319.

Dötterl S, Wolfe LM, Jürgens A. 2005. Qualitative and quantitative analyses of flower scent
in *Silene latifolia*. Phytochemistry **66**, 203–213.

El-Sayed AM. 2011. The pherobase: database of insect pheromones and semiochemicals.
HortResearch, Lincoln, New Zealand,(baza danych). *in press*.

Elzinga JA, Biere A, Harvey A. 2002. The rearing of the gregarious koinobiont
endoparasitoid *Microplitis tristis* (Hymenoptera: Braconidae) on its natural host *Hadena*
bicruris (Lepidoptera: Noctuidae). Proceedings of the Section Experimental and Applied
Entomology of the Netherlands Entomological Society (2002), 109–115.

Fox J, Weisberg S. 2019. *An R Companion to Applied Regression*. SAGE Publications.

Gehring JL, Scoby * Jennifer, Parsons M, Delph LF. 2004. Whole-plant investment in
nectar is greater for males than pollinated females in the dioecious plant *Silene latifolia*.
Evolutionary Ecology Research **6**, 1237–1252.

González-Coloma A, López-Balboa C, Santana O, Reina M, Fraga BM. 2011. Triterpene-
based plant defenses. Phytochemistry Reviews **10**, 245–260.

Gowaty PA, Hubbell SP. 2005. Chance, Time Allocation, and The Evolution of Adaptively
Flexible Sex Role Behavior1. Integrative and Comparative Biology **45**, 931–944.

Hartig F. 2022. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. R package version 0.4.6 **https://CRAN.R-
project.org/package=DHARMa**.

Heil M, McKey D. 2003. Protective Ant-Plant Interactions as Model Systems in Ecological
and Evolutionary Research. Annual Review of Ecology, Evolution, and Systematics **34**, 425–
553.

Hoffmann MA, Nothias L-F, Ludwig M, Fleischauer M, Gentry EC, Witting M, Dorrestein
PC, Dührkop K, Böcker S. 2022. High-confidence structural annotation of metabolites
absent from spectral libraries. Nature Biotechnology **40**, 411–421.

Hossaert-McKey M, Soler C, Schatz B, Proffit M. 2010. Floral scents: their roles in nursery
pollination mutualisms. Chemoecology **20**, 75–88.

https://influentialpoints.com/Gallery/Brachycaudus_lychnidis_Campion_aphids.htm.
2024. *in press*.

Ito K, Suzuki MF, Mochizuki K. 2021. Evolution of honest reward signal in flowers.
Proceedings of the Royal Society B: Biological Sciences **288**, 20202848.

Jessen M-T, Auge H, Harpole WS, Hautier Y, Eskelinen A. 2022. Grazing and light modify
Silene latifolia responses to nutrients and future climate. PLOS ONE **17**, e0276789.

Johnsen S, Kelber A, Warrant E, Sweeney AM, Widder EA, Lee RL, Hernández-Andrés
J. 2006. Crepuscular and nocturnal illumination and its effects on color perception by the
nocturnal hawkmoth *Deilephila elpenor*. Journal of Experimental Biology **209**, 789–800.

Johnson MTJ, Campbell SA, Barrett SCH. 2015. Evolutionary Interactions Between Plant
Reproduction and Defense Against Herbivores. Annual Review of Ecology, Evolution, and
Systematics **46**, 191–213.

Jürgens A, Witt T, Gottsberger G. 1996. Reproduction and Pollination in Central European
Populations of *Silene* and *Saponaria* Species. Botanica Acta **109**, 316–324.

Kessler A, Baldwin IT. 2001. Defensive Function of Herbivore-Induced Plant Volatile
Emissions in Nature. Science **291**, 2141–2144.

Kessler A, Chautá A. 2020. The ecological consequences of herbivore-induced plant
responses on plant–pollinator interactions. Emerging Topics in Life Sciences **4**, 33–43.

Labouche A-M, Bernasconi G. 2010. Male moths provide pollination benefits in the *Silene
latifolia–Hadena bicruris* nursery pollination system. Functional Ecology **24**, 534–544.

Lenth R. 2024. emmeans: Estimated Marginal Means, aka Least-Squares Means.
https://CRAN.R-project.org/package=emmeans **R package version 1.10.2.**

Levin E, McCue MD, Davidowitz G. 2017. More than just sugar: allocation of nectar amino
acids and fatty acids in a Lepidopteran. Proceedings of the Royal Society B: Biological
Sciences **284**, 20162126.

Mamadalieva NZ, Lafont R, Wink M. 2014. Diversity of Secondary Metabolites in the Genus
Silene L. (Caryophyllaceae)—Structures, Distribution, and Biological Properties. Diversity **6**,
415–499.

MassBank consortium and its contributors. 2024. MassBank/MassBank-data: Release
version 2024.11. Zenedo **https://doi.org/10.5281/zenodo.14221628**.

McCall AC, Irwin RE. 2006. Florivory: the intersection of pollination and herbivory. Ecology
Letters **9**, 1351–1365.

Mendelson WB, Basile AS. 2001. The hypnotic actions of the fatty acid amide, oleamide.
Neuropsychopharmacology: Official Publication of the American College of
Neuropsychopharmacology **25**, S36-39.

Meyer MR, Schmitt S, Maurer HH. 2013. Studies on the metabolism and detectability of the
emerging drug of abuse diphenyl-2-pyrrolidinemethanol (D2PM) in rat urine using GC-MS
and LC-HR-MS/MS. Journal of Mass Spectrometry **48**, 243–249.

Moore JC, Pannell JR. 2011. Sexual selection in plants. Current Biology **21**, R176–R182.

Morales-Hojas R. 2017. Molecular ecology of insect pests of agricultural importance: the
case of aphids. Ecological Entomology **42**, 18–27.

Moreira X, Castagneyrol B, Abdala-Roberts L, Traveset A. 2019. A meta-analysis of
herbivore effects on plant attractiveness to pollinators. Ecology **100**, e02707.

Muhlemann JK, Klempien A, Dudareva N. 2014. Floral volatiles: from biosynthesis to
function. Plant, Cell & Environment **37**, 1936–1949.

Muola A, Lucas-Barbosa D, Kessler A. 2022. Editorial: Mechanisms underlying plant-
pollinator-herbivore interactions. Frontiers in Plant Science **13**.

Mustard JA. 2020. Neuroactive nectar: compounds in nectar that interact with neurons.
Arthropod-Plant Interactions **14**, 151–159.

Nalam V, Louis J, Shah J. 2019. Plant defense against aphids, the pest extraordinaire.
Plant Science **279**, 96–107.

Nepi M. 2014. Beyond nectar sweetness: the hidden ecological role of non-protein amino
acids in nectar. Journal of Ecology **102**, 108–115.

Nepi M, Grasso DA, Mancuso S. 2018. Nectar in Plant–Insect Mutualistic Relationships:
From Food Reward to Partner Manipulation. Frontiers in Plant Science **9**, 1063.

Nepi M, Guarnieri M, Pacini E. 2001. Nectar Secretion, Reabsorption, and Sugar
Composition in Male and Female Flowers of *Cucurbita pepo*. International Journal of Plant
Sciences **162**, 353–358.

Nicolson SW. 2022. Sweet solutions: nectar chemistry and quality. Philosophical
Transactions of the Royal Society B: Biological Sciences **377**, 20210163.

Noriega JA, Hortal J, Azcárate FM, *et al.* 2018. Research trends in ecosystem services
provided by insects. Basic and Applied Ecology **26**, 8–23.

Oksanen J, Blanchet FG, Friendly M, *et al.* 2024. Vegan: Community Ecology Package.
https://CRAN.R-project.org/package=vegan **R package version 2.6-8**.

Palmer-**Young EC, Farrell IW, Adler LS, Milano NJ, Egan PA, Junker RR, Irwin RE,**
Stevenson PC. 2019. Chemistry of floral rewards: intra- and interspecific variability of nectar
and pollen secondary metabolites across taxa. Ecological Monographs **89**, e01335.

Pareja M, Qvarfordt E, Webster B, Mayon P, Pickett J, Birkett M, Glinwood R. 2012.
Herbivory by a Phloem-Feeding Insect Inhibits Floral Volatile Production. PLOS ONE **7**,
e31971.

Paul I, Poddar Sarkar M, Bhadoria PBS. 2022. Floral secondary metabolites in context of
biotic and abiotic stress factors. Chemoecology **32**, 49–68.

Peakall R. 2023. Pollination by sexual deception. Current Biology **33**, R489–R496.

R Development Core Team. 2024. R: A Language and Environment for Statistical
Computing. R Foundation for Statistical Computing.

Ranger CM, Winter REK, Rottinghaus GE, Backus EA, Johnson DW. 2005. Mass
spectral characterization of fatty acid amides from alfalfa trichomes and their deterrence
against the potato leafhopper. Phytochemistry **66**, 529–541.

Rasmann S, Köllner TG, Degenhardt J, Hiltpold I, Toepfer S, Kuhlmann U, Gershenzon
J, Turlings TCJ. 2005. Recruitment of entomopathogenic nematodes by insect-damaged
maize roots. Nature **434**, 732–737.

Rivest S, Forrest JRK. 2020. Defence compounds in pollen: why do they occur and how do
they affect the ecology and evolution of bees? New Phytologist **225**, 1053–1064.

Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, Eliceiri KW.
2017. ImageJ2: ImageJ for the next generation of scientific image data. BMC Bioinformatics
18, 529.

Rusman Q, Hooiveld-Knoppers S, Dijksterhuis M, Bloem J, Reichelt M, Dicke M,
Poelman EH. 2022. Flowers prepare thyselves: leaf and root herbivores induce specific
changes in floral phytochemistry with consequences for plant interactions with florivores.
New Phytologist **233**, 2548–2560.

Santamaría L, Rodríguez-Gironés MA. 2015. Are flowers red in teeth and claw?
Exploitation barriers and the antagonist nature of mutualisms. Evolutionary Ecology **29**, 311–
322.

Sargent RD, McKeough AD. 2022. New Evidence Suggests No Sex Bias in Herbivory or
Plant Defense. The American Naturalist **200**, 435–447.

Sasidharan R, Junker RR, Eilers EJ, Müller C. 2023. Floral volatiles evoke partially similar
responses in both florivores and pollinators and are correlated with non-volatile reward
chemicals. Annals of Botany **132**, 1–14.

Schmid R, Heuckeroth S, Korf A, *et al.* 2023. Integrative analysis of multimodal mass
spectrometry data in MZmine 3. Nature Biotechnology **41**, 447–449.

Schrieber K, Stegemann T, Kaltenegger E, Bilger W, Zill K. 2025. Dataset: Aphid
infestation induces plant-sex-specific changes in floral chemistry and pollinator behaviour in
Silene latifolia. Dryad, Dataset. https://doi.org/10.5061/dryad.83bk3jb6q

Schrieber K. 2025. R-script containing code for statistical analyses and graphs. Zenodo.

https://doi.org/10.5281/zenodo.17700170.

**Schrieber K, Glüsing S, Peters L, Eichert B, Althoff M, Schwarz K, Erfmeier A,
Demetrowitsch T**. 2023. Population divergence in heat and drought responses of a coastal
plant: from metabolic phenotypes to plant morphology and growth. (T Obata, Ed.). Journal of
Experimental Botany, erad147.

**Schrieber K, Paul SC, Höche LV, Salas AC, Didszun R, Mößnang J, Müller C, Erfmeier
A, Eilers EJ**. 2021. Inbreeding in a dioecious plant has sex- and population origin-specific
effects on its interactions with pollinators. (MC Aldrich and Y Joo, Eds). eLife **10**, e65610.

Schrieber K, Wolf S, Wypior C, Höhlig D, Hensen I, Lachmuth S. 2017. Adaptive and
non-adaptive evolution of trait means and genetic trait correlations for herbivory resistance
and performance in an invasive plant. Oikos **126**, 572–582.

Schrieber K, Wolf S, Wypior C, Höhlig D, Keller SR, Hensen I, Lachmuth S. 2019.
Release from natural enemies mitigates inbreeding depression in native and invasive *Silene
latifolia* populations. Ecology and Evolution **9**, 3564–3576.

Shykoff JA. 1997. Sex differences in floral nectar production by *Silene latifolia*
(Caryophyllaceae), with reference to susceptibility to a pollinator-borne fungal disease.
Canadian Journal of Botany **75**, 1407–1414.

Smith RN, Martell AE, Motekaitis RJ. 2004. NIST standard reference database. NIST
Critically Selected Stability Constants of Metal Complexes Database **Ver 2**.

Stamp N. 2003. Out Of The Quagmire Of Plant Defense Hypotheses. The Quarterly Review
of Biology **78**, 23–55.

Tang-Martínez Z. 2012. Repetition of Bateman challenges the paradigm. Proceedings of the
National Academy of Sciences of the United States of America **109**, 11476–11477.

Teixeira S, Bernasconi G. 2007. High prevalence of multiple paternity within fruits in natural
populations of *Silene latifolia*, as revealed by microsatellite DNA analysis. Molecular Ecology
16, 4370–4379.

Tew NE, Memmott J, Vaughan IP, Bird S, Stone GN, Potts SG, Baldock KCR. 2021.
Quantifying nectar production by flowering plants in urban and rural landscapes. (I
Bartomeus, Ed.). Journal of Ecology **109**, 1747–1757.

Tonnabel J, David P, Janicke T, Lehner A, Mollet J-C, Pannell JR, Dufay M. 2021. The
Scope for Postmating Sexual Selection in Plants. Trends in Ecology & Evolution **36**, 556–
567.

Toprak U, Hegedus D, Doğan C, Güney G. 2020. A journey into the world of insect lipid
metabolism. Archives of Insect Biochemistry and Physiology **104**, e21682.

Troscianko J, Stevens M. 2015. Image calibration and analysis toolbox – a free software
suite for objectively measuring reflectance, colour and pattern. Methods in Ecology and
Evolution **6**, 1320–1331.

Trunschke J, Lunau K, Pyke GH, Ren Z-X, Wang H. 2021. Flower Color Evolution and the
Evidence of Pollinator-Mediated Selection. Frontiers in Plant Science **12**.

Twayana M, Girija AM, Mohan V, Shah J. 2022. Phloem: At the center of action in plant
defense against aphids. Journal of Plant Physiology **273**, 153695.

van Putten WF, Elzinga JA, Biere A. 2007. Host Fidelity of the Pollinator Guilds of *Silene*
dioica and *Silene latifolia*: Possible Consequences for Sympatric Host Race Differentiation of
a Vectored Plant Disease. International Journal of Plant Sciences **168**, 421–434.

Villacañas de Castro C, Hoffmeister TS. 2020*b*. Friend or foe? A parasitic wasp shifts the
cost/benefit ratio in a nursery pollination system impacting plant fitness. Ecology and
Evolution **10**, 4220–4232.

Waelti MO, Page PA, Widmer A, Schiestl FP. 2009*a*. How to be an attractive male: floral
dimorphism and attractiveness to pollinators in a dioecious plant. BMC Evolutionary Biology
9, 190.

Waelti MO, Page PA, Widmer A, Schiestl FP. 2009*b*. How to be an attractive male: floral
dimorphism and attractiveness to pollinators in a dioecious plant. BMC Evolutionary Biology
9, 190.

Wang M, Carver JJ, Phelan VV, *et al.* 2016. Sharing and community curation of mass
spectrometry data with Global Natural Products Social Molecular Networking. Nature
Biotechnology **34**, 828–837.

Witt T, Jürgens A, Gottsberger G. 2013. Nectar sugar composition of European
Caryophylloideae (Caryophyllaceae) in relation to flower length, pollination biology and
phylogeny. Journal of Evolutionary Biology **26**, 2244–2259.

Wolfe LM, Elzinga JA, Biere A. 2004. Increased susceptibility to enemies following
introduction in the invasive plant *Silene latifolia*. Ecology Letters **7**, 813–820.

Wong DCJ, Pichersky E, Peakall R. 2023. Many different flowers make a bouquet: Lessons
from specialized metabolite diversity in plant–pollinator interactions. Current Opinion in Plant
Biology **73**, 102332.

Yoshinaga N. 2016. Physiological function and ecological aspects of fatty acid-amino acid
conjugates in insects†. Bioscience, Biotechnology, and Biochemistry **80**, 1274–1282.

1141

Tables

Table 1: Overview of statistical analyses investigating the effects of aphid infestation treatment (atm) with *Brachycaudus lychnidis* and *Silene latifolia* plant sex on plant stress indicators, behaviour of the pollinator *Hadena bicruris* and plant traits involved in communication and resource exchange with pollinators.

	(1) Sex	(2) Atm	(3) Cofactor†	(4) Sex × Atm	(5) Sex × Cofactor†	(6) Atm × Cofactor†	(7) Sex × Amt. × Cofactor†	(8) Sex moth	Random effects
Aphid number None, GLMM, NB, none	0.328	-	**0.000**	-	-	0.207	-	-	Plant individual
Aphid weight None, LM, G, none	0.535	-	-	-	-	-	-	-	-
Leaf biomass None, LM, G, none	**0.035**	0.516	-	0.397	-	-	-	-	-
Chlorophyll Log, LMM, G, D~(1-7)	**0.000**	0.774	**0.000**	0.206	0.530	0.495	0.938	-	Plant individual
Absorption by Flavonoids None, LMM, G, none	**0.028**	0.637	**0.000**	0.167	0.421-	0.539	0.448	-	Plant individual
Pollinator choice C-A None, GLMM, B, none	**0.000**	**0.072**	0.986	††	0.301	††	††	**0.097**	Moth individual, trial
Pollinator choice F-M None, GLMM, BB, none	0.380	0.844	0.937	††	††	0.971	††	0.572	Moth individual, trial
Flower number None, GLMM, P, Z~ I D~(1)	**0.000**[ii]	**0.082**[ii]	**0.000**[ii]	0.238[ii]	0.757[ii]	**0.000**[ii]	0.012	-	Plant individual
Flower biomass None, LMM, G, D~(3)	**0.000**[ii]	**0.001**[ii]	**0.000**[ii]	0.123[ii]	**0.000**[ii]	**0.000**[ii]	0.000	-	Plant individual
Flower colour None, LMM, G, D~(1,3,5)	0.394[ii]	**0.000**[ii]	**0.000**[ii]	**0.027**	0.977	**0.016**	0.6908	-	Plant individual
Flower area None, GLMM, NB, D~(1-7)	0.348	0.092	**0.000**	0.645	0.124	0.942	0.53	-	Plant individual
Scent metabolome SQ PERMANOVA (Bray-Curtis)	0.568	**0.087**	-	0.472	-	-	-	-	-
Scent metabolome QO PERMANOVA (Bray-Curtis)	0.625	0.231	-	0.493	-	-	-	-	-
Nectar volume None, LMM, G, none	**0.000**	0.554	-	0.225	-	-	-	-	Plant individual
Nectar metabolome (LC) PERMANOVA (Bray-Curtis)	**0.000**	0.204	-	0.308	-	-	-	-	-

The table provides an overview of the model structures with regard to the included responses (column 1), predictors and random effects (row 1); as well as further model characteristics such as response transformations (log = natural logarithm, none = none), model types (GLMM, LMM, LM, PERMANOVA), the assumed error families (B = binomial, BB = betabinomial, G = Gaussian, NB = overdispersed Poisson with quadratic parametrization, P = Poisson), and additional zero-inflation / dispersion formulas (Z / D~ = account for access of

zeroes in Poisson data / account for predictor-specific variance inhomogeneity, none = none). Moreover, the table provides an overview of p-values from type III ANOVAs (based on Wald χ^2-test for (G)LMMs, F-tests for LMs, and permutation statistics with pseudo-F estimation for PERMANOVA). Significant predictor terms are highlighted with bold lettering. Main effects and lower order interaction terms involved in significant (higher order) interactions are marked with "ii". Effects that were not tested are marked with "-". Please note that (†) the identity of cofactors for each response is described in detail in chapter 3.3.6; and (††) that in pollinator behaviour models, the actual test whether pollinators prefer control (C) over aphid infested (A) plants or female (F) over male (M) plants is tested by assessing whether the model intercept differs from 0.5. This represents the deviation from an equal choice and is not expressed as an interaction term with other predictors. Group-specific patterns are thus evaluated through post-hoc comparisons of the estimated probabilities for the respective factor levels, which are described in the respective text and figures. SQ = semi-quantitative approach, QO = qualitative only approach.

Tables

Table 2: Overview of metabolites highlighted in Figure 5B.

ID Figure 5	Effect(s)	Effect direction(s)	Molecular formula	Compound class	Compound identity
1	Atm	↓	C7H6O	Benzenoids	Benzaldehyde
2	Atm	↓	C9H12O3	Mono-terpenoids	Epoxyoxoisophorone
3	Atm	↓	C10H16O3	Mono-terpenoids	Lilac alcohol formate A
4	Atm	↓	C10H16O3	Mono-terpenoids	Lilac alcohol formate B
5	Atm	↓	C9H16O2	Mono-terpenoids	Lilac alcohol D
6	Atm	↓	C9H14O2	Mono-terpenoids	Lilac aldehyde A
7	Atm	↓	C9H14O2	Mono-terpenoids	Lilac aldehyde B+C
8	Atm	↓	C10H16	Mono-terpenoids	trans-ß-Ocimene
9	Sex*Atm	F↓M↔	C15H14O2	Phenyl-propanoids	Phenylethyl benzoate
10	None	-	C7H8O2	Benzenoids	2-Methoxy phenol
11	None	-	C8H8O2	Benzenoids	Methyl benzoate
12	None	-	C8H8O3	Benzenoids	Methyl salicylate
13	None	-	C8H8O	Benzenoids	Phenylacetaldehyde
14	None	-	C8H10O	Benzenoids	2-Phenylethanol
15	None	-	C8H14O2	Fatty acid derivatives	cis-3-Hexenyl acetate
16	None	-	C9H16O2	Mono-terpenoids	Lilac alcohol A
17	None	-	C9H16O2	Mono-terpenoids	Lilac alcohol B + C
18	None	-	C9H14O2	Mono-terpenoids	Lilac aldehyde D
19	None	-	C5H11NO	N-bearing compounds	anti-3-Methyl-butyl-aldoxime
20	None	-	C5H11NO	N-bearing compounds	syn-3-Methyl-butyl aldoxime

Metabolites highlighted with IDs in Figure 5 were significantly (uncorrected p-value < 0.05) and remarkably ($\log_2$ fold change <-1 or >1) modulated by aphid infestation treatment (atm; ↓ = reduced, ↑ = increased, - = unaffected by aphid infestation) and sex (F=female, M=male) and/or have been shown to elicit behavioural or antennal responses in the pollinator *Hadena bicruris* according to (Dötterl et al., 2006). Metabolites are ordered according to treatment effects, putative compound classes, and putative metabolite identities (alphabetically). See Supplementary Table S1A for a full list of floral VOC.

1142   **Table 3:** Overview of metabolites highlighted in Figure 6C-F.

ID Fig 6	Compund class	Compound identity	Effect direction sex	Effect direction atm	ID S1
1	*Amines*	*Pyrrolidine derivate*	*C:F>M, A:F=M*	*F↓, M↔*	*ID0040*
2	Amino acids, peptides, analogues	† Amino alcohol diester	F>M	-	ID1870
3	Amino acids, peptides, analogues	†Macrocyclic despipeptide	F<M	-	ID1489
4	Amino acids, peptides, analogues	N-(3-Methylbutanoyl)valine	F<M	-	ID0164
5	Amino acids, peptides, analogues	Peptide	F>M	-	ID1655
6	Amino acids, peptides, analogues	Peptide	F>M	-	ID1700
7	Amino acids, peptides, analogues	Peptide	F<M	-	ID0047
8	Amino acids, peptides, analogues	L-PheNH$_2$,Lys-Leu-Val-Phe	F<M	-	ID2889
9	Amino acids, peptides, analogues	DL-Phenylalanine	-	↓	ID0194
10	Amino acids, peptides, analogues	H-Ser-Leu$_5$-OH	F>M	-	ID1690
11	Amino acids, peptides, analogues	L-Tyrosine	F<M	-	ID0179
12	*Benzenesulfonyl compounds*	-	*C:F=M, A:F=M*	*F↓, M↔*	*ID0091*
13	Carbohydrates & conjugates	†Disaccharide glycosylglycerol ester	-	↓	ID0115
14	Carbohydrates & conjugates	D-Fructose	-	-	ID110_125_129
15	Carbohydrates & conjugates	D-Glucose	-	-	ID0000
16	Carbohydrates & conjugates	Geinistin	-	↓	ID0455
17	Carbohydrates & conjugates	2α-Mannobiose	-	-	ID0136
18	Carbohydrates & conjugates	Melezitose	-	-	ID71_142
19	Carbohydrates & conjugates	† Nucleoside analogue	C:F<M, A:F=M	F↔, M↓	ID0054
20	Carbohydrates & conjugates	†Nucleoside analogue	C:F>M, A:F=M	F↔, M↔	ID0111
21	Carbohydrates & conjugates	N-Acetylglucosamine	F>M	-	ID0162
22	Carbohydrates & conjugates	Nystose	-	-	ID0080
23	Carbohydrates & conjugates	Palatinose	-	-	ID0117
24	Carbohydrates & conjugates	†Phenolic acid glycoside	-	↓	ID0033
25	Carbohydrates & conjugates	Stachyose	-	↑	ID0140
26	Carbohydrates & conjugates	Sucrose	-	-	ID60_138
27	Carbohydrates & conjugates	-	-	↓	ID0050
28	Ceramides	† Bis(bishexyloxybiphenyl)amine	F<M	-	ID4068
29	*Ceramides*	*Hydroxylated ceramide*	*C:F>M, A:F=M*	*F↓, M↔*	*ID1697*
30	Ceramides	Palmitoleoyl ethanolamide	C:F<M, A:F=M	F↔, M↔	ID0990
31	*Ceramides*	*-*	*C:F=M, A:F=M*	*F↓, M↔*	*ID1771*
32	Fatty amides	† Aminoglycerol-Derivate	F>M	-	ID1842
33	Fatty amides	Fumaric acid diester	F>M	-	ID1952
34	Fatty amides	† GABA derivate	F<M	-	ID2723
35	Fatty amides	†Glyco-hydroxy fatty amide	F>M	-	ID1787
36	Fatty amides	-	F>M	-	ID1633
37	Flavonoid glycosides	† Spiroflavonoid glycoside	F<M	-	ID0293
38	Flavonoid glycosides	Vitexin-7-Rutinoside	F<M	-	ID0294
39	Flavonoid glycosides	-	F>M	-	ID1611
40	*Flavonoids*	*-*	*C:F>M, A:F=M*	*F↓, M↔*	*ID0131*
41	Glycerophosphates	† PG(O-20:1/17:1)	F>M	-	ID2141
42	Glycerophosphoethanolamines	† Alkyl-PE(C15:0)	C:F>>M, A:F>M	F↔, M↔	ID1683
43	Glycosphingolipids	† β-GalCer (C20mu:0)	F<M	-	ID2724
44	Phosphatidylethanolamine	† PE(16:0/18:sie2)	C:F=M, A:F<M	F↔, M↔	ID3072
45	Polyol	† Trimethyltrihydroxyundecanyl	F>M	-	ID1664
46	Purines and purine derivatives	† Guanosine derivate	F<M	-	ID0165
47	*Purines and purine derivatives*	*Xanthine derivate*	*C:F=M, A:F=M*	*F↓, M↔*	*ID0103*
48	Terpene glycosides	-	F>M	-	ID2302
49	Triterpenoids	† Triterpenoid steroidal glycosides	F>M	-	ID2436
50	Triterpenoids	† Triterpenoid steroidal glycosides	F>M	-	ID2372

Metabolites highlighted with IDs in Figure 6C-F were saccharides (primary nutritional compounds) or putatively identified compounds that were significantly (FDR corrected [bold lettering] / uncorrected p-value <0.05) and remarkably (log$_2$ fold change <-1 or >1) modulated

by plant sex (F=female, M=male, entries only in column 2), aphid infestation treatment (atm; ↓ = reduced, ↑ = increased under aphid infestation, entries only in column 3) or the interaction plant sex × aphid infestation treatment (entries in column 2 and 3). The abundance of features printed in italic correlates with moth behaviour (Figure 3). Metabolites are listed with their compound class, putative identity, and ID in the full dataset Supplementary Table S1B. †: IUPAC names for the regarding metabolites are replaced with simplified names for compact presentation and fully listed S1B.

Figure captions

Figure 1: Plant-pollinator-herbivore study system. (A) *Silene latifolia* (female), (B) *Hadena bicruris* (male), (C) adult *Brachycaudus lychnidis* individual and colony feeding on flowering stems of S. latifolia.

Figure 2: Overview of (A) the experimental setup (abbreviations: FC = control female, FA = aphid infested female, MC = control male, MA = aphid infested male); the data acquisition for (B) the plant stress indicators (i) absorbance by chlorophyll and epidermally located flavonoids and (ii) aphid load in terms of weight and number, (C) pollinator behaviour, (D) visual flower traits related to (i) color, (ii) quantity and (iii) display area, (E) floral scent, and (F) flower nectar; as well as (G) the time line for data acquisition after inoculation of plants with aphids.

Figure 3: Effects of *Silene latifolia* plant sex and infestation by *Brachycaudus lychnidis* on the choice behaviour of the moth pollinator *Hadena bicruris* are analysed and illustrated as proportions of flower visits, feeding attempts and the total time spent between two offered flower types. (A) Illustrates preference (i.e., deviation from a binomial outcome of 0.5 marked by the dashed line) for control (C, deep coloured, outcome > 0.5) versus aphid infested plants (A, light coloured, < outcome 0.5) in *S. latifolia* females (F, red) and males (M, turquoise) (FC/FA: n=11, MC/MA: n=22). (B) Illustrates preference for female (F, red, outcome > 0.5) versus male (M, turquoise, outcome < 0.5) control (C, dark coloured) and aphid infested (A, light coloured) plants (FC/MC: n=19, FA/MA: n=9). Graphs combine Box-Whisker plots of raw data (box: IQR Q1-Q3, whiskers: smallest/largest values within Q1/Q3 -/+ 1.5 × IQR, points: outliers) with the marginal means +/- standard error (black squares with black lines) estimated for all groups by the (G)LMMs. Effects assessed via type III ANOVAs based on Wald χ^2 tests are denoted at the top of the plot, while the results of relevant post-hoc comparisons are denoted within the plots (significance levels: ***p<0.001, **p<0.01, *p<0.05, -: not significant). Asterisks printed on top of lines indicate differences of binomial outcomes among the connected treatment groups, while asterisks printed on top of individual boxes indicate significant deviations of the binomial outcome from 0.5 (i.e., preferences within a specific treatment group).

Figure 4: Combined effects of *Silene latifolia* plant sex and *Brachycaudus lychnidis* aphid infestation treatment on (A) flower numbers, (B) reproductive biomasses, and (C) flower colours. The graphs combine Box-Whisker plots of raw data (box: IQR Q1-Q3, whiskers: smallest/largest values within Q1/Q3 -/+ 1.5 × IQR, points: outliers) for female control (FC, dark red, n = $5_{(A,B)}$/$14_{(C)}$), female infested (FA, light red, n = $5_{(A,B)}$/$7_{(C)}$), male control (MC, dark turquoise, n = $5_{(A,B)}$/$13_{(C)}$), and male infested (MA, light turquoise, n = $5_{(A,B)}$/$13_{(C)}$) plants with the marginal means +/- standard error (black squares) for these groups estimated by the (G)LMMs. Significant effects detected via type III ANOVAs based on Wald χ^2 tests are

denoted at the top of the plot, while the results of relevant post-hoc comparisons are denoted within the plots (significance levels: ***$p<0.001$, **$p<0.01$, *$p<0.05$, - not significant).

Figure 5: Combined effects of *Brachycaudus lychnidis* aphid infestation treatment and plant sex on floral headspace VOC composition in *Silene latifolia.* (A) The NMDS ordination for female control (FC, dark red, n = 4), female infested (FA, light red, n = 4), male control (MC, dark turquoise, n = 5), and male infested (MA, light turquoise, n = 5) plants shows plant individual scores (points), distances to group centroids (spiders), and their standard error (ellipses). (B) Volcano plot summarizing the results of model loops, i.e., the main and interactive effects of plant sex and aphid infestation treatment on the abundance of individual metabolites. Metabolic features modulated by aphid infestation are highlighted ($\log_2$ fold change <-1 or >1 and uncorrected [after FDR correction all effects non-significant] p-value <0.05 = coloured) with their effect direction (increased | decreased under aphid infestation (↑ | ↓) = brown | beige). A single metabolic feature (ID 9) was shaped by the interaction aphid infestation treatment × plant sex and is illustrated with one point for female (F) and male (M) plants. Numbered metabolic features correspond to Table 2. For further details see Supplementary Table S1A.

Figure 6: Combined effects of *Silene latifolia* plant sex and *Brachycaudus lychnidis* aphid infestation treatment on (A) nectar volume, and (B-F) the metabolic composition of a standardized nectar volume. (A) Box-Whisker plots of raw data for female control (FC, dark red, n = 23), female infested (FA, light red, n = 13), male control (MC, dark turquoise, n = 25), and male infested (MA, light turquoise, n = 25) plants with the marginal means +/- standard error (black squares) for these groups estimated by the (G)LMMs. Significant effects detected via type III ANOVAs based on Wald χ^2 tests are denoted at the top of the plot, while the results of relevant post-hoc comparisons are denoted within the plots (significance levels: ***$p<0.001$, **$p<0.01$, *$p<0.05$). (B) The NMDS ordination for FC (n = 5), FA (n = 3), MC (n = 5), and MA (n = 5) plants shows plant individual scores (points), distances to group centroids (spiders), and their standard error (ellipses).(C-F) Volcano plots illustrate the results of model loops, i.e., the main and interactive effects of plant sex and aphid infestation treatment on the abundance of individual metabolic features. Modulated features are highlighted (FDR corrected [uncorrected] p-value <0.05 and $\log_2$ fold change <-1 or >1 = coloured with black margin [coloured without margin]) with their effect direction (plant sex: F>M = red, M>F = turquoise, aphid infestation treatment: increased under aphid infestation (↑) = brown, decreased (↓) = beige). Interaction effect plots adopt the same modulation colour coding for (E) the sex effect within control and aphid infested plants and (F) the aphid infestation effect within females and males. Identical features within these group comparisons are connected by lines. Numbered metabolic features correspond to Table 3. A maximum of 15 identified

features was highlighted for each modulation direction. For further details see Supplementary Table S1B.

Figure 1: Plant-pollinator-herbivore study system. (A) *Silene latifolia* (female), (B) *Hadena bicruris* (male), (C) adult *Brachycaudus lychnidis* individual and colony feeding on flowering stems of *S. latifolia*.

Figure 2: Overview of (A) the experimental setup (abbreviations: FC = control female, FA = aphid infested female, MC = control male, MA = aphid infested male); the data acquisition for (B) the plant stress indicators (i) absorbance by chlorophyll and epidermally located flavonoids and (ii) aphid load in terms of weight and number, (C) pollinator behaviour, (D) visual flower traits related to (i) color, (ii) quantity and (iii) display area, (E) floral scent, and (F) flower nectar; as well as (G) the time line for data acquisition after inoculation of plants with aphids.

Figure 3: Effects of *Silene latifolia* plant sex and infestation by *Brachycaudus lychnidis* on the choice behaviour of the moth pollinator *Hadena bicruris* are analysed and illustrated as proportions of flower visits, feeding attempts and the total time spent between two offered flower types. (A) Illustrates preference (i.e., deviation from a binomial outcome of 0.5 marked by the dashed line) for control (C, deep coloured, outcome > 0.5) versus aphid infested plants (A, light coloured, < outcome 0.5) in S. latifolia females (F, red) and males (M, turquoise) (FC/FA: n=11, MC/MA: n=22). (B) Illustrates preference for female (F, red, outcome > 0.5) versus male (M, turquoise, outcome < 0.5) control (C, dark coloured) and aphid infested (A, light coloured) plants (FC/MC: n=19, FA/MA: n=9). Graphs combine Box-Whisker plots of raw data (box: IQR Q1-Q3, whiskers: smallest/largest values within Q1/Q3 -/+ 1.5 × IQR, points: outliers) with the marginal means +/- standard error (black squares with black lines) estimated for all groups by the (G)LMMs. Effects assessed via type III ANOVAs based on Wald χ^2 tests are denoted at the top of the plot, while the results of relevant post-hoc comparisons are denoted within the plots (significance levels: ***p<0.001, **p<0.01, *p<0.05, -: not significant). Asterisks printed on top of lines indicate differences of binomial outcomes among the connected treatment groups, while asterisks printed on top of individual boxes indicate significant deviations of the binomial outcome from 0.5 (i.e., preferences within a specific treatment group).

Figure 4: Combined effects of *Silene latifolia* plant sex and *Brachycaudus lychnidis* aphid infestation treatment on (A) flower numbers, (B) reproductive biomasses, and (C) flower colours. The graphs combine Box-Whisker plots of raw data (box: IQR Q1-Q3, whiskers: smallest/largest values within Q1/Q3 -/+ 1.5 × IQR, points: outliers) for female control (FC, dark red, n = 5(A,B)/14(C)), female infested (FA, light red, n = 5(A,B)/7(C)), male control (MC, dark turquoise, n = 5(A,B)/13(C)), and male infested (MA, light turquoise, n = 5(A,B)/13(C)) plants with the marginal means +/- standard error (black squares) for these groups estimated by the (G)LMMs. Significant effects detected via type III ANOVAs based on Wald χ^2 tests are denoted at the top of the plot, while the results of relevant post-hoc comparisons are denoted within the plots (significance levels: ***p<0.001, **p<0.01, *p<0.05, - not significant).

Figure 5: Combined effects of *Brachycaudus lychnidis* aphid infestation treatment and plant sex on floral headspace VOC composition in *Silene latifolia*. (A) The NMDS ordination for female control (FC, dark red, n = 4), female infested (FA, light red, n = 4), male control (MC, dark turquoise, n = 5), and male infested (MA, light turquoise, n = 5) plants shows plant individual scores (points), distances to group centroids (spiders), and their standard error (ellipses). (B) Volcano plot summarizing the results of model loops, i.e., the main and interactive effects of plant sex and aphid infestation treatment on the abundance of individual metabolites. Metabolic features modulated by aphid infestation are highlighted (log2 fold change <-1 or >1 and uncorrected [after FDR correction all effects non-significant] p-value <0.05 = coloured) with their effect direction (increased | decreased under aphid infestation (↑ | ↓) = brown | beige). A single metabolic feature (ID 9) was shaped by the interaction aphid infestation treatment × plant sex and is illustrated with one point for female (F) and male (M) plants. Numbered metabolic features correspond to Table 2. For further details see Supplementary Table S1A.

Figure 6: Combined effects of *Silene latifolia* plant sex and *Brachycaudus lychnidis* aphid infestation treatment on (A) nectar volume, and (B-F) the metabolic composition of a standardized nectar volume. (A) Box-Whisker plots of raw data for female control (FC, dark red, n = 23), female infested (FA, light red, n = 13), male control (MC, dark turquoise, n = 25), and male infested (MA, light turquoise, n = 25) plants with the marginal means +/- standard error (black squares) for these groups estimated by the (G)LMMs. Significant effects detected via type III ANOVAs based on Wald χ^2 tests are denoted at the top of the plot, while the results of relevant post-hoc comparisons are denoted within the plots (significance levels: ***p<0.001, **p<0.01, *p<0.05). (B) The NMDS ordination for FC (n = 5), FA (n = 3), MC (n = 5), and MA (n = 5) plants shows plant individual scores (points), distances to group centroids (spiders), and their standard error (ellipses).(C-F) Volcano plots illustrate the results of model loops, i.e., the main and interactive effects of plant sex and aphid infestation treatment on the abundance of individual metabolic features. Modulated features are highlighted (FDR corrected [uncorrected] p-value <0.05 and log2 fold change <-1 or >1 = coloured with black margin [coloured without margin]) with their effect direction (plant sex: F>M = red, M>F = turquoise, aphid infestation treatment: increased under aphid infestation (↑) = brown, decreased (↓) = beige). Interaction effect plots adopt the same modulation colour coding for (E) the sex effect within control and aphid infested plants and (F) the aphid infestation effect within females and males. Identical features within these group comparisons are connected by lines. Numbered metabolic features correspond to Table 3. A maximum of 15 identified features was highlighted for each modulation direction. For further details see Supplementary Table S1B.

Graphical Abstract:

Plant-sex specific effect of aphid infestation on floral traits (number, size, colour, scent composition, nectar quantity and composition) and pollinator behaviour.