Selection of a Lead Long-Acting Formulation of Ivermectin to Target Major Malaria Vectors

in Western Africa: Evaluation of Pharmacokinetics and Mosquitocidal Efficacy in Cattle

under Laboratory Conditions.

Lamidi Zela[1,2], Sié Hermann Pooda[1,3], Angélique Porciani[4], Samuel Beneteau[5], André

Barembaye Sagna[4,6*], Sophie Le Lamer-Déchamps[7], Nicolas Moiroux[4,8], Cheick Oumar

Wendpagnandé Ouédraogo[9,10], Anyirékun Fabrice Somé[9], Cédric Pennetier[4], Christophe

Roberge[7], Adrien Marie Gaston Belem[2], Koumbobr Roch Dabiré[9], Charles Dayo[1], Karine

Mouline[4**]

Affiliations :

1. Centre International de Recherche-Développement sur l'Elevage en zone Subhumide,

13   Bobo-Dioulasso, Burkina Faso

14   2. Université Nazi Boni, Bobo-Dioulasso, Burkina Faso

15   3. Université Daniel Ouezzin COULIBALY, Dédougou

16   4. MIVEGEC, University of Montpellier, IRD, CNRS, Montpellier, France

17   5. Independant Statistician

18   6. IRD, Représentation du Burkina Faso, Antenne de Bobo-Dioulasso, Bobo-Dioulasso,

19   Burkina Faso

20   7. MedinCell, Montpellier, France

21   8. Pôle de Zoologie Médicale, Institut Pasteur de Dakar, Dakar, Sénégal

22   9. Institut de Recherche en Sciences de la Santé, Direction Régionale de l'Ouest, Centre

23   National de Recherche Scientifique et Technologique, Bobo-Dioulasso, Burkina Faso

10. Université Cheikh Anta Diop, Dakar, Sénégal

Lamidi Zela, lamidase@yahoo.fr,

Sié Hermann Pooda, poodasiehermann@yahoo.fr,

Angélique Porciani, angelique.porciani@ird.fr,

Samuel Beneteau, samuel.beneteau@ntymail.com

André Sagna, andre.sagna@ird.fr,

Sophie Le Lamer-Déchamps, sophie.lelamer@medincell.com,

Nicolas Moiroux, nicolas.moiroux@ird.fr

Cheick Omar Wendpagnandé Ouédraogo, cheickoumaroued92@yahoo.fr,

Anyirékun Fabrice Somé, afabricesome@yahoo.fr,

Cédric Pennetie, cedric.pennetier@ird.fr

Christophe Roberge, christophe.roberge@medincell.com,

Adrien M. G. Belem, belemadrien@yahoo.fr,

Roch Dabiré, dabireroch@gmail.com,

Charles Dayo, charlesdayo@yahoo.fr

Karine mouline**, karine.mouline@ird.fr

* present address: ASTRE, CIRAD-INRAE, Univ Montpellier, Centre de Recherche et de Veille

sur les Maladies Vectorielles dans la Caraïbe (CRVC), Petit Bourg, Guadeloupe, France

** corresponding author

ABSTRACT

Background

Ivermectin, a semisynthetic endectocide from the avermectin family, has been used in mass

drug administration (MDA) campaigns since the 1980s to control helminth infections. Its

lethality to *Anopheles* mosquitoes after feeding on treated hosts offers a promising malaria

control strategy, especially for outdoor transmission. However, existing oral formulations

produce short-duration mosquitocidal blood concentrations, limiting epidemiological impact.

The WHO's target product profile for endectocides calls for a hazard ratio >4 and sustained

efficacy for at least one month. In this context, three long-acting injectable ivermectin

formulations (LAIF) based on BEPO® depot technology were evaluated in cattle to evaluate

the *in vivo* performance then to identify the most effective candidate.

Methods

The LAI candidates (mdc-STM-001, -002, and -003) were tested at 0.6 mg/kg, with mdc-STM-

003 also evaluated at 1.5 mg/kg. Each treatment group consisted of five calves, each receiving

single sub-cutaneous injection. Plasma samples intensively collected over 130 days enabled

non-compartmental pharmacokinetic analysis of ivermectin. Two mosquito strains: *Anopheles*

coluzzii (VK5-pyrethroid-resistant) and *Anopheles gambiae* (KIS-Kisumu strain) were allowed

to simultaneously blood-feed on treated animals from 2 to 126 days post-dose. Mosquito

survival was monitored for 30 days. Mortality within 4 and 10 days post-feeding was

emphasized, targeting the window before *Plasmodium falciparum* becomes transmissible.

Efficacy was assessed via hazard ratios, cumulative mortality, lethal concentrations (LC) 50

and 90 values, and duration of effective coverage.

Results

All formulations were well tolerated. mdc-STM-001 showed the most favorable pharmacokinetic profile with a controlled peak (Cmax= 34.5 ± 12.7 ng/mL) and the lowest inter-individual variability (12%). mdc-STM-001 and -003 produced hazard ratios >4 and cumulative mortalities >50% for at least 60 days in both mosquito strains. Surviving mosquitoes had median lifespans below 10 days for at least 90 days post-injection. The 10-day LC50 for resistant mosquitoes (3.66 [2.69-4.97]) was maintained for ≥126 days. mdc-STM-001 provided 100% coverage at this threshold for ≥2 months.

Conclusion

mdc-STM-001 is the lead candidate. A single injection allows killing >50% of biting mosquitoes before they become infectious, even in resistant populations, during at least 2 months. Given ivermectin's established safety profile, its repurposing in LAIFs offers a promising strategy for malaria control. This approach directly targets two major challenges: outdoor transmission and rising insecticide resistance, and warrants further investigation.

Keywords: residual malaria, outdoor transmission, mosquitoes resistance, ivermectin, repurposing, long-acting technology, pharmacokinetics, efficacy, impact

BACKGROUND

Progress in the fight against malaria has stagnated since 2015, following a remarkable 15-years of sustained decrease in disease incidence, largely driven by the widespread deployment of two core insecticide-based vector control tools: long-lasting insecticidal nets (LLINs) and insecticide residual spraying (IRS) [1]. While these methods remain effective,

their impact is increasingly undermined by the large-scale development of physiological

resistance in mosquito vectors to insecticidal compounds [2], as well as by behavioral

avoidance of treated surfaces [3]. New generations of LLINs incorporating dual active

ingredients or synergists have been introduced to address physiological resistances [4] [5].

However, these tools primarily target nocturnal, indoor-biting *Anopheles* mosquitoes are still

bypassed by vectors that feed outdoors and/or at times when people are not protected by

the nets [6][7], which leads to persistent residual malaria transmission.

Malaria therefore remains a leading global health concern with an estimated 263 million

clinical cases and 597,000 deaths in 2023, over 95% of which occurred in Sub-Saharan Africa

[8]. The resurgence in malaria cases in recent years highlights the urgent need for novel and

complementary strategies to reinforce existing interventions and reverse this concerning

trend [8]. In particular, tools targeting the diverse biting behaviors of *Anopheles* vectors,

whether innate or induced by interventions, are essential for addressing this challenge [9].

Mass drug administration (MDA) of endectocides, compounds active against both endo- and

ectoparasites, offers an approach that circumvent the limitations of current vector control

strategies, as it is independent of mosquito feeding time and location [10] : the insecticide is

delivered through the bloodstream of the treated host, making it effective regardless of the

vector's biting behavior. Ivermectin is the leading candidate among endectocides for vector

control, due to its extensive clinical use against lymphatic filariasis and onchocerciasis, with a

strong safety profile [11] [12]. Ivermectin binds to glutamate-gated chloride channels unique

to invertebrates, inducing neuromuscular hyperpolarization, paralysis and flaccid death [13].

This distinct mode of action makes ivermectin effective against mosquitoes physiologically

resistant to standard insecticides [14][15]. To date, it has shown to be lethal to all tested

Anopheles species [12], including nine Southeast Asian malaria vectors recently added to the

list of susceptible species [16]. Besides direct acute effects on survival induced by lethal

plasma concentrations, sub-lethal effects are also interesting to transmission control as they

negatively impact vectors fecundity and fertility [17][18], locomotor and biting behaviors

[19][20], or *Plasmodium* parasite development [21][22].

Standard oral dosing for parasitic disease treatment (150–200 µg/kg) achieves lethal

plasma levels for *Anopheles* for fewer than seven days [23]. The first community-level clinical

trial designed to test repeated ivermectin-MDA used this dosage, administered every 3

weeks during the transmission season [24]. This regimen resulted in a significant 20%

reduction of malaria incidence in children under five, however, it was subsequently

identified that extending ivermectin's duration of action could substantially improve impact

and logistics. Higher oral regimens (1x400 µg/kg or 3x300 µg/kg) have been demonstrated

safe and with extended mosquitocidal effects for at least 10-14 days on laboratory colonies

[25, 26, 27], with modelling suggesting up to 70% reduction in malaria incidence in seasonal

transmission settings [28]. A two-country trial using the 1x400 µg/kg is completed with

results pending [29]. However, two Phase 3 trials testing repeated ivermectin MDA (3x 300

µg/kg) in combination with either dihydroartemisinin-piperaquine (DHAP) [30] or seasonal

malaria chemoprophylaxis (SMC) [15] did not show significant epidemiological effects.

Explanatory factors discussed include underestimated wild mosquito tolerance, shorter-

than-expected ivermectin residence time (as demonstrated in the RIMDAMAL II trial, where

plasma levels dropped below detection in nearly 50% of participants by day 14) [15] and

overlooked aspects of mosquitoes bionomics, such as zoophagy [31].

Long-acting formulations that sustain effective plasma concentrations over extended periods

could overcome these limitations. Commercial veterinary injectables have demonstrated

mosquitocidal effects lasting 1–2 months [16], and BEPO®-based long-acting injectable

formulation (LAIF) prototypes [32] have maintained efficacy for over six months in cattle

after a single injection [33,34]. Comparable durations have been achieved with silicone

implants [35,36]. The only human-targeted long-acting formulation to date—a gastric-

resident oral device—has shown efficacy with ~14-day residence time in pigs [37].

We conducted a preclinical study to characterize the pharmacokinetic profile and

mosquitocidal efficacy of three LAIF based on BEPO® technology [32], developed for MDA

use in malaria endemic regions. The ivermectin formulations were designed to achieve a

molecule residence time of at least 2 months following a single injection, with sustained

plasma concentrations sufficient to ensure mosquitocidal efficacy, achieving a hazard ratio

(HR) > 4 as recommended by the World Health Organization (WHO) for endectocides in

malaria vector control [38]. Efficacy was further assessed via complementary entomological

endpoints, including mosquito median survival time, 4-day and 10-day cumulative mortality,

lethal concentrations 90 and 50 (LC90 and LC50) values, and corresponding duration of

coverage. A cattle-mosquito colony model was used [17][34], exposing both insecticide-

susceptible rand pyrethroid-resistant *Anopheles* strains. Comparative analysis of

pharmacokinetic and efficacy outcomes allowed us to identify the lead candidate for further

development toward clinical evaluation in humans.

METHODS

MOSQUITO COLONIES

Two *Anopheles* mosquito colonies were used in this study:

(i)   ***Anopheles (An.) gambiae* Kisumu (KIS)**: This World Health Organization (WHO)

163   reference strain is fully susceptible to all classes of insecticides. It has been

maintained for over a decade in the insectary at the Institut de Recherche en

Sciences de la Santé (IRSS) in Bobo-Dioulasso, Burkina Faso. The colony is

routinely monitored for contamination by other species and screened for

insecticide resistance using molecular techniques [39,40]. For the experiments,

mosquitoes were transported in meshed cages covered with moist cloth to the

Centre International de Recherche-Développement sur l'Élevage en zone Sub-

humide (CIRDES) using an air-conditioned vehicle to maintain stable temperature.

Transport duration was approximately 10 minutes.

(ii)   ***An. coluzzii* VK5-IMPACT (VK5)**: This colony was established at CIRDES in

November 2020, prior to the start of the experiments in January 2021. A total of

150 blood-fed females were collected from the village of VK5 in the Kou Valley

(~50 km northwest of Bobo-Dioulasso), an area characterized by a high

prevalence of insecticide resistance [41]. Collected females were allowed to

individually oviposit. Species identification was performed post-oviposition by

PCR using established protocols [39] and only progeny from confirmed *An.*

coluzzii—the primary malaria vector in Burkina Faso—were retained. As expected

based on the collection site, all females belonged to the target species [41].

Both colonies were housed in similar environmental conditions and maintenance

protocol. The insectariums temperature was set to 26°C ± 1°C, relative humidity to 75% ±

5%, and photoperiod to 12-hour light/dark. Adult mosquitoes were maintained on 10%

sterile glucose solution provided on cotton balls. For egg production, females were fed

on rabbits two to three times per week. Eggs were placed in trays (32 x 13 x 5 cm) filled

with tap water. After hatching, larvae were transferred to new trays (n=200 larvae /tray)

using Pasteur pipettes and fed ad libitum with commercial alevin food (TetraMin® Baby).

Breeding was conducted in 25 x 25 x 25 cages containing mixed-sex adults. The VK5

colony's insecticide resistance status was routinely monitored by sampling

approximatively 40 females from each generation and screening for the KDR-W mutation

using quantitative PCR (qPCR) ([40]). Routine maintenance included 10-15 cages

containing 100-200 females each to ensure a stable production of experimental

mosquitoes. For each direct skin-feeding assay time point, females were randomly drawn

from at least five different cages to minimize potential cage effects.

LONG ACTING IVERMECTIN FORMULATIONS

The long-acting ivermectin formulations (LAIFs) tested in this study were developed using

BEPO® technology [32], a proprietary long-acting injectable technology (Medincell, Jacou,

France). Following subcutaneous injection, BEPO® forms an *in situ* bioresorbable polymeric

depot via a solvent-exchange mechanism, leading to precipitation of a polymer matrix that

enables sustained drug release. Each formulation consisted of three components: i) a

biocompatible solvent to ensure injectability, ii) ivermectin, as the active pharmaceutical

ingredient, and iii) a mixture of two biocompatible and biodegradable polymers that control

drug release kinetics. These polymers undergo progressive hydrolysis into bioabsorbable by-

products over time, thanks to a simple hydrolysis mechanism. Three ready-to-use LAIFs

solutions were evaluated: **mdc-STM-001**, **mdc-STM-002** and **mdc-STM-003**; the last one

differed in ivermectin loading and each differed in polymer composition and proportions.

All formulations were imported into Burkina Faso with clearance from the Direction

Générale des Services Vétérinaires of Burkina Faso (Permit No 2020/199) and were stored at

room temperature, protected from light, until administration to cattle. The study protocol

was approved by the Institutional Animal Ethics Committee of CIRDES (document reference

number: 004-10/2020 CEB-CIRDES).

CATTLE HOSTS AND FORMULATIONS' ADMINISTRATION

Twenty-five male calves (Métis strain: Fulani zebus x Baoulé crossbreed) with a median

weight of 117 kg (interquartile range [IQR]: 106-129 kg), were purchased from villages in the

Valley du Kou area two months prior to the study to allow for acclimation. Upon arrival at

the CIRDES animal facility, calves were treated for trypanosomiasis and gastrointestinal

parasites using Berenil 2000® and Benzal®, respectively. Animals were housed outdoors

under screened enclosures to prevent exposure to or reinfestation with ectoparasites and

trypamosomoses. They were fed with rice straw and 1 kg/day of cottonseed cake, with

water and sea salt provided ad libitum.

Calves were randomly assigned to five experimental arms (n = 5 calves per arm) using

stratified randomization to balance body weight across groups. Four arms received LAIFs,

and one served as the untreated control. The formulations were administered *via* a single

subcutaneous injection into the loose skin anterior to the shoulder.

- mdc-STM-001 and mdc-STM-002 LAIFs were each injected at a unique dose of 0.6

mg/kg (referred to as mdc-STM-001-0.6 and mdc-STM-002-0.6, respectively).

- mdc-STM-003 was administered at two doses: 0.6 mg/kg and at 1.5 mg/kg (referred

to as mdc-STM-003-0.6 and mdc-STM-003-1.5).

The 1.5 mg/kg dose represents the upper limit of feasible subcutaneous volumes for

human application and was included to assess dose-dependent effects on

pharmacokinetics and mosquitocidal efficacy. Nominal injection volumes per animal

ranged from 0.62 to 2.14 mL (detailed in Table 1). The animals' identification numbers,

their weight upon arrival at CIRDES and on the day of injection, administered

formulations, and injected volumes are provided in the supplementary Table 1

(Appendix). Animal health was monitored throughout the study by assessing clinical

signs, local injection site reactions, body weight, and weight gain.

Table 1: Nominal treatment volumes per formulation to be administered to cattle. N/A:

not applicable.

Number of male cattle	LAI formulation name	Ivermectin dose	LAI nominal administered volume
5	mdc-STM-001	0.6 mg/kg	7.72 μL/kg
5	mdc-STM-002	0.6 mg/kg	7.80 μL/kg
5	mdc-STM-003	0.6 mg/kg	5.85 μL/kg
5	mdc-STM-003	1.5 mg/kg	14.6 μL/kg
5	N/A (control)	0 mg/kg	0 μL/kg

MOSQUITO BLOOD-FEEDING BIOASSAYS

To ensure equivalent ivermectin exposure, mosquitoes from both KIS and VK5 colonies were

simultaneously exposed to the same cattle hosts during blood-feeding. This design enabled

direct comparison of susceptibility across colonies under identical conditions. Direct skin

blood-feeding assays were conducted following previously established protocols [34]. Prior

to feeding, mosquitoes were deprived of sugar for 12 hours, with access to water-soaked

cotton only. At each time point, approximately 100–130 females per colony were introduced

into net-covered bowls, and held against the shaved flank of the selected calf for 20 minutes

using a rubber band. Separate bowls were used for each colony. Fully engorged mosquitoes

were distributed into cups (10 females per cup), with a maximum of 120 cups per colony per

time point. All cups were maintained under controlled conditions in the CIRDES insectary.

Mosquito mortality was recorded daily between 10:00 and 11:00 a.m. for 30 days post-blood

meal. To mitigate positional bias, cups were randomly repositioned on shelves after each

observation.

A total of 16 blood-feeding assays were conducted: once pre-treatment (baseline) and at 2,

7, 17, 21, 28, 42, 56, 70, 84, 91, 98, 105, 112, 119, and 126 Days After Injection (DAI). Due to

limited mosquito availability, KIS mosquitoes were not tested at 7 and 91 DAI. Surplus blood-

fed mosquitoes were stored at –20°C for downstream analyses.

IVERMECTIN BIOANALYSIS AND PHARMACOKINETICS

Jugular venous blood samples were collected from treated calves at each mosquito exposure

time point and at additional time-points (30 samples per animal) to characterize ivermectin

pharmacokinetics (PK). Sampling included pre-dose, 2h, 5h and 12h post-injection, then daily

sampling until DAI 4, every 2-4 days until DAI 21, and weekly until DAI 132-133. Blood

samples were also collected from control animals at pre-dose and at DAI 1, 14, 28, 56, 84

and 111. Blood samples were centrifuged at +4°C (2000 rpm for 10 min) to isolate plasma,

which was stored at -80°C until shipment for bioanalytical analysis. Ivermectin plasma

concentrations were quantified using a liquid-liquid extraction with qualified LC-MS/MS

methods. Two calibration ranges were used (0.2-20 ng/mL low range; 5-500 ng/mL high

range), using K_2-EDTA as anticoagulant and ivermectin-D_2 as internal standard. The quality

controls (0.6, 10, 16 ng/mL low range; 15, 250, 400 ng/mL high range) were within the

acceptance criteria (precision $\leq$ 15.00%, deviation $\pm$ 15.00%). Incurred sample reanalysis

validated ivermectin stability for at least 680 days at -20°C. Plasma samples collected from

cattle were stored at -80°C no longer than 162 days before analysis, validating the

ivermectin concentrations measured for PK analysis.

At study completion, control animals were returned to stock, while treated animals were

euthanized. Injection sites and depot with surrounding tissues were sampled to quantify the

residual ivermectin and copolymers. These analyses were performed using an ultra-

performance liquid chromatography (limit of quantification: 0.9 µg/mL) for ivermectin and

nuclear magnetic resonance analysis for copolymers. Both methods were developed for

exploratory purposes only.

OUTCOMES AND ANALYSIS

Unless specified otherwise, analysis were performed using the R version 4.4.3 (2025-02-

28) [42]. Model fit was assessed using the "performance" [43] and DHARMa [44] packages.

All estimates are reported with 95% confidence intervals (CI95%) and/or p-values.

288
Cattle weight gain

Differences in weight gain between treatment groups were analyzed using the Kruskal–Wallis

rank sum test, followed by pairwise two-sided exact Wilcoxon rank sum tests with 95%

confidence intervals. *p*-values were adjusted using the Holm method ([45]).

Pharmacokinetics

PK parameters of ivermectin were calculated for each LAIF using a non-compartmental

analysis (Phoenix® WinNonlin® version 8.1; Certara USA, Inc., Princeton, NJ). These included

the rate of systemic exposure (Cmax), the time to reach Cmax (Tmax), the extent of systemic

exposure over the study duration (aera under the curve, AUClast), over 28 DAI (AUC0-28d)

and over 91 DAI (AUC0-91d), concentrations at 28 DAI (C28d), 56 days (C56d), 91 DAI (C91d)

and at study end (Clast), and the apparent terminal phase half-life (T1/2) when calculable.

The systemic exposure parameters were normalized to the actual administered dose,

calculated for each cattle based on the weight of each syringe before and after injection. The

inter-individual variability in plasma concentrations and PK parameters was expressed as the

coefficient of variation (CV %) for each experimental group.

Mosquito related Outcomes

<u>Mosquito blood feeding rate</u>

For each colony, we compared the mosquitoes' blood-feeding rates across treatment arms

using a generalized linear mixed model (GLMM [46, 47]) with a Poisson error distribution.

The outcome variable was the number of blood-fed mosquitoes, with the total number of

mosquitoes included as an offset. The model included an interaction term between

treatment and DAI, as well as a random intercept for cattle. The model was fitted using

restricted maximum likelihood.

<u>Mosquitocidal efficacy</u>

Multiple complementary outcomes were used to assess the efficacy of each LAIF and to

allow comparison between formulations and mosquito strains. Depending on the outcome,

values are given for a 4-day or a 10-day follow up period. Outcomes for 30-day follow-up

period are given in the Appendix when specified in the results section.

Outcomes provided for the 4 days post-blood feeding period reflect the acute effect of the

ivermectin concentration absorbed by the mosquitoes, as the direct lethal effect of the

molecule typically manifests within 2-3 days post-ivermectin blood meal. In relation to the

extrinsic incubation period of *Plasmodium* parasites, we also measured efficacy outcomes

for the 10-day post-blood feeding period, in order to assess the treatment's potential to

prevent the build-up of infectious mosquitoes' populations. Extended 30-day follow-up

captures the residual effects of ivermectin on surviving mosquitoes, which may include both

indirect sublethal effects and potential interactions with the mosquitoes' age.

Cumulative Mortality
327
328   Proportions of dead mosquitoes at 4- and 10-days post-feeding were computed for each

329   colony, formulation, and DAI.

330   *Survival Analysis and Hazard Ratios*
331
Daily mosquito mortality was recorded for up to 30 days post blood-feeding. For each

mosquito colony and DAI, survival data were used to generate Kaplan–Meier survival curves,

stratified by treatment group.

Per colony, survival patterns at each time-point were compared to the control group using

Cox proportional hazards models (mixed-effects regression model [48]). The models

included "treatment group" and "DAI" as fixed effects, with random intercepts for individual

calf ID and mosquito cup (nested random effects) to account for variability within treatment

groups. Hazard ratios (HR) at different time points post-blood feeding are given to assess the

impact of ivermectin on mosquito mortality risks and HR variations over time. Post hoc

pairwise comparisons of HRs between time points were adjusted for multiple testing using

the Tukey method, implemented via the "emmeans" package [42]

Dose–Response Analysis and coverage duration

Dose–response relationships were analyzed using the "drc" package in R [49], modeling 4-

day and 10-day mosquito mortality as a function of host ivermectin plasma concentrations.

A four-parameter log-logistic model was fitted with the upper limit fixed at 1. Confidence

intervals were computed using the tfls method (t from logarithm scale).

RESULTS

Cattle follow up

A mild reaction (agitation) was observed in most animals after the injection, regardless of

dose or LAIF administered (data not shown). There were no unscheduled cattle deaths, and

no clinical signs were observed in any of the treated groups throughout the study. The major

local reaction was swelling, which occurred in all animals, as expected with this LAIF

technology. Median weight intake was unaffected by the test item treatment (Table 2). For

each experimental group, monthly weight measures are given in the supplementary Table 2

of the Appendix.

359
360   Table 2: Median weight intake (in kg) per group with interquartile range (IQR). Q1 and Q3

361   values are given in brackets. The p-value is the result of the Kruskal-Wallis rank sum test.

Treatment Group	Control N = 5	mdc-STM-001-0.6 N = 5	mdc-STM-002-0.6 N = 5	mdc-STM-003-0.6 N = 5	mdc-STM-003-1.5 N = 5	*p-value*
Median weight intake (kg)	27.4 (25.8, 31.6)	37.2 (35.8, 44.2)	39.6 (36.6, 41.6)	35.3 (31.2, 36.2)	30.0 (28.6, 40.0)	0.075

Ivermectin pharmacokinetic

A total of 635 blood samples were collected, including 600 samples from LAIF-treated cattle

(4 treatment groups, 5 cattle per group, n=30 per cattle) and 35 samples from control cattle

(1 control group, 5 cattle, n=7 per cattle). Concentration-time patterns were analyzed using

data until DAI = 132 for mdc-STM-001-0.6, mdc-STM-002-0.6 and DAI 133 for mdc-STM-003-

0.6 and mdc-STM-003-1.5. Ivermectin plasma concentrations were analyzed according to the

treatment (formulation and dose). Individual concentration-time profiles of ivermectin after

LAIFs injection are illustrated in Figure 1. Additional figures, such as the mean concentration-

time profiles over the study duration and those focused on the first 7 days to better visualize

the burst (i.e., peak plasma level), are provided in the supplementary figures 1A and 1B

(Appendix).

Figure 1: Individual kinetic profiles of ivermectin in cattle plasma after single sub-cutaneous

administration of candidate LAIFs at 0.6 or 1.5 mg/kg

Ivermectin was not detected in any of the samples collected in the control group during the

study period nor in any samples collected before ivermectin injection in the treated groups.

Following sub-cutaneous injection, ivermectin plasma concentrations were generally

measured in all samples collected during the study, with quantifiable levels still present at

the final study time-point (*i.e.,* DAI 132 or 133).

After a single sub-cutaneous administration of LAIFs, the plasma levels of ivermectin increased

to reach the maximal level (*i.e.,* the burst, known to occur with LAIs [32]) at 24h or 48h post-

dose (median Tmax). This was followed by two phases of decrease: a first phase until Day 56

(*i.e.,* 1344h post dose) then a slower phase with a regular decrease of ivermectin plasma levels

until the study end. Among the formulations, mdc-STM-001-0.6 showed the lowest rate of

decrease during the first phase and the steadiest levels during the second phase. In contrast,

the mdc-STM-002-0.6 exhibited the fastest decrease compared to other LAIFs. The inter-

animal variability in ivermectin plasma levels (expressed as coefficient of variation CV%) was

moderate. The lowest variability was seen with mdc-STM-001-0.6 (mean CV%: 41%, range: 12-

70%), followed by mdc-STM-003-1.5 (mean: 54%, range: 22-87). The highest variability was

observed for mdc-STM-002-0.6 (mean CV%: 62%, range: 25-106%) and mdc-STM-003-0.6

(mean: 72%, range: 34-98%).

The mean ivermectin PK parameters of each LAIF after injection are reported in Table 3.

Additional parameters are given in the supplementary Table 3 of the Appendix.

Table 3: Mean (SD) pharmacokinetic parameters of ivermectin in plasma cattle after single

sub-cutaneous administration of LAIF at 0.6 or 1.5 mg/kg

Parameters	mdc-STM-001-0.6	mdc-STM-002-0.6	mdc-STM-003-0.6	mdc-STM-003-1.5
Tmax* (h)	24 (24; 48)	48 (24; 168)	24 (24; 48)	24 (24; 168)
Cmax (ng/mL)	35.4 (12.7)	50.3 (13.7)	58.1 (25.2)	81.2 (32.2)
Cmax/Dose (ng/mL/mg/kg)	58.3 (22.0)	84.3 (22.6)	99.4 (42.2)	56.1 (22.4)
Clast (ng/mL)	3.93 (2.75)	1.03 (0.70)	3.43 (2.74)	6.30 (4.35)
AUC(0-91d) (h.ng/mL)	25055 (2936)	21121 (8866)	28890 (18668)	41820 (13286)
AUC(0-91d)/Dose (h.ng/mL/mg/kg)	40990 (5664)	35479 (14839)	49483 (31994)	28903 (9300)
C28d (ng/mL)	13.7 (1.6)	8.90 (4.85)	11.1 (8.3)	15.5 (5.1)
C56d (ng/mL)	6.70 (1.67)	2.92 (2.63)	7.34 (7.14)	10.3 (6.9)
C91d (ng/mL)	3.48 (1.85)	1.11 (0.84)	4.33 (3.99)	5.55 (3.72)

*Median (min; max).

At a dose of 0.6 mg/kg, PK differences emerged across formulations when considering both

the entire study duration or the 91-day period (*i.e.,* 3 months post-injection), which is the

best targeted release duration. These differences involved performance, quantification

levels over time, and inter-animal variability. The peak plasma concentration (Cmax) for

mdc-STM-001-0.6 was at least 30% lower compared to the other LAIFs, with mdc-STM-003-

0.6 showing the highest Cmax. This suggests that mdc-STM-001-0.6 provided a safer burst

compared to the other LAIFs. The inter-animal variability in Cmax was moderate for all

formulations, with the lowest variability observed for mdc-STM-002-0.6 (CV% = 27%)

followed by mdc-STM-001-0.6 (CV%=36%) and mdc-STM-003-0.6 (CV%=43%). The overall

systemic exposure over DAI 91 (AUC_{0-91d}) was quite similar for all formulations, with the

lowest value for mdc-STM-002-0.6 and the highest value for mdc-STM-003-0.6. Inter-animal

variability in AUC was much lower for mdc-STM-001-0.6 (CV% = 12%), compared to mdc-

STM-002-0.6 (CV% = 42%) and mdc-STM-003-0.6 (CV = 66%). The mean ivermectin plasma

concentrations after each month post sub-cutaneous injection (C28d, C56d, C91d, Clast)

decreased slowly until the end of the study for all formulations, remaining close to or above

3.5 ng/mL for mdc-STM-001-0.6 and mdc-STM-003-0.6, and around 1 ng/mL for mdc-STM-

002-0.6. There was moderate to high inter-animal variability, ranging from 53% for mdc-

STM-001-0.6 to 92% for mdc-STM-003-0.6. Additionally, the ratio of Cmax to the level

measured after each month (C28d, C56d, C91d) was up to 2-fold lower for mdc-STM-001-0.6

compared to other LAIFs at the dose of 0.6 mg/kg.

For mdc-STM-003, both tested doses (0.6 and 1.5 mg/kg) exhibited a hypo-proportional

increase in Cmax and AUC, approximately 40% less than proportionality. However, these

findings should be considered cautiously due to the high inter-animal variability observed,

particularly at the low dose (Table 3 and Supplementary Table 2 of the Appendix).

Because the second phase of the PK profile was relatively flat, the apparent terminal phase

half-life (T1/2) was accurately estimated for only some animals, with a mean T1/2 ranging

from 20 days for mdc-STM-003-1.5 to 33 days for mdc-STM-001-0.6.

The analysis of each animal's depot collected at the end of the study (DAI 132 or 133)

showed that the remaining amount of ivermectin not released from the depot was the

lowest for mdc-STM-001-0.6 and mdc-STM-003-0.6 with an average of 23.0% of the injected

amount compared to 39% for mdc-STM-002-0.6 at the same dose of 0.6 mg/kg. A similar

trend was observed for the remaining amounts of copolymers which were the lowest mdc-

STM-001-0.6 (approximately 19% of the injected amount) compared to 26% for mdc-STM-

003-0.6 and 34% for mdc-STM-002-0.6. For mdc-STM-003 at the dose of 1.5 mg/kg, the

remaining amount of ivermectin and copolymers were 33% and 34% of the injected amount,

respectively (Supplementary Tables 4 and 5, Appendix).

Mosquitocidal activity evaluation

<u>Knock-down resistance (Kdr) status of the VK5 mosquito lots</u>

444
Generations F2 to F8 of the VK5 colony were used in this experiment. Mosquito batches

exposed at DAI 2, 7, 17, 21, 28, and 42 consisted of 74–80% individuals exhibiting

pyrethroid-resistant phenotypes (i.e., carrying RR or RS genotypes, [40]). The remaining

batches (exposed at DAI 56, 70, 84, 91, 98, 105, 112, 119 and 126) contained 50–54%

resistant mosquitoes. Detailed qPCR results by generation and batch are given in the

Appendix (p10).

451
<u>Blood feeding rate</u>

High blood-feeding rates were consistently observed across all time points for both *An.*

gambiae Kisumu (KIS) and *An. coluzzii* (VK5) colonies (Table 4). No statistically significant

differences in feeding success were observed between treatment groups or formulations at

any time point (data not shown), indicating uniform host attractiveness and feeding

opportunity across experimental conditions.

462 Table 4: Blood feeding rates and the number of fed mosquitoes out of the total exposed to

463 calves, either controls or those receiving the LAIFs.

Mosquito Colony	Cattle treatments					
	Control	mdc-STM-001-0.6	mdc-STM-002-0.6	mdc-STM-003-0.6	mdc-STM-003-1.5	Total
KIS	0.900 (3908/4341)	0.921 (4128/4481)	0.897 (4206/4487)	0.892 (4080/4572)	0.831 (3675/4421)	0.889 (19817/22302)
VK5	0.966 (5047/5224)	0.945 (5005/5295)	0.935 (5091/5447)	0.961 (5160/5371)	0.968 (5174/5343)	0.955 (25477/26680)

Efficacy outcomes

The total number of mosquitoes monitored for survival by colony and treatment group is

presented in Table 5. Their detailed distribution by individual cattle, cup, and day after

injection (DAI) is provided in Supplementary Tables 6A (KIS) and 6B (VK5). Due to limited

availability of KIS mosquitoes, no feeding assays were conducted at DAI 7 and DAI 28.

Mosquito husbandry issues affected the same colony at DAI 105 and DAI 119; these data

points were excluded from the analysis. Unexpected high mortality rates during the first 4

days post blood feeding (> 0.20) were detected for KIS mosquitoes fed on the control cattle

B328 at DAI 42 (mortality rate = 0.437) and B321 at DAI 112 (mortality rate = 0.21).

Therefore, these mosquito lots were excluded from efficacy analyses.

Table 5. Number of KIS and VK5 mosquitoes followed for survival experiments after blood-

feeding on cattle treated with the different LAIFs or on controls. Numbers in brackets

represent the proportion for each treatment relative to the total number of mosquitoes

followed.

Treatment	KIS Ntotal= 10.399	VK5 Ntotal = 15.942
Control	1.926 (19%)	3.161 (20%)
mdc-STM-001-0.6	2.122 (20%)	3.206 (20%)
mdc-STM-002-0.6	2.123 (20%)	3.186 (20%)
mdc-STM-003-0.6	2.135 (21%)	3.207 (20%)
mdc-STM-003-1.5	2.093 (20%)	3.182 (20%)

Mortality rates

A.

Figure 2.

Cumulative mortality rates of KIS and VK5 mosquitoes for the 4 day- (A) and 10 day- (B)

periods post-feeding on cattle treated with three candidates of long-acting formulations at

the dose of 0.6 mg/kg (mdc-STM-001, mdc-STM-002, mdc-STM-003) or mdc-STM-003 at 1.5

mg/kg (i.e., mdc-STM-004 in the legend). Direct skin blood feeding experiments were

performed on different days after injection (DAI) of the formulations. For each plot, the

smooth line represents mean values estimated by the LOESS (locally estimated scatterplot

smoothing) method and grey area shows CI95% [42].

Acute toxic effects of ivermectin were assessed using 4-day cumulative mortality rates for

both *An. gambiae* Kisumu (KIS) and *An. coluzzii* VK5 mosquitoes (Figure 2A). For KIS

mosquitoes, mortality exceeded 50% for up to DAI = 50 with mdc-STM-002 and extended to

approximately 90 days with mdc-STM-001. For VK5 mosquitoes, mortality rates were

generally lower at equivalent DAIs. However, mdc-STM-001-0.6 and mdc-STM-003-1.5

maintained >50% mortality through 63 and 66 days post-injection, respectively. In contrast,

mdc-STM-002-0.6 and mdc-STM-003-0.6 exceeded the 50% mortality threshold only until

around DAI 30.

A similar trend was observed for 10-day cumulative mortality rates (Figure 2B). mdc-STM-

001-0.6 showed the highest efficacy in KIS mosquitoes, comparable to mdc-STM-003-1.5.

From DAI 2 to approximately DAI 100, over 75% of KIS mosquitoes died within 10 days after

feeding cattle treated with mdc-STM-001-0.6, and mortality rates remained above 50%

beyond this period. For VK5 mosquitoes, mdc-STM-001-0.6 maintained ≥50% mortality

between DAI 2 and 80, whereas other formulations at the dose of 0.6 mg/kg were less

effective. The high-dose mdc-STM-003-1.5 achieved the longest duration of efficacy for VK5

mosquitoes, with mortality rates >50% from DAI 2 to 110.

Among all formulations, mdc-STM-001-0.6 exhibited the most consistent efficacy profile.

Mosquito mortality data for this LAIF showed the lowest inter-animal variability, with fewer

deviations from the mean trend across all sampling points.

517   *Survival and Hazard Ratios*

No significant differences in mosquito survival were observed across treatment groups prior

to LAIF administration for either KIS or VK5 colonies (Supplementary Figures 2, 3;

Supplementary Table 7; Figure 3), confirming no confounding effects from host variability.

Following treatment, all formulations administered at the dose of 0.6 mg/kg (mdc-STM-001,

mdc-STM-002, mdc-STM-003) and at 1.5 mg/kg (mdc-STM-003) induced a significant

mosquitocidal effect. The magnitude and duration of these effects varied according to

mosquito strain and formulation but remained statistically significant through the full follow-

up period (DAI 126) in most cases.

Hazard ratios (HRs), derived from Cox proportional hazards models (Figure 3) quantified

treatment effects on mosquito survival relative to the control group. When benchmarked

against the WHO-recommended HR threshold of 4 for endectocidal efficacy against malaria

vectors [38], the different formulations—particularly mdc-STM-001-0.6 and mdc-STM-003-

1.5—met or exceeded this target across extended time intervals.

Figure 3: Heat maps illustrating the lower 95% confidence limit (LCL) of mortality hazard

ratios for KIS (A) and VK5 (B) colony mosquitoes fed on cattle treated with the formulations

mdc-STM-001-0.6, mdc-STM-002-0.6, mdc-STM-003-0.6, mdc-STM-003-1.5 relative to the

controls. Parallel direct skin feeding experiments were conducted for KIS and VK5 on each

cattle. Experiments were performed at different days after injection (DAI) of the

formulations. For each DAI, mosquito survival was monitored over a 30-day period following

blood feeding. Hazard ratios are provided for follow-up periods of 4 and 10 days. HR for 30-

days follow up periods are given in the supplementary Figure 2 of the Appendix. Colored

cells indicate a significant difference in survival between mosquitoes fed on treated versus

control calves (p < 0.05), while white cells represent time points where no significant

difference in survival was observed (p > 0.05). HR and *P-values* for each DAI are available in

supplementary table 7 of the Appendix. BI is the timepoint before injection.

546
547
The mdc-STM-002-0.6 formulation was clearly the least effective, with HRs ≥ 4 for the

shortest duration regardless of colony or follow-up period (4 or 10 days post blood-feeding).

It was therefore excluded from further efficacy comparisons.

For KIS mosquitoes, both mdc-STM-001-0.6 and mdc-STM-003-0.6 sustained 4-day HRs ≥ 4

across nearly the entire study period (up to DAI 126), except at DAI 84 (HRs between 2 and

4). Ten-days HR ≥ 4 was sustained through 70 post-dose for mdc-STM-001-0.6 and through

DAI 21 (and again at DAI 56 and 126) for mdc-STM-003-0.6. For both formulations, 30-day

HRs remained above 4 until DAI 21 (Supplementary Figure 3). The higher-dose formulation

mdc-STM-003-1.5 showed a similar pattern to mdc-STM-001-0.6, with the added benefit of

10-day HRs >4 at an additional time point (DAI 112). For DAI >21, its 30-day HRs remained

between 2 and 4.

For VK5 mosquitoes, both mdc-STM-001-0.6 and mdc-STM-003-0.6 maintained 4-day HR ≥ 4

through 91 days post-injection, except at DAI 84 (HRs was between 2 and 4 for mdc-STM-

001-0.6). Ten-day HRs remained >4 until DAI 70 and 30-day HRs until DAI 7 for both

formulations. The higher-dose LAIF mdc-STM-003-1.5 induced 4-day HRs >4 throughout the

experiment, except of DAI 112, where HR dropped slightly below the threshold (but

remained >2). Ten-day HRs were comparable in duration (up to DAI 70) to the smaller dose

of 0.6 mg/kg, and 30-day HRs exceeded 4 for a period of 21 days.

Median survival times

Median survival times were consistently greater than 10 days for KIS and VK5 mosquitoes

fed on control calves (Figure 4) for all the DAI tested.

For KIS mosquitoes, median survival times remained consistently below 10 days (the

minimum average extrinsic incubation period (EIP) of *Plasmodium falciparum*) throughout

the entire follow-up (up to DAI 126) when fed on cattle treated with mdc-STM-001-0.6, mdc-

STM-003-0.6 and mdc-STM-003-1.5 (Figure 4). In contrast, mdc-STM-002-0.6, induced

median survival times equal to or greater than the 10-day threshold from DAI 70 onward

(Figure 4).

For VK5 mosquitoes, median survival times remained ≤ 10 days up to 91 days post-

treatment, and again at DAI 126, when exposed to cattle treated with mdc-STM-001-0.6,

mdc-STM-003-0.6, and mdc-STM-003-1.5 (Figure 4). The LAIF mdc-STM-002-0.6 maintained

median survival times below or equal to the EIP threshold for only 40 days post-dose.

Notably, mdc-STM-001-0.6 and mdc-STM-003-1.5 induced particularly short survival

durations, with median survival times <4 days for nearly 100 days in KIS mosquitoes and

approximately 60 days in VK5.

Figure 4: Median survival times of mosquitoes from both KIS and VK5 colonies fed at

different days after injection of the LAIFs mdc-STM-001-0.6, mdc-STM-002-0.6, mdc-STM-

003-06 and mdc-STM-003-1.5 (i.e, mdc-STM-004 in the legend). The dashed horizontal line

represents the average extrinsic incubation period (EIP) for *Plasmodium falciparum*. The

dashed vertical lines represent 2- and 3-months post injections. Median values +/- 95% CI

are given.

Lethal concentrations and coverage duration

Concentration/response relationships were examined for each colony and for each LAIF,

using both 4- and 10-day cumulative mortality outcome. No significant differences in LC50 or

LC90 values were found between LAIFs, regardless of the mosquito colony (Supplementary

Figures 4 and 5; Supplementary Tables 8–11 of the Appendix). A unified concentration–

response model, including mosquito colony as explanatory variable was thus developed

using the full dataset (Figure 5). LC50 and LC90 values are summarized in Table 6.

Figure 5. Ivermectin concentrations in relation to cumulative mortalities for KIS and VK5

colony mosquitoes. Cumulative mortalities are considered over either a 4-day (A) or a 10-day

period (B).

Table 6: Ivermectin concentrations inducing 50% and 90% KIS and VK5 mosquito mortalities

over a 4-day or a 10-day period. Mean concentrations are given with lower and upper

confidence intervals (95% CI). Concentration values for KIS and VK5, along with P-values for

comparisons between strains, were obtained using the drc model.

Cumulative mortality window (days)	Mortality %	KIS mosquitoes	VK5 mosquitoes	*P-value* (between colony comparison)
4	50	2.89 [2.27-3.67]	5.26 [4.36-6.35]	1.0e-07
4	90	9.2 [6.72-12.59]	17.65 [12.86-24.22]	0.0e+00
10	50	1.51 [1.1-2.07]	3.66 [2.69-4.97]	1.0e-07
10	90	5.63 [3.98-7.96]	12.28 [8.8-17.14]	1.5e-06

A significant difference in cumulative mortalities between KIS and VK5 was observed in

relation to ivermectin concentrations, with both LC50 and LC90 values consistently lower for

KIS than for VK5, irrespective of 4- or 10-day mortality window. As expected, LC90 values

were consistently higher than LC50, and lethal concentrations associated with 4-day

cumulative mortality were greater than those for the 10-day period. Table 7 summarizes the

time intervals during which each formulation maintained the mean, upper and lower bounds

of LC50 and LC90 values sufficient to kill KIS and VK5 mosquitoes. For consistency with

existing literature and to facilitate direct comparison with previous studies, only LC50 values

were presented in detail in the following sections.

Table 7: Time period during which the LAIF candidates reach Ivermectin Plasma

Concentration values above lethal concentrations inducing 50 and 90 % mortalities, over a 4-

day period or a 10-day period. The time period corresponding to mean LC values are given

and those corresponding to upper and lower confidence intervals (95%), respectively, are

given in brackets. KIS and VK5 are respectively the *An. gambiae* insecticide-susceptible and

the *An. coluzzii* resistant Anopheles colonies**.**

KIS	VK5

FORMULATION	LC (%)	4-Day	10-Day	4-Day	10-Day
MDC-STM-001-0.6	50	132 [132-132]	132 [132-132]	65 [59-69]	132 [67-132]
	90	49 [32-55]	64 [52-73]	20 [8-31]	33 [21-50]
MDC-STM-002-0.6	50	56 [50-63]	73 [65-119]	38 [33-43]	50 [39-58]
	90	27 [21-33]	36 [30-47]	16 [12-20]	21 [16-28]
MDC-STM-003-0.6	50	132 [127-132]	132 [132-132]	75 [67-89]	128 [78-132]
	90	34 [25-65]	71 [53-111]	18 [13-25]	26 [19-49]
MDC-STM-003-1.5	50	132 [132-132]	132 [132-132]	132 [85-132]	132 [132-132]
	90	69 [38-83]	132 [76-132]	26 [20-37]	39 [26-71]

Inter-animal variability was also assessed, as it directly informs the coverage potential of

each formulation, *i.e.* the proportion of treated animals maintaining mosquitocidal

ivermectin plasma concentrations above the LC50 over time. For mdc-STM-001-0.6, 80% of

treated calves sustained ivermectin concentrations above the KIS 10-day LC50 threshold for

at least 4 months post-injection, compared to 60% for mdc-STM-003-0.6. The contrast was

even more pronounced when considering VK5 mosquitoes: with mdc-STM-001-0.6, 60% of

the calves maintained plasma concentrations above the VK5 10-day LC50 for at least 4

months, while the remaining animals sustained this threshold for at least 2 months (64 and

71 days, respectively). In contrast, for mdc-STM-003-0.6, only two calves maintained VK5 10-

day LC50 levels for 4 months, while the remaining three dropped below this threshold after

29, 41 and 50 days post-dose. Taken together, these results indicate that mdc-STM-001-0.6

achieved 100% coverage of VK5 10-day LC50 for a duration of at least 2 months post-

treatment.

DISCUSSION

This study represents the first evaluation of candidate long-acting injectable formulations of

ivermectin specifically developed for human use in malaria control. These formulations are

issued from unique long lasting delivery technology and are designed to meet, with a single

subcutaneous injection, the World Health Organization's minimum target product profile for

endectocides against malaria: achieving, with a single administration, at least one month of

efficacy, defined by hazard ratios (HR) for *Plasmodium* vector survival exceeding 4. The

primary objective of this work was to identify the most promising formulation for further

development, including progression to Phase 1 clinical trial. To guide formulation selection,

we employed a comprehensive approach integrating multiple, complementary criteria

related to mosquitocidal efficacy, pharmacokinetic (PK) characteristics, and inter-individual

variability, all critical for achieving effective and sustained vector control in real-world

conditions and settings.

The lead formulation is mdc-STM-001

All three LAI candidate formulations were well tolerated, and the increased dose of

mdc-STM-003 at 1.5 mg/kg did not produce any observable adverse effects. Based on

pharmacokinetics and efficacy results comparison, mdc-STM-001 emerged as the lead

formulation for further development toward human use. When administered at a single

dose of 0.6 mg/kg, mdc-STM-001 demonstrated several critical features associated with

optimal effectiveness:

i)   An area under the curve (AUC) primarily driven by a longer duration above the

LC50 threshold rather than a higher Cmax. This indicates improved control of the

initial plasma concentration burst and resulted in the lowest observed Cmax

values (mean [range]: 23.0 ng/mL [11.9-32.0]), which is a favorable outcome from

a safety standpoint. Notably, this Cmax is at least eight times lower than the

average safe Cmax (248 ng/mL) reported in healthy volunteers receiving

escalating oral doses of ivermectin tablets up to 120 mg (equivalent to 1404-2000

µg/kg) ([50]);

ii)   Lowest inter-individual variability in ivermectin plasma levels and systemic

exposure, enabling the highest proportion of treated animals to maintain LC50

plasma concentrations over a predefined period. This enhances the reliability of

achieving effective coverage at the population level;

iii)   Mosquitocidal efficacy equal to or exceeding that of other formulations,

sustained for more than two months across all evaluated endpoints. Hazard

ratios (HRs) for both 4- and 10-day follow-up periods remained above 4 for at

least two months post-treatment in both insecticide-susceptible (KIS) and

resistant (VK5) mosquito strains. Median mosquito survival times remained

below 10 days post-blood feeding for a minimum of two months in VK5 and

throughout the entire 126-day observation period for KIS. Additionally, LC50

values were maintained for a mean duration of at least 132 days (95% CI: 67–132)

for VK5, and consistently for (at least) 132 days (95% CI: 132–132) for KIS.

In both magnitude and duration, the mosquitocidal efficacy of a single subcutaneous

injection of mdc-STM-001 formulation is compelling. It would likely reduce Anopheles vector

populations by at least 50% during the first 4 to 10 days after each blood meal, and this

would be sustained over 2 months (VK5) to at least 4 months (KIS) depending on insecticide

resistance status. Transmission potential would be further curtailed because infectious

mosquitoes typically require multiple blood meals across at least 2–3 gonotrophic cycles to

become competent vectors. Moreover, mosquitoes exposed to mdc-STM-001 had median

life spans insufficient to complete the average 10-day Plasmodium falciparum extrinsic

incubation period (EIP)**,** functionally rendering them unable to transmit the parasite.

Thus, mass drug administration (MDA) campaigns incorporating mdc-STM-001, as a

complement to existing malaria control strategies, could substantially and sustainably

reduce Plasmodium transmission, even in regions with high levels of insecticide resistance.

Additional sub-lethal effects of ivermectin—on vector capacity (e.g., fecundity, feeding

behavior) and competence (parasite development)—not assessed in this study, would likely

enhance this impact further and reinforce the formulation's epidemiological impact.

However, several concerns may arise regarding communities' mass treatments using

the lead long-lasting formulation.

Exclusion Criteria and Targeted Population Coverage

In ivermectin-based malaria control strategy, a cornerstone for impact lies in

achieving effective coverage of the human blood reservoir from which mosquitoes feed. This

requires that a sufficient proportion of the targeted population maintains ivermectin plasma

concentrations above mosquitocidal thresholds for a sustained period.

Currently, MDAs of ivermectin for onchocerciasis and lymphatic filariasis exclude

pregnant women, nursing mothers, and children under 15 kg from treatment [47,48].

Applying the same exclusion criteria to malaria-targeted interventions, modeling studies

have estimated that 70–80% population coverage is necessary to achieve meaningful

epidemiological impact with monthly high-dose oral regimens (300 µg/kg/day for 3 days,

maintaining efficacy for ~14 days)[28]. However, two recent Phase 3 randomized controlled

trials using this oral regimen failed to demonstrate reductions in malaria incidence, with

insufficient effective coverage cited as one likely explanation [15,30].

In the Rimdamal II study by Somé et al., PK data revealed that 50% of treated

individuals had plasma ivermectin concentrations below the limit of quantification just two

weeks post-treatment [15]. This indicates that effective drug exposure over the 14-day

period was substantially lower than anticipated, likely contributing to the lack of measurable

impact on malaria incidence. In comparison to oral regimens, long-acting injectable

formulations also require the exclusion of women of childbearing age for safety reasons [53],

further reducing the proportion of the population eligible for treatment. However, this

limitation is counterbalanced by the formulation's prolonged maintenance of mosquitocidal

plasma levels. The extended remanence of ivermectin, combined with low inter-individual

PK variability, increases the cumulative number of person-days with vector-lethal

concentrations. As a result, the effective population coverage during the intervention

window is enhanced, offering greater potential to interrupt transmission compared to oral

formulations [23]. Previous modeling studies have projected that long-lasting systemic

endectocides could significantly reduce malaria incidence across a range of epidemiological

contexts [54]. When administered alongside SMC, a formulation achieving at least two

months of efficacy could result in a 30–90% reduction in malaria burden, depending on the

achieved coverage (from 50% to full, excluding children under 15 kg and women of

childbearing age). Notably, the mdc-STM-001 formulation met and exceeded this two-month

efficacy threshold, depending on the resistance profile of the local mosquito population,

underscoring its high potential as a vector control tool. Ongoing modeling analyses based on

the present data will help assess the projected epidemiological impact of mdc-STM-001

across diverse transmission settings and deployment scenarios, including comparisons to

various oral regimens and incorporating the coverage limitations now documented.

Resistances selection and mitigation

Prolonged systemic exposure to ivermectin may exert selective pressure on both

endoparasites typically targeted by ivermectin treatments, as well as on Anopheles

mosquitoes and other ectoparasites [55,56]. However, since current exclusion criteria

prevent treatment of women of childbearing age, pregnant or lactating women, and children

under 15 kg, large untreated subpopulations would serve as genetic refuges. These refuges

help preserve parasite and vector genetic diversity, substantially mitigating the risk of

resistance selection [57]. As such, the presence of these refuges supports the long-term

sustainability of this long-acting strategy [34,55]. Nevertheless, behavioral resistance

remains a potential concern, particularly through the selection of more zoophagic or

opportunistic mosquito populations that avoid feeding on treated humans. This could

compromise the overall effectiveness of the intervention. Therefore, future implementation

should incorporate regular monitoring of mosquito biting behavior and the early

development of mitigation strategies to address possible shifts in vector ecology [55].

Community acceptability considerations

Mass drug administration campaigns involving an injectable product that provides

community-level protection against malaria, rather than direct individual benefit, may

encounter acceptability challenges. However, previous studies have shown high community

acceptance of altruistic strategies such as injectable transmission-blocking vaccines [58]. For

our approach, acceptability may be further reinforced by the additional health benefits of

ivermectin, which will remain effective against several neglected diseases still prevalent in

malaria-endemic regions. Notably, past MDAs have been associated with reductions in

childhood anemia [15], a previously unknown benefit, as well as decreases in jiggers

infestations and bedbug nuisances [59]. These positive side effect may contribute to

improved community acceptability and adherence.

The single injection regimen provides additional advantages for both acceptability and

compliance when compared to monthly oral regimen requiring repeated dosing. The

formulation volumes to be injected subcutaneously will be small in human (around 8 µL/kg,

max 1.5 mL per human) and should not represent a limiting factor for the approach to be

implemented. Nonetheless, proactive efforts to identify and address potential acceptability

challenges specific to long-acting ivermectin MDA should be initiated early in the

development process alongside further effectiveness evaluation. Community engagement

will be essential; for sustained adoption, administration strategies must be co-developed

with local stakeholders, including affected populations and health authorities [60].

Physiological resistance of mosquitoes and ivermectin mosquitocidal efficacy

Our findings reveal distinct differences in ivermectin susceptibility between the two

mosquito colonies, with VK5 requiring higher plasma lethal concentrations than KIS. This

suggests the presence of ivermectin tolerance mechanisms in the VK5 strain, potentially

linked to genetic factors as species-specific traits or cross-resistance provided by the

knockdown resistance (kdr) mutation, associated with pyrethroid resistance, identified in

this colony. Variability in ivermectin susceptibility have also been documented among

different Anopheles species (e.g. [16]), as well as among different pyrethroid-resistant Aedes

strains, supporting the potential for shared or overlapping resistance mechanisms [61]. The

persistence of metabolic resistance is unlikely, as such mechanisms typically vanish rapidly

under laboratory conditions, especially in the absence of ivermectin exposure.

Tolerance mechanisms, possibly selected through repeated ivermectin MDA

campaigns targeting lymphatic filariasis and onchocerciasis, have been proposed as a

contributing factors to the limited success of some MDA interventions in reducing Anopheles

vector densities [15]. While such underlying mechanisms have been observed in other

species, they remained poorly studied in Anopheles [56]. Further investigations, including

experimental selection studies, are needed to better characterize the potential for

ivermectin tolerance in malaria vectors [62].

Differential susceptibility among malaria vector populations represents a significant

challenge to intervention effectiveness and must be considered when designing appropriate

dosing regimens for future strategies. For example, our data have shown that increasing the

dose of the mdc-STM-003 formulation extended its efficacy against VK5-tolerant

mosquitoes, achieving outcomes comparable to those observed in the more susceptible KIS

strain.

Additional threats to efficacy may arise from incomplete understanding of mosquito

bionomics in target areas, particularly regarding seasonal population dynamics and blood-

feeding behavior [31]. Incomplete coverage of mosquitoes blood sources, due to untreated

human individuals or the presence of domestic animals in the case of zoophagic behaviour,

could sustain vector populations at levels compatible with continued transmission.

Furthermore, subcutaneous ivermectin formulations may alter host odor profiles, potentially

affecting human attractiveness to mosquitoes. As children under 15kg and women of

childbearing age will not receive treatment, the possibility of their increased exposure to

mosquito bites warrants further investigation [55]. Ongoing studies are exploring potential

ivermectin-induced changes in the attractiveness of calves, but similar research addressing

this question in humans is needed.

CONCLUSION

The LAIF mdc-STM-001 emerged as the lead candidate formulation for further development

toward human use against malaria. A single injection can kill >50% of biting mosquitoes

before they become infectious, even in insecticide-resistant populations, for a duration of at

least 2 months. These findings strongly support the need for in-human pharmacokinetic and

pharmacodynamic studies to characterize the formulation's specific behavior in humans and

evaluate its real-world potential for malaria control.

To ensure relevance for vector control, mosquitocidal efficacy should be rigorously

assessed using hazard ratios, LC50 values, and mortality rates, with mosquito follow-up

limited to 10 days post-exposure. In other words, prioritizing acute ivermectin-induced

lethality in efficacy assessments is essential to suppress the buildup of infectious vector

populations. Epidemiological modeling studies must account for the most extreme

scenarios, notably regarding key variables such as human biting rate, human population

coverage and heterogeneous demographic structures. Crucially, understanding local

mosquito bionomics—including the diversity and proportion of blood sources—is essential

for optimizing deployment strategies and assessing whether complementary treatment of

domestic animals is warranted.

With the drug and the BEPO® long-acting delivery technology already available, ivermectin's

safety profile well established, and key challenges to ivermectin-MDA effectiveness both

identified and inherently addressed by the long-lasting formulations approach, we are

confident that mdc-STM-001 could be a game changer for controlling residual malaria when

used alongside existing tools. Crucially, it remains one of the very few approaches capable of

targeting outdoor transmission, a persistent gap in current malaria control strategies.

Evaluating the pharmacokinetics and efficacy of mdc-STM-001 in humans, following a single

injection across varying doses, is now the critical next step in determining its suitability for

broader deployment. Given the strength of these preliminary results—and the additional

public health value of ivermectin in treating co-endemic parasitic infections—it would be a

missed opportunity not to advance mdc-STM-001 into Phase 1 clinical trials. In our view, the

case for further investigation is both timely and compelling.

Competing interests

All authors and contributors currently or formerly employed at Medincell S.A. did not

influence data collection, data analysis, manuscript writing, decision to publish, or

conclusions presented in this paper. The other authors are not shareholders of Medincell

S.A. and declare no competing interests.

Funding

This study was funded by Unitaid under the IMPACT project consortium.
853
854

Authors' contribution

LZ performed the experiments at CIRDES facilities, preliminary statistical analysis and

provided the first original version of the manuscript. SHP, KM, NM, CP, RKD designed the

protocol. SHP coordinated experimental work at CIRDES facilities. AS managed the

molecular biology and coordinated the manuscript drafting. SB and AP performed statistical

analysis. SB produced the final figures. KM was the scientific coordinator for the IMPACT

project. All co-authors contributed to revision of the manuscript, approved the final version

of the manuscript for publication. All authors read and approved the final manuscript.

Data, code, and supplementary information availability

Supplementary information, data, analysis and figures are given in the Appendix.

Data and codes are available online (datasud link, ongoing)

REFERENCES

1. Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of
malaria control on \emphPlasmodium falciparum in Africa between 2000 and 2015. Nature.
2015;526:207–11.
2. Riveron Miranda J, Tchouakui M, Mugenzi L, Menze B, Chiang M-C, Wondji C. Insecticide
Resistance in Malaria Vectors: An Update at a Global Scale. 2018.
3. Carrasco D, Lefèvre T, Moiroux N, Pennetier C, Chandre F, Cohuet A. Behavioural
adaptations of mosquito vectors to insecticide control. Current Opinion in Insect Science.
2019;34:48–54.
4. Martin JL, Mosha FW, Lukole E, Rowland M, Todd J, Charlwood JD, et al. Personal
protection with PBO-pyrethroid synergist-treated nets after 2 years of household use against
pyrethroid-resistant Anopheles in Tanzania. Parasit Vectors. 2021;14:150.
5. Accrombessi M, Cook J, Dangbenon E, Yovogan B, Akpovi H, Sovi A, et al. Efficacy of
pyriproxyfen-pyrethroid long-lasting insecticidal nets (LLINs) and chlorfenapyr-pyrethroid
LLINs compared with pyrethroid-only LLINs for malaria control in Benin: a cluster-
randomised, superiority trial. The Lancet. 2023;401:435–46.
6. Moiroux N, Gomez MB, Pennetier C, Elanga E, Djenontin A, Chandre F, et al. Changes in
Anopheles funestus biting behavior following universal coverage of long-lasting insecticidal
nets in Benin. J Infect Dis. 2012;206:1622–9.
7. Sangbakembi-Ngounou C, Costantini C, Longo-Pendy NM, Ngoagouni C, Akone-Ella O,
Rahola N, et al. Diurnal biting of malaria mosquitoes in the Central African Republic indicates
residual transmission may be "out of control." Proceedings of the National Academy of
Sciences. 2022;119:e2104282119.
8. World malaria report 2024 [Internet]. [cited 2025 Mar 22]. Available from:
https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2024
9. Takken W, Charlwood D, Lindsay SW. The behaviour of adult Anopheles gambiae, sub-
Saharan Africa's principal malaria vector, and its relevance to malaria control: a review.
Malaria Journal. 2024;23:161.
10. Foy BD, Kobylinski KC, da Silva IM, Rasgon JL, Sylla M. Endectocides for malaria control.
Trends Parasitol. 2011;27:423–8.
11. Campbell WC, Benz GW. Ivermectin: a review of efficacy and safety. Journal of Veterinary
Pharmacology and Therapeutics. 1984;7:1–16.
12. Maia M, Kobylinski K. Ivermectin for the Control and Elimination of Malaria. Practical
Control of Mosquitoes Disease Vectors [Internet]. 2025 [cited 2025 Mar 22]. p. 139–65.
Available from: https://www.cabidigitallibrary.org/doi/10.1079/9781789248845.0007
13. Campbell WC, editor. Ivermectin and Abamectin [Internet]. New York, NY: Springer; 1989
[cited 2025 Mar 22]. Available from: http://link.springer.com/10.1007/978-1-4612-3626-9
14. Soma DD, Zogo BM, Some A, Tchiekoi BN, Hien DFD, Pooda HS, et al. Anopheles
bionomics, insecticide resistance and malaria transmission in southwest Burkina Faso : a pre-
intervention study. PLoS One. 2020;15:e0236920 [21 p.].
15. Somé AF, Somé A, Sougué E, Ouédraogo COW, Da O, Dah SR, et al. Safety and efficacy of
repeat ivermectin mass drug administrations for malaria control (RIMDAMAL II): a phase 3,

double-blind, placebo-controlled, cluster-randomised, parallel-group trial. The Lancet Infectious Diseases [Internet]. 2025 [cited 2025 Mar 22];0. Available from: https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(24)00751-5/abstract

16. Kobylinski KC, Satoto TBT, Nurcahyo W, Nugraheni YR, Testamenti VA, Winata IPBA, et al. Impact of standard and long-lasting ivermectin formulations in cattle and buffalo on wild Anopheles survival on Sumba Island, Indonesia. Sci Rep. 2024;14:29770.

17. Pooda SH, Hien DF de S, Pagabeleguem S, Heinrich AP, Porciani A, Sagna AB, et al. Impact of blood meals taken on ivermectin-treated livestock on survival and egg production of the malaria vector Anopheles coluzzii under laboratory conditions. PLoS One. 2024;19:e0308293.

18. Mekuriaw W, Balkew M, Messenger LA, Yewhalaw D, Woyessa A, Massebo F. The effect of ivermectin((R)) on fertility, fecundity and mortality of Anopheles arabiensis fed on treated men in Ethiopia. Malar J. 2019;18:357.

19. Sampaio V de S, Rivas GB da S, Kobylinski K, Pinilla YT, Pimenta PFP, Lima JBP, et al. What does not kill it makes it weaker: effects of sub-lethal concentrations of ivermectin on the locomotor activity of Anopheles aquasalis. Parasites & Vectors. 2017;10:623.

20. Kobylinski KC, Deus KM, Butters MP, Hongyu T, Gray M, da Silva IM, et al. The effect of oral anthelmintics on the survivorship and re-feeding frequency of anthropophilic mosquito disease vectors. Acta Trop. 2010;116:119–26.

21. Kobylinski KC, Foy BD, Richardson JH. Ivermectin inhibits the sporogony of Plasmodium falciparum in Anopheles gambiae. Malar J. 2012;11:381.

22. Pinilla YT, S CPL, V SS, Andrade FS, Melo GC, Orfano AS, et al. Promising approach to reducing Malaria transmission by ivermectin: Sporontocidal effect against Plasmodium vivax in the South American vectors Anopheles aquasalis and Anopheles darlingi. PLoS Negl Trop Dis. 2018;12:e0006221.

23. Chaccour C, Hammann F, Rabinovich NR. Ivermectin to reduce malaria transmission I. Pharmacokinetic and pharmacodynamic considerations regarding efficacy and safety. Malaria Journal. 2017;16:161.

24. Foy BD, Alout H, Seaman JA, Rao S, Magalhaes T, Wade M, et al. Efficacy and risk of harms of repeat ivermectin mass drug administrations for control of malaria (RIMDAMAL): a cluster-randomised trial. Lancet. 2019;393:1517–26.

25. Dabira ED, Soumare HM, Conteh B, Ceesay F, Ndiath MO, Bradley J, et al. Mass drug administration of ivermectin and dihydroartemisinin-piperaquine against malaria in settings with high coverage of standard control interventions: a cluster-randomised controlled trial in The Gambia. Lancet Infect Dis. 2022;22:519–28.

26. Smit MR, Ochomo EO, Aljayyoussi G, Kwambai TK, Abong'o BO, Chen T, et al. Safety and mosquitocidal efficacy of high-dose ivermectin when co-administered with dihydroartemisinin-piperaquine in Kenyan adults with uncomplicated malaria (IVERMAL): a randomised, double-blind, placebo-controlled trial. Lancet Infect Dis. 2018;18:615–26.

27. Kamau Y, Tuwei M, Wanjiku C, Ominde K, Ngama M, Karisa J, et al. Mosquitocidal efficacy and pharmacokinetics of single-dose ivermectin versus three-day dose regimen for malaria vector control compared with albendazole and no treatment: An open-label randomized controlled trial. International Journal of Infectious Diseases [Internet]. 2024 [cited 2025 Mar 22];148. Available from: https://www.ijidonline.com/article/S1201-9712(24)00307-2/fulltext

28. Slater HC, Foy BD, Kobylinski K, Chaccour C, Watson OJ, Hellewell J, et al. Ivermectin as a novel complementary malaria control tool to reduce incidence and prevalence: a modelling

957 study. The Lancet Infectious Diseases. 2020;20:498–508.

958 29. Chaccour C, Casellas A, Hammann F, Ruiz-Castillo P, Nicolas P, Montaña J, et al.
959 BOHEMIA: Broad One Health Endectocide-based Malaria Intervention in Africa—a phase III
960 cluster-randomized, open-label, clinical trial to study the safety and efficacy of ivermectin
961 mass drug administration to reduce malaria transmission in two African settings. Trials.
962 2023;24:128.

963 30. Hutchins H, Pretorius E, Bradley J, Silva ET da, Vasileva H, Ndiath MO, et al. Adjunctive
964 ivermectin mass drug administration for malaria control on the Bijagos Archipelago of
965 Guinea-Bissau (MATAMAL): a quadruple-blinded, cluster-randomised, placebo-controlled
966 trial. The Lancet Infectious Diseases [Internet]. 2024 [cited 2025 Mar 22];0. Available from:
967 https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(24)00580-2/fulltext

968 31. Mouline K, Costantini C. Is ivermectin surviving expectations in residual malaria control?
969 Lancet Infect Dis. 2024;S1473-3099(24)00652-2.

970 32. Roberge C, Cros J-M, Serindoux J, Cagnon M-E, Samuel R, Vrlinic T, et al. BEPO®:
971 Bioresorbable diblock mPEG-PDLLA and triblock PDLLA-PEG-PDLLA based in situ forming
972 depots with flexible drug delivery kinetics modulation. J Control Release. 2020;319:416–27.

973 33. Boussinesq M, Enyong P, Chounna-Ndongmo P, Njouendou A-J, Pion SD, Rech A, et al.
974 Effects of an injectable long-acting formulation of ivermectin on Onchocerca ochengi in zebu
975 cattle. Parasite. 2020;27:36.

976 34. Heinrich AP, Pooda SH, Porciani A, Zéla L, Schinzel A, Moiroux N, et al. An
977 ecotoxicological view on malaria vector control with ivermectin-treated cattle. Nat Sustain.
978 2024;7:724–36.

979 35. Chaccour C, Barrio A, Royo A, Urbistondo D, Slater H, Hammann F, et al. Screening for an
980 ivermectin slow-release formulation suitable for malaria vector control. Malar J.
981 2015;14:102.

982 36. Chaccour CJ, Ngha'bi K, Abizanda G, Irigoyen Barrio A, Aldaz A, Okumu F, et al. Targeting
983 cattle for malaria elimination: marked reduction of Anopheles arabiensis survival for over six
984 months using a slow-release ivermectin implant formulation. Parasit Vectors. 2018;11:287.

985 37. Bellinger AM, Jafari M, Grant TM, Zhang S, Slater HC, Wenger EA, et al. Oral, ultra-long-
986 lasting drug delivery: Application toward malaria elimination goals. Sci Transl Med.
987 2016;8:365ra157.

988 38. Endectocide and ectocide products for malaria transmission control [Internet]. [cited
989 2022 Jul 12]. Available from: https://www.who.int/publications/i/item/9789240052512

990 39. Santolamazza F, Mancini E, Simard F, Qi Y, Tu Z, della Torre A. Insertion polymorphisms
991 of SINE200 retrotransposons within speciation islands of Anopheles gambiae molecular
992 forms. Malar J. 2008;7:163.

993 40. Bass C, Nikou D, Donnelly MJ, Williamson MS, Ranson H, Ball A, et al. Detection of
994 knockdown resistance (kdr) mutations in Anopheles gambiae: a comparison of two new
995 high-throughput assays with existing methods. Malar J. 2007;6:111.

996 41. Toé KH, Jones CM, N'Fale S, Ismail HM, Dabiré RK, Ranson H. Increased Pyrethroid
997 Resistance in Malaria Vectors and Decreased Bed Net Effectiveness, Burkina Faso. Emerg
998 Infect Dis. 2014;20:1691–6.

999 42. R: The R Project for Statistical Computing [Internet]. [cited 2024 Jan 25]. Available from:
1000 https://www.r-project.org/

1001 43. Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D. performance: An R Package
1002 for Assessment, Comparison and Testing of Statistical Models. JOSS. 2021;6:3139.

1003 44. Hartig F. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression

Models. R package version 0.3.3.0. 2020.

45. Holm S. A Simple Sequentially Rejective Multiple Test Procedure. Scandinavian Journal of Statistics. 1979;6:65–70.

46. Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al. glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. The R Journal. 2017;9:378–400.

47. McGillycuddy M, Popovic G, Bolker BM, Warton DI. Parsimoniously Fitting Large Multivariate Random Effects in glmmTMB. Journal of Statistical Software. 2025;112:1–19.

48. Therneau TM. coxme: Mixed Effects Cox Models [Internet]. 2024 [cited 2025 May 21]. Available from: https://cran.r-project.org/web/packages/coxme/index.html

49. Ritz C, Baty F, Streibig JC, Gerhard D. Dose-Response Analysis Using R. Xia Y, editor. PLoS ONE. 2015;10:e0146021.

50. Guzzo CA, Furtek CI, Porras AG, Chen C, Tipping R, Clineschmidt CM, et al. Safety, tolerability, and pharmacokinetics of escalating high doses of ivermectin in healthy adult subjects. J Clin Pharmacol. 2002;42:1122–33.

51. Crompton DWT, World Health Organization. Preventive chemotherapy in human helminthiasis : coordinated use of anthelminthic drugs in control interventions : a manual for health professionals and programme managers. Chimioprévention des helminthiases chez l'homme : utilisation coordonnée des médicaments anthelminthiques pour les interventions de lutte : manuel à l'intention des professionnels de la santé et des administrateurs de programmes. 2006;62.

52. World Health Organization. WHO Community-Directed treatment with Ivermectin (CDTI) [Internet]. [cited 2025 Mar 26]. Available from: https://iris.who.int/bitstream/handle/10665/275546/275546-eng.pdf

53. Nicolas P, Maia MF, Bassat Q, Kobylinski KC, Monteiro W, Rabinovich NR, et al. Safety of oral ivermectin during pregnancy: a systematic review and meta-analysis. The Lancet Global Health. 2020;8:e92–100.

54. Selvaraj P, Suresh J, Wenger EA, Bever CA, Gerardin J. Reducing malaria burden and accelerating elimination with long-lasting systemic insecticides: a modelling study of three potential use cases. Malaria Journal. 2019;18:307.

55. Sagna AB, Zéla L, Ouedraogo COW, Pooda SH, Porciani A, Furnival-Adams J, et al. Ivermectin as a novel malaria control tool: Getting ahead of the resistance curse. Acta Tropica. 2023;245:106973.

56. Furnival-Adams J, Kiuru C, Sagna AB, Mouline K, Maia M, Chaccour C. Ivermectin resistance mechanisms in ectoparasites: a scoping review. Parasitol Res. 2024;123:221.

57. van Wyk JA. Refugia--overlooked as perhaps the most potent factor concerning the development of anthelmintic resistance. Onderstepoort J Vet Res. 2001;68:55–67.

58. White SE, Harvey SA, Meza G, Llanos A, Guzman M, Gamboa D, et al. Acceptability of a herd immunity-focused, transmission-blocking malaria vaccine in malaria-endemic communities in the Peruvian Amazon: an exploratory study. Malaria Journal. 2018;17:179.

59. Onyango T, Nuru K, Kazungu K, Wangari W, Chaccour C, Regina Rabinovich N, et al. Through the eyes of the participant: using photovoice to understand the experiences and effects of ivermectin MDA in the context of the BOHEMIA clinical trial in Kwale, Kenya. Malaria Journal. 2025;24:79.

60. Orok AB, Ajibaye ,Olusola, Aina ,O. Oluwagbemiga, Iboma ,Godswill, Adagyo Oboshi ,Sunday, and Iwalokun B. Malaria interventions and control programes in Sub-Saharan Africa: A narrative review. Shiri R, editor. Cogent Medicine. 2021;8:1940639.

61. Deus KM, Saavedra-Rodriguez K, Butters MP, Black WC th, Foy BD. The effect of
ivermectin in seven strains of Aedes aegypti (Diptera: Culicidae) including a genetically
diverse laboratory strain and three permethrin resistant strains. J Med Entomol.
2012;49:356–63.
62. OUEDRAOGO COW, SOME FA, SAGNA AB, SOUGUE E, SOMA DD, NDIAYE M, et al.
Comparative susceptibility of wild and laboratory-reared Aedes and Anopheles larvae to
Ivermectin® [Internet]. Research Square; 2024 [cited 2025 May 20]. Available from:
https://www.researchsquare.com/article/rs-5408919/v1
1059

Appendix

Selection of a Lead Long-Acting Formulation of Ivermectin to Target Major Malaria Vectors in Western Africa: Evaluation of Pharmacokinetics and Mosquitocidal Efficacy in Cattle under Laboratory Conditions.

Lamidi Zela[1,2], Sié Hermann Pooda[1,3], Angélique Porciani[4], Samuel Beneteau[5], André Barembaye Sagna[4,6*], Sophie Le Lamer-Déchamps[7], Nicolas Moiroux[4,8], Cheick Oumar Wendpagnandé Ouédraogo[9,10], Anyirékun Fabrice Somé[9], Cédric Pennetier[4], Christophe Roberge[7], Adrien M.G. Belem[2], Koumbobr Roch Dabiré[9], Charles Dayo[1], Karine Mouline[4**]

Affiliations :

1. Centre International de Recherche-Développement sur L'Elevage en zone Subhumide, Bobo-Dioulasso, Burkina Faso

2. Université Nazi Boni, Bobo-Dioulasso, Burkina Faso

3. Université Ouezzin COULIBALY, Dédougou

4. MIVEGEC, University of Montpellier, IRD, CNRS, Montpellier, France

5. Independant Statistician

6. IRD, Représentation du Burkina Faso, Antenne de Bobo-Dioulasso, Bobo-Dioulasso, Burkina Faso

7. MedinCell, Montpellier, France

8. Pôle de Zoologie Médicale, Institut Pasteur de Dakar, Dakar, Sénégal

9. Institut de Recherche en Sciences de la Santé, Direction Régionale de l'Ouest, Centre National de Recherche Scientifique et Technologique, Bobo-Dioulasso, Burkina Faso

10. Université Cheikh Anta Diop, Dakar, Sénégal

Table of contents

Methods
Injection volumes per cattle

Supplementary table 1
Injected volumes (mL) per animal and per formulation, given the body weight (BW) measured the day of the treatment (kg).

Animal ID	BW at arrival (kg)	BW at injection (kg)	Formulation ID	Injected volume (mL)
B122	141.6	153.4	mdc-STM-001	1.18
B126	116.8	140.2	mdc-STM-001	1.08
B128	109.4	119.8	mdc-STM-001	0.92
B332	119.0	126.6	mdc-STM-001	0.98
B335	99.2	115.2	mdc-STM-001	0.89
B121	107.2	120.8	mdc-STM-002	0.94
B130	128.8	137.6	mdc-STM-002	1.07
B324	97.4	115.2	mdc-STM-002	0.90
B330	138.0	144.6	mdc-STM-002	1.13
B334	142.0	151.4	mdc-STM-002	1.18
B123	101.2	122.3	mdc-STM-003	0.72
B124	124.4	130.6	mdc-STM-003	0.76
B125	124.6	142.2	mdc-STM-003	0.83
B326	140.8	160.8	mdc-STM-003	0.94
B333	94.2	106.6	mdc-STM-003	0.62
B322	127.4	146.8	mdc-STM-003	2.14
B323	108.6	120.0	mdc-STM-003	1.75
B327	129.4	144.6	mdc-STM-003	2.11
B329	99.8	115.8	mdc-STM-003	1.69
B331	106.4	124.8	mdc-STM-003	1.82

Results
Cattle Weight
Supplementary table 2
Median weight (Q1, Q3) in kg for each experimental cattle group at each weighing date.

Experimental arm					
	Control N=5[1]	mdc-STM-001 N=5[1]	mdc-STM-002 N=5[1]	mdc-STM-003 N=5[1]	mdc-STM-004 N=5[1]
Weighing date					
2020-11-25	114 (110,125)	117 (109,119)	129 (107,138)	124 (101,125)	109 (106,127)
2020-12-24	116 (116,126)	118 (114, 123)	134 (105,136)	121 (109,126)	113 (112,133)
2021-01-14	129 (123,140)	127 (120,140)	138 (121,145)	131 (122, 142)	125 (120, 145)
2021-02-15	138 (138,151)	135 (134,153)	157 (134,162)	146 (138,159)	139 (134,157)
2021-03-15	144 (141,152)	141 (141,158)	164 (134,169)	150 (138,162)	142 (140,163)
2021-04-15	162 (155,162)	160 (152,170)	177 (149,178)	166 (148,181)	163 (149,178)
2021-05-15	161 (155,167)	164 (161,177)	183 (157,186)	167 (158,182)	170 (149,173)

Pharmacokinetics
Supplementary Figure 1
A. Mean concentration-time profiles of ivermectin in plasma cattle after single subcutaneous administration of LAIFs at 0.6 mg/kg (Red: mdc-STM-001; purple: mdc-STM-002; yellow: mdc-STM-003).
B. Mean concentration-time profiles of ivermectin in plasma cattle after single subcutaneous administration mdc-STM-003 formulation at 0.6 and 1.5 mg/kg

Supplementary table 3

Mean (SD) pharmacokinetic parameters of ivermectin in plasma cattle after single subcutaneous administration of LAIFs at 0.6 or 1.5 mg/kg

Parameters	mdc-STM-001 0.6 mg/kg	mdc-STM-002 0.6 mg/kg	mdc-STM-003 0.6 mg/kg	mdc-STM-003 1.5 mg/kg
Tmax* (h)	24 (24; 48)	48 (24; 168)	24 (24; 48)	24 (24; 168)
Cmax (ng/mL)	35.4 (12.7)	50.3 (13.7)	58.1 (25.2)	81.2 (32.2)
Cmax/Dose (ng/mL/mg/kg)	58.3 (22.0)	84.3 (22.6)	99.4 (42.2)	56.1 (22.4)
Clast (ng/mL)	3.93 (2.75)	1.03 (0.70)	3.43 (2.74)	6.30 (4.35)
AUC(0-28d) (h.ng/mL)	14227 (3853)	16253 (5465)	18085 (9789)	26248 (5534)
AUC(0-28d)/Dose (h.ng/mL/mg/kg)	23378 (6862)	27298 (9171)	30956 (16689)	18116 (3822)
AUC(0-91d) (h.ng/mL)	25055 (2936)	21121 (8866)	28890 (18668)	41820 (13286)
AUC(0-91d)/Dose (h.ng/mL/mg/kg)	40990 (5664)	35479 (14839)	49483 (31994)	28903 (9300)
AUClast (h.ng/mL)	28534 (2316)	22322 (9059)	32851 (21691)	47192 (15724)
AUClast/Dose (h.ng/mL/mg/kg)	46589 (3960)	37497 (15165)	56269 (37147)	32630 (11042)
T1/2 (h)	784 (244) (n=4)	731 (579) (n=4)	728 (399) (n=3)	489 (n=1)
C28d (ng/mL)	13.7 (1.6)	8.90 (4.85)	11.1 (8.3)	15.5 (5.1)
C56d (ng/mL)	6.70 (1.67)	2.92 (2.63)	7.34 (7.14)	10.3 (6.9)
C91d (ng/mL)	3.48 (1.85)	1.11 (0.84)	4.33 (3.99)	5.55 (3.72)

SD: standard deviation.

*Median (min; max). Tmax: time to reach Cmax. Cmax: maximal plasma concentration. Clast : last quantifiable plasma concentration. AUC(0-28d): area under curve from time 0 to 28 days. AUC(0-28d)/Dose: dose-normalized AUC(0-28d). AUC(0-91d): area under curve from time 0 to 91 days. AUC(0-91d)/Dose: dose-normalized AUC(0-91d). AUClast: area under curve from time 0 to last quantifiable plasma level. AUClast/Dose: dose-normalized AUClast. T1/2: apparent terminal phase half-life. C28d: plasma concentration 28 days post injection. C56d: plasma concentration 56 days post injection. C91d: plasma concentration 91 days post injection.

Copolymers extraction at study end
Supplementary table 4
Ivermectin extraction results in each depot collected at study end

Treatment Dose (mg/kg)	Animal ID	IVM injected dose (mg)	IVM recovered dose (mg)		IVM recovered dose (%)	
			Indiv.	Mean (SD)	Indiv.	Mean (SD)
F31-0.6	1	100.4	38.6	19.4 (12.9)	38.5	23.0 (12.7)
	2	85.0	18.3		21.6	
	3	73.3	2.5		3.3	
	4	76.3	20.6		27.0	
	5	67.9	16.8		24.8	
F35-0.6	1	76.1	39.0	32.3 (17.7)	51.3	39.1 (19.6)
	2	84.7	6.1		7.2	
	3	69.0	23.4		33.9	
	4	87.7	41.7		47.6	
	5	92.0	51.1		55.5	
F73-0.6	1	70.6	5.4	17.8 (14.4)	7.7	23.1 (17.6)
	2	76.4	14.3		18.7	
	3	81.1	42.6		52.6	
	4	96.4	12.0		12.5	
	5	61.6	14.8		24.0	
F73-1.5	1	211.6	69.8	59.5 (35.5)	33.0	32.7 (20.0)
	2	173.8	55.5		32.0	
	3	212.3	0.5		0.2	
	4	164.8	79.7		48.4	
	5	182.9	91.8		50.2	

IVM : ivermectin. Indiv. : individual. SD: standard deviation.

Supplementary table 5
Copolymers extraction results in each depot collected at study end

Treatment Dose (mg/kg)	Animal ID	Copolymer injected (mg)	Copolymer recovered (mg)	Copolymer recovered (%)	
				Indiv.	Mean (SD)
F31-0.6	1	442.2	120.1	27.2	18.8 (8.5)
	2	374.4	90.0	24.0	
	3	323.1	17.3	5.4	
	4	336.0	55.5	16.5	
	5	299.1	63.3	21.1	
F35-0.6	1	217.6	105.8	48.6	34.1 (15.9)
	2	242.2	19.5	8.0	
	3	197.4	77.3	39.2	
	4	250.8	108.8	43.4	
	5	263.2	81.6	31.0	
F73-0.6	1	155.6	25.0	16.1	25.9 (25.0)
	2	168.4	11.4	6.8	
	3	178.6	39.5	22.1	
	4	212.4	32.0	15.1	
	5	135.6	94.2	69.5	
F73-1.5	1	466.2	196.4	42.1	34.1 (18.8)
	2	382.8	164.3	42.9	
	3	467.6	3.5	0.8	
	4	363.0	167.4	46.1	
	5	403.0	155.6	38.6	

Indiv. : individual. SD: standard deviation.

Mosquitocidal activity evaluation
VK5 resistance status
Sample sizes
Supplementary Table 6A and 6B

Number of mosquitoes followed for their survival before injection (BI) and after blood feeding on the calves at different days either for the control arm or after injection (DAI) of the LAI formulations for the 4 treatment arms. A. KIS mosquitoes B. VK5 mosquitoes

A.

Mosquito Colony	Treatment	Cattle ID	BI	Day after injection (DAI)														
				2	7	17	21	28	42	56	70	84	91	98	105	112	119	126
kis	Control	B127	40	31	0	37	39	0	19	29	41	40	0	40	37	38	39	39
		B129	40	34	0	31	41	0	38	31	41	41	0	36	32	38	40	40
		B321	40	34	0	38	34	0	39	38	41	40	0	0	0	39	43	41
		B328	41	20	0	35	37	0	32	38	42	40	0	42	42	41	41	38
		B336	38	34	0	37	42	0	19	32	41	40	0	39	39	41	39	40
	mdc-STM-001-0.6	B122	40	38	0	39	40	20	40	37	40	39	0	40	37	40	41	42
		B126	39	40	0	40	40	21	40	38	38	36	0	40	40	42	38	36
		B128	38	28	0	39	41	9	40	41	41	40	0	0	0	40	37	41
		B332	40	41	0	34	39	21	38	42	40	44	0	38	40	43	39	38
		B335	40	38	0	40	40	24	32	39	39	43	0	39	41	41	40	41
	mdc-STM-002-0.6	B121	38	38	0	40	43	10	37	44	38	40	0	39	36	41	40	40
		B130	40	39	0	40	40	21	40	42	32	40	0	38	15	39	38	39
		B324	40	39	0	40	40	21	40	39	40	40	0	37	40	36	39	37
		B330	40	41	0	41	40	24	31	40	39	40	0	40	38	39	39	39
		B334	40	38	0	40	40	18	40	38	39	43	0	0	0	42	40	38
	mdc-STM-003-0.6	B123	40	40	0	41	38	10	40	40	40	39	0	38	37	40	39	38
		B124	38	39	0	37	39	6	40	40	40	42	0	39	42	38	37	40
		B125	40	40	0	38	40	25	39	38	40	41	0	0	0	38	39	41
		B326	40	37	0	39	40	16	40	38	40	42	0	39	37	39	40	40
		B333	40	42	0	39	39	4	41	36	43	38	0	40	33	40	40	42
	mdc-STM-003-1.5	B322	40	39	0	39	40	20	40	39	40	42	0	41	37	38	42	40
		B323	40	40	0	39	40	18	39	32	40	40	0	25	37	40	42	44
		B327	40	35	0	38	40	17	41	24	38	41	0	43	26	36	39	41
		B329	38	38	0	40	40	0	35	40	41	42	0	41	36	40	41	39
		B331	40	38	0	38	40	6	30	39	39	40	0	0	0	42	38	39
Total			990	921	0	959	992	311	910	934	993	1013	0	774	722	991	990	993

B.

Mosquito Colony	Treatment	Cattle ID	BI	Days after injection (DAI)														
				2	7	17	21	28	42	56	70	84	91	98	105	112	119	126
vk5	Control	B127	40	41	31	39	41	38	39	40	38	40	42	40	41	39	40	41
		B129	40	39	34	40	40	39	40	39	41	41	41	40	41	38	39	39
		B321	40	38	36	42	34	36	40	41	37	40	39	41	40	39	39	38
		B328	40	40	38	38	39	37	40	42	40	42	39	42	41	39	40	41
		B336	41	37	39	36	37	41	40	42	37	42	42	43	42	40	41	42
	F31-0.6	B122	40	41	39	40	40	41	40	40	40	41	40	39	43	41	42	41
		B126	40	41	40	39	41	40	39	39	39	42	40	40	43	40	40	40
		B128	39	39	40	39	40	38	41	41	39	41	40	40	40	39	42	41
		B332	40	40	37	40	40	38	41	41	40	40	38	39	41	41	42	39
		B335	41	41	40	39	41	41	40	38	38	40	40	42	40	38	40	40
	F35-0.6	B121	38	41	40	41	40	40	38	37	38	38	37	38	40	39	42	42
		B130	40	38	37	40	40	41	38	39	40	46	39	40	45	39	39	39
		B324	40	40	40	40	40	42	39	41	39	42	40	43	42	39	39	40
		B330	40	39	39	40	39	40	37	40	36	40	40	38	43	40	43	37
		B334	40	39	41	41	40	38	40	40	40	39	38	39	43	39	43	40
	F73-0.6	B123	40	40	40	40	39	39	39	40	40	40	40	41	41	42	40	43
		B124	39	40	39	40	39	43	36	41	39	42	40	41	43	40	43	39
		B125	40	39	38	42	42	41	39	40	40	40	41	39	43	41	40	41
		B326	40	40	39	40	41	34	40	39	40	40	40	43	43	39	40	39
		B333	40	40	41	40	38	40	38	38	40	39	40	40	42	41	40	39
	F73-1.5	B322	40	39	40	40	40	41	39	39	40	38	38	40	40	40	39	39
		B323	39	39	39	38	39	37	41	40	40	40	41	43	40	43	39	41
		B327	40	40	40	40	40	39	38	39	37	43	40	39	40	40	42	40
		B329	40	41	40	39	39	37	37	40	41	40	40	39	40	41	43	41
		B331	40	40	39	40	39	41	39	41	40	39	40	38	41	39	41	39
Total			997	992	966	993	988	982	978	997	979	1015	995	1007	1038	996	1018	1001

Kaplan-Meier curves

Supplementary Figure 2

Kaplan-Meier plots illustrating the mosquitocidal effects of the candidate long-lasting ivermectin formulations administered to calves at doses of 0.6 mg/kg (mdc-STM-001, mdc-STM-002 and mdc-STM-003) or 1.5 mg/kg (mdc-STM-003 only). The plots depict the results from each direct skin feeding assay conducted at the indicated time points (Days After Injection, DAI). Mosquito survival was monitored daily for up to 30 days post-blood meal. Pannel A: KIS colony, Pannel B: VK5 colony.

Supplementary Figure 3

Heat maps illustrating the lower 95% confidence limit (LCL) of mortality hazard ratios for KIS (A) and VK5 (B) colony mosquitoes fed on cattle treated with the formulations mdc-STM-001-0.6, mdc-STM-002-0.6, mdc-STM-003-0.6, mdc-STM-003-1.5 (here mdc-ST004) relative to the controls. Parallel direct skin feeding experiments were conducted for KIS and VK5 on each cattle. Experiments were performed at different days after injection (DAI) of the formulations. For each DAI, mosquito survival was monitored over a 30-day period following blood feeding. Hazard ratios are provided for follow-up periods of 30 days. Colored cells indicate a significant difference in survival between mosquitoes fed on treated versus control calves (p < 0.05), while white cells represent time points where no significant difference in survival was observed (p > 0.05). HR and *P-values* for each DAI are available in the supplementary table 6. BI is the timepoint before injection.

HR values and corresponding p-values
Supplementary table 7

Hazard ratios (HR) and their 95% CI with the corresponding p-values for mosquitoes fed on cattle treated with LAIFs of ivermectin and on controls. HR were computed for 4, 10 and 30-days of survival follow-up after mosquitoes blood feeding, before injection (BI) and for each day after injection (DAI). Hazard ratios and p-values are given for mosquitoes fed on the calves treated with LAIFs, relatively to the controls. A. KIS mosquitoes. B. VK5 mosquitoes.

A.

Formulation	Treatment arm	DAI	Follow-up (days)	HR (CI 95%)	p.value
(MDC-STM-001-0.6)	Control	BI	4	1 (0.1-17.2)	1.0000000
(MDC-STM-002-0.6)	Control	BI	4	1.9 (0.2-22.7)	0.9587058
(MDC-STM-003-0.6)	Control	BI	4	0.8 (0-14.3)	0.9997884

Formulation	Treatment arm	DAI	Follow-up (days)	HR (CI 95%)	p.value
(MDC-STM-003-1.5)	Control	BI	4	1.9 (0.2-22.4)	0.9618095
(MDC-STM-001-0.6)	Control	2	4	32.2 (9.8-106.3)	0.0000000
(MDC-STM-002-0.6)	Control	2	4	59.3 (18-195.1)	0.0000000
(MDC-STM-003-0.6)	Control	2	4	58.3 (17.7-191.8)	0.0000000
(MDC-STM-003-1.5)	Control	2	4	57.3 (17.4-188.9)	0.0000000
(MDC-STM-001-0.6)	Control	17	4	56 (17-184.3)	0.0000000
(MDC-STM-002-0.6)	Control	17	4	53.4 (16.2-175.7)	0.0000000
(MDC-STM-003-0.6)	Control	17	4	50.4 (15.3-165.8)	0.0000000
(MDC-STM-003-1.5)	Control	17	4	60.6 (18.4-199.5)	0.0000000
(MDC-STM-001-0.6)	Control	21	4	25.5 (9.3-70.3)	0.0000000
(MDC-STM-002-0.6)	Control	21	4	23.4 (8.5-64.4)	0.0000000
(MDC-STM-003-0.6)	Control	21	4	27.7 (10.1-76.4)	0.0000000
(MDC-STM-003-1.5)	Control	21	4	27.7 (10-76.2)	0.0000000
(MDC-STM-001-0.6)	Control	42	4	41.1 (10.2-165.9)	0.0000000
(MDC-STM-002-0.6)	Control	42	4	24 (5.9-97.5)	0.0000000
(MDC-STM-003-0.6)	Control	42	4	28.2 (7-114.1)	0.0000000
(MDC-STM-003-1.5)	Control	42	4	40.3 (10-162.9)	0.0000000
(MDC-STM-001-0.6)	Control	56	4	55.9 (14-223.3)	0.0000000
(MDC-STM-002-0.6)	Control	56	4	23.6 (5.8-95)	0.0000000
(MDC-STM-003-0.6)	Control	56	4	52.2 (13-209.7)	0.0000000
(MDC-STM-003-1.5)	Control	56	4	81.4 (20.3-326.2)	0.0000000
(MDC-STM-001-0.6)	Control	70	4	26.4 (8.2-84.9)	0.0000000
(MDC-STM-002-0.6)	Control	70	4	6.2 (1.9-20.7)	0.0003562
(MDC-STM-003-0.6)	Control	70	4	19.8 (6.1-64)	0.0000000
(MDC-STM-003-1.5)	Control	70	4	34.4 (10.7-110.5)	0.0000000
(MDC-STM-001-0.6)	Control	84	4	7.9 (2.9-21.6)	0.0000002
(MDC-STM-002-0.6)	Control	84	4	2.2 (0.8-6.2)	0.2699699

Formulation	Treatment arm	DAI	Follow-up (days)	HR (CI 95%)	p.value
(MDC-STM-003-0.6)	Control	84	4	6.8 (2.5-18.8)	0.0000023
(MDC-STM-003-1.5)	Control	84	4	12.8 (4.7-34.9)	0.0000000
(MDC-STM-001-0.6)	Control	98	4	38.4 (8.7-169.4)	0.0000000
(MDC-STM-002-0.6)	Control	98	4	2.7 (0.5-13.6)	0.4587964
(MDC-STM-003-0.6)	Control	98	4	21.2 (4.7-94.6)	0.0000003
(MDC-STM-003-1.5)	Control	98	4	36.7 (8.3-162.4)	0.0000000
(MDC-STM-001-0.6)	Control	112	4	25.7 (6.4-103.6)	0.0000000
(MDC-STM-002-0.6)	Control	112	4	5.6 (1.3-23.6)	0.0092896
(MDC-STM-003-0.6)	Control	112	4	23.5 (5.8-95.2)	0.0000000
(MDC-STM-003-1.5)	Control	112	4	26.8 (6.6-108)	0.0000000
(MDC-STM-001-0.6)	Control	126	4	160.9 (9.3-2793.8)	0.0000118
(MDC-STM-002-0.6)	Control	126	4	29.5 (1.6-528.9)	0.0120573
(MDC-STM-003-0.6)	Control	126	4	156.2 (9-2712.6)	0.0000137
(MDC-STM-003-1.5)	Control	126	4	215.2 (12.4-3727.6)	0.0000028
(MDC-STM-001-0.6)	Control	BI	10	0.6 (0.2-2.3)	0.8747113
(MDC-STM-002-0.6)	Control	BI	10	0.5 (0.1-1.8)	0.5163317
(MDC-STM-003-0.6)	Control	BI	10	0.5 (0.2-1.9)	0.6559403
(MDC-STM-003-1.5)	Control	BI	10	0.4 (0.1-1.5)	0.2850212
(MDC-STM-001-0.6)	Control	2	10	22.3 (8.1-61.5)	0.0000000
(MDC-STM-002-0.6)	Control	2	10	44.4 (16.1-122.5)	0.0000000
(MDC-STM-003-0.6)	Control	2	10	44.1 (16-121.6)	0.0000000
(MDC-STM-003-1.5)	Control	2	10	40.8 (14.8-112.8)	0.0000000
(MDC-STM-001-0.6)	Control	17	10	22.1 (8.7-56.1)	0.0000000
(MDC-STM-002-0.6)	Control	17	10	19.9 (7.9-50.5)	0.0000000
(MDC-STM-003-0.6)	Control	17	10	19.5 (7.7-49.5)	0.0000000
(MDC-STM-003-1.5)	Control	17	10	22.9 (9-57.9)	0.0000000
(MDC-STM-001-0.6)	Control	21	10	19.4 (7.7-48.6)	0.0000000

Formulation	Treatment arm	DAI	Follow-up (days)	HR (CI 95%)	p.value
(MDC-STM-002-0.6)	Control	21	10	16.7 (6.7-42)	0.0000000
(MDC-STM-003-0.6)	Control	21	10	20.7 (8.3-52)	0.0000000
(MDC-STM-003-1.5)	Control	21	10	21.3 (8.5-53.5)	0.0000000
(MDC-STM-001-0.6)	Control	42	10	12.1 (4.6-31.9)	0.0000000
(MDC-STM-002-0.6)	Control	42	10	7 (2.6-18.4)	0.0000005
(MDC-STM-003-0.6)	Control	42	10	8.9 (3.4-23.5)	0.0000000
(MDC-STM-003-1.5)	Control	42	10	12.3 (4.7-32.5)	0.0000000
(MDC-STM-001-0.6)	Control	56	10	14.4 (5.6-37.2)	0.0000000
(MDC-STM-002-0.6)	Control	56	10	7.2 (2.8-18.7)	0.0000002
(MDC-STM-003-0.6)	Control	56	10	14.5 (5.6-37.7)	0.0000000
(MDC-STM-003-1.5)	Control	56	10	21.2 (8.2-54.9)	0.0000000
(MDC-STM-001-0.6)	Control	70	10	12.3 (4.8-31.6)	0.0000000
(MDC-STM-002-0.6)	Control	70	10	3.2 (1.2-8.5)	0.0087904
(MDC-STM-003-0.6)	Control	70	10	9 (3.5-23.2)	0.0000000
(MDC-STM-003-1.5)	Control	70	10	15.7 (6.1-40.3)	0.0000000
(MDC-STM-001-0.6)	Control	84	10	8 (3.2-20.3)	0.0000000
(MDC-STM-002-0.6)	Control	84	10	2.4 (1-6.3)	0.0734027
(MDC-STM-003-0.6)	Control	84	10	5.7 (2.2-14.4)	0.0000038
(MDC-STM-003-1.5)	Control	84	10	10.1 (4-25.4)	0.0000000
(MDC-STM-001-0.6)	Control	98	10	10.1 (3.9-26.2)	0.0000000
(MDC-STM-002-0.6)	Control	98	10	2.1 (0.8-5.5)	0.2703929
(MDC-STM-003-0.6)	Control	98	10	5.7 (2.2-15.1)	0.0000082
(MDC-STM-003-1.5)	Control	98	10	9.6 (3.7-25)	0.0000000
(MDC-STM-001-0.6)	Control	112	10	10.9 (4-29.9)	0.0000000
(MDC-STM-002-0.6)	Control	112	10	3.3 (1.2-9.3)	0.0131335
(MDC-STM-003-0.6)	Control	112	10	10.7 (3.9-29.5)	0.0000000
(MDC-STM-003-1.5)	Control	112	10	11.5 (4.2-31.6)	0.0000000

Formulation	Treatment arm	DAI	Follow-up (days)	HR (CI 95%)	p.value
(MDC-STM-001-0.6)	Control	126	10	34.9 (9.3-131.3)	0.0000000
(MDC-STM-002-0.6)	Control	126	10	10.5 (2.7-40.3)	0.0000181
(MDC-STM-003-0.6)	Control	126	10	33.2 (8.8-125.1)	0.0000000
(MDC-STM-003-1.5)	Control	126	10	45.5 (12.1-170.7)	0.0000000
(MDC-STM-001-0.6)	Control	BI	30	1.1 (0.5-2.3)	0.9974111
(MDC-STM-002-0.6)	Control	BI	30	1 (0.5-2.1)	1.0000000
(MDC-STM-003-0.6)	Control	BI	30	0.8 (0.4-1.8)	0.9663867
(MDC-STM-003-1.5)	Control	BI	30	0.9 (0.4-1.9)	0.9941915
(MDC-STM-001-0.6)	Control	2	30	7.4 (3.5-15.7)	0.0000000
(MDC-STM-002-0.6)	Control	2	30	18.3 (8.6-38.9)	0.0000000
(MDC-STM-003-0.6)	Control	2	30	18.4 (8.7-39.1)	0.0000000
(MDC-STM-003-1.5)	Control	2	30	12.6 (5.9-26.8)	0.0000000
(MDC-STM-001-0.6)	Control	17	30	10.6 (5.1-22.3)	0.0000000
(MDC-STM-002-0.6)	Control	17	30	9.4 (4.5-19.7)	0.0000000
(MDC-STM-003-0.6)	Control	17	30	9.4 (4.5-19.6)	0.0000000
(MDC-STM-003-1.5)	Control	17	30	11.8 (5.6-24.7)	0.0000000
(MDC-STM-001-0.6)	Control	21	30	10.1 (4.8-21.2)	0.0000000
(MDC-STM-002-0.6)	Control	21	30	7.4 (3.5-15.4)	0.0000000
(MDC-STM-003-0.6)	Control	21	30	10.7 (5.1-22.4)	0.0000000
(MDC-STM-003-1.5)	Control	21	30	11.6 (5.6-24.3)	0.0000000
(MDC-STM-001-0.6)	Control	42	30	5.9 (2.7-12.7)	0.0000000
(MDC-STM-002-0.6)	Control	42	30	3.8 (1.8-8.2)	0.0000160
(MDC-STM-003-0.6)	Control	42	30	4.7 (2.2-10.2)	0.0000003
(MDC-STM-003-1.5)	Control	42	30	6.8 (3.2-14.6)	0.0000000
(MDC-STM-001-0.6)	Control	56	30	5.5 (2.6-11.7)	0.0000000
(MDC-STM-002-0.6)	Control	56	30	3.1 (1.5-6.6)	0.0002827
(MDC-STM-003-0.6)	Control	56	30	6.8 (3.2-14.3)	0.0000000

Formulation	Treatment arm	DAI	Follow-up (days)	HR (CI 95%)	p.value
(MDC-STM-003-1.5)	Control	56	30	7.1 (3.4-15.1)	0.0000000
(MDC-STM-001-0.6)	Control	70	30	3.5 (1.7-7.2)	0.0000345
(MDC-STM-002-0.6)	Control	70	30	1.4 (0.7-2.9)	0.7146990
(MDC-STM-003-0.6)	Control	70	30	2.7 (1.3-5.7)	0.0018361
(MDC-STM-003-1.5)	Control	70	30	4.8 (2.3-9.9)	0.0000001
(MDC-STM-001-0.6)	Control	84	30	3.6 (1.7-7.6)	0.0000153
(MDC-STM-002-0.6)	Control	84	30	1.5 (0.7-3.2)	0.5079043
(MDC-STM-003-0.6)	Control	84	30	2.5 (1.2-5.1)	0.0070421
(MDC-STM-003-1.5)	Control	84	30	4.2 (2-8.7)	0.0000010
(MDC-STM-001-0.6)	Control	98	30	3.8 (1.8-8.1)	0.0000099
(MDC-STM-002-0.6)	Control	98	30	1.4 (0.6-2.9)	0.7948357
(MDC-STM-003-0.6)	Control	98	30	2.8 (1.3-6)	0.0015562
(MDC-STM-003-1.5)	Control	98	30	4.4 (2.1-9.3)	0.0000008
(MDC-STM-001-0.6)	Control	112	30	3.7 (1.8-7.8)	0.0000166
(MDC-STM-002-0.6)	Control	112	30	1.5 (0.7-3.2)	0.5637848
(MDC-STM-003-0.6)	Control	112	30	3.7 (1.8-7.9)	0.0000133
(MDC-STM-003-1.5)	Control	112	30	4.4 (2.1-9.3)	0.0000006
(MDC-STM-001-0.6)	Control	126	30	3.9 (1.9-8.3)	0.0000053
(MDC-STM-002-0.6)	Control	126	30	1.7 (0.8-3.6)	0.3188277
(MDC-STM-003-0.6)	Control	126	30	4.1 (2-8.7)	0.0000023
(MDC-STM-003-1.5)	Control	126	30	4.6 (2.2-9.7)	0.0000002

B.

Formulation	Reference	DAI	Follow-up (days)	HR (CI 95%)	p.value
(MDC-STM-001-0.6)	Control	BI	4	1 (0.1-17.2)	1.0000000
(MDC-STM-002-0.6)	Control	BI	4	1.9 (0.2-22.7)	0.9587058
(MDC-STM-	Control	BI	4	0.8 (0-14.3)	0.9997884

Formulation	Reference	DAI	Follow-up (days)	HR (CI 95%)	p.value
003-0.6)					
(MDC-STM-003-1.5)	Control	BI	4	1.9 (0.2-22.4)	0.9618095
(MDC-STM-001-0.6)	Control	2	4	32.2 (9.8-106.3)	0.0000000
(MDC-STM-002-0.6)	Control	2	4	59.3 (18-195.1)	0.0000000
(MDC-STM-003-0.6)	Control	2	4	58.3 (17.7-191.8)	0.0000000
(MDC-STM-003-1.5)	Control	2	4	57.3 (17.4-188.9)	0.0000000
(MDC-STM-001-0.6)	Control	17	4	56 (17-184.3)	0.0000000
(MDC-STM-002-0.6)	Control	17	4	53.4 (16.2-175.7)	0.0000000
(MDC-STM-003-0.6)	Control	17	4	50.4 (15.3-165.8)	0.0000000
(MDC-STM-003-1.5)	Control	17	4	60.6 (18.4-199.5)	0.0000000
(MDC-STM-001-0.6)	Control	21	4	25.5 (9.3-70.3)	0.0000000
(MDC-STM-002-0.6)	Control	21	4	23.4 (8.5-64.4)	0.0000000
(MDC-STM-003-0.6)	Control	21	4	27.7 (10.1-76.4)	0.0000000
(MDC-STM-003-1.5)	Control	21	4	27.7 (10-76.2)	0.0000000
(MDC-STM-001-0.6)	Control	42	4	41.1 (10.2-165.9)	0.0000000
(MDC-STM-002-0.6)	Control	42	4	24 (5.9-97.5)	0.0000000
(MDC-STM-003-0.6)	Control	42	4	28.2 (7-114.1)	0.0000000
(MDC-STM-003-1.5)	Control	42	4	40.3 (10-162.9)	0.0000000
(MDC-STM-001-0.6)	Control	56	4	55.9 (14-223.3)	0.0000000
(MDC-STM-002-0.6)	Control	56	4	23.6 (5.8-95)	0.0000000
(MDC-STM-003-0.6)	Control	56	4	52.2 (13-209.7)	0.0000000
(MDC-STM-003-1.5)	Control	56	4	81.4 (20.3-326.2)	0.0000000
(MDC-STM-001-0.6)	Control	70	4	26.4 (8.2-84.9)	0.0000000
(MDC-STM-002-0.6)	Control	70	4	6.2 (1.9-20.7)	0.0003562
(MDC-STM-003-0.6)	Control	70	4	19.8 (6.1-64)	0.0000000
(MDC-STM-003-1.5)	Control	70	4	34.4 (10.7-110.5)	0.0000000
(MDC-STM-001-0.6)	Control	84	4	7.9 (2.9-21.6)	0.0000002
(MDC-STM-	Control	84	4	2.2 (0.8-6.2)	0.2699699

Formulation	Reference	DAI	Follow-up (days)	HR (CI 95%)	p.value
002-0.6)					
(MDC-STM-003-0.6)	Control	84	4	6.8 (2.5-18.8)	0.0000023
(MDC-STM-003-1.5)	Control	84	4	12.8 (4.7-34.9)	0.0000000
(MDC-STM-001-0.6)	Control	98	4	38.4 (8.7-169.4)	0.0000000
(MDC-STM-002-0.6)	Control	98	4	2.7 (0.5-13.6)	0.4587964
(MDC-STM-003-0.6)	Control	98	4	21.2 (4.7-94.6)	0.0000003
(MDC-STM-003-1.5)	Control	98	4	36.7 (8.3-162.4)	0.0000000
(MDC-STM-001-0.6)	Control	112	4	25.7 (6.4-103.6)	0.0000000
(MDC-STM-002-0.6)	Control	112	4	5.6 (1.3-23.6)	0.0092896
(MDC-STM-003-0.6)	Control	112	4	23.5 (5.8-95.2)	0.0000000
(MDC-STM-003-1.5)	Control	112	4	26.8 (6.6-108)	0.0000000
(MDC-STM-001-0.6)	Control	126	4	160.9 (9.3-2793.8)	0.0000118
(MDC-STM-002-0.6)	Control	126	4	29.5 (1.6-528.9)	0.0120573
(MDC-STM-003-0.6)	Control	126	4	156.2 (9-2712.6)	0.0000137
(MDC-STM-003-1.5)	Control	126	4	215.2 (12.4-3727.6)	0.0000028
(MDC-STM-001-0.6)	Control	BI	10	0.6 (0.2-2.3)	0.8747113
(MDC-STM-002-0.6)	Control	BI	10	0.5 (0.1-1.8)	0.5163317
(MDC-STM-003-0.6)	Control	BI	10	0.5 (0.2-1.9)	0.6559403
(MDC-STM-003-1.5)	Control	BI	10	0.4 (0.1-1.5)	0.2850212
(MDC-STM-001-0.6)	Control	2	10	22.3 (8.1-61.5)	0.0000000
(MDC-STM-002-0.6)	Control	2	10	44.4 (16.1-122.5)	0.0000000
(MDC-STM-003-0.6)	Control	2	10	44.1 (16-121.6)	0.0000000
(MDC-STM-003-1.5)	Control	2	10	40.8 (14.8-112.8)	0.0000000
(MDC-STM-001-0.6)	Control	17	10	22.1 (8.7-56.1)	0.0000000
(MDC-STM-002-0.6)	Control	17	10	19.9 (7.9-50.5)	0.0000000
(MDC-STM-003-0.6)	Control	17	10	19.5 (7.7-49.5)	0.0000000
(MDC-STM-003-1.5)	Control	17	10	22.9 (9-57.9)	0.0000000
(MDC-STM-	Control	21	10	19.4 (7.7-48.6)	0.0000000

Formulation	Reference	DAI	Follow-up (days)	HR (CI 95%)	p.value
001-0.6)					
(MDC-STM-002-0.6)	Control	21	10	16.7 (6.7-42)	0.0000000
(MDC-STM-003-0.6)	Control	21	10	20.7 (8.3-52)	0.0000000
(MDC-STM-003-1.5)	Control	21	10	21.3 (8.5-53.5)	0.0000000
(MDC-STM-001-0.6)	Control	42	10	12.1 (4.6-31.9)	0.0000000
(MDC-STM-002-0.6)	Control	42	10	7 (2.6-18.4)	0.0000005
(MDC-STM-003-0.6)	Control	42	10	8.9 (3.4-23.5)	0.0000000
(MDC-STM-003-1.5)	Control	42	10	12.3 (4.7-32.5)	0.0000000
(MDC-STM-001-0.6)	Control	56	10	14.4 (5.6-37.2)	0.0000000
(MDC-STM-002-0.6)	Control	56	10	7.2 (2.8-18.7)	0.0000002
(MDC-STM-003-0.6)	Control	56	10	14.5 (5.6-37.7)	0.0000000
(MDC-STM-003-1.5)	Control	56	10	21.2 (8.2-54.9)	0.0000000
(MDC-STM-001-0.6)	Control	70	10	12.3 (4.8-31.6)	0.0000000
(MDC-STM-002-0.6)	Control	70	10	3.2 (1.2-8.5)	0.0087904
(MDC-STM-003-0.6)	Control	70	10	9 (3.5-23.2)	0.0000000
(MDC-STM-003-1.5)	Control	70	10	15.7 (6.1-40.3)	0.0000000
(MDC-STM-001-0.6)	Control	84	10	8 (3.2-20.3)	0.0000000
(MDC-STM-002-0.6)	Control	84	10	2.4 (1-6.3)	0.0734027
(MDC-STM-003-0.6)	Control	84	10	5.7 (2.2-14.4)	0.0000038
(MDC-STM-003-1.5)	Control	84	10	10.1 (4-25.4)	0.0000000
(MDC-STM-001-0.6)	Control	98	10	10.1 (3.9-26.2)	0.0000000
(MDC-STM-002-0.6)	Control	98	10	2.1 (0.8-5.5)	0.2703929
(MDC-STM-003-0.6)	Control	98	10	5.7 (2.2-15.1)	0.0000082
(MDC-STM-003-1.5)	Control	98	10	9.6 (3.7-25)	0.0000000
(MDC-STM-001-0.6)	Control	112	10	10.9 (4-29.9)	0.0000000
(MDC-STM-002-0.6)	Control	112	10	3.3 (1.2-9.3)	0.0131335
(MDC-STM-003-0.6)	Control	112	10	10.7 (3.9-29.5)	0.0000000
(MDC-STM-	Control	112	10	11.5 (4.2-31.6)	0.0000000

Formulation	Reference	DAI	Follow-up (days)	HR (CI 95%)	p.value
003-1.5)					
(MDC-STM-001-0.6)	Control	126	10	34.9 (9.3-131.3)	0.0000000
(MDC-STM-002-0.6)	Control	126	10	10.5 (2.7-40.3)	0.0000181
(MDC-STM-003-0.6)	Control	126	10	33.2 (8.8-125.1)	0.0000000
(MDC-STM-003-1.5)	Control	126	10	45.5 (12.1-170.7)	0.0000000
(MDC-STM-001-0.6)	Control	BI	30	1.1 (0.5-2.3)	0.9974111
(MDC-STM-002-0.6)	Control	BI	30	1 (0.5-2.1)	1.0000000
(MDC-STM-003-0.6)	Control	BI	30	0.8 (0.4-1.8)	0.9663867
(MDC-STM-003-1.5)	Control	BI	30	0.9 (0.4-1.9)	0.9941915
(MDC-STM-001-0.6)	Control	2	30	7.4 (3.5-15.7)	0.0000000
(MDC-STM-002-0.6)	Control	2	30	18.3 (8.6-38.9)	0.0000000
(MDC-STM-003-0.6)	Control	2	30	18.4 (8.7-39.1)	0.0000000
(MDC-STM-003-1.5)	Control	2	30	12.6 (5.9-26.8)	0.0000000
(MDC-STM-001-0.6)	Control	17	30	10.6 (5.1-22.3)	0.0000000
(MDC-STM-002-0.6)	Control	17	30	9.4 (4.5-19.7)	0.0000000
(MDC-STM-003-0.6)	Control	17	30	9.4 (4.5-19.6)	0.0000000
(MDC-STM-003-1.5)	Control	17	30	11.8 (5.6-24.7)	0.0000000
(MDC-STM-001-0.6)	Control	21	30	10.1 (4.8-21.2)	0.0000000
(MDC-STM-002-0.6)	Control	21	30	7.4 (3.5-15.4)	0.0000000
(MDC-STM-003-0.6)	Control	21	30	10.7 (5.1-22.4)	0.0000000
(MDC-STM-003-1.5)	Control	21	30	11.6 (5.6-24.3)	0.0000000
(MDC-STM-001-0.6)	Control	42	30	5.9 (2.7-12.7)	0.0000000
(MDC-STM-002-0.6)	Control	42	30	3.8 (1.8-8.2)	0.0000160
(MDC-STM-003-0.6)	Control	42	30	4.7 (2.2-10.2)	0.0000003
(MDC-STM-003-1.5)	Control	42	30	6.8 (3.2-14.6)	0.0000000
(MDC-STM-001-0.6)	Control	56	30	5.5 (2.6-11.7)	0.0000000
(MDC-STM-002-0.6)	Control	56	30	3.1 (1.5-6.6)	0.0002827
(MDC-STM-	Control	56	30	6.8 (3.2-14.3)	0.0000000

Formulation	Reference	DAI	Follow-up (days)	HR (CI 95%)	p.value
003-0.6)					
(MDC-STM-003-1.5)	Control	56	30	7.1 (3.4-15.1)	0.0000000
(MDC-STM-001-0.6)	Control	70	30	3.5 (1.7-7.2)	0.0000345
(MDC-STM-002-0.6)	Control	70	30	1.4 (0.7-2.9)	0.7146990
(MDC-STM-003-0.6)	Control	70	30	2.7 (1.3-5.7)	0.0018361
(MDC-STM-003-1.5)	Control	70	30	4.8 (2.3-9.9)	0.0000001
(MDC-STM-001-0.6)	Control	84	30	3.6 (1.7-7.6)	0.0000153
(MDC-STM-002-0.6)	Control	84	30	1.5 (0.7-3.2)	0.5079043
(MDC-STM-003-0.6)	Control	84	30	2.5 (1.2-5.1)	0.0070421
(MDC-STM-003-1.5)	Control	84	30	4.2 (2-8.7)	0.0000010
(MDC-STM-001-0.6)	Control	98	30	3.8 (1.8-8.1)	0.0000099
(MDC-STM-002-0.6)	Control	98	30	1.4 (0.6-2.9)	0.7948357
(MDC-STM-003-0.6)	Control	98	30	2.8 (1.3-6)	0.0015562
(MDC-STM-003-1.5)	Control	98	30	4.4 (2.1-9.3)	0.0000008
(MDC-STM-001-0.6)	Control	112	30	3.7 (1.8-7.8)	0.0000166
(MDC-STM-002-0.6)	Control	112	30	1.5 (0.7-3.2)	0.5637848
(MDC-STM-003-0.6)	Control	112	30	3.7 (1.8-7.9)	0.0000133
(MDC-STM-003-1.5)	Control	112	30	4.4 (2.1-9.3)	0.0000006
(MDC-STM-001-0.6)	Control	126	30	3.9 (1.9-8.3)	0.0000053
(MDC-STM-002-0.6)	Control	126	30	1.7 (0.8-3.6)	0.3188277
(MDC-STM-003-0.6)	Control	126	30	4.1 (2-8.7)	0.0000023
(MDC-STM-003-1.5)	Control	126	30	4.6 (2.2-9.7)	0.0000002

Lethal Concentrations

Supplementary Figure 4

4-days mortality rates of mosquitoes fed on cattle treated with different LAIFs as a function of Ivermectin Plasmatic concentration. A. KIS colony mosquitoes. B. VK5 colony mosquitoes.

Supplementary Figure 5

10-days mortality rates of mosquitoes fed on cattle treated with different LAIFs as a function of Ivermectin Plasmatic concentration. A. KIS colony mosquitoes. B. VK5 colony mosquitoes.

Supplementary table 8

Between formulations comparisons of 4-day LC50 and L90 values for KIS mosquitoes fed on treated cattle. Model outputs are given by the drc package (ref)

Comparison	Estimate	Std. Error	t-value	p-value
001-0.6/002-0.6:50/50	0.8867454	0.5245928	-0.2158904	0.8290732
001-0.6/003-0.6:50/50	0.9276367	0.5590509	-0.1294396	0.8970098
001-0.6/003-1.5:50/50	0.9299036	0.5664743	-0.1237415	0.9015200
002-0.6/003-0.6:50/50	1.0461138	0.3575767	0.1289621	0.8973877
002-0.6/003-1.5:50/50	1.0486704	0.3703614	0.1314131	0.8954485
003-0.6/003-1.5:50/50	1.0024438	0.3723224	0.0065637	0.9947629
001-0.6/002-0.6:90/90	1.3316348	0.7287634	0.4550651	0.6490624
001-0.6/003-0.6:90/90	1.1868356	0.5802706	0.3219801	0.7474678
001-0.6/003-1.5:90/90	0.9649750	0.4975064	-0.0704012	0.9438744
002-0.6/003-0.6:90/90	0.8912621	0.4205113	-0.2585850	0.7959554
002-0.6/003-1.5:90/90	0.7246544	0.3618711	-0.7608942	0.4467203
003-0.6/003-1.5:90/90	0.8130654	0.3533909	-0.5289739	0.5968236

001 = mdc-STM-001
002 = mdc-STM-002
003 = mdc-STM-003

Supplementary table 9

Between formulations comparisons of 10-day LC50 and LC90 values for KIS mosquitoes fed on treated cattle. Model outputs are given by the drc package (ref)

Comparison	Estimate	Std. Error	t-value	p-value
001-0.6/002-0.6:50/50	0.8503147	0.4901133	-0.3054095	0.7600542
001-0.6/003-0.6:50/50	0.8484447	0.4406361	-0.3439467	0.7308864
001-0.6/003-1.5:50/50	0.9622920	0.5826786	-0.0647150	0.9484009
002-0.6/003-0.6:50/50	0.9978008	0.4234921	-0.0051931	0.9958566
002-0.6/003-1.5:50/50	1.1316892	0.5956824	0.2210728	0.8250358
003-0.6/003-1.5:50/50	1.1341835	0.5253544	0.2554152	0.7984025
001-0.6/002-0.6:90/90	0.9526083	0.5708619	-0.0830179	0.9338373
001-0.6/003-0.6:90/90	1.1594824	0.5445723	0.2928580	0.7696307
001-0.6/003-1.5:90/90	1.0292734	0.5106321	0.0573278	0.9542841
002-0.6/003-0.6:90/90	1.2171660	0.6993343	0.3105324	0.7561561
002-0.6/003-1.5:90/90	1.0804792	0.6443631	0.1248973	0.9006048
003-0.6/003-1.5:90/90	0.8877008	0.4136421	-0.2714888	0.7860151

Supplementary table 10

Between formulations comparisons of 4-day LC50 and LC90 values for VK5 mosquitoes fed on treated cattle. Model outputs are given by the drc package (ref).

Comparison	Estimate	Std. Error	t-value	p-value
001-0.6/002-0.6:50/50	0.8629191	0.2962409	-0.4627344	0.6435547
001-0.6/003-0.6:50/50	0.9532895	0.2969513	-0.1573003	0.8750082
001-0.6/003-1.5:50/50	0.9981696	0.3163292	-0.0057865	0.9953830
002-0.6/003-0.6:50/50	1.1047263	0.3647694	0.2871029	0.7740335
002-0.6/003-1.5:50/50	1.1567359	0.3878491	0.4041157	0.6861277
003-0.6/003-1.5:50/50	1.0470792	0.3169059	0.1485588	0.8819018
001-0.6/002-0.6:90/90	0.8442851	0.4320073	-0.3604451	0.7185143
001-0.6/003-0.6:90/90	0.9480422	0.4348192	-0.1194928	0.9048849
001-0.6/003-1.5:90/90	1.0899361	0.4614506	0.1948987	0.8454723
002-0.6/003-0.6:90/90	1.1228935	0.5912852	0.2078414	0.8353528
002-0.6/003-1.5:90/90	1.2909575	0.6405087	0.4542600	0.6496417
003-0.6/003-1.5:90/90	1.1496704	0.5072949	0.2950363	0.7679661

Supplementary table 11

Between formulations comparisons of 10-day LC50 and LC90 values for VK5 mosquitoes fed on treated cattle. Model outputs are given by the drc package (ref).

Comparison	Estimate	Std. Error	t-value	p-value
001-0.6/002-0.6:50/50	0.9513449	0.4334657	-0.1122468	0.9106277
001-0.6/003-0.6:50/50	0.9647061	0.4190361	-0.0842264	0.9328765
001-0.6/003-1.5:50/50	0.9823391	0.4071531	-0.0433767	0.9654013
002-0.6/003-0.6:50/50	1.0140446	0.4935095	0.0284586	0.9772964
002-0.6/003-1.5:50/50	1.0325793	0.4842848	0.0672731	0.9463643
003-0.6/003-1.5:50/50	1.0182780	0.4565668	0.0400337	0.9680663
001-0.6/002-0.6:90/90	0.9165698	0.5121319	-0.1629076	0.8705911
001-0.6/003-0.6:90/90	1.0141546	0.4742174	0.0298484	0.9761880
001-0.6/003-1.5:90/90	1.0997614	0.5007874	0.1992090	0.8420993
002-0.6/003-0.6:90/90	1.1064674	0.6179795	0.1722831	0.8632150
002-0.6/003-1.5:90/90	1.1998665	0.6578991	0.3037950	0.7612841
003-0.6/003-1.5:90/90	1.0844119	0.4934883	0.1710515	0.8641833