Species specific marker genes for systemic defence and stress responses to leaf
wounding and flagellin stimuli in hybrid aspen and silver birch

Karlis Trevors Blums[1], Baiba Krivmane[2], Maryna Ramanenka[2], Roberts Matisons[2], Dainis
Edgars Rungis[2], Martins Zeps[2], Zigmunds Orlovskis[1]*

[1] Latvian Biomedical Research and Study Centre, Ratsupites str. 1 k-1, Riga, Latvia, LV-
1067
[2] Latvian State Forest Research Institute "Silava", Rigas str. 111, Salaspils, Latvia, LV-2169
* Corresponding author: zigmunds.orlovskis@biomed.lu.lv

ORCID
Kārlis Trevors Blūms https://orcid.org/0009-0009-5290-3846
Baiba Krivmane https://orcid.org/0000-0001-9849-6806
Maryna Ramanenka https://orcid.org/0000-0002-2365-8732
Roberts Matisons https://orcid.org/0000-0002-4042-0689
Dainis Ruņģis https://orcid.org/0000-0001-5173-2912
Mārtiņš Zeps https://orcid.org/0000-0001-5064-9156
Zigmunds Orlovskis https://orcid.org/0000-0001-8583-9814

Abstract:
In Northern Europe, climate warming is driving the northward expansion of deciduous tree
species such as aspen and silver birch. At the same time, it is intensifying biotic stresses from
pests and pathogens, necessitating the development of innovative biotechnological
approaches to enhance forest resistance and resilience. However, progress in understanding
the genetic basis of stress responses in temperate deciduous trees remains limited, largely due
to the absence of reproducible laboratory model systems and well-characterized molecular
stress markers. In this study, we aimed to identify and validate known plant defense gene
markers to support future functional research on stress responses in hybrid aspen and silver
birch. We used sequence mining and phylogenetic analyses to identify homologues of biotic
stress-response genes in the genomes of both tree species. We then employed *in vitro*
propagated tree clones to test defense gene activation in distal leaves following systemic
signal induction via leaf wounding and bacterial flagellin application, assessed at 4- and 24-

hours post-treatment. We present a short communication of a novel *in vitro* methodological approach for the preliminary evaluation of molecular responses to systemic signaling in birch and aspen. The findings represent a methodological advancement for studies of systemic signalling and molecular defense responses in ecologically and economically important tree species and offer a pipeline for gene homologue discovery for the future research on tree-microbe interactions and resilience.

Keywords: tree stress responses, systemic signals, defence markers, *Populus tremuloides* × *P. tremula* L., *Betula pendula* Roth, clonal plantations, *in vitro* assays, forest productivity

1. Introduction

Accelerating climate change, intensifying weather extremes and their legacy effects, such as outbreaks of pests and diseases (Meier *et al*., 2022; Allen *et al*., 2015), are subjecting trees to increasingly intense stresses (Seidl *et al*., 2017). These stresses can exceed the adaptive capacity (i.e., phenotypic plasticity) of local tree populations (Aitken and Bemmels, 2016), emphasizing the urgent necessity for adaptive management strategies (Sousa-Silva *et al*., 2018) to cope with growing environmental challenges (Isaac-Renton *et al*., 2018). Given the long life cycle of trees and the stresses associated with large-scale environmental changes (Moran *et al*., 2017; Seidl *et al*., 2017), effective adaptive measures should be self-sustaining and grounded in a thorough understanding of ecological, ecophysiological, and biochemical processes (Jansson *et al*., 2017). Examples of such measures include locally finetuned management (Murray *et al*., 2025; Forrester and Bauhus, 2016), as well as the inoculation of plants with beneficial microbes, such as mycorrhizal fungi, which have been suggested as promising strategies to enhance forest resilience against increasing environmental stresses (Lutter *et al*., 2023; Ciadamidaro *et al*., 2017; Bois *et al*., 2005).

In the eastern Baltic region, climate-driven shifts in forest species distribution are anticipated, with deciduous species expected to push conifers northward (Buras and Menzel, 2019). Under these conditions, the prevalence of silver birch (*Betula pendula*) and aspen (*Populus tremula*) — both currently of high commercial importance — is projected to increase (Dubois *et al*., 2020; Tullus *et al*., 2012). However, both species are sensitive to climate (Matisons *et al*., 2022; Šēnhofa *et al*., 2015), and thus, an intensification of biotic legacy effects, such as insect pest outbreaks following climatic disturbances, is also expected (Gely *et al*., 2020; Seidl *et al*., 2017). Additionally, the use of specific, highly productive genotypes — such as the most economically valuable birch clones or superior hybrid aspens

— is recommended for commercial forestry (Bāders *et al*., 2024; Tullus *et al*., 2012). This
underscores the need to identify innovative biotechnological approaches for improving stress
tolerance in selected high yielding genotypes and to enhance understanding of tree stress
responses at a molecular level (Estravis-Barcala *et al*., 2020; Bradshaw *et al*., 2019; Lelu-
Walter *et al*., 2013).

Molecular stress responses have been extensively studied in herbaceous crop plants
and model species such as *Arabidopsis thaliana*, however, trees with their large and complex
genomes have been comparatively understudied and often limited to genus *Populus*
(Harfouche, 2014). This has hitherto limited functional studies on the genetic mechanisms
involved in biotic stress responses in other economically important tree species. Nevertheless,
recent advances in sequencing the genomes of important forestry species such as silver birch
(Salojarvi *et al*., 2017) or several aspen species (Lin *et al*., 2018) have catalysed recent
progress in elucidating the molecular basis of tree responses to abiotic stresses (Estravis-
Barcala *et al*. 2020). Similar to *Arabidopsis,* tree responses to chilling (frost), heat and
drought, are likely mediated by abscisic acid (ABA), ROS and Ca^{2+} signalling, mitogen-
activating protein kinases (MAPK), heat-shock proteins (HSP), MYB and bZIP family
transcription factors (among others), resulting in antioxidant production, changes in
membrane fluidity and repair (reviewed by Estravis-Barcala *et al*., 2020; Harfouche *et al*.,
2014). Still, markers for tree responses to herbivore attack or microbial infections remain to
be disclosed.

The study aimed to identify plant defence gene homologues in hybrid aspen and silver
birch and to use *in vitro* propagated tree clones for characterisation of tree leaf responses to
systemic wounding and immunogenic peptide flagellin that mimic the mechanical damage by
chewing herbivores and bacterial elicitor recognition by plant PTI receptor FLS2 (Chinchila,
2006), respectively. To this end we mined hybrid aspen and silver birch genomes for plant
wound response genes in the jasmonic acid pathway, including *LOX, JAZ* (Fürstenberg-Hägg,
2013), as well as response genes associated with salicylic acid defence responses to bacterial
elicitors, including *FRK1, NHL10, MPK3, PAD4 and MYB51* (Boudsockq *et al.*, 2010; Cui *et*
al., 2017). For the first time we employed *in vitro* propagated tree clones to identify tree
defence marker genes and characterise their activation in distal leaves after the induction of
systemic signals by leaf wounding and treatment of bacterial flagellin at different time-points.
Here we present a novel *in vitro* methodological approach for the initial testing of molecular
responses to systemic signals in birch and aspen for future downstream application in intra-
and inter-tree signalling research.

2. Materials and Methods

2.1. In-vitro propagation of tree clones

Silver birch and hybrid aspen clone cultures were obtained from the clonal collection of the LSFRI "Silava" plant physiology laboratory. For hybrid aspen *Populus tremuloides × tremula*, clone "44" was selected, as it shows superior field performance (Šēnhofa *et al*., 2016). For silver birch, clone "54-146-143", which shows above average field performance and exceeds wild populations by 7–30 % in volume growth (Gailis *et al*., 2020; Jansson *et al*., 2017), was used. The clones were cultivated *in vitro* on 1X Murashige and Skoog (MS) media, supplemented with MS vitamins, MS micronutrients, 20 g/L sucrose, 6 g/L of Agar and 0.1 mg/L indole-3-butyric acid (IBA) at pH=5.8 (Kondratovičs *et al*. 2022).

Each clone (plantlet) was cultivated individually in a 150 mL jar filled with 15 mL of the media. During the cultivation, the jars were covered with an aluminium foil cap to prevent microbial contamination whilst enabling air flow and easy access to the plantlet for manipulation and stress treatments. To ensure controlled environmental conditions, all the clone plantlets were grown in a climate chamber maintained at 30-40% relative humidity and 25 □ on four multi-layer shelf systems equipped with luminaries. All clones were grown under the same illumination with a photon flux density of 110 ± 10 µmol m^{-2} s^{-1}, for the wavelengths ranging from 400 to 750 nm with 16 h light and 8 h dark photoperiod to imitate long-day conditions of growing period. For each tree species, a total of 32 plantlets were cultivated for the assessment of stress responses in orthogonal time-course experimental design, allowing four biological replicates per treatment and observation combination.

2.2. Stimulation of local leaves and sample collection

To assess the defence-related gene expression, the plantlets were subjected to three different treatments: a chemical stimulus with 22 amino acid fragments of bacterial flagellin (*flg22*), mechanical injury to leaves and the combination of both. These stimuli were chosen to imitate a herbivore attack (mechanical injury) or the attack of a bacterial pathogen (*flg22*) to activate defence responses related to jasmonic and salicylic acid pathways, respectively (Felix *et al*., 1999; Vlot *et al*., 2009; Fürstenberg-Hägg, 2013). The mechanical injury was applied by squeezing the leaf with tweezers in three parallel lines, while the flagellin stimulus was applied by pipetting 5 µL of 1 µM *flg22* solution (in water), on the abaxial surface of the leaf. For the combined treatment, *flg22* was applied on the wounded abaxial surface of the leaf. As the control, 5 µl of distilled nuclease free water was used.

The stimuli, including the control, were applied to the third fully unfolded leaf from the apex, marking the stimulated leaf on the outside of the jar. Plantlets were treated in a sterile environment of laminar flow before returning to the climate chamber. For the gene expression analysis, a non-stimulated (naïve) leaf - second fully unfolded leaf from the apex - distal from the treated leaf was collected 4 and 24 h after the treatment to characterise the short-term dynamics of systemic plant responses. To preserve RNA, the samples were flash-frozen in liquid nitrogen and stored at -80 °C.

2.3. RNA extraction and cDNA synthesis

To extract total RNA from the leaf samples, a modified protocol from Rubio-Piña and Zapata-Pérez (2011) was used. Prior to the extraction, all labware was treated with RNaseZap™ tissues to eliminate any traces of contaminants. The 1.5- and 2-mL test tubes were submerged in 0.1% DEPC solution for 24 h and autoclaved three times at 121 °C for 40 min, using saturated steam with pressure of at least 103 kPa (15 psi), to ensure the degradation of DEPC. Leaf samples were homogenized in liquid nitrogen using a pestle and mortar before transferring the material into 2 mL test tubes with 900 µl CTAB extraction buffer (2-cetyltrimethylammonium bromide; 0.1M Tris-HCl (pH 8); 1.4M NaCl; 20mM Ethylenediaminetetraacetic acid (EDTA) (pH 8); 2% polyvinylpolypyrrolidone (PVPP)) and 100 µl of β-mercaptoethanol). The mixture was shaken for 30 s and incubated in a thermoblock for 10 min at 65 □. After incubation, 800 µl of chloroform were added to the mixture, which was followed by repeated shaking for 30 s. The mixture was then centrifuged for 10 min at 4 □ and 10'000 rpm.

The supernatant was transferred to a new 2 mL test tube together with 800 µl of a phenol/chloroform/isoamyl alcohol mixture (50:49:1) and shaken for 30 s. Samples were then centrifuged for 10 min at 10'000 rpm and 4 □ and the supernatant was transferred to a new 2 mL test tube together with an equal volume of a chloroform and isoamyl alcohol mix (24:1). The samples were then shaken for 30 s and centrifuged at 10'000 rpm at 4 □. The supernatant was transferred to a new 1.5 mL test tube together with a third of the supernatant volume of 8M LiCl. Subsequently the samples were incubated for 24 h at -20□, after which they were centrifuged for 20 min at 4 □ and 10000 rpm.

The sediment pellets were then washed with 1 mL 96% and 70% ethanol and centrifuged for 5 min at 10000 rpm. After centrifugation, ethanol was discarded, and the samples were dried at room temperature. The sediment pellets were then dissolved in 15 µl of 1x TE buffer with 1 µl of 1u/µl RNasin® Plus ribonuclease inhibitor solution (Promega). The

mix was then treated with RQ1 Rnase free Dnase kit (Promega) according to manufacturer
instructions. This process was followed by RNA extraction using a
phenol/chloroform/isoamyl alcohol mixture (50:49:1). For convenience, 230 μl of 1x TE
buffer were added to 10 μl of the sample. The phenol/chloroform/isoamyl alcohol mixture
was added at a ratio of 1:1 to the sample solution.

After the extraction, RNA was sedimented by adding 0.1 volume of 3M sodium
acetate and 2 volumes of 96% ethanol, which was followed by 24 hours of incubation at -20
⬚. Following the incubation period, the mixture was centrifuged for 15 min at 13'000 rpm
and 4 ⬚ and subsequently washed with 70% ethanol. The RNA sediment pellets were dried at
room temperature and dissolved in 20 μl of TE buffer together with 1 μl of RNase inhibitor.
The mRNA samples were stored at -80 ⬚. Total mRNA sample concentration and purity
were measured using a "NanoDrop 2000" spectrophotometer (Thermo Scientific).

2.4. rt-qPCR analysis

To carry out qPCR the extracted total mRNA was used as a template for cDNA
synthesis. Reverse transcription reaction mix was prepared using the Thermo Scientific
Maxima First Strand cDNA synthesis kit according to manufacturer instructions using 500 ng
of total RNA. The reaction was carried out using a BIO-RAD T100^{TM} thermal cycler and the
reaction mix was subsequently diluted to 800 μl. The qPCR reactions were carried out using
the Thermo Scientific Maxima SYBR Green/ROX qPCR Master Mix (2X) kit, with each
reaction containing 2 μl of 0.3 μM primer solution (containing both forward and reverse
primer), 5.5 μl of sample cDNA solution and 7.5 μl of the Maxima SYBR Green/ROX qPCR
Master Mix (2X). For each sample, three qPCR reactions (technical replicates) were carried
out with each primer pair. The resulting average Ct value was then used in further
calculations.

Primer efficiency (E%) was calculated using the formula $\left(10^{\frac{-1}{slope}} - 1\right) \times 100$, with slope
being obtained by plotting the $\log_{10}$ values of dilution factors (0-7) in a serial dilution of the
gene specific qPCR product on the x axis and average Ct value of three technical replicates
on the y axis. The expression of each gene was normalized to the expression of the chosen
reference gene *ACT2* using the formula $\frac{E_r^{Ct_r}}{E_g^{Ct_g}}$ where E_r - the amplification factor of the
reference gene, E_g - the amplification factor of the gene of interest, Ct - the average Ct value
of the three qPCR technical replicates for the reference gene and the gene of interest

respectively. In total, four expression values were obtained for each gene, for all stimuli
groups and at each time point for both birch and hybrid aspen.

2.5. Primer design

In order to observe potential differences in defence related gene expression in naïve
systemic plant tissue (distal leaf), 8 different genes with a well described role in *Arabidopsis*
thaliana defense mechanisms were chosen as the qPCR targets (**Supplementary material,**
Table S1); *EIN2, FRK1, PAD4, LOX2, MPK3, NHL10*, *JAZ10, MYB51*, with *ACT2* chosen
as a constitutively expressed reference gene for both taxa. Using gene specific protein
sequences obtained from The Arabidopsis Information Resource (TAIR), BLAST was used
to find potential defense gene homologues in the genomes of both tree taxa. To increase the
chances of finding optimal gene homologues, BLAST search was performed in two databases
for each taxon: the National Centre for Biotechnology Information (NCBI) database was used
for both taxa (*Populus tremula x tremuloides taxid:47664; Betula pendula taxid:3505)*, while
Comparative Genomics (CoGe) and PlantGenIE were used for birch (*Betula pendula scaffold*
assembly id35079, id35080) and hybrid aspen (*Populus tremula x tremuloides v10*),
respectively. The BLAST search results with the highest homology identifiers (query
coverage, percent identity and e-value) were chosen as the potential homologues of each gene
and subjected to reciprocal BLAST against *A. thaliana* and other taxa included in the NCBI
database.

A complementary phylogenetic analysis of the selected tree homologues was
performed using the MEGA (v11.0.13) Maximum likelihood algorithm (bootstrap=500). In
addition to the potential tree homologues, additional *A. thaliana*, *Medicago truncatula* and
Marchantia polymorpha sequences from NCBI and MarpolBase were included in the
phylogenetic trees. Based on the *in silico* gathered data, a single potential homologue was
selected for each gene. Using NCBI Primer-BLAST, a primer pair was designed for each
selected gene sequence. All primer pairs were designed with T_m of around 60 ☐ (with ±1 ☐
as the maximal deviation) and PCR product size of 100-300 nt.

2.6. Statistical analysis

To assess differences in gene expression between treatment groups, a two-way, single
factor analysis of variance (ANOVA) was performed along with Tukey HSD post-hoc tests in
cases of significant differences, using base R. In cases where ANOVA assumptions were not
met, non-parametric Kruskal-Wallis tests and subsequent post-hoc Dunn tests were

performed. Kruskal-Wallis tests were done using base R, while the "dunn.test" package

(Dinno A, 2024, R package version 1.3.6.) was used for the Dunn post-hoc tests. A

significance level α of 0.05 was used for all hypothesis significance tests. Heatmaps were

generated using the *pheatmap* function from the "pheatmap" package (Kolde R., 2019, R

package version 1.0.12) in R with centroid linkage clustering based on Euclidean distance. To

assess the interaction between wound or *flg22* treatments and the different time-points of

gene-expression across the 6 gene panel, a multivariate linear model was used where gene

Fold Change ~ Treatment * GeneID * Time-point. The statistical models were checked for

compliance with assumptions via diagnostic plots. Data analysis was conducted in R version

4.5.0. (R Core Team).

3. Results

3.1. Identification of plant stress response gene homologs and primer design

To identify plant defence gene homologs in silver birch and hybrid aspen, the BLAST

hits from *Arabidopsis thaliana* query protein sequence were used in an additional reciprocal

BLAST against all NCBI plant taxa. Candidate tree sequences with 99% coverage to any

plant protein with a matching gene annotation were shortlisted by the employed BLAST

algorithms. To confirm the homology, a phylogenetic analysis using all candidate

homologues for a gene of interest (GOI) from hybrid aspen (**Supplementary material,**

Figure S1) and silver birch (**Supplementary material, Figure S2**) were compared alongside

well annotated herbaceous plant (*A. thaliana* or *M. truncatula)* homologues and liverwort *M.*

polymorpha homologue as an outgroup. For each GOI, tree sequences with highest coverage

and identity to *A. thaliana* and *M. truncatula* and different from *M. polymorpha* were selected

for primer design (**Supplementary material, Table S1**). Primer pairs displaying a single-

peak melting curve, as well as yielding the desired amplicon (**Supplementary material,**

Figure S3 A, B) were further tested for their PCR efficiency (E%) which typically ranged

between 90-110% (**Supplementary material, Table S1**). Tree *ACT2* homologue was

selected as a reference gene as it displayed earlier amplification (Ct=18) and more uniform

expression across different treatments compared to *TUB*, *SAND* or *GAPDH* (**Supplementary**

material, Figure S3 C).

3.2. Characterisation of systemic response marker expression

The tested wounding and *flg22* stimuli and their combination demonstrated

significantly different effects (relative to the control) on gene expression patterns in silver

birch (p<0.01), as well as hybrid aspen (p=0.03), indicating clear inducible systemic
responses in the distal leaves (**Supplementary material, Table S2**). The gene expression
patterns showed strong dependency on time after stimulation of the local leaf, highlighting
species specific early and late response markers of biotic stress (**Figure 1**). Birch showed
significant early induction of *LOX2* (p=0.03), *MPK3* (p<0.01) and *EIN2* (p=0.03) 4h after
wounding. The effect, however, was clearly transitory as it disappeared after 24h. In contrast,
flg22 treatment did not display any significant gene induction in systemic tissues neither 4 h
nor 24 h after stimulation of the local leaves. 4h after the application of the combined
wounding and *flg22* treatment only *MPK3* expression was significantly elevated (p=0.03),
indicating potential antagonistic responses to wounding and *flg22* treatment in silver birch.

In contrast to silver birch, hybrid aspen showed significant interaction between
treatment type, identity of the activated genes and timepoints after stimulation (p<0.001)
(**Supplementary material, Table S2**), indicating complex control. Only the combined
wounding and *flg22* treatment induced significant aspen marker gene activation at 4 h and 24
h (**Figure 1**). *JAZ10* was significantly (p=0.01) induced 4h after the combined wounding and
flg22 treatment and displayed increased expression relative to the water control in the
wounded plants as well, albeit not significantly. Surprisingly, the wound-responsive JA
pathway marker *LOX2* showed no significant induction in hybrid aspen at neither 4 h nor 24
h after the stimulation. The ethylene signalling gene *EIN2* displayed significant suppression
(p=0.01) 24 h after the combined wounding and *flg22* treatment. Still, it demonstrated a
tendency to be induced after wounding, suggesting potential opposing effects of *flg22* and
wounding in the systemic responses of hybrid aspen leaves (**Figure 1**).

Figure 1. Experimental setup and relative marker gene expression in systemic leaves of hybrid aspen
(column A) and silver birch (column B) clones in response to wounding and *flg22* treatment of local
leaves. The heatmaps demonstrate $\log_2$ fold changes in gene expression (x-axis) for each treatment group (y-
axis) relative to the water control at 4 hours and 24 hours post stimulation. The green rectangles indicate
statistically significant (p<0.05) differences in gene expression in systemic leaves based on ANOVA post-hoc
tests. Local leaf stimuli are *flg22* - flagellin; *w* - wounding; *flg22 + w* - combination of flagellin with wounding.
Response similarity to treatments based on centroid linkage clustering by Euclidean distance. Full relative
expression data along with baseline expression in control treatments are displayed in **Figure S4 (Supplemental**
material).

4. Discussion

4.1. In vitro tree clones as models for plant systemic signalling research

The present study is the first to our knowledge to employ *in vitro* propagated tree clones as a model for studying intra-tree responses to stress signals and offers a scalable screening platform for design and discovery of tree stress response marker genes to diverse biotic stressors and their combinations. The applied methods provide certain potential advantages, including rapid propagation time, genetic uniformity due to clonal propagation, controlled growth and media conditions, sterile environment absent from uncontrolled pest or pathogen infections and the ability to precisely subject trees to selected defence elicitors or individual stress elements, potentially overcoming many challenges in future elucidation of molecular mechanisms for tree biotic and abiotic stress tolerance associated with heterogeneity and interaction of multiple environmental and stress factors in greenhouse and field studies (Estravis-Barcala, 2020; Harfouche *et al*., 2014). The early identification of stress markers via the *in vitro* system would accelerate follow-up functional studies in saplings grown on soil in growth chambers, greenhouse or field conditions for a range of different biological questions.

The experimental system herein allowed for optimal selection of tree stress inducing treatments and their response time estimation. For example, we discovered that only the combined treatment of wounding and *flg22* application to local leaves triggered significant systemic activation of JA-response gene *JAZ10* after 4h in hybrid aspen (**Figure 1**), while the individual treatments failed to induce consistent early responses. In contrast, leaf wounding displayed the strongest early response in silver birch compared to the other treatments. Selection of reliable and reproducible stress inducing treatments and their marker genes could be a key primer for several applications such as mechanistic studies on inter-tree signals and responses (Orlovskis *et al*., 2024) in the future.

4.2. Species specificity of SA or JA marker responses upon wounding or flg22 application

While certain markers of early responses in pattern triggered immunity (PTI) have been well described in *Arabidopsis* such as early response genes *NHL10 (NDR1/HIN1-LIKE 10)* or MAPK-specific target gene *FRK1 (FLG22-INDUCED RECEPTOR KINASE1)* (Boudsocq *et al*., 2010), no significant gene expression differences compared to the water control treatment were observed for these markers in *flg22* treatment in birch or aspen, potentially indicating different timing or species specific differences in the mechanisms of

early FLS2-mediated PTI signalling and their responses. These can be attributed to species-specific differences in the recognition of shortened *flg22* ligands by FLS2 receptor- co-receptor complexes (Mueller *et al.,* 2012) or, alternatively, heterogeneity of flagellin fragments (Colaianni *et al.,* 2021) produced by leaf endophyte communities that may interfere with responses to exogenously applied pathogenic flagellin.

PAD4 (PHYTOALEXIN DEFICIENT4) and *EDS1 (ENHANCED DISEASE SUSCEPTIBILITY1)* constitute a well-established module in PTI (Pruitt *et al.*, 2021), as well as abiotic stress (Szechynska-Hebda *et al.*, 2016) signalling and a positive regulator of salicylic acid (SA) pathway in *Arabidopsis* (Wiermer *et al.*, 2005). However, *PAD4* did not display significant response to *flg22* or combined wound and *flg22* treatments of the tree species, suggesting a potentially different *flg22* perception and signalling mechanisms than *Arabidopsis*. Both *PAD4* and plant transcription factor *MYB51* are also involved in glucosinolate and callose metabolism in innate immune responses to *flg22* (Bednarek *et al.*, 2009; Clay *et al.*, 2009) and mediate plant responses to phloem feeding insects (Kim & Jander, 2007; Luis & Shah, 2015). When tested in hybrid aspen, *MYB51* did not show any significant responses to wounding or *flg22* treatments.

LOX2 (LIPOXIGENASE2) encodes a key enzyme in jasmonic acid (JA) biosynthesis (Wasternack *et al.*, 2013) while *JAZ10 (JASMONATE ZIM-Domain10)* is an important component in downstream JA signalling (Chini *et al.*, 2007) in wound responses and control of glucosinolate production in *Arabidopsis* (Fürstenberg-Hägg *et al.*, 2013). Interestingly, *LOX2* was a good early wounding marker in birch but not in aspen (**Figure 1**), indicating species specific responses or response timing to wounding. However, *JAZ10* appeared to be a good marker for the combined wounding and *flg22* treatment in aspen and displayed upregulation upon wounding as well. Since the JA and SA pathways are considered antagonistic (Gimenez-Ibanez & Solano, 2013), this may potentially illustrate a more complex interaction between SA and JA pathways in hybrid aspen compared to *Arabidopsis*.

MAPK (MITOGEN ACTIVATED PROTEIN KINASES) are important regulators of herbivory associated local wounding as well as plant systemic responses (Hettenhausen *et al.*, 2015), and appear to be a useful marker for the systemic combined wounding and *flg22* responses in silver birch (**Figure 1**). *MPK3* and *MPK6* are known activators of ethylene (ET) biosynthesis in *Arabidopsis* (Liu & Zhang, 2006; Broekgaarden *et al*. 2015). As reviewed in Broekgaarden *et al*. (2015), ET signalling also regulates *flg22*–triggered PTI and hormonal crosstalk between SA-mediated responses to biotrophic pathogens and JA-mediated responses to wounding and chewing herbivores. *ETHYLENE INSENSITIVE2 (EIN2)* is an

important ET signalling component (Yoo *et al*., 2009) and was induced in silver birch upon
wounding, consistent with the observed *LOX2* and *MPK3* responses and involvement in
wound signalling (Broekgaarden *et al*. 2015). Interestingly, the tendency for the opposite
regulation in *EIN2* during wounding and the combined wounding and *flg22* treatment in
hybrid aspen may be indicative of the potential crosstalk between *flg22*-dependent SA
pathway and wound-inducible JA pathway in hybrid aspen.
374
Conclusion
We have identified *BpLOX2, BpMPK3* and *BpEIN2* as suitable early response
markers for wounding responses in silver birch, while *PttJAZ10* and *PttEIN2* as strong
markers for hybrid aspen response to combined wounding and flagellin application. Selection
of biotic stress treatments and response marker identification will aid future studies on intra-
and inter-tree systemic stress responses using *in vitro* and soil propagated tree clones.
381

382 **Author Contributions:** Conceptualization: ZO; methodology: KB, BK, MR, ZO;
383 investigation: KB, ZO, MB, BK; formal analysis: KB, ZO, RM; data curation: KB, ZO;
384 writing - original draft preparation: ZO, KB; writing - review and editing: ZO, KB, RM;
385 visualization: ZO, KB; supervision: ZO, DR, MZ; project administration: ZO, MZ; funding
386 acquisition: ZO. All authors have read and agreed to the published version of the manuscript.
387

388 **Funding:** This work was supported by Latvian Council of Science Fundamental and Applied
389 Research grant lzp-2022/1-0283.
390

391 **Conflicts of Interest:** The authors declare no conflicts of interest.
392

References:

Aitken, S.N., Bemmels, J.B. (2016). Time to get moving: assisted gene flow of forest trees. Evolutionary Applications, 9, 271–290. https://doi.org/10.1111/eva.12293

Allen, C.D., Breshears, D.D., McDowell, N.G. (2015). On underestimation of global vulnerability to tree mortality and forest die□off from hotter drought in the Anthropocene. Ecosphere, 6, 1–55. https://doi.org/10.1890/ES15-00203.1

Bāders, E., Zeltiņš, P., Elferts, D., Ruņģis, D., Gailis, A., & Jansons, Ā. (2024). Trends in genetic diversity of silver birch: Insights from varied planting scenarios. Baltic Forestry, 30(2), id759. https://doi.org/10.46490/BF759

Bednarek P, Pislewska-Bednarek M, Svatos A, Schneider B, Doubsky J, Mansurova M, Humphry M, Consonni C, Panstruga R, Sanchez-Vallet A, *et al*. 2009. A glucosinolate metabolism pathway in living plant cells mediates broad-spectrum antifungal defense. Science 323: 101–106. https://doi.org/10.1126/science.1163732

Bois, G., Piché, Y., Fung, M. Y. P., Khasa, D. P. (2005). Mycorrhizal inoculum potentials of pure reclamation materials and revegetated tailing sands from the Canadian oil sand industry. Mycorrhiza, 15, 149-158. https://doi.org/10.1007/s00572-004-0315-4

Boudsocq M, Willmann MR, McCormack M, Lee H, Shan L, He P, Bush J, Cheng S-H, Sheen J. 2010. Differential innate immune signalling via Ca2+ sensor protein kinases. Nature 464: 418–422. https://doi.org/10.1038/nature08794

Bradshaw, R. H., Ingvarsson, P. K., & Rosvall, O. (2019). The ecological consequences of using clones in forestry. Scandinavian Journal of Forest Research, 34(5), 380-389. https://doi.org/10.1080/02827581.2018.1557246

Broekgaarden C, Caarls L, Vos IA, Pieterse CMJ, Van Wees SCM. 2015. Ethylene: traffic controller on hormonal crossroads to defense. Plant Physiology: pp.01020.2015. https://doi.org/10.1104/pp.15.01020

Buras, A., Menzel, A. (2019). Projecting tree species composition changes of European forests for 2061–2090 under RCP 4.5 and RCP 8.5 scenarios. Frontiers in Plant Science 9, N1986. https://doi.org/10.3389/fpls.2018.01986

Chinchilla D, Bauer Z, Regenass M, Boller T, Felix G. 2006. The Arabidopsis Receptor Kinase FLS2 Binds flg22 and Determines the Specificity of Flagellin Perception. The Plant Cell 18: 465–476. https://doi.org/10.1105/tpc.105.036574

Chini A, Fonseca S, Fernández G, Adie B, Chico JM, Lorenzo O, García-Casado G, López-Vidriero I, Lozano FM, Ponce MR, *et al*. 2007. The JAZ family of repressors is the missing link in jasmonate signalling. Nature 448: 666–671. https://doi.org/10.1038/nature06006

Ciadamidaro, L., Girardclos, O., Bert, V., Zappelini, C., Yung, L., Foulon, J., ... Chalot, M. (2017). Poplar biomass production at phytomanagement sites is significantly enhanced by mycorrhizal inoculation. Environmental and Experimental Botany, 139, 48-56. https://doi.org/10.1016/j.envexpbot.2017.04.004

Clay NK, Adio AM, Denoux C, Jander G, Ausubel FM. 2009. Glucosinolate metabolites required for an Arabidopsis innate immune response. Science 323: 95–101. https://doi.org/10.1126/science.1164627

Colaianni NR, Parys K, Lee H-S, Conway JM, Kim NH, Edelbacher N, Mucyn TS, Madalinski M, Law TF, Jones CD, et al. 2021. A complex immune response to flagellin epitope variation in commensal communities. Cell Host & Microbe 29: 635-649.e9. https://doi.org/10.1016/j.chom.2021.02.006

Cui H, Gobbato E, Kracher B, Qiu J, Bautor J, Parker JE. 2017. A core function of EDS1 with PAD4 is to protect the salicylic acid defense sector in Arabidopsis immunity. The New Phytologist 213: 1802–1817. https://doi.org/10.1111/nph.14302

Dubois, H., Verkasalo, E., Claessens, H. (2020). Potential of birch (Betula pendula Roth and B. pubescens Ehrh.) for forestry and forest-based industry sector within the changing climatic and socio-economic context of Western Europe. Forests, 11, N336. https://doi.org/10.3390/f11030336.

Estravis-Barcala M, Mattera MG, Soliani C, Bellora N, Opgenoorth L, Heer K, Arana MV. 2020. Molecular bases of responses to abiotic stress in trees (R Gutiérrez, Ed.). Journal of Experimental Botany 71: 3765–3779. https://doi.org/10.1093/jxb/erz532

Felix G, Duran JD, Volko S, Boller T. 1999. Plants have a sensitive perception system for the most conserved domain of bacterial flagellin: Plants perceive a conserved domain of bacterial flagellin. The Plant Journal 18: 265–276. https://doi.org/10.1046/j.1365-313X.1999.00265.x

Forrester, D. I., & Bauhus, J. (2016). A review of processes behind diversity—productivity relationships in forests. Current Forestry Reports, 2, 45-61. https://doi.org/10.1007/s40725-016-0031-2

Fürstenberg-Hägg J, Zagrobelny M, Bak S. 2013. Plant Defense against Insect Herbivores. International Journal of Molecular Sciences 14: 10242–10297. https://doi.org/10.3390/ijms140510242

Gely, C., Laurance, S. G., Stork, N. E. (2020). How do herbivorous insects respond to drought stress in trees? Biological Reviews, 95(2), 434-448. https://doi.org/10.1111/brv.12571

Gimenez-Ibanez S, Solano R. 2013. Nuclear jasmonate and salicylate signaling and crosstalk in defense against pathogens. Frontiers in Plant Science 4. https://doi.org/10.3389/fpls.2013.00072

Harfouche A, Meilan R, Altman A. 2014. Molecular and physiological responses to abiotic stress in forest trees and their relevance to tree improvement. Tree Physiology 34: 1181–1198. https://doi.org/10.1093/treephys/tpu012

Hettenhausen C, Schuman MC, Wu J. 2015. MAPK signaling: A key element in plant defense response to insects: MAPK signaling in plant-insect interactions. Insect Science 22: 157–164. https://doi.org/10.1111/1744-7917.12128

Isaac-Renton, M., Montwé, D., Hamann, A., Spiecker, H., Cherubini, P., Treydte, K. (2018). Northern forest tree populations are physiologically maladapted to drought. Nature communications, 9(1), 5254. https://doi.org/10.1038/s41467-018-07701-0.

Yoo S-D, Cho Y, Sheen J. 2009. Emerging connections in the ethylene signaling network. Trends in Plant Science 14: 270–279. https://doi.org/10.1016/j.tplants.2009.02.007

Jansson, G., Hansen, J. K., Haapanen, M., Kvaalen, H., Steffenrem, A. (2017). The genetic and economic gains from forest tree breeding programmes in Scandinavia and Finland.

Scandinavian Journal of Forest Research, 32(4), 273-286.
https://doi.org/10.1080/02827581.2016.1242770

Kim JH, Jander G. 2007. Myzus persicae (green peach aphid) feeding on Arabidopsis induces
the formation of a deterrent indole glucosinolate. The Plant Journal: For Cell and Molecular
Biology 49: 1008–1019. https://doi.org/10.1111/j.1365-313X.2006.03019.x

Lelu-Walter, M. A., Thompson, D., Harvengt, L., Sanchez, L., Toribio, M., Paques, L. E.
(2013). Somatic embryogenesis in forestry with a focus on Europe: state-of-the-art, benefits,
challenges and future direction. Tree Genetics & Genomes, 9(4), 883-899.
https://doi.org/10.1007/s11295-013-0620-1

Liu Y, Zhang S. 2004. Phosphorylation of 1-aminocyclopropane-1-carboxylic acid synthase
by MPK6, a stress-responsive mitogen-activated protein kinase, induces ethylene
biosynthesis in Arabidopsis. The Plant Cell 16: 3386–3399.
https://doi.org/10.1105/tpc.104.026609

Louis J, Shah J. 2015. Plant defence against aphids: the PAD4 signalling nexus. Journal of
Experimental Botany 66: 449–454. https://doi.org/10.1093/jxb/eru454

Louis J, Shah J. 2015. Plant defence against aphids: the PAD4 signalling nexus. Journal of
Experimental Botany 66: 449–454. https://doi.org/10.1093/jxb/eru454

Lutter, R., *et al*., (2023). Soil fungal diversity of birch plantations on former agricultural land
resembles naturally regenerated birch stands on agricultural and forest land. Forest Ecology
and Management, 542, 121100. https://doi.org/10.1016/j.foreco.2023.121100.

Matisons, R., Jansone, D., Elferts, D., Schneck, V., Kowalczyk, J., Wojda, T., Jansons, Ā.
(2022). Silver birch shows nonlinear responses to moisture availability and temperature in the
eastern Baltic Sea region. Dendrochronologia, 76, 126003.
https://doi.org/10.1016/j.dendro.2022.126003.

Meier, H. M., Kniebusch, M., Dieterich, C., Gröger, M., Zorita, E., Elmgren, R., ... Zhang,
W. (2022). Climate change in the Baltic Sea region: a summary. Earth System Dynamics,
13(1), 457–593. https://doi.org/10.5194/esd-13-457-2022

Moran, E., Lauder, J., Musser, C., Stathos, A., Shu, M. (2017). The genetics of drought
tolerance in conifers. New Phytologist, 216(4), 1034-1048.
https://doi.org/10.1111/nph.14774.

Mueller K, Bittel P, Chinchilla D, Jehle AK, Albert M, Boller T, Felix G. 2012. Chimeric
FLS2 Receptors Reveal the Basis for Differential Flagellin Perception in Arabidopsis and
Tomato. The Plant Cell 24: 2213–2224. https://doi.org/10.1105/tpc.112.096073

Murray, S. J., Will, R. E., Gholizadeh, H., Joshi, O., Zhai, L. (2025). Structural diversity is an
important predictor of forest productivity responses to drought. Journal of Applied Ecology.
https://doi.org/10.1111/1365-2664.14883

Pruitt RN, Locci F, Wanke F, Zhang L, Saile SC, Joe A, Karelina D, Hua C, Fröhlich K, Wan
W-L, *et al*. 2021. The EDS1–PAD4–ADR1 node mediates Arabidopsis pattern-triggered
immunity. Nature 598: 495–499. https://doi.org/10.1038/s41586-021-03829-0

Rähn, E., *et al*., 2024. Soil mycobiomes in native European aspen forests and hybrid aspen
plantations have a similar fungal richness but different compositions, mainly driven by
edaphic and floristic factors. Frontiers in Microbiology, 15, 1372938.
https://doi.org/10.3389/fmicb.2024.1372938.

Rubio-Piña JA, Zapata-Pérez O. 2011. Isolation of total RNA from tissues rich in
polyphenols and polysaccharides of mangrove plants. *Electronic Journal of Biotechnology*
14: 11–11. https://doi.org/ 10.2225/vol14-issue5-fulltext-8

Seidl, R., *et al*., (2017). Forest disturbances under climate change. Nature Climate Change,
7(6), 395-402. https://doi.org/10.1038/nclimate3303

Sousa-Silva, R., *et al*., (2018). Adapting forest management to climate change in Europe:
Linking perceptions to adaptive responses. Forest Policy and Economics 90, 22–30.
https://doi.org/10.1016/j.forpol.2018.01.004

Szechyńska-Hebda M, Czarnocka W, Hebda M, Bernacki MJ, Karpiński S. 2016. PAD4,
LSD1 and EDS1 regulate drought tolerance, plant biomass production, and cell wall
properties. Plant Cell Reports 35: 527–539. https://doi.org/10.1007/s00299-015-1901-y

Šēnhofa, S., Zeps, M., Gailis, A., Kāpostiņš, R., & Jansons, Ā. (2016). Development of stem
cracks in young hybrid aspen plantations. Forestry Studies, 65(1), 16.
https://doi.org/10.1515/fsmu-2016-0008

Šēnhofa, S., Zeps, M., Matisons, R., Smilga, J., Lazdiņa, D., Jansons, Ā. (2015). Effect of
climatic factors on tree-ring width of Populus hybrids in Latvia. Silva Fennica, 50(1), 1442.
https://doi.org/10.14214/sf.1442

Tullus, A., Lukason, O., Vares, A., Padari, A., Lutter, R., Tullus, T., ... Tullus, H. (2012).
Economics of hybrid aspen (Populus tremula L.× P. tremuloides Michx.) and silver birch
(Betula pendula Roth.) plantations on abandoned agricultural lands in Estonia. Baltic
Forestry, 18(2), 288–298.

Vlot AC, Dempsey DA, Klessig DF. 2009. Salicylic Acid, a Multifaceted Hormone to
Combat Disease. Annual Review of Phytopathology 47: 177–206.
https://doi.org/10.1146/annurev.phyto.050908.135202

Wasternack C, Hause B. 2013. Jasmonates: biosynthesis, perception, signal transduction and
action in plant stress response, growth and development. An update to the 2007 review in
Annals of Botany. Annals of Botany 111: 1021–1058. https://doi.org/10.1093/aob/mct067

Wiermer M, Feys BJ, Parker JE. 2005. Plant immunity: the EDS1 regulatory node. Current
Opinion in Plant Biology 8: 383–389. https://doi.org/10.1016/j.pbi.2005.05.010

Supplementary materials

Table S1. List of silver birch and hybrid aspen marker genes used in the study. Primer
sequences are provided alongside the corresponding gene ID from TAIR, GOGE, PlantGene
databases. Amplification factor was calculated based on 10-fold template DNA dilution series
ranging from 100ag to 1ng and displayed for all genes used in **Figure 1**. Genes that did not
display a specific single band in the electrophoresis and did not produce consistent melting
curves (**Figure S3**), the amplification factor was not calculated.

Table S2. Generalized linear model results for testing the differences in gene expression between treatments across the panel of genes used in Figure 1. The model tests for the interaction of wound or *flg22* treatments with 4h and 24h timepoints and individual expression patterns of each gene (Fold Change ~ Treatment * GeneID * Time-point).

Silver birch	Df	Sum Sq	Mean Sq	F value	Pr(>F)	Significance
Treatment1	2	8,428	4,2139	5,4797	0,00545	**
Gene	5	9,33	1,866	2,4266	0,04002	*
Timepoint	1	26,904	26,9036	34,9851	4,20E-08	***
Treatment1:Gene	10	6,048	0,6048	0,7865	0,64162	
Treatment1:Timepoint	2	0,677	0,3385	0,4401	0,64513	
Gene:Timepoint	5	9,334	1,8668	2,4275	0,03995	*
Treatment1:Gene:Timepoint	10	2,668	0,2668	0,347	0,96565	
Residuals	105	80,745	0,769			
Adj. R2	0,2531					
Hybrid aspen	Df	Sum Sq	Mean Sq	F value	Pr(>F)	Significance
Treatment1	2	15,368	7,6839	3,7558	0,02661	*
Gene	5	78,781	15,7561	7,7013	3,41E-06	***
Timepoint	1	0,037	0,0367	0,0179	0,89371	

Treatment1:Gene	10	32,845	3,2845	1,6054	0,11515	
Treatment1:Timepoint	2	6,501	3,2503	1,5887	0,20911	
Gene:Timepoint	5	68,361	13,6722	6,6827	1,95E-05	***
Treatment1:Gene:Timepoint	10	112,282	11,2282	5,4881	1,63E-06	***
Residuals	104	212,774	2,0459			
Adj. R2	0,4603					

 Figure S1. Cladograms of putative *Populus tremula x tremuloides* marker gene
homologs. Phylogeny was based on translated protein sequences of *JAZ10, MYB51, EIN2,*
PAD4, LOC2, NHL10, FRK1, MPK3, and reference gene *ACT2* from *Populus tremula x*
tremuloides (PTT), *Arabidopsis thaliana* (AT), *Medicago truncatula* (MT), *Marchantia*
polymorpha (MarPol). In case of multiple spliced variants, all sequences were included for
tree construction using the Maximum likelihood method. Green box denotes the homolog
used for primer design in the study.

Figure S2. Cladograms of putative *Betula pendula* marker gene homologs. Phylogeny
was based on translated protein sequences of *JAZ10, MYB51, EIN2, PAD4, LOC2, NHL10,*
FRK1, MPK3, and reference gene *ACT2* from *Betula pendula* (BP), *Arabidopsis thaliana*
(AT), *Medicago truncatula* (MT), *Marchantia polymorpha* (MarPol). In case of multiple
spliced variants, all sequences were included for tree construction using the Maximum
likelihood method. Green box denotes the homolog used for primer design in the study.

Figure S3. Electrophoresis gels of stress marker gene amplicons in *Betula pendula* (A) and *Populus tremula x tremuloides* (B) as well as marker gene amplification curves for tree reference gene selection (C). 35-cycle PCR amplicons were used to check for primer specificity, target size and absence of double bands. rt-qPCR analysis was performed for all primers with a visible single band to check for a single uniform melting-curve. Reference gene *ACT2* was selected based on earlier amplification and uniform expression across the different stress treatments in the qPCR amplification curve **(C).** deltaRn indicates the intensity of SYBR signal corresponding to target amplification during 40 qPCR cycles.

Figure S4. Marker gene expression relative to reference gene in *Betula pendula* (A) and *Populus tremula x tremuloides* (B). Boxplots represent the interquartile range (IQR), the line indicates the median expression, the whiskers - variance within 1.5x IQR. Summary results of ANOVA or Kruskal-Wallis and corresponding post-hoc tests are provided for each marker gene.

A — Populus tremula x tremuloides
Systemic response / Local stimulus

B — Betula pendula
Systemic response / Local stimulus

4 hours

log₂ fold change

flg22
w
flg22 + w

JAZ10 MYB51 EIN2 PAD4 LOX2 NHL10

4 hours

log₂ fold change

flg22
w
flg22 + w

LOX2 PAD4 MPK3 NHL10 EIN2 FRK1

24 hours

w
flg22
flg22 + w

EIN2 JAZ10 MYB51 NHL10 LOX2 PAD4

24 hours

w
flg22
flg22 + w

LOX2 PAD4 EIN2 FRK1 MPK3 NHL10