Response of UBR-box E3 ubiquitin ligases and protein quality control pathways to perturbations in protein synthesis and skeletal muscle size

Leslie M. Baehr[1], Luis Gustavo Oliveira de Sousa[1], Craig A. Goodman[2], Adam P. Sharples[3], David S. Waddell[4], Sue C. Bodine[1] & David C. Hughes[1*]

[1] Aging and Metabolism Research Program, Oklahoma Medical Research Foundation, Oklahoma City, OK, USA

[2] Centre for Muscle Research (CMR), Department of Anatomy and Physiology, The University of Melbourne, Victoria, AUS.

[3] Institute for Physical Performance, Norwegian School of Sport Sciences (NiH), Oslo, Norway

[4] University of North Florida, Department of Biology, 1 UNF Drive, Jacksonville, Florida, 32224, USA.

*Corresponding Author: D.C Hughes (david-hughes@omrf.org). Aging and Metabolism Research Program, Oklahoma Medical Research Foundation, 825 NE 13th St. Oklahoma City, OK 73104.

Abstract

The N-degron pathway contributes to proteolysis by targeting N-terminal residues of destabilized proteins via E3 ligases that contain a UBR-box domain. Emerging evidence suggests the UBR-box family of E3 ubiquitin ligases (UBR1-7) are involved in the positive regulation of skeletal muscle mass. The purpose of this study was to explore the role of UBR-box E3 ubiquitin ligases under enhanced protein synthesis and skeletal muscle growth conditions. Cohorts of adult male mice were electroporated with constitutively active Akt (Akt-CA) or UBR5 RNAi constructs with a rapamycin diet intervention for 7 and 30 days, respectively. In addition, the UBR-box family was studied during the regrowth phase post nerve crush induced inactivity. Skeletal muscle growth with Akt-CA or regrowth following inactivity increased protein abundance of UBR1, UBR2, UBR4, UBR5 and UBR7. This occurred with corresponding increases in Akt-mTORC1/S6K and MAPK/p90RSK signaling and protein synthesis. The increases in UBR-box E3s, ubiquitination, and proteasomal activity occurred independently of mTORC1 activity and were associated with increases in markers related to autophagy, ER-stress, and protein quality control pathways. Finally, while UBR5 knockdown (KD) evokes atrophy, it occurs together with hyperactivation of mTORC1 and protein synthesis. In UBR5 KD muscles, we identified an increase in protein abundance for UBR2, UBR4 and UBR7, which may highlight a compensatory response to maintain proteome integrity. Future studies will seek to understand the role of UBR-box E3s towards protein quality control in skeletal muscle plasticity.

Key Words: hypertrophy, UBR5, proteasome system, N-degron pathway

Running head: UBR-box E3 ligases and skeletal muscle size

New and Noteworthy: Novel UBR-box E3 ubiquitin ligases are responsive to heightened protein synthesis and alterations in skeletal muscle mass and fiber size, in order to maintain proteome integrity.

Introduction

Skeletal muscle is a highly adaptable tissue that can respond to various stimuli (e.g., inactivity, disease, and exercise) and has the ability to increase (i.e. hypertrophy) or decrease (i.e. atrophy) in size when required (1-3). The regulation of skeletal muscle mass is orchestrated by the activity of key signaling pathways that control protein breakdown and synthesis within myofibers (2). Early seminal work highlighted the critical role for the Akt-mTORC1-p70S6K1 signaling pathway in skeletal muscle mass regulation (4, 5). Since this discovery, our understanding of growth signals in skeletal muscle has evolved towards mTORC1-dependent and mTORC1-independent mechanisms, along with other processes that may contribute to hypertrophy, such as ribosome biogenesis (6-13). Genetic models, such as transgenic constitutively active Akt (Akt-CA) and TSC1 knockout mice, have expanded our knowledge of the mTORC1 signaling pathway and the regulation of muscle mass (14-19). Indeed, the transgenic Akt mice display a phenotype of increased skeletal muscle fiber size and protein synthesis, while electroporation-mediated transfection of an Akt-CA construct elicits similar effects on skeletal muscle growth (4, 19, 20). Previously published evidence from the TSC1 knockout mouse has observed mTORC1 to be an inhibitor of autophagy and regulate the ubiquitin proteasome system (UPS) (19, 21). The regulation of the UPS by mTORC1 directly remains to be determined as the UPS may be active in response to mTORC1-mediated activation of translation, where the UPS is needed to maintaining proteome integrity. However, given that the characterization of the UPS has predominantly centered on atrophy models, and only a handful of E3 ubiquitin ligases have been studied to date (e.g. MuRF1, MAFbx, and MUSA1) (22-24), very little is known about the UPS in skeletal muscle remodeling and growth. Thus, our understanding of the UPS, and other protein quality control systems (e.g. unfolded protein response, autophagy), could be expanded by characterizing new targets or exploring different contexts and scenarios where these systems are influential in tissue health, such as skeletal muscle growth and remodeling.

The UBR-box family of E3 ubiquitin ligases, which consists of HECT, RING and F-box E3 ubiquitin ligases named UBR1 through UBR7, encompasses part of the ubiquitin proteasome system, and their action on protein substrate recognition forms the

N-degron pathway (25, 26). The N-degron pathway (also known as the N-end rule

pathway) identifies degradation signals on destabilized proteins through the presence of

ubiquitin and acetyl motifs (25, 27). As part of the N-degron pathway, the UBR-box

family of E3 ubiquitin ligases identify unacetylated protein substrates for ubiquitination

via the UBR-box domain.(25, 27-29). Studies have highlighted that UBR-box E3s play

diverse roles in a range of cellular processes, with protein quality control and protein

clearance typically being referenced (30-33). However, very little is known about this

family of E3 ubiquitin ligases in skeletal muscle. In an early study from the Goldberg lab

(34), various inhibitors targeting E3α (also known as UBR1) showed an ~60%

contribution of this proteolytic pathway to the turnover of endogenous proteins in

mammalian skeletal muscle. More recently, studies have identified UBR4, UBR5 and

UBR7 to be responsive to exercise stimuli (35-39) and UBR2 for being important in the

development of cancer cachexia (40, 41). Our previous studies have identified a

potential role for UBR5 in skeletal muscle growth and recovery following disuse (42).

Further, UBR5 knockdown in mouse skeletal muscle perturbed Akt-mTORC1-p70S6K1

and MAPK/p90RSK signaling pathways, that coincided with a loss of skeletal muscle

mass and fiber cross-sectional area (CSA) (43). Thus, it has been observed that UBR-

box E3 ubiquitin ligases may be important for skeletal muscle health and the regulation

of skeletal muscle mass (30, 36, 37, 42-44).

Therefore, we sought to investigate if UBR-box E3 ubiquitin ligases and protein

quality control systems are responsive to genetic manipulation of Akt-mTORC1

signaling and the associated increase in protein synthesis and skeletal muscle size. In

addition, we wanted to ascertain if these effects were dependent on mTORC1 activity.

Lastly, given our previous observations of UBR5 in models of skeletal muscle regrowth

following a period of atrophy (42), we hypothesized that UBR1, UBR2, UBR4 and UBR7

might display a similar temporal profile to UBR5 during the regrowth phase of the nerve

crush injury model.

Materials and methods

Study approval for animal experiments

The mice used in these studies were males from the C57BL/6N strain obtained from
Charles River laboratories at 5-6 months of age and used for experiments within 2
weeks of their arrival. Animals were housed in ventilated cages maintained in a room at
21 °C with 12-h light/ dark cycles and had *ad libitum* access to standard chow (LabDiet
PicoLab® 5053) and water throughout the study. All animal procedures were approved
by the Institutional Animal Care and Use Committee at the Oklahoma Medical Research
Foundation.

Rapamycin Diet Intervention
Male C57BL/6 mice received either a control diet (PicoLab® 5053 (LabDiet®) containing
Eudragit) or a diet containing 42 mg/kg chow active encapsulated rapamycin (Rapa;
Rapamycin Holdings, Inc., San Antonio, TX) which was provided ad libitum. The
Eudragit in the chow serves as a true experimental control, given that the rapamycin is
encapsulated by this material in order to provide effective release of the drug. For the
Akt overexpression experiments, age-matched cohorts of mice were randomized into
control or rapamycin diet interventions for 7 days prior to the electroporation procedure
and remained on the respective diet until completion (total time equaled 14 days)
(Figure 1). For the UBR5 knockdown experiments, age-matched cohorts of mice were
electroporated and randomized into control or rapamycin diet interventions and
remained on the diets for a total of 30 days (Figure 1). Chronic treatment of rapamycin
has been observed to target mTORC1 and mTORC2 for inhibition (45). The rapamycin
dose implemented has been reported to inhibit mTORC1 activity (46-48) and we
confirmed mTORC1 inhibition by assessing p70S6K1 and rpS6 phosphorylation as
downstream markers for mTORC1 activity. Body weight was assessed throughout the
dietary interventions, and no difference was observed between control and rapamycin-
treated mice over the 14 to 30 days intervention (data not shown).

Akt and UBR5 Plasmid Constructs
For the Akt experiments, the open reading frame of mouse Akt1 (Akt-WT) was cloned
and ligated into the pCMV5 expression plasmid and fused with a hemagglutinin (HA) tag
at the carboxyl-terminus. To create the constitutively active (Akt-CA) construct, a
consensus myristylation sequence (MGSSKSKPKSR) was fused to the amino terminus

of HA-tagged, wild-type mouse Akt1 in the pCMV5 expression plasmid. For RNAi

experiments, the negative control RNAi plasmid was described previously (42, 49) and

encodes emerald green fluorescent protein (EmGFP) and a non-targeting pre-miRNA

under bicistronic control of the cytomegalovirus (CMV) promoter in the

pcDNA6.2GW/EmGFP-miR plasmid (Invitrogen, Carlsbad, CA). The UBR5 RNAi

plasmid encodes EmGFP and an artificial pre-miRNA targeting the full-length gene of

mouse UBR5 under bicistronic control of the CMV promoter; it was generated by

ligating the Mmi571982 oligonucleotide duplex (Invitrogen, Carlsbad, CA) into the

pcDNA6.2GW/EmGFP-miR plasmid. The successful knockdown of UBR5 mRNA

expression and protein levels in mouse skeletal muscle was previously screened and

confirmed at approximately 50-60% (42, 43).

In vivo electroporation

Transfection of mouse skeletal muscle with plasmid DNA was performed in mice under

isoflurane (2-4% inhalation) anesthesia as previously described (50-52). Briefly, after a

2h pre-treatment with 0.4 units/µl of bovine placental hyaluronidase (Sigma)

resuspended in sterile 0.9% saline, 20µg of plasmid DNA was injected into the tibialis

anterior (TA) muscle. The hind limbs were placed between two-paddle electrodes and

subjected to 10 pulses (20 msec) of 175 V/cm using an ECM-830 electroporator (BTX

Harvard Apparatus, Holliston, MA). For Akt overexpression studies, an additional 2 µg

of emerald-GFP plasmid was electroporated for identification of GFP positive fibers.

Following transfection, mice were returned to their cages to resume normal activities

until tissue collection.

Nerve Crush Injury Model

Gastrocnemius complex (GSTC) muscle samples from our previously published study

(42) were reanalyzed for protein abundance of UBR-box E3 ubiquitin ligases (UBR1,

UBR2, UBR4, UBR5 and UBR7) via immunoblotting. Targeted nerve crush in the lower

limb muscles of the right leg was accomplished via acute 10 s crushing of the sciatic

nerve in the midthigh region of mice using forceps. The crushing of the sciatic nerve

was performed by the same investigator for all cohorts of mice with firm pressure being

applied for the 10 s time period. The procedure was performed under isoflurane

anesthesia (3% inhalation) using aseptic surgical techniques. Male mice (C57BL/6; 12

weeks old) purchased from Charles River Laboratories (Wilmington, MA) were given an

analgesic (buprenorphine, 0.1 mg kg^{-1}) immediately, as well as for 48 h following

surgery, and returned to their cage following recovery. Following completion of the

appropriate time point (n=6/group), mice were anesthetized with 3% isoflurane, and the

gastrocnemius complex (GSTC) muscles were excised, weighed, frozen in liquid

nitrogen and stored at −80°C for later analysis. Prior to tissue collection, a sciatic nerve

test was applied via a single electrical pulse to the sciatic nerve through electrodes to

observe a twitch response in the hindlimb muscles. Between 21 and 60 days, all

animals displayed a twitch, suggesting that neural activity to the hindlimb muscles had

returned at these time points. A separate untreated cohort of animals (n = 4/group) was

used as the relevant control. We assessed UBR-box E3 ubiquitin ligase protein

abundance at time points 21-, 28-, 45-, and 60-days post nerve crush injury which

coincided with reinnervation and reflective of skeletal muscle regrowth following a

period of inactivity.

Tissue Collection

Following completion of the appropriate time period following the electroporation

procedure (i.e. 7 or 30 days), mice were anesthetized with isoflurane, the TA muscles

were excised, weighed, frozen in liquid nitrogen, and stored at −80°C for later analysis.

A subset of muscles was collected for histology and processed as described below. On

completion of tissue removal, mice were euthanized by exsanguination.

Histology

Harvested muscles were immediately fixed in 4% (w/v) paraformaldehyde for 16h at

4°C. Following a sucrose gradient incubation period (10%, 20%, and 30% for 2-3 hours

each), the TA muscles were embedded in Tissue Freezing Medium (Triangle

Biomedical Sciences), and a Thermo HM525 cryostat was used to prepare 10-μm serial

sections from the muscle midbelly. All sections were examined and photographed using

a Nikon Eclipse Ti automated inverted microscope equipped with NIS-Elements BR

digital imaging software at 10x and 20x magnification for laminin and Hematoxylin &
Eosin Stain respectively.

Laminin Stain: TA muscle sections were permeabilized in PBS with 1% triton for 10 min
at room temperature. After washing with PBS, sections were blocked with 5% goat
serum for 15 min at room temperature. Sections were incubated with Anti-Laminin
(1:500, Sigma Aldrich, Cat no. L9393) in 5% goat serum for 2 h at room temperature,
followed by two 5-min washes with PBS. Goat-anti-rabbit AlexaFluor® 555 secondary
(1:333, Invitrogen Cat no. A28180) in 5% goat serum was then added for 1 h at room
temperature. Slides were cover slipped using ProLong Gold Antifade reagent (Life
Technologies). Image analysis was performed using Myovision software (53, 54).
Skeletal muscle fiber size was analyzed by measuring ≥ 400 transfected muscle fibers
per muscle (GFP-positive), per animal (10x magnification). In some muscles the
transfection frequency was so high that there were few GFP-negative fibers within the
same region as the GFP-positive fibers, as has been observed previously (43, 50, 51).
Therefore, for Akt-CA overexpression and UBR5 knockdown (RNAi) studies, fiber size
comparisons were made between the GFP-positive fibers in the EV controls and Akt-CA
or UBR5 RNAi transected muscles with and without the rapamycin diet.

Hematoxylin & Eosin Stain: A standard H&E protocol was performed (adapted from
Parlee et al. (55)). Briefly, TA muscle sections at room temperature were incubated in
deionized H_2O for 2 min on a rocker at 30 rpm (Labnet International Inc). Next, slides
were incubated with Mayer's hematoxylin solution (EMS SKU: 26043-05) for 10 min to
stain the nuclei, rinsed in running H_2O until clear, dipped 3 times in blue solution, and
rinsed under running H_2O for 2 min. This was followed by incubation in 80% ethanol for
2 min. TA sections were stained with 0.1% Eosin (Sigma E4382-25G) solution for 10 s,
followed by sequential immersion in graded ethanol concentrations for 2 min each (1 x
80%, 2 x 95%, 2 x 100%). Slides were then incubated in xylene solution (three times, 3
min each). A coverslip was applied using Permount™ media (Thermo Fisher Scientific,
Waltham, MA). Slides were placed horizontally at room temperature to air dry for at
least 24 hours before imaging. The percentage of aberrant myofibers was calculated by
dividing the number of fibers displaying aberrant/vacuole structures by the total number

of fibers (19, 56, 57). An average of 1500-1800 myofibers were counted per animal (n =
4-5/group), and quantification was performed using ImageJ software (National Institutes
of Health, Bethesda, MD, USA).

Muscle Protein Synthesis (Puromycin Incorporation)
Changes in muscle protein synthesis (MPS) were assessed in TA muscles transfected
for 7 and 30 days by measuring the incorporation of exogenous puromycin into nascent
peptides as described previously (6, 58). Puromycin (EMD Millipore, Billerica, MA; Cat.
No. 540222) was dissolved in sterile saline and delivered (0.02 µmol/g body wt by ip
injection) 30 min before muscle collection. Protein synthesis was measured under fed
conditions and studied in the light cycle.

Immunoblotting
Frozen TA muscles were homogenized in sucrose lysis buffer (50 mM Tris pH 7.5, 250
mM sucrose, 1 mM EDTA, 1 mM EGTA, 1% Triton X 100, 50 mM NaF). The
supernatant was collected following centrifugation at 8,000 *g* for 10 min and protein
concentrations were determined using the 660 protein assay (Thermo Fisher Scientific,
Waltham, MA). Twenty micrograms of protein were subjected to SDS-PAGE on 4-20%
Criterion TGX stain-free gels (Bio-Rad, Hercules, CA) and transferred to polyvinylidene
diflouride membranes (PVDF, Millipore, Burlington, MA). Membranes were blocked in
3% nonfat milk in Tris-buffered saline with 0.1% Tween-20 added for one hour and then
probed with primary antibody overnight at 4°C. Membranes were washed and incubated
with HRP-conjugated secondary antibodies at 1:10,000 for one hour at room
temperature (Cell Signaling technology Cat no. 7076 and Cat no. 7074). Immobilon
Western Chemiluminescent HRP substrate was then applied to the membranes prior to
image acquisition. Image acquisition and band quantification were performed using the
ChemiDoc MP System and Image Laboratory 6.1 software (Bio-Rad), respectively.
Total protein loading of the membranes captured from images using stain-free gel
technology was used as the normalization control for all blots. The following antibodies
were used at 1:1000 concentration unless otherwise stated: Total Ubiquitin (Millipore,
RRID:AB_2043482 and Cell Signaling, RRID:AB_2799235), UBR1 (ProteinTech, Cat

no. 260069), UBR2 (Abcam, Cat no. 217069), UBR4 (Abcam, Cat no. 86738), UBR5 (Protein Tech, Cet no. 66937), UBR7 (Novus Biologicals, Cat no. NBP1-88409), VCP (Protein Tech, RRID:AB_2214635), p62 (Sigma, RRID:AB_1841064), LC3B (Sigma, Cat no. L7543), eIF2A (Protein Tech, RRID:AB_2096489), NDRG1 (Protein Tech, RRID:AB_2880676). Cell Signaling Technologies (Danvers, MA) – K48 Ub-linkage (RRID:AB_10859893), phospho-p44/42 MAPK[Thr202/Tyr204] (RRID:AB_2315112), phospho-p90RSK[Ser380] (RRID:AB_2687613), p90RSK (RRID:AB_659900), phospho-Akt[Ser473] (RRID:AB_2315049), phospho-Akt[Thr308] (RRID:AB_2629447), Akt (RRID:AB_329827), phospho-NDRG1[Thr346] (RRID:AB_10693451), Raptor (RRID:AB_561245), phospho-p70S6K Thr[389] (RRID:AB_330944), p70S6K (RRID:AB_331676), phospho-rpS6[Ser240/244] (RRID:AB_10694233), rpS6 (RRID:AB_331355), phospho-4EBP1 [Thr37/46] (RRID: AB_560835), 4EBP1 (RRID:AB_2097841), eIF4E (RRID:AB_823488), eIF4G (RRID:AB_2096025), BiP (RRID:AB_2119845), PDI (RRID:AB_2156433), CHOP (RRID:AB_2089254). EMD Millipore–puromycin (RRID:AB_2566826). Anti-rabbit IgG, HRP-linked (1:10 000, Cell Signaling, Cat no. 7074) and anti-mouse IgG, HRP-linked (1:10 000, Cell Signaling, Cat no. 7076) were used as secondary antibodies.

Proteasome Activity Assays

20S and 26S proteasome activities were performed as previously described (59, 60). The 26S ATP-dependent assays were performed in homogenization buffer (50 mM Tris, 150 mM NaCl, 1 mM EDTA, 5 mM $MgCl_2$ [pH 7.5] and 0.5mM dithiothreitol (DTT)) with the addition of 100 µM ATP. The 20S ATP-independent assays were carried out in assay buffer containing 25 mM HEPES, 0.5 mM EDTA, and 0.001% SDS (pH 7.5). Proteasome activities were determined by adding substrates at 100 µM: Z-Leu-Leu-Glu-MCA (Peptide Institute, catalog no. 3179-v), Boc-Leu-Ser-Thr-Arg-AMC (Bachem, catalog no. I-1940), or succinyl-Leu-Leu-Val-Tyr-7-AMC (Bachem, catalog no. I-1395), for β_1- (caspase-like), β_2- (trypsin-like), and β_5-subunits (chymotrypsin-like), respectively. Each assay was conducted in the absence and presence of the proteasome inhibitor bortezomib (Cell Signaling, catalog no. 2204) at a final concentration of 2 mM (β_5) or 10 mM (β_1 and β_2). All proteasome assays were conducted using 10 µg of protein/well on a 96-well plate (Greiner Bio-One, catalog no.

655076), and each sample was loaded in triplicate on the plate. The activity of the 20S

and 26S proteasome was measured by calculating the difference between fluorescence

units recorded with or without the inhibitor in the reaction medium. Fluorescence was

measured using a Spectra Max M2 Fluorescent Microplate reader (Molecular Devices,

Sunnyvale, CA; excitation wavelength, 360 nm; emission wavelength, 460 nm) at 15-

min intervals for 2 h. To compare all groups on the same plate e.g. control vs rapamycin

diet and empty vector vs Akt-CA constructs, we utilized a sample of N=4 per group to

allow for technical replicates on the same assay plate.

Gene expression by Quantitative RT-PCR in skeletal muscle

Frozen muscle powder was homogenized using RNAzol RT reagent (Sigma-Aldrich, St

Louis, MO) in accordance with the manufacturer's instructions. cDNA was synthesized

using a reverse transcription kit (High-Capacity cDNA synthesis kit; Applied Biosystems,

Waltham, MA) from 1 μg of total RNA. PCR reactions (10 μL) were set up as: 2 μL of

cDNA, 0.5 μL (10 μM stock) forward and reverse primer, 5 μL of Power SYBR Green

master mix (Thermo Fisher Scientific, Waltham, MA) and 2 μL of RNA/DNA free water.

Gene expression analysis was then performed by quantitative PCR on a Quantstudio 6

Flex Real-time PCR System (Applied Biosystems, Waltham, MA) using the mouse

primers shown in Table 1. PCR cycling comprised: hold at 50°C for 2 min, 10 min hold

at 95°C, before 40 PCR cycles of 95°C for 15 s followed by annealing temp (see Table

1) for 30 s, and extension at 72°C for 30 s). Melt curve analysis at the end of the PCR

cycling protocol yielded a single peak. As a result of reference gene instability, gene

expression was normalized to tissue weight and subsequently reported as the fold

change relative to empty vector control muscles, as described previously (61). This type

of analysis has previously been used extensively by our group (62-64).

Statistical Analysis

Results are presented as mean ± standard error of measure (SEM) for all experiments.

Statistical differences for in vivo overexpression studies were determined using a paired

students' t-test on muscle mass and fiber CSA data sets. For data sets with the Akt-Wt

and Akt-CA construct comparisons, a one-way analysis of variance (ANOVA) was

performed followed by Dunnett's test for multiple comparisons. For data sets generated

with diet (control and rapa) and construct experiments, a two-way ANOVA was

performed. When applicable, a Šídák's post hoc test was conducted for multiple

comparisons across groups. Potential outliers within data sets were identified and

excluded using a Grubbs test. All statistical analyses were performed using GraphPad

Prism (GraphPad Software, Inc., La Jolla, CA). Results were considered significant

when $P \leq 0.05$.

Results

Effect of differing levels of Akt activation on mTORC1 signaling and skeletal muscle

mass and fiber cross-sectional area

To explore the role of protein quality control systems in response to skeletal muscle

growth, we utilized an Akt overexpression model where either a wild-type (WT) or

constitutively active (CA) Akt construct was electroporated into male TA skeletal

muscles for 7 days (Figure 1A). We observed a significant increase in TA muscle mass

with the Akt-WT (+4%) and Akt-CA (+12%) constructs when compared to the respective

empty vector (EV) control (Figure 2A; $P \leq 0.05$). Skeletal muscle fiber CSA was

assessed in EV control and Akt-CA transfected TA muscles with GFP-positive fibers

being analyzed (Figure 2B). We observed a shift in GFP-positive fiber CSA distribution

towards both smaller and larger fiber sizes with the Akt-CA construct verses the EV

control (Figure 2B). Lastly, overexpression of Akt-WT and Akt-CA increased Akt

phosphorylation and phosphorylation of downstream mTORC1 substrates, rpS6 and

p70S6K1, compared to the EV control (Figure 2C and 2E). In addition, we saw a

significant increase in protein abundance for eIF4E, eIF4G and puromycin labeling,

indicating elevated protein synthesis with Akt-CA overexpression (Figure 2C and 2E).

Changes in UBR-box E3 ubiquitin ligase protein abundance and mRNA expression in

Akt-transfected skeletal muscles

Given the proposed role for UBR-box E3 ubiquitin ligases in protein quality control and

previous literature observing UBR5 to be elevated in models of skeletal muscle growth

(35, 42), we assessed UBR-box E3 ubiquitin ligase expression in experimental models

where protein synthesis and skeletal muscle mass are increased. In our Akt-WT and

Akt-CA overexpression models, we observed increases in the abundance of UBR1,

UBR2, UBR4, UBR5 and UBR7 proteins (Figure 2D and 2E). In addition, we observed

increased levels of protein ubiquitination only in the Akt-CA transfected muscles

compared to the EV control (Figure 2D). At the mRNA level, we found increased

expression of UBR2, UBR4 and UBR7 in Akt-CA transfected muscles (Figure 2F). We

observed no changes in UBR1, UBR5, MuRF1 or MAFbx mRNA expression with either

Akt-WT or Akt-CA compared to the EV control (Figure 2F).

UBR-box E3 ubiquitin ligases are elevated during a period of skeletal muscle regrowth

following neural inactivity

To extend our observations, we analyzed UBR-box E3 ubiquitin ligase protein

abundance in gastrocnemius muscle from a previously published nerve crush injury

model (42). Between 21 and 60 days post nerve crush injury, skeletal muscle regrowth

occurs following a period of neural inactivity (42). All the UBR-box E3 ubiquitin ligases

displayed increased protein abundance between 21 and 45 days post-surgery (Figure

3A and 3B). Only UBR1 and UBR7 protein remained elevated above control tissue

levels at the 60 day time point (Figure 3B). The changes in protein abundance for

UBR1, UBR2, UBR4 and UBR7 correspond with our previously published observations

for increased UBR5 protein abundance in the nerve crush injury model (42) during a

reinnervation period when skeletal muscle mass and the protein synthetic machinery

are remodeling.

Effect of rapamycin on Akt-induced skeletal muscle growth, rapamycin-sensitive and

insensitive signaling

Next, we investigated whether changes in the content of these ubiquitin ligases with Akt

overexpression were dependent on mTORC1 activity (Figure 1B), through the

implementation of a rapamycin diet. In the control diet, there was a significant increase

in TA muscle mass with Akt-CA overexpression, whereas the rapamycin diet blunted

Akt-CA induced growth (Figure 4A). At the fiber CSA level, GFP-positive fibers were

distributed towards larger and smaller size compared to the EV transfected muscles

under the control diet with Akt overexpression (Figure 4B and 4C). However, the GFP-
positive fibers in the rapamycin diet condition displayed only smaller fiber size in the
Akt-CA transfected muscles, with a reduction in the percentage of larger fibers being
observed (Figure 4B and 4C).

In terms of Akt-mTORC1 signaling, we observed increased phosphorylation of
Akt, S6K and rpS6 under control diet conditions with Akt-CA overexpression (Figure 4D
and 4E). Conversely, these downstream targets of mTORC1 were either non-detectable
or blunted in the rapamycin diet condition (Figure 4D). However, rapamycin did not
impact 4E-BP1 phosphorylation, suggesting that mTORC1 signals might not have been
fully blocked. Further, we report increased protein abundance for raptor and eIF4E
under control and rapamycin diet conditions with Akt-CA overexpression. Given that Akt
is upstream of mTORC1 activation and can initiate signaling of other independent
pathways, we assessed mTORC2 and p44/42 MAPK activity. Phosphorylation of
NDRG1 (mTORC2 substrate), p44/42 MAPK and p90RSK were all significantly elevated
irrespective of the diet conditions in transfected TA muscles with Akt-CA (Figure 4E).
Lastly, puromycin-labeling for nascent peptides was increased in Akt-CA
overexpression muscles irrespective of mTORC1 activity, highlighting the contribution of
rapamycin-insensitive mechanisms to increases in the protein synthetic machinery
(Figure 4D and 4E). It should be noted that the magnitude of increase for the
puromycin-labelling was greater in the control group verses the rapamycin treated
groups.

Changes in UBR-box E3 ubiquitin ligases, ubiquitination and proteasomal activity are
independent of mTORC1 activity
In control and rapamycin diet conditions, we assessed UBR-box E3 ubiquitin ligase
protein levels to ascertain if changes occurred in a rapamycin-sensitive or insensitive
manner. Protein levels for UBR2, UBR4, UBR5 and UBR7 significantly increased in the
Akt-CA overexpression muscles independent of the diet condition (Figure 5A and 5B).
Interestingly, UBR1 increased in the Akt-CA overexpression muscles, but also appeared
to be responsive to rapamycin treatment as we observed elevated levels of UBR1 in the
EV control plus rapamycin condition (Figure 5B). To further assess the UPS, we

measured total ubiquitinated proteins, K48-specific linkage levels and proteasome
activity for 20S and 26S subunits (Figure 5A, 5C and 5D). Independent of diet condition,
we observed significantly increased levels of total ubiquitinated proteins and K48-
specific linkage in Akt-CA overexpression muscles compared to the EV control (Figure
5C). For the 20S subunit proteasomal activity, we observed little change in activity with
Akt-CA overexpression, except for 20S β1 activity under the rapamycin diet background
(Figure 5D). In comparison, 26S proteasomal activity displayed significant elevations in
β1 and β2 subunit activity in Akt-CA overexpression muscles independent of the diet
condition (Figure 5D). For the 26S β5 subunit, we observed a near significant increase
in activity with the Akt-CA plus rapamycin condition compared to the control diet
background (Figure 5D; $P = 0.051$). Altogether, these data highlight changes in UBR-
box E3 ubiquitin ligases, ubiquitination, and proteasomal activity that occur in a
condition of low mTORC1 activity.

Markers for autophagy and ER-stress are elevated in Akt transfected muscles with
suppressed mTORC1 activity
Given the reported association between the UBR-box E3 ubiquitin ligases and their
involvement in protein quality control (27, 30, 31), we sought to assess changes in
markers related to autophagy and ER-stress pathways in Akt-CA overexpression TA
muscles. For VCP, LC3B I and LC3B II, we observed significant increased protein levels
in Akt-CA overexpression muscles that were independent of diet background (Figure 6A
and 6B). Interestingly for p62 protein abundance we observed significant increases in
the control diet condition only with Akt-CA overexpression (Figure 6B). However, similar
to UBR1, we observed a rapamycin-induced increase in basal p62 protein levels in the
EV control muscles (Figure 6B). For markers related to the ER stress response, there
was no change in protein levels for BiP (Figure 6C and 6D). However, for PDI, CHOP
and eIF2α protein levels, there was a significant increase in Akt-CA transfected muscles
that was present independent of the diet conditions (Figure 6D). Lastly, similar to Akt
transgenic and TSC1/2 knockout rodent models (19), we observed the presence of
'aberrant' muscle fibers containing multiple vacuole-like structures in the Akt-CA
overexpression muscles, which were also present in the rapamycin diet condition

(Figure 6E and 6F). Overall, these data highlight the response of protein quality control pathways to Akt induction, and that these pathways may be responding to protein clearance and removal of damaged/misfolded proteins that are driven by hyperactive protein synthesis, leading to potential increased errors of translation.

UBR5 knockdown results in loss of muscle mass and fiber size under control and rapamycin conditions

We have previously observed UBR5 knockdown to result in loss of muscle mass and fiber size which was accompanied by mTORC1 hyperactivation, as measured by phosphorylation of downstream substrates p70S6K and rpS6 at multiple time points (43). However, it remained to be determined if the response of mTORC1 activity was adaptive or maladaptive and could be contributing to the development of skeletal muscle atrophy with UBR5 knockdown. Therefore, we sought to utilize the rapamycin diet in conjunction with UBR5 knockdown for 30 days post electroporation (Figure 1C). In the control diet group, we observed significant loss of skeletal muscle mass and fiber size, as measured by a reduction in muscle weight and a shift toward smaller CSA size for GFP-positive fibers for the UBR5 RNAi construct compared to the contralateral EV muscle (Figure 7A-C). These observations confirmed our previous findings for the effect of UBR5 knockdown on skeletal muscle mass size (43). Interestingly, we also observed significant reductions in muscle mass and fiber CSA size in UBR5 RNAi transfected muscles compared to the contralateral EV in the presence of rapamycin (Figure 7B and 7C).

At the biochemical level, similar to our previously published data (43), we observed increased phosphorylation of Akt, p70S6K1, rpS6 and p90RSK in UBR5 RNAi transfected muscles in the control diet condition compared to the EV muscles (Figure 7D). The phosphorylation of p70S6K at Thr389 was suppressed in the EV and UBR5 RNAi transfected muscles confirming the inhibitory effect of the rapamycin diet on mTORC1 activity (Figure 7D and 7E). Interestingly, rapamycin did not inhibit the UBR5 RNAi-induced phosphorylation of Akt, rpS6 and p90RSK sites compared to the EV transfected muscle (Figure 7D and 7E). The dissociation between p70S6K1 and rpS6 phosphorylation is interesting and might suggest the potential involvement of either

another rpS6 kinase (e.g. PKA) or the inhibition of a rpS6 phosphatase (65-67). This
observation warrants further investigation in the UBR5 knockdown model. Lastly, we
wanted to observe how the protein abundance for UBR1, UBR2, UBR4, and UBR7
might change with UBR5 Knockdown. In the control diet, we observed significant
increases in total ubiquitin, UBR2, UBR4, and UBR7 when UBR5 protein was
suppressed 50-60% (Figure 7F and 7G) In the rapamycin diet condition, basal UBR1
and UBR7 proteins were significantly increased in EV transfected muscles compared to
control diet EV transfected muscles, while only UBR2 remained significantly elevated in
the UBR5 RNAi tissues compared to the EV transfected muscle (Figure 7F and 7G).

Discussion
Skeletal muscle can regulate its size in response to internal and external cues
throughout the life span (2, 3, 68). Homeostasis of skeletal muscle size occurs through
a balance between protein synthesis and breakdown (2). In the present study, we
sought to further our understanding of skeletal muscle proteostasis under growth and
remodeling conditions by using a short term Akt overexpression model, in conjunction
with a rapamycin diet intervention. Our results highlight a novel group of E3 ubiquitin
ligases (UBR1, UBR2, UBR4, UBR5, and UBR7) that appear to be responsive under
conditions of heightened protein synthesis and when other protein quality control
pathways are active. In addition, the change in protein abundance for UBR-box E3
ubiquitin ligases was still evident with rapamycin treatment, suggesting that the UBR-
box E3 ubiquitin ligases might act in a rapamycin-insensitive manner and maybe
dependent on upstream Akt signals and mTORC1-independent control. Indeed, we
observed the activation of p44/p42 MAPK, p90RSK and mTORC2 signaling that can
contribute to protein synthesis and skeletal muscle growth (69-73). Future studies are
required to ascertain a physiological role of UBR-box E3 ubiquitin ligases with Akt-
mediated protein synthesis and turnover in the context of skeletal muscle growth and
remodeling.

As the UBR-box E3 ubiquitin ligases are part of the N-degron pathway, they contain a
UBR-domain which can recognize destabilized proteins and protein fragments for
protein degradation (25, 27). The N-degron pathway serves as an important protein

quality control mechanism (25, 31) and limited studies have been performed in skeletal

muscle exploring this pathway. However, recent observations from us and other

laboratories have implicated UBR4, UBR5, and UBR7 in the regulation of the skeletal

muscle proteome under growth and exercise stimuli (35-39). For instance, UBR5 and

UBR4 have each been observed to be exercise responsive E3 ubiquitin ligases in

human and mouse skeletal muscle under resistance and aerobic exercise models (35,

36, 38), with a change in phosphorylation status for UBR5 noted in multiple different

exercise modalities (38). Further, UBR5 has been observed to increase in other models

of mechanical loading (42) and UBR5 activity may be rapamycin insensitive (74) which

compliments data collected in the current study in our Akt overexpression and UBR5

knockdown models. Our understanding of the transcriptional regulation of UBR5 and

other UBR-related E3 ligases in skeletal muscle is limited. Previous literature has

highlighted UBR5 DNA methylation status to coincide with changes in gene expression

in human skeletal muscle under exercise training (35). Future studies are warranted to

understand the transcriptional regulation of the UBR E3 ubiquitin ligases under different

cellular conditions. To note, UBR5 has been reported to interact with p90RSK signaling

via p90RSK phosphorylation sites on UBR5 (75), and we saw p90RSK activation with

UBR5 suppression or increased UBR5 protein abundance in the presence of p90RSK

phosphorylation, highlighting the potential responsiveness of UBR5 and UBR-box E3

ubiquitin ligases to changes in the protein synthetic machinery. These observations

extend our understanding of E3 ubiquitin ligases in protein quality control and the

regulation of skeletal muscle size.

Although very little is known about the substrates for the UBR-box E3 ubiquitin ligases,

we observed an increase in K48-linked ubiquitinated proteins, total ubiquitinated protein

levels and 26S proteasome activity under control and rapamycin diet conditions in the

Akt overexpression model. Recent evidence by Kaiser and colleagues (19) has alluded

to the proteasome system being controlled by mTORC1 via feedback inhibition of

PKB/Akt. The authors utilized transgenic Akt-CA mice and TSC1 knockout models to

manipulate mTORC1 activation and observed changes in proteasome biogenesis and

activity as well as atrophy-related gene expression for specific E3 ubiquitin ligases in

these models. It could be postulated that the increase in mTORC1-mediated translation

results in a buildup of error containing peptides and misfolded proteins, which
inadvertently increases the demand on the UPS for targeted protein degradation. These
ideas warrant further investigation, and are highlighted in the literature drawing on
observations reported using other cell and tissue models (76). In our current study, Akt
overexpression and subsequent mTORC1 activation was induced for a shorter period
of time compared to previous studies (15, 19, 20), and notably, we observed UPS-
induction and markers of other protein quality control pathways (e.g. ER stress and
autophagy) to be elevated under conditions where mTORC1 activity is low (e.g.
rapamycin diet). Given, that it has previously been proposed that mTORC1 may control
the UPS system and proteasome activity in skeletal muscle (19), the current study's
findings extend our observations for the UPS and protein quality control system being
controlled under rapamycin-sensitive and insensitive signals. It is important to note that
the chronic use of rapamycin can inhibit mTORC1 and mTORC2 (45). However, we
observed increased phosphorylation of p90RSK and NDRG1 (mTORC2 substrate) with
Akt overexpression and thus the activation of alternative protein synthetic pathways
may require a response from a variety of protein quality control mechanisms such as
the N-degron pathway in order to maintain proteome integrity.

Recent studies have alluded to the importance of the N-degron pathway being critical
for protein quality control via its ability to recognize destabilized proteins that display a
degron signal (31-33). Interestingly, with UBR5 knockdown, we observed an increase in
protein abundance for UBR2, UBR4 and UBR7 which may highlight a compensatory
response in order to maintain proteostasis due to increased protein synthesis in UBR5
KD transfected muscles. The response in UBR E3 ubiquitin ligases to skeletal muscle
regrowth (21d to 60d) is notable given that all the UBR E3s measured in the current
study were elevated in protein abundance as the tissue recovered following a period of
inactivity. We recently hypothesized that the UBR E3 ubiquitin ligases work as a protein
surveillance system under growth conditions in order to maintain proteome integrity
during production of newly synthesized proteins (39). The observations from the current
study are interesting because UBR-box E3 ligases all share the UBR-domain, but the
mechanisms for ubiquitin transfer differs between RING, F-box and Hect E3 ubiquitin
ligases which are all comprised in the UBR class of E3 ligases (24, 29, 77). The

substrates for the UBR-box E3 ligases remain to be identified in vivo. Previous literature

has suggested that different UBRs may elicit their function for ubiquitination under

different cellular compartments (27, 31, 78). For instance, UBR1 and UBR2 are reported

to target misfolded proteins for degradation in the cytosolic compartment (31) whereas

the action of UBR5 might be dominant in the nucleus and regulating nuclear receptors

and transcription factors (33, 79-81). However, these previously published observations

remain to be confirmed in vivo and in the context of skeletal muscle plasticity.

Limitations

The data reported within the current study is not without limitations. The use of in vivo

electroporation allows for a targeted skeletal muscle approach to understand the

molecular mechanisms of hypertrophy and atrophy (50). However, transfection

efficiency can play a role in observations made at the biochemical level, as a "dilution"

effect can occur whereby changes in gene/protein expression are diluted by a large

proportion of non-transfected fibers, leading to an underestimation of the effect of the

gene of interest (50). Our electroporation protocol has been optimized to maximize

transfection efficiency whilst also limiting the level of potential muscle damage that

might be observed (50, 51). Our current observations for the UBR-box E3 ligases in

skeletal muscle remodeling are limited to only the tibialis anterior muscle, although we

have performed similar electroporation experiments in the gastrocnemius muscle with

the UBR5 RNAi construct and observed similar levels of muscle mass and fiber CSA

decreases with UBR5 suppression (data not shown). In the current study, we observed

changes in UBR-box E3 ligase protein abundance during the recovery of muscle mass

following atrophy in the gastrocnemius complex muscles (Fig. 3), which might imply that

their role is similar in all hindlimb skeletal muscles, but future studies are warranted. It

should be noted that the rapamycin dose used in the current study is not without

potential side effects at the whole body or physiological level with alterations in body

weight and glucose metabolism being reported (82, 83). However, in the current study,

we utilized the rapamycin diet intervention for a shorter period of time, compared to

studies focused on rapamycin as an intervention for lifespan and healthspan (82, 83).

Lastly, protein turnover and quality control are dynamic processes for maintaining

proteostasis (76, 84, 85) and thus caution must be applied given that a single snapshot

in time was measured for changes in protein quality control mechanisms when protein
synthesis was genetically enhanced to induce skeletal muscle growth.

Conclusions

The present study highlights a novel group of E3 ubiquitin ligases that respond to
heightened protein synthesis under skeletal muscle growth or atrophy conditions via
genetic manipulation or recovery following nerve crush injury. Little is known about the
N-degron pathway in tissue health, and our initial observations provide important steps
towards understanding this pathway. Future studies are warranted to identify protein
substrates for these E3 ubiquitin ligases and explore their potential role in other tissues
(e.g. heart, brain) given that they are ubiquitous E3 ligases. Overall, the UBR-box E3
ubiquitin ligases could be critical in the adaptive response to cellular stress where
perturbations in proteostasis and tissue remodeling might occur in the context of aging,
disease, and exercise.

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Author Contributions

Conception and design of the experiments: SCB and DCH. Collection, analysis and interpretation of data: LMB, LGO, CAG, APS, DSW, SCB and DCH. Drafting the article or revising it critically for important intellectual content: LMB, LGO, CAG, APS, DSW, SCB and DCH. All authors read and approved the final version of the manuscript, and all authors listed qualify for authorship.

Conflict of Interest

SCB is on the scientific advisory board for Emmyon Inc. All other authors declare that they do not have a conflict of interest.

Funding

D.C. Hughes was supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under Award Numbers K01AR077684 and R03AR083980. Additional support was provided by a pilot project award to D.C. Hughes from the CoBRE Center for Cellular Metabolism Research (NIGMS P20GM139763).

Figure Legends

Figure 1. Overview of the experimental designs for studies using in vivo electroporation
of the tibialis anterior (TA) muscle with dietary manipulation. Mice were electroporated with
an empty vector control and either the wild type (Akt-WT) or constitutively active (Akt-CA) Akt
constructs for 7 days (**A**). Next, cohorts of mice were randomly assigned to a control or
rapamycin diet for 7 days. The mice were then electroporated with EV and Akt-CA constructs
and collected after 7 days. Total time on the dietary intervention was 14 days (**B**). Lastly,
cohorts of mice were electroporated with EV and UBR5 RNAi constructs and then placed on a
control or rapamycin diet for 30 days (**C**). All studies utilized adult male mice (4-5 months of
age). Figure was created with Biorender.com.

Figure 2. Differing degrees of Akt activity result in skeletal muscle growth and activation
of mTORC1 signaling, translation and increased UBR-box E3 ubiquitin ligase abundance.
Muscle mass (**A**) measurements (milligrams) for muscles transfected with either the empty
vector (EV) or Akt-WT, and Akt-CA plasmids after 7 days (n=5-10/group). Representative
microscope images (**B**) for green fluorescent protein (GFP) and laminin (red) in TA muscles
transfected with EV or Akt-CA constructs (x10 magnification; scale bar = 100µm), highlighting
transfection efficiency and muscle morphology after 7 days. Muscle cross-sectional area (CSA)
distribution for EV and Akt-CA transfected muscles. GFP-positive fibers were measured for
CSA, with ≥ 450 transfected fibers analyzed per animal, per muscle (n=5/group). Total number
of fibers analyzed per group are reported in parentheses. For CSA data, fibers presented as
percentage of fibers between 0 and 4800 µm plus size. Representative immunoblot images (**C**)
for Akt/mTORC1 signaling and puromycin. Representative immunoblot images (**D**) for UBR-box
E3 ubiquitin ligases and total ubiquitin. Quantification of Akt/mTORC1 signaling, puromycin-
labelling, UBR-box E3 ubiquitin ligases (UBR1, UBR2, UBR4, UBR5 and UBR7) and total
ubiquitin (**E**) in EV, Akt-WT, and Akt-CA transfected muscles (n=5/group). Total protein loading
was used as the normalization control for all blots. In EV, Akt-WT, and Akt-CA transfected
muscles, mRNA expression (**F**) was determined via quantitative PCR for UBR1, UBR2, UBR4,
UBR5, UBR7, MuRF1 (Trim63), and MAFbx (FBXO32). N=4-5/group. Data presented as means
± SEM. *$P<0.05$; **$P<0.01$.

Figure 3. Response of UBR-box E3 ligases to skeletal muscle regrowth following a period
of neural inactivity. Representative immunoblot images (**A**) for UBR-box E3 ubiquitin ligases
post nerve crush injury in gastrocnemius complex (GSTC) skeletal muscle. We have previously

published the UBR5 protein response in the GSTC muscle following nerve crush injury (NCI)
and the 21 to 60-day time period corresponds to the recovery (regrowth) of skeletal muscle
mass following a period of disuse. (**B**) Quantification of UBR-box E3 ubiquitin ligases (UBR1,
UBR2, UBR4, UBR5, and UBR7) in GSTC muscle from control and time points (21, 28 ,45, and
60 days post NCI) corresponding to skeletal muscle regrowth (n=4-5/group). Data presented as
means ± SEM. *$P<0.05$, **$P<0.01$ vs control.

Figure 4. Effect of rapamycin on Akt-induced skeletal muscle growth, mTORC1
dependent and independent signaling. Male mice were provided either a control or
rapamycin diet seven days prior to the electroporation procedure and kept on the respective diet
for the remainder of the experiment. Muscle mass (**A**) measurements (milligrams) for muscles
transfected with either the empty vector (EV) or Akt-CA plasmids for the control and rapamycin
interventions after 7 days (n=10/group; left panel). The percentage difference in muscle mass
compared to the EV transfected muscle for the control and rapamycin diets (A; right panel).
Representative microscope images (**B**) for green fluorescent protein (GFP) and Laminin (red) in
TA muscles transfected with EV or Akt-CA plasmid on the control or rapamycin diet (x10
magnification; scale bar = 100µm), highlighting transfection efficiency and muscle morphology
after 7 days. Muscle cross-sectional area (CSA) distribution for EV and Akt-CA transfected
muscles on the control or rapamycin diet (**C**). GFP-positive fibers were measured for CSA, with
≥ 450 transfected fibers analyzed per animal, per muscle (n=5/group). Total number of fibers
analyzed per group are reported in parentheses. For CSA data, fibers presented as percentage
of fibers between 0 and 4800 µm plus size. Representative immunoblot images (**D**) for
Akt/mTORC1 signaling and puromycin from EV and Akt-CA plasmids with control or rapamycin
diet (n=5/group). Quantification of puromycin and phosphorylation status of rpS6, NDRG1 and
p90RSK in control and rapamycin conditions with Akt overexpression (**E**). Data presented as
means ± SEM. *$P<0.05$; **$P<0.01$.

Figure 5. Increase in UBR-box E3 ligase abundance and proteasome activity in Akt
transfected skeletal muscles under suppressed mTORC1 activity. Representative
immunoblot images (**A**) for UBR-box E3 ubiquitin ligases, total ubiquitin, and K48-specific
ubiquitin linkage. Quantification of UBR-box E3 ubiquitin ligases (UBR1, UBR2, UBR4, UBR5,
and UBR7) (**B**), total ubiquitin, and K48 specific ubiquitin (**C**) in empty vector (EV) and Akt-CA
transfected muscles under control or rapamycin diet conditions (n=5/group). Total protein
loading was used as the normalization control for all blots. Proteasome activity assays were

performed for 20S and 26S subunits (β1, β2, and β5) in transfected muscles (n=4/group; **D**).

Data presented as means ± SEM. *P<0.05; **P<0.01.

Figure 6. Changes in autophagy-related and ER stress-related proteins with Akt

transfected skeletal muscle independent of mTORC1 activity. Representative immunoblot

images (**A**) for autophagy-related markers, VCP, p62, and LC3B. Quantification of autophagy-

related markers (**B**) in empty vector (EV) and Akt-CA transfected muscles under control or

rapamycin diet conditions (n=5/group). Representative immunoblot images (**C**) for ER stress-

related markers, BiP, PDI, CHOP, and eIF2α. Quantification of ER stress-related markers (**D**) in

EV and Akt-CA transfected muscles under control or rapamycin diet conditions (n=5/group).

Total protein loading was used as the normalization control for all blots. Representative

microscope images for hematoxylin-eosin (H&E) in TA mouse muscles transfected with EV and

Akt-CA plasmids under control and rapamycin diet conditions (×20 magnification; scale

bar□=□50 μm) highlighting muscle morphology after 7 days post electroporation (**E**). Examples

of aberrant skeletal muscle fibers are identified on H&E images with yellow arrows and were

quantified in EV and Akt-CA transfected muscles under control and rapamycin conditions

(*n*=5/group) (**F**). Data presented as means ± SEM. *P<0.05; **P<0.01.

Figure 7. Loss of skeletal muscle mass and fiber size with UBR5 knockdown occurs

independent of mTORC1 activity. Representative microscope images (**A**) for green

fluorescent protein (GFP) and Laminin (red) in TA muscles transfected with EV or UBR5 RNAi

plasmid on the control or rapamycin diet (x10 magnification; scale bar = 100μm), highlighting

transfection efficiency and muscle morphology after 30 days. TA muscle mass (**B**)

measurements (milligrams) for muscles transfected with either the EV or UBR5 RNAi plasmids

for the control and rapamycin interventions after 30 days (n=9-10/group). Mean GFP-positive

fiber size (**B**) was quantified for TA muscles transfected with EV or UBR5 RNAi under control or

rapamycin diets (n=3-4/group). Muscle fiber CSA distribution for EV and UBR5 RNAi

transfected muscles (**C**). GFP-positive fibers were measured for CSA, with ≥ 450 transfected

fibers analyzed per animal, per muscle (n=3-4/group). Total number of fibers analyzed per

group are reported in parentheses and represent Fiber CSA data for TA transfected muscles.

For CSA data, fibers presented as percentage of fibers between 0 and 4800 μm. Representative

immunoblot images (**D**) for Akt/mTORC1 signaling and puromycin from EV and UBR5 RNAi

plasmids with control or rapamycin diet (n=5-6/group). Quantification of puromycin and

phosphorylation status of Akt, p70S6K1, rpS6, and p90RSK in control and rapamycin conditions

with UBR5 knockdown (**E**). Representative immunoblot images (**F**) for UBR-box E3 ubiquitin

ligases. Quantification of UBR-box E3 ubiquitin ligases (UBR1, UBR2, UBR4, UBR5, and
UBR7) and total ubiquitin (**G**) in EV and UBR5 RNAi transfected muscles (n=5-6/group) under
control and rapamycin diet conditions. $*P<0.05$ vs. EV. Total protein loading was used as the
normalization control for all blots. Data presented as means ± SEM. $*P<0.05$; $**P<0.01$.

References

1.   **Hughes DC, Ellefsen S, and Baar K**. Adaptations to endurance and strength training. *Cold Spring Harbor perspectives in medicine* a029769, 2017.

2.   **Baehr LM, Hughes DC, Waddell DS, and Bodine SC**. SnapShot: Skeletal muscle atrophy. *Cell* 185: 1618-1618. e1611, 2022.

3.   **Sharples AP, Hughes DC, Deane CS, Saini A, Selman C, and Stewart CE**. Longevity and skeletal muscle mass: the role of IGF signalling, the sirtuins, dietary restriction and protein intake. *Aging cell* 14: 511-523, 2015.

4.   **Bodine SC, Stitt TN, Gonzalez M, Kline WO, Stover GL, Bauerlein R, Zlotchenko E, Scrimgeour A, Lawrence JC, and Glass DJ**. Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nature cell biology* 3: 1014-1019, 2001.

5.   **Baar K, and Esser K**. Phosphorylation of p70S6k correlates with increased skeletal muscle mass following resistance exercise. *American Journal of Physiology-Cell Physiology* 276: C120-C127, 1999.

6.   **West DW, Baehr LM, Marcotte GR, Chason CM, Tolento L, Gomes AV, Bodine SC, and Baar K**. Acute resistance exercise activates rapamycin-sensitive and-insensitive mechanisms that control translational activity and capacity in skeletal muscle. *The Journal of physiology* 594: 453-468, 2016.

7.   **Lee Hamilton D, Philp A, MacKenzie MG, Patton A, Towler MC, Gallagher IJ, Bodine SC, and Baar K**. Molecular brakes regulating mTORC1 activation in skeletal muscle following synergist ablation. *American Journal of Physiology-Endocrinology and Metabolism* 307: E365-E373, 2014.

8.   **You J-S, Dooley MS, Kim C-R, Kim E-J, Xu W, Goodman CA, and Hornberger TA**. A DGKζ-FoxO-ubiquitin proteolytic axis controls fiber size during skeletal muscle remodeling. *Science signaling* 11: 2018.

9.   **Frey JW, Jacobs BL, Goodman CA, and Hornberger TA**. A role for Raptor phosphorylation in the mechanical activation of mTOR signaling. *Cellular Signalling* 26: 313-322, 2014.

10.   **Goodman CA**. Role of mTORC1 in mechanically induced increases in translation and skeletal muscle mass. *Journal of applied physiology* 127: 581-590, 2019.

11.   **Kim H-G, Guo B, and Nader GA**. Regulation of ribosome biogenesis during skeletal muscle hypertrophy. *Exercise and sport sciences reviews* 47: 91-97, 2019.

12.   **Figueiredo VC, and McCarthy JJ**. Regulation of ribosome biogenesis in skeletal muscle hypertrophy. *Physiology* 34: 30-42, 2019.

13.   **Wallace MA, Hughes DC, and Baar K**. mTORC1 in the Control of Myogenesis and Adult Skeletal Muscle Mass. In: *Molecules to Medicine with mTOR* Elsevier, 2016, p. 37-56.

14.   **Castets P, Rion N, Théodore M, Falcetta D, Lin S, Reischl M, Wild F, Guérard L, Eickhorst C, Brockhoff M, Guridi M, Ibebunjo C, Cruz J, Sinnreich M, Rudolf R, Glass DJ, and Rüegg MA**. mTORC1 and PKB/Akt control the muscle response to denervation by regulating autophagy and HDAC4. *Nat Commun* 10: 3187, 2019.

15.   **Marabita M, Baraldo M, Solagna F, Ceelen Judith Johanna M, Sartori R, Nolte H, Nemazanyy I, Pyronnet S, Kruger M, Pende M, and Blaauw B**. S6K1 Is Required for Increasing Skeletal Muscle Force during Hypertrophy. *Cell Reports* 17: 501-513, 2016.

16.   **Bentzinger CF, Lin S, Romanino K, Castets P, Guridi M, Summermatter S, Handschin C, Tintignac LA, Hall MN, and Rüegg MA**. Differential response of skeletal muscles to mTORC1 signaling during atrophy and hypertrophy. *Skeletal muscle* 3: 1-16, 2013.

17.   **Tang H, Inoki K, Brooks SV, Okazawa H, Lee M, Wang J, Kim M, Kennedy CL, Macpherson PC, and Ji X**. mTORC1 underlies age-related muscle fiber damage and loss by inducing oxidative stress and catabolism. *Aging cell* 18: e12943, 2019.

18. **Lai K-MV, Gonzalez M, Poueymirou WT, Kline WO, Na E, Zlotchenko E, Stitt TN, Economides**
AN, Yancopoulos GD, and Glass DJ. Conditional activation of akt in adult skeletal muscle induces rapid
hypertrophy. *Molecular and cellular biology* 24: 9295-9304, 2004.
19. **Kaiser MS, Milan G, Ham DJ, Lin S, Oliveri F, Chojnowska K, Tintignac LA, Mittal N, Zimmerli CE,**
and Glass DJ. Dual roles of mTORC1-dependent activation of the ubiquitin-proteasome system in muscle
proteostasis. *Communications biology* 5: 1141, 2022.
20. **Blaauw B, Canato M, Agatea L, Toniolo L, Mammucari C, Masiero E, Abraham R, Sandri M,**
Schiaffino S, and Reggiani C. Inducible activation of Akt increases skeletal muscle mass and force
without satellite cell activation. *The FASEB journal* 23: 3896-3905, 2009.
21. **Castets P, and Rüegg MA**. MTORC1 determines autophagy through ULK1 regulation in skeletal
muscle. *Autophagy* 9: 1435-1437, 2013.
22. **Bodine SC, Latres E, Baumhueter S, Lai VK-M, Nunez L, Clarke BA, Poueymirou WT, Panaro FJ,**
Na E, and Dharmarajan K. Identification of ubiquitin ligases required for skeletal muscle atrophy.
Science 294: 1704-1708, 2001.
23. **Milan G, Romanello V, Pescatore F, Armani A, Paik J-H, Frasson L, Seydel A, Zhao J, Abraham R,**
and Goldberg AL. Regulation of autophagy and the ubiquitin–proteasome system by the FoxO
transcriptional network during muscle atrophy. *Nature communications* 6: 6670, 2015.
24. **Hughes DC, Goodman CA, Baehr LM, Gregorevic P, and Bodine SC**. A critical discussion on the
relationship between E3 ubiquitin ligases, protein degradation, and skeletal muscle wasting: it's not that
simple. *American Journal of Physiology-Cell Physiology* 325: C1567-C1582, 2023.
25. **Varshavsky A**. N-degron and C-degron pathways of protein degradation. *Proceedings of the*
National Academy of Sciences 116: 358-366, 2019.
26. **Varshavsky A**. The N-end rule pathway and regulation by proteolysis. *Protein science* 20: 1298-
1345, 2011.
27. **Sherpa D, Chrustowicz J, and Schulman BA**. How the ends signal the end: Regulation by E3
ubiquitin ligases recognizing protein termini. *Molecular Cell* 82: 1424-1438, 2022.
28. **Choi WS, Jeong B-C, Joo YJ, Lee M-R, Kim J, Eck MJ, and Song HK**. Structural basis for the
recognition of N-end rule substrates by the UBR box of ubiquitin ligases. *Nature structural & molecular*
biology 17: 1175-1181, 2010.
29. **Tasaki T, Mulder LC, Iwamatsu A, Lee MJ, Davydov IV, Varshavsky A, Muesing M, and Kwon**
YT. A family of mammalian E3 ubiquitin ligases that contain the UBR box motif and recognize N-degrons.
Molecular and cellular biology 2005.
30. **Hunt LC, Schadeberg B, Stover J, Haugen B, Pagala V, Wang Y-D, Puglise J, Barton ER, Peng J,**
and Demontis F. Antagonistic control of myofiber size and muscle protein quality control by the
ubiquitin ligase UBR4 during aging. *Nature Communications* 12: 1418, 2021.
31. **Nillegoda NB, Theodoraki MA, Mandal AK, Mayo KJ, Ren HY, Sultana R, Wu K, Johnson J, Cyr**
DM, and Caplan AJ. Ubr1 and Ubr2 function in a quality control pathway for degradation of unfolded
cytosolic proteins. *Molecular biology of the cell* 21: 2102-2116, 2010.
32. **Shimshon A, Dahan K, Israel-Gueta M, Olmayev-Yaakobov D, Timms RT, Bekturova A, Makaros**
Y, Elledge SJ, and Koren I. Dipeptidyl peptidases and E3 ligases of N-degron pathways cooperate to
regulate protein stability. *Journal of Cell Biology* 223: e202311035, 2024.
33. **Tsai JM, Aguirre JD, Li Y-D, Brown J, Focht V, Kater L, Kempf G, Sandoval B, Schmitt S, and**
Rutter JC. UBR5 forms ligand-dependent complexes on chromatin to regulate nuclear hormone receptor
stability. *Molecular cell* 83: 2753-2767. e2710, 2023.
34. **Solomon V, Lecker SH, and Goldberg AL**. The N-end rule pathway catalyzes a major fraction of
the protein degradation in skeletal muscle. *Journal of Biological Chemistry* 273: 25216-25222, 1998.

35.   **Seaborne RA, Strauss J, Cocks M, Shepherd S, O'Brien TD, van Someren KA, Bell PG,**
Murgatroyd C, Morton JP, Stewart CE, and Sharples AP. Human Skeletal Muscle Possesses an
Epigenetic Memory of Hypertrophy. *Sci Rep* 8: 1898, 2018.
36.   **Chambers TL, Dimet-Wiley A, Keeble AR, Haghani A, Lo WJ, Kang G, Brooke R, Horvath S, Fry**
CS, and Watowich SJ. Methylome–proteome integration after late-life voluntary exercise training
reveals regulation and target information for improved skeletal muscle health. *The Journal of Physiology*
603: 211-237, 2025.
37.   **Roberts MD, Ruple BA, Godwin JS, McIntosh MC, Chen S-Y, Kontos NJ, Agyin-Birikorang A,**
Michel M, Plotkin DL, and Mattingly ML. A novel deep proteomic approach in human skeletal muscle
unveils distinct molecular signatures affected by aging and resistance training. *Aging (Albany NY)* 16:
6631, 2024.
38.   **Blazev R, Carl CS, Ng Y-K, Molendijk J, Voldstedlund CT, Zhao Y, Xiao D, Kueh AJ, Miotto PM,**
and Haynes VR. Phosphoproteomics of three exercise modalities identifies canonical signaling and
C18ORF25 as an AMPK substrate regulating skeletal muscle function. *Cell Metabolism* 34: 1561-1577.
e1569, 2022.
39.   **Hughes DC, and Bodine SC**. UBR5: A New Player in Protein Quality Control for Skeletal Muscle
Growth and Remodeling. *Exercise and Sport Sciences Reviews* 53: 205-213, 2025.
40.   **Gao S, Zhang G, Zhang Z, Zhu JZ, Li L, Zhou Y, Rodney GG, Jr., Abo-Zahrah RS, Anderson L,**
Garcia JM, Kwon YT, and Li YP. UBR2 targets myosin heavy chain IIb and IIx for degradation: Molecular
mechanism essential for cancer-induced muscle wasting. *Proc Natl Acad Sci U S A* 119: e2200215119,
2022.
41.   **Kwak KS, Zhou X, Solomon V, Baracos VE, Davis J, Bannon AW, Boyle WJ, Lacey DL, and Han**
HQ. Regulation of protein catabolism by muscle-specific and cytokine-inducible ubiquitin ligase E3alpha-
II during cancer cachexia. *Cancer Res* 64: 8193-8198, 2004.
42.   **Seaborne RA, Hughes DC, Turner DC, Owens DJ, Baehr LM, Gorski P, Semenova EA, Borisov**
OV, Larin AK, Popov DV, Generozov EV, Sutherland H, Ahmetov II, Jarvis JC, Bodine SC, and Sharples
AP. UBR5 is a novel E3 ubiquitin ligase involved in skeletal muscle hypertrophy and recovery from
atrophy. *The Journal of Physiology* 597: 3727-3749, 2019.
43.   **Hughes DC, Turner DC, Baehr LM, Seaborne RA, Viggars M, Jarvis JC, Gorski PP, Stewart CE,**
Owens DJ, Bodine SC, and Sharples AP. Knockdown of the E3 Ubiquitin ligase UBR5 and its role in
skeletal muscle anabolism. *American Journal of Physiology-Cell Physiology* 320: C45-C56, 2021.
44.   **Hunt LC, Stover J, Haugen B, Shaw TI, Li Y, Pagala VR, Finkelstein D, Barton ER, Fan Y, and**
Labelle M. A key role for the ubiquitin ligase UBR4 in myofiber hypertrophy in Drosophila and mice. *Cell*
reports 28: 1268-1281. e1266, 2019.
45.   **Sarbassov DD, Ali SM, Sengupta S, Sheen JH, Hsu PP, Bagley AF, Markhard AL, and Sabatini**
DM. Prolonged rapamycin treatment inhibits mTORC2 assembly and Akt/PKB. *Mol Cell* 22: 159-168,
2006.
46.   **Ham DJ, Börsch A, Chojnowska K, Lin S, Leuchtman AB, Ham AS, Thürkauf M, Delezie J, Furrer**
R, Burri D, Sinnreich M, Handschin C, Tintignac LA, Zavolan M, Mittal N, and Rüegg MA. Distinct and
additive effects of calorie restriction and rapamycin in aging skeletal muscle. *Nature Communications*
13: 2025, 2022.
47.   **Ham DJ, Börsch A, Lin S, Thürkauf M, Weihrauch M, Reinhard JR, Delezie J, Battilana F, Wang**
X, and Kaiser MS. The neuromuscular junction is a focal point of mTORC1 signaling in sarcopenia. *Nature*
Communications 11: 1-21, 2020.
48.   **Chiao YA, Kolwicz SC, Basisty N, Gagnidze A, Zhang J, Gu H, Djukovic D, Beyer RP, Raftery D,**
and MacCoss M. Rapamycin transiently induces mitochondrial remodeling to reprogram energy
metabolism in old hearts. *Aging (Albany NY)* 8: 314, 2016.

49.   **Ebert SM, Dyle MC, Kunkel SD, Bullard SA, Bongers KS, Fox DK, Dierdorff JM, Foster ED, and**
Adams CM. Stress-induced skeletal muscle Gadd45a expression reprograms myonuclei and causes
muscle atrophy. *J Biol Chem* 287: 27290-27301, 2012.
50.   **Hughes DC, Hardee JP, Waddell DS, and Goodman CA**. CORP: Gene delivery into murine skeletal
muscle using in vivo electroporation. *Journal of Applied Physiology* 133: 41-59, 2022.
51.   **Hughes DC, Baehr LM, Driscoll JR, Lynch SA, Waddell DS, and Bodine SC**. Identification and
characterization of Fbxl22, a novel skeletal muscle atrophy-promoting E3 ubiquitin ligase. *Am J Physiol*
Cell Physiol 319: C700-c719, 2020.
52.   **Baehr LM, Hughes DC, Lynch SA, Van Haver D, Maia TM, Marshall AG, Radoshevich L, Impens**
F, Waddell DS, and Bodine SC. Identification of the MuRF1 skeletal muscle ubiquitylome through
quantitative proteomics. *Function* 2: zqab029, 2021.
53.   **Wen Y, Murach KA, Vechetti IJ, Jr., Fry CS, Vickery C, Peterson CA, McCarthy JJ, and Campbell**
KS. MyoVision: software for automated high-content analysis of skeletal muscle immunohistochemistry.
J Appl Physiol (1985) 124: 40-51, 2018.
54.   **Viggars MR, Wen Y, Peterson CA, and Jarvis JC**. Automated cross-sectional analysis of trained,
severely atrophied, and recovering rat skeletal muscles using MyoVision 2.0. *J Appl Physiol (1985)* 132:
593-610, 2022.
55.   **Parlee SD, Lentz SI, Mori H, and MacDougald OA**. Quantifying size and number of adipocytes in
adipose tissue. *Methods Enzymol* 537: 93-122, 2014.
56.   **Castets P, Lin S, Rion N, Di Fulvio S, Romanino K, Guridi M, Frank S, Tintignac LA, Sinnreich M,**
and Rüegg MA. Sustained activation of mTORC1 in skeletal muscle inhibits constitutive and starvation-
induced autophagy and causes a severe, late-onset myopathy. *Cell Metab* 17: 731-744, 2013.
57.   **Ching JK, Elizabeth SV, Ju JS, Lusk C, Pittman SK, and Weihl CC**. mTOR dysfunction contributes
to vacuolar pathology and weakness in valosin-containing protein associated inclusion body myopathy.
Hum Mol Genet 22: 1167-1179, 2013.
58.   **Goodman CA, and Hornberger TA**. Measuring protein synthesis with SUnSET: a valid alternative
to traditional techniques? *Exerc Sport Sci Rev* 41: 107-115, 2013.
59.   **Baehr LM, Tunzi M, and Bodine SC**. Muscle hypertrophy is associated with increases in
proteasome activity that is independent of MuRF1 and MAFbx expression. *Front Physiol* 5: 69, 2014.
60.   **Gomes AV, Waddell DS, Siu R, Stein M, Dewey S, Furlow JD, and Bodine SC**. Upregulation of
proteasome activity in muscle RING finger 1-null mice following denervation. *The FASEB Journal* 26:
2986-2999, 2012.
61.   **Heinemeier KM, Olesen JL, Haddad F, Schjerling P, Baldwin KM, and Kjaer M**. Effect of
unloading followed by reloading on expression of collagen and related growth factors in rat tendon and
muscle. *Journal of applied physiology* 106: 178-186, 2009.
62.   **Baehr LM, West DWD, Marcotte G, Marshall AG, De Sousa LG, Baar K, and Bodine SC**. Age-
related deficits in skeletal muscle recovery following disuse are associated with neuromuscular junction
instability and ER stress, not impaired protein synthesis. *Aging* 8: 127-146, 2016.
63.   **Baehr LM, West DWD, Marshall AG, Marcotte GR, Baar K, and Bodine SC**. Muscle-specific and
age-related changes in protein synthesis and protein degradation in response to hindlimb unloading in
rats. *J Appl Physiol (1985)* 122: 1336-1350, 2017.
64.   **Hughes DC, Marcotte GR, Baehr LM, West DW, Marshall AG, Ebert SM, Davidyan A, Adams**
CM, Bodine SC, and Baar K. Alterations in the muscle force transfer apparatus in aged rats during
unloading and reloading: impact of microRNA-31. *The Journal of physiology* 596: 2883-2900, 2018.
65.   **Biever A, Puighermanal E, Nishi A, David A, Panciatici C, Longueville S, Xirodimas D,**
Gangarossa G, Meyuhas O, Hervé D, Girault J-A, and Valjent E. PKA-Dependent Phosphorylation of
Ribosomal Protein S6 Does Not Correlate with Translation Efficiency in Striatonigral and Striatopallidal
Medium-Sized Spiny Neurons. *The Journal of Neuroscience* 35: 4113-4130, 2015.

66.   **Bonito-Oliva A, Pallottino S, Bertran-Gonzalez J, Girault J-A, Valjent E, and Fisone G**.
Haloperidol promotes mTORC1-dependent phosphorylation of ribosomal protein S6 via dopamine- and
cAMP-regulated phosphoprotein of 32 kDa and inhibition of protein phosphatase-1.
Neuropharmacology 72: 197-203, 2013.
67.   **Roux PP, Shahbazian D, Vu H, Holz MK, Cohen MS, Taunton J, Sonenberg N, and Blenis J**.
RAS/ERK signaling promotes site-specific ribosomal protein S6 phosphorylation via RSK and stimulates
cap-dependent translation. *J Biol Chem* 282: 14056-14064, 2007.
68.   **Sartori R, Romanello V, and Sandri M**. Mechanisms of muscle atrophy and hypertrophy:
Implications in health and disease. *Nature Communications* 12: 1-12, 2021.
69.   **Ogasawara R, and Suginohara T**. Rapamycin-insensitive mechanistic target of rapamycin
regulates basal and resistance exercise-induced muscle protein synthesis. *The FASEB Journal* 32: 5824-
5834, 2018.
70.   **Ogasawara R, Knudsen JR, Li J, Ato S, and Jensen TE**. Rapamycin and mTORC2 inhibition
synergistically reduce contraction-stimulated muscle protein synthesis. *The Journal of physiology* 598:
5453-5466, 2020.
71.   **You J-S, McNally RM, Jacobs BL, Privett RE, Gundermann DM, Lin K-H, Steinert ND, Goodman
CA, and Hornberger TA**. The role of raptor in the mechanical load-induced regulation of mTOR signaling,
protein synthesis, and skeletal muscle hypertrophy. *The FASEB Journal* 33: 4021, 2018.
72.   **Burd NA, Holwerda AM, Selby KC, West DW, Staples AW, Cain NE, Cashaback JG, Potvin JR,
Baker SK, and Phillips SM**. Resistance exercise volume affects myofibrillar protein synthesis and anabolic
signalling molecule phosphorylation in young men. *The Journal of physiology* 588: 3119-3130, 2010.
73.   **Williamson D, Gallagher P, Harber M, Hollon C, and Trappe S**. Mitogen-activated protein kinase
(MAPK) pathway activation: effects of age and acute exercise on human skeletal muscle. *The Journal of
physiology* 547: 977-987, 2003.
74.   **Steinert ND, Potts GK, Wilson GM, Klamen AM, Lin KH, Hermanson JB, McNally RM, Coon JJ,
and Hornberger TA**. Mapping of the contraction-induced phosphoproteome identifies TRIM28 as a
significant regulator of skeletal muscle size and function. *Cell Rep* 34: 108796, 2021.
75.   **Cho JH, Kim SA, Seo Y-S, Park SG, Park BC, Kim J-H, and Kim S**. The p90 ribosomal S6 kinase–
UBR5 pathway controls Toll-like receptor signaling via miRNA-induced translational inhibition of tumor
necrosis factor receptor–associated factor 3. *Journal of Biological Chemistry* 292: 11804-11814, 2017.
76.   **Pla-Prats C, and Thomä NH**. Quality control of protein complex assembly by the ubiquitin–
proteasome system. *Trends in cell biology* 32: 696-706, 2022.
77.   **Komander D, and Rape M**. The Ubiquitin Code. *Annual Review of Biochemistry* 81: 203-229,
2012.
78.   **Hogan AK, Sathyan KM, Willis AB, Khurana S, Srivastava S, Zasadzińska E, Lee AS, Bailey AO,
Gaynes MN, and Huang J**. UBR7 acts as a histone chaperone for post-nucleosomal histone H3. *The
EMBO journal* 40: e108307, 2021.
79.   **Taherbhoy AM, and Daniels DL**. Harnessing UBR5 for targeted protein degradation of key
transcriptional regulators. *Trends in Pharmacological Sciences* 2023.
80.   **Duan C-y, Li Y, Zhi H-y, Tian Y, Huang Z-y, Chen S-P, Zhang Y, Liu Q, Zhou L, and Jiang X-g**. E3
ubiquitin ligase UBR5 modulates circadian rhythm by facilitating the ubiquitination and degradation of
the key clock transcription factor BMAL1. *Acta Pharmacologica Sinica* 45: 1793-1808, 2024.
81.   **Mark KG, Kolla S, Aguirre JD, Garshott DM, Schmitt S, Haakonsen DL, Xu C, Kater L, Kempf G,
and Martínez-González B**. Orphan quality control shapes network dynamics and gene expression. *Cell*
186: 3460-3475. e3423, 2023.
82.   **Baghdadi M, Nespital T, Monzó C, Deelen J, Grönke S, and Partridge L**. Intermittent rapamycin
feeding recapitulates some effects of continuous treatment while maintaining lifespan extension. *Mol
Metab* 81: 101902, 2024.

83. **Miller RA, Harrison DE, Astle CM, Fernandez E, Flurkey K, Han M, Javors MA, Li X, Nadon NL,**
Nelson JF, Pletcher S, Salmon AB, Sharp ZD, Van Roekel S, Winkleman L, and Strong R. Rapamycin-
mediated lifespan increase in mice is dose and sex dependent and metabolically distinct from dietary
restriction. *Aging Cell* 13: 468-477, 2014.
84. **O'Reilly CL, Bodine SC, and Miller BF**. Current limitations and future opportunities of tracer
studies of muscle ageing. *The Journal of Physiology* 603: 7, 2023.
85. **Brandman O, Stewart-Ornstein J, Wong D, Larson A, Williams CC, Li G-W, Zhou S, King D, Shen**
PS, and Weibezahn J. A ribosome-bound quality control complex triggers degradation of nascent
peptides and signals translation stress. *Cell* 151: 1042-1054, 2012.

A)

Empty Vector (EV)

Overexpression

Akt-WT or Akt-CA

7d

B)

-7d

Rapamycin or control diet

Empty Vector (EV)

Overexpression

Akt-CA

7d

C)

Empty Vector (EV)

UBR5 RNAi

Rapamycin or control diet

30d

A)

Nerve Crush Injury Model

	Control	21d	28d	45d	60d	Kda
UBR1						— 150
UBR2						— 150
UBR4						— 250
UBR5						— 250
UBR7						— 25

B)

Protein Level (% Control) vs E3 Ligase (UBR1, UBR2, UBR4, UBR5, UBR7)

Control, 21d, 28d, 45d, 60d — Regrowth Phase

Specialization of UBR E3 ligases to perturbations in skeletal muscle size

RESULTS

CONCLUSION Novel UBR-box E3 ubiquitin ligases are responsive to heightened protein synthesis and alterations in skeletal muscle mass and fiber size.