Continental-scale genomic surveillance of *Plasmodium falciparum* malaria with rapid nanopore sequencing

Mulenga Mwenda[1]**, Karolina Mosler**[2]**, Bernd Bohmeier**[2]**, Miriam Chomba**[3]**, Welmoed Van Loon**[4]**, Brenda Mambwe**[1]**, Amy Gaye**[5]**, Adedolapo Olorunfemi**[6]**, Salma Suliman**[6]**, Nassandba Julien Yanogo**[7]**, Djiby Sow**[5]**, Bassirou Ngom**[5]**, Oumou Aïcha Zeïna Zoure**[7]**, Yssimini Nadege Guillene Tibiri**[7]**, Fiyinfoluwa Ojeniyi**[6]**, Arsène Zongo**[8]**, Etilé A. Anoh**[9]**, Vincent Achi**[9]**, Carol Chiyesu**[1]**, Sheila Otieno**[3]**, Bixa Ogola**[3]**, Moussa Niangaly**[10,2]**, Manuela Carrasquilla**[2]**, Dagaga Kenea Goboto**[11,12]**, Torsten Feldt**[11,13]**, Tafese Beyene Tufa**[11,12,13]**, Rafael Oliveira**[4]**, Emma Schallenberg**[4]**, Yuhana Sogoba**[10]**, Christina Ntalla**[2]**, Oumarou Ouedraogo**[7]**, Kephas Otieno**[3]**, Oluyinka Opaleye**[6,14]**, Adekunle Olowe**[6,14]**, Marley Gibbons**[1]**, Chris Drakeley**[1]**, Grit Schubert**[8]**, Frank P. Mockenhaupt**[4]**, Silvia Portugal**[2]**, Awa B. Deme**[5]**, Issiaka Soulama**[7]**, Daouda Ndiaye**[5]**, Olusola Ojurongbe**[6,14]**, Simon Kariuki**[3]**, Ya Ping Shi**[15]**, Jonathan S. Schultz**[15]**, Moonga Hawela**[16]**, Daniel J. Bridges**[1]**, and Jason A. Hendry**[2,*]

[1]PATH, Lusaka, Zambia
[2]Max Planck Institute for Infection Biology, Berlin, Germany
[3]Centre for Global Health Research, Kenya Medical Research Institute, Kisumu, Kenya
[4]Charité – Universitaetsmedizin Berlin, Charité Center for Global Health, Institute of International Health, Berlin, Germany
[5]Centre International de recherche, de Formation en Genomique Appliquee et de Surveillance Sanitaire (CIGASS), Dakar, Senegal
[6]Center for Emerging and Re-emerging Infectious Diseases, Ladoke Akintola University of Technology, Ogbomoso, Nigeria
[7]Department of Biomedical and Public Health, Health Sciences Research Institute (IRSS), National Center for Scientific and Technological Research (CNRST), Ouagadougou, Burkina Faso
[8]Robert Koch Institute, Centre for International Health Protection (ZIG), Berlin, Germany
[9]Centre Hospitalier et Universitaire de Bouaké, Bouaké, Côte d'Ivoire
[10]Parasites and Microbes Research and Training Center (MRTC), University of Science, Techniques and Technologies (USTTB), Bamako, Mali
[11]Hirsch Institute of Tropical Medicine, Asella, Ethiopia
[12]College of Health Sciences, Arsi University, Asella, Ethiopia
[13]Department of Gastroenterology, Hepatology and Infectious Diseases, University Hospital Düsseldorf, Heinrich Heine University Düsseldorf, Germany
[14]Department of Medical Microbiology and Parasitology, Ladoke Akintola University of Technology, Ogbomoso, Nigeria
[15]Division of Parasitic Diseases and Malaria, National Center for Emerging and Zoonotic Diseases, Centers for Disease Control and Prevention, Atlanta, GA, USA
[16]National Malaria Elimination Centre, Chainama, Lusaka, Zambia
[*]hendry@mpiib-berlin.mpg.de

ABSTRACT

In sub-Saharan Africa, continental-scale genomic surveillance of *Plasmodium falciparum* malaria is needed to track the spread of antimalarial drug resistance and diagnostic test evasion, as well as to monitor parasite evolutionary responses to vaccine rollout. Yet implementation of malaria genomic surveillance at a continental-scale is hindered by resource constraints, the vastness of the continent, and the lack of sequencing protocols suitable for most local laboratories. To address this, we developed an approach to enable a decentralized scale-up of *P. falciparum* genomic surveillance and established it in six African countries in one year, locally sequencing 1,065 samples. The approach includes a rapid (∼5 hours) and cost-efficient (<$25 USD/sample) nanopore sequencing protocol that provides surveillance of a panel of drug resistance-associated genes, *hrp2/3* gene deletions, the vaccine target *csp*, and the highly diverse gene *ama1*. We coupled this to a bioinformatics dashboard that runs offline on a laptop and displays mapping and variant calling results in real-time. We demonstrate robust sequencing coverage across parasitemia levels and laboratories, accurate detection of *hrp2/3* deletions in field samples and accurate identification of antimalarial drug resistance markers. Our approach will accelerate genomic surveillance of *P. falciparum* malaria across sub-Saharan Africa at a time of urgent need.

Introduction

The endeavor to eliminate *Plasmodium (P.) falciparum* malaria from sub-Saharan Africa – where it causes over half a million deaths annually – relies heavily on accurate diagnosis with rapid diagnostic tests (RDTs) and effective treatment with artemisinin-based combination therapies (ACTs)[1]. *P. falciparum*, however, is evolving to undermine both of these tools. For RDTs, *P. falciparum* strains that have deletions of the *hrp2* and *hrp3* genes evade diagnosis by producing false-negative test results[2–6]. For ACTs, mutations in the *kelch13* gene[7] can cause delayed parasite clearance[8] and, if combined with partner drug resistance, treatment failure[9]. Since at least 2016, these *kelch13* mutations have been spreading in East Africa[10–13], where they have begun to co-circulate with *hrp2/3* deletions[14]. They are now being described in Southern[15] and Central Africa[16, 17]. Public health institutions are urgently exploring a variety of responses, such as switching to alternative RDTs when the *hrp2/3* deletion frequency surpasses a threshold regionally[18]; or deploying multiple first-line therapies (MFT) to mitigate the spread of *kelch13* mutations[19, 20]. However, essential to coordinating these responses are high-quality, granular and timely malaria surveillance data, which are lacking in many parts of Africa.

The need to increase malaria surveillance in Africa could be met using genomics. Genomic methods like amplicon sequencing[21–23] and molecular inversion probes (MIPs)[24, 25] can interrogate tens to thousands of *P. falciparum* genes in parallel, enabling reporting on many control-relevant genetic markers at once. This makes them more efficient than conventional molecular methods, which typically only report on a single gene or mutation. Moreover, many genomic methods require just a single dried-blood spot (DBS) as the sample, making them less invasive and labour intensive than standard surveillance approaches like therapeutic efficacy studies. In principle, these genomics approaches could affordably and routinely generate high-quality malaria surveillance data. But scaling *P. falciparum* genomic surveillance to meet the growing need across sub-Saharan Africa is an immense challenge. First, the region is vast and diverse, spanning over 24 million km^2 and encompassing 49 countries. Perhaps 50,000 to 100,000 samples would need to be sequenced annually to have adequate statistical power to detect emerging mutations across the entire region. Second, access to Next-Generation Sequencing (NGS) platforms is limited in many areas. Where NGS platforms do exist, maintenance and the procurement of reagents is invariably unreliable

and expensive. Similarly, the computational infrastructure (and/or internet connectivity) required to process large volumes of genomic data is frequently lacking. As a consequence, most *P. falciparum* genomic data have been generated by shipping samples out of Africa, a slow process that also impedes local capacity development and ownership. More recently, centralised sequencing facilities located within Africa are beginning to generate malaria genomic data domestically and regionally, but questions over capacity and country ownership persist.

The possibility of scaling *P. falciparum* genomic surveillance across sub-Saharan Africa in a decentralised way has been relatively neglected. The decentralised approach would leverage the portable and low-cost MinION device from Oxford Nanopore Technologies (ONT)[26] to conduct genomic surveillance from a network of laboratories that span the continent. This would circumvent the need for large capital investments associated with centralised facilities and Illumina platforms, and instead engage more of the existing funding, infrastructure and human potential latent in the thousands of conventional laboratories across Africa. These laboratories could serve their local geography by interacting closely with nearby clinics and public health officials; while simultaneously coordinating activities and sharing data to improve national and continental understanding. Beneficially, a network of decentralised sequencing laboratories would be more robust to disruption and could grow autonomously and exponentially. Until now, the major drawback for this paradigm has been that nanopore sequencing protocols for *P. falciparum* genomic surveillance, although increasing in number[27–30], are less mature and comprehensive than Illumina-based ones[21–23,31]. For example, state-of-the-art Illumina-based amplicon sequencing protocols[23] concurrently provide information on antimalarial drug resistance, *hrp2/3* deletions, the vaccine target *csp* and multiple genetic diversity markers. In contrast, existing nanopore sequencing protocols typically provide information on only one[27,28] or two[29,30] of these outputs. An exception is the NOMADS16 protocol[32] but its focus on generating long-read data (3–4 kbp) results in it being substantially less sensitive than short-read approaches.

Here, we have filled this gap by developing a novel nanopore sequencing protocol for *P. falciparum* malaria that is rapid, cost-effective, sensitive, and comprehensive — providing information on a panel of genes associated with antimalarial drug resistance (*crt*, *dhfr*, *dhps*, *kelch13*, *mdr1*), as well as *hrp2/3* deletions, the vaccine target *csp*, and the highly polymorphic gene *ama1*. To enable on-site analysis, we developed a real-time bioinformatics pipeline and dashboard that can run on the same laptop used for sequencing. Using these tools, we locally sequenced and analysed 1,065 *P. falciparum* DBS samples from six laboratories spanning sub-Saharan Africa in one year. In addition, we validated critical aspects of protocol performance across mock and field samples.

Results

A protocol to rapidly scale-up decentralised malaria genomic surveillance

We developed an optimal nanopore sequencing protocol for scaling decentralised *P. falciparum* genomic surveillance across sub-Saharan Africa (Fig. 1a). First, we designed a novel amplicon panel using our open-source multiplex PCR primer design software, *Multiply*[32] (Methods). We targeted nine genes that would jointly provide information on antimalarial drug resistance,

hrp2/3 deletions, vaccine target evolution and complexity of infection (COI) (Table 1). Each gene was targeted with a single amplicon, except *mdr1* for which separate N- and C-terminal amplicons were designed. The amplicons vary in size from 604 bp to 1,405 bp and amplify a combined 10,860 bp in a single reaction. A total of 45 validated and candidate World Health Organisation (WHO) antimalarial drug resistance-associated single-nucleotide polymorphisms (SNPs) are genotyped (Table 1). The amplicons targeting *hrp2* and *hrp3* both have reverse primers annealing in exon 1, and forward primers upstream of exon 2, which enables detection of partial or complete hrp gene deletions[33]. The *csp* amplicon spans from codon 19 to the end of the gene, which includes the entire central repeat region and C-terminal domain used in the RTS,S/AS01 and R21/Matrix-M vaccines[34,35]. A high diversity, 826 bp window of *apical membrane antigen 1* (*ama1*) is covered to support COI estimation. We named this amplicon panel the NOMADS Minimal Viable Panel (MVP).

No.	Reference ID	Gene Name	Amplicon (bp)	Codons Spanned	Key Regions
1	PF3D7_1133400	*ama1*	826	$74 - 384$	
2	PF3D7_0709000	*crt*	692	$14 - 125$	K76T
3	PF3D7_0304600	*csp*	1,273	$19 - 398^{\dagger}$	RTS,S and R21 epitopes
4	PF3D7_0417200	*dhfr*	1,405	$1 - 410$	A16V, N51I, C59R, S108M, I164L
5	PF3D7_0810800	*dhps*	1,327	$317 - 707^{\dagger}$	S436A/F, A437G, K540E, A581G, A613T/S
6	PF3D7_0831800	*hrp2*	1,296	$14 - 306^{\dagger}$	
7	PF3D7_1372200	*hrp3*	1,216	$14 - 276^{\dagger}$	
8	PF3D7_1343700	*kelch13*	1,286	$383 - 727^{\dagger}$	BTB/Propeller
9	PF3D7_0523000	*mdr1*(n)	935	$46 - 245$	N86Y/F, Y184F
10	PF3D7_0523000	*mdr1*(c)	604	$968 - 1278$	S1034C, N1024D, D1246Y

Table 1. Amplicons included in NOMADS-Minimal Viable Panel (MVP). For *kelch13* all WHO candidate and validate markers are spanned[36]. *mdr1*(n) and *mdr1*(c) indicate N- and C-terminal *mdr1* amplicons respectively. bp, base pairs. †Amplicon spans C-terminal codon.

To promote widespread adoption, we developed a nanopore sequencing protocol for the NOMADS-MVP amplicons that minimises costs, complexity, and time. We initially explored two versions of the protocol: one that included a selective whole genome amplification (sWGA)[37] step to preamplify bulk *P. falciparum* DNA, and one where the multiplex PCR was performed directly on extracted DNA. We reduced the run time of the sWGA step from 17 hours[37] to 1 hour while still providing robust yield by using the EquiPhi29 polymerase (Methods). We hypothesised that including sWGA might increase sensitivity, although at additional protocol cost, complexity and time. While this was confirmed with mock DBS samples created from laboratory strains, we saw no benefit on field DBS samples and discontinued use of sWGA in August 2024. We optimised the NOMADS-MVP multiplex PCR using gradient PCR and several full factorial experimental designs: varying template DNA, primer and polymerase concentrations, annealing temperature, and the number of thermal cycles (Supplementary Fig. 1). After optimisation, multiplex PCR conditions maximised sensitivity (with amplification on mock DBS samples down to 100 parasites/μL) and simplicity. The final PCR only requires pipetting three reagents (primers, KAPA HiFi ReadyMix buffer, and sample), thus reducing contamination risk and increasing consistency. Finally, we adopted a rapid barcoding approach that dramatically reduces library preparation time and complexity, and simplifies procurement as no additional third-party reagents are required.

Starting from extracted DNA, the final optimised protocol costs $21 USD/sample, assuming 48 samples are sequenced in a batch and including all plasticware ($5.47 USD/sample, 26.2%) and reagents ($15.45 USD/sample, 78.3%; Supplementary Table 1). The three most costly items are the flow cells ($7.2 USD/sample, 34.4%), the KAPA HiFi polymerase ($3.19 USD/sample, 15.3%) and the 20 μL multichannel pipette tips ($2.50 USD/sample, 12.0%). All the required equipment (including a laptop suitable for analysis) costs an estimated $12,500 USD (Supplementary Table 2). To objectively measure protocol complexity, we quantified the number of pipetting steps from extracted DNA to sequencing (1 step = aspirating and dispensing a reagent or sample). For a batch of 48 samples, our protocol requires 172 pipetting steps, compared to 292 steps for our previous long-read protocol[32] or over 400 steps for two state-of-the-art Illumina-based protocols[23,31] (Supplementary Table 3). The total incubation time for all steps in the NOMADS-MVP protocol is approximately 3.5 hours. After gaining familiarity, the workflow from extracted DNA to sequencing takes approximately 5 hours (Fig. 1a, Supplementary Fig. 2).

A real-time, point-of-use bioinformatics pipeline and dashboard

We developed a real-time, on-site bioinformatics pipeline and analysis dashboard which accompanies the NOMADS-MVP protocol, called *Nomadic* (Fig. 1b). *Nomadic* maps reads, computes quality control statistics for each sample and amplicon, performs preliminary variant calling and annotation, and presents this information in a graphical dashboard while sequencing is ongoing. It allows laboratory scientists to make informed decisions regarding sample quality and amplicon performance, determine when to stop sequencing, and provides an immediate first-look at mutations present across samples. The dashboard is easy to install and is launched with a single command from the terminal, enabling use by scientists without significant bioinformatics training. Beyond what is required to perform real-time basecalling with *MinKNOW* (a laptop with a CUDA-enabled graphics processor unit), *Nomadic* requires no additional computational infrastructure and runs offline. In addition, *Nomadic* produces human-readable summary files that contain all key information about sequencing performance and detected mutations. These files are small enough (<20 Mbp) to be easily shared even over slow internet connections and are sufficient to reopen the graphical dashboard after the experiment is completed. *Nomadic* was used to process all the data described in this manuscript at the point of sequencing. Further details on *Nomadic* use and implementation are provided in the Methods.

Implementation at scale across sub-Saharan Africa

From April 2024 to July 2025, a total of 1,404 dried-blood spot (DBS) samples from eight countries across sub-Saharan Africa were sequenced using the NOMADS-MVP protocol (Fig. 1c-e). Of these, 1,065 (65.8%) were sequenced locally, with the protocol being established in six countries: Nigeria (April 2024), Kenya (July 2024), Zambia (August 2024), Senegal (March 2025), Burkina Faso (April 2025), and Côte d'Ivoire (May 2025) (Fig. 1c). Overall, 13 local scientists independently conducted 28 sequencing runs across these six countries (Fig. 1d). The NOMADS-MVP protocol enabled periods of high sample processing throughput. For example, in Kenya two scientists processed 190 samples in 7 days at the end of July 2024 (Fig. 1d); and in late 2024, 320 samples were processed in 24 days by two scientists in Zambia.

Implementation of the protocol was feasible across laboratories that varied considerably with respect to prior sequencing

Figure 1. Overview of sequencing approach and implementation across sub-Saharan Africa. (a) The laboratory protocol uses *P. falciparum* positive dried-blood spots (DBS) as source material for DNA extraction. From extracted DNA to sequencing takes approximately 5 hours. (b) Data analysis occurs in real-time on a laptop. While sequencing is ongoing, quality control (QC) and variant calling results are displayed to an interactive dashboard. GPU: graphics processor unit. Sequencing time depends on flow cell quality and number of samples. (c) DBS samples were processed from 8 countries across sub-Saharan Africa. For six countries, all sequencing occurred locally (filled circles); for two countries, samples were sequenced internationally (Mali, Ethiopia; open circles). (d) Timeline of sequencing runs. Size of point indicates number of samples per run. Total sequencing runs for each country is shown in a box at right. (e) Barplot displaying total number of samples sequenced per country. Overall, 1,065/1,404 (65.8%) of samples were processed in Africa.

experience and available infrastructure. One team had both Illumina and ONT sequencing experience (CIGASS, Senegal); three teams had previous nanopore sequencing experience (NMEC, Zambia; HRH-CERID, Nigeria; CHUB, Côte d'Ivoire); and for the remainder our protocol was the first experience with NGS. Infrastructure ranged from well-equipped regional sequencing hubs (CIGASS, Senegal) to small container laboratories with only the essentials for nanopore sequencing (NMEC, Zambia). For five countries, an initial in-person training was conducted by a core team (typically 1–3 weeks in duration) and followed by remote support. In Côte d'Ivoire, the protocol was established locally through a collaboration with the Robert Koch Institute. Overall, these results demonstrate that our approach is amenable to large-scale implementation across a wide range of contexts in sub-Saharan Africa.

Robust sequencing coverage across countries and parasitemia levels

We examined the sequencing coverage generated by the NOMADS-MVP protocol on a set of 1,283 samples, comprising 1,154 field DBS samples (1,154/1,283, 89.9%) and 129 mock DBS samples (129/1,283, 10.1%), all processed without sWGA (Fig. 2). Parasitemia data were available for 950 samples (950/1,283, 74.0%), which together had a median of 3,645 parasites/μL (IQR: 631–11,540 parasites/μL). To evaluate sequencing performance, we calculated two per-sample metrics: (1) the mean coverage across all amplicons; (2) the fold-difference in coverage between the most- and least-abundant amplicon (excluding the *hrp2/3* amplicons), a measure of uniformity where a value of 1 indicates perfectly uniform coverage across amplicons (see Methods).

Across all samples, the median per-sample mean coverage was 1,881$\times$ (IQR: 1,064–3,086$\times$), with 76.3% (881/1,154) of field DBS samples and 87.5% (113/129) mock DBS samples achieving a mean coverage of greater than 1,000$\times$. Across countries, the median per-sample mean coverage ranged from 544$\times$ in Côte d'Ivoire ($n = 29$) to 4,161$\times$ in Burkina Faso ($n = 88$), and 6 of 8 countries had a median per-sample mean coverage exceeding 1,000$\times$ (Fig. 2a). The median fold-difference in coverage across all samples was 10.2$\times$ (IQR: 4.8–53.6$\times$), with samples sequenced in Senegal having the most uniform coverage across amplicons ($n = 110$, median: 2.97$\times$, IQR: 2.14–4.95$\times$) and those from Zambia had the least uniform coverage ($n = 457$, median: 44.4$\times$, IQR: 9.5–197.4$\times$).

Next, for each sample we quantified the number of amplicons with greater than 50$\times$ coverage, which is sufficient for high-accuracy variant calling. Overall, 96.1% (124/129) of mock DBS samples and 87.9% (1,015/1,154) of field DBS samples had at least 8 amplicons with greater than 50$\times$ coverage. In 68.2% (88/129) of mock DBS samples and 64.5% of field DBS samples (745/1,154) all 10 amplicons exceeded 50$\times$ coverage. By country, the percentage of samples with ≥ 8 amplicons exceeding 50$\times$ coverage ranged from 69% in Côte d'Ivoire to 95% in Kenya (Fig. 2b). Combined, these results demonstrate that NOMADS-MVP can generate robust sequencing coverage across a diversity of sample sets and laboratories in sub-Saharan Africa.

Two major factors influencing sequencing performance are DNA quality and sample parasitemia. We observed a moderate positive correlation between parasitemia and per-sample mean coverage for both field DBS samples (Spearman's $\rho = 0.32$, $p < 0.001$) and mock DBS samples ($\rho = 0.37$, $p < 0.001$). For example, in field DBS samples, the mean coverage at 1,000 parasites/μL (2,099$\times$) was about twice that at 100 parasites/μL (1,011$\times$). A stronger, negative correlation was observed between parasitemia and the per-sample fold-difference in coverage across amplicons (field DBS, $\rho = -0.43$, $p < 0.001$; mock DBS, $\rho = -0.88$, $p < 0.001$); this increased imbalance in coverage at lower parasitemia levels was driven primarily by decline (and occasional drop-out) of the two longest amplicons, which target *dhps* and *dhfr* (Supplementary Fig. 3). Finally, we evaluated the percentage of field DBS samples that had 8, 9, or 10 amplicons with greater than 50$\times$ coverage as a function of parasitemia. These percentages were consistently high for samples with $\geq$1,000 parasites/μL, declining steadily below this threshold. For example, at 1,000 parasites/μL greater than 90% of samples had at least 8 amplicons with 50$\times$ coverage, compared to approximately 75% of samples at 100 parasites/μL. Including field DBS samples from Ethiopia, some of which had *hrp2/3* deletions, resulted in a lower percentage of samples with 9 or 10 amplicons exceeding 50$\times$ coverage. Interestingly,

Figure 2. Sequencing coverage across countries and parasitemia levels. (a) Strip plots of parasitemia and sequencing coverage data for 1,283 samples processed with the NOMADS-MVP protocol, grouped by country. Each point represents a sample and black circles indicate country-level medians. The left subpanel shows sample parasitemia (parasites/μL); the middle subpanel shows mean per-sample coverage across amplicons; and the right subpanel shows per-sample coverage fold-difference between the least and most abundant amplicon (see Methods). All x-axes are on a logarithmic scale. (b) Bar plot showing, for each country, the percentage of samples with $> 50\times$ coverage for 8 amplicons (lightest shade), 9 amplicons (intermediate), or all 10 amplicons (darkest); bar height indicates the total percentage of samples with $> 50\times$ coverage for ≥ 8 amplicons. At right, percentages for 10 amplicons and ≥ 8 amplicons passing $> 50\times$ coverage is annotated. (c) Scatter plots of parasitemia against mean coverage (left subpanel); and coverage fold-difference (right subpanel). Each point is a sample, coloured by country. Black lines are a locally weighted scatterplot smoothing (LOWESS). Spearman's ρ for the field (ρ_{field}) and lab (ρ_{lab}) samples is annotated at top left. Both axes use logarithmic scales. (d) Relationship between parasitemia and samples passing (%). Field samples were ordered by parasitemia, and rolling means were computed using windows of 80 samples. Lines represent percentage of samples with $> 50\times$ coverage for ≥ 8 amplicons (thinnest line), ≥ 9 amplicons (medium), and for all 10 amplicons (thickest). The top subpanel shows the proportion of samples from each country (by colour) across the rolling parasitemia windows. Only field samples were included.

the mock DBS samples generated from laboratory strains performed consistently well even at 100 parasites/μL (Supplementary Fig. 4). One explanation is that while mock DBS samples recapitulate absolute parasitemia, they represent a best-case scenario in terms of DNA quality, which may be challenging to achieve in the field.

Rapid sequencing produces sufficiently long reads

The Rapid Barcoding Kit (SQK-RBK114.96) from ONT leverages transposome-based chemistry that cleaves sample DNA during barcode ligation. The barcoding reaction takes just four minutes, but the sample DNA is fragmented in the process. To characterise the impact on read lengths, we randomly selected 80 samples from across four different sequencing experiments, including field DBS samples from Kenya ($n = 20$), Ethiopia ($n = 20$) and Mali ($n = 20$), as well as a set of mock DBS samples created from laboratory strains in Germany ($n = 20$). Across these samples, a total of 3.45 million reads mapped uniquely to the *P. falciparum* reference genome and their median length was 567 bp (IQR: 376–808 bp). There was considerably more read length variation within individual samples than between samples or experiments (Fig. 3a). The consistency across four independent sample sets and experiments suggests that the read length distribution is robust to technical and biological factors. Similarly, in optimisation experiments we found that read lengths were insensitive to variation in the barcoding reaction time or input DNA mass (Supplementary Fig. 5). The within-sample read length variation was driven by the different amplicons in our panel, with, as expected, longer amplicons generating longer reads (Fig. 3b). In particular, when grouped by the amplicon to which they mapped, reads had a median length of roughly half of their amplicon length (mean: 59%, range: 45%–69%), consistent with a single cleavage and barcoding event per amplicon (Fig. 3b). Across amplicons, the longest median read length was for the *kelch13* amplicon at 872 bp and the shortest for the *mdr1* N-terminal amplicon at 397 bp.

Figure 3. Read lengths generated by the NOMADS-MVP protocol. (a) Read length distributions for 80 samples from four countries. The top subpanel shows the percentage of reads (y-axis) longer than a given length (x-axis). Each line represents one sample; colours indicate country. The bottom subpanel shows box plots of read length distributions by country. Boxes indicate the interquartile range, black lines mark the median. (b) Same as (a), but reads are grouped by target amplicon rather by than sample. In the bottom subpanel, the transparent bar behind each box indicates the full amplicon length. (c) Strip plot showing the percentage of sequencing reads fully spanning a given genomic region of interest for each sample. Genomic regions are defined by gene and codon range. 4CAST, region of *ama1* amplified in *La Verriere et al.*[22]; TSR, thrombospondin-like type I repeat; KREP, Kelch-repeat propeller. Black circles indicate the region-level medians.

Longer reads support physical phasing of heterozygous variants and the investigation of repetitive regions, but only when the variants or regions are fully spanned by individual reads. To this end, we examined whether our data contained reads that

fully spanned regions of interest for *P. falciparum* genomic surveillance (Fig. 3c). All regions investigated had fully-spanning reads at appreciable frequencies across the eighty samples. For example, a median of 29.4% (IQR: 26.4%–33.5%) of reads per sample overlapping the Kelch-repeat propeller domain fully spanned the region. For *dhps*, which mediates Sulfadoxine resistance, a median of 40.0% (IQR: 35.9–43.0%) of reads spanned from codon 436 to 613, enabling joint interrogation of the key resistance markers (S436A, A437G, K540E, A581G and A613S). A high-diversity region of *ama1*[22] was spanned by a median of 76.5% (IQR: 72.1–80.6%) of reads per sample. Finally, key regions of the RTS,S/AS01 and R21/Matrix-M vaccine target *csp* also had fully-spanning reads, including the entire portion used in the vaccine (codons 199–397, median: 34.6% of reads fully-spanning) and the entire NANP central repeat region (codons 105–272, median: 40.2%).

SNPs are called accurately in clonal infections

To investigate the ability of NOMADS-MVP to identify known markers of antimalarial drug resistance, we sequenced six mock clonal *P. falciparum* infections. These included three mock samples created from laboratory strains (3D7, Dd2 and HB3) for which high-quality genome assemblies exist[38] and can be used to benchmark accuracy. Furthermore, we included three mock samples created from cultured *P. falciparum* strains from Cambodia which carry artemisinin-resistance associated *Kelch13* mutations[39]. The samples were sequenced in duplicate on two R10.4.1 flowcells and SNPs were called using *bcftools*[40], which has been shown to perform well on nanopore data in other applications[41]. For all six mock *P. falciparum* infections, we observed all expected antimalarial resistance markers and no false-positives in both replicates (Fig. 4a).

Next, we explored the effect of sequencing coverage on SNP calling accuracy by downsampling sequencing data from the mock *P. falciparum* samples *in silico*. For each mock sample, we downsampled from $10 - 100\times$ coverage in triplicate producing thirty-six replicates at each coverage-level (6 mock samples sequenced in duplicate, by 3 downsampling replicates). We compared to the true variants using *hap.py*[42] and evaluated performance using the F_1-score, which jointly summarises the precision and recall (Methods). With $20\times$ coverage or greater, the mean F_1-score was 1.0 (indicating perfect precision and recall) in all but two regions. The first was in *dhfr*, where we observed reduced F_1-scores ($0.77 - 0.97$) across nearly all mean coverage levels, but outside of the coding sequence. Further investigation revealed that these reduced F_1-scores were caused by false-positive mutations upstream of *dhfr*, likely driven by a large low-complexity region comprised of A+T repeats nearby (Supplementary Fig. 6a). The second reduction in F_1-score was in exon 2 of *crt*, exclusively in the *Dd2* mock samples carrying the CVIET haplotype (codons 72–76) associated with chloroquine resistance. When reads carrying the CVIET haplotype are aligned to the 3D7 reference genome (which carries the CVMNK haplotype) they produce an insertion, deletion and substitution mutation in close proximity (Supplementary Fig. 6b). As indel calling with *bcftools* is not recommended on nanopore data, the insertion and deletion are ignored. However, typically a small percentage of reads ($\sim$1%) contain sequencing errors that can cause them to align a mismatched nucleotide to the position carrying the deletion, which gets called as a substitution by *bcftools*, due to the lack of incorporation of information about the vast majority of reads carrying deletions. Enforcing a minimum coverage threshold can mitigate this issue and, in future, using a haplotype calling tool, or a variant caller that performs well on indels, will solve it. Regardless, the K76T mutation was called correctly in all replicates.

Figure 4. SNP calling accuracy for clonal mock samples. (a) SNP calling results for antimalarial drug-resistance associated SNPs covered by NOMADS-MVP, across six mock *P. falciparum* samples were sequenced in duplicate (R1: replicate 1, R2: replicate 2). The most common *kelch13* mutations circulating in Africa[13] are included in the panel (bold). Color indicates homozygous reference (white), heterozygous (light red) or homozygous alternative (dark red) SNP call. Strains prefaced with 'IPC' are from Cambodia. (b) Heatmaps showing the mean F_1-score (harmonic mean of precision and recall) at different mean coverage levels (x-axis) and over different regions (y-axis). The F_1-score is averaged over $n = 18$ replicates produced by downsampling to the indicated depth in triplicate for each mock sample (see Methods). The left pane is the F_1-score over the entire sequenced amplicon for each region; whereas the right pane is only over the coding sequence (CDS) of each amplicon. All: all amplicons included in NOMADS-MVP. Note that the F_1-scores are rounded to two decimal places, so values > 0.995 are displayed as 1.00.

High accuracy inference of *hrp2/3* deletions

We evaluated the ability of NOMADS-MVP to detect *hrp2* and *hrp3* deletions, which can cause false-negative RDT results and are currently prevalent in the Horn of Africa[2,3,5,6]. We used a set of 149 DBS samples collected from malaria patients in central Ethiopia, which had been previously assayed for *hrp2/3* deletions using recommended conventional PCR assays performed in duplicate[43]. We sequenced the 149 samples across four MinION sequencing runs, including in each run mock DBS samples created from the laboratory strains 3D7 (*hrp2+/hrp3+*), Dd2 (*hrp2−/hrp3+*) and HB3 (*hrp2+/hrp3−*), at parasitemia levels

ranging from 100–10,000 parasites/μL; as well as mock *P.falciparum*-negative samples. We inferred *hrp2/3* deletions using a statistical model that estimates sample quality (from the mean abundance of other amplicons in the panel) and the rate of background sequencing contamination (from the negative controls), before calculating a deletion probability for *hrp2* and *hrp3*[32].

Figure 5. **Validation of NOMADS-MVP *hrp2/3* deletion detection on mock and field DBS samples**. (a) Data for 36 mock DBS samples of different *hrp2/3* genotypes and mock *P.f.* samples, which contain only human DNA (Methods). From top to bottom, subpanels show: lab strains used in mock samples; scatter plot of parasitemia; bar plot of mean coverage; heatmap with rows indicating different amplicons in NOMADS-MVP, columns indicating samples, and color indicating mean coverage, normalised to the mean coverage for the experiment. (b) Heatmap showing probability of deletion for *hrp2* and *hrp3* from statistical model. Note uncertainty (i.e. probability in 0.2–0.8) for negative controls, due to low or absence of coverage. (c) Scatter plot showing mean coverage over *hrp2* for all mock samples. (d) Same as (c), but for *hrp3*. (e) Same as (a), but showing data for 149 Ethiopian DBS samples. Vertical lines delineate *hrp2/3* deletion status as estimated by NOMADS-MVP. (f) Estimated *hrp2/3* deletion status by NOMADS-MVP (top subpanel) and the conventional PCR-based assays (bottom). Samples marked with numerals in red font indicate discrepancies (ii, iii) or residual coverage (i).

NOMADS-MVP correctly estimated the *hrp2/3* deletion status for all mock samples (Fig. 5a). Mean amplicon coverage tended to decline with parasitemia and low levels of sequencing contamination were evident in some *P.f.*-negative controls. Notwithstanding, for both *hrp2* and *hrp3*, the difference in coverage between mock samples with and without deletions

consistently exceeded 100-fold. For example, the mean *hrp2* amplicon coverage in Dd2 versus 3D7 mock samples was $6.6\times$ versus $2,340\times$ at 10,000 parasites/μL; $1.8\times$ versus $1,968\times$ at 1,000 parasites/μL; and $1.1\times$ versus $984\times$ at 100 parasites/μL (Fig. 5b). Similarly, for *hrp3* the mean amplicon coverage in HB3 versus 3D7 was $8.8\times$ versus $4,910\times$ at 10,000 parasites/μL; $3.3\times$ versus $3,716\times$ at 1,000 parasites/μL; and $2.0\times$ versus $2,262\times$ at 100 parasites/μL (Fig. 5c).

Of the 149 field DBS samples from Ethiopia, 138 (92.6%) passed quality control. Among these, the joint *hrp2/3* deletion results from conventional PCR matched those from NOMADS-MVP in 136 samples (98.6%; Fig. 5d). The two discrepancies included one sample that was *hrp2−/hrp3−* by NOMADS-MVP but *hrp2+/hrp3−* by conventional PCR (Fig. 5c, indicated by 'ii'). This sample had a mean coverage of $425\times$ over the *hrp2* amplicon; a much higher mean coverage than other samples predicted as *hrp2−* (median: $0.3\times$, IQR: 0.0–$0.9\times$), but lower than those predicted *hrp2+* (median: $1,468\times$, IQR: 934–$3,433$). Heterozygosity over *ama1* indicated that the sample was a mixed infection with at least two *P.f.* clones. A possible explanation for this discrepancy is that one clone carries an *hrp2* deletion that is detected by the statistical model, but still yields a positive result by conventional PCR. Similarly, the second discrepancy (*hrp2−/hrp3−* by NOMADS-MVP, *hrp2−/hrp3+* by conventional PCR) had non-negligible coverage over *hrp3* ($78\times$) and *ama1* diversity suggesting a mixed infection (Fig. 5c, indicated by 'iii'). We also noted one other sample where, although conventional PCR and NOMADS-MVP both predicted an *hrp3* deletion, considerable coverage over the *hrp3* ($479\times$) was present (Fig. 5c, indicated by 'i'). None of the discordand samples had particularly low parasitemia (2,960 parasites/μL, i; 4,960 parasites/μL, ii; 9,080 parasites/μL, iii) or mean coverage ($1,505\times$, i; $1,752\times$, ii; $3,753\times$, iii), which might be expected if insufficient sensitivity was an underlying cause. All samples whose parasitemia or mean coverage was in the lowest decile (229–960 parasites/μL; or 323–$647\times$) had corresponding *hrp2/3* deletion calls between the PCR-based assays and NOMADS-MVP. Overall, these results demonstrate that NOMADS-MVP is a robust method for *hrp2/3* deletion detection, yielding results that align closely with conventional PCR assays in widespread use.

Discussion

Genomic surveillance of *P. falciparum* malaria can generate valuable public health data, but implementation across sub-Saharan Africa has been limited by a lack of sequencing approaches suitable for most local laboratories. Here, we described the development of a novel nanopore sequencing protocol for *P. falciparum* genomic surveillance which targets ten genomic regions, collectively providing information on a panel of antimalarial drug resistance-associated genes, *hrp2/3* deletions, the vaccine target *csp*, and the high diversity gene *ama1*. In one year, we implemented the protocol in Senegal, Burkina Faso, Côte d'Ivoire, Nigeria, Zambia and Kenya — locally sequencing and analysing 1,065 DBS samples — and demonstrating the feasibility of continental-scale decentralised *P. falciparum* genomic surveillance.

After genomic data are generated, difficulties performing bioinformatic analysis in sub-Saharan Africa often create another barrier to timely surveillance. We addressed this by developing a real-time bioinformatics dashboard called *Nomadic*, which has two main advantages. First, it can be run with minimal bioinformatics expertise, enabling a wide range of users to independently

and immediately derive insights from their data. Second, it runs offline on the same laptop used for sequencing, making it affordable and invulnerable to the internet connectivity issues which can bottleneck workflows reliant on transferring data to remote servers[30]. While analysis occurs on-site and raw data can remain local, *Nomadic* produces compact summary files that can be easily shared. These summary files are useful for collaborative troubleshooting and, in the future, could facilitate regional data integration when decentralised genomic surveillance becomes widespread.

Our protocol does have limitations that make it less well suited for some malaria genomic surveillance activities. The longer amplicons (500–1,500 bp) generated allow contiguous coverage of target genes and simpler multiplexing, but also require samples to have higher parasitemia and/or DNA quality for robust amplification. For clinical cases of malaria we have demonstrated robust performance with NOMADS-MVP across a wide range of studies. However, for asymptomatic or submicroscopic cases short-read sequencing approaches are likely to have more consistent performance and less dependence on DNA quality, although we note that several studies processed here contained asymptomatic cases and performed well (including dry-season samples from Mali and community-collected samples from Zambia). Similarly, for studies distinguishing *P. falciparum* recrudescence from reinfection, short-read approaches may enable more sensitive minor clone detection[23,30,31]. Beyond sensitivity, our assay lacks some outputs generated by other assays, such as between sample relatedness estimates. Relatedness and identity-by-descent (IBD) are best estimated by leveraging a larger number of unlinked variant sites (10s to 1000s) spread across the genome. Highly multiplexed short-read amplicon sequencing[21–23], MIPs[24,25], SNP barcodes[44] or whole-genome sequencing (WGS) methods are better suited to make these estimates for studies that require them. Accurate COI estimation will be possible leveraging the *ama1* and *csp* amplicons and work to provide this output is underway.

The limitations of NOMADS-MVP protocol are counterbalanced by its ability to detect important threats to malaria control and case management with simplicity and rapidity — thereby meeting an urgent public health need. It is feasible to sequence a moderate sized batch of samples (24–64) and use *Nomadic* to generate results within a single day, and with substantially less laboratory manipulation than other approaches. The protocol can be easily implemented even in small laboratories with limited infrastructure, internet connectivity, or prior sequencing experience. Because it takes only one or two days to complete, scientists both process sample sets faster and, in the context of training, gain mastery faster, repeating the protocol and accruing experience quickly. Reducing the set of required reagents makes procurement for the protocol easier, simplifies troubleshooting, and ultimately makes outputs more consistent. Moreover, with a rapid protocol the consequences of a disruption (e.g. by a power outage) are less profound, and scheduling sequencing around other activities is easier. Altogether, these features will accelerate decentralised *P. falciparum* genomic surveillance across sub-Saharan Africa — improving both the quantity and quality of data that are generated, while simultaneously strengthening local capacity and country ownership.

Methods

Sample collection and ethics

All blood samples were collected from patients with *P. falciparum* malaria, with informed consent from the patient or from a parent or guardian. In all cases, consent allowed for the samples to be used for purposes such as this study. This study was conducted using the collected samples only; there was no human subject contact.

All studies received ethical approval from an appropriate research ethics committee: in Senegal, ethical approval for the SEN19/49 study was granted by the Comité National d'Ethique pour la Recherche en Santé (000317/MSAS/CNERS/SP); in Torodo, Mali, ethical approval for a cohort study in 2022 was granted by the Ethics Committee of Charité (EA2/264/21) and the Faculty of Medicine, Pharmacy and Odontostomatology (FMPOS) at the University of Bamako (N°2022/20/CE/USTTB/24.01.22); in Burkina Faso, ethical approval for the AMTIP study was granted by the Comité d'Ethique Institutionnel / Institut National de Santé Publique (2023-10/MSHP/SG/INSP/CEI); in Côte d'Ivoire, ethical approval for the study "Surveillance génomique de Plasmodium falciparum aux CTA à Bouaké, Côte d'ivoire" was granted by the Direction Médicale et Scientifique du aux Centre Hospitalier et Universitaire (192MSHPCMU/CHU-B/DG/DMS/ONAR/24) and by the Ethikkommission der Charité Universitätsmedizin (EA2/171/24); in Nigeria, ethical approval for samples collected in 2023–2024 was granted by the Ethical Committee of Ladoke Akintola University of Technology Teaching Hospital, Ogbomoso (LTH/OGB/EC/2022/304); in Zambia, ethical approval for the 2024 Malaria Indicator Survey (MIS2024) and the 2024–2025 *hrp2/3* Surveillance Study was granted by the Research Ethics Committee at the University of Zambia (5055-20241) and the Tropical Diseases Research Centre Ethics Review Committee (TRC/C4//03/2024), respectively; in Kenya, ethical approval for samples collected in the ATSB study was granted by the KEMRI Scientific and Ethics Review Unit (KEMRI/SERU/CGHR/368/4189; CDC Project ID 0900f3eb81d7ec3c and 0900f3eb82546323); in Ethiopia, ethical approval for the ARSUNA study was granted by the Arsi University institutional review board (AU/HSC/ST-129/5494) and the Federal Democratic Republic of Ethiopia, Ministry of Education (17/256/476/24).

Mock DBS samples

Creation of DBS

For validation of our protocol we created mock *P. falciparum* positive dried-blood spots (DBS) as follows. We cultured the *P. falciparum* laboratory strains 3D7, Dd2 and HB3 at 5% hematocrit in commercially available red blood cells (RBCs) obtained from DRK-Blutspendedienst Nord-Ost gGmbH, as previously described[45]. We brought each culture to ∼5% parasitemia and then synchronised to ring-stage parasites using 5% Sorbitol (PanReac AppliChem, #A2222). After synchronisation, three technicians independently measured the parasitemia of each culture by microscopy and used a Neubauer Counting Chamber to determine the number of RBCs per microlitre. We used the average of the parasitemia and RBCs/μL measurements to standardise each culture to 100,000 parasites/μL at 50% hematocrit. We performed 10-fold serial dilutions of the 100,000 parasites/μL stocks in 80 μL of whole human blood to produce clonal mock samples down to 1 parasites/μL. The DBS were stored at

−20°C in individual plastic bags with a desiccant until use.

DNA extraction

For each sample we used a single (6mm) punch from the DBS for DNA extraction. DNA was extracted by using QIAamp DNA Mini Kit (Qiagen, #51306) according to the manufacturer's instructions with the following modifications: elution was performed by using 2x40 μL of nuclease free water heated to 50 °C. Each elution was incubated for 3 min at room temperature.

NOMADS-MVP Multiplex PCR

Primer design

We designed primers for the NOMADS-MVP (Minimum Viable Panel) amplicon panel using our open-source software *Multiply* (https://github.com/JasonAHendry/multiply). In brief, *Multiply* takes as input a user-specified "design file" defining the target organism, regions of interest, acceptable amplicon size range, and other parameters. From the design file, it generates a large set of candidate primers for each target region and finds an optimal multiplex PCR primer set by minimising primer dimers, offtarget annealing, and primer overlap with known genetic variants. For moderate target numbers (> 20), it typically takes a few minutes to run. For the ten targets of NOMADS-MVP (Table 1), *Multiply* produced a total of 453 candidate primer pairs (median of 47 per target); across which it identified 4,096 potential offtarget binding sites, 77 primers overlapping common variants (>5% minor allele frequency (MAF) in any population defined in the Pf6 Project[46] or resistance-associated codons), and 2,028 high-affinity primer dimers. The greedy search algorithm minimised these factors to select a set of 3 candidate multiplexes (from a theoretical 22 thousand trillion possible combinations) for the ten targets. After pilot sequencing experiments, we selected the multiplex with the highest coverage over all ten targets as our final panel (Supplementary Fig. 1d).

PCR optimisation

We optimised our multiplex PCR using KAPA HiFi HotStart ReadyMix (Roche Diagnostics, #KK2602) formulation which performs well for (A+T)-rich genomes like *P. falciparum*. Using mock samples of varying parasitemia, we searched for a reaction optimum in several full factorial experimental designs. For simplicity, we maintained a 2-step PCR program and 25 μL reaction volume throughout, but explored varying 5 other factors (template DNA amount, number of cycles, primer concentration, polymerase concentration, and annealing temperature). By agarose gel, we observed that increasing the template amount increased sensitivity up to 8 μL after which there was no clear benefit; and that a program with 35 cycles was more sensitive than 30 cycles (Supplementary Fig. 1a). A gradient PCR indicated an ideal annealing temperature of between 58–60 °C (Supplementary Fig. 1b). Increasing the amount of primer increased sensitivity but also increased visible background amplification, whereas increasing the amount of polymerase seemed to increase sensitivity without visibly affecting background (Supplementary Fig. 1c). Using this information, we selected four candidate optimal conditions, and sequenced seven mock samples under each (Supplementary Fig. 1d). From these, we selected the multiplex PCR conditions to maximize the percentage of reads mapping to *P. falciparum* malaria and mean coverage across amplicons. The final conditions were: 95 °C, 3 min; followed by 35-cycles of 98 °C 20 s, 60 °C 3 min; and a final at 60 °C for 10 min.

NOMADS-MVP Sequencing Protocol

The complete laboratory protocol, including materials and primer sequences, is available online at protocols.io (English: *NOMADS-MVP: Rapid Genomic Surveillance of Malaria*; French: *Surveillance Génomique du Paludisme par la méthode Nanopore: Protocole Rapide NOMADS-MVP*). In brief, 8 μL of extracted DNA is used as template in a 25 μL multiplex PCR using 15.5 μL of Kapa HiFi HotStart ReadyMix (Roche Diagnostics, #KK2602) and 1.5 μL of the NOMADS-MVP primer pool. Multiplex PCR products are cleaned using a 0.5X ratio of AMPure XP Beads (Beckman Coulter, A63881) and eluted in 15 μL of nuclease-free water. DNA elute is quantified using the Qubit dsDNA HS Assay Kit (ThermoFisher Scientific, #Q33231) and between 200–800 ng of DNA is taken forward for barcoding and sequencing. We use Rapid Barcoding Kit 96 (SQK-RBK114.96) from Oxford Nanopore Technologies (ONT) following the associated ONT protocol with the following exceptions: we use 1 μL (rather than 1.5 μL) of each barcode during rapid barcoding; after barcoding we purify the pooled samples at a 0.5X ratio and elute in 15 μL; we use 800 ng of DNA for adapter ligation, with a master mix of 1.0 μL of RA and 2.3 μL ABD (rather than 1.5 μL RA, 3.5 μL ADB). All sequencing was done using R10.4.1 flow cells on MinION Mk1B devices.

Exclusion of selective whole genome amplification (sWGA)

We developed a modified version of the selective whole-genome amplification (sWGA) protocol[37] that substituted the phi29 DNA polymerase (New England Biolabs, #M0269S) with EquiPhi29 (ThermoFisher Scientific, #A39390) as follows: 2 μL EquiPhi29 10x buffer, 0.2 μL 0.1M DTT, 2 μL 500 μM sWGA primer pool, 1 EquiPhi29 DNA polymerase, 12.8 μL DNA; 45 °C 1hr, 60 °C 10min; dilute before PCR with 160 μL nuclease-free water. We were able to obtain robust amplification with these conditions despite the substantial reduction in incubation time relative to phi29 DNA polymerase. However, despite both field and mock DBS samples typically yielding stronger bands on an agarose gel if sWGA was used, we saw no consistent improvement in the number of samples passing our quality control ($\geq$8 amplicons with $\geq 50\times$ mean coverage). In other words, samples that pass with using sWGA also passed without sWGA; and using sWGA to process samples did not rescue samples that failed without it. At the same time, we saw evidence that using sWGA skewed the proportions of strains within polyclonal infections, as has been documented previously[22]. Given that it also adds cost and complexity to the protocol, we removed it in August 2024.

Real-time bioinformatics pipeline and dashboard

We developed a real-time bioinformatics pipeline and dashboard for nanopore sequencing, called *Nomadic*. The source code is publicly available on GitHub (https://github.com/JasonAHendry/nomadic) and user documentation is hosted on GitHub Pages (https://jasonahendry.github.io/nomadic/). *Nomadic* was designed for *P. falciparum* genomic surveillance with the NOMADS-MVP protocol, but was coded flexibily and supports other amplicon panels or organisms. Both the bioinformatics pipeline and dashboard are implemented in Python.

Running a real-time analysis

Briefly, to run *Nomadic*, a user first starts a nanopore sequencing run using *MinKNOW*, ensuring to: (1) assign a unique experiment name, hereafter referred to as `<expt_name>`; (2) enable real-time basecalling; (3) enable sample demultiplexing with the appropriate barcoding kit (for NOMADS-MVP, SQK-RBK114.96). The user must prepare a comma-separated values (CSV) metadata table, which maps the barcodes to sample identifiers, and save it in a *Nomadic* workspace folder with the name `<expt_name>.csv`. The user can then launch *Nomadic* from a terminal window with the command `nomadic realtime <expt_name>`.

Implementation and bioinformatics details

As sequencing proceeds, *MinKNOW* writes batches of basecalled reads to disk every ten minutes as FASTQ files. These are demultiplexed by *MinKNOW* such that each barcode possesses its own folder containing associated FASTQ files. *Nomadic* keeps track of the FASTQ files in these folders and, when a new FASTQ file has been generated for a given barcode, launches a per-sample bioinformatics pipeline to process it and update results for the sample. In brief, the pipeline maps reads to the *P. falciparum* 3D7 reference genome (PlasmoDB release 67) using *Minimap2*[47] (v2.28), summarizes the output of mapping (i.e. fraction of reads mapped) and coverage of the target amplicons using *samtools* (v1.17), performs variant calling with *bcftools call*[40] (v1.17), and annotates variants using *bcftools csq* (v1.17). Once a per-sample bioinformatics pipeline has been completed, a set of summary CSV files describing the results across all samples in the experiment is updated to reflect the incorporation of the new reads.

The dashboard runs in a separate thread and creates a graphical summary of these summary CSV files which updates every minute. The plots are interactive and users can hover over features of interest to open tooltips with additional information, zoom into areas of interest, or export static versions of the plots as PNG files. We implemented it using the Dash library (v2.17.1) from plotly. In our experience, sequencing and basecalling are often slower than the time *Nomadic* takes to process batches of FASTQ files, which allows the dashboard to remain up-to-date throughout the sequencing run. Once sequencing is completed, *MinKNOW* and *Nomadic* are stopped by the user. The final summary CSV files contain key results and are sufficient to reopen the dashboard, as described below.

Viewing the dashboard for a previous experiment

The *Nomadic* dashboard can be reopened after a sequencing experiment is completed by running the command `nomadic dashboard <expt_name>` in a terminal window.

Sequencing coverage summary statistics

We used summary files from *Nomadic* to compute per-sample summary statistics of sequencing coverage and evaluate performance across countries and parasitemia levels. In particular, we computed the mean coverage across all amplicons using the `mean_cov` column in the `summary.bedcov.csv` files produced by *Nomadic*. To summarise coverage uniformity across amplicons, we compute a fold-difference between the most- and least-abundant amplicon per sample, excluding the

hrp2/3 amplicons. In particular, defining m_{ij} as the mean coverage in sample i of amplicon j, and a is the set of all amplicons excluding *hrp2* and *hrp3*, then we compute:

$$\frac{\max_{j \in a}(m_{ij})}{\min_{j \in a}(m_{ij})}.$$

Excluding *hrp2* and *hrp3* makes the statistic robust to *hrp2/3* deletions; and in practice, for NOMADS-MVP *hrp2* or *hrp3* are rarely the lowest abundance amplicon in the absence of deletion and so do not contribute to the statistic's value.

Analysis of read lengths

As each experiment typically contains millions of reads, for computational simplicity we included only four experiments in this analysis: from Kenya (sequenced on 2024/07/30), Mali (2024/10/25), Ethiopia (2024/11/27) and a mock sample experiment (2024/11/26). For each experiment, we randomly sampled 20 barcodes (samples) without replacement from the those that: (i) had 8 or more amplicons with greater than $50\times$ coverage; (ii) had >100 parasites/μL; (iii) did not use sWGA; (iv) were not positive or negative controls. For these barcodes, the BAM file generated by *Nomadic* was processed using *Pysam* (v0.22.1) to determine read lengths from the 'query length' field. Genomic regions of interest were delineated with a custom python script that converted gene codon numbers of interest into genomic coordinates using the *P. falciparum* 3D7 reference genome (PlasmoDB release 67) and associated general feature format (GFF) file, and then were manually verified using the Integrative Genome Viewer (IGV) (v2.5.0). We determined the percentage of reads overlapping by first using the intersect command from *bedtools*[48] (v2.31) which enabled us to filter the BAM file to only reads overlapping the region of interest; and then comparing the number of mapped reads in this BAM file against the original BAM file using *samtools* (v1.17).

Inference of *hrp2/3* deletions

We used a set of 219 previously published samples from Asella, Ethiopia to evaluate concordance between NOMADS-MVP and a gold-standard approach for *hrp2/3* deletion detection[43]. Recommended[49] conventional PCRs for *hrp2* and *hrp3*[50] were performed in duplicate and visualised by agarose gel electrophoresis. For discordant results, we called a sample positive if there was a clear band of the appropriate length in at least one of the replicates[49]. We excluded from analysis 46 samples that failed a *kelch13* PCR, suggesting low DNA quality or parasitemia and 24 samples that contained other *Plasmodium* species, leaving 149 samples. Of the 149 samples, 138 (92.6%) had 8 or more amplicons exceeding $50\times$ coverage after NOMADS-MVP nanopore sequencing and were used for the comparison.

We predicted *hrp2/3* deletions from NOMADS-MVP sequencing data using a previously developed Bayesian statistical model[32]. In brief, it works as follows. For each sequencing run, the statistical model first estimates the rate of background contamination / barcode misclassification using included negative controls. Next, it estimates the quality of each sample and average variation in quality across samples using the mean coverage over all amplicons in the multiplex PCR, excluding the *hrp2* and *hrp3* amplicons. Finally, across all samples the posterior probability of *hrp2* and *hrp3* deletion are estimated separately using Markov Chain Monte Carlo (MCMC). Here, each chain was run for 50,000 iterations with a prior deletion probability of 0.5 for *hrp2* and *hrp3*. Although a lower prior might be justified for hrp2 given prevalence of deletions in the region, in practice

our estimates were insensitive to the prior. For Figure 5a, we directly show the posterior probabilities from the model. For Figure 5b, we are showing the *maximum a posteriori* (MAP) estimate of deletion. The statistical model and Bayesian MCMC were implemented in Python.

SNP calling accuracy evaluation

Methodology for evaluating SNP calling accuracy with respect to sequencing coverage was described in a previous publication[32]. Below we provide a brief summary.

Downsampling reads.

We randomly downsample our sequencing data to a specific mean coverage equally over each amplicon using *samtools*[51] (v1.22). In particular, we downsample BAM files for each amplicon with the `samtools view` command and `-s/--subsample` flag to achieve the intended mean coverage, ranging from 10–100×, ensuring to sample a different random seed for each replicate. The procedure was repeated for all six mock samples. For the Cambodian strains, only data over *Kelch13* was used to evaluate the F_1-score. In Figure 4b, the 'All' category is computed across the entire amplicon (left pane) panel or coding sequencing (CDS, right pane).

Creating a set of true variants.

3D7, Dd2, HB3 have been sequenced using Pacific Bioscience Sequencing SMRT technology and assembled with a FASTA sequences available on *PlasmoDB*[52]. To identify variants in these assemblies with respect to the 3D7 reference genome, we simulated high-quality (Phred 60) error-free reads *in silico* from the FASTA files, mapped them to the 3D7 reference genome with *minimap2* (v2.28), and then identified variants using the *bcftools*[40] (v1.22) `mpileup` and `call` commands. We simulated 60 error-free reads, half forward and half reverse strand, for each amplicon in our NOMADS-MVP panel by extracting the FASTA sequence spanning +/-4 kbp of the target, based on GFF files for 3D7, Dd2 and HB3.

For each Cambodian strain, information about the *Kelch13* amino acid changes were available from BEI resources (IPC3445, C580Y; IPC4912, I543T; IPC5202, R539T); however, we found no information about corresponding nucleotide changes. Therefore, we performed Sanger sequencing using the NOMADS-MVP *Kelch13* amplicon as a singleplex PCR to determine the *Kelch13* nucleotide sequence for each strain. We used *tracy*[53] to convert the chromatogram file into a VCF with the `tracy decompose` command.

Evaluating accuracy.

We used *hap.py*[42] from a *Docker* image (jmcdani20/hap.py:v0.3.12) to compute measures of variant calling accuracy. We used the *P. falciparum* 3D7 GFF file (release 67) to define coding regions for each gene. In *hap.py*, the F_1-score is computed as the harmonic mean of the precision and recall. For amplicons where there are no variants for a particular mock sample (e.g. *Kelch13* for 3D7, Dd2 and HB3), the precision and recall are considered to be undefined; unless false-positive mutations are present, in which case the precision is zero.

across different target regions in comparison to the true variant set described above. To restrict accuracy measure analysis

to coding sequence, we subset the 3D7 GFF to only CDS features, identified the rows pertaining to our targets, and output the chromosome, start, and stop positions as a BED file. We used then used the `--stratifications` flag of *hap.py* to compute measures over these intervals. We used the annotated VCF files produced by *hap.py* to generate positional plots of false- and true-positive rate across targets in Python.

References

1. World Health Organization (WHO). *World Malaria Report 2024*. World Health Organization, Geneva (2024).

2. Menegon, M. *et al.* Identification of *Plasmodium falciparum* isolates lacking histidine-rich protein 2 and 3 in Eritrea. *Infect. Genet. Evol.* **55**, 131–134, DOI: 10.1016/j.meegid.2017.09.004 (2017).

3. Berhane, A. *et al.* Major Threat to Malaria Control Programs by Plasmodium falciparum Lacking Histidine-Rich Protein 2, Eritrea - Volume 24, Number 3—March 2018 - Emerging Infectious Diseases journal - CDC. DOI: 10.3201/eid2403. 171723.

4. Funwei, R. *et al.* Molecular surveillance of pfhrp2 and pfhrp3 genes deletion in Plasmodium falciparum isolates and the implications for rapid diagnostic tests in Nigeria. *Acta Trop.* **196**, 121–125, DOI: 10.1016/j.actatropica.2019.05.016 (2019).

5. Golassa, L., Messele, A., Amambua-Ngwa, A. & Swedberg, G. High prevalence and extended deletions in Plasmodium falciparum hrp2/3 genomic loci in Ethiopia. *PLOS ONE* **15**, e0241807, DOI: 10.1371/journal.pone.0241807 (2020).

6. Feleke, S. M. *et al.* Plasmodium falciparum is evolving to escape malaria rapid diagnostic tests in Ethiopia. *Nat. Microbiol.* **6**, 1289–1299, DOI: 10.1038/s41564-021-00962-4 (2021).

7. Ariey, F. *et al.* A molecular marker of artemisinin-resistant Plasmodium falciparum malaria. *Nature* **505**, 50–55, DOI: 10.1038/nature12876 (2014).

8. Miotto, O. *et al.* Genetic architecture of artemisinin-resistant Plasmodium falciparum. *Nat. Genet.* **47**, 226–234, DOI: 10.1038/ng.3189 (2015).

9. World Health Organization (WHO). *Strategy to respond to antimalarial drug resistance in Africa.* World Health Organization, Geneva (2022).

10. Uwimana, A. *et al.* Emergence and clonal expansion of in vitro artemisinin-resistant Plasmodium falciparum kelch13 R561H mutant parasites in Rwanda. *Nat. Medicine* **26**, 1602–1608, DOI: 10.1038/s41591-020-1005-2 (2020).

11. Balikagala, B. *et al.* Evidence of Artemisinin-Resistant Malaria in Africa. *New Engl. J. Medicine* **385**, 1163–1171, DOI: 10.1056/NEJMoa2101746 (2021).

12. van Loon, W. *et al.* Escalating Plasmodium falciparum K13 marker prevalence indicative of artemisinin resistance in southern Rwanda. *Antimicrob. Agents Chemother.* **68**, e01299–23, DOI: 10.1128/aac.01299-23 (2023).

13. Rosenthal, P. J. *et al.* The emergence of artemisinin partial resistance in Africa: how do we respond? *The Lancet Infect. Dis.* **24**, e591–e600, DOI: 10.1016/S1473-3099(24)00141-5 (2024). Publisher: Elsevier.

14. Fola, A. A. *et al.* Plasmodium falciparum resistant to artemisinin and diagnostics have emerged in Ethiopia. *Nat. Microbiol.* **8**, 1911–1919, DOI: 10.1038/s41564-023-01461-4 (2023).

15. Eloff, L. *et al.* High Prevalence of Molecular Markers Associated with Artemisinin, Sulphadoxine and Pyrimethamine Resistance in Northern Namibia, DOI: 10.1101/2025.01.09.25320247 (2025).

16. Fola, A. A. *et al.* Temporal genomic analysis of Plasmodium falciparum reveals increased prevalence of mutations associated with delayed clearance following treatment with artemisinin-lumefantrine in Choma District, Southern Province, Zambia, DOI: 10.1101/2024.06.05.24308497 (2024).

17. Aranda-Díaz, A. *et al.* Plasmodium falciparum genomic surveillance reveals a diversity of kelch13 mutations in Zambia, DOI: 10.1101/2025.02.19.25322554 (2025).

18. World Health Organization (WHO). *Response plan to pfhrp2 gene deletions, second edition*. World Health Organization, Geneva (2024).

19. World Health Organization (WHO). *Multiple first-line therapies as part of the response to antimalarial drug resistance*. World Health Organization, Geneva (2024).

20. Zupko, R. J. *et al.* Modeling policy interventions for slowing the spread of artemisinin-resistant pfkelch R561H mutations in Rwanda. *Nat. Medicine* **29**, 2775–2784, DOI: 10.1038/s41591-023-02551-w (2023).

21. Jacob, C. G. *et al.* Genetic surveillance in the Greater Mekong subregion and South Asia to support malaria control and elimination. *eLife* **10**, e62997, DOI: 10.7554/eLife.62997 (2021).

22. LaVerriere, E. *et al.* Design and implementation of multiplexed amplicon sequencing panels to serve genomic epidemiology of infectious disease: A malaria case study. *Mol. Ecol. Resour.* **22**, 2285–2303, DOI: 10.1111/1755-0998.13622 (2022).

23. Aranda-Díaz, A. *et al.* Sensitive and modular amplicon sequencing of Plasmodium falciparum diversity and resistance for research and public health. *Sci. Reports* **15**, 10737, DOI: 10.1038/s41598-025-94716-5 (2025).

24. Verity, R. *et al.* The impact of antimalarial resistance on the genetic structure of Plasmodium falciparum in the DRC. *Nat. Commun.* **11**, 2107, DOI: 10.1038/s41467-020-15779-8 (2020).

25. Niaré, K. *et al.* Highly multiplexed molecular inversion probe panel in Plasmodium falciparum targeting common SNPs approximates whole-genome sequencing assessments for selection and relatedness. *Front. Genet.* **16**, DOI: 10.3389/fgene.2025.1526049 (2025).

26. Loman, N. J. & Watson, M. Successful test launch for nanopore sequencing. *Nat. Methods* **12**, 303–304, DOI: 10.1038/nmeth.3327 (2015).

27. Runtuwene, L. R. *et al.* Nanopore sequencing of drug-resistance-associated genes in malaria parasites, Plasmodium falciparum. *Sci. Reports* **8**, 8286, DOI: 10.1038/s41598-018-26334-3 (2018).

28. Sabin, S. *et al.* Portable and cost-effective genetic detection and characterization of Plasmodium falciparum hrp2 using the MinION sequencer. *Sci. Reports* **13**, 2893, DOI: 10.1038/s41598-022-26935-z (2023).

29. Girgis, S. T. *et al.* Drug resistance and vaccine target surveillance of Plasmodium falciparum using nanopore sequencing in Ghana. *Nat. Microbiol.* **8**, 2365–2377, DOI: 10.1038/s41564-023-01516-6 (2023).

30. Holzschuh, A. *et al.* Using a mobile nanopore sequencing lab for end-to-end genomic surveillance of Plasmodium falciparum: A feasibility study. *PLOS Glob. Public Heal.* **4**, e0002743, DOI: 10.1371/journal.pgph.0002743 (2024).

31. Sadler, J. M. *et al.* Application of a new highly multiplexed amplicon sequencing tool to evaluate Plasmodium falciparum antimalarial resistance and relatedness in individual and pooled samples from Dschang, Cameroon. *medRxiv* DOI: 10.1101/2024.10.03.24314715 (2024). Pages: 2024.10.03.24314715.

32. de Cesare, M. *et al.* Flexible and cost-effective genomic surveillance of P. falciparum malaria with targeted nanopore sequencing. *Nat. Commun.* **15**, 1413, DOI: 10.1038/s41467-024-45688-z (2024).

33. Cheng, Q. *et al.* Plasmodium falciparum parasites lacking histidine-rich protein 2 and 3: A review and recommendations for accurate reporting. *Malar. J.* **13**, 283, DOI: 10.1186/1475-2875-13-283 (2014).

34. White, M. T. *et al.* Immunogenicity of the RTS,S/AS01 malaria vaccine and implications for duration of vaccine efficacy: Secondary analysis of data from a phase 3 randomised controlled trial. *The Lancet Infect. Dis.* **15**, 1450–1458, DOI: 10.1016/S1473-3099(15)00239-X (2015).

35. Datoo, M. S. *et al.* Efficacy of a low-dose candidate malaria vaccine, R21 in adjuvant Matrix-M, with seasonal administration to children in Burkina Faso: A randomised controlled trial. *The Lancet* **397**, 1809–1818, DOI: 10.1016/S0140-6736(21)00943-0 (2021).

36. World Health Organization (WHO). *Report on antimalarial drug efficacy, resistance and response: 10 years of surveillance (2010-2019)*. World Health Organization, Geneva (2020).

37. Oyola, S. O. *et al.* Whole genome sequencing of Plasmodium falciparum from dried blood spots using selective whole genome amplification. *Malar. J.* **15**, 597, DOI: 10.1186/s12936-016-1641-7 (2016).

38. Otto, T. D. *et al.* Long read assemblies of geographically dispersed plasmodium falciparum isolates reveal highly structured subtelomeres. *Wellcome Open Res* **3**, 52, DOI: 10.12688/wellcomeopenres.14571.1 (2018).

39. Straimer, J. *et al.* Drug resistance. k13-propeller mutations confer artemisinin resistance in plasmodium falciparum clinical isolates. *Science* **347**, 428–31, DOI: 10.1126/science.1260867 (2015).

40. Li, H. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinforma. (Oxford, England)* **27**, 2987–2993, DOI: 10.1093/bioinformatics/btr509 (2011).

41. Hall, M. B. *et al.* Evaluation of Nanopore sequencing for Mycobacterium tuberculosis drug susceptibility testing and outbreak investigation: a genomic analysis. *The Lancet Microbe* **4**, e84–e92, DOI: 10.1016/S2666-5247(22)00301-9 (2023). Publisher: Elsevier.

42. Krusche, P. *et al.* Best practices for benchmarking germline small-variant calls in human genomes. *Nat. Biotechnol.* **37**, 555–560, DOI: 10.1038/s41587-019-0054-x (2019).

43. Tufa, T. B. *et al.* Plasmodium falciparum Kelch13 Variants and Histidine-Rich Protein 2/3 Deletions in Central Ethiopia, 2023. *The Am. J. Trop. Medicine Hyg.* **-1**, DOI: 10.4269/ajtmh.25-0114 (2025). Publisher: The American Society of Tropical Medicine and Hygiene Section: The American Journal of Tropical Medicine and Hygiene.

44. Schaffner, S. F. *et al.* Malaria surveillance reveals parasite relatedness, signatures of selection, and correlates of transmission across Senegal. *Nat. Commun.* **14**, 7268, DOI: 10.1038/s41467-023-43087-4 (2023). Publisher: Nature Publishing Group.

45. Trager, W. & Jensen, J. B. Human malaria parasites in continuous culture. *Science* **193**, 673–5, DOI: 10.1126/science.781840 (1976).

46. , M. *et al.* An open dataset of Plasmodium falciparum genome variation in 7,000 worldwide samples [version 2; peer review: 2 approved]. *Wellcome Open Res.* **6**, DOI: 10.12688/wellcomeopenres.16168.2 (2021).

47. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100, DOI: 10.1093/bioinformatics/bty191 (2018).

48. Quinlan, A. R. & Hall, I. M. Bedtools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842, DOI: 10.1093/bioinformatics/btq033 (2010).

49. Parr, J. B., Anderson, O., Juliano, J. J. & Meshnick, S. R. Streamlined, PCR-based testing for pfhrp2- and pfhrp3-negative Plasmodium falciparum. *Malar. J.* **17**, 137, DOI: 10.1186/s12936-018-2287-4 (2018).

50. Baker, J. *et al.* Genetic Diversity of Plasmodium falciparum Histidine-Rich Protein 2 (PfHRP2) and Its Effect on the Performance of PfHRP2-Based Rapid Diagnostic Tests. *The J. Infect. Dis.* **192**, 870–877, DOI: 10.1086/432010 (2005).

51. Li, H. *et al.* The sequence alignment/map format and samtools. *Bioinformatics* **25**, 2078–2079, DOI: 10.1093/bioinformatics/btp352 (2009).

52. Aurrecoechea, C. *et al.* PlasmoDB: a functional genomic database for malaria parasites. *Nucleic Acids Res.* **37**, D539–D543, DOI: 10.1093/nar/gkn814 (2009).

53. Rausch, T., Fritz, M. H.-Y., Untergasser, A. & Benes, V. Tracy: basecalling, alignment, assembly and deconvolution of sanger chromatogram trace files. *BMC Genomics* **21**, 230, DOI: 10.1186/s12864-020-6635-8 (2020).

Acknowledgements

The NOMADS project is funded by the Bill Melinda Gates Foundation (INV-048316 to M.M., D.J.B. and J.A.H.). Sample collection and *hrp2/3* deletion investigation in Ethiopia was supported by the ARSUNA project, funded by the German Federal

Ministry for Economic Cooperation and Development and the Else Kröner-Fresenius Foundation (EKFS) via the Hospital Partnerships Programme (project 21Ac01060). We thank Bob Verity for helping to estimate the number of samples required to detect mutations spreading across Africa with adequate power; Olaf Kostbahn, Wendy Vienneau, and Sossena Assefa for critical financial and legal administration; and Estée Török for fruitful scientific discussions, feedback and encouragement throughout the project's duration. We are grateful to all those who participated in the studies conducted across Africa, especially the health workers and patients who made the project possible.

Author contributions statement

M.M.: Conceptualization, Methodology, Validation, Investigation, Data curation, Funding acquisition. K.M.: Methodology, Validation, Investigation. B.B.: Methodology, Software, Validation, Formal analysis. M.C.: Investigation. W.V.L.: Investigation, Resources, Data curation. B.M.: Investigation. A.G.: Investigation. A.O.: Investigation. S.S.: Investigation. N.J.Y.: Investigation. D.S.: Investigation. B.N.: Data curation. D.Z.: Investigation. Y.N.G.T.: Investigation. F.O.: Investigation, Data curation. A.Z.: Investigation. E.A.A.: Investigation. C.C.: Investigation. S.O.: Investigation. B.O.: Investigation. M.N.: Resources. D.K.G.: Resources. T.F.: Resources. T.B.T.: Resources. R.O.: Investigation. E.S.: Investigation. Y.S.: Resources. C.N.: Resources. O. Ouedraogo.: Resources. K.O.: Resources. O.Opaleye.: Resources. A.Olowe.: Resources. M.G.: Project administration. C.D.: Supervision. G.S.: Resources, Data curation, Supervision. F.P.M.: Resources, Supervision. S.P.: Resources, Supervision. A.D.: Resources, Supervision. I.S.: Resources, Supervision. D.N.: Resources, Supervision. O.Ojurongbe.: Resources, Supervision. S.K.: Resources, Supervision. Y.P.S.: Conceptualization, Resources, Data curation, Supervision. J.S.S.: Resources, Data curation, Supervision. M.H.: Resources, Supervision. D.J.B.: Conceptualization, Data curation, Supervision, Funding acquisition. J.A.H.: Conceptualization, Methodology, Software, Validation, Formal analysis, Data curation, Supervision, Funding acquisition, Writing - Original Draft. All authors reviewed and approved the final version of the manuscript.

Additional information

The authors declare no competing interests. **CDC Disclaimer:** The opinions expressed by the authors do not necessarily reflect the official views of the CDC or the U.S. Department of Health and Human Services.