Comparative analysis of shared and unique mechanisms important for diverse strains

of *Pasteurella multocida* to cause systemic infection in mice

Thomas R. Smallman[1], Xiaochu Wang[1], Xenia Kostoulias[1,2,3], Amy Wright[1], Faye C.

Morris[1,2], Marina Harper[1†] and John D. Boyce[1†#]

[1] Infection and Immunity Program, Monash Biomedicine Discovery Institute and Department

of Microbiology, Monash University, Clayton, Victoria

[2] Centre to Impact AMR, Monash University, Clayton, Victoria, Australia

[3] Department of Infectious Diseases, Alfred Health and School of Translational Medicine,

Monash University, Melbourne, Victoria, Australia

[#] Corresponding author: John Boyce, john.boyce@monash.edu

[†] Joint senior authors

Running title: TraDIS investigation of *P. multocida* pathogenesis

Abstract

Pasteurella multocida is a Gram-negative bacterium that causes a range of distinct diseases

in livestock animals. Different *P. multocida* diseases are associated with different capsule

and lipopolysaccharide (LPS) types, but little else is known about what underpins this

disease specificity. In this study, we utilised transposon-directed insertion site sequencing

(TraDIS) to identify genes in two diverse *P. multocida* strains that are required for growth in

rich media, and genes important for survival during systemic infections in BALB/c mice.

Analysis of growth in heart infusion broth showed that the fowl cholera isolate VP161

(capsule type A and LPS type L1) and the bovine haemorrhagic septicaemia isolate M1404

(capsule type B and LPS type L2) shared 461 genes essential for growth in rich media, with

95% of these present in all publicly available closed *P. multocida* genomes. *In vivo* fitness

analysis identified 63 and 94 genes important for VP161 and M1404 survival in BALB/c

mice, respectively. Only 35 homologs were identified as important for survival in both strains,

showing that conserved biological systems can be differentially important for different

P. multocida strains. Investigation of proteins involved in the catabolite response showed that

an active cyclic-adenosine monophosphate (cAMP) receptor protein (CRP) was required for

maximal fitness in M1404. Furthermore, disrupting CRP or cAMP production also reduced

capsule production in M1404, but increased capsule production in VP161, demonstrating

that these *P. multocida* strains have different regulatory systems for crucial virulence factors.

Author Summary

Pasteurella multocida is an important livestock pathogen, causing several distinct severe

diseases in many different livestock animals. *P. multocida* can spread rapidly throughout

animal populations, with peracute infections causing death within 48 h, resulting in large

outbreaks with high mortality. Host predilection and disease presentation often correlate with

the capsule and lipopolysaccharide type produced by the causative strain. However, the

processes that allow certain strains to cause a particular disease are not well understood. In

this study, we have comprehensively identified genes required for two *P. multocida* strains to

cause systemic infection in mice, showing that these diverse strains have differential

requirements for survival during a systemic infection. This information is crucial for

understanding *P. multocida* diseases and for the development of new strategies to combat

infection.

Introduction

Pasteurella multocida is an upper respiratory tract commensal of many mammal and bird

species [1-3], but the bacterium can also cause serious disease, including fowl cholera in

birds, haemorrhagic septicaemia in ungulates, atrophic rhinitis in pigs, and pneumonia in

most livestock animals [4]. Strains of *P. multocida* are commonly classified based on capsule

type (namely A, B, D, E and F) and/or lipopolysaccharide (LPS) genotype (namely L1-L9) [5-

7]. *P. multocida* disease presentation and host predilection often correlates with genotype of

surface structures produced by the infecting strain [4]. For instance, fowl cholera is caused

primarily by strains with capsule type A or F, and LPS type L1, L3 or L6 [5, 8-10], while

haemorrhagic septicaemia is only caused by strains with capsule type B or E, and LPS type

L2 or L5 [11-13]. Apart from capsule and LPS type, very little is known about the pathogenic

mechanisms that are unique to strains causing these different diseases or what pathogenic

mechanisms are shared by such strains.

P. multocida disease often progresses rapidly and death can occur within 48 h of infection

[14, 15], making antibiotic treatment in these cases rarely successful. *P. multocida* is readily

transmitted between animals within a population, either directly or via environmental

contamination. Large *P. multocida* disease outbreaks occur in both wild and livestock animal

populations [16-20], with increased susceptibility linked to environmental stresses such as

increased temperature and humidity, as well as animal crowding [21]. Outbreaks of

P. multocida disease in livestock result in high mortality and large economic losses.

Vaccination is an important preventative measure against *P. multocida* disease. However,

while several *P. multocida* vaccines are currently available [22], most only provide protection

against strains producing the same capsule and LPS structures as the vaccine strain and the

duration of immunity is generally short [23]. Currently, there is no vaccine available that

protects against the full range of diseases in different hosts. We predict that to improve the

breadth of protection of *P. multocida* vaccines, it is important to understand what pathogenic

mechanisms are shared or unique across *P. multocida* strains that cause distinct diseases.

In this study, we investigated the pathogenic mechanisms shared between two different

P. multocida strains, VP161, a chicken fowl cholera isolate belonging to capsule serotype A

and LPS genotype L1, and M1404, a bison haemorrhagic septicaemia isolate belonging to

capsule serotype B and LPS genotype L2. These strains are genetically diverse, but both

can cause lethal systemic infection in mice [24-26]. We performed transposon-directed

insertion site sequencing (TraDIS, also called Tn-Seq) in both strains to identify all genes in

each strain that were essential for growth on rich media, and those genes important for *in*

vivo fitness in a mouse systemic infection model. The TraDIS analysis identified a core-set of

447 genes that we predict are essential for the *P. multocida* species. We also identified

shared pathways and processes important for both strains VP161 and M1404 to survive *in*

vivo (35 genes), and processes important only for strain VP161 (28 genes) or only for strain

M1404 (59 genes) to survive *in vivo*. These data identify multiple unique requirements for

each of these strains to cause disease in mice.

Results

Production of a *P. multocida* strain M1404 *Himar1* mutant library, systemic infections
in mice, and TraDIS library production

We previously constructed a *Himar1* transposon mutant library in the *P. multocida* strain

VP161 (capsule type A and LPS type L1) [27], which was also used in this study. For TraDIS

analysis of strain M1404 (capsule type B and LPS type L2), we generated a comparable

Himar1 transposon mutant library. A total of ~1.12 x 10^6 M1404 *Himar1* transposon mutants

were pooled to produce the final library. To generate a high-quality M1404 reference genome

for TraDIS analysis, we used a combination of Nanopore and Illumina sequencing that

allowed for the assembly of a 2.42 Mbp genome that contained two contigs (see S1_Text for

full assembly methodology).

The M1404 and VP161 *Himar1* mutant libraries were each used to infect BALB/c mice at a

dose of ~2 x 10^7 CFU via intraperitoneal injection (project overview described in S1 Fig and

infectious doses are given in S1 Table; note that all Supplemental Tables, Supplemental

Figures and Supplemental Figure Legends are in S1_Text). All mice succumbed to systemic

infection, reaching humane endpoints at five hours (S2A Fig). Surviving *P. multocida* mutants

were recovered from the blood, liver, and spleen by plating *in vivo*-derived samples onto rich

media. Greater than 4 x 10^6 CFU of *P. multocida* were recovered from each site for all

infections (S2B Fig). Four TraDIS library replicates were generated using genomic DNA

isolated from the recovered mutants representing each strain (VP161 and M1404) and

condition (rich media input culture, recovered mutants from each tissue following infection in

mice). Each library was sequenced on an Illumina NextSeq and analysed using the Bio-

TraDIS pipeline and related scripts (full TraDIS data in S1_Data). Combined, the four TraDIS

library replicates generated for each strain and condition had high transposon-DNA junction

read coverage across the genome and a small bp distance between unique insertion sites

(UIS) (S3 Fig); the VP161 libraries had between 31 and 33 bp distance between each UIS,

and the M1404 libraries had between 40 and 43 bp between each UIS (S2 Table in

S1_Text). Principal component analysis using UIS or reads per gene showed that in all

conditions the rich media TraDIS libraries clustered together and separated away from the

mouse TraDIS libraries (S4 Fig).

Identification of core genes essential for growth in rich media

We used the input mutant libraries, grown on solid heart infusion media overnight, to identify

genes essential for growth in rich media for both *P. multocida* strain VP161 and M1404.

Essential genes were identified by comparing normalised UIS counts per gene (Fig 1A, S5

Fig, see S2_Data for the list of all essential genes for growth on rich media). There were 585

genes, of which 542 were coding sequences (CDS), identified as essential for strain VP161.

There were also 585 genes, of which 540 were CDS, essential for strain M1404. Although a

similar number of genes were identified as essential in both strains, only 461 were homologs

(Fig 1B), as determined using eggNOG or BlastP. The majority of these 461 shared essential

proteins were involved in housekeeping processes. Analysis using the STRING database

identified several processes and pathways that were enriched in the essential protein set,

including: DNA replication (false discovery rate [FDR] = 4 x 10^{-4}), transcription (FDR =

0.047), translation (FDR = 2.04 x 10^{-16}) including ribosomal structural proteins (FDR = 2.7 x

10^{-5}) and tRNA amino acylation (FDR = 7.83 x 10^{-5}), cell division and regulation of cell shape

(FDR = 0.015), fatty acid biosynthesis (FDR = 4 x 10^{-6}), peptidoglycan (FDR = 0.004) and

lipid A biosynthesis (FDR = 0.019), glycolysis (FDR = 0.026), purine nucleoside biosynthesis

(FDR = 0.0026), heme/porphyrin biosynthesis (FDR = 0.027), isoprenoid biosynthesis (FDR

= 0.027), and ATP production (FDR = 0.0033).

We then assessed if these shared essential genes were also conserved across the entire

P. multocida species. We analysed 167 publicly available complete *P. multocida* genomes,

together with VP161 and M1404, and identified a total of 1,564 genes present in all strains

(the 100% core-genome). A core-genome phylogeny generated from this set of 169

complete genomes was highly similar to a previously constructed *P. multocida* core-genome

phylogeny generated using 402 *P. multocida* strains that included complete and incomplete

genomes (S6 Fig) [7], indicating that this subset of 169 strains is representative of the entire

P. multocida species. Of the 461 essential genes identified in both strain VP161 and strain

M1404, 447 (97%) were present in the 100% core-genome (Fig 1B). We also compared the

100% core-genome and conserved rich media essential genes to our curated *P. multocida*-

specific virulence factor and antibiotic resistance database (PastyVRDB) [7]. This analysis

identified 53 genes associated with virulence that are present in all *P. multocida* strains, and

included the genes *hpbA*, *oma87*, *spoT*, *fur*, Pm0442, and *kdtA*, which were all essential in

both strains (S3 Table in S1_Text).

There were 75 genes essential in VP161 that had a non-essential homolog in M1404, and

65 genes essential in M1404 that had a non-essential homolog in VP161 (Fig 1B). Genes

identified as essential in VP161 but not M1404 included *crp*, *crr* and *relA*, and genes

identified as essential in M1404 but not VP161 included several genes involved in the Tol-

Pal system (*tolBQR* and *pal*), DNA repair (*ruvBC* and *aroQ*), and sugar transport (*ptsH*).

There were also six essential VP161 genes that had no homolog in M1404, and 14 essential

genes in M1404 that had no homolog in VP161 (Fig 1B). Most of the unique essential M1404

genes encoded hypothetical proteins; however, M1404_00858 and M1404_00859 contained

antitoxin-like domains, and M1404_01327 contained a putative peptidase toxin domain.

**

P. multocida **strains VP161 and M1404 have both shared and unique genes required**

for virulence in mice

To identify genes that were important for *P. multocida* fitness during murine systemic

infections, we compared the read count per gene between the mouse and rich media TraDIS

libraries. Any gene with a four-fold decrease in *Himar1* insertions and a *q*-value < 0.01 was

identified as resulting in decreased *in vivo* fitness when disrupted, and any gene with a four-

fold increase in *Himar1* insertions and a *q*-value < 0.01 was identified as showing increased

in vivo fitness when disrupted (see S3_data and S4_data for full fitness-associated data).

Overall, 134 genes were *in vivo* fitness-associated in at least one type of tissue and one

strain (Table 1, Fig 2A). There were 63 and 94 genes that resulted in decreased *in vivo*

fitness of strain VP161 and M1404, respectively (Fig 2B, S4 Table in S1_Text). For both

VP161 and M1404, most of these genes were identified as important for *in vivo* fitness in all

three tissues (Fig 2B). The genes were then compared to identify homologs between the two

strains. Of the genes that had a homolog in the other strain, 35 were important for fitness in

both strains, 23 were important for fitness only in VP161, and 47 were important for fitness

only in M1404 (Fig 2C). Furthermore, there were five genes important for fitness in VP161

that did not have a homolog in M1404, and 12 genes important for fitness in M1404 that did

not have a homolog in VP161 (Fig 2C). In total, there were seven genes that increased

fitness when disrupted in either VP161 or M1404, with only two genes, *yeeZ* and *speB*,

common to both strains (S5 Table in S1_Text).

Common pathways and processes were identified as important for both VP161 and M1404

during infection in mice (Fig 3A, S7 Fig and S8 Fig). In both strains, disruption of almost all

capsule biosynthesis genes had a severe fitness-cost. Both *fis* and *hfq*, which encode

transcriptional regulators known to positively regulate capsule biosynthesis genes in strain

VP161, were important for fitness in VP161 but not in M1404, suggesting these strains used

different regulatory mechanisms. Several genes encoding proteins involved in LPS core-

polysaccharide synthesis were identified as important for *in vivo* fitness. Genes from the

ADP-L-glycero-β-D-manno-heptose synthesis pathway (*hldDE-gmhAB*), that encode

proteins that synthesise a modified heptose that is added to the LPS core-polysaccharide,

were either important for *in vivo* fitness or essential in both strains. The gene *hptA*, which

encodes an LPS inner-core heptosyltransferase, was identified as important for fitness in

both strains. Additionally, genes encoding the LPS inner-core heptosyltransferase genes

hptC and *hptD*, the LPS inner-core glucosyltransferase *gtcD*, and the LPS outer-core

galactosyltransferase *gatD* were important for fitness of strain M1404. Several genes

identified as important for *in vivo* fitness in both strains encoded proteins involved in amino

acid synthesis and transport, or protein translation (Fig 3). These included an

aminotransferase involved in aspartate, phenylalanine, and tyrosine synthesis (*aspC*), an

asparagine synthase (*asnA*) and regulator of asparagine synthase expression (*asnC*), a

putative alanine transporter (*alsT_1*), and proteins involved in translation efficiency (*prmC*

and *epmA*) (Fig 3). Two genes in the L-serine biosynthesis pathway (*serAB*) and a putative

tyrosine transporter (*tyrP_1*) were important for the *in vivo* fitness of VP161 but not M1404.

Conversely, elongation factor Tu subunit genes (*tufB_1* and *tufB_2*) were important for the *in*

vivo fitness of strain M1404 but not strain VP161. Other systems important for *in vivo* fitness

in both strains included the purine biosynthesis pathway, the Ton-B energy transduction

system (*tonB-exbBD*), methionine uptake (*plpB-metPN*), and *aroA* from the chorismate

pathway.

Several pathways were identified as important for *in vivo* fitness of M1404 but not VP161

(Fig 3A). This included genes encoding two components of the sialic acid uptake system

(*nanP* and *nanU*), sialic acid metabolism (*nanE*) and the N-acetylneuraminate CMP

transferase (*neuA_1*), a sensor histidine kinase (*cpxA*), the sorbitol-specific EIIA (*srlB*), and

genes encoding the Mla phospholipid transport system (*mlaBDEF*). The genes encoding

adenylate cyclase (*cyaA*) and cAMP receptor protein (*crp*) were also identified as important

for *in vivo* fitness of M1404 but not VP161; however, *crp* was identified as essential for strain

VP161 to grow in rich media. The gene M1404_01811, which encodes a *P. multocida*

capsule type B- and E-specific surface lipoprotein (SLP) PmSLP-3 [28], and the adjacent

gene M1404_01812 that encodes a surface lipoprotein assembly modulator (Slam) required

for PmSLP-3 export, were both identified as important for M1404 *in vivo* fitness.

Comparatively, other putative SLPs (PlpE and PmVP161_1666 in strain VP161,

M1404_01814 in strain M1404), and the Slam exporter in VP161 (PmVP161_1663) were not

important for *in vivo* fitness. Most of the genes important for *in vivo* fitness of M1404 that did

not have a homolog in VP161 were required for production of the type B capsule polymer

(Fig 3B). Genes required for *in vivo* fitness of VP161 but not for M1404 included genes

encoding components of the ubiquinone biosynthesis pathway (*ubiGHE_3X*), a heme

utilisation protein (*hutZ_1*), components of the aminodeoxychorismate synthase complex

(*pabAB*), a putative sodium proton antiporter (PmVP161_0183), a putative tyrosine importer

(*tyrP_1*)*,* and *aroE* from the chorismate pathway (Fig 3A). All of the genes important for *in*

vivo fitness in VP161 that did not have a homolog in M1404 were required for production of

the type A capsule polymer (Fig 3C).

The genes *alsT_1*, *crp* and *cyaA* are important for strain M1404 fitness *in vivo*

We performed time to euthanasia experiments with selected individual *P. multocida* directed

mutants to confirm that a sub-set of genes identified by TraDIS were individually important

for *in vivo* survival. ClosTron mutagenesis was used to generate individual *alsT_1*, *crp* and

cyaA mutants in strain M1404. Wild-type M1404 and each mutant strain was used to

inoculate separate groups of BALB/c mice via intraperitoneal injection on two different days

(see S1 Table in S1_Text for infectious doses). Signs of systemic infection were monitored

and used to determine time to euthanasia. Mice infected with wild-type M1404 reached the

humane endpoint on average 6.5 hours post infection (hpi) (Fig 4). Mice infected with the *crp*

or *cyaA* mutants reached the humane endpoint at an average 11 and 10.3 hpi, respectively

(Fig 4). Mice infected with the *alsT_1* mutants did not reach the humane endpoint and

showed low total clinical scores over the 12-hour infection period (Fig 4). All strains had a

significant increase in time to euthanasia when compared to wild-type M1404, confirming the

TraDIS data that *alsT_1*, *crp* and *cyaA* are important for *in vivo* fitness of strain M1404.

The carbon catabolite repression system controls capsule production differently in
***P. multocida* strains VP161 and M1404**

Carbon catabolite repression was important for *P. multocida* fitness and is controlled by the

cyclic-adenosine monophosphate (cAMP) receptor protein (CRP, also called catabolite

activator protein (CAP) or virulence factor regulator (VFR) in other species). CRP was

important for *in vivo* fitness of strain M1404, and essential for strain VP161 to grow in rich

media. While CRP has not been well characterised in *P. multocida*, in other bacterial species

CRP family proteins are global transcriptional regulators that have increased affinity for DNA

targets when bound to cAMP [29]. Thus, CRP activity is modulated by the intracellular levels

of cAMP. The level of intracellular cAMP is controlled by the adenylate cyclase CyaA, which

synthesises cAMP from ATP [30], and the cAMP phosphodiesterase CpdA, which degrades

cAMP to AMP [31]. As shown above, both *crp* and *cyaA* were identified as important for *in*

vivo fitness of M1404 by TraDIS analysis and individual mutants in these genes showed

reduced virulence in mice. Furthermore, the TraDIS data showed that inactivation of *cpdA* in

M1404 slightly improved fitness during *in vivo* growth (average $\log_2$ fold-change of between

0.93 and 1.34, see S1_data). Collectively, these data suggest that cAMP-dependent

activation of CRP is important for M1404 *in vivo* fitness. CRP family proteins are known to

regulate a wide range of targets in different bacterial species [32-34], and transcriptomic

analysis of a *P. multocida* strain 0818 *crp* mutant identified reduced capsule biosynthesis

gene expression [35]. To determine if inactivating the catabolite repression system affected

capsule production in M1404, capsule production was assessed using cells grown overnight

on solid media. Both the M1404 *crp* and M1404 *cyaA* mutants harbouring empty vector

produced barely detectable absorbance readings, indicating that these strains are essentially

acapsular (Fig 5A). Providing the *crp* mutant with a wild-type copy of *crp in trans* led to a

modest, but significant increase in absorbance readings, compared to the *crp* mutant

harbouring empty vector (Fig 5A). Providing the *cyaA* mutant with the *cyaA* complementation

plasmid restored absorbance readings, and therefore capsule production, to wild-type levels

(Fig 5A). These data confirm that adenylate cyclase is required for capsule production in

M1404 and strongly suggest that this is a CRP-dependent mechanism.

The role of active CRP and catabolite repression in VP161 was different from that observed

in M1404. While *crp* was essential for strain VP161 growth in rich media, the TraDIS data

showed that inactivation of *cyaA* had no impact on *in vivo* fitness of VP161, and inactivation

of *cpdA* slightly reduced fitness (average log$_2$ fold-change of between -1.75 and -1.93, see

S1_Data). In our previous TraDISort study, *cpdA* was identified as important for *in vitro*

capsule production in strain VP161 [27]. Collectively, these data suggest that active CRP-

cAMP may act to repress capsule production in strain VP161. As CRP was essential in

VP161, we attempted to inactivate *cpdA* and *cyaA* using ClosTron mutagenesis; however,

only a *cyaA* mutant was successfully generated. Capsule absorbance assays were

performed using wild-type VP161 and the *cyaA* mutant harbouring empty vector, and the

complemented *cyaA* mutant. Capsule was extracted from cells during early stationary phase,

when cAMP levels would be expected to be high. The VP161 *cyaA* mutant produced

significantly more capsule than wild-type VP161, with complementation restoring normal

levels of capsule production (Fig 5). These data indicate that inactivation of adenylate

cyclase, which should reduce the level of cAMP and therefore CRP activity, de-represses

capsule production in VP161. This strongly suggests that an active CRP-cAMP represses

capsule in strain VP161, which is the opposite of the role of CRP-cAMP in M1404.

Discussion

In this study, we performed TraDIS analysis of two distinct *P. multocida* strains, the poultry

isolate VP161 (A:L1) and the bovine isolate M1404 (B:L2). These strains produce different

capsule and LPS structures. Initially, we identified genes required for both VP161 and

M1404 to grow in rich media, with 585 genes identified as essential in both strains. We have

previously reported that 509 genes were essential for VP161 to grow in rich media [27], 95%

of which were also identified as essential in this study. The increase in rich media essential

genes is likely due to the mutant libraries being grown for ~48 h longer than the previous

study, allowing for further reduction of mutants with reduced viability from the population.

Previous analysis of random transposon mutant viability has shown that the time required for

a mutation to lead to a replication defect determines at what time point the gene is identified

as important for survival [36]. Strains VP161 and M1404 shared a set of 461 protein-

encoding genes that were essential for growth in rich media, most of which were involved in

housekeeping processes. Pan-genome analysis of 169 complete *P. multocida* genomes

identified 1,564 genes present in all genomes. Of these 1,564 core genes, 447 were

identified as essential in both strain VP161 and strain M1404 for growth in rich media. Thus,

97% of the 461 essential genes identified in strains VP161 and M1404 are present in all

P. multocida genomes examined and are likely to be required for all *P. multocida* strains to

grow in rich media. While most of these genes encode housekeeping genes involved in

replication, transcription, translation and energy metabolism, some encode surface-

associated proteins and therefore could be considered potential targets for vaccine

development.

TraDIS analysis of VP161 and M1404 mutants recovered from BALB/c mice following

systemic infection identified 134 genes associated with *in vivo* fitness in at least one tissue

and one strain. To validate these results, we generated directed *alsT_1*, *crp* and *cyaA*

mutants in strain M1404. Separate infection studies with these mutants showed reduced

virulence in mice, confirming each is important for *in vivo* fitness. Additionally, several genes

and pathways identified in the TraDIS analysis performed in this study have previously been

identified as important for *P. multocida* virulence. Infection studies conducted using mice or

poultry have shown that inactivation of *aroA* in *P. multocida* strains X-73, P-1059, and 85020

[37, 38], *tonB*, *exbB*, and *exbD* in strain PM25 [39], *plpB* in strain VP161 [40], and *nanP* and

nanU in strain P-1059 [41] all resulted in reduced virulence. Furthermore, several

P. multocida aroA mutants are commercially available for use as live-attenuated vaccines

[37, 38]. Signature-tagged mutagenesis studies using strain UC6731 and strain VP161

identified *purN*, *purF, guaB*, *nanE* and *nanU* as important for virulence in mice and/or

chickens [24, 42]. Transcriptomics analysis of strain X-73 cells recovered from systemic

infections in chickens showed there was increased expression *in vivo* of *purH*, *aspC* and

asnA [43], and transcriptomic analysis of Pm70 grown in media with limited iron showed

increased expression of *hutZ_1* [44]. These previous studies strongly support that our

TraDIS analysis has comprehensively identified genes required for *P. multocida* survival *in*

vivo. However, surface structures that have previously been proposed to play a role in

P. multocida in vivo fitness and/or virulence, including filamentous haemagglutinin, fimbriae,

and Flp-pili, were not identified as important for *in vivo* fitness in our TraDIS analysis [24, 42,

45, 46]. These surface structures are likely required for initial colonisation of the upper

respiratory tract but not when *P. multocida* is delivered via intramuscular, intraperitoneal or

intravenous injection. Indeed, *P. multocida* filamentous haemagglutinin mutants have a

higher infectious dose (ID_{50}) compared to wild-type parent strains in birds but only when

introduced intranasally and not via intramuscular or intravenous route [45, 46]. The TraDIS

data presented in this study identified several processes that have not previously been

shown to be important for *P. multocida* disease, including ubiquinone biosynthesis,

aminodeoxychorismate synthesis, the Mla phospholipid transport system and putative

sodium proton antiporters. Importantly, we showed that there were clear differences between

pathways and processes required for the two different *P. multocida* strains to survive *in vivo*.

The ubiquinone biosynthesis pathway and the putative sodium proton antiporter

(PmVP161_0183) were only important for *in vivo* fitness of strain VP161. Conversely,

components of the sialic uptake and utilisation pathway, the Mla ABC-transport system, the

adenylate cyclase CyaA, the SLP protein PmSLP-3 and its cognate Slam exporter were only

important for *in vivo* fitness of strain M1404. This crucially shows that although these

P. multocida strains share these systems, their importance for survival *in vivo* is strain

dependent.

Different *P. multocida* strains typically encode two or more SLPs that are exported to the cell

surface by a Slam protein [28, 47]. Subunit vaccines formulated using *P. multocida* SLPs

(PlpE, PmSLP-3 and PmSLP-1) have been shown to elicit a protective immune response

against *P. multocida* strains that produce the cognate SLP [28, 48-50]. Despite both VP161

and M1404 encoding two SLPs, our TraDIS analysis only identified PmSLP-3 in M1404 as

important for survival in BALB/c mice. PmSLP-1, an SLP that is encoded by a subclade of

capsule type A strains isolated from cattle, has recently been shown to bind to complement

factor I and block complement-mediated killing [51]. Previous studies have shown that

capsule was important for blocking complement-mediated killing in strain VP161 but not for

strain M1404 [25, 26]. As such, it is possible that PmSLP-3 is important for blocking

complement-mediated killing in M1404. Several SLPs from *Neisseria*, *Actinobacillus*,

Hemophilus and *Moraxella* spp. have a similar predicted protein fold to *P. multocida* SLPs

but have diverse functions, including binding to human iron-binding proteins, lactoferrin and

factor H [47]. Thus, despite SLPs having similar structure, each protein can have a different

biological function and role in pathogenesis. Given *P. multocida* can persist in several

different hosts, it is possible that different SLAM-dependent SLPs are required for survival in

different hosts.

Most capsule biosynthesis genes and known regulators of capsule biosynthesis were

identified as important for *in vivo* fitness, supporting many previous studies demonstrating

that capsule is important for *P. multocida* virulence in both type A and B strains [25, 26, 42,

52-54]. Additionally, several other genes previously identified in strain VP161 as important

for capsule production were also identified by TraDIS as important for *in vivo* fitness,

including *prmC*, *epmA*, *epmB* and *ubiX*. However, our data indicated that capsule production

is regulated differently in the two strains. The genes encoding the global regulator Fis and

the RNA chaperone protein Hfq, previously shown to be important for capsule production in

strain VP161 [53, 54], were also identified in this study as important for *in vivo* fitness of

VP161 but not M1404. Furthermore, the TraDIS analysis and capsule absorbance assays

together showed that CRP-cAMP controls capsule production in these *P. multocida* strains,

but in different ways. Disrupting CRP-cAMP by inactivating either *cyaA* or *crp*, abrogated

capsule production in strain M1404, showing that CRP-cAMP positively regulates capsule

production in this strain. Conversely, disrupting CRP-cAMP by inactivating *cyaA* increased

capsule production in strain VP161, indicating that CRP-cAMP represses capsule production

in strain VP161. Together, these data show that capsule expression responds differently to

CRP-cAMP in these two strains. The TraDIS data also suggested that activation of the

catabolite response in strain M1404 may depend on the level of sorbitol. Adenylate cyclase

activity is commonly controlled by phosphoenolpyruvate(PEP):carbohydrate

phosphotransferase system (PTS) proteins. The PTS system couples carbohydrate import

and phosphorylation, with the energy required for import and the donated phosphate

provided by PEP [55, 56]. The phosphate is transferred to the carbohydrate via a protein

phosphate transfer chain consisting of the general PTS proteins EI and HPr, and the sugar

specific EII, with EII generally comprised of three proteins/domains (EIIA, EIIB, EIIC) [57].

Different proteins in the PTS system can allosterically regulate different protein targets

depending on whether they are phosphorylated. In *E. coli*, phosphorylated EIIA[Glc] is known

to be involved in inducing adenylate cyclase activity [58, 59], thus allowing cAMP production

and activation of the catabolite repressor system when the levels of glucose are low. EIIA[Glc],

encoded by *crr*, was not important for *in vivo* fitness of M1404. However, the sorbitol-specific

EIIA protein, encoded by *srlB*, was identified as important for *in vivo* fitness of M1404. This

suggests that the absence of sorbitol increases the amount of phosphorylated sorbitol-

specific EIIA, which in turn induces adenylate cyclase activity and activates the catabolite

response.

Several LPS biosynthesis genes were important for fitness in both strains. The LPS of

P. multocida lacks an O-antigen, making the core-polysaccharide the most distal component

of the LPS [60]. The core-polysaccharide is comprised of a conserved inner-core and a

variable outer core, and is synthesised by stepwise addition of sugars by different

glycosyltransferases [60]. As the function of LPS transferases have been determined for LPS

type L1 and L2 strains [61-64], the putative structures required for survival in BALB/c mice

can be predicted based on which transferases had reduced fitness in the TraDIS analysis. All

P. multocida strains examined to date simultaneously produce two inner-core structures

(namely glycoforms A and B), with the relative levels of each glycoform determined by the

competing activity of KdkA and KdtA. KdkA adds a phosphate to the first 3-deoxy-D-manno-

oct-2-ulosonic acid (Kdo) leading to glycoform A production, while KdtA adds a Kdo II to Kdo

I leading to glycoform B production [61]. The nascent inner-core is then extended by the

heptosyltransferases HptA (specific for glycoform A) and HptB (specific for glycoform B) [61].

The TraDIS analysis identified *hptA*, but not *hptB*, as important for fitness in both strain

VP161 and M1404. Inactivation of *hptA*, but not *hptB*, has previously been shown to

attenuate virulence of strain VP161 in chickens [61]. The VP161 *hptA* mutants have severely

truncated LPS as the Lipid A – phosphorylated Kdo I cannot be extended in this mutant [61].

The presence of a large amount of severely truncated LPS on the surface make the

bacterium vulnerable to the innate immune system [61]. No other LPS transferases were

identified as important for fitness of VP161, supporting previous data that showed a full

length LPS is not important for fitness of this strain in mice [24]. In strain M1404, the inner-

core transferases *hptC*, *hptD* and *gctA* were all important for fitness, indicating that a

glycoform A structure of at least Kdo I - Hep I [Hep II – Hep III] – Glucose (Glc) I is required

for *in vivo* survival. Additionally, inactivation of the outer-core galactosyltransferase gene

gatD significantly reduced fitness of M1404 in BALB/c mice. The LPS outer-core structure

produced by M1404 is Hep IV – Hep V – N-acetyl glucosamine (GlcNAc) I – Galactose (Gal)

I [64]. Identification of *gatD* but not other outer-core transferases as important for fitness

suggests that removal of the terminal galactose therefore leaving the GlcNAc as the terminal

sugar reduces the viability of M1404 in BALB/c mice, but truncations terminating the outer-

core at either Hep IV, Hep V or GlcNAc I do not impact fitness. Previous analysis of several

P. multocida type L3 strains has shown that inactivating mutations in transferase genes that

result in different LPS outer-core structures affect virulence in chickens at different levels

[65].

In this study, we have determined the 100% core-genome for *P. multocida* and identified a

set of core genes likely essential for all *P. multocida* strains. Furthermore, we have

comprehensively identified genes important for *in vivo* fitness of the type A strain VP161 and

the type B strain M1404 during systemic infections in BALB/c mice, with these strains having

shared and unique pathways that are important for fitness. Genes identified as important for

both strains to survive *in vivo* and conserved across the entire *P. multocida* species

represent ideal vaccine targets for generating a vaccine that provides protection against all

P. multocida strains. Finally, we have shown that capsule production is regulated differently

between these two strains, and that a full length LPS is more important in M1404 than

VP161 for *in vivo* fitness in mice.

Materials and methods

Bacterial strains and growth conditions

The bacterial strains and plasmids used in this study are listed in S6 Table (in S1_Text).

E. coli strains were cultured in lysogeny broth (LB), and *P. multocida* strains were cultured in

heart infusion (HI) broth (Oxoid). Solid media was generated by adding 1.5% agar (w/v) to

liquid media. When required, antibiotics were added to the media at the following

concentrations: kanamycin 50 µg/mL, spectinomycin 50 µg/mL, tetracycline 1.25 µg/mL,

ampicillin 100 µg/mL. When required, diaminopimelic acid (DAP) was added to media at a

concentration of 100 µM.

Oligonucleotides, DNA extraction, DNA manipulation and DNA sequencing

All oligonucleotides used in this study were synthesised by Sigma, and the sequences of all

oligonucleotides are given in S7 Table (in S1_Text). Genomic DNA was extracted using the

HiYield Genomic DNA Mini Kit (Real Biotech Corporation) as per the manufacturer's

instructions for extraction from Gram-negative bacteria or extracted using Genomic Tip

100/G (QIAGEN) as per manufacturer's instructions. Plasmids were extracted using the

NucleoSpin Plasmid Mini Kit (Machery-Nagel) as per the manufacturer's instructions.

Polymerase chain reactions (PCR) were performed using either *Taq* DNA polymerase

(Roche) or Phusion High-Fidelity DNA polymerase (New England Biolabs), as per the

manufacturer's instructions for each polymerase. DNA digests were performed using

restriction enzymes sourced from New England Biolabs as per manufacturer's instructions.

Ligations were performed using T4 DNA ligase (Roche) as per manufacturer's instructions.

DNA was purified using the NucleoSpin Gel and PCR Clean-up Mini Kit (Machery-Nagel).

Sanger sequencing reactions were performed as previously described [65]. For whole-

genome sequencing of *P. multocida* strain M1404, genomic DNA was submitted to The

Ramaciotti Centre for Genomics (University of New South Wales) and sequenced using both

an Illumina MiSeq v3, and a Nanopore GridION X5 Mk1 using a GridION flow cell (FLO-

MIN112) with DNA prepared for sequencing using a Native Barcoding Expansion Kit (SQK-

NBD112.24). Additionally, M1404 genomic DNA was also submitted to Plasmidsaurus and

sequenced using a Nanopore PromethION using a PromethION flow cell (FLO-PRO114M)

with DNA prepared for sequencing using a Rapid Barcoding Kit (SQK-RBK114.96). The

M1404 genome was assembled as previously described [7]; see supplementary materials for

additional information.

Transformation, site-directed mutagenesis using ClosTron and construction of

complementation plasmids

Electrocompetent *P. multocida* cells were prepared and transformed as previously described

[25]. *P. multocida* site-directed mutants were generated by a modified version of the

ClosTron system (also called TargeTron) in the pAL953 plasmid as previously described [65,

66]. Intron insertions into each of the genes of interest were confirmed by PCR-amplification

using primers that flanked the gene of interest, and by Sanger sequencing directly from

genomic DNA extracted from putative mutants with oligonucleotide BAP6544, which binds

within the intron and directs sequencing into the adjacent region of the genome. For

complementation, the wild-type gene was amplified using Phusion polymerase and cloned

into pAL99S under the control of the *tpiA* promoter. The sequence of the cloned gene and

promoter were confirmed by Sanger sequencing. Complementation plasmids or pAL99S

empty vector were used to transform the corresponding mutant strains to produce the

complemented mutant and empty vector strains, respectively.

Conjugation conditions and generation of the *P. multocida* strain M1404 *Himar1* transposon mutant library

Conjugation between an *E. coli* donor and *P. multocida* recipient was performed by filter mating as previously described [27], with minor modifications. The donor used in this study was *E. coli* strain JKE201 [67], a diaminopimelic acid (DAP) auxotrophic strain that cannot grow without the addition of DAP in the media. Filters containing the conjugation mix were incubated on HI agar plates containing DAP for one hour. The conjugation mix was recovered from the filter by vortexing the filter in 1 mL of HI broth. The conjugation mix was plated onto LB agar with ampicillin and DAP to recover donor *E. coli*, HI agar to recover all *P. multocida* recipients, and HI agar with spectinomycin to recover *P. multocida Himar1* mutants. *P. multocida Himar1* transconjugants were recovered from initial selection plates after overnight growth and replated on fresh HI agar containing spectinomycin to ensure selection against mutants with insertions in essential genomic regions, and to increase culture density for library storage. Following the second overnight growth, the *P. multocida Himar1* mutants were recovered in glycerol broth (30% glycerol in HI broth) and stored at -80°C. A total of ~1.12×10^6 M1404 *Himar1* mutants were pooled to generate the final M1404 mutant library.

Mouse systemic infections

Animal experiments were performed according to Victorian State Government regulations, approved by the Monash University Animal Ethics Committee (Monash University AEC no. MARP/2023/36738). Each *P. multocida* systemic infection was performed with an equal number of male and female 6–10-week-old BALB/c mice (housed separately) supplied by Monash Animal Research Platform (Monash University), with mice housed in the Animal Research Laboratory (Monash University). *P. multocida* strains used for systemic infections were grown in HI medium to early-exponential phase (OD_{600} ~0.25). Mice were each injected intraperitoneally using a 27-gauge needle with a 200 μL total volume containing one of the following: ~1×10^7 CFU of the *P. multocida* VP161 transposon mutant library, ~1×10^7 CFU

of the *P. multocida* strain M1404 transposon mutant library, ~5 x 10^4 CFU of wild-type strain

M1404, or ~1 x 10^5 CFU of one of the directed M1404 mutants (S1 Table in S1_Text). Mice

were monitored using a clinical score monitoring system for systemic infections in mice [68];

mice were monitored hourly and reached humane endpoints when they had a clinical score

of 4 in a single category, or a cumulative clinical score of 10 or more. Mice were humanely

euthanised using CO_2 asphyxiation followed by cervical dislocation. Blood samples were

collected via cardiac puncture, and liver and spleen samples were recovered and

homogenized. Tissue and blood samples were plated at the appropriate dilutions onto HI

agar to obtain single colonies and determine the bacterial burden in the bloodstream or

tissue. For TraDIS analysis, to ensure there were enough cells for genomic DNA extraction,

blood and homogenized tissue samples were plated neat onto heart infusion agar and grown

overnight before bacteria were recovered. Samples of the VP161 and M1404 *Himar1* mutant

libraries used to inoculate the mice were also plated neat onto HI agar in the same way as

the bacteria recovered from mice. For DNA isolation, bacterial samples recovered from solid

media using 1 x PBS were centrifuged at 13,000 x *g* then the cell pellet was resuspended 1

x PBS to wash cells before being centrifuged again as above. The cell pellets generated

were then used to extract genomic DNA.

Transposon-directed insertion site sequencing library production, Illumina
sequencing and bioinformatic analysis of TraDIS libraries

TraDIS libraries were prepared for Illumina sequencing as previously described (29). TraDIS

libraries were sequenced by The Ramaciotti Centre for Genomics at the University of New

South Wales using an Illumina NextSeq1000 P1 with 100 bp-single end reads. TraDIS data

was analysed using the Bio-TraDIS toolkit and modified scripts, and replicates compared by

principal component analysis, as previously described [27]. Briefly, genes essential for

growth in rich media were identified by comparing normalized UIS per gene plotted on a

histogram. Genes important for fitness were identified by comparison of read count per gene

between each mouse TraDIS library and the input rich medium TraDIS library. Genes with

log_2 fold-change < -2 and a *q*-value < 0.01 were called as having a fitness cost, and genes

with a $\log_2$ fold-change > 2 and a *q*-value < 0.01 as having a fitness benefit. Full data sheets

are provided as Supplemental Material S1_Data, S2_Data, S3_Data and S4_Data.

To identify protein homologs between strains, all coding sequences from *P. multocida* strains

VP161 and M1404 were analysed using eggNOG mapper 5.0 [69] and then matched by

seed ortholog value; proteins with the same seed ortholog value were taken as homologs.

Proteins without a match in eggNOG and non-coding sequences were compared using the

Basic Local Alignment Search Tool (BLAST) [70], with homologs called when there was

>90% amino acid/nucleotide identity over >90% of the query. Venn diagrams were generated

using https://eulerr.co/. Heatmaps were generated in R using the ggplot2 package [71].

String was used to identify genes involved in the same pathway or processes, and to identify

pathways and processes enriched in each dataset [72]. Roary analysis was performed using

the VP161 and M1404 genomes, and an additional 167 complete *P. multocida* genomes

obtained from the BV-BRC database (in April 2024). Pan-genome analysis and maximum-

likelihood core-genome phylogenies were generated using Roary and IQ-TREE as

previously described [7].

Capsule absorbance assays

Capsule absorbance assays were performed as previously described [7]. For assessing

capsule production in strain VP161 and derivative strains, cells were cultured in liquid broth

and samples taken to assess capsule production at early stationary phase (7 h, OD_{600} ~2.2).

For strain VP161, the amount of hyaluronic acid was determined against a standard curve of

absorbance values obtained using different amounts of purified hyaluronic acid (Sigma

catalogue number H5388). For strain M1404 and derivative strains, cells were cultured

overnight on heart infusion agar and then recovered by adding 1 x PBS to the plate and

scraping. As purified polysaccharide equivalent to the *P. multocida* type B capsule is not

commercially available to use as a standard, the absorbance values were normalised

against the optical density values of the culture used for each capsule extraction. Statistical

comparisons between strains were performed using a Mann-Whitney U-test.

Data availability

The M1404 genome was submitted to Genbank with the biosample accession number

SAMN49532731. The TraDIS data analysis output files have been uploaded to the NCBI

Gene Expression Omnibus and are available under accession number GSE300598.

Supporting information

S1_Text: Additional materials and methods and supporting tables, figures and figure legends.

S2_Text: Newick format maximum-likelihood core-genome phylogeny of 169 *P. multocida* strains.

S3_Text: Annotated assembly of *P. multocida* strain M1404 used for TraDIS analysis in this study.

S1Fig: S1 Figure

S2Fig: S2 Figure

S3Fig: S3 Figure

S4Fig: S4 Figure

S5Fig: S5 Figure

S6Fig: S6 Figure

S7Fig: S7 Figure

S8Fig: S8 Figure

S9Fig: S9Figure

S1_Data: All TraDIS data from this study.

S2_Data: TraDIS data for genes identified as essential for *P. multocida* strains VP161 and M1404 growth in rich media.

S3_Data: TraDIS data for genes identified as having an *in vivo* fitness-cost.

S4_Data: TraDIS data for genes identified as having an *in vivo* fitness-benefit.

Acknowledgements

We would like to acknowledge the Monash Bioinformatics Platform for assistance with

bioinformatic analysis. We would like to acknowledge Abbey Allgood and Irene Alevizos for

assistance provided while plating bacteria during the mouse infections.

594 **Tables**

595 **Table 1.** Number of genes identified in *P. multocida* strain VP161 or M1404 that when

596 disrupted by a *Himar1* transposon insertion resulted in decreased or increased *in vivo*

597 fitness.

	VP161				M1404			
	Blood	Liver	Spleen	Total	Blood	Liver	Spleen	Total
Decreased fitness	59	57	50	63	85	75	74	94
Increased fitness	4	6	6	7	6	5	4	7

Figures

Fig 1. Identification and comparison of genes essential for *P. multocida* strains VP161 and

M1404 to grow in rich media. Both the VP161 and M1404 *Himar1* mutant libraries were

grown to mid-exponential phase in heart infusion (HI) broth, before being plated on HI agar.

Viable mutants were recovered, used to generate TraDIS libraries that were sequenced by

Illumina sequencing, and the data analysed using Bio-TraDIS and related scripts. **A.**

Insertion indexes (number of unique *Himar1* transposon insertion sites divided by gene

length) for all genes following growth in HI broth. The cut-off for essentiality is marked by a

grey line for each dataset. **B.** Comparison of essential protein encoding genes present in

both strain VP161 and M1404, or only one of the strains. Protein homologs between strains

were identified using eggNOG matches, or by having > 90% amino acid identity. A 100%

P. multocida core-genome was identified in strains VP161 and M1404 along with 167

complete, publicly available *P. multocida* genomes using Roary, and compared using a >

90% amino acid identity cut-off to the essential proteins identified in both strains VP161 and

M1404.

Fig 2. Identification and comparison of genes associated with *in vivo* fitness for *P. multocida* strain VP161 and M1404 during systemic infections in mice. The *P. multocida* strain VP161 and M1404 *Himar1* libraries were individually used in systemic infections in BALB/c mice. Surviving mutants were recovered from the bloodstream, liver and spleen, and then plated on heart infusion agar. Genomic DNA was extracted from surviving mutants and used to generate TraDIS libraries. TraDIS libraries were sequenced via Illumina sequencing and the data analysed using Bio-TraDIS and related scripts. **A.** Volcano plots showing the $\log_2$ fold-change in read count per gene and $-\log_{10}$ *q*-value for all genes in the bloodstream (red), liver (blue), and spleen (gold) TraDIS libraries compared to the rich media TraDIS library. Genes were identified as *in vivo* fitness-associated if they had a four-fold increase or decrease in read count, with a *q*-value of < 0.01 (cut-offs shown as grey lines). **B.** Venn diagrams showing the number of genes identified as important for *in vivo* fitness that were shared between the bloodstream, liver, and spleen TraDIS libraries. **C.** Comparison of *in vivo* fitness genes between strain VP161 and M1404. Protein homologs between strains were identified using matching eggNOG IDs, or by having > 90% amino acid identity and > 90% coverage.

Fig 3. Heat maps showing the log$_2$ fold-change in reads for *in vivo* fitness-associated genes.

The log$_2$ fold-change in reads per gene was identified using the fitness.comparison R-script,

using the rich media TraDIS data as the control, and each of the bloodstream, liver, and

spleen TraDIS libraries as the condition. Genes were included in the heatmap if they were

identified as *in vivo* fitness-associated in any one of the TraDIS libraries. The gene *clcA*,

which is known not to impact *in vivo* fitness [61], was included as a control. **A.** *In vivo* fitness-

associated genes that are present in both strains. For genes that are annotated with different

names or are identified only by different locus tags, both names or locus tags are given, with

the VP161 name or locus tag first followed by the M1404 name or locus tag. For the locus

tags, the prefix (PmVP161_ or M1404_) has been omitted. **B.** *In vivo* fitness-associated

genes in M1404 with no homolog in VP161. **C.** *In vivo* fitness-associated genes in VP161

with no homolog in M1404.

Fig 4. Kaplan-Meier survival curves for mice infected with *P. multocida* wild-type M1404 or

an M1404 directed mutant. Groups of six BALB/c mice (three male and three female) were

inoculated via intraperitoneal injection with wild-type strain M1404, the M1404 *alsT_1*

mutant, the M1404 *crp* mutant, or the M1404 *cyaA* mutant (colony forming units ranged

between 2 x 10^4 and 2 x 10^5). Statistical differences in survival between strains was

determined using a log-rank Mantel-Cox test; *** *P* < 0.001.

Fig 5. Capsule produced by VP161 and M1404 wild-type and mutant strains. Capsule

absorbance assays were performed by mixing capsule extracted from cells with stains-all

solution and measuring the absorbance at 640 nm. **A.** Capsule measured from wild-type

M1404 harbouring empty vector, the M1404 *crp* mutant and M1404 *cyaA* mutant harbouring

either empty vector or the appropriate complementation plasmid. Capsule was extracted

from cells recovered from overnight growth on heart infusion agar. **B.** Capsule measured

from wild-type VP161 harbouring empty vector, and the VP161 *cyaA* mutant harbouring

empty vector or an expression plasmid containing an intact copy of the VP161 *cyaA* gene.

Capsule was extracted from cells grown in heart infusion broth to early stationary phase (7

h). Error bars represent mean ± standard error of the mean, and statistical differences in

capsule production between samples were determined using Mann-Whitney U-tests; * *P* <

0.05.

References

1. Dewhirst FE, Klein EA, Thompson EC, Blanton JM, Chen T, Milella L, et al. The canine oral microbiome. PLoS One. 2012;7:e36067. doi:10.1371/journal.pone.0036067.

2. Lowe BA, Marsh TL, Isaacs-Cosgrove N, Kirkwood RN, Kiupel M, Mulks MH. Defining the "core microbiome" of the microbial communities in the tonsils of healthy pigs. BMC Microbiol. 2012;12:20. doi:10.1186/1471-2180-12-20.

3. Muhairwa AP, Christensen JP, Bisgaard M. Investigations on the carrier rate of *Pasteurella multocida* in healthy commercial poultry flocks and flocks affected by fowl cholera. Avian Pathol. 2000;29:133-42. doi:10.1080/03079450094162.

4. Wilkie IW, Harper M, Boyce JD, Adler B. *Pasteurella multocida*: diseases and pathogenesis. Curr Top Microbiol Immunol. 2012;361:1-22. doi:10.1007/82_2012_216.

5. Harper M, John M, Turni C, Edmunds M, St Michael F, Adler B, et al. Development of a rapid multiplex PCR assay to genotype *Pasteurella multocida* strains by use of the lipopolysaccharide outer core biosynthesis locus. J Clin Microbiol. 2015;53:477-85. doi:10.1128/jcm.02824-14.

6. Townsend KM, Boyce JD, Chung JY, Frost AJ, Adler B. Genetic organization of *Pasteurella multocida* cap loci and development of a multiplex capsular PCR typing system. J Clin Microbiol. 2001;39:924-9. doi:10.1128/jcm.39.3.924-929.2001.

7. Smallman TR, Perlaza-Jiménez L, Wang X, Korman TM, Kotsanas D, Gibson JS, et al. Pathogenomic analysis and characterization of *Pasteurella multocida* strains recovered from human infections. Microbiol Spectr. 2024;12:e0380523. doi:10.1128/spectrum.03805-23.

8. Hirsh DC, Jessup DA, Snipes KP, Carpenter TE, Hird DW, McCapes RH. Characteristics of *Pasteurella multocida* isolated from waterfowl and associated avian species in California. J Wildl Dis. 1990;26:204-9. doi:10.7589/0090-3558-26.2.204.

9. Shivachandra SB, Kumar AA, Gautam R, Saxena MK, Chaudhuri P, Srivastava SK. Detection of multiple strains of *Pasteurella multocida* in fowl cholera outbreaks by polymerase chain reaction-based typing. Avian Pathol. 2005;34:456-62. doi: 10.1080/03079450500367963.

10. Strugnell BW, Dagleish MP, Bayne CW, Brown M, Ainsworth HL, Nicholas RA, et al. Investigations into an outbreak of corvid respiratory disease associated with *Pasteurella multocida*. Avian Pathol. 2011;40:329-36. doi: 10.1080/03079457.2011.571659.

11. Orynbayev M, Sultankulova K, Sansyzbay A, Rystayeva R, Shorayeva K, Namet A, et al. Biological characterization of *Pasteurella multocida* present in the saiga population. BMC Microbiol. 2019;19:37. doi: 10.1186/s12866-019-1407-9.

12. Davies RL. Genetic diversity among *Pasteurella multocida* strains of avian, bovine, ovine and porcine origin from England and Wales by comparative sequence analysis of the 16S rRNA gene. Microbiol. 2004;150:4199-210. doi: 10.1099/mic.0.27409-0.

13. Biswas A, Shivachandra SB, Saxena MK, Kumar AA, Singh VP, Srivastava SK. Molecular variability among strains of *Pasteurella multocida* isolated from an outbreak of haemorrhagic septicaemia in India. Vet Res Commun. 2004;28:287-98. doi: 10.1023/b:verc.0000026656.77847.5b.

14. Petersen KD, Christensen JP, Permin A, Bisgaard M. Virulence of *Pasteurella multocida* subsp. *multocida* isolated from outbreaks of fowl cholera in wild birds for domestic poultry and game birds. Avian Pathol. 2001;30:27-31. doi: 10.1080/03079450020023168.

15. Wilkie IW, Grimes SE, O'Boyle D, Frost AJ. The virulence and protective efficacy for chickens of *Pasteurella multocida* administered by different routes. Vet Microbiol. 2000;72:57-68. doi: 10.1016/s0378-1135(99)00187-x.

16. Woo YK, Kim JH. Fowl cholera outbreak in domestic poultry and epidemiological properties of *Pasteurella multocida* isolate. J Microbiol. 2006;44:344-53. doi.

17. Blackall PJ, Pahoff JL, Marks D, Fegan N, Morrow CJ. Characterisation of *Pasteurella multocida* isolated from fowl cholera outbreaks on turkey farms. Aust Vet J. 1995;72:135-8. doi: 10.1111/j.1751-0813.1995.tb15033.x.

18. Crawford R, Allwright D, He C. High mortality of Cape Cormorants (Phalacrocorax capensis) off western South Africa in 1991 caused by *Pasteurella multocida*. Waterbirds. 1992:236-8. doi.

19. Fereidouni S, Freimanis GL, Orynbayev M, Ribeca P, Flannery J, King DP, et al. Mass die-off of saiga antelopes, Kazakhstan, 2015. Emerg Infect Dis. 2019;25:1169-76. doi: 10.3201/eid2506.180990.

20. Eriksen L, Aalbaek B, Leifsson PS, Basse A, Christiansen T, Eriksen E, et al. Hemorrhagic septicemia in fallow deer (*Dama dama*) caused by *Pasteurella multocida*. J Zoo Wildl Med. 1999;30:285-92. doi.

21. Kock RA, Orynbayev M, Robinson S, Zuther S, Singh NJ, Beauvais W, et al. Saigas on the brink: Multidisciplinary analysis of the factors influencing mass mortality events Sci Adv. 2018;4:eaao2314. doi: 10.1126/sciadv.aao2314.

22. Mostaan S, Ghasemzadeh A, Sardari S, Shokrgozar MA, Nikbakht Brujeni G, Abolhassani M, et al. *Pasteurella multocida* vaccine candidates: a systematic review. Avicenna J Med Biotechnol. 2020;12:140-7. doi.

23. Harper M, Smallman TR, Boyce JD. Pasteurella. In: John F. Prescott, Janet I. MacInnes, Filip Van Immerseel, John D. Boyce, Andrew N. Rycroft, José A. Vázquez-Boland, editors. Pathogenesis of Bacterial Infections in Animals.5th ed. Wiley Blackwell; 2022. p. 221-243.

24. Harper M, Boyce JD, Wilkie IW, Adler B. Signature-tagged mutagenesis of *Pasteurella multocida* identifies mutants displaying differential virulence characteristics in mice and chickens. Infect Immun. 2003;71:5440-6. doi: 10.1128/iai.71.9.5440-5446.2003.

25. Chung JY, Wilkie I, Boyce JD, Townsend KM, Frost AJ, Ghoddusi M, et al. Role of capsule in the pathogenesis of fowl cholera caused by *Pasteurella multocida* serogroup A. Infect Immun. 2001;69:2487-92. doi: 10.1128/iai.69.4.2487-2492.2001.

26. Boyce JD, Adler B. The capsule is a virulence determinant in the pathogenesis of *Pasteurella multocida* M1404 (B:2). Infect Immun. 2000;68:3463-8. doi: 10.1128/iai.68.6.3463-3468.2000.

27. Smallman TR, Williams GC, Harper M, Boyce JD. Genome-wide investigation of *Pasteurella multocida* identifies the stringent response as a negative regulator of hyaluronic acid capsule production. Microbiol Spectr. 2022;10:e0019522. doi: 10.1128/spectrum.00195-22.

28. Islam EA, Fegan JE, Tefera TA, Curran DM, Waeckerlin RC, Ng D, et al. Reverse vaccinology-based identification of a novel surface lipoprotein that is an effective vaccine antigen against bovine infections caused by *Pasteurella multocida*. PLoS Pathog. 2023;19:e1011249. doi: 10.1371/journal.ppat.1011249.

29. Youn H, Carranza M. cAMP activation of the cAMP receptor protein, a model bacterial transcription factor. J Microbiol. 2023;61:277-87. doi: 10.1007/s12275-023-00028-6.

30. Danchin A, Guiso N, Roy A, Ullmann A. Identification of the *Escherichia coli cya* gene product as authentic adenylate cyclase. J Mol Biol. 1984;175:403-8. doi:10.1016/0022-2836(84)90356-5.

31. Imamura R, Yamanaka K, Ogura T, Hiraga S, Fujita N, Ishihama A, et al. Identification of the *cpdA* gene encoding cyclic 3',5'-adenosine monophosphate phosphodiesterase in *Escherichia coli*. J Biol Chem. 1996;271:25423-9. doi:10.1074/jbc.271.41.25423.

32. Ritzert JT, Minasov G, Embry R, Schipma MJ, Satchell KJF. The cyclic AMP receptor protein regulates quorum sensing and global gene expression in *Yersinia pestis* during planktonic growth and growth in biofilms. mBio. 2019;10. doi:10.1128/mBio.02613-19.

33. Gosset G, Zhang Z, Nayyar S, Cuevas WA, Saier MH, Jr. Transcriptome analysis of Crp-dependent catabolite control of gene expression in *Escherichia coli*. J Bacteriol. 2004;186:3516-24. doi:10.1128/jb.186.11.3516-3524.2004.

34. Suh SJ, Runyen-Janecky LJ, Maleniak TC, Hager P, MacGregor CH, Zielinski-Mozny NA, et al. Effect of *vfr* mutation on global gene expression and catabolite repression control of *Pseudomonas aeruginosa*. Microbiol Mol Biol Rev. 2002;148:1561-9. doi:10.1099/00221287-148-5-1561.

35. Zhao X, Liu Q, Xiao K, Hu Y, Liu X, Li Y, et al. Identification of the *crp* gene in avian *Pasteurella multocida* and evaluation of the effects of *crp* deletion on its phenotype, virulence and immunogenicity. BMC Microbiol. 2016;16:125. doi:10.1186/s12866-016-0739-y.

36. Gallagher LA, Bailey J, Manoil C. Ranking essential bacterial processes by speed of mutant death. Proc Natl Acad Sci U S A. 2020;117:18010-7. doi:10.1073/pnas.2001507117.

37. Homchampa P, Strugnell RA, Adler B. Cross protective immunity conferred by a marker-free *aroA* mutant of *Pasteurella multocida*. Vaccine. 1997;15:203-8. doi:10.1016/s0264-410x(96)00139-9.

38. Tabatabaei M, Liu Z, Finucane A, Parton R, Coote J. Protective immunity conferred by attenuated *aroA* derivatives of *Pasteurella multocida* B:2 strains in a mouse model of hemorrhagic septicemia. Infect Immun. 2002;70:3355-62. doi:10.1128/iai.70.7.3355-3362.2002.

39. Bosch M, Garrido E, Llagostera M, Pérez de Rozas AM, Badiola I, Barbé J. *Pasteurella multocida exbB*, *exbD* and *tonB* genes are physically linked but independently transcribed. FEMS Microbiol Lett. 2002;210:201-8. doi:10.1111/j.1574-6968.2002.tb11181.x.

40. Hatfaludi T, Al-Hasani K, Boyce JD, Adler B. Outer membrane proteins of *Pasteurella multocida*. Vet Microbiol. 2010;144:1-17. doi:10.1016/j.vetmic.2010.01.027.

41. Tatum FM, Tabatabai LB, Briggs RE. Sialic acid uptake is necessary for virulence of *Pasteurella multocida* in turkeys. Microb Pathog. 2009;46:337-44. doi:10.1016/j.micpath.2009.04.003.

42. Fuller TE, Kennedy MJ, Lowery DE. Identification of *Pasteurella multocida* virulence genes in a septicemic mouse model using signature-tagged mutagenesis. Microb Pathog. 2000;29:25-38. doi:10.1006/mpat.2000.0365.

43. Boyce JD, Wilkie I, Harper M, Paustian ML, Kapur V, Adler B. Genomic scale analysis of *Pasteurella multocida* gene expression during growth within the natural chicken host. Infect Immun. 2002;70:6871-9. doi:10.1128/iai.70.12.6871-6879.2002.

44. Paustian ML, May BJ, Kapur V. *Pasteurella multocida* gene expression in response to iron limitation. Infect Immun. 2001;69:4109-15. doi:10.1128/iai.69.6.4109-4115.2001.

45. Guo D-C, Yan S, Zhang A-Q, Liu J-S, Yan L, Liu P-X, et al. Construction and virulence of filamentous hemagglutinin protein B1 mutant of *Pasteurella multocida* in chickens. J Integr Agric. 2014;13:2268-75. doi: 10.1016/S2095-3119(14)60844-3

46. Tatum FM, Yersin AG, Briggs RE. Construction and virulence of a *Pasteurella multocida fhaB2* mutant in turkeys. Microb Pathog. 2005;39:9-17. doi:10.1016/j.micpath.2005.05.003.

47. Hooda Y, Lai CCL, Moraes TF. Identification of a large family of Slam-dependent surface lipoproteins in Gram-negative bacteria. Front Cell Infect Microbiol. 2017;7:207. doi:10.3389/fcimb.2017.00207.

48. Fegan JE, Waeckerlin RC, Tesfaw L, Islam EA, Deresse G, Dufera D, et al. Developing a PmSLP3-based vaccine formulation that provides robust long-lasting protection against hemorrhagic septicemia-causing serogroup B and E strains of *Pasteurella multocida* in cattle. Front Immunol. 2024;15:1392681. doi:10.3389/fimmu.2024.1392681.

49. Wu JR, Shien JH, Shieh HK, Chen CF, Chang PC. Protective immunity conferred by recombinant *Pasteurella multocida* lipoprotein E (PlpE). Vaccine. 2007;25:4140-8. doi:10.1016/j.vaccine.2007.03.005.

50. Hatfaludi T, Al-Hasani K, Gong L, Boyce JD, Ford M, Wilkie IW, et al. Screening of 71 *P. multocida* proteins for protective efficacy in a fowl cholera infection model and characterization of the protective antigen PlpE. PLoS One. 2012;7:e39973. doi:10.1371/journal.pone.0039973.

51. Nguyen QH, Lai CHR, Norris MJ, Ng D, Shah M, Lai CC, et al. A surface lipoprotein on Pasteurella multocida binds complement factor I to promote immune evasion. bioRxiv. 2024. doi:10.1101/2024.10.21.619360.

52. Li N, Feng T, Wang Y, Li P, Yin Y, Zhao Z, et al. A single point mutation in the *hyaC* gene affects *Pasteurella multocida* serovar A capsule production and virulence. Microb Pathog. 2021;159:105145. doi:10.1016/j.micpath.2021.105145.

53. Mégroz M, Kleifeld O, Wright A, Powell D, Harrison P, Adler B, et al. The RNA-binding chaperone Hfq is an important global regulator of gene expression in *Pasteurella multocida* and plays a crucial role in production of a number of virulence factors, including hyaluronic acid capsule. Infect Immun. 2016;84:1361-70. doi:10.1128/iai.00122-16.

54. Steen JA, Steen JA, Harrison P, Seemann T, Wilkie I, Harper M, et al. Fis is essential for capsule production in *Pasteurella multocida* and regulates expression of other important virulence factors. PLoS Pathog. 2010;6:e1000750. doi:10.1371/journal.ppat.1000750.

55. Deutscher J, Francke C, Postma PW. How phosphotransferase system-related protein phosphorylation regulates carbohydrate metabolism in bacteria. Microbiol Mol Biol Rev. 2006;70:939-1031. doi:10.1128/mmbr.00024-06.

56. Postma PW, Lengeler JW, Jacobson GR. Phosphoenolpyruvate:carbohydrate phosphotransferase systems of bacteria. Microbiol Rev. 1993;57:543-94. doi:10.1128/mr.57.3.543-594.1993.

57. Tchieu JH, Norris V, Edwards JS, Saier MH, Jr. The complete phosphotransferase system in *Escherichia coli*. J Mol Microbiol Biotechnol. 2001;3:329-46. doi.

58. Lévy S, Zeng GQ, Danchin A. Cyclic AMP synthesis in *Escherichia coli* strains bearing known deletions in the pts phosphotransferase operon. Gene. 1990;86:27-33. doi:10.1016/0378-1119(90)90110-d.

59. Reddy P, Kamireddi M. Modulation of *Escherichia coli* adenylyl cyclase activity by catalytic-site mutants of protein IIA(Glc) of the phosphoenolpyruvate:sugar phosphotransferase system. J Bacteriol. 1998;180:732-6. doi:10.1128/jb.180.3.732-736.1998.

60. Harper M, Boyce JD. The myriad properties of *Pasteurella multocida* lipopolysaccharide. Toxins (Basel). 2017;9. doi:10.3390/toxins9080254.

61. Harper M, Boyce JD, Cox AD, St Michael F, Wilkie IW, Blackall PJ, et al. Pasteurella multocida expresses two lipopolysaccharide glycoforms simultaneously, but only a single form is required for virulence: identification of two acceptor-specific heptosyl I transferases. Infect Immun. 2007;75:3885-93. doi:10.1128/iai.00212-07.

62. Harper M, Cox AD, St Michael F, Wilkie IW, Boyce JD, Adler B. A heptosyltransferase mutant of *Pasteurella multocida* produces a truncated lipopolysaccharide structure and is

attenuated in virulence. Infect Immun. 2004;72:3436-43. doi:10.1128/iai.72.6.3436-3443.2004.

63. Harper M, St Michael F, John M, Vinogradov E, Adler B, Boyce JD, et al. *Pasteurella multocida* Heddleston serovars 1 and 14 express different lipopolysaccharide structures but share the same lipopolysaccharide biosynthesis outer core locus. Vet Microbiol. 2011;150:289-96. doi:10.1016/j.vetmic.2011.01.028.

64. St Michael F, Harper M, Parnas H, John M, Stupak J, Vinogradov E, et al. Structural and genetic basis for the serological differentiation of *Pasteurella multocida* Heddleston serotypes 2 and 5. J Bacteriol. 2009;191:6950-9. doi:10.1128/jb.00787-09.

65. Harper M, St Michael F, John M, Vinogradov E, Steen JA, van Dorsten L, et al. *Pasteurella multocida* Heddleston serovar 3 and 4 strains share a common lipopolysaccharide biosynthesis locus but display both inter- and intrastrain lipopolysaccharide heterogeneity. J Bacteriol. 2013;195:4854-64. doi:10.1128/jb.00779-13.

66. Heap JT, Kuehne SA, Ehsaan M, Cartman ST, Cooksley CM, Scott JC, et al. The ClosTron: Mutagenesis in *Clostridium* refined and streamlined. J Microbiol Methods. 2010;80:49-55. doi:10.1016/j.mimet.2009.10.018.

67. Harms A, Liesch M, Körner J, Québatte M, Engel P, Dehio C. A bacterial toxin-antitoxin module is the origin of inter-bacterial and inter-kingdom effectors of *Bartonella*. PLoS Genet. 2017;13:e1007077. doi:10.1371/journal.pgen.1007077.

68. Shrum B, Anantha RV, Xu SX, Donnelly M, Haeryfar SM, McCormick JK, et al. A robust scoring system to evaluate sepsis severity in an animal model. BMC Res Notes. 2014;7:233. doi:10.1186/1756-0500-7-233.

69. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, et al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res. 2019;47:D309-d14. doi:10.1093/nar/gky1085.

70. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990;215:403-10. doi:10.1016/s0022-2836(05)80360-2.

71. Villanueva RAM, Chen ZJ. ggplot2: elegant graphics for data analysis. Taylor & Francis; 2019. https://doi.org/10.1080/15366367.2019.1565254

72. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Res. 2023;51:D638-d46. doi:10.1093/nar/gkac1000.