Aggressive Cholesterol Lowering Normalizes Atherosclerosis Regression in *Jak2^V617F^*

Clonal Hematopoiesis

Brian D. Hardaway (MPhil)[1], Trevor P. Fidler (PhD)[2], Mojdeh Tavallaie (PharmD-PhD)[1],

Kleopatra Avrampou (PhD)[1], Cheng-Chieh Hsu (PhD)[1], Sandra Schiavone (MA)[1], Tong Xiao

(MS)[1], Nan Wang (PhD)[1*], Alan R. Tall (MD)[1*]

[1]Division of Molecular Medicine, Department of Medicine, Columbia University Irving Medical

Center, New York, NY, USA.

[2]Cardiovascular Research Institute, University of California San Francisco, San Francisco, CA,

USA.

Short Title: Atherosclerosis Regression in *Jak2^V617F^* CH

[*]These authors contributed equally to this work.

Correspondence to:

Alan R. Tall: art1@cumc.columbia.edu

Abstract

Background - Clonal hematopoiesis (CH) has emerged as an important risk factor for atherosclerotic cardiovascular disease (ACVD). Mouse studies have established a causal role of CH in atherosclerosis progression and have defined macrophage inflammatory responses as a key underlying mechanism. We undertook the present study to assess the hypothesis that ongoing inflammation would impede atherosclerosis regression in $Jak2^{V617F}$ ($Jak2^{VF}$) CH mice.

Methods and Results - Chimeric $Jak2^{VF/WT}$ or control WT/WT bone marrow was transplanted into $Ldlr^{-/-}$ mice and, following 13-16 weeks of Western diet-induced atherosclerosis progression, cholesterol was lowered either moderately (to 200-300 mg/dl) or markedly (to 100 mg/dl). With moderate cholesterol lowering there was impaired regression in $Jak2^{VF}$ CH mice compared to controls. However, with marked cholesterol lowering, regression was similar in $Jak2^{VF}$ CH and control mice.

Two mechanisms of low-density lipoprotein (LDL) lowering-induced suppression of inflammation in plaques were implicated: 1) reversal of increased proliferation, DNA damage and Absent in Melanoma 2 (AIM2) inflammasome activation specifically in $Jak2^{VF}$ macrophages and 2) markedly increased macrophage triggering receptor expressed on myeloid cells 2 (TREM2), c-myc expressing macrophages in both $Jak2^{VF}$ and control mice.

Conclusions - Aggressive LDL lowering reverses inflammasome activation and induces pro-resolving changes in macrophages in $Jak2^{VF}$ CH, halting atherosclerosis progression and promoting features of plaque stabilization. These findings suggest that aggressive LDL cholesterol lowering could effectively reverse ACVD risk in individuals with $JAK2^{VF}$ clonal hematopoiesis.

Translational Perspective

Clonal hematopoiesis increases the risk of atherosclerotic cardiovascular disease. Amongst mutations associated with clonal hematopoiesis, $Jak2^{V617F}$ confers the greatest increased risk of atherosclerotic cardiovascular disease. Our findings demonstrate that aggressive LDL lowering normalizes atherosclerosis regression in $Jak2^{V617F}$ clonal hematopoiesis by reversing inflammasome activation and inducing pro-resolving changes in macrophages. These changes lead to increased features of plaque stability, including decreased macrophage burden, increased collagen, and improved efferocytosis. These results suggest that aggressive LDL-C lowering in $Jak2^{V617F}$ clonal hematopoiesis patients would mitigate the increased atherosclerotic cardiovascular disease risk.

1. Introduction

Despite therapeutic advances, ACVD remains the leading cause of death in the developed world.[1] Although inflammation plays a key role in atherosclerosis progression, cholesterol accumulation remains a prerequisite for lesion development, and LDL lowering has consistently shown a benefit in many clinical trials.[2-4]

The role of inflammation in increasing ACVD risk has also now been well established. The LoDoCo 2 trial found that the risk of cardiovascular events was significantly lower in patients with chronic coronary disease who were given 0.5 mg of colchicine daily.[5] Based on these and other results, the FDA approved the use of low dose colchicine as the first anti-inflammatory drug for heart disease.[6] The CANTOS trial involving the use of interleukin-1β (IL-1β) antibodies showed reduced cardiovascular events in high-risk patients with mean LDL cholesterol levels of 81 mg/dl - 84 mg/dl and a high-sensitivity C-reactive protein (hs-CRP) level of at least 2 mg/L.

However, there was a small increase in fatal infections, and the FDA did not approve this treatment for marketing.[7] Many observations suggest that hypercholesterolemia is linked to inflammatory risk. For example, statins lower hs-CRP levels[8] and preclinical studies have shown that cholesterol accumulation in plaque macrophages promotes inflammasome activation.[9-12] However, the mechanisms linking hypercholesterolemia, macrophage cholesterol accumulation and inflammation remain incompletely understood, especially in the clinically important context of atherosclerosis regression induced by LDL lowering.[13-19]

Clonal hematopoiesis (CH) arises from somatic mutations in hematopoietic stem cells (HSC) that confer a selective advantage and lead to clonal expansion of blood cells. CH commonly arises from mutations in genes that mediate epigenetic modifications or cytokine signaling.[20,21] Amongst CH mutations, $Jak2^{VF}$, which increases signaling by hematopoietic cytokines, confers the greatest risk of ACVD.[22-24] Moreover, $JAK2^{VF}$ has been detected in 3-4% of a general European population,[25] and increases thrombotic risk and coronary artery plaque erosion even at low variable allele frequency().[26,27,28] We showed in a hypercholesterolemic mouse model that $Jak2^{VF}$ CH exacerbates atherosclerosis via AIM2 inflammasome activation in mice[29] with genetic studies suggesting a similar role in humans.[30] The AIM2 inflammasome is activated by double stranded DNA fragments in the cytosol.[31] We proposed two potential mechanisms of DNA damage underlying AIM2 inflammasome activation in $Jak2^{VF}$ CH: increased proliferation of $Jak2^{VF}$ macrophages leading to DNA replication stress or increased mitochondrial glycolytic metabolism, reactive oxygen species (ROS) formation and oxidative DNA damage. Based on these findings, we hypothesized that LDL lowering-induced atherosclerosis regression would be impaired in $Jak2^{VF}$ CH mice due to persistence of the underlying mechanisms promoting inflammasome activation. Contrary to our hypothesis, we found that cholesterol lowering

81 reversed inflammasome associated changes in *Jak2^{VF}* macrophages and that marked cholesterol

82 lowering induced regression similar to control mice. These studies suggest the importance of

83 rigorous control of LDL cholesterol levels in individuals with *JAK2^{VF}* clonal hematopoiesis.

2. Methods

85 Several experimental procedures were performed as previously described[16,29]

2.1 Mice

All mice used for these studies were male and on a C57BL/6J background. Mice were housed in

a pathogen-free facility under standard conditions of temperature (23 °C) with a 12-hour light-

dark with ad libitum food and water access. Cages and water were changed every 7-14 days. All

mouse experiments were approved by the Institutional Animal Care and Use Committee of

Columbia University under protocol number AABG0561, and were conducted in accordance

with NIH guidelines and the Guide for the Care and Use of Laboratory Animals. Mice

heterozygous for the *Jak2^{VF}* conditional knock-in allele were generated as previously

described.[32] *Jak2^{VF}* mice were then crossed to *Mx1-Cre* (The Jackson Laboratory, B6.Cg-

Tg(*Mx1-Cre*)1*Cgn*/J (003556) or *Scl-Cre* (The Jackson Laboratory, C57BL/6-Tg(Tal1-

cre/ERT)42-056Jrg/J (037466) to generate *Mx1-CreJak2^{VF}* and *Scl-CreJak2^{VF}* mice respectively.

Scl-Cre-Jak2^{VF} mice were then crossed with ZsGreen reporter mice (The Jackson Laboratory,

B6.Cg-Gt(ROSA)26Sor^{tm6(CAG-ZsGreen1)Hze}/J (Ai6(RCL-ZsGreen) (007906) to generate *Scl-

CreJak2^{VF}ZsGreen* mice. *Ldlr^{-/-}* recipient mice for the *Mx1-CreJak2^{VF}* moderate cholesterol

lowering model that included *Myh11-Cre^{ERT2}* and *ZsGreen* alleles from a previous lineage-tracing

study were obtained from a collaborator and have been previously described.[33] However, these

alleles were not relevant to the experimental outcomes assessed and were not analyzed. Half of

the recipients were *Ldlr^{-/-} Myh-11-CreZsGreen* while the other half either lacked the *ZsGreen*

allele or both the *ZsGreen* allele and the *Myh11-Cre*. For the *Mx1-CreJak2^{VF}* and *Scl-CreJak2^{VF}*

aggressive cholesterol lowering studies, *Ldlr^{-/-}* recipient mice were purchased from The Jackson

Laboratory (B6.129S7-*Ldlr*tm1Her/J (002207)). Littermate control mice for studies involving

Mx1-CreJak2^{VF} mice contained either an allele with *Mx1-Cre* or the *Jak2^{VF}* transgene, but not

both. Littermate control mice for studies involving *Scl-CreJak2^{VF}ZsGreen* mice contained the

alleles for *Scl-Cre* and *ZsGreen* but not the *Jak2^{VF}* transgene.

2.2 Bone Marrow Transplantation

Bone marrow transplantations (BMT) were conducted as previously described with the following

exceptions.[10] For the *Mx1-CreJak2^{VF}* moderate cholesterol lowering study, donor mice were 8-12

weeks old while recipients were 11-14 weeks old. For the *Mx1-CreJak2^{VF}* effective cholesterol

lowering study, donor mice were 7-16 weeks old while recipients were 7 weeks old. For the *Scl-*

CreJak2^{VF} study, donor mice were 6-12 weeks old while recipients were 8 weeks old. For the

Mx1-CreJak2^{VF} moderate cholesterol lowering study, recipients were lethally irradiated once

with 10.5 Gy from a cesium gamma source. Due to decommissioning of the cesium gamma

source, recipients from the *Mx1-CreJak2^{VF}* and *Scl-CreJak2^{VF}* effective cholesterol lowering

studies were irradiated with 10.5 Gy using a Multirad 350 X-ray Irradiator from Precision X-

Ray. Within 24 hours of irradiation, bone marrow was isolated from donors with the indicated

genotypes. Total bone marrow cell number was quantified using an iN CYTO C-Chip

hemocytometer (DHC-N01) according to the manufacturer instructions. Irradiated mice were

randomized to treatment groups, anesthetized with isoflurane before receiving 3×10^6 total bone

marrow cells via intravenous (i.v.) injection at a total final volume of 100 µl. For the *Mx1-*

CreJak2^{VF} studies, recipients received either 6×10^5 *CD45.1Mx1-CreJak2^{VF/WT}* or 6×10^5

CD45.1 control bone marrow cells (20%) combined with 2.4 x 10^6 CD45.2Jak2$^{WT/WT}$ cells (80%).

For the *Scl-CreJak2^{VF}* study, recipients received either 6 x 10^5 *CD45.1CD45.2Scl-*

CreJak2$^{VF/WT}$ZsGreen or 6 x 10^5 *CD45.1CD45.2Scl-CreJak2$^{WT/WT}$ZsGreen* bone marrow cells

(20%) combined with 2.4 x 10^6 *CD45.1Jak2$^{WT/WT}$* cells (80%). For the *Mx1-CreJak2^{VF}* studies,

mice were allowed to recover for four weeks after BMT before being injected intraperitonially

(i.p.) with 50 µg/mouse/day polyinosinic:polycytidylic acid (pIpC) two times, 48 hours apart.

2.3 Atherosclerosis Studies

For the *Mx1-CreJak2^{VF}* studies, power analysis using GPower3.1 indicated that with a total

sample size of 80 (n = 20 per group), the study had 80% power to detect an effect size of f = 0.32

at α = 0.05, using two-way analysis of variance (ANOVA) across two genotypes and two

treatments. For the *Scl-CreJak2^{VF}* study, power analysis using GPower3.1 indicated that with a

total sample size of 80 (n = 16 per group), the study had 80% power to detect an effect size of f =

0.40 at α = 0.05, using one-way ANOVA across five groups. Four weeks after BMT. *Ldlr$^{-/-}$*

recipient mice were fed a western diet (ENVIGO, cat. no. TD.88137) for the indicated times. For

the final 10 days of the western diet period of the *Scl-CreJak2^{VF}* study, mice were switched to a

western diet containing tamoxifen at 500 mg/kg by weight (ENVIGO, cat. no. TD130889). To

induce LDL lowering in the *Mx1-CreJak2^{VF}* study of Figure 1A, mice were administered 2 x 10^{11}

viral particles per mouse in 200 µl of 1X PBS via tail vein injection of an hDAD-*hLDLr* viral

vector (Gene Vector Core Laboratory at the Baylor College of Medicine) and switched to chow

diet. To induce LDL lowering in the *Mx1-CreJak2^{VF}* study of Figure 4A and the *Scl-CreJak2^{VF}*

study of Figure 6A, mice were administered 2.5 x 10^{11} genome copies per mouse in 100 µl of 1X

PBS via tail vein injection of an AAV8-TBG-*mLdlr* vector (AAV-263355, Vector Biolabs) and

switched to chow diet. Mice were anesthetized using 5% isoflurane via inhalation at a flow rate

of approximately 3L/min oxygen using a calibrated vaporizer. Anesthetic depth was assessed

continuously using respiratory rate monitoring and the toe-pinch reflex. Isoflurane was

administered acutely during procedures requiring anesthesia, including blood collection and

euthanasia. Blood collection was performed via retro-orbital sinus for plasma, complete blood

counts, and flow cytometric analysis at the indicated timepoints. For euthanasia, mice were first

rendered deeply unconscious with isoflurane, after which cervical dislocation was performed as a

secondary method to ensure death, in accordance with AVMA guidelines. Following euthanasia,

aortic roots were collected and fixed with 4% paraformaldehyde (PFA) (Electron Microscopy

Sciences 15710; 16% PFA diluted 1:4 in 1.33X PBS) for 24 hours at 4 $^{\circ}$C , then embedded in

paraffin.

2.4 Histological Analysis

Lesion area and necrotic core area were quantified as previously described.[22,29] Researchers

blinded to genotype sectioned paraffin-embedded aortic root lesions at similar points between

mice over a 150 μm span for a total of 50 sections spaced 3 μm apart with 2 sections per slide on

25 slides. Six total slides spaced 30 μm apart (slides 1,5,10,15,20,25) were then stained with

hematoxylin & eosin (H&E) (Sigma MHS32-1L; Abcam ab246824) and imaged with an

Olympus DP27 camera attached to an Olympus CX43 biological microscope set to the 10X

objective using Olympus cellSens Entry 3.1 software. Lesion area was then quantified in a

blinded fashion and the average of the six slides was used to determine lesion area. The Necrotic

core area was similarly quantified in a blinded fashion using the same six H&E sections.

Collagen was stained using the Polysciences Picrosirius Red Stain Kit (cat. no. 24901) according

to the manufacturer's instructions. The same slide between mice was stained and the collagen

quantified by applying a consistent color threshold to calculate the collagen-positive area as a

percentage of total lesion area. Picrosirius red was imaged via the same system used for H&E

imaging. All lesion analysis was conducted using FIJI software.

2.5 Immunofluorescence

Identical slides of aortic roots between mice were baked in a dry incubator at 60 °C for 30

minutes and then deparaffinized in Histo-clear (National Diagnostics HS-200/50-899-90147)

three times for 10 minutes. Histo-clear was removed by 20 shakes in 100% ethanol twice

followed by 20 shakes in Histoprep Reagent Alcohol (fisher HC-600-1GAL) before being

washed in tap water 7 times. Antigen retrieval was performed by placing slides either in citric

acid-based antigen retrieval solution (Vector Laboratories, H3300) or tris-based antigen retrieval

solution (Vector Laboratories, H3301) and pressure cooking for 15 minutes with a 5-minute

natural release in an Instant Pot IP-LUX pressure cooker. Citric acid-based antigen retrieval was

performed for all antibodies except for the antibody against cleaved GasD. After pressure

cooking, slides were cooled for 10 minutes and then washed 3 times for 5 minutes in 1X

phosphate buffered saline (PBS) (Boston Bioproducts BM-220; diluted 1:10). Hydrophobic

circles were drawn using a liquid blocker pen (Sigma Z377821). For antibodies sourced from

mice, mouse on mouse blocking was performed by incubating sections in 10% goat serum

(ThermoFisher 10000C) in 1X PBS (Corning 21-040-CM) with 1 drop of mouse-on-mouse block

per 1.25ml of solution (Vector Laboratories NC9290646) for 1 hour at room temperature in a

humidified chamber. Slides were then washed in 1X PBS containing tween 20 (Fisher BP337-

500) at a concentration of 0.01% (PBS-T) twice for 5 minutes and then in 1X PBS once for 5

minutes. For antibodies not sourced from mice, sections were blocked by incubating in Normal

Serum Block (Biolegend 927503) for 1 hour at room temperature in a humidified chamber.

Following blocking, sections were incubated with the following primary antibodies at the

indicated dilutions or concentrations overnight at 4 °C in a humidified chamber: AIM2 (Abcam,

ab119791, 1:250), 8-hydroxydeoxyGuanosine (Abcam, ab183395, 1:100), Cleaved GasD (Cell

Signaling, 10137, 10μg/ml), c-myc (Cell Signaling, 5605, 5.76 μg/ml), Ki67 (Abcam, ab15580,

9 μg/ml), MAC2 conjugated to Alexa Fluor 488 (Cedarlane, CL8942AF4, 1 μg/ml), MAC2

(Cedarlane, CL8942AP, 1 μg/ml), MerTK (R&D, BAF591, 2 μg/ml), pγH2AX (Cell Signaling,

9718, 0.74 μg/ml), ZsGreen (ThermoFisher, TA180002, 10 μg/ml), TREM2 (Denali, 4D9

DC1847, 1:100). The following IgG antibodies were used as negative controls for primary

antibodies sourced from the same species at the same concentration: Mouse IgG (ChromPure,

015-000-003), Rabbit IgG (Abcam, ab172730), Rabbit IgG (Novus Biologicals, NBP2-24891),

Rabbit IgG (ChromPure 011-000-003), Rat IgG (ChromPure, 012-000-003), Rat IgG (Biolegend,

400431), Rat IgG conjugated to Alexa Fluor 488 (Invitrogen, 53-4321-80). Slides were then

washed twice with PBS-T for 5 minutes and once with 1X PBS for 5 minutes. The following

secondary antibodies were used at a dilution of 1:200 for 1 hour at room temperature: Rat

conjugated to Alexa Fluor 488 (Invitrogen A11006), Rat conjugated to Alexa Fluor 568

(Invitrogen A11077), Rat conjugated to Alexa Fluor 647 (Invitrogen A-21247), Rabbit

conjugated to Alexa Fluor 568 (Invitrogen A11011), Rabbit conjugated to Alexa Fluor 647

(A27040). The following secondary antibodies were used at a dilution of 1:800 for 1 hour at

room temperature: Mouse conjugated to Alexa Fluor 647 (Invitrogen A21237), Streptavidin

conjugated to Alexa Fluor 568 (Invitrogen S11226), Streptavidin conjugated to Alexa Fluor 647

(Invitrogen S32357). DAPI (4',6-Diamidino-2-Phenylindole, Dilactate) (Biolegend, 422801) was

included during incubation with secondary antibodies at a concentration of 5 μg/ml. Slides were

then washed twice with PBS-T for 5 minutes and once with 1X PBS for 5 minutes. After

washing, slides were mounted using Prolong Gold Antifade Reagent with DAPI (Invitrogen

P36935). TUNEL staining was performed using the In Situ Cell Death Detection Kit TMR Red

(Roche, 12156792910) according to the manufacturer's instructions. Slides were imaged on a

Leica Fluorescent DMI 6000B wide-field microscope or a Nikon Ti Eclipse inverted microscope

with high-sensitivity confocal imaging at the Confocal and Specialized Microscopy Shared

Resource (CSMSR) Core at Columbia University. All images were analyzed in a blinded fashion

using FIJI software.

2.6 Statistics

Statistical analyses were conducted using GraphPad Prism v10.5. Data are presented as the mean

$\pm$ s.e.m., with all measurements obtained from distinct biological samples. All statistical tests

were performed using two-sided analyses. Normality was tested using the D'Agostino-Pearson

omnibus and, where appropriate, log transformation was applied to approximate normality.

Formal outliers were removed using a two-tailed Grubbs' test ($\alpha = 0.05$). Repeated measures data

were analyzed using two-way ANOVA with the Geisser-Greenhouse correction for sphericity and

Tukey's multiple comparisons test with individual variance estimation. For datasets involving

two independent variables, two-way ANOVA with Tukey's multiple comparisons test or Kruskal-

Wallis with Dunn's multiple comparisons test were used based on distribution. For datasets

involving five experimental groups, one-way ANOVA with Holm-Sidak's multiple comparisons

test or Kruskal-Wallis with Dunn's multiple comparisons test were used based on distribution.

Details of blood analysis can be found in the supplement.

3. Results

3.1 Atherosclerosis Regression is Impaired with Moderate Cholesterol Lowering in $Jak2^{VF}$

CH

To assess the effect of $Jak2^{VF}$ CH on atherosclerosis regression, we placed $Jak2^{VF}$ CH male mice

and controls on a western diet for 16 weeks and then induced cholesterol lowering using a

Helper-dependent Adenoviral vector containing the human LDL receptor gene (hDAD-hLDLR)

combined with a switch to chow diet for an additional 5 weeks (Figure 1A). Prior studies were

performed in female mice.[16,29] $Jak2^{VF}$ CH male mice exhibited splenomegaly, lower body

weight, and increased RBCs, hematocrit, red blood cell distribution width (RDWc), and

neutrophils, mirroring previous findings in females (Supplemental Figure 1A-I).[29] In this model,

CD45.1+ cells are $Jak2^{VF}$ in experimental mice and WT in controls. CD45.1+ neutrophils and

monocytes were elevated while CD45.1+ lymphocytes were reduced (Supplemental Figure 1J-

L). Plasma cholesterol was reduced in $Jak2^{VF}$ CH mice on western diet (Figure 1B) similar to

previous findings in female mice and humans.[24,29,34] Plasma cholesterol remained moderately

elevated 10 days post-virus treatment but returned to pre-diet levels (~200-300 mg/dl) by the end

of the regression period (Figure 1B). At baseline, lesion area and necrotic core area were

significantly increased in the aortic root of male $Jak2^{VF}$ CH mice compared to controls,

confirming that $Jak2^{VF}$ CH worsens atherosclerosis in both male and female mice (Figure 1C-

E).[29] Moderate cholesterol lowering halted progression of lesion and necrotic core area in control

mice whereas $Jak2^{VF}$ CH mice developed significantly larger necrotic cores and displayed a

strong trend toward increased lesion area (p=0.052) (Figure 1C-E). Consistent with previous

regression studies,[35,36] lesional collagen was increased in control mice after cholesterol lowering

with no significant change in $Jak2^{VF}$ CH mice, further indicating impaired regression in the

$Jak2^{VF}$ CH mice (Figure 1F,G). The most consistent phenotype found in atherosclerosis

regression studies has been a decrease in macrophage burden.[13,17-19,36] Macrophages were

significantly decreased in control mice but not in $Jak2^{VF}$ CH mice after cholesterol lowering

(Figure 1H,I). These results indicate that atherosclerosis regression is impaired in $Jak2^{VF}$ CH

with moderate cholesterol lowering.

We next assessed whether impaired regression in $Jak2^{VF}$ CH mice was associated with persistent

inflammatory features in plaques. Necrotic, inflammatory cell death is mediated by activated

Gasdermin D (GasD) downstream of inflammasome activation.[37] We used an antibody to the

active N-terminal fragment of GasD to assess pyroptosis in plaques.[38] Total cleaved GasD in

plaques as well as cleaved GasD in the necrotic core were increased in $Jak2^{VF}$ CH lesions at

baseline and were not reduced by cholesterol lowering (Figure 2A-C). However, macrophage-

specific cleaved GasD was decreased by cholesterol lowering in $Jak2^{VF}$ CH lesions, suggesting

that macrophage pyroptosis had been suppressed by the time of harvest (Figure 2D,E).

Consistent with earlier findings in females, AIM2+ macrophages and phosphorylated histone

H2A.X (pγH2AX), an indicator of double stranded DNA break formation, were increased in

$Jak2^{VF}$ CH lesions at baseline and were decreased by cholesterol lowering, suggesting reversal of

AIM2 inflammasome activation (Figure 2F-I).[29] Additionally, macrophage proliferation trended

higher at baseline in $Jak2^{VF}$ CH mice and was significantly reduced by cholesterol lowering

(Figure 2J,K). Macrophage oxidized DNA was increased in control lesions and unchanged by

cholesterol lowering in $Jak2^{VF}$ CH lesions (Figure 2L,M). These data suggest reversal of

macrophage AIM2 inflammasome activation and pyroptosis by moderate cholesterol lowering in

$Jak2^{VF}$ CH mice, correlating with decreased macrophage proliferation and decreased DNA

damage response but not with oxidative DNA damage changes. The persistence of cleaved GasD

in the necrotic core area may reflect ongoing pyroptosis during the early phase of regression

when cholesterol levels were persistently elevated and could explain impaired regression in the

$Jak2^{VF}$ mice.

3.2 Cholesterol Lowering Restores Impaired Efferocytosis and increases TREM2+

Macrophages in *Jak2^{VF}* CH Lesions

Atherosclerosis regression has been associated with improvements in pro-resolving features such

as increased macrophage efferocytosis, mer proto-oncogene tyrosine kinase (MerTK)

expression,[13,39,40] increased numbers of TREM2+ macrophages and limitation of necrotic core

formation in plaques.[16,41] During atherosclerosis progression, macrophage MerTK and TREM2

were suppressed in *Jak2^{VF}* CH mouse plaques as a result of interleukin-1 (IL-1) signaling from

mutant to wild-type (WT) cells.[38] Consistent with these studies, we found that macrophage

MerTK was decreased in *Jak2^{VF}* CH lesions at baseline compared to controls (Figure 3A,B).

MerTK levels were completely restored by cholesterol lowering (Figure 3A,B). Additionally, we

found that cholesterol lowering dramatically increased the percentage of macrophages positive

for TREM2 in both control and *Jak2^{VF}* CH lesions (Figure 3C,D). Consistent with these findings,

macrophage-associated TUNEL+ nuclei as a percentage of total lesional TUNEL+ nuclei, an

estimate of in situ efferocytosis,[14] were significantly decreased in *Jak2^{VF}* CH lesions at baseline

and restored by cholesterol lowering, (Figure 3E,F). Together, these data suggest that cholesterol

lowering restores impaired efferocytosis in *Jak2^{VF}* CH lesions and increases the proportion of

inflammation-resolving macrophages in lesions. Thus, macrophage inflammasome activation and

associated defects in efferocytosis were reversed by cholesterol lowering in *Jak2^{VF}* CH mice. The

increased necrotic core and decreased collagen content in these mice could reflect limited

resolution of hypercholesterolemia during the earlier phases of regression.

3.3 More Effective Cholesterol Lowering Restores Atherosclerosis Regression

To test this hypothesis, we performed a similar regression study but with more effective LDL

cholesterol lowering. Murine LDL does not bind as strongly to the human LDLR (employed in

309 Figure 1) as to the mouse LDLR.[42,43] Therefore, we employed an Adeno-Associated Virus

310 serotype 8 vector (AAV8) containing the mouse *Ldlr* gene which more effectively and more

311 quickly lowered cholesterol levels (Figure 4A). In this model, CD45.1+ cells are *Jak2^{VF}* in

312 experimental mice and WT in controls. Again, *Jak2^{VF}* CH mice exhibited splenomegaly, lower

313 body weight, and increased RBCs, hematocrit, and RDWc while CD45.1+ neutrophils and

314 monocytes were elevated, and these parameters were not appreciably changed by cholesterol

315 lowering (Supplemental Figure 2, Figure 4B). Consistent with the first study, lesion area and

316 necrotic core area were significantly increased in *Jak2^{VF}* CH mice at baseline. However, in

317 contrast to the first study, while both groups showed similar upwards trends, there was no

318 significant change in lesion area or necrotic core area in either group after aggressive LDL

319 lowering (Figure 4C-E). Furthermore, aggressive cholesterol lowering significantly increased

320 lesional collagen and decreased macrophage burden in both groups (Figure 4F-I). Analysis of

321 cleaved GasD showed an increase in macrophages in *Jak2^{VF}* mice at baseline that was reversed

322 by cholesterol lowering, and in contrast to the first study a decrease in cleaved GasD in the

323 necrotic core that appeared to involve both control and *Jak2^{VF}* CH mice (Supplemental Figure

324 3A-D). Consistent with the first study, double stranded DNA breaks (pγH2AX) were increased in

325 Jak2VF CH lesions at baseline and more strongly decreased by aggressive cholesterol lowering

326 along with a significant decrease in macrophage proliferation (Supplemental Figure 3E-H).

327 Additionally, macrophage MerTK was increased by aggressive cholesterol lowering in a manner

328 consistent with the first study (Supplemental Figure 3I,J). These results indicate that more

329 effective cholesterol lowering completely reversed inflammatory features and induced similar

330 levels of atherosclerosis regression in control and *Jak2^{VF}* CH lesions.

3.4 Cholesterol Lowering Suppresses Proliferation of *Jak2*^VF Macrophages and increases c-Myc+TREM2+ Macrophages

The first two studies indicated that effective cholesterol lowering could reverse impaired regression and the underlying cellular defects that promote AIM2 inflammasome activation. However, in these studies *Jak2*^VF was active during progression and lesion area was increased at baseline, potentially influencing the results. To simulate the situation where the *Jak2*^VF mutation was acquired later in life after atherosclerosis had already developed, we performed a third study in which *Jak2*^VF was activated in HSCs at the start of regression by using a tamoxifen-inducible Scl-driven cre recombinase along with a floxed ZsGreen reporter allele (*Jak2*^VF *Scl* CH) to assess cell genotype specific changes (Figure 5A). In addition to the two cholesterol lowering groups (regression groups) we also included control and *Jak2*^VF mice that remained on the western diet (progression groups). As expected, plasma cholesterol remained elevated in the progression groups but was markedly and rapidly decreased in both regression groups (Figure 5B). In this model, cells double positive for CD45.1 and CD45.2 are *Jak2*^VF in experimental mice and WT in controls. The *Jak2*^VF phenotype was partially apparent as indicated by increased spleen weights, RDWc, and hematocrit as well as expansion of CD45.1+CD45.2+ blood monocytes and neutrophils, but less pronounced than in the earlier studies when *Jak2*^VF was activated prior to Western diet feeding (Supplemental Figure 4).

Compared to the earlier models, the impact of *Jak2*^VF on lesion progression was less pronounced which likely reflects the shorter period of *Jak2*^VF activation (Supplemental Figure 5A-C). However, consistent with the earlier studies, the necrotic core significantly worsened in the *Jak2*^VF *Scl* CH progression mice but not in the control *Scl* mice compared to baseline, while the necrotic core was not significantly changed in either regression group compared to baseline

(Supplemental Figure 5C). Additionally, macrophage burden was similarly decreased in both

regression groups, (Supplemental Figure 5D,E). These studies further support that aggressive

cholesterol lowering similarly promotes regression in control and *Jak2*[VF] CH mice, compared to

progression or baseline groups, in mice with *Jak2*[VF] only activated at the beginning of the

regression period.

Cleaved GasD was increased in *Jak2*[VF] CH lesions compared to baseline and trended lower

(p=.06) after cholesterol lowering, consistent with aggressive cholesterol lowering reversing

pyroptosis in lesions (Figure 5C,D). In the *Jak2*[VF] CH progression group, proliferation was

significantly increased in *Jak2*[VF] (green) macrophages but not in WT (non-green) macrophages

indicating a mutant cell specific effect, as seen previously.[29] Notably, this was reversed by

cholesterol lowering indicating that proliferation of the *Jak2*[VF] cells was highly dependent on

LDL cholesterol levels (Figure 5E-G). Consistently, the DNA damage response (pγH2AX) was

significantly increased in *Jak2*[VF] cells, and this was reversed by cholesterol lowering while WT

cells showed no significant changes within *Jak2*[VF] CH lesions (Figure 5H-J). This suggests that

cell autonomous proliferation and associated formation of double stranded DNA breaks is

reversed by cholesterol lowering in lesions of *Jak2*[VF] CH mice.

Given that a proliferative efferocytic population with high c-Myc expression has been described

during regression,[13,39] we assessed c-Myc expression along with TREM2. In contrast to the

decreases in Ki67, macrophage c-Myc and c-Myc+TREM2[Hi] macrophages were increased

during regression in both control and *Jak2*[VF] mice, suggesting a selective increase of TREM2+,

anti-inflammatory macrophages despite a decrease in total macrophages (Figure 6).

4. Discussion

We previously showed that the combination of LDL lowering and *Jak2[VF]* genetic inactivation

induced fibrous cap thickening and increased TREM2[Hi] macrophages in *Jak2[VF]* CH mice.[16]

Given the propensity of plaque *Jak2[VF]* mutant cells to proliferation, DNA damage, and AIM2

inflammasome activation,[29] we hypothesized that LDL lowering alone would be insufficient to

induce plaque regression in *Jak2[VF]* mice. While this proved to be the case with moderate LDL

lowering, aggressive cholesterol lowering produced plaque regression (or halted progression)

similarly in controls and *Jak2[VF]* CH mice. Surprisingly, LDL lowering reversed the major

cellular changes in *Jak2[VF]* macrophages that have been associated with athero-progression. This

included reduced AIM2 expression and pyroptosis and increased MerTK, TREM2 and in situ

efferocytosis that have all been mechanistically linked in prior studies of *Jak2[VF]* mice.[38] Whereas

we previously considered glycolytic mitochondrial ROS production and oxidative DNA damage

as a likely mechanism underlying AIM2 inflammasome activation, the reversal of pyroptosis by

cholesterol lowering was not associated with reduced oxidized DNA. Rather, decreased

macrophage proliferation and reduced replication stress appears to be the major mechanism

underlying reversal of pyroptosis.

Studies of atherosclerosis regression induced by LDL lowering in mouse models have revealed

multiple underlying mechanisms[44] including macrophage emigration,[45] decreased monocyte

entry[46] and a key role of acquired immunity and T-regs.[17] Similar to macrophage proliferation

being a major factor accounting for increased macrophage number during advanced

atherosclerosis progression,[47] suppression of proliferation is the predominant mechanism

decreasing macrophage burden during regression.[35] Our studies suggest that decreased

proliferation of *Jak2[VF]* macrophages is also a key mechanism decreasing macrophage burden

during atherosclerosis regression. Moreover, decreased proliferation was associated with reduced

DNA damage response, AIM2 expression and pyroptosis in $Jak2^{VF}$ CH lesions, suggesting

decreased replication induced DNA damage as the key underlying mechanism responding to

cholesterol lowering.

Paradoxically, despite the overall reduction in macrophage burden, during regression there may

also be a relative increase in a proliferative set of efferocytotic macrophages that also express c-

myc expression and contribute to plaque stability.[13,14] However, in addition to a potential role in

cell proliferation c-myc has an additional role in Akt signaling and cell survival.[48,49] Previously,

we observed a 4-fold increase in TREM2[Hi] macrophages during atherosclerosis regression

induced by both LDL lowering and $Jak2^{VF}$ inactivation in $Jak2^{VF}$ CH mice.[16] In the present

study, we found that LDL lowering alone led to a marked increase in TREM2 staining in

macrophages, as well as elevated c-myc expression specifically within TREM2[Hi] macrophages,

in both control and mutant mice. This contrasts with the decreased proliferation and DNA

damage response found specifically in the $Jak2^{VF}$ macrophages. Our findings together suggest

that during regression there is an increase in TREM2[Hi], c-myc and MerTK expressing

macrophages either through increased proliferation or increased survival of these subsets that

then have beneficial effects on efferocytosis and plaque stability. Given that inflammasome

activation and IL-1 release have been linked to reduced MerTK and TREM2 as a result of A

Disintegrin And Metalloproteinase 17 (ADAM17)-mediated cleavage,[39,50] the increase in this

pro-resolving macrophage population may in part be secondary to reduced cholesterol-induced

inflammasome activation. Although IL-1β antibodies have been shown to inhibit atherosclerosis

regression in $Apoe^{-/-}$ mice,[15] IL-1β antibodies improved lesion stability in hypercholesterolemic

$Jak2^{VF}$ CH mice.[16,29] Elevated IL-1β signaling as a result of increased pyroptosis in $Jak2^{VF}$ CH

lesions is decreased in the present study as a result of LDL lowering. Combined, these results

suggest that excessive IL-1β activation is pro-atherogenic in the context of hypercholesterolemia

while a basal level of IL-1β activation may be required for lesion resolution after cholesterol

lowering which is blocked when IL-1β antibodies are present.

The history of LDL lowering guidelines has been to recommend progressively lower targets

based on clinical trial evidence. Current recommendations suggest an LDL cholesterol target

below 70 mg/dl for patients with established ACVD or in certain high-risk groups.[51] Statin use

has been associated with improved outcomes in human myeloproliferative neoplasm (MPN)

patients and was associated with a 37% reduction in the risk of thrombosis in patients who had

polycythemia vera or essential thrombocythemia as well as a 22% reduction in all-cause

mortality.[52] While data from mice should be extrapolated to humans with caution, the AIM2

inflammasome seems to increase ACVD risk in both mice and humans carrying the $Jak2^{VF}$

mutation.[30] The present preclinical study adds mechanistic findings to support the idea that

aggressive LDL-C lowering in $Jak2^{VF}$ MPN patients would reduce ACVD risk.

Acknowledgements

The authors thank Wenli Liu for assistance and support during the course of this study. This

study used the Confocal and Specialized Microscopy Shared Resource of the Herbert Irving

Comprehensive Cancer Center at Columbia University, funded in part through NIH/NCI Cancer

Center Support Grant P30CA013696. Additionally, this study used the resources of the Herbert

Irving Comprehensive Cancer Center Flow Cytometry Shared Resources funded in part through

Center Grant P30CA013696.

Sources of Funding

443   This work was supported by grants from the National Institutes of Health and The National

444   Heart, Lung, and Blood Institute: HL155431, HL107653HL170157-02, P01HL172741 to A.R.T

445   and HL148071 to N.W.

Disclosures

447   A.R.T. is a consultant for CSL Behring and is on the scientific advisory board of Beren

448   Therapeutics. The remaining authors declare no competing financial interests.

Author Contribution Statement

450   B.H., A.T., N.W., T.F. made substantial contributions to conception, design, data acquisition,

451   analysis, and interpretation. They also drafted and critically reviewed the manuscript for

452   intellectual content. M.T., K.A., C.H., S.S., and T.X. contributed to data acquisition and analysis.

Data Availability Statement

454   The datasets generated for this study are available from the corresponding author upon

455   reasonable request.

References

1. Martin SS, Aday AW, Almarzooq ZI, Anderson CAM, Arora P, Avery CL, Baker-Smith CM, Barone Gibbs B, Beaton AZ, Boehme AK, Commodore-Mensah Y, Currie ME, Elkind MSV, Evenson KR, Generoso G, Heard DG, Hiremath S, Johansen MC, Kalani R, Kazi DS, Ko D, Liu J, Magnani JW, Michos ED, Mussolino ME, Navaneethan SD, Parikh NI, Perman SM, Poudel R, Rezk-Hanna M, Roth GA, Shah NS, St-Onge MP, Thacker EL, Tsao CW, Urbut SM, Van Spall HGC, Voeks JH, Wang NY, Wong ND, Wong SS, Yaffe K, Palaniappan LP; American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. 2024 Heart Disease and Stroke Statistics: A Report of US and Global Data From the American Heart Association. *Circulation* 2024;**149**:e347–e913.

2. Vallejo-Vaz AJ, Fayyad R, Boekholdt SM, Hovingh GK, Kastelein JJP, Melamed S, Barter P, Waters DD, Ray KK. Triglyceride-rich lipoprotein cholesterol and risk of cardiovascular events among patients receiving statin therapy in the TNT trial. *Circulation* 2018;**138**:770–781.

3. Cannon CP, Blazing MA, Giugliano RP, McCagg A, White JA, Theroux P, Darius H, Lewis BS, Ophuis TO, Jukema JW, De Ferrari GM, Ruzyllo W, De Lucca P, Im K, Bohula EA, Reist C, Wiviott SD, Tershakovec AM, Musliner TA, Braunwald E, Califf RM; IMPROVE-IT Investigators. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med* 2015;**372**:2387–2397.

4. Sabatine MS, Giugliano RP, Keech AC, Honarpour N, Wiviott SD, Murphy SA, Kuder JF, Wang H, Liu T, Wasserman SM, Sever PS, Pedersen TR; FOURIER Steering

Committee and Investigators. Evolocumab and clinical outcomes in patients with

cardiovascular disease. *N Engl J Med* 2017;**376**:1713–1722.

5. Nidorf SM, Fiolet ATL, Mosterd A, Eikelboom JW, Schut A, Opstal TSJ, The SHK, Xu

XF, Ireland MA, Lenderink T, Latchem D, Hoogslag P, Jerzewski A, Nierop P, Whelan A,

Hendriks R, Swart H, Schaap J, Kuijper AFM, van Hessen MWJ, Saklani P, Tan I,

Thompson AG, Morton A, Judkins C, Bax WA, Dirksen M, Alings M, Hankey GJ,

Budgeon CA, Tijssen JGP, Cornel JH, Thompson PL; LoDoCo2 Trial Investigators.

Colchicine in patients with chronic coronary disease. *N Engl J Med* 2020;**383**:1838–

1847.

6. U.S. Food and Drug Administration. Drug approval package: Lodoco (colchicine) tablets,

0.5 mg. Approval letter. Published June 16, 2023. Accessed July 5, 2025.

https://www.accessdata.fda.gov/drugsatfda_docs/nda/2024/215727Orig1s000Approv.pdf

7. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne CM, Fonseca

F, Nicolau J, Koenig W, Anker SD, Kastelein JJP, Cornel JH, Pais P, Pella D, Genest J,

Cifkova R, Lorenzatti A, Forster T, Kobalava Z, Vida-Simiti L, Flather M, Shimokawa H,

Ogawa H, Dellborg M, Rossi PRF, Troquay RPT, Libby P, Glynn RJ; CANTOS Trial

Group. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J*

Med 2017;**377**:1119–1131.

8. Ridker PM. From C-reactive protein to interleukin-6 to interleukin-1: moving upstream to

identify novel targets for atheroprotection. *Circ Res* 2016;**118**:145–156.

9. Yalcinkaya M, Liu W, Xiao T, Abramowicz S, Wang R, Wang N, Westerterp M, Tall AR.

Cholesterol trafficking to the ER leads to the activation of CaMKII/JNK/NLRP3 and

promotes atherosclerosis. *J Lipid Res* 2024;**65**:100534.

10. Westerterp M, Fotakis P, Ouimet M, Bochem AE, Zhang H, Molusky MM, Wang W,

Abramowicz S, la Bastide-van Gemert S, Wang N, Welch CL, Reilly MP, Stroes ES,

Moore KJ, Tall AR. Cholesterol efflux pathways suppress inflammasome activation,

NETosis, and atherogenesis. *Circulation* 2018;**138**:898–912.

11. Yalcinkaya M, Fotakis P, Liu W, Endo-Umeda K, Dou H, Abramowicz S, Xiao T, Libby

P, Wang N, Tall AR, Westerterp M. Cholesterol accumulation in macrophages drives

NETosis in atherosclerotic plaques via IL-1β secretion. *Cardiovasc Res* 2023;**119**:969–

981.

12. Sheedy FJ, Grebe A, Rayner KJ, Kalantari P, Ramkhelawon B, Carpenter SB, Becker CE,

Ediriweera HN, Mullick AE, Golenbock DT, Stuart LM, Latz E, Fitzgerald KA, Moore

KJ. CD36 coordinates NLRP3 inflammasome activation by facilitating intracellular

nucleation of soluble ligands into particulate ligands in sterile inflammation. *Nat*

Immunol 2013;**14**:812–820.

13. Gerlach BD, Ampomah PB, Yurdagul A Jr, Liu C, Lauring MC, Wang X, Kasikara C,

Kong N, Shi J, Tao W, Tabas I. Efferocytosis induces macrophage proliferation to help

resolve tissue injury. *Cell Metab* 2021;**33**:2445–2463.e8.

14. Yurdagul A Jr, Subramanian M, Wang X, Crown SB, Ilkayeva OR, Darville L, Kolluru

GK, Rymond CC, Gerlach BD, Zheng Z, Kuriakose G, Kevil CG, Koomen JM,

Cleveland JL, Muoio DM, Tabas I. Macrophage metabolism of apoptotic cell-derived

arginine promotes continual efferocytosis and resolution of injury. *Cell Metab*

2020;**31**:518–533.e10.

15. Karnewar S, Karnewar V, Deaton RA, Shankman LS, Benavente ED, Williams CM,

Bradley X, Alencar GF, Bulut GB, Kirmani S, Baylis RA, Zunder ER, den Ruijter HM,

Pasterkamp G, Owens GK. IL-1β inhibition partially negates the beneficial effects of diet-induced atherosclerosis regression in mice. *Arterioscler Thromb Vasc Biol* 2024;**44**:1379–1392.

16. Fidler TP, Dunbar A, Kim E, Hardaway B, Pauli J, Xue C, Abramowicz S, Xiao T, O'Connor K, Sachs N, Wang N, Maegdefessel L, Levine R, Reilly MP, Tall AR. Suppression of IL-1β promotes beneficial accumulation of fibroblast-like cells in atherosclerotic plaques in clonal hematopoiesis. *Nat Cardiovasc Res* 2024;**3**:60–75.

17. Sharma M, Schlegel MP, Afonso MS, Brown EJ, Rahman K, Weinstock A, Sansbury BE, Corr EM, van Solingen C, Koelwyn GJ, Shanley LC, Beckett L, Peled D, Lafaille JJ, Spite M, Loke P, Fisher EA, Moore KJ. Regulatory T cells license macrophage pro-resolving functions during atherosclerosis regression. *Circ Res* 2020;**127**:335–353.

18. Zhang X, Rotllan N, Canfrán-Duque A, Sun J, Toczek J, Moshnikova A, Malik S, Price NL, Araldi E, Zhong W, Sadeghi MM, Andreev OA, Bahal R, Reshetnyak YK, Suárez Y, Fernández-Hernando C. Targeted suppression of miRNA-33 using pHLIP improves atherosclerosis regression. *Circ Res* 2022;**131**:77–90.

19. Basu D, Hu Y, Huggins LA, Mullick AE, Graham MJ, Wietecha T, Barnhart S, Mogul A, Pfeiffer K, Zirlik A, Fisher EA, Bornfeldt KE, Willecke F, Goldberg IJ. Novel reversible model of atherosclerosis and regression using oligonucleotide regulation of the LDL receptor. *Circ Res* 2018;**122**:560–567.

20. Tall AR, Fuster JJ. Clonal hematopoiesis in cardiovascular disease and therapeutic implications. *Nat Cardiovasc Res* 2022;**1**:116–124.

21. Jaiswal S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG, Lindsley RC, Mermel CH, Burtt N, Chavez A, Higgins JM, Moltchanov V, Kuo FC, Kluk MJ,

Henderson B, Kinnunen L, Koistinen HA, Ladenvall C, Getz G, Correa A, Banahan BF, Gabriel S, Kathiresan S, Stringham HM, McCarthy MI, Boehnke M, Tuomilehto J, Haiman CA, Groop L, Atzmon G, Wilson JG, Neuberg D, Altshuler D, Ebert BL. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med* 2014;**371**:2488–2498.

22. Jaiswal S, Natarajan P, Silver AJ, Gibson CJ, Bick AG, Shvartz E, McConkey M, Gupta N, Gabriel S, Ardissino D, Baber U, Mehran R, Fuster V, Danesh J, Frossard P, Saleheen D, Melander O, Sukhova GK, Neuberg D, Libby P, Kathiresan S, Ebert BL. Clonal hematopoiesis and risk of atherosclerotic cardiovascular disease. *N Engl J Med* 2017;**377**:111–121.

23. Kar SP, Quiros PM, Gu M, Jiang T, Mitchell J, Langdon R, Iyer V, Barcena C, Vijayabaskar MS, Fabre MA, Carter P, Petrovski S, Burgess S, Vassiliou GS. Genome-wide analyses of 200 453 individuals yield new insights into the causes and consequences of clonal hematopoiesis. *Nat Genet* 2022;**54**:1155–1166.

24. Bick AG, Weinstock JS, Nandakumar SK, Fulco CP, Bao EL, Zekavat SM, Szeto MD, Liao X, Leventhal MJ, Nasser J, Chang K, Laurie C, Burugula BB, Gibson CJ, Lin AE, Taub MA, Aguet F, Ardlie K, Mitchell BD, Barnes KC, Moscati A, Fornage M, Redline S, Psaty BM, Silverman EK, Weiss ST, Palmer ND, Vasan RS, Burchard EG, Kardia SLR, He J, Kaplan RC, Smith NL, Arnett DK, Schwartz DA, Correa A, de Andrade M, Guo X, Konkle BA, Custer B, Peralta JM, Gui H, Meyers DA, McGarvey ST, Chen IY, Shoemaker MB, Peyser PA, Broome JG, Gogarten SM, Wang FF, Wong Q, Montasser ME, Daya M, Kenny EE, North KE, Launer LJ, Cade BE, Bis JC, Cho MH, Lasky-Su J, Bowden DW, Cupples LA, Mak ACY, Becker LC, Smith JA, Kelly TN, Aslibekyan S,

Heckbert SR, Tiwari HK, Yang IV, Heit JA, Lubitz SA, Johnsen JM, Curran JE, Wenzel SE, Weeks DE, Rao DC, Darbar D, Moon JY, Tracy RP, Buth EJ, Rafaels N, Loos RJF, Durda P, Liu Y, Hou L, Lee J, Kachroo P, Freedman BI, Levy D, Bielak LF, Hixson JE, Floyd JS, Whitsel EA, Ellinor PT, Irvin MR, Fingerlin TE, Raffield LM, Armasu SM, Wheeler MM, Sabino EC, Blangero J, Williams LK, Levy BD, Sheu WH, Roden DM, Boerwinkle E, Manson JE, Mathias RA, Desai P, Taylor KD, Johnson AD; NHLBI Trans-Omics for Precision Medicine Consortium; Auer PL, Kooperberg C, Laurie CC, Blackwell TW, Smith AV, Zhao H, Lange E, Lange L, Rich SS, Rotter JI, Wilson JG, Scheet P, Kitzman JO, Lander ES, Engreitz JM, Ebert BL, Reiner AP, Jaiswal S, Abecasis G, Sankaran VG, Kathiresan S, Natarajan P. Inherited causes of clonal haematopoiesis in 97,691 whole genomes. *Nature* 2020;**586**:763–768.

25. Cordua S, Kjaer L, Skov V, Pallisgaard N, Hasselbalch HC, Ellervik C. Prevalence and phenotypes of JAK2 V617F and calreticulin mutations in a Danish general population. *Blood* 2019;**134**:469–479.

26. Wang S, Luo X, Hu S, Zhao C, Sun Q, Zeng M, Bao X, Wang Y, Wu F, Yang Y, Lv Y, Bai X, Hao W, Liu M, Yi B, Chen Y, Meng W, Li J, Li M, Huang J, Wu T, Zhao Y, Zhang Z, An J, Libby P, Jia H, Yu B. Plaque erosion risk and JAK2 V617F variant. *Eur Heart J* 2025;**46**:2205–2219.

27. Molinaro R, Sellar RS, Vromman A, Sausen G, Folco E, Sukhova GK, McConke ME, Corbo C, Ebert BL, Libby P. The clonal hematopoiesis mutation Jak2V617F aggravates endothelial injury and thrombosis in arteries with erosion-like intimas. *Int J Cardiol* 2024;**409**:132184.

28. Liu W, Pircher J, Schuermans A, Ul Ain Q, Zhang Z, Honigberg MC, Yalcinkaya M,

Nakao T, Pournamadri A, Xiao T, Hajebrahimi MA, Wasner L, Stegner D, Petzold T,

Natarajan P, Massberg S, Tall AR, Schulz C, Wang N. Jak2 V617F clonal hematopoiesis

promotes arterial thrombosis via platelet activation and cross talk. *Blood* 2024;**143**:1539–

1550.

29. Fidler TP, Xue C, Yalcinkaya M, Hardaway B, Abramowicz S, Xiao T, Liu W, Thomas

DG, Hajebrahimi MA, Pircher J, Silvestre-Roig C, Kotini AG, Luchsinger LL, Wei Y,

Westerterp M, Snoeck HW, Papapetrou EP, Schulz C, Massberg S, Soehnlein O, Ebert

BL, Levine RL, Reilly MP, Libby P, Wang N, Tall AR. The AIM2 inflammasome

exacerbates atherosclerosis in clonal haematopoiesis. *Nature* 2021;**592**:296–301.

30. Yu Z, Fidler TP, Ruan Y, Vlasschaert C, Nakao T, Uddin MM, Mack T, Niroula A,

Heimlich JB, Zekavat SM, Gibson CJ, Griffin GK, Wang Y, Peloso GM, Heard-Costa N,

Levy D, Vasan RS, Aguet F, Ardlie KG, Taylor KD, Rich SS, Rotter JI, Libby P, Jaiswal

S, Ebert BL, Bick AG, Tall AR, Natarajan P. Genetic modification of inflammation- and

clonal hematopoiesis-associated cardiovascular risk. *J Clin Invest* 2023;**133**:e168597.

31. Rathinam VA, Fitzgerald KA. Inflammasome complexes: emerging mechanisms and

effector functions. *Cell* 2016;**165**:792–800.

32. Mullally A, Lane SW, Ball B, Megerdichian C, Okabe R, Al-Shahrour F, Paktinat M,

Haydu JE, Housman E, Lord AM, Wernig G, Kharas MG, Mercher T, Kutok JL, Gilliland

DG, Ebert BL. Physiological Jak2V617F expression causes a lethal myeloproliferative

neoplasm with differential effects on hematopoietic stem and progenitor cells. *Cancer*

Cell 2010;**17**:584–596.

33. Pan H, Xue C, Auerbach BJ, Fan J, Bashore AC, Cui J, Yang DY, Trignano SB, Liu W, Shi J, Ihuegbu CO, Bush EC, Worley J, Vlahos L, Laise P, Solomon RA, Connolly ES, Califano A, Sims PA, Zhang H, Li M, Reilly MP. Single-cell genomics reveals a novel cell state during smooth muscle cell phenotypic switching and potential therapeutic targets for atherosclerosis in mouse and human. *Circulation* 2020;**142**:2060–2075.

34. Kar SP, Quiros PM, Gu M, Jiang T, Mitchell J, Langdon R, Iyer V, Barcena C, Vijayabaskar MS, Fabre MA, Carter P, Petrovski S, Burgess S, Vassiliou GS. Genome-wide analyses of 200 453 individuals yield new insights into the causes and consequences of clonal hematopoiesis. *Nat Genet* 2022;**54**:1155–1166.

35. Härdtner C, Kornemann J, Krebs K, Ehlert CA, Jander A, Zou J, Starz C, Rauterberg S, Sharipova D, Dufner B, Hoppe N, Dederichs TS, Willecke F, Stachon P, Heidt T, Wolf D, von Zur Mühlen C, Madl J, Kohl P, Kaeser R, Boettler T, Pieterman EJ, Princen HMG, Ho-Tin-Noé B, Swirski FK, Robbins CS, Bode C, Zirlik A, Hilgendorf I. Inhibition of macrophage proliferation dominates plaque regression in response to cholesterol lowering. *Basic Res Cardiol* 2020;**115**:78.

36. Peled M, Nishi H, Weinstock A, Barrett TJ, Zhou F, Quezada A, Fisher EA. A wild-type mouse-based model for the regression of inflammation in atherosclerosis. *PLoS One* 2017;**12**:e0173975.

37. Broz P. Pyroptosis: molecular mechanisms and roles in disease. *Cell Res* 2025;**35**:334–344.

38. Liu W, Hardaway BD, Kim E, Pauli J, Wettich JL, Yalcinkaya M, Hsu CC, Xiao T, Reilly MP, Tabas I, Maegdefessel L, Schlepckow K, Haass C, Wang N, Tall AR. Inflammatory

crosstalk impairs phagocytic receptors and aggravates atherosclerosis in clonal

hematopoiesis in mice. *J Clin Invest* 2024;**135**:e182939.

39. Ngai D, Schilperoort M, Tabas I. Efferocytosis-induced lactate enables the proliferation

of pro-resolving macrophages to mediate tissue repair. *Nat Metab* 2023;**5**:2206–2219.

40. Sukka SR, Ampomah PB, Darville LNF, Ngai D, Wang X, Kuriakose G, Xiao Y, Shi J,

Koomen JM, McCusker RH, Tabas I. Efferocytosis drives a tryptophan metabolism

pathway in macrophages to promote tissue resolution. *Nat Metab* 2024;**6**:1736–1755.

41. Patterson MT, Xu Y, Hillman H, Osinski V, Schrank PR, Kennedy AE, Barrow F, Zhu A,

Tollison S, Shekhar S, Stromnes IM, Tassi I, Wu D, Revelo XS, Binstadt BA, Williams

JW. Trem2 agonist reprograms foamy macrophages to promote atherosclerotic plaque

stability — brief report. *Arterioscler Thromb Vasc Biol* 2024;**44**:1646–1657.

42. Corsini A, Mazzotti M, Villa A, Maggi FM, Bernini F, Romano L, Romano C, Fumagalli

R, Catapano AL. Ability of the LDL receptor from several animal species to recognize

the human apo B binding domain: studies with LDL from familial defective apo B-100.

Atherosclerosis 1992;**93**:95–103.

43. Innerarity TL, Pitas RE, Mahley RW. Disparities in the interaction of rat and human

lipoproteins with cultured rat fibroblasts and smooth muscle cells: requirements for

homology for receptor binding activity. *J Biol Chem* 1980;**255**:11163–11172.

44. Lin JD, Nishi H, Poles J, Niu X, McCauley C, Rahman K, Brown EJ, Yeung ST, Vozhilla

N, Weinstock A, Ramsey SA, Fisher EA, Loke P. Single-cell analysis of fate-mapped

macrophages reveals heterogeneity, including stem-like properties, during atherosclerosis

progression and regression. *JCI Insight* 2019;**4**:e124574.

45. Feig JE, Pineda-Torra I, Sanson M, Bradley MN, Vengrenyuk Y, Bogunovic D, Gautier

EL, Rubinstein D, Hong C, Liu J, Wu C, van Rooijen N, Bhardwaj N, Garabedian M,

Tontonoz P, Fisher EA. LXR promotes the maximal egress of monocyte-derived cells

from mouse aortic plaques during atherosclerosis regression. *J Clin Invest*

2010;**120**:4415–4424.

46. Nagareddy PR, Murphy AJ, Stirzaker RA, Hu Y, Yu S, Miller RG, Ramkhelawon B,

Distel E, Westerterp M, Huang LS, Schmidt AM, Orchard TJ, Fisher EA, Tall AR,

Goldberg IJ. Hyperglycaemia promotes myelopoiesis and impairs the resolution of

atherosclerosis. *Cell Metab* 2013;**17**:695–708.

47. Robbins CS, Hilgendorf I, Weber GF, Theurl I, Iwamoto Y, Figueiredo JL, Gorbatov R,

Sukhova GK, Gerhardt LM, Smyth D, Zavitz CC, Shikatani EA, Parsons M, van Rooijen

N, Lin HY, Husain M, Libby P, Nahrendorf M, Weissleder R, Swirski FK. Local

proliferation dominates lesional macrophage accumulation in atherosclerosis. *Nat Med*

2013;**19**:1166–1172.

48. Miller DM, Thomas SD, Islam A, Muench D, Sedoris K. c-Myc and cancer metabolism.

Clin Cancer Res 2012;**18**:5546–5553.

49. Wise DR, DeBerardinis RJ, Mancuso A, Sayed N, Zhang XY, Pfeiffer HK, Nissim I,

Daikhin E, Yudkoff M, McMahon SB, Thompson CB. Myc regulates a transcriptional

program that stimulates mitochondrial glutaminolysis and leads to glutamine addiction.

Proc Natl Acad Sci U S A 2008;**105**:18782–18787.

50. Endo-Umeda K, Kim E, Thomas DG, Liu W, Dou H, Yalcinkaya M, Abramowicz S, Xiao

T, Antonson P, Gustafsson JÅ, Makishima M, Reilly MP, Wang N, Tall AR. Myeloid

LXR (liver X receptor) deficiency induces inflammatory gene expression in foamy

macrophages and accelerates atherosclerosis. *Arterioscler Thromb Vasc Biol* 2022;**42**:719–731.

51. Virani SS, Newby LK, Arnold SV, Bittner V, Brewer LC, Demeter SH, Dixon DL, Fearon WF, Hess B, Johnson HM, Kazi DS, Kolte D, Kumbhani DJ, LoFaso J, Mahtta D, Mark DB, Minissian M, Navar AM, Patel AR, Piano MR, Rodriguez F, Talbot AW, Taqueti VR, Thomas RJ, van Diepen S, Wiggins B, Williams MS; Peer Review Committee Members. 2023 AHA/ACC/ACCP/ASPC/NLA/PCNA guideline for the management of patients with chronic coronary disease: a report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *Circulation* 2023;**148**:e9–e119.

52. Podoltsev NA, Wang R, Shallis RM, Stempel JM, Di M, Neparidze N, Zeidan AM, Huntington SF, Giri S, Hull SC, Gore SD, Ma X. Statin use, survival and incidence of thrombosis among older patients with polycythemia vera and essential thrombocythemia. *Cancer Med* 2023;**12**:18889–18900.

Figure 1

Figure 1. Impaired Atherosclerosis Regression in *Jak2^VF* CH Mice with Moderate Cholesterol Lowering. A. Study Design. **B.** Plasma cholesterol (n = 34, 55, 55, 54, 16, 16 for Control mice, n=31, 55, 49, 39, 17, 16 for *Jak2^VF* mice, for weeks 0, 5, 11, 15, 17.5, 21 respectively). **C.** H&E images of aortic root lesions. Black lines, necrotic core. Scale bar, 200 μm. **D.** Lesion area, n = 15-23. **E.** Necrotic core area, n = 15-23. **F.** Picrosirius red-stained aortic root lesions. Scale bar, 278 μm **G.** Collagen area as a percentage of lesion area, n = 15-24. **H.** Images of aortic root lesions stained for MAC2 (Green) and DAPI (Blue). Scale bar, 100 μm. **I.** Macrophage area, n = 15-24. All quantifications shown as mean ± s.e.m. Two-way ANOVA with the Geisser-Greenhouse correction for sphericity and Tukey's multiple comparisons test (B). Two-way ANOVA with Tukey's multiple comparisons test (D,E,G,I).

Figure 2

Figure 2. Moderate Cholesterol Lowering Reverses Macrophage AIM2 Inflammasome Activation, DNA Damage, and Proliferation in *Jak2^{VF}* CH Lesions. A. Images of aortic root lesions stained for MAC2 (Green), Cleaved GasD (Red), and DAPI (Blue). Scale bar, 127 μm. **B.** $\log_{10}$ transformed cleaved GasD mean fluorescence intensity (MFI) in lesions with the addition of constant 1, n = 15-24. **C.** $\log_{10}$ transformed cleaved GasD mean fluorescence intensity (MFI) in necrotic cores with the addition of the constant 1, n = 15-24. **D.** Images of aortic root lesions stained for MAC2 (Green), Cleaved GasD (Red), and DAPI (Blue). Scale bar, 40 μm. White arrows, cleaved GasD+ macrophages. **E.** $\log_{10}$ transformed cleaved GasD positive macrophages per section with the addition of constant 1, n = 15-24. **F.** Images of aortic root lesions stained for MAC2 (Green), AIM2 (Red), and DAPI (Blue). Scale bar, 75 μm. White arrows, AIM2 positive macrophages. **G.** $\log_{10}$ transformed AIM2 positive macrophages per section with the addition of constant 1, n = 15-23. **H.** Images of aortic root lesions stained for MAC2 (Green), pγH2AX (Red), and DAPI (Blue). Scale bar, 46 μm. White arrows, pγH2AX positive cells. **I.** $\log_{10}$ transformed pγH2AX positive macrophages per section with the addition of constant 1, n = 15-23. **J.** Images of aortic root lesions stained for MAC2 (Green), Ki67 (Red), and DAPI (Blue). Scale bar, 25 μm. White arrows, Ki67 positive macrophages. **K.** $\log_{10}$ transformed Ki67 positive macrophages per section with the addition of constant 1, n = 15-23. **L.** Images of aortic root lesions stained for MAC2 (Green), 8OHdG (Red), and DAPI (Blue). Scale bar, 78 μm. **M.** Percentage of MAC2 positive area double positive for 8OHdG and MAC2, n = 14-24. MAC2 intensity differences reflect independent staining and imaging sessions across panels. All quantifications shown as mean ± s.e.m. Two-way ANOVA with Tukey's multiple comparisons test (B,C). Kruskal-Wallis test with Dunn's multiple comparisons test (E,G,I,K,M).

Figure 3

Figure 3. Moderate Cholesterol Lowering Reverses Impaired Efferocytosis in *Jak2^{VF}* CH

Lesions While Increasing TREM2^{Hi} Macrophages in Control and *Jak2^{VF}* CH Lesions A.

Images of aortic root lesions stained for MAC2 (Green), MerTK (Red), and DAPI (Blue). Scale

bar, 102 μm. **B.** Percentage of MAC2 positive area double positive for MerTK and MAC2, n =

15-23. **C.** Images of aortic root lesions stained for MAC2 (Green), TREM2 (Red), and DAPI

(Blue). Scale bar, 100 μm. **D.** Percentage of MAC2 positive area double positive for TREM2 and

MAC2, n = 15-24. **E.** Images of in situ efferocytosis in aortic root lesions: MAC2 (Green),

TUNEL (Red), and DAPI (Blue). Scale bar, 40 μm. White arrows, nuclei double positive for

TUNEL and MAC2. White wedges, nuclei positive for TUNEL but negative for MAC2. **F.**

Percentage of TUNEL positive nuclei also positive for MAC2. MAC2 intensity differences

reflect independent staining and imaging sessions across panels, n = 13-24. All quantifications

shown as mean ± s.e.m. Two-way ANOVA with Tukey's multiple comparisons test (D). Kruskal-

Wallis test with Dunn's multiple comparisons test (B,F).

Figure 4

Figure 4. Aggressive Cholesterol Lowering Normalizes Regression in *Jak2^{VF}* CH Lesions. A.

Study Design. **B.** Plasma cholesterol (n = 5, 38, 19, 18, 18 for Control mice, n = 5, 37, 15, 16, 15

for Jak2VF mice, for weeks 0, 4, 13, 15, 18 respectively). **C.** H&E images of aortic root lesions.

Black lines, necrotic core. Scale bar, 200 μm. **D.** Lesion area, n = 13-20. **E.** Necrotic core area, n

= 13-20. **F.** Picrosirius red-stained aortic root lesions. Scale bar, 219 μm **G.** Collagen area as a

percentage of lesion area, n = 13-20. **H.** Images of aortic root lesions stained for MAC2 (Green)

and DAPI (Blue). Scale bar, 193 μm. **I.** Macrophage area, n = 13-20. All quantifications shown

as mean ± s.e.m. Two-way ANOVA with Tukey's multiple comparisons test (D,G,I). Two-way

ANOVA with the Geisser-Greenhouse correction for sphericity and Tukey's multiple

comparisons test (B). Kruskal-Wallis test with Dunn's multiple comparisons test (E)

Figure 5

Figure 5. Cholesterol Lowering Suppresses *Jak2^VF* Macrophage Proliferation and DNA Damage. A. Study design. **B.** Plasma cholesterol (n = 80, 79, 79 for Baseline mice for weeks 3, 11, 15 respectively; n = 13, 15 for Control Progression mice, n = 16, 16 for Control Regression mice, n = 15, 15 for *Jak2^VF* Progression mice, n = 16, 16 for *Jak2^VF* Regression mice, for weeks 18, 23 respectively) **C.** Images of aortic root lesions for MAC2 (Green), Cleaved GasD (Red), and DAPI (Blue). Scale bar, 98 µm. **D.** Log_{10} transformed cleaved GasD mean fluorescence intensity (MFI) in lesions with the addition of constant 1, n = 14-16. **E.** Images of aortic root lesions for MAC2 (Red), Ki67 (White), and ZsGreen (Green). Scale bar, 133 µm. White arrows, macrophages double positive for Ki67 and ZsGreen. **F.** Macrophages positive for both Ki67 and ZsGreen per section, n = 14-15. **G.** Macrophages positive for Ki67 but negative for ZsGreen per section, n = 13-16. **H.** Images of aortic root lesions stained for pγH2AX (Red), ZsGreen (Green), and DAPI (Blue). Scale bar, 75 µm. White arrows, pγH2AX positive cells. **I.** Log_{10} transformed cells double positive for pγH2AX and ZsGreen in lesions with the addition of constant 1, n = 13-16. **J.** Log_{10} transformed cells positive for pγH2AX but negative for ZsGreen in lesions with the addition of constant 1, n = 13-16. All quantifications shown as mean ± s.e.m. One-way ANOVA with Holm-Sidak's multiple comparisons test (D). Kruskal-Wallis test with Dunn's multiple comparison's test (F,G,I,J).

Figure 6

Figure 6. Cholesterol Lowering Increases c-myc in TREM2^{Hi} Macrophages. A. Images of

aortic root lesions stained for MAC2 (Green), c-myc (Red), and DAPI (Blue). Scale bar, 121 μm.

B. Percentage of macrophages positive for c-myc, n = 14-16. **C.** Images of aortic root lesions

stained for MAC2 (Green), c-myc (Blue), and TREM2 (Red). Scale bar, 121 μm. **D.** Percentage

of TREM2 positive macrophages positive for c-myc, n = 14-16. All quantifications shown as

mean ± s.e.m. One-way ANOVA with Holm-Sidak's multiple comparisons test (B). Kruskal-

Wallis test with Dunn's multiple comparisons test (D).

44