Aggressive Cholesterol Lowering Normalizes Atherosclerosis Regression in *Jak2^{V617F}* Mice

Brian D. Hardaway (MPhil)[1], Trevor P. Fidler (PhD)[2], Mojdeh Tavallaie (PharmD-PhD)[1], Kleopatra

Avrampou (PhD)[1], Cheng-Chieh Hsu (PhD)[1], Sandra Schiavone (MA)[1], Tong Xiao (MS)[1], Nan Wang

(PhD)[1], Alan R. Tall (MD)[1]

[1]Division of Molecular Medicine, Department of Medicine, Columbia University Irving Medical Center,

New York, NY, USA.

[2]Cardiovascular Research Institute, University of California San Francisco, San Francisco, CA, USA.

Brian D. Hardaway: bdh2138@cumc.columbia.edu

Trevor P. Fidler: Trevor.Fidler@ucsf.edu

Mojdeh Tavallaie: mt3591@cumc.columbia.edu

Kleopatra Avrampou: ka2897@cumc.columbia.edu

Cheng-Chieh Hsu: ch3863@cumc.columbia.edu

Sandra Schiavone: sa2876@cumc.columbia.edu

Tong Xiao: tongxiao4009@gmail.com

Nan Wang: nw30@cumc.columbia.edu

Alan R. Tall: art1@cumc.columbia.edu

Correspondence to Alan R. Tall:

630 W. 168th St. Room 8-401

New York, NY 10032

Tel: 1-212-305-5789

Email: art1@cumc.columbia.edu

Running Title: Atherosclerosis Regression in *Jak2*^{V617F} Mice

Sources of Funding

This work was supported by grants from the National Institutes of Health and The National Heart, Lung,

and Blood Institute: HL155431, HL107653HL170157-02, P01HL172741 to Alan R. Tall and HL148071

to Nan Wang.

Abbreviations:

AAV8 - Adeno-Associated Virus serotype 8 vector

acLDL - acetylated LDL

ACVD - atherosclerotic cardiovascular disease

ADAM17 - a disintegrin and metalloproteinase 17

AIM2 - absent in melanoma 2

BMDM - bone marrow-derived macrophages

BMT - bone marrow transplantation

CH - clonal hematopoiesis

CHIP - clonal hematopoiesis of indeterminate potential

FIJI - FIJI Is Just ImageJ

GasD - gasdermin D

hDAD - helper-dependent adenoviral vector

HSC - hematopoietic stem cell

IL-1 - interleukin-1

IL-1β - interleukin-1β

Jak2^{VF} - Jak2^{V617F}

MerTK - mer proto-oncogene tyrosine kinase

MitoTEMPO - mitochondria-targeted TEMPO (a triphenylphosphonium-conjugated nitroxide)

MPN - myeloproliferative neoplasms

MβCD - methyl-β-cyclodextrin

oxPAPC - oxidized 1-palmitoyl-2-arachidonoyl-sn-glycero-3-phosphocholine

pIpC - polyinosinic:polycytidylic acid

pγH2AX - phosphorylated histone H2A.X

TREM2 - triggering receptor expressed on myeloid cells 2

Abstract

Background - The $Jak2^{V617F}$ ($Jak2^{VF}$) mutation is an important cause of both clonal hematopoiesis of indeterminate potential (CHIP) and myeloproliferative neoplasms (MPN). Mouse models of $Jak2^{VF}$ CHIP and MPN show accelerated atherosclerosis progression, driven by macrophage inflammasome activation. We undertook the present study to assess the hypothesis that ongoing inflammation would impede atherosclerosis resolution in $Jak2^{VF}$ mice.

Methods and Results - Chimeric $Jak2^{VF/WT}$ or control WT/WT bone marrow was transplanted into $Ldlr^{-/-}$ mice and, following 13-16 weeks of Western diet-induced atherosclerosis progression, cholesterol was lowered either moderately (to 200-300 mg/dl) or markedly (to 100 mg/dl). With moderate cholesterol lowering, there was impaired resolution of lesions in $Jak2^{VF}$ MPN mice compared to controls. However, with marked cholesterol lowering, progression of lesions was halted in both $Jak2^{VF}$ MPN and control mice while macrophage burden was decreased and lesional collagen was increased similarly in $Jak2^{VF}$ MPN and control mice.

Two mechanisms of low-density lipoprotein (LDL) lowering-induced suppression of inflammation in plaques were implicated: 1) reversal of increased proliferation, DNA damage and Absent in Melanoma 2 (AIM2) inflammasome activation specifically in $Jak2^{VF}$ macrophages and 2) markedly increased macrophage triggering receptor expressed on myeloid cells 2 (TREM2), c-myc expressing macrophages in both $Jak2^{VF}$ and control mice.

Conclusions - Aggressive LDL lowering reverses inflammasome activation and induces pro-resolving changes in macrophages in $Jak2^{VF}$ MPN, halting atherosclerosis progression and promoting features of plaque stabilization. These findings suggest that aggressive LDL cholesterol lowering could reverse atherosclerotic cardiovascular disease (ACVD) risk in individuals with $JAK2^{VF}$ CHIP or MPN.

Keywords: foam cells, inflammation, bone marrow, LDL, vascular biology, inflammasome, atherosclerotic plaque regression, absent in melanoma 2, clonal hematopoiesis, $Jak2^{V617F}$ mutation

1. Introduction

Despite therapeutic advances, atherosclerotic cardiovascular disease (ACVD) remains the leading cause of death in the developed world (1). Although inflammation plays a key role in atherosclerosis progression, cholesterol accumulation remains a prerequisite for lesion development, and low-density lipoprotein (LDL) lowering has consistently shown a benefit in clinical trials (2,3,4).

The role of inflammation in increasing ACVD risk has also been well established. The LoDoCo 2 trial found that the risk of cardiovascular events was reduced in patients with chronic coronary disease treated with colchicine (5). Based on these and other results, the U.S. Food and Drug Administration approved the use of low dose colchicine as the first anti-inflammatory drug for heart disease (6). The CANTOS trial involving the use of interleukin-1β (IL-1β) antibodies showed reduced cardiovascular events in high-risk patients with mean LDL cholesterol levels of 81 mg/dl - 84 mg/dl and a high-sensitivity C-reactive protein (hs-CRP) level of at least 2 mg/L. However, there was a small increase in fatal infections, and the FDA did not approve this treatment for marketing (7). Many observations suggest that hypercholesterolemia is linked to inflammatory risk. For example, statins lower hs-CRP levels (8) and preclinical studies have shown that cholesterol accumulation in plaque macrophages promotes inflammasome activation (9,10,11,12). However, the mechanisms linking hypercholesterolemia, macrophage cholesterol accumulation and inflammation remain incompletely understood, especially in the clinically important context of atherosclerosis regression induced by LDL lowering (13,14,15,16,17,18,19).

Clonal hematopoiesis (CH) arises from leukemogenic mutations in hematopoietic stem cells (HSC) that confer a selective advantage and lead to clonal expansion of blood cells. CH commonly arises from mutations in genes that mediate epigenetic modifications or cytokine signaling (20,21). Clonal hematopoiesis of indeterminate potential (CHIP) is defined as having a mutant allele burden >2% in blood cells in the absence of changes associated with hematopoietic malignancy (20). Amongst CHIP mutations, $JAK2^{VF}$, which increases signaling by hematopoietic cytokines, confers the greatest risk of

ACVD (22,23,24). Moreover, $JAK2^{VF}$ has been detected in 3-4% of a general European population (25).

$JAK2^{VF}$ CHIP and myeloproliferative neoplasms (MPN) appear to form a disease spectrum, with both

increasing atherosclerotic and thrombotic risk (26,27). Similarly, both $Jak2^{VF}$ MPN and low allele burden

$Jak2^{VF}$ CHIP in mice that lack changes in blood cell counts or splenomegaly increase thrombosis and

atherosclerosis progression involving leukocyte mediated inflammatory mechanisms (28,29).

Additionally, we showed in a hypercholesterolemic mouse model that $Jak2^{VF}$ CHIP or MPN exacerbates

atherosclerosis via inflammasome activation (30) and genetic studies suggest a similar role in humans

(31). Based on these findings, we hypothesized that LDL lowering-induced changes in plaques such as

halted progression and beneficial remodeling would be impaired in $Jak2^{VF}$ mice due to persistence of the

underlying mechanisms promoting inflammasome activation. Contrary to our hypothesis, we found that

cholesterol lowering reversed inflammasome associated changes in $Jak2^{VF}$ macrophages and that marked

cholesterol lowering induced beneficial lesional changes similarly in $Jak2^{VF}$ MPN mice and controls.

These studies suggest the importance of rigorous control of LDL cholesterol levels in individuals with

$JAK2^{VF}$ CHIP or MPN.

2. Methods

Several experimental procedures were performed as previously described (16,30).

2.1 Anesthesia and Euthanasia

Mice were anesthetized using 5% isoflurane via inhalation at a flow rate of approximately 3L/min oxygen

using a calibrated vaporizer. Anesthetic depth was assessed using respiratory rate monitoring and the toe-

pinch reflex. Isoflurane was administered acutely during procedures requiring anesthesia, including blood

collection and euthanasia. Blood collection was performed via retro-orbital sinus for plasma, complete

blood counts, and flow cytometric analysis at the indicated timepoints. For euthanasia, mice were first

rendered deeply unconscious with isoflurane, after which cervical dislocation was performed as a

secondary method to ensure death, in accordance with institutional IACUC approval, NIH guidelines, and

the AVMA Guidelines for the Euthanasia of Animals (2020 Edition).

2.2 Mice

All mice used for these studies were male and on a C57BL/6J background. Mice were housed in a

pathogen-free facility under standard conditions of temperature (23°C) with a 12-hour light-dark with ad

libitum food and water access. Cages and water were changed every 7-14 days. All mouse experiments

were approved by the Institutional Animal Care and Use Committee of Columbia University under

protocol number AABG0561 and were conducted in accordance with NIH guidelines and the Guide for

the Care and Use of Laboratory Animals. Mice heterozygous for the *Jak2^{VF}* conditional knock-in allele

were generated as previously described (32). *Jak2^{VF}* mice were crossed to *Mx1-Cre* (The Jackson

Laboratory, B6.Cg-Tg(*Mx1-Cre*)1*Cgn*/J (003556) or *Scl-Cre* (The Jackson Laboratory, C57BL/6-Tg(Tal1-

cre/ERT)42-056Jrg/J (037466) to generate *Mx1-CreJak2^{VF}* and *Scl-CreJak2^{VF}* mice respectively.

CD45.1*Mx1-CreJak2^{VF}* mice were generated by crossing *Mx1-CreJak2^{VF}* mice to CD45.1 (The Jackson

Laboratory B6.SJL-Ptprca Pepcb/BoyJ (002014)) mice. CD45.1/CD45.2*Scl-CreJak2^{VF}* mice were

generated by crossing Scl-*CreJak2^{VF}* mice to CD45.1 (The Jackson Laboratory B6.SJL-Ptprca

Pepcb/BoyJ (002014)) mice. CD45.1/CD45.2*Scl-Cre-Jak2^{VF}* mice were crossed with ZsGreen reporter

mice (The Jackson Laboratory, B6.Cg-Gt(ROSA)26Sor$^{tm6(CAG-ZsGreen1)Hze}$/J (Ai6(RCL-ZsGreen) (007906)

to generate CD45.1/CD45.2*Scl-CreJak2^{VF}ZsGreen* mice. *Ldlr$^{-/-}$* recipient mice for the *Mx1-CreJak2^{VF}*

moderate cholesterol lowering model that included *Myh11-Cre^{ERT2}* and *ZsGreen* alleles from a previous

lineage-tracing study were obtained from a collaborator and were previously described (33). However,

these alleles were not relevant to the experimental outcomes and were not analyzed. Half of recipients

were *Ldlr$^{-/-}$ Myh-11-CreZsGreen* while the other half either lacked the *ZsGreen* allele or both the *ZsGreen*

allele and the *Myh11-Cre*. For the *Mx1-CreJak2^{VF}* and *Scl-CreJak2^{VF}* aggressive cholesterol lowering

studies, *Ldlr$^{-/-}$* recipient mice were purchased from The Jackson Laboratory (B6.129S7-*Ldlr*tm1Her/J

(002207)). Littermate control mice for studies involving *Mx1-CreJak2^{VF}* mice contained either an allele

with *Mx1-Cre* or the *Jak2*[VF] transgene, but not both. Littermate control mice for studies involving *Scl-*

CreJak2[VF]*ZsGreen* mice contained alleles for *Scl-Cre* and *ZsGreen* but not the *Jak2*[VF] transgene.

2.3 Bone Marrow Transplantation

Bone marrow transplantations (BMT) were conducted as previously described with the following

exceptions (10). For the *Mx1-CreJak2*[VF] moderate cholesterol lowering study, donor mice were 8-12

weeks old while recipients were 11-14 weeks old. For the *Mx1-CreJak2*[VF] effective cholesterol lowering

study, donor mice were 7-16 weeks old while recipients were 7 weeks old. For the *Scl-CreJak2*[VF] study,

donor mice were 6-12 weeks old while recipients were 8 weeks old. For the *Mx1-CreJak2*[VF] moderate

cholesterol lowering study, recipients were lethally irradiated once with 10.5 Gy from a cesium gamma

source. Due to decommissioning of the cesium gamma source, recipients from the *Mx1-CreJak2*[VF] and

Scl-CreJak2[VF] effective cholesterol lowering studies were irradiated with 10.5 Gy using a Multirad 350

X-ray Irradiator from Precision X-Ray. Within 24 hours of irradiation, bone marrow was isolated from

donors with the indicated genotypes. Total bone marrow cell number was quantified using an iN CYTO

C-Chip hemocytometer (DHC-N01) according to the manufacturer instructions. Irradiated mice were

randomized to treatment groups, anesthetized with isoflurane before receiving 3×10^6 total bone marrow

cells via intravenous (i.v.) injection at a total final volume of 100 µl. For the *Mx1-CreJak2*[VF] studies,

recipients received either 6×10^5 *CD45.1Mx1-CreJak2*[VF/WT] or 6×10^5 *CD45.1* control bone marrow cells

(20%) combined with 2.4×10^6 *CD45.2Jak2*[WT/WT] cells (80%). For the *Scl-CreJak2*[VF] study, recipients

received either 6×10^5 *CD45.1CD45.2Scl-CreJak2*[VF/WT]*ZsGreen* or 6×10^5 *CD45.1CD45.2Scl-*

CreJak2[WT/WT]*ZsGreen* bone marrow cells (20%) combined with 2.4×10^6 *CD45.1Jak2*[WT/WT] cells (80%).

For the *Mx1-CreJak2*[VF] studies, mice were allowed to recover for four weeks after BMT before being

injected intraperitonially (i.p.) with 50 µg/mouse/day polyinosinic:polycytidylic acid (pIpC) two times, 48

hours apart.

2.4 Atherosclerosis Studies

For the *Mx1-CreJak2*[VF] studies, power analysis using GPower3.1 indicated that with a total sample size of

80 (n = 20 per group), the study had 80% power to detect an effect size of f = 0.32 at α = 0.05, using two-

way analysis of variance (ANOVA) across two genotypes and two treatments. For the *Scl-CreJak2*[VF]

study, power analysis using GPower3.1 indicated that with a total sample size of 80 (n = 16 per group),

the study had 80% power to detect an effect size of f = 0.40 at α = 0.05, using one-way ANOVA across

five groups. Four weeks after BMT, *Ldlr*[-/-] recipient mice were fed a western diet (ENVIGO, cat. no.

TD.88137) for the indicated times. For the final 10 days of the western diet in the *Scl-CreJak2*[VF] study,

mice were switched to a western diet containing tamoxifen at 500 mg/kg (ENVIGO, cat. no. TD130889).

To induce LDL lowering in the *Mx1-CreJak2*[VF] study of Figure 1A, mice were administered 2 x 10^{11} viral

particles per mouse in 200 µl of 1X PBS via tail vein injection of an hDAD-*hLDLR* vector (Gene Vector

Core Laboratory at the Baylor College of Medicine) and switched to chow diet. To induce LDL lowering

in the *Mx1-CreJak2*[VF] study of Figure 4A and the *Scl-CreJak2*[VF] study of Figure 6A, mice were

administered 2.5 x 10^{11} genome copies per mouse in 100 µl of 1X PBS via tail vein injection of an AAV8-

TBG-*mLdlr* vector (AAV-263355, Vector Biolabs) and switched to chow diet. Following euthanasia,

aortic roots were collected and fixed with 4% paraformaldehyde (PFA) (Electron Microscopy Sciences

15710; 16% PFA diluted 1:4 in 1.33X PBS) for 24 hours at 4°C, then embedded in paraffin.

2.5 Histological Analysis

Lesion area and necrotic core area were quantified as previously described (22,30). Blinded researchers

sectioned paraffin-embedded aortic root lesions at similar points between mice over a 150 µm span for a

total of 50 sections spaced 3 µm apart with 2 sections per slide on 25 slides. Six total slides spaced 30 µm

apart were then stained with hematoxylin & eosin (H&E) (Sigma MHS32-1L; Abcam ab246824) and

imaged with an Olympus DP27 camera attached to an Olympus CX43 biological microscope set to the

10X objective using Olympus cellSens Entry 3.1 software. Lesion area was quantified in a blinded

fashion, and the average of the six slides was used to determine lesion area. The Necrotic core area was

similarly quantified in a blinded fashion using the same six H&E sections. Collagen was stained using the

Polysciences Picrosirius Red Stain Kit (cat. no. 24901) according to the manufacturer's instructions. The

same slide between mice was stained and the collagen quantified by applying a consistent color threshold

to calculate the collagen-positive area as a percentage of total lesion area. Picrosirius red was imaged via

the same system used for H&E imaging. All lesion analysis was conducted using FIJI software.

2.6 Immunofluorescence

Identical slides of aortic roots between mice were baked in an incubator at 60°C for 30 minutes and

deparaffinized in Histo-clear (National Diagnostics HS-200/50-899-90147) three times for 10 minutes.

Histo-clear was removed by 20 shakes in 100% ethanol twice followed by 20 shakes in Histoprep

Reagent Alcohol (fisher HC-600-1GAL) before being washed in tap water 7 times. Antigen retrieval was

performed by placing slides either in citric acid-based antigen retrieval solution (Vector Laboratories,

H3300) or tris-based antigen retrieval solution (Vector Laboratories, H3301) and pressure cooking for 15

minutes with a 5-minute natural release in an Instant Pot IP-LUX pressure cooker. Citric acid-based

antigen retrieval was performed for all antibodies except for the antibody against cleaved GasD. After

pressure cooking, slides were cooled for 10 minutes and then washed 3 times for 5 minutes in 1X

phosphate buffered saline (PBS) (Boston Bioproducts BM-220; diluted 1:10). Hydrophobic circles were

drawn using a liquid blocker pen (Sigma Z377821). For antibodies sourced from mice, mouse-on-mouse

blocking was performed by incubating sections in 10% goat serum (ThermoFisher 10000C) in 1X PBS

(Corning 21-040-CM) with 1 drop of mouse-on-mouse block per 1.25ml of solution (Vector Laboratories

NC9290646) for 1 hour at room temperature in a humidified chamber. Slides were then washed in 1X

PBS containing tween 20 (Fisher BP337-500) at a concentration of 0.01% (PBS-T) twice for 5 minutes

and then in 1X PBS once for 5 minutes. For other antibodies, sections were blocked by incubating in

Normal Serum Block (Biolegend 927503) for 1 hour at room temperature in a humidified chamber.

Following blocking, sections were incubated with the following primary antibodies at the indicated

concentrations overnight at 4°C in a humidified chamber: Absent in melanoma 2 (AIM2) (Abcam,

ab119791, 1:250), Cleaved GasD (Cell Signaling, 10137, 10µg/ml), c-myc (Cell Signaling, 5605, 5.76

µg/ml), Ki67 (Abcam, ab15580, 9 µg/ml), MAC2 conjugated to Alexa Fluor 488 (Cedarlane,

CL8942AF4, 1 µg/ml), MAC2 (Cedarlane, CL8942AP, 1 µg/ml), mer proto-oncogene tyrosine kinase

(MerTK) (R&D, BAF591, 2 µg/ml), pγH2AX (Cell Signaling, 9718, 0.74 µg/ml), ZsGreen

(ThermoFisher, TA180002, 10 µg/ml), triggering receptor expressed on myeloid cells 2 (TREM2)

(Denali, 4D9 DC1847, 1:100). The following IgG antibodies were used as negative controls for primary

antibodies at the same concentration: Mouse IgG (ChromPure, 015-000-003), Rabbit IgG (Abcam,

ab172730), Rabbit IgG (Novus Biologicals, NBP2-24891), Rabbit IgG (ChromPure 011-000-003), Rat

IgG (ChromPure, 012-000-003), Rat IgG (Biolegend, 400431), Rat IgG conjugated to Alexa Fluor 488

(Invitrogen, 53-4321-80). Slides were then washed twice with PBS-T for 5 minutes and once with 1X

PBS for 5 minutes. The following secondary antibodies were used at a dilution of 1:200 for 1 hour at

room temperature: Rat conjugated to Alexa Fluor 488 (Invitrogen A11006), Rat conjugated to Alexa Fluor

568 (Invitrogen A11077), Rat conjugated to Alexa Fluor 647 (Invitrogen A-21247), Rabbit conjugated to

Alexa Fluor 568 (Invitrogen A11011), Rabbit conjugated to Alexa Fluor 647 (A27040). The following

secondary antibodies were used at a dilution of 1:800 for 1 hour at room temperature: Mouse conjugated

to Alexa Fluor 647 (Invitrogen A21237), Streptavidin conjugated to Alexa Fluor 568 (Invitrogen S11226),

Streptavidin conjugated to Alexa Fluor 647 (Invitrogen S32357). DAPI (4',6-Diamidino-2-Phenylindole,

Dilactate) (Biolegend, 422801) was included during incubation with secondary antibodies at a

concentration of 5 µg/ml. Slides were then washed twice with PBS-T for 5 minutes and once with 1X

PBS for 5 minutes. After washing, slides were mounted using Prolong Gold Antifade Reagent with DAPI

(Invitrogen P36935). TUNEL staining was performed using the In Situ Cell Death Detection Kit TMR

Red (Roche, 12156792910) according to the manufacturer's instructions. Slides were imaged on a Leica

Fluorescent DMI 6000B wide-field microscope or a Nikon Ti Eclipse inverted microscope with high-

sensitivity confocal imaging at the Confocal and Specialized Microscopy Shared Resource (CSMSR)

Core at Columbia University. All images were analyzed in a blinded fashion using FIJI software.

2.7 Blood Isolation and Analysis

Blood was collected at the indicated timepoints via retro-orbital sinus using heparinized microhematocrit

capillary tubes (Fisherbrand 22-362566) into EDTA-coated microvette tubes (Sarstedt 20.1278.100).

Complete blood cell counts were quantified using a VetScan HM5 Hematology system (Zoetis/Abaxis).

Plasma was collected by centrifuging blood at 13,000 x g for 10 minutes at 4°C. Plasma cholesterol was

quantified using the Cholesterol E kit (Fujifilm Wako Pure Chemical Corporation, #999-

02601/NC9138103) according to the manufacturer's instructions. Flow cytometric analysis for blood

leukocytes was performed by lysing in RBC lysis buffer (Biolegend 420302) for 2 minutes at room

temperature before centrifuging at 3,000 x g for 2 minutes at 4°C. Cells were washed in ice cold MACS

Buffer (0.5% BSA and 2mM EDTA in 1X PBS) (FischerScientific BP9706-100, Invitrogen 15575-038)

and centrifuged at 800 x g for 10 minutes at 4°C before staining with antibodies at a 1:100 dilution in ice

cold MACS buffer for 20 minutes. Cells were washed once more in ice cold MACS Buffer, centrifuged at

800 x g for 10 minutes at 4°C before being resuspended in 250 µl of MACS Buffer. Antibodies used in

each experiment are listed in Supplementary Tables S1–S3. Cells were captured using a BD Fortessa

Flow Cytometer running BD FACSDiva software at the Flow Cytometry Shared Core of the Columbia

Center for Translational Immunology (CCTI) and Herbert Irving Comprehensive Cancer Center

(HICCC). Flow cytometric data were analyzed using FCS Express 7 Research Edition.

2.8 Bone Marrow-Derived Macrophage Cultures

Hindlimbs were isolated and stored in ice cold DMEM. The ends of the bones were cut and the bones

placed into 600 µl microcentrifuge tubes with a hole punctured at the bottom and sitting inside of 1.5 ml

microcentrifuge tubes containing 100 µl of ice cold DMEM. The bones were pulse centrifuged at 4°C for

8 seconds. The pellet was then suspended in 1.5 ml of ice cold DMEM before being filtered through a 40

µm cell strainer (Fisher 22-363-547) into 50 ml of ice cold DMEM. Suspensions were centrifuged at 800

x g for 10 minutes at 4°C and the supernatant aspirated. The pellet was resuspended in M-CSF complete

media for the *Jak2^{VF}Aim2^{-/-}* experiment (DMEM + 20 ng/ml M-CSF + 10% FBS + 1% PENSTREP) (M-

CSF from Cell Signaling, 33444S or LCM complete media (all other experiments) (DMEM + 20% LCM

+ 10% FBS + 1% PENSTREP) and plated into tissue culture treated dishes (Corning CLS430167). Cells

were differentiated into macrophages by incubating for 5 days at 37°C and 5% CO_2. After 5 days, the

media was aspirated and cells washed in 20 ml of 1X PBS before being stripped by incubation in 5 ml of

Corning Cell Stripper (Corning 25-056-CL) for 5 minutes before being collected using 5 ml of complete

media. Cells were then centrifuged at 800 x g for 10 minutes at 4°C and the supernatant aspirated. Cells

were counted using an iN CYTO C-Chip hemocytometer (DHC-N01) according to the manufacturer

instructions. Cells were plated into well plates of various sizes depending on experiment at a density of

333,333 cells/ml (3 ml of suspension for 6-well plates, 2 ml of suspension for 12-well plates). Cells were

allowed to reattach overnight before proceeding with further experiments.

2.9 acLDL Cell Culture Experiments

Preparation of acetyl-LDL (acLDL) was performed as previously described (34). Bone marrow-derived

macrophages (BMDMs) were differentiated and plated as described in section 2.8 and then incubated in

M-CSF-free media containing either vehicle or 25 µg/ml acLDL overnight at 37°C and 5% CO_2.

2.10 oxPAPC Cell Culture Experiments

BMDMs were differentiated and plated as described in section 2.8 and then incubated in M-CSF-free

media containing either vehicle or 50 µg/ml oxidized 1-palmitoyl-2-arachidonoyl-sn-glycero-3-

phosphocholine (oxPAPC) overnight at 37°C and 5% CO_2.

2.11 MβCD-Cholesterol Cell Culture Experiments

BMDMs were differentiated and plated as described in 2.8. Cells were then primed in complete media

containing 100 ng/ml lipopolysaccharide (LPS) (Cell Signaling 14011) for 3 hours at 37°C and 5% CO_2

before treating with complete media containing vehicle, (2-(2,2,6,6-tetramethylpiperidin-1-oxyl-4-

ylamino)-2-oxoethyl)triphenylphosphonium chloride (mitoTEMPO) (MedChemExpress HY-112879) (10

µM), methyl-β-cyclodextrin-cholesterol (MβCD-Cholesterol) (Sigma C4951) (30 µg/ml) or a

combination of mitoTEMPO (10 µM) and MβCD -Cholesterol (30 µg/ml) for 4 hours at 37°C and 5%

CO_2. LPS remained in the media for the duration of these treatments at 100 ng/ml.

2.12 Bone Marrow-Derived Macrophage Proliferation Quantification

BMDM proliferation was measured using the Invitrogen Click-iT™ Plus EdU Alexa Fluor™ 488 Flow

Cytometry Assay Kit according to the manufacturer's instructions. After incubation with EdU, cells were

washed in 1X PBS before being stripped by incubation in 500 µl of Corning Cell Stripper (Corning 25-

056-CL) for 5 minutes. Cells were captured using a BD Fortessa Flow Cytometer running BD FACSDiva

software at the Flow Cytometry Shared Core of the Columbia Center for Translational Immunology

(CCTI) and Herbert Irving Comprehensive Cancer Center (HICCC). Flow cytometric data were analyzed

using FCS Express 7 Research Edition.

2.13 Immunoblotting

Cells were lysed in radioimmunoprecipitation assay (RIPA) buffer containing protease and phosphatase

inhibitors (Invitrogen 87786, ThermoFisher 78420) for 30 minutes at 4°C on a rocker. Lysates were then

frozen at -80°C to enhance lysis. Lysates were then thawed at 4°C, centrifuged at max speed for 5 minutes

at 4°C before being sonicated at 50% power for 30 seconds. Total protein concentration was determined

by h bicinchoninic acid (BCA) analysis (ThermoFisher Scientific, 23225) according to the manufacturer's

instructions. Equal protein amounts were then loaded into polyacrylamide gels (Biorad 5671094) using

1X Running Buffer (Boston Bioproducts BP-150) and transferred to nitrocellulose membranes (Biorad

1620115) using transfer buffer (Boston Bioproducts BP-190) with 20% ethanol (Fisher 2701).

Membranes were washed in Tris-Buffered Saline with Tween 20 (TBS-T) (Boston Bioproducts IBB-

180X) for 5 minutes at room temperature before blocking for 1 hour at room temperature in 3% bovine

serum albumin (BSA) (Fisher BP9706-100) in TBS-T while gently rocking. After blocking, membranes

were incubated overnight at 4°C in 1:1000 primary antibody solution in 3% BSA in TBS-T with the

following antibodies: pYH2AX (Cell Signaling 9718) and total GasD (Cell Signaling 38754) or for 20

minutes at room temperature in 1:80,000 primary antibody solution with the following antibodies: β-

Actin (Cell Signaling 12262). After incubation with non-HRP-conjugated primary antibodies, membranes

were washed in TBS-T 5 times for 5 minutes before incubation in a horseradish peroxidase (HRP)

conjugated anti-rabbit secondary antibody (7074 Cell Signaling Technology) in 3% BSA in TBS-T for 1

hour at room temperature while gently rocking. After HRP-conjugated primary or secondary antibody

incubation, membranes were washed 5 times for 5 minutes before incubation for 30 seconds with a

chemiluminescent substrate (ThermoFisher 34578). Membranes were then exposed in a dark room using

film (Fisher XAR ALF 2025). Densitometric quantification was performed using FIJI software.

2.14 IL1β ELISA Detection

IL1β detection was quantified using the Mouse IL-1 beta/IL-1F2 DuoSet enzyme-linked immunosorbent

assay (ELISA) (R&D Systems DY401) according to the manufacturer's instructions. The plate was

detected using a SpectraMax M2 (Molecular Devices).

2.15 Statistics

Statistical analyses were conducted using GraphPad Prism v10.5. Data are presented as the mean $\pm$ s.e.m.

All statistical tests were performed using two-sided analyses. Normality was tested using the D'Agostino-

Pearson omnibus and, where appropriate, log transformation was applied to approximate normality.

Repeated measures data were analyzed using two-way analysis of variance (ANOVA) with the Geisser-

Greenhouse correction for sphericity and Tukey's multiple comparisons test with individual variance

estimation. For datasets involving two independent variables, two-way ANOVA with Tukey's multiple

comparisons test or Kruskal-Wallis with Dunn's multiple comparisons test were used based on

distribution. For datasets involving five experimental groups, one-way ANOVA with Holm-Sidak's

multiple comparisons test or Kruskal-Wallis with Dunn's multiple comparisons test were used based on

distribution. Formal outliers were removed using a two-tailed Grubbs' test ($\alpha = 0.05$). For experiments

with technical replicates from a single mouse, 3–6 wells were plated per condition. Wells were excluded

as technical outliers when 6 wells were plated if the coefficient of variation (%CV) among wells for a

given condition exceeded 20% (predefined QC threshold).

3. Results

3.1 Atherosclerosis Regression is Impaired with Moderate Cholesterol Lowering in *Jak2^VF* Mice

To assess the effect of *Jak2^VF* MPN on atherosclerosis regression, we placed *Jak2^VF* MPN male mice and

controls on a western diet for 16 weeks and then induced cholesterol lowering using an hDAD vector

containing the human LDL receptor gene (hDAD-*hLDLR*) combined with a switch to chow diet for an

additional 5 weeks (Figure 1A). Prior studies were performed in female mice (16,30). *Jak2^VF* MPN male

mice exhibited splenomegaly, lower body weight, and increased RBCs, hematocrit, red blood cell

distribution width (RDWc), and neutrophils, mirroring previous findings in females (Supplemental Figure

1A-I) (30). In this model, CD45.1+ cells are *Jak2^VF* in experimental mice and WT in controls. CD45.1+

neutrophils and monocytes were elevated while CD45.1+ lymphocytes were reduced (Supplemental

Figure 1J-L). Plasma cholesterol was reduced in *Jak2^VF* MPN mice on Western diet (Figure 1B) similar to

previous findings in female mice and humans (24,30,35). Plasma cholesterol remained moderately

elevated for 10 days post-virus treatment but returned to pre-diet levels (~200-300 mg/dl) by the end of

the LDL lowering period (Figure 1B). At baseline, lesion area and necrotic core area were significantly

increased in the aortic root of male *Jak2^VF* MPN mice compared to controls, confirming that *Jak2^VF* MPN

worsens atherosclerosis in both male and female mice (Figure 1C-E) (30). Moderate cholesterol lowering

halted progression of lesion and necrotic core area in control mice whereas *Jak2^VF* MPN mice developed

significantly larger necrotic cores and displayed a trend toward increased lesion area (p=0.052) (Figure

1C-E). Consistent with previous studies (36,37), lesional collagen was increased in control mice after

cholesterol lowering with no significant change in *Jak2^VF* MPN mice, further indicating impaired

resolution in the *Jak2^VF* MPN mice (Figure 1F,G). The most consistent phenotype found in atherosclerosis

regression studies has been a decrease in macrophage burden (13,17,18,19,37). Macrophages were

significantly decreased in control mice but not in *Jak2^VF* MPN mice after cholesterol lowering (Figure

1H,I). These results indicate that atherosclerosis resolution is impaired in *Jak2*[VF] MPN with moderate

cholesterol lowering.

We next assessed inflammatory features in plaques. Necrotic, inflammatory cell death is mediated by

activated Gasdermin D (GasD) downstream of inflammasome activation (38). We used an antibody to the

active N-terminal fragment of GasD to assess pyroptosis in plaques (29). Total cleaved GasD in plaques

as well as cleaved GasD in the necrotic core were increased in *Jak2*[VF] MPN lesions at baseline and were

not reduced by cholesterol lowering (Figure 2A-C). However, macrophage-specific cleaved GasD was

decreased by cholesterol lowering in *Jak2*[VF] MPN lesions, suggesting that macrophage pyroptosis had

been suppressed by the time of harvest (Figure 2D,E). Consistent with earlier findings in females, absent

in melanoma 2 (AIM2)+ macrophages and phosphorylated histone H2A.X (pγH2AX), an indicator of

double stranded DNA break formation, were increased in *Jak2*[VF] MPN lesions at baseline and were

decreased by cholesterol lowering, suggesting reversal of AIM2 inflammasome activation (Figure 2F-I)

(30). Additionally, macrophage proliferation trended higher at baseline in *Jak2*[VF] MPN mice and was

significantly reduced by cholesterol lowering (Figure 2J,K). These data suggest reversal of macrophage

AIM2 inflammasome activation and pyroptosis by moderate cholesterol lowering in *Jak2*[VF] MPN mice,

correlating with decreased macrophage proliferation and decreased DNA damage response. The

persistence of cleaved GasD in the necrotic core area may reflect ongoing pyroptosis during the early

phase of LDL lowering when cholesterol levels were persistently elevated.

3.2 Cholesterol Lowering Restores Impaired Efferocytosis and increases TREM2+ Macrophages in

***Jak2*[VF] mice.**

LDL Lowering has been associated with improvements in pro-resolving features of lesions such as

increased macrophage efferocytosis, mer proto-oncogene tyrosine kinase (MerTK) expression (13,39,40),

increased numbers of triggering receptor expressed on myeloid cells 2 (TREM2)+ macrophages and

limitation of necrotic core formation in plaques (16,41). During atherosclerosis progression, macrophage

MerTK and TREM2 were suppressed in *Jak2*[VF] CH mouse plaques as a result of interleukin-1 (IL-1)

signaling from mutant to wild-type (WT) cells (29). Consistent with these studies, we found that

macrophage MerTK was decreased in $Jak2^{VF}$ MPN lesions at baseline compared to controls (Figure

3A,B). MerTK levels were completely restored by cholesterol lowering (Figure 3A,B). Additionally, we

found that cholesterol lowering dramatically increased the percentage of macrophages positive for

TREM2 in both control and $Jak2^{VF}$ MPN lesions (Figure 3C,D). Consistent with these findings,

macrophage-associated TUNEL+ nuclei as a percentage of total lesional TUNEL+ nuclei, an estimate of

in situ efferocytosis (14), were significantly decreased in $Jak2^{VF}$ MPN lesions at baseline and restored by

cholesterol lowering, (Figure 3E,F). Together, these data suggest that cholesterol lowering restores

impaired efferocytosis in $Jak2^{VF}$ MPN lesions and increases the proportion of inflammation-resolving

macrophages in lesions. Thus, macrophage inflammasome activation and associated defects in

efferocytosis were reversed by cholesterol lowering in $Jak2^{VF}$ MPN mice. The increased necrotic core and

decreased collagen content in these mice could reflect limited resolution of hypercholesterolemia during

the earlier phases of LDL lowering.

3.3 More Effective Cholesterol Lowering Restores Atherosclerosis Regression

To test this hypothesis, we performed a similar study but with more effective LDL cholesterol lowering.

Murine LDL does not bind as strongly to the human LDLR (employed in Figure 1) as to the mouse LDLR

(42,43). Therefore, we employed an AAV8 containing the mouse *Ldlr* gene which more effectively and

more quickly lowered cholesterol levels (Figure 4A). Blood parameters were similar to the first

experiment and not appreciably changed by cholesterol lowering (Supplemental Figure 2, Figure 4B).

Consistent with the first study, lesion area and necrotic core area were significantly increased in $Jak2^{VF}$

MPN mice at baseline. However, in contrast to the first study, while both groups showed non-significant

upward trends of 1.29-fold (Ctrl) and 1.25-fold ($Jak2^{VF}$) in lesion area, there was no change of necrotic

core area in either group after aggressive LDL lowering (Figure 4C-E). Furthermore, aggressive

cholesterol lowering significantly increased lesional collagen and decreased macrophage burden in both

groups (Figure 4F-I). Analysis of cleaved GasD showed an increase in macrophages in $Jak2^{VF}$ mice at

baseline that was reversed by cholesterol lowering, and in contrast to the first study a decrease in cleaved

GasD in the necrotic core that appeared similar in control and $Jak2^{VF}$ MPN mice (Supplemental Figure

3A-D). Similarly, cleaved GasD was increased in CD11b+ splenocytes of $Jak2^{VF}$ MPN mice at baseline

and reversed by cholesterol lowering (Supplemental Figure 3E,F). Consistent with the first study, double

stranded DNA breaks (pγH2AX) were increased in $Jak2^{VF}$ MPN lesions at baseline and more strongly

decreased by aggressive cholesterol lowering along with a significant decrease in macrophage

proliferation (Supplemental Figure 3G-J). Additionally, macrophage MerTK was increased by aggressive

cholesterol lowering (Supplemental Figure 3K,L). These results indicate that more effective cholesterol

lowering completely reversed inflammatory features and induced similar levels of atherosclerosis

resolution in control and $Jak2^{VF}$ MPN lesions.

3.4 Cholesterol Lowering Suppresses Proliferation of $Jak2^{VF}$ Macrophages and increasesTREM2+

Macrophages

In the previous studies $Jak2^{VF}$ was active during progression and lesion area was increased at baseline. To

simulate the situation where the $Jak2^{VF}$ mutation was acquired later in life after atherosclerosis had

already developed, we performed a third study in which $Jak2^{VF}$ was activated in HSCs at the time of LDL

lowering by using a tamoxifen-inducible Scl-driven cre recombinase along with a floxed ZsGreen

reporter allele ($Jak2^{VF}$ Scl CH) to assess cell genotype specific changes (Figure 5A). In addition to the two

LDL lowering groups, we included control and $Jak2^{VF}$ mice that remained on the western diet

(Progression groups). As expected, plasma cholesterol remained elevated in the Progression groups but

was markedly and rapidly decreased in both LDL lowering groups (Figure 5B). The $Jak2^{VF}$ MPN

phenotype was partially apparent as indicated by increased spleen weights, RDWc, and hematocrit as well

as expansion of $Jak2^{VF}$ (CD45.1+CD45.2+) blood monocytes and neutrophils, but less pronounced than in

the earlier studies when $Jak2^{VF}$ was activated prior to Western diet feeding (Supplemental Figure 4).

Compared to the earlier studies, the impact of $Jak2^{VF}$ on lesion progression was less pronounced

reflecting the shorter period of $Jak2^{VF}$ activation (Supplemental Figure 5A-C). However, consistent with

the earlier studies, the necrotic core significantly worsened in the $Jak2^{VF}$ high LDL mice but not in the

control mice compared to baseline, while the necrotic core was not significantly changed in either LDL

lowering group compared to baseline (Supplemental Figure 5C). Additionally, macrophage burden was

similarly decreased in both LDL lowering groups, (Supplemental Figure 5D,E). These studies further

support that aggressive cholesterol lowering similarly promotes resolution in control and $Jak2^{VF}$ mice

with acquisition of $Jak2^{VF}$ in the context of established atherosclerosis.

Cleaved GasD was increased in $Jak2^{VF}$ lesions compared to baseline and trended lower (p=.06) after

cholesterol lowering, consistent with aggressive cholesterol lowering reversing pyroptosis in lesions

(Figure 5C,D). In the $Jak2^{VF}$ Progression group, proliferation was significantly increased in $Jak2^{VF}$

(green) macrophages but not in WT (non-green) macrophages indicating a mutant cell-specific effect, as

seen previously (30). Importantly, this increased proliferation of $Jak2^{VF}$ cells was reversed by cholesterol

lowering, (Figure 5E-G). The DNA damage response (pγH2AX) was significantly increased in $Jak2^{VF}$

cells and was reversed by cholesterol lowering while WT cells showed no significant proliferative

changes or changes in the DNA damage response within $Jak2^{VF}$ lesions (Figure 5H-J).

Given that a proliferative efferocytic population with high c-Myc expression has been described with

LDL lowering (13,39), we assessed c-Myc expression along with TREM2. In contrast to the decreases in

Ki67, macrophage c-Myc and c-Myc+TREM2^{Hi} macrophages were increased by aggressive LDL

lowering in both control and $Jak2^{VF}$ mice, suggesting a selective increase of TREM2+, anti-inflammatory

macrophages despite a decrease in total macrophages (Figure 6).

3.5 Cholesterol Loading Increases AIM2 Inflammasome Activation and the DNA Damage Response

in Macrophages.

The parallel decrease in proliferation and DNA damage response in $Jak2^{VF}$ macrophages (Fig 5H-J)

suggested that reversal of proliferation could be a key mechanism to explain reduced inflammasome

activation in response to cholesterol lowering. To further assess the relationship between proliferation, the

DNA damage response and inflammasome activation, we carried out mechanistic studies in bone marrow-

derived macrophages (BMDMs). We found that cholesterol loading of macrophages with acetylated LDL

(acLDL), which promotes entry of cholesterol via the scavenger receptor A/endo-lysosomal system (44),

caused a modest increase in proliferation (Figure 7A). While $Jak2^{VF}$ macrophages showed significantly

increased pγH2AX compared to controls, there was no effect of acLDL loading on pγH2AX (Figure

7B,C). We also showed that oxidized 1-palmitoyl-2-arachidonoyl-sn-glycero-3-phosphocholine

(oxPAPC), a major component of oxidized LDL, stimulated macrophage proliferation, but this was

associated with a reduced level of pγH2AX (Figure 7D-F). We next employed methyl-β-cyclodextrin

(MβCD)-cholesterol, a distinctive mode of cholesterol loading that increases mitochondrial cholesterol

content (45), promoting mitochondrial reactive oxygen species (ROS) formation and AIM2

inflammasome activation (46). MβCD-cholesterol increased IL-1β secretion into cell culture media

especially in $Jak2^{VF}$ macrophages, and this was reversed by the mitochondrial-specific antioxidant

mitochondria-targeted TEMPO (a triphenylphosphonium-conjugated nitroxide) (MitoTEMPO) (Figure

7G), suggesting that the effect was dependent on mitochondrial ROS production, consistent with prior

studies (46). The increase in IL-1β secretion in $Jak2^{VF}$ macrophages in response to MβCD-cholesterol was

reversed by AIM2 deficiency implicating the AIM2 inflammasome (Figure 7H). MβCD-cholesterol

increased pγH2AX in control and $Jak2^{VF}$ macrophages, with a larger effect in the latter (two-way

ANOVA) (Figure 7I,J). In contrast, MβCD-cholesterol loading did not increase macrophage proliferation

(Figure 7K). Together these findings suggest that increased AIM2 inflammasome activation in response to

MβCD-cholesterol loading may largely reflect increased mitochondrial ROS and DNA damage, rather

than changes in cell proliferation.

4. Discussion

Based on prior studies (30), we hypothesized that LDL lowering alone would not prevent plaque

progression in $Jak2^{VF}$ mice. While this proved to be the case with moderate LDL lowering, aggressive

cholesterol lowering halted progression and induced beneficial remodeling changes, including decreased

macrophage burden and increased lesional collagen content similarly in controls and $Jak2^{VF}$ MPN mice.

Surprisingly, LDL lowering reversed the major cellular changes in $Jak2^{VF}$ macrophages that have been

associated with athero-progression. This included reduced AIM2 expression and pyroptosis and increased

MerTK, TREM2 and in situ efferocytosis that have all been mechanistically linked in prior studies of

$Jak2^{VF}$ mice (29). Our prior studies have shown that inflammasome activation leads to reduced levels of

MerTK and TREM2 (29), suggesting that increases in these factors during cholesterol lowering may be

secondary to reduced inflammasome activation.

Prior studies have shown that LDL lowering in mice with established atherosclerosis may induce

atherosclerosis regression, limitation of progression, or beneficial remodeling involving multiple

mechanisms (47) including macrophage emigration (48), decreased monocyte entry (49) and a role of

acquired immunity and T-regs (17). Our studies using a cleaved GasD antibody to monitor pyroptosis in

lesions, indicated that reversal of inflammasome activation may also be a key mechanism underlying

beneficial remodeling of plaques to a more stable phenotype, especially in settings where there is

underlying increased inflammasome activation as occurs in $Tet2$ and $Jak2VF$ CH. Similar to macrophage

proliferation being a major factor accounting for increased macrophage number during advanced

atherosclerosis progression (50), suppression of proliferation may be an important mechanism decreasing

macrophage burden with LDL lowering (36). Our studies suggest that reversal of the proliferative

tendency of $Jak2^{VF}$ macrophages is also a key mechanism decreasing macrophage burden with LDL

lowering in $Jak2^{VF}$ mice. The mechanism of proliferation could involve uptake of modified LDL by

scavenger receptors into the endo-lysosomal system (44) consistent with our observations. The decreased

proliferation of $Jak2VF$ macrophages signifies that there are fewer inflammatory macrophages in lesions,

consistent with the virtual disappearance of $Jak2^{VF}$ macrophages in lesions with aggressive cholesterol

lowering.

The parallel decrease in macrophage proliferation and DNA damage response suggested that decreased

replication induced DNA damage could be a mechanism to explain decreased AIM2 inflammasome

activation and pyroptosis. However, mechanistic studies in $Jak2^{VF}$ BMDMs indicated that macrophage

proliferation could be clearly dissociated from DNA damage responses and AIM2 inflammasome

activation that were rather dependent on mitochondrial ROS generation and DNA damage as suggested by

earlier studies (46,51). The increased DNA damage and inflammasome activation in $Jak2^{VF}$ macrophages

compared to controls likely reflects increased glycolysis, mitochondrial ROS formation and oxidative

mitochondrial damage (30,52). However, the decreased $Jak2^{VF}$ macrophage burden, as well as decreased

AIM2 expression and decreased cholesterol-induced AIM2 inflammasome activation together explain

why effective LDL lowering was sufficient to reverse the adverse effects of $Jak2^{VF}$ on atherosclerosis.

Paradoxically, despite the overall reduction in macrophage burden, during resolution there may also be a

relative increase in a proliferative set of efferocytotic macrophages that have increased c-myc expression

and contribute to plaque stability (13,14). In addition to a potential role in cell proliferation, c-myc

increases Akt signaling and cell survival (53,54). Previously, we observed a 4-fold increase in TREM2^{Hi}

macrophages induced by both LDL lowering and $Jak2^{VF}$ inactivation in $Jak2^{VF}$ CH mice (16). In the

present study, we found that LDL lowering alone led to a marked increase in TREM2 staining in

macrophages, as well as elevated c-myc expression specifically within TREM2^{Hi} macrophages, in both

control and mutant mice. This contrasts with the decreased proliferation and DNA damage response found

specifically in the $Jak2^{VF}$ macrophages. Our findings together suggest that there is an increase in

TREM2^{Hi}, c-myc and MerTK expressing macrophages induced by LDL lowering either through increased

proliferation or increased survival of these subsets that then have beneficial effects on efferocytosis and

plaque stability. Given that inflammasome activation and IL-1 release have been linked to reduced

MerTK and TREM2 as a result of a disintegrin and metalloproteinase 17 (ADAM17)-mediated cleavage

(39,55), the increase in this pro-resolving macrophage population may in part be secondary to reduced

cholesterol-induced inflammasome activation.

LDL lowering guidelines have recommended progressively lower targets based on clinical trial evidence.

Current recommendations suggest an LDL cholesterol target below 70 mg/dl for patients with established

ACVD or in certain high-risk groups (56). Statin use has been associated with improved outcomes in

human MPN patients and was associated with a 37% reduction in the risk of thrombosis in patients who

had polycythemia vera or essential thrombocythemia as well as a 22% reduction in all-cause mortality

(57). While data from mice should be extrapolated to humans with caution, the AIM2 inflammasome

seems to increase ACVD risk in both mice and humans carrying the $Jak2^{VF}$ mutation (31). The present

preclinical study adds mechanistic findings to support the idea that aggressive LDL-C lowering in $Jak2^{VF}$

MPN patients would reduce ACVD risk.

Acknowledgements

The authors thank Wenli Liu for assistance and support during the course of this study. This study used

the Confocal and Specialized Microscopy Shared Resource of the Herbert Irving Comprehensive Cancer

Center at Columbia University, funded in part through NIH/NCI Cancer Center Support Grant

P30CA013696. Additionally, this study used the resources of the Herbert Irving Comprehensive Cancer

Center Flow Cytometry Shared Resources funded in part through Center Grant P30CA013696.

Disclosures

Alan R. Tall is a consultant for CSL Behring and is on the scientific advisory board of Beren

Therapeutics. The remaining authors declare no competing financial interests.

Data Availability

The datasets generated for this study are available from the corresponding author upon reasonable

request.

References

1. Martin SS, Aday AW, Almarzooq ZI, Anderson CAM, Arora P, Avery CL, Baker-Smith CM,

Barone Gibbs B, Beaton AZ, Boehme AK, Commodore-Mensah Y, Currie ME, Elkind MSV,

Evenson KR, Generoso G, Heard DG, Hiremath S, Johansen MC, Kalani R, Kazi DS, Ko D, Liu

J, Magnani JW, Michos ED, Mussolino ME, Navaneethan SD, Parikh NI, Perman SM, Poudel R,

Rezk-Hanna M, Roth GA, Shah NS, St-Onge MP, Thacker EL, Tsao CW, Urbut SM, Van Spall

HGC, Voeks JH, Wang NY, Wong ND, Wong SS, Yaffe K, Palaniappan LP; American Heart

Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics

Subcommittee. 2024 Heart Disease and Stroke Statistics: A Report of US and Global Data From

the American Heart Association. *Circulation* 2024;**149**:e347–e913.

2. Vallejo☐Vaz AJ, Fayyad R, Boekholdt SM, Hovingh GK, Kastelein JJP, Melamed S, Barter P,

Waters DD, Ray KK. Triglyceride-rich lipoprotein cholesterol and risk of cardiovascular events

among patients receiving statin therapy in the TNT trial. *Circulation* 2018;**138**:770–781.

3. Cannon CP, Blazing MA, Giugliano RP, McCagg A, White JA, Theroux P, Darius H, Lewis BS,

Ophuis TO, Jukema JW, De Ferrari GM, Ruzyllo W, De Lucca P, Im K, Bohula EA, Reist C,

Wiviott SD, Tershakovec AM, Musliner TA, Braunwald E, Califf RM; IMPROVE-IT

Investigators. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med*

2015;**372**:2387–2397.

4. Sabatine MS, Giugliano RP, Keech AC, Honarpour N, Wiviott SD, Murphy SA, Kuder JF, Wang

H, Liu T, Wasserman SM, Sever PS, Pedersen TR; FOURIER Steering Committee and

Investigators. Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl*

J Med 2017;**376**:1713–1722.

5. Nidorf SM, Fiolet ATL, Mosterd A, Eikelboom JW, Schut A, Opstal TSJ, The SHK, Xu XF,

Ireland MA, Lenderink T, Latchem D, Hoogslag P, Jerzewski A, Nierop P, Whelan A, Hendriks R,

Swart H, Schaap J, Kuijper AFM, van Hessen MWJ, Saklani P, Tan I, Thompson AG, Morton A,

Judkins C, Bax WA, Dirksen M, Alings M, Hankey GJ, Budgeon CA, Tijssen JGP, Cornel JH,

Thompson PL; LoDoCo2 Trial Investigators. Colchicine in patients with chronic coronary

disease. *N Engl J Med* 2020;**383**:1838–1847.

6. U.S. Food and Drug Administration. Drug approval package: Lodoco (colchicine) tablets, 0.5 mg.

Approval letter. Published June 16, 2023. Accessed July 5, 2025.

https://www.accessdata.fda.gov/drugsatfda_docs/nda/2024/215727Orig1s000Approv.pdf

7. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne CM, Fonseca F,

Nicolau J, Koenig W, Anker SD, Kastelein JJP, Cornel JH, Pais P, Pella D, Genest J, Cifkova R,

Lorenzatti A, Forster T, Kobalava Z, Vida-Simiti L, Flather M, Shimokawa H, Ogawa H,

Dellborg M, Rossi PRF, Troquay RPT, Libby P, Glynn RJ; CANTOS Trial Group.

Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med*

2017;**377**:1119–1131.

8. Ridker PM. From C-reactive protein to interleukin-6 to interleukin-1: moving upstream to

identify novel targets for atheroprotection. *Circ Res* 2016;**118**:145–156.

9. Yalcinkaya M, Liu W, Xiao T, Abramowicz S, Wang R, Wang N, Westerterp M, Tall AR.

Cholesterol trafficking to the ER leads to the activation of CaMKII/JNK/NLRP3 and promotes

atherosclerosis. *J Lipid Res* 2024;**65**:100534.

10. Westerterp M, Fotakis P, Ouimet M, Bochem AE, Zhang H, Molusky MM, Wang W, Abramowicz

S, la Bastide-van Gemert S, Wang N, Welch CL, Reilly MP, Stroes ES, Moore KJ, Tall AR.

Cholesterol efflux pathways suppress inflammasome activation, NETosis, and atherogenesis.

Circulation 2018;**138**:898–912.

11. Yalcinkaya M, Fotakis P, Liu W, Endo-Umeda K, Dou H, Abramowicz S, Xiao T, Libby P, Wang

N, Tall AR, Westerterp M. Cholesterol accumulation in macrophages drives NETosis in

atherosclerotic plaques via IL-1β secretion. *Cardiovasc Res* 2023;**119**:969–981.

12. Sheedy FJ, Grebe A, Rayner KJ, Kalantari P, Ramkhelawon B, Carpenter SB, Becker CE,

Ediriweera HN, Mullick AE, Golenbock DT, Stuart LM, Latz E, Fitzgerald KA, Moore KJ. CD36

coordinates NLRP3 inflammasome activation by facilitating intracellular nucleation of soluble

ligands into particulate ligands in sterile inflammation. *Nat Immunol* 2013;**14**:812–820.

13. Gerlach BD, Ampomah PB, Yurdagul A Jr, Liu C, Lauring MC, Wang X, Kasikara C, Kong N,

Shi J, Tao W, Tabas I. Efferocytosis induces macrophage proliferation to help resolve tissue

injury. *Cell Metab* 2021;**33**:2445–2463.e8.

14. Yurdagul A Jr, Subramanian M, Wang X, Crown SB, Ilkayeva OR, Darville L, Kolluru GK,

Rymond CC, Gerlach BD, Zheng Z, Kuriakose G, Kevil CG, Koomen JM, Cleveland JL, Muoio

DM, Tabas I. Macrophage metabolism of apoptotic cell-derived arginine promotes continual

efferocytosis and resolution of injury. *Cell Metab* 2020;**31**:518–533.e10.

15. Karnewar S, Karnewar V, Deaton RA, Shankman LS, Benavente ED, Williams CM, Bradley X,

Alencar GF, Bulut GB, Kirmani S, Baylis RA, Zunder ER, den Ruijter HM, Pasterkamp G,

Owens GK. IL-1β inhibition partially negates the beneficial effects of diet-induced

atherosclerosis regression in mice. *Arterioscler Thromb Vasc Biol* 2024;**44**:1379–1392.

16. Fidler TP, Dunbar A, Kim E, Hardaway B, Pauli J, Xue C, Abramowicz S, Xiao T, O'Connor K,

Sachs N, Wang N, Maegdefessel L, Levine R, Reilly MP, Tall AR. Suppression of IL-1β promotes

beneficial accumulation of fibroblast-like cells in atherosclerotic plaques in clonal hematopoiesis.

Nat Cardiovasc Res 2024;**3**:60–75.

17. Sharma M, Schlegel MP, Afonso MS, Brown EJ, Rahman K, Weinstock A, Sansbury BE, Corr

EM, van Solingen C, Koelwyn GJ, Shanley LC, Beckett L, Peled D, Lafaille JJ, Spite M, Loke P,

Fisher EA, Moore KJ. Regulatory T cells license macrophage pro-resolving functions during

atherosclerosis regression. *Circ Res* 2020;**127**:335–353.

18. Zhang X, Rotllan N, Canfrán-Duque A, Sun J, Toczek J, Moshnikova A, Malik S, Price NL,

Araldi E, Zhong W, Sadeghi MM, Andreev OA, Bahal R, Reshetnyak YK, Suárez Y, Fernández-

Hernando C. Targeted suppression of miRNA-33 using pHLIP improves atherosclerosis

regression. *Circ Res* 2022;**131**:77–90.

19. Basu D, Hu Y, Huggins LA, Mullick AE, Graham MJ, Wietecha T, Barnhart S, Mogul A, Pfeiffer

K, Zirlik A, Fisher EA, Bornfeldt KE, Willecke F, Goldberg IJ. Novel reversible model of

atherosclerosis and regression using oligonucleotide regulation of the LDL receptor. *Circ Res*

2018;**122**:560–567.

20. Tall AR, Fuster JJ. Clonal hematopoiesis in cardiovascular disease and therapeutic implications.

Nat Cardiovasc Res 2022;**1**:116–124.

21. Jaiswal S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG, Lindsley RC, Mermel

CH, Burtt N, Chavez A, Higgins JM, Moltchanov V, Kuo FC, Kluk MJ, Henderson B, Kinnunen

L, Koistinen HA, Ladenvall C, Getz G, Correa A, Banahan BF, Gabriel S, Kathiresan S,

Stringham HM, McCarthy MI, Boehnke M, Tuomilehto J, Haiman CA, Groop L, Atzmon G,

Wilson JG, Neuberg D, Altshuler D, Ebert BL. Age-related clonal hematopoiesis associated with

adverse outcomes. *N Engl J Med* 2014;**371**:2488–2498.

22. Jaiswal S, Natarajan P, Silver AJ, Gibson CJ, Bick AG, Shvartz E, McConkey M, Gupta N,

Gabriel S, Ardissino D, Baber U, Mehran R, Fuster V, Danesh J, Frossard P, Saleheen D,

Melander O, Sukhova GK, Neuberg D, Libby P, Kathiresan S, Ebert BL. Clonal hematopoiesis

and risk of atherosclerotic cardiovascular disease. *N Engl J Med* 2017;**377**:111–121.

23. Kar SP, Quiros PM, Gu M, Jiang T, Mitchell J, Langdon R, Iyer V, Barcena C, Vijayabaskar MS,

Fabre MA, Carter P, Petrovski S, Burgess S, Vassiliou GS. Genome-wide analyses of 200 453

individuals yield new insights into the causes and consequences of clonal hematopoiesis. *Nat*

Genet 2022;**54**:1155–1166.

24. Bick AG, Weinstock JS, Nandakumar SK, Fulco CP, Bao EL, Zekavat SM, Szeto MD, Liao X,

Leventhal MJ, Nasser J, Chang K, Laurie C, Burugula BB, Gibson CJ, Lin AE, Taub MA, Aguet

F, Ardlie K, Mitchell BD, Barnes KC, Moscati A, Fornage M, Redline S, Psaty BM, Silverman

EK, Weiss ST, Palmer ND, Vasan RS, Burchard EG, Kardia SLR, He J, Kaplan RC, Smith NL,

Arnett DK, Schwartz DA, Correa A, de Andrade M, Guo X, Konkle BA, Custer B, Peralta JM,

Gui H, Meyers DA, McGarvey ST, Chen IY, Shoemaker MB, Peyser PA, Broome JG, Gogarten

SM, Wang FF, Wong Q, Montasser ME, Daya M, Kenny EE, North KE, Launer LJ, Cade BE, Bis

JC, Cho MH, Lasky-Su J, Bowden DW, Cupples LA, Mak ACY, Becker LC, Smith JA, Kelly TN,

Aslibekyan S, Heckbert SR, Tiwari HK, Yang IV, Heit JA, Lubitz SA, Johnsen JM, Curran JE,

Wenzel SE, Weeks DE, Rao DC, Darbar D, Moon JY, Tracy RP, Buth EJ, Rafaels N, Loos RJF,

Durda P, Liu Y, Hou L, Lee J, Kachroo P, Freedman BI, Levy D, Bielak LF, Hixson JE, Floyd JS,

Whitsel EA, Ellinor PT, Irvin MR, Fingerlin TE, Raffield LM, Armasu SM, Wheeler MM, Sabino

EC, Blangero J, Williams LK, Levy BD, Sheu WH, Roden DM, Boerwinkle E, Manson JE,

Mathias RA, Desai P, Taylor KD, Johnson AD; NHLBI Trans-Omics for Precision Medicine

Consortium; Auer PL, Kooperberg C, Laurie CC, Blackwell TW, Smith AV, Zhao H, Lange E,

Lange L, Rich SS, Rotter JI, Wilson JG, Scheet P, Kitzman JO, Lander ES, Engreitz JM, Ebert

BL, Reiner AP, Jaiswal S, Abecasis G, Sankaran VG, Kathiresan S, Natarajan P. Inherited causes

of clonal haematopoiesis in 97,691 whole genomes. *Nature* 2020;**586**:763–768.

25. Cordua S, Kjaer L, Skov V, Pallisgaard N, Hasselbalch HC, Ellervik C. Prevalence and

phenotypes of JAK2 V617F and calreticulin mutations in a Danish general population. *Blood*

2019;**134**:469–479.

26. Wang S, Luo X, Hu S, Zhao C, Sun Q, Zeng M, Bao X, Wang Y, Wu F, Yang Y, Lv Y, Bai X, Hao

W, Liu M, Yi B, Chen Y, Meng W, Li J, Li M, Huang J, Wu T, Zhao Y, Zhang Z, An J, Libby P,

Jia H, Yu B. Plaque erosion risk and JAK2 V617F variant. *Eur Heart J* 2025;**46**:2205–2219.

27. Molinaro R, Sellar RS, Vromman A, Sausen G, Folco E, Sukhova GK, McConke ME, Corbo C,

Ebert BL, Libby P. The clonal hematopoiesis mutation Jak2V617F aggravates endothelial injury

and thrombosis in arteries with erosion-like intimas. *Int J Cardiol* 2024;**409**:132184.

28. Liu W, Pircher J, Schuermans A, Ul Ain Q, Zhang Z, Honigberg MC, Yalcinkaya M, Nakao T,

Pournamadri A, Xiao T, Hajebrahimi MA, Wasner L, Stegner D, Petzold T, Natarajan P, Massberg

S, Tall AR, Schulz C, Wang N. Jak2 V617F clonal hematopoiesis promotes arterial thrombosis

via platelet activation and cross talk. *Blood* 2024;**143**:1539–1550.

29. Liu W, Hardaway BD, Kim E, Pauli J, Wettich JL, Yalcinkaya M, Hsu CC, Xiao T, Reilly MP,

Tabas I, Maegdefessel L, Schlepckow K, Haass C, Wang N, Tall AR. Inflammatory crosstalk

impairs phagocytic receptors and aggravates atherosclerosis in clonal hematopoiesis in mice. *J*

Clin Invest 2024;**135**:e182939.

30. Fidler TP, Xue C, Yalcinkaya M, Hardaway B, Abramowicz S, Xiao T, Liu W, Thomas DG,

Hajebrahimi MA, Pircher J, Silvestre-Roig C, Kotini AG, Luchsinger LL, Wei Y, Westerterp M,

Snoeck HW, Papapetrou EP, Schulz C, Massberg S, Soehnlein O, Ebert BL, Levine RL, Reilly

MP, Libby P, Wang N, Tall AR. The AIM2 inflammasome exacerbates atherosclerosis in clonal

haematopoiesis. *Nature* 2021;**592**:296–301.

31. Yu Z, Fidler TP, Ruan Y, Vlasschaert C, Nakao T, Uddin MM, Mack T, Niroula A, Heimlich JB,

Zekavat SM, Gibson CJ, Griffin GK, Wang Y, Peloso GM, Heard-Costa N, Levy D, Vasan RS,

Aguet F, Ardlie KG, Taylor KD, Rich SS, Rotter JI, Libby P, Jaiswal S, Ebert BL, Bick AG, Tall

AR, Natarajan P. Genetic modification of inflammation- and clonal hematopoiesis-associated

cardiovascular risk. *J Clin Invest* 2023;**133**:e168597.

32. Mullally A, Lane SW, Ball B, Megerdichian C, Okabe R, Al-Shahrour F, Paktinat M, Haydu JE,

Housman E, Lord AM, Wernig G, Kharas MG, Mercher T, Kutok JL, Gilliland DG, Ebert BL.

Physiological Jak2V617F expression causes a lethal myeloproliferative neoplasm with

differential effects on hematopoietic stem and progenitor cells. *Cancer Cell* 2010;**17**:584–596.

33. Pan H, Xue C, Auerbach BJ, Fan J, Bashore AC, Cui J, Yang DY, Trignano SB, Liu W, Shi J,

Ihuegbu CO, Bush EC, Worley J, Vlahos L, Laise P, Solomon RA, Connolly ES, Califano A,

Sims PA, Zhang H, Li M, Reilly MP. Single-cell genomics reveals a novel cell state during

smooth muscle cell phenotypic switching and potential therapeutic targets for atherosclerosis in

mouse and human. *Circulation* 2020;**142**:2060–2075.

34. Yalcinkaya M, Liu W, Thomas LA, Olszewska M, Xiao T, Abramowicz S, Papapetrou EP,

Westerterp M, Wang N, Tabas I, Tall AR. BRCC3-mediated NLRP3 deubiquitylation promotes

inflammasome activation and atherosclerosis in Tet2 clonal hematopoiesis. *Circulation*

2023;**148**:1764–1777.

35. Kar SP, Quiros PM, Gu M, Jiang T, Mitchell J, Langdon R, Iyer V, Barcena C, Vijayabaskar MS,

Fabre MA, Carter P, Petrovski S, Burgess S, Vassiliou GS. Genome-wide analyses of 200 453

individuals yield new insights into the causes and consequences of clonal hematopoiesis. *Nat*

Genet 2022;**54**:1155–1166.

36. Härdtner C, Kornemann J, Krebs K, Ehlert CA, Jander A, Zou J, Starz C, Rauterberg S,

Sharipova D, Dufner B, Hoppe N, Dederichs TS, Willecke F, Stachon P, Heidt T, Wolf D, von Zur

Mühlen C, Madl J, Kohl P, Kaeser R, Boettler T, Pieterman EJ, Princen HMG, Ho-Tin-Noé B,

Swirski FK, Robbins CS, Bode C, Zirlik A, Hilgendorf I. Inhibition of macrophage proliferation

dominates plaque regression in response to cholesterol lowering. *Basic Res Cardiol* 2020;**115**:78.

37. Peled M, Nishi H, Weinstock A, Barrett TJ, Zhou F, Quezada A, Fisher EA. A wild-type mouse-

based model for the regression of inflammation in atherosclerosis. *PLoS One* 2017;**12**:e0173975.

38. Broz P. Pyroptosis: molecular mechanisms and roles in disease. *Cell Res* 2025;**35**:334–344.

39. Ngai D, Schilperoort M, Tabas I. Efferocytosis-induced lactate enables the proliferation of pro-

resolving macrophages to mediate tissue repair. *Nat Metab* 2023;**5**:2206–2219.

40. Sukka SR, Ampomah PB, Darville LNF, Ngai D, Wang X, Kuriakose G, Xiao Y, Shi J, Koomen

JM, McCusker RH, Tabas I. Efferocytosis drives a tryptophan metabolism pathway in

macrophages to promote tissue resolution. *Nat Metab* 2024;**6**:1736–1755.

41. Patterson MT, Xu Y, Hillman H, Osinski V, Schrank PR, Kennedy AE, Barrow F, Zhu A, Tollison

S, Shekhar S, Stromnes IM, Tassi I, Wu D, Revelo XS, Binstadt BA, Williams JW. Trem2 agonist

reprograms foamy macrophages to promote atherosclerotic plaque stability — brief report.

Arterioscler Thromb Vasc Biol 2024;**44**:1646–1657.

42. Corsini A, Mazzotti M, Villa A, Maggi FM, Bernini F, Romano L, Romano C, Fumagalli R,

Catapano AL. Ability of the LDL receptor from several animal species to recognize the human

apo B binding domain: studies with LDL from familial defective apo B-100. *Atherosclerosis*
1992;**93**:95–103.

43. Innerarity TL, Pitas RE, Mahley RW. Disparities in the interaction of rat and human lipoproteins
with cultured rat fibroblasts and smooth muscle cells: requirements for homology for receptor
binding activity. *J Biol Chem* 1980;**255**:11163–11172.

44. Dubland JA, Francis GA. Lysosomal acid lipase: at the crossroads of normal and atherogenic
cholesterol metabolism. *Front Cell Dev Biol* 2015;**3**:3.

45. de Dios C, Bartolessis I, Roca-Agujetas V, Barbero-Camps E, Mari M, Morales A, Colell A.
Oxidative inactivation of amyloid beta-degrading proteases by cholesterol-enhanced
mitochondrial stress. *Redox Biol* 2019;**26**:101283.

46. Dang EV, McDonald JG, Russell DW, Cyster JG. Oxysterol restraint of cholesterol synthesis
prevents AIM2 inflammasome activation. *Cell* 2017;**171**:1057–1071.e11.

47. Lin JD, Nishi H, Poles J, Niu X, McCauley C, Rahman K, Brown EJ, Yeung ST, Vozhilla N,
Weinstock A, Ramsey SA, Fisher EA, Loke P. Single-cell analysis of fate-mapped macrophages
reveals heterogeneity, including stem-like properties, during atherosclerosis progression and
regression. *JCI Insight* 2019;**4**:e124574.

48. Feig JE, Pineda-Torra I, Sanson M, Bradley MN, Vengrenyuk Y, Bogunovic D, Gautier EL,
Rubinstein D, Hong C, Liu J, Wu C, van Rooijen N, Bhardwaj N, Garabedian M, Tontonoz P,
Fisher EA. LXR promotes the maximal egress of monocyte-derived cells from mouse aortic
plaques during atherosclerosis regression. *J Clin Invest* 2010;**120**:4415–4424.

49. Nagareddy PR, Murphy AJ, Stirzaker RA, Hu Y, Yu S, Miller RG, Ramkhelawon B, Distel E,
Westerterp M, Huang LS, Schmidt AM, Orchard TJ, Fisher EA, Tall AR, Goldberg IJ.
Hyperglycaemia promotes myelopoiesis and impairs the resolution of atherosclerosis. *Cell Metab*
2013;**17**:695–708.

50. Robbins CS, Hilgendorf I, Weber GF, Theurl I, Iwamoto Y, Figueiredo JL, Gorbatov R, Sukhova
GK, Gerhardt LM, Smyth D, Zavitz CC, Shikatani EA, Parsons M, van Rooijen N, Lin HY,

Husain M, Libby P, Nahrendorf M, Weissleder R, Swirski FK. Local proliferation dominates lesional macrophage accumulation in atherosclerosis. *Nat Med* 2013;**19**:1166–1172.

51. Bae JH, Jo SI, Kim SJ, Lee JM, Jeong JH, Kang JS, Cho NJ, Kim SS, Lee EY, Moon JS. Circulating cell-free mtDNA contributes to AIM2 inflammasome-mediated chronic inflammation in patients with type 2 diabetes. *Cells* 2019;**8**:328.

52. Rao TN, Hansen N, Hilfiker J, Rai S, Majewska JM, Leković D, Gezer D, Andina N, Galli S, Cassel T, Geier F, Delezie J, Nienhold R, Hao-Shen H, Beisel C, Di Palma S, Dimeloe S, Trebicka J, Wolf D, Gassmann M, Fan TW, Lane AN, Handschin C, Dirnhofer S, Kröger N, Hess C, Radimerski T, Koschmieder S, Čokić VP, Skoda RC. JAK2-mutant hematopoietic cells display metabolic alterations that can be targeted to treat myeloproliferative neoplasms. *Blood* 2019;**134**:1832–1846.

53. Miller DM, Thomas SD, Islam A, Muench D, Sedoris K. c-Myc and cancer metabolism. *Clin Cancer Res* 2012;**18**:5546–5553.

54. Wise DR, DeBerardinis RJ, Mancuso A, Sayed N, Zhang XY, Pfeiffer HK, Nissim I, Daikhin E, Yudkoff M, McMahon SB, Thompson CB. Myc regulates a transcriptional program that stimulates mitochondrial glutaminolysis and leads to glutamine addiction. *Proc Natl Acad Sci U S A* 2008;**105**:18782–18787.

55. Endo-Umeda K, Kim E, Thomas DG, Liu W, Dou H, Yalcinkaya M, Abramowicz S, Xiao T, Antonson P, Gustafsson JÅ, Makishima M, Reilly MP, Wang N, Tall AR. Myeloid LXR (liver X receptor) deficiency induces inflammatory gene expression in foamy macrophages and accelerates atherosclerosis. *Arterioscler Thromb Vasc Biol* 2022;**42**:719–731.

56. Virani SS, Newby LK, Arnold SV, Bittner V, Brewer LC, Demeter SH, Dixon DL, Fearon WF, Hess B, Johnson HM, Kazi DS, Kolte D, Kumbhani DJ, LoFaso J, Mahtta D, Mark DB, Minissian M, Navar AM, Patel AR, Piano MR, Rodriguez F, Talbot AW, Taqueti VR, Thomas RJ, van Diepen S, Wiggins B, Williams MS; Peer Review Committee Members. 2023 AHA/ACC/ACCP/ASPC/NLA/PCNA guideline for the management of patients with chronic

coronary disease: a report of the American Heart Association/American College of Cardiology

Joint Committee on Clinical Practice Guidelines. *Circulation* 2023;**148**:e9–e119.

57. Podoltsev NA, Wang R, Shallis RM, Stempel JM, Di M, Neparidze N, Zeidan AM, Huntington

SF, Giri S, Hull SC, Gore SD, Ma X. Statin use, survival and incidence of thrombosis among

older patients with polycythemia vera and essential thrombocythemia. *Cancer Med*

2023;**12**:18889–18900.

Figure 1

Figure 1. Impaired Atherosclerosis Regression in *Jak2^{VF}* MPN Mice with Moderate Cholesterol

Lowering. A. Study Design, created with BioRender.com. **B.** Plasma cholesterol (n = 34, 55, 55, 54, 16,

16 for Ctrl mice, n=31, 55, 49, 39, 17, 16 for *Jak2^{VF}* mice, for weeks 0, 5, 11, 15, 17.5, 21 respectively). *P*

= 0.028, <0.0001, <0.0001, <0.0001, 0.0005, <0.0001 (Ctrl vs *Jak2^{VF}* at weeks 0, 5, 11, 15, 17.5, 21

respectively). **C.** H&E images of aortic root lesions. Black lines, necrotic core. Scale bar, 200 µm. **D.**

Lesion area, n = 15-23. *P* < 0.0001 (Ctrl Baseline vs *Jak2^{VF}* Baseline; Ctrl LDL Lowering vs. *Jak2^{VF}* LDL

Lowering), *P* = 0.052 (*Jak2^{VF}* Baseline vs LDL Lowering). **E.** Necrotic core area, n = 15-23. *P* = 0.0079

(Ctrl Baseline vs *Jak2^{VF}* Baseline), *P* = 0.0002 (Ctrl LDL Lowering vs. *Jak2^{VF}* LDL Lowering), *P* =

0.0003 (*Jak2^{VF}* Baseline vs LDL Lowering). **F.** Picrosirius red-stained aortic root lesions. Scale bar, 278

µm **G.** Collagen area as a percentage of lesion area, n = 15-24. *P* = 0.02 (Ctrl Baseline vs LDL

Lowering). **H.** Images of aortic root lesions stained for MAC2 (Green) and DAPI (Blue). Scale bar, 100

µm. **I.** Macrophage area, n = 15-24. *P* = 0.006 (Ctrl Baseline vs LDL Lowering). All quantifications

shown as mean ± s.e.m. Two-way ANOVA with the Geisser-Greenhouse correction for sphericity and

Tukey's multiple comparisons test (B). Two-way ANOVA with Tukey's multiple comparisons test

(D,E,G,I).

Figure 2

Figure 2. Moderate Cholesterol Lowering Reverses Macrophage AIM2 Inflammasome Activation,

DNA Damage, and Proliferation in *Jak2^{VF}* MPN Lesions. A. Images of aortic root lesions stained for

MAC2 (Green), Cleaved GasD (Red), and DAPI (Blue). Scale bar, 127 µm. **B.** $\log_{10}$ transformed cleaved

GasD mean fluorescence intensity (MFI) in lesions with the addition of constant 1, n = 15-24. $P = 0.021$

(Ctrl Baseline vs *Jak2^{VF}* Baseline). **C.** $\log_{10}$ transformed cleaved GasD mean fluorescence intensity

(MFI) in necrotic cores with the addition of the constant 1, n = 15-24. $P = 0.0008$ (Ctrl Baseline vs *Jak2^{VF}*

Baseline), $P = 0.032$ (Ctrl LDL Lowering vs *Jak2^{VF}* LDL Lowering). **D.** Images of aortic root lesions

stained for MAC2 (Green), Cleaved GasD (Red), and DAPI (Blue). Scale bar, 40 µm. White arrows,

cleaved GasD+ macrophages. **E.** $\log_{10}$ transformed cleaved GasD positive macrophages per section with

the addition of constant 1, n = 15-24. $P = 0.03$ (*Jak2^{VF}* Baseline vs LDL Lowering). **F.** Images of aortic

root lesions stained for MAC2 (Green), AIM2 (Red), and DAPI (Blue). Scale bar, 75 µm. White arrows,

AIM2 positive macrophages. **G.** $\log_{10}$ transformed AIM2 positive macrophages per section with the

addition of constant 1, n = 15-23. $P = 0.0001$ (Ctrl Baseline vs *Jak2^{VF}* Baseline), $P = 0.03$ (*Jak2^{VF}*

Baseline vs LDL Lowering). **H.** Images of aortic root lesions stained for MAC2 (Green), pγH2AX (Red),

and DAPI (Blue). Scale bar, 46 µm. White arrows, pγH2AX positive cells. **I.** $\log_{10}$ transformed pγH2AX

positive macrophages per section with the addition of constant 1, n = 15-23. $P = 0.0019$ (Ctrl Baseline vs

Jak2^{VF} Baseline), $P = 0.045$ (*Jak2^{VF}* Baseline vs LDL Lowering). **J.** Images of aortic root lesions stained

for MAC2 (Green), Ki67 (Red), and DAPI (Blue). Scale bar, 25 µm. White arrows, Ki67 positive

macrophages. **K.** $\log_{10}$ transformed Ki67 positive macrophages per section with the addition of constant

1, n = 15-23. $P = 0.0019$ (*Jak2^{VF}* Baseline vs LDL Lowering). **L.** Images of aortic root lesions stained for

MAC2 (Green), 8OHdG (Red), and DAPI (Blue). Scale bar, 78 µm. **M.** Percentage of MAC2 positive

area double positive for 8OHdG and MAC2, n = 14-24. $P = 0.023$ (Ctrl Baseline vs LDL Lowering).

MAC2 intensity differences reflect independent staining and imaging sessions across panels. All

quantifications shown as mean ± s.e.m. Two-way ANOVA with Tukey's multiple comparisons test (B,C).

Kruskal-Wallis test with Dunn's multiple comparisons test (E,G,I,K,M).

Figure 3

Figure 3. Moderate Cholesterol Lowering Reverses Impaired Efferocytosis in *Jak2^{VF}* MPN Lesions

While Increasing TREM2^{Hi} Macrophages in Control and *Jak2^{VF}* MPN Lesions A. Images of aortic

root lesions stained for MAC2 (Green), MerTK (Red), and DAPI (Blue). Scale bar, 102 µm. **B.**

Percentage of MAC2 positive area double positive for MerTK and MAC2, n = 15-23. $P = 0.02$ (Ctrl

Baseline vs *Jak2^{VF}* Baseline), $P < 0.0001$ (*Jak2^{VF}* Baseline vs LDL Lowering). **C.** Images of aortic root

lesions stained for MAC2 (Green), TREM2 (Red), and DAPI (Blue). Scale bar, 100 µm. **D.** Percentage of

MAC2 positive area double positive for TREM2 and MAC2, n = 15-24. $P < 0.0001$ (Ctrl and *Jak2^{VF}*

Baseline vs LDL Lowering). **E.** Images of in situ efferocytosis in aortic root lesions: MAC2 (Green),

TUNEL (Red), and DAPI (Blue). Scale bar, 40 µm. White arrows, nuclei double positive for TUNEL and

MAC2. White wedges, nuclei positive for TUNEL but negative for MAC2. **F.** Percentage of TUNEL

positive nuclei also positive for MAC2. $P = 0.011$ (Ctrl Baseline vs *Jak2^{VF}* Baseline), $P = 0.021$ (*Jak2^{VF}*

Baseline vs LDL Lowering). MAC2 intensity differences reflect independent staining and imaging

sessions across panels, n = 13-24. All quantifications shown as mean ± s.e.m. Two-way ANOVA with

Tukey's multiple comparisons test (D). Kruskal-Wallis test with Dunn's multiple comparisons test (B,F).

Figure 4

Figure 4. Aggressive Cholesterol Lowering Normalizes Regression in *Jak2^{VF}* MPN Lesions. A. Study

Design, created with BioRender.com. **B.** Plasma cholesterol (n = 5, 38, 19, 18, 18 for Ctrl mice, n = 5, 37,

15, 16, 15 for *Jak2^{VF}* mice, for weeks 0, 4, 13, 15, 18 respectively). *P* = 0.0041 for genotype effect by

two-way ANOVA with Geisser-Greenhouse correction, *P* = 0.011, 0.048, 0.056, 0.0008 (Ctrl vs *Jak2^{VF}* at

weeks 0, 4, 13, 15 respectively). **C.** H&E images of aortic root lesions. Black lines, necrotic core. Scale

bar, 200 µm. **D.** Lesion area, n = 13-20. *P* < 0.0001 (Ctrl Baseline vs *Jak2^{VF}* Baseline; Ctrl LDL Lowering

vs. *Jak2^{VF}* LDL Lowering). **E.** Necrotic core area, n = 13-20. *P* = 0.0003 (Ctrl Baseline vs *Jak2^{VF}*

Baseline), *P* = 0.0006 (Ctrl LDL Lowering vs. *Jak2^{VF}* LDL Lowering), *P* = 0.90 (*Jak2^{VF}* Baseline vs LDL

Lowering). **F.** Picrosirius red-stained aortic root lesions. Scale bar, 219 µm **G.** Collagen area as a

percentage of lesion area, n = 13-20. *P* < 0.0001 (Ctrl Baseline vs LDL Lowering), *P* = 0.0002 (*Jak2^{VF}*

Baseline vs LDL Lowering), *P* = 0.02 (Ctrl LDL Lowering vs. *Jak2^{VF}* LDL Lowering). **H.** Images of

aortic root lesions stained for MAC2 (Green) and DAPI (Blue). Scale bar, 193 µm. **I.** Macrophage area, n

= 13-20. *P* < 0.0001 (Ctrl and *Jak2^{VF}* Baseline vs LDL Lowering). All quantifications shown as mean ±

s.e.m. Two-way ANOVA with Tukey's multiple comparisons test (D,G,I). Two-way ANOVA with the

Geisser-Greenhouse correction for sphericity and Tukey's multiple comparisons test (B). Kruskal-Wallis

test with Dunn's multiple comparisons test (E).

Figure 5

Figure 5. Cholesterol Lowering Suppresses *Jak2*^{VF} Macrophage Proliferation and DNA Damage. A.

Study design, created with BioRender.com. **B.** Plasma cholesterol (n = 80, 79, 79 for Baseline mice for

weeks 3, 11, 15 respectively; n = 13, 15 for Ctrl Progression mice, n = 16, 16 for Ctrl LDL Lowering

mice, n = 15, 15 for *Jak2*^{VF} Progression mice, n = 16, 16 for *Jak2*^{VF} LDL Lowering mice, for weeks 18,

23 respectively) **C.** Images of aortic root lesions for MAC2 (Green), Cleaved GasD (Red), and DAPI

(Blue). Scale bar, 98 µm. **D.** Log$_{10}$ transformed cleaved GasD mean fluorescence intensity (MFI) in

lesions with the addition of constant 1, n = 14-16. *P* = 0.014 (Baseline vs *Jak2*^{VF} Progression), *P* = 0.06

(*Jak2*^{VF} Progression vs *Jak2*^{VF} LDL Lowering). **E.** Images of aortic root lesions for MAC2 (Red), Ki67

(White), and ZsGreen (Green). Scale bar, 133 µm. White arrows, macrophages double positive for Ki67

and ZsGreen. **F.** Macrophages positive for both Ki67 and ZsGreen per section, n = 14-15. *P* = 0.0028

(Baseline vs *Jak2*^{VF} Progression), *P* = 0.0051 (Ctrl Progression vs *Jak2*^{VF} Progression), *P* = 0.0005

(*Jak2*^{VF} Progression vs *Jak2*^{VF} LDL Lowering). **G.** Macrophages positive for Ki67 but negative for

ZsGreen per section, n = 13-16. **H.** Images of aortic root lesions stained for pγH2AX (Red), ZsGreen

(Green), and DAPI (Blue). Scale bar, 75 µm. White arrows, pγH2AX positive cells. **I.** Log$_{10}$ transformed

cells double positive for pγH2AX and ZsGreen in lesions with the addition of constant 1, n = 13-16. *P* =

0.0024 (Baseline vs *Jak2*^{VF} Progression), *P* = 0.0019 (*Jak2*^{VF} Progression vs *Jak2*^{VF} LDL Lowering). **J.**

Log$_{10}$ transformed cells positive for pγH2AX but negative for ZsGreen in lesions with the addition of

constant 1, n = 13-16. All quantifications shown as mean ± s.e.m. One-way ANOVA with Holm-Sidak's

multiple comparisons test (D). Kruskal-Wallis test with Dunn's multiple comparison's test (F,G,I,J).

Figure 6

Figure 6. Cholesterol Lowering Increases c-myc in TREM2^Hi Macrophages. A. Images of aortic root

lesions stained for MAC2 (Green), c-myc (Red), and DAPI (Blue). Scale bar, 121 µm. **B.** Percentage of

macrophages positive for c-myc, n = 14-16. $P < 0.0001$ (Baseline vs Ctrl Progression; Ctrl Progression vs

LDL Lowering; *Jak2^VF* Progression vs LDL Lowering), $P = 0.028$ (Baseline vs Ctrl LDL Lowering), $P =$

0.0009 (Baseline vs *Jak2^VF* Progression). **C.** Images of aortic root lesions stained for MAC2 (Green), c-

myc (Blue), and TREM2 (Red). Scale bar, 121 µm. **D.** Percentage of TREM2 positive macrophages

positive for c-myc, n = 14-16. $P = 0.012$ (Baseline vs Ctrl Progression), $P < 0.0001$ (Ctrl Progression vs

LDL Lowering), $P = 0.0037$ (*Jak2^VF* Progression vs LDL Lowering). All quantifications shown as mean $\pm$

s.e.m. One-way ANOVA with Holm-Sidak's multiple comparisons test (B). Kruskal-Wallis test with

Dunn's multiple comparisons test (D).

Figure 7

 Figure 7. Cholesterol Increases AIM2 Inflammasome Activation and Nuclear dsDNA Breaks via

 Increased Mitochondrial ROS in *Jak2^{VF}* Macrophages. A. EdU+ BMDMs as a percentage of total

 BMDMs after overnight incubation with acLDL (25 µg/ml). n = 4 biological replicates. $P = 0.032$ for

 acLDL effect. **B.** Representative immunoblot analysis of pγH2AX in BMDMs after overnight incubation

 with acLDL (25 µg/ml). **C.** Densitometric quantification of pγH2AX normalized to β-actin from B. Each

 data point represents a cell-well technical replicate from one mouse (1 biological replicate, 3 technical

 replicates). $P = 0.022$ (Ctrl Vehicle vs. *Jak2^{VF}* Vehicle), $P = 0.038$ (Ctrl acLDL vs. *Jak2^{VF}* acLDL).

 Statistical comparisons in this panel were performed across cell-well technical replicates from a single

 mouse (1 biological replicate; n = 3 wells per condition) to assess within-experiment reproducibility. **D.**

 EdU+ BMDMs as a percentage of total BMDMs after overnight incubation with oxPAPC (50 µg/ml).

 Each data point represents a cell-well technical replicate from one mouse (1 biological replicate, 3

 technical replicates). $P = 0.022$ (Ctrl Vehicle vs. Ctrl oxPAPC), $P < 0.0001$ (*Jak2^{VF}* Vehicle vs. *Jak2^{VF}*

 oxPAPC), $P = 0.0004$ (Ctrl oxPAPC vs. *Jak2^{VF}* oxPAPC). Statistical comparisons in this panel were

 performed across cell-well technical replicates from a single mouse (1 biological replicate; n = 3 wells per

 condition) to assess within-experiment reproducibility. **E.** Representative immunoblot analysis of

 pγH2AX in BMDMs after overnight incubation with oxPAPC (50 µg/ml). **F.** Densitometric quantification

 of pγH2AX normalized to β-Actin from E. Each data point represents a cell-well technical replicate from

 one mouse (1 biological replicate, 3 technical replicates). $P = 0.026$ for oxPAPC effect. Statistical

 comparisons in this panel were performed across cell-well technical replicates from a single mouse (1

 biological replicate; n = 3 wells per condition) to assess within-experiment reproducibility. **G.** ELISA

 quantification of total IL1β in cell culture media of BMDMs incubated with MβCD-cholesterol (30

 µg/ml) alone, mitoTEMPO (10 µM) alone, or MβCD-cholesterol (30 µg/ml) in combination with

 mitoTEMPO (10 µM) for 4 hours after priming with LPS (100 ng/ml) for 3 hours. n = 4 cell-well

 technical replicates from a single mouse (1 biological replicate). Wells failing prespecified quality-control

 criteria were excluded (see Statistical Analysis). Statistical comparisons in this panel were performed

 across cell-well technical replicates from a single mouse (1 biological replicate; n = 4 wells per condition)

to assess within-experiment reproducibility. **H.** ELISA quantification of total IL1β in cell culture media of

BMDMs incubated with MβCD-cholesterol (30 µg/ml) for 4 hours after priming with LPS (100 ng/ml)

for 3 hours. n = 6 cell-well technical replicates from a single mouse (1 biological replicate). No biological

replicates were available. P = 0.0017 (*Jak2^{VF}* Vehicle vs. *Jak2^{VF}* MβCD-cholesterol), P = 0.0026 (*Jak2^{VF}*

MβCD-cholesterol vs. *Jak2^{VF}Aim2$^{-/-}$* MβCD-cholesterol). Statistical comparisons in this panel were

performed across cell-well technical replicates from a single mouse (1 biological replicate; n = 6 wells per

condition) to assess within-experiment reproducibility. **I.** Representative immunoblot analysis of pγH2AX

in BMDMs after a 4-hour incubation with MβCD-cholesterol (30 µg/ml). **J.** Densitometric quantification

of pγH2AX normalized to β-Actin from H. n = 8 biological replicates for the vehicle groups and n = 4

biological replicates for the MβCD-cholesterol groups. P = 0.0019 (Ctrl Vehicle vs. Ctrl MβCD-

cholesterol), P = 0.0028 (*Jak2^{VF}* Vehicle vs. *Jak2^{VF}* MβCD-cholesterol). P = 0.019 for genotype effect. P

< 0.0001 for MβCD-cholesterol effect. Ctrl Vehicle data include earlier assessments of this endpoint in

untreated cells, prior to the initiation of the treatment arm. All samples were processed and analyzed using

the same protocol. **K.** EdU+ BMDMs as a percentage of total BMDMs after a 4-hour incubation with

MβCD-cholesterol (30 µg/ml). n = 3 biological replicates. P = 0.043 for genotype effect. All

quantifications shown as mean ± s.e.m. Two-way ANOVA with Tukey's multiple comparisons test

(A,C,D,F,H,J,K). Kruskal-Wallis test with Dunn's multiple comparisons test (G).