1 **Two classes of amine/glutamate multi-transmitter neurons innervate Drosophila**
2 **internal male reproductive organs**
3

4 Martha Chaverra[1], John Paul Toney[1], Lizetta D. Dardenne-Ankringa[1], Jace Tolleson
5 Knee[1], Ann R. Morris[1], Joseph B. Wadhams[1], Sarah J. Certel[2], and R. Steven
6 Stowers[1,3]
7
8 [1]Montana State University, Department of Microbiology and Cell Biology, Bozeman, MT
9 59717
10
11 [2]The University of Montana, Division of Biological Sciences, Missoula, MT 59812
12
13 [3]corresponding author

Abstract

    The essential outcome of a successful mating is the transfer of genetic material from males to females in sexually reproducing animals from insects to mammals. In males, this culminates in ejaculation, a precisely timed sequence of organ contractions driven by the concerted activity of interneurons, sensory neurons, and motor neurons. Although central command circuits that trigger copulation have been mapped, the motor architecture and the chemical logic that couple specific neuronal subclasses to organ specific contractility, seminal fluid secretion, and sperm emission remain largely uncharted. This gap in knowledge limits our ability to explain how neural circuits adapt to varying contexts and how their failure contributes to infertility. Here we present an in-depth anatomical and functional analysis of the motor neurons that innervate the internal male reproductive tract of Drosophila melanogaster. We identify two classes of multi-transmitter motor neurons based on neurotransmitter usage, namely octopamine and glutamate neurons (OGNs) and serotonin and glutamate neurons (SGNs), each with a biased pattern of innervation: SGNs predominate in the accessory glands, OGNs in the ejaculatory duct, with equal contributions of each to the seminal vesicles. Both classes co-express vesicular transporters for glutamate (vGlut) and amines (vMAT), confirming their dual chemical identity. Their target organs differentially express receptors for glutamate, octopamine, and serotonin, suggesting combinatorial neuromodulation of contractility. Functional manipulations show that SGNs are essential for male fertility but OGNs are dispensable. Glutamatergic transmission from both classes is also dispensable for fertility. These findings provide the first high-resolution map linking multi-transmitter motor neurons to specific reproductive organs, reveal an unexpected division of labor between serotonergic and octopaminergic signaling pathways, and establish a framework for dissecting conserved neural principles that govern ejaculation and male fertility.

Introduction

    Reproduction is a biological imperative of all species. For most sexually reproducing animal species, this involves locating a mate, courtship, and most essentially, the transfer of genetic material via copulation. The endpoint of copulation, ejaculation, consists of two coordinated phases, emission and expulsion (ALWAAL et al. 2015; SONI et al. 2022). During the emission phase, sperm and seminal fluid are relocated from their site of origin to a central canal for transfer to the female during the subsequent expulsion phase. The series of events that culminate in ejaculation are regulated by a spinal ejaculation generator located in the lumbar L3-L4 region of the spinal cord of mammals that executes a motor program coordinating a series of muscle contractions and relaxations of the internal and external male genitalia (TRUITT AND COOLEN 2002; CARRO-JUAREZ et al. 2003; CHEHENSSE et al. 2017).

    While male external genitalia exhibit remarkable evolutionary variation (EBERHARD 1985), internal male reproductive organs exhibit remarkable evolutionary conservation over the last 600 million years since the last common ancestor of insects and mammals (GUERRERO et al. 2004; GOMES et al. 2012; BEGUELINI et al. 2013; TOMAS et al. 2019; FROMME et al. 2021; OLIVEIRA et al. 2021; YARTSEV AND EVSEEVA 2021; MACIEL et al. 2024). These evolutionarily conserved structures of the male reproductive system

include paired testes, seminal vesicles (SVs), and semen producing organs (prostate
glands in humans, accessory glands (AGs) in insects), as well as a singular duct
(urethra in humans, ejaculatory duct (ED) in insects) through which sperm and seminal
fluid flow during mating.
Drosophila melanogaster has been instrumental in revealing fundamental principles
of male reproductive behavior and physiology that extend beyond insects, including how
neural circuits control mating drive, courtship, and the mechanics and duration of
copulation (ACEBES et al. 2004; BILLETER et al. 2006a; VILLELLA AND HALL 2008; TAYLER
et al. 2012; CRICKMORE AND VOSSHALL 2013; PAVLOU et al. 2016; JOIS et al. 2018; BAKER
et al. 2024). Among the findings of these studies is the existence of neural circuitry in
the ventral nerve cord that regulates ejaculation similar to the spinal ejaculation
generator circuitry of mammals, suggesting an ancient evolutionary origin. How the
motor neurons directly innervating the internal male reproductive organs control the
events leading to ejaculation is less well understood. It is also known in Drosophila and
other species, including humans, that male ejaculate quality is plastic and can be
adapted to environmental conditions, especially the level of male-male competition
(GAGE AND BAKER 1991; WEDELL AND COOK 1999; PIZZARI et al. 2003; POUND AND GAGE
2004; BRETMAN et al. 2009; GARBACZEWSKA et al. 2013; HOPKINS et al. 2019; DELECCE et
al. 2025), but the neural mechanisms underlying these adaptations remain to be
identified.
Previous studies have determined there are at least three types of motor neurons
innervating the Drosophila internal male reproductive system distinguished by
neurotransmitter usage including serotonergic (LEE AND HALL 2001), octopaminergic
(PAULS et al. 2018), and glutamatergic (PAVLOU et al. 2016) motor neurons, although
there may be at least some overlap in neurotransmitter usage as serotonin/glutamate
multi-transmitter motor neurons have been reported in the seminal vesicle
(CASTELLANOS et al. 2013). How these neurons coordinate the motor program to elicit
ejaculation is an open question, as is their anatomical organization relative to each
other. Clarifying the targets and anatomy of these motor neuron classes will uncover
critical principles for how neuronal coordination of internal male reproductive organ
function contributes to male fertility. To address these questions, we performed a
comprehensive analysis of the neurons innervating each internal organ, defined their
neurotransmitter complements and receptor profiles, and tested their necessity for
successful sperm transfer and fecundity.
94
95 **Results**
96
Drosophila male reproductive system anatomy
The Drosophila male reproductive system consists of paired testes, SVs, and AGs,
as well as a singular ED and ejaculatory bulb (Figure 1) (DEMEREC 1950). The testes
are surrounded by smooth muscle with no neuronal innervation, while the SVs, AGs,
and ED are ensheathed by neuronally innervated skeletal muscles (LEE AND HALL 2001;
BILLETER *et al.* 2006b; SUSIC-JUNG *et al.* 2012). The ejaculatory bulb also possesses
neuronally innervated skeletal muscle for pumping sperm out of the ED to the female.
The SVs fuse just prior to termination in the ED where they form a narrow sphincter that
opens during mating to allow sperm to flow. The AGs, insect equivalent of mammalian

prostate glands, produce the bulk of seminal fluid (WOLFNER 1997; MAJANE et al. 2022)
and remain separate before terminating in the ED (BAIRATI 1968). Underlying the
muscles of the internal male reproductive system organs are a layer of secretory
epithelial cells (BAIRATI 1968) that are the cellular source of the seminal fluid
components (SEPIL et al. 2019; WIGBY et al. 2020).
111
Neurotransmitter identity of neurons innervating internal male reproductive
organs
To confirm previous reports of innervation of the Drosophila male reproductive
system by octopaminergic, serotonergic, and glutamatergic neurons, GAL4 drivers for
the octopamine (OA) neurotransmitter synthesis enzyme tyrosine decarboxylase 2
(Tdc2), the serotonin (5-Hydroxytryptamine, 5-HT) neurotransmitter synthesis enzyme
tryptophan hydroxylase (TRH) and the vesicular glutamate transporter (vGlut) were
used to drive expression of a *UAS-CD8-mCherry* plasma membrane reporter. As
expected, all three drivers exhibited expression in neurons of the male reproductive
system, but with distinguishable differences in expression patterns. *Tdc2-GAL4*
exhibited dense innervation of the SV and ED with sparse innervation of the AG (Figure
2, A). *TRH-GAL4* exhibited dense innervation of the SV and AGs, with modest
innervation of the ED (Figure 2, B), and *vGlut-GAL4* exhibited dense innervation of all
three organs (Figure 2, C).
126
OA/glutamate neurons are distinct from 5-HT/glutamate neurons
To determine if the observed neuronal expression of the GAL4 drivers for Tdc2, TRH,
and vGlut are coincident or exclusive, two separate approaches were implemented. In
the first approach, split-GAL4 drivers specific for each neurotransmitter were tested in
pairwise combinations. Both *Tdc2-AD* paired with *vGlut-GAL4-DBD* (Figure 2, D) and
vGlut-AD paired with *Tdc2-GAL4-DBD* (Figure 2, E) exhibited expression patterns very
similar to *Tdc2-GAL4* with dense innervation of the SV and ED, and sparse innervation
of the AGs. These results suggest that all the Tdc2 neurons innervating the male
reproductive system are OA/glutamate dual neurotransmitter neurons (OGNs).
Similarly, both *TRH-AD* paired with *vGlut-GAL4-DBD* (Figure 2, F) and *vGlut-AD* paired
with *TRH-GAL4-DBD* (Figure 2, G) exhibited expression patterns very similar to *TRH-*
GAL4 with dense innervation of the SVs and AGs with modest innervation of the ED.
These results suggest that all the serotonergic neurons innervating the male
reproductive system are 5-HT/glutamate dual neurotransmitter neurons (SGNs). To
determine if there is overlap between the OGNs and SGNs, *Tdc2-AD* was paired with
TRH-GAL4-DBD with the result that no neuronal expression was observed (Figure 2,
H). This result suggests the OGNs and SGNs are exclusive and non-overlapping.
A second approach to assess overlap between octopaminergic, serotonergic, and
glutamatergic neurons of the male reproductive system utilized pairwise combinations of
GAL4 and LexA drivers specific for each neurotransmitter. Pairing *TRH-GAL4* and
Tdc2-LexA together with *UAS-6XGFP* and *LexAop-6XmCherry* reporters revealed no
overlap in either the SVs (Figure 3, A-F), ED (Figure 3, G-L), or AGs (Figure 3, M-R).
These results are consistent with the split-GAL4 result above of no intersection between
octopaminergic and serotonergic neurons of the male reproductive system, although
both neuron types are often tightly fasciculated, especially in the SVs. Pairing *vGlut-*

LexA with *TRH-GAL4* revealed a high degree of overlap in the SVs (Figures 3S1, A-F),
partial overlap in the ED (Figures 3S1, G-L), and near complete overlap in the AGs
(Figures 3S1, M-R). Pairing *vGlut-LexA* with *Tdc2-GAL4* revealed significant overlap in
the SVs (Figures 3S2, A-F), majority overlap in the ED (Figures 3S2, G-L), and limited
overlap in the AGs (Figures 3S2, M-R). The results of the split-GAL4 intersectional
strategy and the GAL4/LexA overlap strategy are entirely consistent with each other.
Taken together, they demonstrate that the Drosophila internal male reproductive system
receives input from two distinct and nonoverlapping motor-neuron populations, one
population that co-expresses octopamine and glutamate (OGNs) and another distinct
population that co-expresses serotonin and glutamate (SGNs). The SV receives roughly
equal innervation from both classes, the AG is innervated more extensively by SGNs,
and the ED is innervated more extensively by OGNs.
164
OGNs are *fru +*/*dsx +* while SGNs are *fru+*/*dsx -*
Neurons that innervate the male reproductive tract are expected to be sexually
dimorphic (Billeter and Goodwin 2004; Yamamoto 2007; Siwicki and Kravitz 2009;
Verhulst and van de Zande 2015). To determine whether they express the canonical
sex-determination genes *fruitless* (*fru*) and/or *doublesex* (*dsx*), *fru-GAL4* and *vGlut-*
LexA were paired with *UAS-6XGFP* and *LexAop-6XmCherry*. This experiment revealed
complete overlap: every vGlut-positive neuron in the reproductive system—both OGNs
and SGNs—was also *fru*-positive (Fig. 3S3, A–C). To examine *dsx* expression, the
octopaminergic driver *Tdc2-AD* was combined with *dsx-GAL4-DBD*. The resulting
pattern (dense terminals in the SVs and ED, sparse arbors in the AGs; Fig. 3S3, D)
matched that of *Tdc2-GAL4* alone, indicating that all OGNs are *dsx*-positive. In contrast,
pairing the serotonergic *TRH-AD* with *dsx-GAL4-DBD* produced no signal in the
reproductive tract (Fig. 3S3, E), showing that SGNs are *dsx*-negative. Together, these
experiments establish that OGNs are *fru+*/*dsx+*, whereas SGNs are *fru+*/*dsx–*,
underscoring a molecular distinction between the two neuromodulatory inputs to the
male reproductive system.
181
Non-overlapping expression of TβH and Tdc2 with 5-HT confirms exclusivity of
OGNs and SGNs
Several predictions follow from the finding that non-overlapping sets of OGNs and
SGNs innervate the male reproductive system. Among these are that the
neurotransmitter synthesis enzymes for OA, Tyramine-β-hydroxylase (TβH), and
Tyrosine decarboxylase 2 (Tdc2) should be exclusively expressed in the OGNs along
with the sole Drosophila vesicular glutamate transporter (vGlut), while 5-HT should be
exclusively expressed in the SGNs along with vGlut-40XV5. To test these predictions,
vGlut-40XV5 expression was separately assessed in combination with TβH or Tdc2. 5-
HT expression, visualized with an anti-5-HT antibody, was also included in this analysis
to distinguish SGNs. As predicted, extensive vGlut-40XV5 expression was observed
throughout the male reproductive system except for the testes (Figures 4, A and G) with
a sharp boundary of neuronal innervation at the testicular duct (arrows, Figure 4, A).
Also as predicted, TβH-GFP and Tdc2 were observed in the male reproductive system
with expression patterns similar to that of *Tdc2-GAL4* (Figures 4, B and H, respectively).

Pairwise combinations of overlapping expression of vGlut-40XV5, 5-HT, and either TβH
(Figures 4, D-F) or Tdc2 (Figures 4, J-L) are also shown.
Higher resolution images of vGlut-40XV5, TβH-GFP, Tdc2, and 5-HT in the male
reproductive system allow an assessment of overlapping expression. In the SV, broad
robust expression was observed for vGlut-40XV5 (Figures 4S1, A, G, M, and S) and 5-
HT (Figures 4S1, C, I, O, and U), with somewhat more restricted expression for TβH-
GFP (Figure 4S1, B and H) and Tdc2 (Figure 4S1, N, T). Pairwise overlapping
expression is also shown for vGlut-40XV5 and 5-HT with TβH-GFP (Figures 4S1, D-F
and J-L) and Tdc2 (Figures 4S1, P-R and V-X). In the highest resolution images,
coincident expression is evident between vGlut-40XV5 with TβH-GFP (Figure 4S1, J)
and vGlut-40XV5 with Tdc2 (Figure 4WS1, V). Consistent with results above,
exclusivity of expression is apparent between TβH-GFP and 5-HT (Figure 4S1, L) and
between Tdc2 and 5-HT (Figure 4S1, X), although the tight fasciculation of OGNs and
SGNs is evident.
Similar results were obtained in higher resolution images of the ED and AGs
although the extent of innervation of the different amines varies between these tissues
with more OGNs than SGNs in the ED and the reciprocal in the AGs. In higher
resolution images of the ED, strong extensive expression was observed for vGlut-
40XV5 (Figures 4S2, A, G, M, and S) with moderately less expression for TβH-GFP
(Figure 4S2, B and H) and Tdc2 (Figure 4S2, N, T), with significantly less expression of
5-HT (Figures 4S2, C, I, O, and U. Pairwise overlapping expression is also shown for
vGlut-40XV5 and 5-HT with TβH-GFP (Figures 4S2, D-F and J-L) and Tdc2 (Figures
4S2, P-R and V-X). In the highest resolution images, coincident expression is evident
between vGlut-40XV5 with TβH-GFP (Figure 4S2, J) and vGlut-40XV5 with Tdc2
(Figure 4S2, V). As in the SV, exclusivity of expression is apparent between TβH-GFP
and 5-HT (Figure 4S2, L) and between Tdc2 and 5-HT (Figure 4S2, X), also with tight
fasciculation of the OGNs and SGNs.
In higher resolution images of the AGs, robust widespread expression was observed
for vGlut-40XV5 (Figures 4S3, A, G, M, and S) and 5-HT (Figures 4S3, C, I, O, and U,
with no expression of TβH-GFP (Figure 4S3, B and H), and sparse expression of Tdc2
(Figure 4S3, N, T). Pairwise overlapping expression is also shown for vGlut-40XV5 and
5-HT with TβH-GFP (Figures 4S3, D-F and J-L) and Tdc2 (Figures 4S3, P-R and V-X).
In the highest resolution images, coincident expression is evident between vGlut-40XV5
and Tdc2 (Figure 4S3, V). As in the SV and ED, exclusivity of expression is apparent
between Tdc2 and 5-HT (Figure 4S3, X).
The expression of vGlut-40XV5, TβH-GFP, and 5-HT was also assessed at the
junction of the SVs and AGs with the ED. vGlut-40XV5 innervation of both the SV and
AG is particularly dense (Figure 4S4, A). Significant expression of TβH-GFP and 5-HT
was also observed (Figure 4S4, B and C, respectively). Pairwise overlapping
expression for vGlut-40XV5, TβH-GFP, and 5-HT is also shown (Figure 4S4, D-F).
These experiments showing no overlap in expression of TβH-GFP and Tdc2 with 5-HT
provide additional support to the exclusivity of OGNs and SGNs. Collectively, these
findings confirm that octopaminergic vGlut-positive and serotonergic vGlut-positive
motor neurons form distinct, nonoverlapping populations that differentially innervate the

Drosophila male reproductive tract, underscoring strict transmitter exclusivity within this
circuit.
243

Expression of vGlut and vMAT in both OGNs and SGNs further confirms multi-
neurotransmitter usage
Co-labeling of vGlut-40XV5 with TβH / Tdc2 in OGNs and with 5-HT in SGNs implies
both neuron types are multi-transmitter and predicts they will express not only vGlut but
also the sole Drosophila monoamine vesicular transporter vMAT known to package both
OA and 5-HT (DESHPANDE *et al.* 2020). Since OGNs and SGNs often fasciculate tightly
together, co-conditional expression of vGlut and vMAT was implemented to visualize
expression separately in each population. This was achieved using either the *TRH-*
GAL4 or *Tdc2-GAL4* drivers in combination with the *B2RT-STOP-B2RT-vGlut-40XMYC*,
RSRT-STOP-RSRT-6XV5-vMAT, *UAS-B2* and *UAS-R* recombinase transgenes. Using
the *TRH-GAL4* driver to restrict conditional expression to SGNs, as predicted, both
vGlut-40XMYC (Figures 5, A, E) and 6XV5-vMAT (Figures 5, B, F) were detected in the
SV. Tdc2 expression was also included in this experiment (Figures 5, C and G) as was
a *UAS-CD8-mCherry* reporter to outline SGNs (Figures 5, D and H). A high degree of
overlap was observed between vGlut-40XMYC and 6XV5-vMAT (Figure5, I) with no
overlap between Tdc2 and the other markers (Figures 5, J-L).
In a reciprocal experiment in the SV using the *Tdc2-GAL4* driver and 5-HT
immunostaining, both vGlut-40XMYC (Figure 5, M and Q) and 6XV5-vMAT (Figure 5, N
and R) were detected with a mostly overlapping distribution (Figure 5, U). 5-HT
immunostaining was in close proximity but did not overlap with any of the other markers
(Figure 5, V-X). It is also notable that the morphology of the SGNs is distinct from the
OGNs with the SGNs being larger and more extensively branched compared to the
OGNs (SGNs-Figures 5, D and H; OGNs-Figures 5, P and T), suggesting SGNs may
assert more regulatory control of SV muscle activity than OGNs.
Similar results were obtained in these co-conditional experiments for the ED and
AGs. With the *TRH-GAL4* experiment, vGlut-40XMYC (Figure 5S1, A and E) and 6XV5-
vMAT (Figure 5S1, B and F) were both present with a highly overlapping distribution
(Figure 5S1, I). Tdc2 did not overlap with any of the other markers (Figure 5S1, J-L).
With the *Tdc2-GAL4* experiment, vGlut-40XMYC (Figure 5S1, M and Q) and 6XV5-
vMAT (Figure 5S1, N and R) were both observed in a highly overlapping distribution
(Figures 5S1, U). 5-HT immunostaining did not overlap with any of the other markers
(Figures 5S1, V-X). Note that morphological differences between the SGNs and OGNs
in the ED are not as easily distinguished as in the SV.
With the *TRH-GAL4* experiment in the AGs, both vGlut-40XMYC (Figures 5S2, A and
E) and 6XV5-vMAT (Figures 5S2, B and F) are present in a largely overlapping
distribution (Figure 5S2, I). As with the SV and ED, Tdc2 expression did not overlap
with any of the other markers (Figures 5S2, J-L). With the *Tdc2-GAL4* experiment in
the AG, both vGlut-40XMYC (Figures 5S2, M and Q) and 6XV5-vMAT (Figures 5S2, N
and R) are present in an overlapping configuration (Figure 5S2, U). The distribution of
5-HT is distinct from any of the other markers (Figure 5S2, V-X). In the AGs the
morphology of the SGNs (Figures 5S2, D and H) is distinct from that of the OGNs
(Figures 5S2, P and T).

Since OGNs and SGNs are intermingled in tightly fasciculated neuron bundles, we
used intersectional labeling to visualize the glutamate transporter vGlut-40XV5 and the
monoamine transporter 6XV5-vMAT in each class separately. Together, the results
show that both vesicular transporters are expressed in OGNs and SGNs in an
overlapping distribution while the OGNs and SGNs remain anatomically distinct
throughout the male reproductive tract.

Expansion microscopy reveals prominent co-occurrence of vGlut and vMAT on
ED synaptic vesicles of both OGNs and SGNs
Although there is apparent overlap between vGlut-40XMYC and 6XV5-vMAT in both
SGNs and OGNs it is not possible to resolve whether they reside on the same or
different synaptic vesicles using standard microscopy methods. This constraint is due
to the ~250nm diffraction limit of light at visible wavelengths (ABBE 1873). To overcome
this limitation, expansion microscopy was performed on EDs using co-conditional
expression of vGlut-40XMYC and 6XV5-vMAT separately in SGNs and OGNs as above.
The protocol utilized in these experiments reported 10X expansion in a single round
(DAMSTRA et al. 2023) and this agreed well with our empirically determined expansion
factor of 10.47X based on the size comparison of nuclei area in muscles of the
unexpanded and expanded EDs (Figure 5S5, A and B, respectively). In a
representative example of expanded SGNs, 6XV5-vMAT (Figure 5S3, A) and vGlut-
40XMYC (Figure 5S3, B) revealed a significantly overlapping distribution (Figure 5S3,
C). To quantify the co-occurrence of vGlut-40XMYC and 6XV5-vMAT the BIOP JACop
ImageJ plug-in was used to calculate a Mander's co-efficient, a metric that calculates
the fraction of one signal that co-occurs with another. The output of the BIOP JACop
plug-in revealed substantial co-occurrence of 6XV5-vMAT and vGlut-40XMYC (Figures
5S3, D-I) including a graphical representation (Figure 5S3, J). For the SGNs, the
numerical Mander's coefficients for 6XV5-vMAT co-occurrence with vGlut-40XMYC
were 67.9% and vGlut-40XMYC co-occurrence with 6XV5-vMAT were 57.2% (Figure
5S3, K).
Expanded OGNs yielded similar results. In a representative example, 6XV5-vMAT
(Figure 5S4, A) and vGlut-40XMYC (Figure 5S4, B) exhibited substantial overlap
(Figure 5S4, C). The output of the BIOP JACop plug-in also revealed majority co-
occurrence in image (Figure 5S4, D-I) and graphical form (Figure 5S4, J). For the
OGNs, the numerical values of the Mander's coefficients were 67.4% for 6XV5-vMAT
co-occurrence with vGlut-40XMYC and 75.6% for vGlut-40XMYC co-occurrence with
6XV5vMAT (Figure 5S4, K).
The results of the expansion microscopy experiments indicate that in both SGNs and
OGNs the majority of ED synaptic vesicles are positive for both vGlut-40XMYC and
6XV5-vMAT and thereby they co-release glutamate and either OA (OGNs) or 5-HT
(SGNs). This demonstration of a neurotransmitter co-release mechanism may be a
conserved strategy for synchronizing rapid gland and muscle activity during copulation
that has general implications across species for how co-transmitter neurons mediate
complex motor behaviors.
328
OGNs and SGNs express synaptic markers common to other neuromuscular
junctions

In addition to providing information on whether a given synapse is primarily
excitatory or modulatory, identifying vesicular transporters and synaptic markers allows
mapping of active release sites to determine how pre- and post-synaptic specializations
align. Such information is essential for (i) elucidating how each neurotransmitter
influences muscle contractility and secretory function, and (ii) predicting how synaptic
efficacy might change during prolonged copulation or in response to neuromodulatory
feedback. To further characterize the synapses of the neurons innervating the male
reproductive system immunostaining was performed with vGlut-40XMYC, 6XV5-vMAT,
and either of three well-established Drosophila synaptic markers including Synapsin
(Syn)-synaptic vesicles, Bruchpilot (Brp)-active zones, and Discs-large (Dlg-Drosophila
homolog of mammalian PSD-95)-post-synaptic density. In the SV, strong Syn staining
was observed (Figures 6, C and F) that exhibited extensive co-localization with vGlut-
40XMYC (Figure 6, H) and 6XV5-vMAT (Figure 6, I). Similar results were obtained in
the ED and AGs. Abundant Syn expression was observed (Figure 6, C' and F') that
exhibited near precise overlap with vGlut-40XMYC (Figure 6, H') and 6XV5-vMAT
(Figure 6H, I'). For the AG, robust Syn expression was observed (Figures 6A, C" and
F") that co-localized strongly with vGlut-40XMYC (Figure 6A, H") and 6XV5-vMAT
(Figure 6A, I"). It is notable that a significantly higher density of Syn expression was
observed in the SV (Figure 6A, F) compared with the AGs (Figure 6A, F') and ED
(Figure 6A. F"), presumably indicating greater neurotransmitter output.
The active zone marker Brp exhibited strong expression in the SV (Figure 6S1, C
and G) that overlapped almost entirely with 6XV5-vMAT. Brp expression in the ED and
AG was noticeably sparser as compared to the SV. The sparse Brp expression in the
ED (Figure 6S1, K and O) overlapped with 6XV5-vMAT. Similarly, in the AGs, Brp
expression was sparse and overlapped with 6XV5-vMAT (Figure 6S1, S and W).
The post-synaptic density marker Dlg was expressed broadly throughout the male
reproductive system (Figure 6S2, C). In the SV, the bulk of Dlg expression is
predominantly in the epithelial cells with lower levels of expression in the muscles
(Figure 6S2, G and K) and no preferential post-synaptic accumulation (Figures 6S2, H
and L). Similar results were observed in the ED and AGs. In the ED at the level of the
muscle Dlg expression is diffuse (Figure 6S2, O) and does not accumulate post-
synaptically opposite synapses in the muscles (Figure 6S2, P). In the AGs, Dlg
expression is also diffuse in the muscles (Figure 6S2, T) and does not accumulate post-
synaptically opposite synapses in the muscle (Figure 6C, U). Images of Dlg expression
in the epithelial layer of the ED (Figure 6C, Q) and AG (Figure 6C, V) reveal apparent
membrane localization in a honeycomb pattern. The lack of post-synaptic accumulation
of Dlg opposite the synapses of SGNs and OGNs in the male reproductive system is
reminiscent of larval type II OGNs and distinct from larval type I glutamatergic neurons
that exhibit prominent post-synaptic concentration of Dlg (PARNAS et al. 2001). It is also
notable that there is no vGlut-40XMYC that does not co-localize with 6XV5-vMAT,
indicating there are no neurons innervating the SVs, AGs, or ED that uses glutamate as
their sole small molecule neurotransmitter. The absence of synapses with post-synaptic
Dlg accumulation, a characteristic of type I glutamate-only larval neurons, also supports
this assertion. Defining the molecular architecture of these synapses is critical for
understanding how they package, release, and receive neurotransmitters to shape the
strength and specificity of communication between motor neurons and target organs.

LDCVs are present in OGNs and SGNs

In addition to synaptic transmission via synaptic vesicles, neurons are also known to communicate using neuropeptides released via Large Dense Core Vesicles (LDCVs). To determine whether neurons innervating the male reproductive system contain LDCVs, immunostaining was performed with the LDCV marker IA2-GFP (Yu et al. 2025) in combination with vGlut-40XMYC and 6XV5-vMAT. Abundant IA2-GFP expression was observed throughout the male reproductive system (Figure 6S3, C). In the SVs, IA2-GFP (Figure 6S3, F and L) localizes to the same pre-synaptic regions of the neurons as vGlut-40XMYC (Figures 6S3, D and J) and 6XV5-vMAT (Figures 6S3, E and K). In the highest resolution images, there appears to be some overlap between IA2-GFP and vGlut-40XMYC (Figure 6S3, N) as well as between IA2-GFP and 6XV5-vMAT (Figure 6S3, O), but a substantial fraction of IA2-GFP distributes in close proximity to, but is distinct from, vGlut-40XMYC and 6XV5-vMAT. This distribution pattern is not entirely surprising as it is consistent with the prior observation that vesicular neurotransmitter transporters have been detected on LDCVs (Yu et al. 2025).

Similar results were observed in the ED and AGs. In the ED, prominent IA2-GFP expression was evident (Figures 6S4, C and I) that partially co-localized with vGlut-40XMYC (Figures 6S4, E and K) and 6XV5-vMAT (Figures 6S4, F and L). Considerable IA2-GFP expression was also apparent in the AGs (Figures 6S4, O and U) that partially overlapped with vGlut-40XMYC (Figures 6S4, Q and W) and 6XV5-vMAT (Figures 6S4, R and X). This finding of pronounced expression of IA2-GFP throughout the pre-synaptic terminals of neurons innervating the male reproductive system suggest substantial LDCVs are present in both SGNs and OGNs, and that these neurons are thus communicating not just via small molecule neurotransmitters inside synaptic vesicles but also by neuropeptides contained within LDCVs.

No cholinergic or GABAergic transmission at OGN or SGN synapses

Although to our knowledge there have been no motor neurons reported in Drosophila that use acetylcholine or GABA as a neurotransmitter, to rule out these possibilities for the male reproductive system, Drosophila containing genome-edited 7XMYC-vAChT and 9XV5-vGAT were immunostained for their corresponding epitope tags. Not unexpectedly, no expression of 7XMYC-vAChT was observed in the SVs, ED, or AGs (Figures 6S5, C, G, and K, respectively). Similarly, no expression of 9XV5-vGAT was observed in the SVs, ED, or AGs (Figures 6S5, D, H, and L, respectively), indicating that neither acetylcholine nor GABA are used as neurotransmitters in the OGNs and SGNs of the Drosophila internal male reproductive system.

GAL4 drivers for OA, 5-HT, and glutamate receptors express in the muscles of male reproductive organs

Given that OGNs and SGNs innervate the male reproductive system, an obvious prediction that follows is that neurotransmitter receptors for glutamate, OA, and 5-HT should be expressed in the muscles of the male reproductive system. As a first step in testing this prediction, GAL4 drivers for each of these neurotransmitters were paired with the *UAS-CD8-mCherry* reporter and assessed for expression. As it is well established that the GluRIIA-E/kainate type glutamate receptors mediate synaptic

transmission at Drosophila NMJs (DiAntonio 2006; He and Dickman 2025), GAL4
drivers for all five were assessed. GAL4 drivers for GluRIIA, GluRIID, and GluRIIE
showed no expression in the male reproductive system, but they also exhibited no
expression in other adult or larval muscles where they are known to be expressed.
These results suggest that the GAL4 drivers for these three GluRII receptors are simply
non-functional and are therefore non-informative. However, GluRIIB-GAL4 and
GluRIIC-GAL4 exhibited broad expression in the musculature of the male reproductive
system (Figures 7, A and B, respectively). This suggests glutamate is used for signaling
throughout the male reproductive system, consistent with the widespread expression of
vGlut in pre-synaptic terminals.
OA receptor GAL4 drivers for OAMB, OAα2R, Oct-TyrR, OAβ1R, OAβ2R, and
OAβ3R were assessed for expression in the male reproductive system. OAMB-GAL4
exhibited expression in the muscles of the SVs and ED, as well as epithelial cells of the
AGs (Figure 7S1, A). OAα2R-GAL4 displayed strong expression in the muscles of the
SVs, weak expression in the muscles of the AGs, and no expression in the ED (Figure
7S1, B). Oct-TyrR-GAL4 showed strong expression in non-muscle cells of the distal
AG, muscle expression in the AGs, ED, and ejaculatory bulb, expression in the testes,
but no expression in the muscles of the SVs (Figure 7S1, C). OAβ2R-GAL4 exhibited
strong expression in the epithelial cells of the ED, muscle expression in the SVs, ED,
and ejaculatory bulb, expression in the testes, but no muscle expression in the AGs
(Figure 7S1, E). OAβ1R-GAL4 and OAβ3R-GAL4 exhibited broad expression in the
neurons innervating the male reproductive system (Figures 7S1, D and F, respectively)
but no muscle expression, suggesting they are not mediating OA signaling in the
muscles. OAβ3R-GAL4 also displayed expression in the testes.   These results suggest
OA receptors play a prominent role in the functioning of the male reproductive system
and differentially mediate OA signaling in specific parts of the male reproductive system
based on their restricted, non-uniform expression patterns.
5-HT receptor GAL4 drivers for 5-HT1A, 5-HT2A, 5-HT-1B, 5-HT2B, and 5-HT7 were
also assessed for expression in the male reproductive system. 5-HT1A-GAL4, 5-HT1B-
GAL4, and 5-HT2B-GAL4 exhibited broad expression in the neurons innervating the
male reproductive system, but no muscle expression (Figures 7S2, A, C, and D,
respectively), suggesting these receptors are not mediating 5-HT signaling in the
muscles. 5-HT2A exhibited expression specifically in the epithelial cells of the SVs
(Figure 7S2, B). 5-HT7 showed substantial expression in neurons, but most strikingly,
strong expression in the muscles of the distal SVs, including the sphincter region, with a
discrete boundary of expression (arrows, Figure 7S2, E).
459
GluRIIA is expressed in the SVs and AGs, but not the ED
While informative for determining cellular expression, GAL4 driven reporters for
genes of interest do not provide critical mechanistic information about the subcellular
distribution of the corresponding proteins. To gain further insight into the role of OA, 5-
HT, and glutamate receptors in the function of the male reproductive system the
expression patterns of several receptors were assessed using epitope-tagging. To
determine the subcellular distribution of the ionotropic GluRII/kainate-type glutamate
receptors known to mediate fast excitatory glutamatergic transmission at other
Drosophila NMJs (DiAntonio 2006; He and Dickman 2025), the expression of a

previously characterized genome-edited GluRIIA-GFP fly strain in which GFP has been
inserted near the carboxy-terminus of GluRIIA at its endogenous genomic location was
evaluated in the male reproductive system. In the SV, GluRIIA-GFP (Figure 8, B and E)
exhibited punctate expression that paralleled, but was complementary to, that of the
glutamatergic synaptic vesicle marker vGlut-40XV5 (Figure 8, A and D). This
distribution of GluRIIA-GFP is not unlike its distribution at the NMJ of third instar larva,
optimally localized post-synaptically to bind glutamate released from pre-synaptic
terminals (BECKERS *et al.* 2024). Surprisingly, no GluRIIA-GFP was observed in the ED
(Figures 8, H and K) despite abundant expression of vGlut-40XV5 (Figures 8, G and J).
Expression of GluRIIA-GFP in the AGs (Figures 8, N and Q) was similar to that in the
SVs in that it closely paralleled vGlut-40XV5 expression (Figure 8, M and P) but was
complementary to it. Finding GluRIIA-GFP expression in the SVs and AGs, but not the
ED, and the expression of GluRIIB in all three tissues based on the *GluRIIB-GAL4*
expression pattern, suggest the SVs and AGs use a mixture of glutamate receptors
containing one or the other of the mutually exclusive GluRIIA or GluRIIB subunits in
combination with the common GluRIIC, D, and E subunits, while all glutamate receptors
in the ED solely utilize GluRIIB. As GluRII receptors containing GluRIIA generate higher
currents than those that contain GluRIIB (DIANTONIO et al. 1999), the lower GluRIIB
currents are apparently sufficient for a functional ED.
488
OA receptors exhibit distinct expression patterns in the muscles, neurons, and
epithelial cells of the male reproductive system
OA receptors are modulatory G-protein coupled receptors (GPCRs) that mediate
slower-acting but longer-lasting effects than ionotropic receptors. To visualize
expression of the OA receptors OAMB, OAα2R, and OAβ2R, CRISPR/Cas9 genome
editing was used to create conditional alleles of each receptor fused at their carboxy
termini to a multimerized epitope tag for enhanced detection sensitivity. These genome
edits are at the endogenous genomic location of each gene and their expression is thus
under the control of their complete endogenous regulatory regions. For visualizing
expression in the male reproductive system, germline inversion or excision variants
were generated that converted conditional expression to constitutive expression.
OAMB-10XV5 exhibited strong expression in the epithelial cells of the ED and AGs
with limited expression in muscles (Figure 8S1, A). Co-immunostaining with the
synaptic vesicle marker Synapsin was conducted to visualize pre-synaptic terminals
(Figure 8S1, B). In the SVs, higher resolution images revealed low level expression of
OAMB-10XV5 in muscles as well as obvious expression in pre-synaptic neurons
(Figures 8S1, D and G) that largely overlapped with Syn (Figures 8S1, F and I). OAMB-
10XV5 was similarly distributed in the ED with low-level expression in muscle and
clearly discernible expression in pre-synaptic neurons (Figure 8S1, J) based on
substantial overlap with Syn (Figure 8S1, L).
Strong OAMB-10XV5 expression was also observed in the epithelial cells of the ED
in a honeycomb pattern (Figure 8S1, M and P) reminiscent of Dlg and presumably
associated with the plasma membrane. OAMB-10XV5 expression in the epithelial cells
of the AGs was similar to its expression in the epithelial cells of the ED with prominent
signal in a honeycomb pattern (Figure 8S2, A and D). OAMB-10XV5 also exhibited low-
level expression in the muscles of the AG and expression in the pre-synaptic neurons

(Figure 8S2, G) by virtue of significant overlap with Syn (Figure 8S2, I). Prominent
OAMB-10XV5 expression was also observed in the ejaculatory bulb. Low-level
expression of OAMB-10XV5 was detected in the muscles of the ejaculatory bulb as was
expression in the pre-synaptic neurons (Figure 8S2, J and M) as evidenced by mostly
overlapping expression with Syn (Figure 8S2, L and O). In the non-muscular portion of
the ejaculatory bulb, OAMB-10XV5 expression was readily discernable in the epithelial
cells in a honeycomb pattern (Figure 8S2, J and P).
OAα2R-20XV5 exhibited strong expression in the muscles of both the SVs and
more distal regions of the proximal ED near where it connects with the ejaculatory bulb,
as well as lower-level expression in the AGs (Figure 8S3, A). Higher resolution images
of OAα2R-20XV5 in the SV indicate expression is restricted to the muscle (Figure 8S3,
G) as there is no apparent overlap with Syn (Figure 8S3, I). A cross-section of the SV
muscle reveals OAα2R-20XV5 is localized to the membrane of the muscle (Figure 8S3,
J), as would be expected for a GPCR. In the AGs, OAα2R-20XV5 is expressed in the
muscles (Figure 8S3, M and P) as well as in the pre-synaptic neurons as evidenced by
co-localization with Syn (Figure 8S3, O and R). OAα2R-20XV5 expression ends
abruptly just after the SVs fuse into the sphincter that regulates the movement of sperm
into the ED (Figure 8S4, A and D). OAα2R is expressed in the pre-synaptic neurons
innervating the sphincter with little to no expression in the sphincter muscles (Figure
8S4, C and F). In the proximal region of the anterior ED prominent OAα2R-20XV5
expression is observed throughout the innervating neurons with little to no OAα2R-
20XV5 in the muscles (Figures 8S4, G, I, J, and L). In contrast, in the distal region of
the anterior ED, robust expression of OAα2R-20XV5 is detected in the muscles (Figures
8S4, M, P, and S) but not in the pre-synaptic neurons (Figures 8E, O, R, and U). This
complementary pattern of OAα2R-20XV5 in neurons but not muscles in the proximal ED
and muscles but not neurons in the distal ED suggests these different regions of the ED
are exhibiting a differential response to OA. As in the SV, OAα2R-20XV5 localizes to
the membrane of the muscles as evident in a cross-sectional image (Figure 8E, S).
Interestingly in this region, Syn expression is inside the muscles of the ED (Figure 8E,
T), potentially suggesting the muscles are being stimulated from the inside, or
alternatively, that the neurons are stimulating the underlying epithelial cells. As pre-
synaptic neuronal expression of adrenoreceptors in mammals has been shown to be
release-inhibiting (BROWN AND GILLESPIE 1957; STARKE 1972), including in the urinary
bladder (SOMOGYI AND DE GROAT 1990) and vas deferens (O'CONNOR et al. 1999;
SCHEIBNER et al. 2001), OAα2R expression in the pre-synaptic neurons of the internal
male reproductive system may also function via inhibitory auto-reception.
OAβ2R-40XV5 exhibits strong expression in the epithelial cells of the ED, the
muscles of the ejaculatory bulb, the muscles of the SVs near the point of fusion, the
testes, with lower-level muscle expression in the SVs, ED, and AGs (Figure 8S5, A).
Higher resolution images of the SV reveal obvious expression in the muscle (Figure
8S5, D and G) but not in the pre-synaptic neurons as there is no apparent overlap with
Syn (Figures 8S5, F and I). A cross-sectional image of the SV reveals OAβ2R-40XV5 is
localized to the plasma membrane of muscle cells (Figure 8S5, J). In higher resolution
images of the ED, moderate levels of OAβ2R-40XV5 are present in the muscles
(Figures 8S5, M and P) but not in the innervating pre-synaptic neurons as there is no

obvious overlap with Syn (Figures 8S5, O and R). At the junction where the SVs fuse
into a single duct, OAβ2R-40XV5 expression is strong prior to fusion but drops off
abruptly at the point of full fusion (arrow, Figure 8S6, A). The strongest levels of
OAβ2R-40XV5 expression throughout the entire male reproductive system is in what
appears to be the epithelial cells that line the ED (Figure 8S6, A). A cross-section of the
ED reveals the distribution of OAβ2R-40XV5 in epithelial cells is exclusively at the
apical surface (Figure 8S6, D). Also notable, there is another boundary of OAβ2R-
40XV5 expression in the SVs somewhat prior to fusion (solid arrow, Figure 8S6, G).
Higher resolution images of OAβ2R-40XV5 expression at the point of SV fusion shows
the sharp dip in expression at the point of full fusion (arrow, Figure 8S6, J). A cross-
sectional image also reveals the drop in expression and membrane localization of
OAβ2R-40XV5 in the muscle (Figure 8S6, M). Similar to the proximal anterior ED, the
more distal region of the anterior ED exhibits strong expression in the muscles, and
especially the underlying epithelial cells (Figures 8S7, A and D), In the AGs, OAβ2R-
40XV5 expression is manifest in the muscles (Figures 8S7, G and J) but does not
overlap with the pre-synaptic neurons as evidenced by the lack of overlap with Syn
(Figures 8S7, I and L). Prominent OAβ2R-40XV5 expression was also observed in the
ejaculatory bulb (Figure 8S7, M and P). As β2 adrenoreceptors relax bladder muscles
in mice (WUEST et al. 2009), OAβ2R may relax muscles in the male reproductive
system.
Octopamine receptor expression in the internal organs of the male reproductive tract
was predicted based on the dense OGN innervation of these tissues. Although OA
receptor presence in these muscles aligns with their role in modulating contractile
strength and rhythm, the organ-specific differences in receptor subtype abundance
could only be established empirically. Epithelial expression implies that octopamine
influences non-neuronal functions such as protein secretion or barrier permeability, but
the precise cellular responses remain to be defined. Mapping OAMB, OAα2R, and
OAβ2R to discrete muscle, neuronal, and epithelial compartments provides mechanistic
information about how octopamine fine-tunes contraction, secretion, and presynaptic
feedback across the male reproductive tract, offering an evolutionary framework for
adrenergic control of fertility and pinpointing cell-specific targets for modulating
reproductive output.
592
5-HT7 is widely expressed in the male reproductive system but most prominently
in the sphincter region
Since the *5-HT7-GAL4* reporter delineates a sharp posterior boundary in the SV,
endogenous 5-HT7 receptor expression was evaluated to pinpoint where 5-HT may
directly modulate reproductive tract motility. To assess the expression pattern of the 5-
HT7 receptor, a 20XV5 tag was added to the carboxy-terminus by CRISPR/Cas9
genome editing at its endogenous genomic location. 5-HT7-20XV5 localizes to the
muscles of the SV just before and after the junction, the epithelial cells of the AGs, and
the ejaculatory bulb (Figures 8S8, A). Higher resolution images of the SV reveal low
level expression in muscles (Figures 8S8, D and G), as well as expression in the pre-
synaptic neurons innervating the male reproductive system as evidenced by co-
localization with Syn (Figures 8S8, F and I). In the ED, 5-HT7-20XV5 is expressed in
the muscles (Figures 8S8, J and M), but unlike the SV, it is not expressed the pre-

synaptic neurons as it does not appear to co-localize with Syn (Figures 8S8, L and O).
Expression levels of 5-HT7-20XV5 increase in the SV near the point of fusion and are
maximal from just before the junction all the way to the terminus in the ED (Figure 8S9,
A). Expression levels of 5-HT7-20XV5 appear to increase at a discrete boundary point
on the SVs prior to fusion (arrow, Figure 8S9, D). This boundary of increasing
expression at this same region of the SV is highly reminiscent of OAβ2R-40XV5 (Figure
8S6, G). A higher resolution image of the sphincter region of the SV post-fusion reveals
high levels of 5-HT7-20XV5 expression (Figure 8S9, G) but not in the innervating pre-
synaptic neurons based on the absence of co-localization with Syn (Figure 8S9, I). A
cross-section of this same region reveals the 5-HT7-20XV5 protein localizes to muscle
membranes (Figure 8S9, J). A high-resolution image of the terminus of the sphincter
confirms 5-HT7-20XV5 expression extends all the way to the end (Figure 8S9, M). This
observation of 5-HT7-20XV5 expression throughout the fused sphincter region of the
SV contrasts with that of OAα2R-20XV5 (Figure 8S4, D) and OAβ2R-40XV5 (Figure
8S6, A), both of which end at the point of full SV fusion and suggests 5-HT plays a more
important role than OA in sphincter regulation. 5-HT7-20XV5 expression is also
elevated in the muscles at the terminus of the AGs (Figure 8S9, P). In the AGs, 5-HT7-
20XV5 is most prominently expressed in the underlying epithelial cells (Figures 8S10, A
and D). An AG cross-section reveals 5-HT7-20XV5 localizes exclusively to the apical
surface of the epithelial cells (Figure 8S10, G), highly reminiscent of OAβ2R-40XV5
localization to the apical surface of epithelial cells of the ED (Figure 8S6, D). Nearer the
surface of the AGs, low-level expression of 5-HT7-20XV5 is detected in the muscle
(Figure 8S10, J) and obvious expression is observed in the innervating pre-synaptic
neurons based on overlapping expression with Syn (Figure 8S10, L). 5-HT7-20XV5
expression was also observed in the ejaculatory bulb (Figures 8S10, M). Interestingly,
none of the OA or serotonin receptors exhibited preferential accumulation in the post-
synaptic muscle opposite the pre-synaptic terminals, as observed for GluRIIA. This
observation implies volume transmission may play a significant role in modulating
muscle activity in the male reproductive system. Neurotransmitter signaling via volume
transmission has previously been proposed for the Drosophila female reproductive
system based on expression patterns of OA receptors(ROHRBACH *et al.* 2024a).
637
SGNs are essential for male fertility but OGNs are dispensable
Having established that both OGNs and SGNs innervate the male reproductive
system an obvious question follows: what are their roles in male fertility?　It has
previously been reported that OA-deficient flies are male fertile and female sterile
(MONASTIRIOTI et al. 1996; COLE et al. 2005). However, in Tβh and Tdc2 mutant flies
that fail to synthesize OA, glutamate signaling in the OGNs innervating the male
reproductive system is still intact. To determine if simultaneously silencing the OGNs
innervating the male reproductive system for both glutamate and OA signaling causes
sterility, *Tdc2-GAL4-DBD* was combined with *AbdB-AD* and *UAS-BONT-C* to block all
synaptic transmission in these neurons. This intersectional combination results in the
silencing of only the Tdc2 neurons in the posterior VNC by virtue of the restricted
expression of the AbdB-AD hemidriver to the posterior few segments of the VNC. The
result was the same as eliminating OA signaling alone, males were fertile, and females
were sterile (Table 1), thus establishing that neither OA nor glutamate signaling is

required for fertility in the OGNs innervating the male reproductive system. The
observed female sterility validates the silencing function of the UAS-BONT-C transgene.
To assess the requirement of the SGNs innervating the male reproductive system for
fertility, flies containing TRH-DBD, AbdB-AD, and UAS-BONT-C were generated. Both
males and females of this genotype were sterile (Table 1). The observation that
females were also sterile, while no serotonergic innervation of the female reproductive
system has been reported, suggests the male sterility is due to 5-HT neurons common
to both males and females, and thus the male sterility cannot be unambiguously
attributed to the SGNs innervating the male reproductive system.
In an attempt to identify a split-GAL4 combination that more specifically targets
SGNs innervating the male reproductive system, several dozen hemi-drivers that
express in neurons whose cell bodies are located in the most posterior region of the
VNC were screened for male sterility in combination with UAS-BONT-C and either *TRH-*
AD or *TRH-GAL4-DBD*, as appropriate. One line, *VT019028-GAL4-DBD* was identified
that resulted in male sterility and female fertility, suggesting silencing of male-specific
sexually dimorphic neurons, and not neurons common to both males and females, were
responsible for the male sterility. The *TRH-AD/VT019028-GAL4-DBD* driver in
combination with a *UAS-His2A-GFP* nuclear reporter results in expression in sparse
neurons in the brain and VNC as well as around two dozen neurons at the posterior tip
of the VNC (Figure 9, A). Female flies of the same genotype exhibited a similar sparse
pattern of expression in the brain and VNC with the exception that there were noticeably
fewer His2A-GFP expressing neurons at the tip of the VNC (Figure 9, B). The neurons
most likely responsible for male sterility are those unique to males. The *VT019028-*
GAL4-DBD driver was also assessed for fertility in combination with *AbdB-AD* and *UAS-*
BONT-C with the same result of male sterility and female fertility (Table 1). The
expression patterns of *VT019028-GAL4-DBD* with *AbdB-AD* in combination with *UAS-*
His2A-GFP reveals restricted expression to the tip of the ventral nerve cord with
noticeably more neurons in the male (Figure 9A, C and D). These results with *AbdB-AD*
narrow the neurons responsible for the male sterility to the tip of the VNC. The
expression patterns of 5-HT and Tdc2 in these immunostains are also shown (Figures
9, E-H, and I-L, respectively).
To determine if any of the *VT019028-GAL4-DBD* neurons intersecting with *TRH-AD*
or *AbdB-AD* are serotonergic or octopaminergic, higher resolution images of the tip of
the VNC were examined. In the full confocal stack image, Three His2A-GFP neurons
are clearly visible that overlap with 5-HT (Figure 9S1, D). However, it has previously
been shown that there are both dorsal (~10 neurons) and ventral (~9 neurons) clusters
of serotonergic neurons innervating the male reproductive system (BILLETER et al.
2006b) and some neurons may thus be obscured from a dorsal perspective. A partial
stack containing the dorsal cluster reveals the same three neurons (Figure 9S1, I).
However, a ventral view of these neurons shows a fourth neuron overlapping with 5-HT
(Figure 9S1, K). A partial stack of the ventral cluster reveals three His2A-GFP neurons
overlapping with 5-HT (Figure 9S1, O). There also appears to be one or two Tdc2
neurons overlapping with His2A-GFP. However, examination of three-dimensional
images revealed they are only in close proximity but not overlapping.
Similar results were obtained with *VT019028-GAL4-DBD* neurons intersecting with
AbdB-AD. Four His2A-GFP neurons overlap with 5-HT in a full confocal stack view

(Figure 9S2, D). These four neurons are visible in the dorsal serotonergic cluster
(Figure 9S2, I), while three additional His2A-GFP neurons overlapping with 5-HT are
apparent in the ventral serotonergic neuron cluster (Figure 9S2, N). One His2A-GFP
neuron appears to overlap with Tdc2 (Figure 9S2, E), but this apparent overlap is not
reproduced in the dorsal (Figure 9S2, J) or ventral (Figure 9S2. O) partial stacks. The
innervation of *VT019028-GAL4-DBD* neurons with *TRH-AD* and *AbdB-AD* visualized
with the *UAS-CD8-mCherry* reporter is similarly broad in the SVs, ED, and AGs (Figure
9S3, A and B, respectively). Taken together, these results strongly suggest synaptic
transmission in the SGNs included in the *VT019028-GAL4-DBD* intersections with *TRH-*
AD and *AbdB-AD* that innervate the male reproductive organs are required for male
fertility, although a role for the small number of other interneurons at the tip of the VNC
in common between both combinations cannot be excluded.
To rule out the possibility that the male sterility observed in the *VT019028-GAL4-*
DBD intersections with *UAS-BONT-C* and either *TRH-AD* and *AbdB-AD* is due to failure
to mate, mating assays were performed where single males and single females of the
different genotypes were placed together in a well of a 12-well plate and videotaped.
The mating success rates of *VT019028-GAL4-DBD/TRH-AD* and *VT019028-GAL4-*
DBD/AbdB-AD were 81.25% (n=16) and 57.9% (n=28), respectively. These results
demonstrate the observed male sterility is not due to failure to copulate.
In these mating assays, mating duration was also assessed along with several
control genotypes. Mean mating durations for *VT019028-GAL4-DBD/TRH-AD* and
VT019028-GAL4-DBD/AbdB-AD were 28.74 and 24.75 minutes, respectively (Figure
10, A). This was not significantly different from *Canton-S* wildtype controls (26.91), and
other controls missing one or more of the components necessary for silencing *AbdB-*
AD, *UAS-BONT-C* (27.07) and *UAS-BONT-C* alone (28.75) (Figures 10, B and C).
These results are in line with previously reported mating durations for Drosophila
(CRICKMORE AND VOSSHALL 2013). Interestingly, silencing all 5-HT neurons (*TRH-GAL4-*
DBD ∩ *AbdB-AD*) or all OA neurons (*TRH-GAL4-DBD* ∩ *AbdB-AD*) innervating the
male reproductive system resulted in statistically significant reductions in mating
duration, 19.11 (Figure 10B, D) and 21.8 (Figure 10B, E), respectively.
To assess the distribution of sperm in the reproductive system during mating by
VT019028-GAL4-DBD/TRH-AD > UAS-BONT-C sterile males, a Protamine-GFP
reporter that fluorescently labels sperm nuclei was incorporated into the genotype. For
this experiment, control and experimental males were placed in petri dishes with virgin
females, separated 10 minutes after the initiation of mating, immediately dissected, and
the isolated male reproductive systems video recorded using a stereofluorescent
microscope. In the control males, ProtB-GFP sperm were readily visible in the ED
(Figure 10S1, A, Video 1). In contrast, in the experimental males, no ProtB-GFP sperm
were observed (Figure 10S1, B, Video 2). These results demonstrate that the male
sterility is due to a failure of the sphincter that regulates the flow of sperm between the
SV and ED to open and thus the defect is with the emission phase of ejaculation.
739
Glutamate signaling in OGNs and SGNs is dispensable for male fertility
In the *VT019028-GAL4-DBD/TRH-AD > UAS-BONT-C* sterile males, 5-HT and
glutamate signaling are both abolished due to the BONT-C blockade of synaptic
transmission. To determine if glutamate signaling alone is required in these neurons, a

conditional allele of vGlut was incorporated into this genotype such that specifically
glutamate signaling was eliminated while serotonergic signaling remained intact. These
males were determined to be fertile. To ascertain whether there were changes in the
level of fertility, fertility assays were conducted on 5-day post-eclosion males. No
changes in fertility as compared to controls were observed (Figure 10S2, A).
Immunostaining the AGs of control and experimental flies confirmed the effectiveness of
the conditional vGlut allele by the presence of vGlut in control flies and its absence in
experimental flies (Figure 10S3, A and D, respectively). As glutamate has been shown
to promote neuron survival as neurons age (SHEN et al. 2018; BUCK et al. 2023), to
determine if fertility might be altered in older males, the same 5-day old males used in
the previous fertility experiment were held in isolation from females until they were 30
days post-eclosion and the experiment was repeated. While there was not
unexpectedly a modest decrease in fertility with age, no differences in fertility were
apparent between the control and experimental genotypes (Figure 10S2, B). These
results establish that glutamatergic signaling is not required in *VT019028-GAL4-*
DBD/TRH-AD neurons for fertility. To determine if more extensive elimination of
glutamate signaling in the OGNs and SGNs would alter fertility, glutamate was
specifically eliminated in both classes of neurons. No reductions in fertility were
observed in 5-day post-eclosion males (Figure 10S2, A). These results indicate
glutamate is not required in either OGNs or SGNs for fertility.
764
The sphincter region of the SV initiates ED peristaltic waves
Neuron-independent peristaltic waves are known to occur spontaneously in the male
ED of Drosophila that persist even when all nerves innervating the male reproductive
system have been severed (NORVILLE et al. 2010). To gain a better understanding of
the muscle activity occurring during these waves in both the ED and throughout the rest
of the male reproductive system, a fly strain was generated that expressed the Ca++
sensor GCAMP8m specifically in muscles under control of the myosin heavy chain
promoter (*MHC-GCAMP8m*). This fly strain allows visualization of the changes in Ca++
levels that occur in the muscles during these peristaltic waves. As expected, waves of
fluorescence are visualizable coincident with peristaltic waves of muscle contractions in
the ED (video 3). These peristaltic waves originate not in the ED but in the sphincter
region of the SV just prior to and after the point of SV fusion (video 4), although the
region of the SV prior to the fusion point that is involved in initiation varies (video 5).
Muscle contractions in other regions of the male reproductive system are also apparent.
These contractions are weak and seemingly random in the SV (video 6), AG (video 7),
and testes (video 8), but occasionally in the AG strong coordinated contractions are
observed (video 9). Together, these imaging data reveal that the seminal-vesicle
sphincter acts as an autonomous pacemaker, launching neuron-independent peristaltic
waves that drive ejaculatory-duct motility.
784
785 **Discussion**
786
787 　　The male reproductive tract of flies and mammals have obvious morphological
788 differences, yet they share many essential anatomical features and a common
789 organizational logic. During copulation Drosophila males deliver seminal fluid and

~4000-6000 sperm to females during a tightly choreographed ~20-minute window (KAUFMAN 1942; GILBERT 1981). This involves the emission of sperm and seminal fluid from their sites of origin into the ED where they mix and are subsequently expelled into the female reproductive tract. Despite the critical importance of ejaculation, its neuronal mechanisms have remained poorly defined, hindering broader insights into the neural logic of male fertility. Here we close this gap by demonstrating that two distinct yet intermingled classes of amine/glutamate neurons, OGNs and SGNs, coordinate the ejaculatory sequence that ensures successful reproduction. Their transmitter pairs confer the capacity to co-release both fast excitatory and slower acting aminergic modulatory neurotransmitters. Differential expression of ionotropic glutamate receptors and aminergic metabotropic neurotransmitter receptors combine in post-synaptic muscle across the male reproductive system to confer an additional level of complexity to coordinately regulate fluid secretion, organ contractility, and directional sperm movement, to the end of optimizing male fertility.

This system can serve as a genetically tractable model to gain general insights into how multi-transmitter neurons coordinate and modulate interrelated action sequences critical for complex organ function. Lessons learned from this system may have implications for understanding how males of Drosophila and other species adapt their reproductive strategies to a dynamic environment. Additionally, understanding the mechanisms by which this system operates has the potential to identify neuromodulation strategies for male infertility and prostate cancer as multi-transmitter sympathetic and parasympathetic neurons innervating human male reproductive organs have been implicated in these conditions (BURNSTOCK 2009; MAGNON *et al.* 2013; BURNSTOCK 2014; DROBNIS AND NANGIA 2017; DWIVEDI *et al.* 2021).

Mechanisms of multi-transmitter neuron-mediated transfer of sperm and seminal fluid

Movement of sperm and seminal fluid through the male reproductive tract requires an alternating series of coordinated muscle contractions and relaxations. Based on the observed co-packaging of vMAT and vGlut in the majority of synaptic vesicles in the ED of both the OGNs and SGNs, combined with the observed non-uniform distribution of neurotransmitter receptors for OA, 5-HT, and glutamate in neurons and muscles, potential pre- and post-synaptic mechanisms for regulating the activity of the muscles of the internal male reproductive system are possible.

Pre-synaptic expression of the α-type OA receptors OAMB and OAα2R may facilitate alternating bursts of action potential-dependent neurotransmitter release by diminishing pre-synaptic neurotransmitter release via an auto-receptive mechanism. As α-type adrenergic and noradrenergic receptors have been consistently shown to inhibit neurotransmitter release in multiple rodent species in both the peripheral and central nervous system including the vas deferens, heart, and brain when expressed in pre-synaptic neurons (STARKE 1972; TRENDELENBURG et al. 1994; MIYAZAKI et al. 1998; SCHEIBNER et al. 2001; TRENDELENBURG et al. 2003), the presence of OAMB and OAα2R in the pre-synaptic terminals of neurons innervating the internal male reproductive organs may have the effect of suppressing neurotransmitter output from pre-synaptic terminals and shortening the duration of muscle contractions. Thus, pre-

synaptic expression of OAMB and OAα2R may have the effect of increasing the cycling
frequency of muscle contractions and relaxations.
As glutamate is acting via post-synaptic GluRII-type receptors, its effect will almost
certainly be excitatory with the result of eliciting fast muscle contractions. In contrast,
the functional consequences of OA and 5-HT are not so clear cut. Either of these
amines could enhance muscle contractions or facilitate muscle relaxation, depending on
the neurotransmitter receptor(s) receiving the signal. In muscles of the mammalian
prostate and lower urinary tract, α-type adrenergic and noradrenergic receptors have
been demonstrated to augment muscle contractions while β-type receptors have been
found to facilitate muscle relaxation (HONDA et al. 1985; MICHEL AND VRYDAG 2006;
WUEST et al. 2009). In contrast, OAMB and OAβ2R have been shown to enhance
muscle relaxation and contraction, respectively in the Drosophila female reproductive
system (DESHPANDE et al. 2022).
5-HT7 receptors are known to promote smooth muscle relaxation in the urinary tract,
gut, oviduct, and blood vessels of mammals (CARTER et al. 1995; LEUNG et al. 1996;
PRINS et al. 1999; INOUE et al. 2003; RECIO et al. 2009; CHANG CHIEN et al. 2015;
MATSUMOTO-MIYAI et al. 2015). Although muscle contraction receives preferential focus,
muscle relaxation is also critically important for the execution of motor programs
(GOULDING et al. 2014; WONG AND LANGE 2014; GOWDA et al. 2018; HIRAMOTO et al.
2021) and 5-HT7 most likely plays a role in mediating muscle relaxation in the internal
male reproductive system. An additional confounding factor is that OAα2R has been
shown to respond not just to OA but also to 5-HT (QI et al. 2017). The observations of
the presence of both muscle contraction enhancing, and muscle relaxation promoting,
OA and 5-HT receptors in the same muscles of the internal male reproductive organs
indicates the mechanisms of neuronal regulation of their activity are complex and may
even involve synergistic effects that can only be established empirically.
861
A potential timing mechanism mediated by multi-transmitter neurons
Pre-synaptic release of glutamate acting on post-synaptic ionotropic GluRII-type
glutamate receptors will cause near immediate, short-term, contraction of muscles. At
least in the ED, our results show that the majority of synaptic vesicles co-package either
OA or 5-HT along with glutamate, and thus simultaneous co-release is being used as a
mechanism of neurotransmitter signaling. The amines OA or 5-HT acting on post-
synaptic metabotropic OA and 5-HT receptors will have the effect of either prolonging
the length of, or augmenting the relaxation of, muscles, dependent on the specific
aminergic neurotransmitter receptors involved. These latter effects will be delayed
relative to the glutamate-induced fast muscle contractions due to the discrepancy in the
time course of action between ionotropic and metabotropic receptors. Thus, co-
releasing a fast-acting excitatory neurotransmitter acting through an ionotropic receptor
simultaneous with a modulatory aminergic neurotransmitter acting through slower acting
GPCRs in the same multi-transmitter neuron may create a sequential timing mechanism
whereby fast glutamate-mediated muscle contractions always precede either amine-
mediated prolonging of muscle contraction or facilitation of muscle relaxation. A single
neuron co-releasing one fast and one slow neurotransmitter may thus have the
advantage over two distinct neurons separately releasing these transmitters in that the

former can more effectively synchronize, and at the same time, stagger events that
require precise timing.
882
Potential for adaptation to environment
The complexity of neuronal control of male reproductive function revealed in this
investigation exposes a variety of possibilities for adaptive regulation to dynamic
environmental conditions. Polygynous Drosophila males are well-established to alter
the quantity and quality of their ejaculate based on environmental circumstances,
especially the level of competition with other males (BRETMAN et al. 2009; MOATT et al.
2014), but also due to nutrient availability (MACARTNEY et al. 2021) and the presence of
pathogens (LIAO et al. 2024). The male reproductive system may adapt to
environmental conditions by tuning octopamine and serotonin signaling, adjusting the
levels of their biosynthetic enzymes, the vesicular monoamine transporter vMAT, or
relevant receptors to alter the amount of sperm and seminal fluid released during
copulation. A potential mechanism for altering OA and/or 5-HT levels is extra-synaptic
volume transmission whereby the level of OA or 5-HT secreted into the hemolymph in
locations anatomically distant from the male reproductive system. Thus, global levels of
OA and 5-HT in the hemolymph could be adjusted based on environmental conditions
and this signal could then impact the male reproductive system to optimize ejaculate
quality and quantity. Finally, the presence of OAMB and 5-HT7 on epithelial cells
suggest the potential for the regulation of the quality of seminal fluid as the expression
of these receptors implies, they could differentially alter gene expression in the epithelial
cells in response to changing levels of OA and 5-HT. The results of this study create a
foundation for mechanistic investigations into the molecular basis by which male
reproductive resources are adaptively modulated.
905
Potential for signaling via neuropeptides
The broad abundant expression of the LDCV reporter IA2 in the neurons innervating
the internal organs of the male reproductive system suggests a significant role for
neuropeptide signaling since neuropeptides are known to be packaged into LDCVs
(PELLETIER AND LECLERC 1979; KREINER et al. 1986). Bolstering the prospect of this
possibility, numerous neuropeptides across evolutionary distant insect species have
been shown to alter the frequency and/or amplitude of the spontaneous peristaltic
waves in the ED of male insect reproductive systems (MARCINIAK et al. 2017; CIZMAR et
al. 2019; LANGE et al. 2023). Multiple neuropeptides have also been implicated in the
function of the internal male reproductive organs of mammals, including humans,
(SEGAWA et al. 1978; ALLEN et al. 1982; TAINIO 1995). Given these considerations, it
thus seems likely that neuropeptides outside the scope of this investigation also play a
role in regulating Drosophila male reproduction.
919
The SV/ED junction is a control center
Immediately proximal to the ED, SVs fuse into a single duct that forms the sphincter
regulating sperm flow from the SV to the ED during mating. The differential expression
patterns of OAα2R, OAβ2R, and 5-HT7 in this region of the SV is intriguing. The extent
of OAα2R and OAβ2R expression ends very shortly after the merge, at precisely the
point where full SV fusion occurs and there is no longer a partition separating the ducts

(BAIRATI 1968). Only 5-HT7 expression extends into the fully fused SV that constitutes
the sphincter separating the SV from the ED. This expression pattern implies a
prominent role for 5-HT7 in sphincter regulation. Previous reports of 5-HT7 mediating
relaxation of smooth muscle in the mammalian gut (CARTER et al. 1995; PRINS et al.
1999), urinary bladder neck (RECIO et al. 2009), cardiovascular system (LEUNG et al.
1996; CHANG CHIEN et al. 2015), lymphatic system (CHAN AND VON DER WEID 2003), and
oviduct (INOUE et al. 2003) raises the possibility it may function to relax the sphincter
muscles during copulation to permit sperm flow. Additionally, expression of OAβ2R and
the likely relaxation-promoting 5-HT7 receptors increases sharply in the SV just prior to
the point of fusion. Perhaps not coincidentally, this limit of expression corresponds with
the distal-most extent of sperm localization in the SV where relaxation of the SV
muscles must also occur for sperm to move along the duct to the ED. It is also notable,
as revealed in the Ca++ imaging experiments, that the sphincter region is the site of
initiation of the peristaltic waves of muscle contractions that are promulgated through
the ED. The SV/ED junction is thus a critical control center for both regulation of sperm
flow into the ED during mating and initiating neuron-independent peristaltic waves in the
ED.
943
Molecular continuity of neurotransmitter signaling between males and females?
The expression of 5-HT7 and OAβ2R on the apical surface of the epithelial cells
lining the AG and ED, respectively, is surprising because it implies that levels of 5-HT
and OA are fluctuating in the lumen of these organs. The origin of these
neurotransmitters is not obvious but if OA and 5-HT are present in the lumen of the AG
and ED, another intriguing possibility is that these neurotransmitters are transferred to
the female at mating for signaling in order to modulate female behavior and/or
reproductive activity. A recent study detailed the molecular continuity of proteins in the
seminal fluid of males functionally interacting with proteins in the female reproductive
tract (MCCULLOUGH et al. 2022). The potential presence of OA and 5-HT in the lumen of
the AG and ED raises the possibility that this molecular continuity could extend to
neurotransmitters in male seminal fluid signaling to neurotransmitter receptors in the
female reproductive tract to convey information such as health, internal state, or
environmental conditions to regulate female reproductive output. A detailed
examination of the expression patterns of OA receptors in the female reproductive tract
indicates this is indeed a possibility, at least for OA signaling (ROHRBACH *et al.* 2024b).
960
961 **Materials and methods**
962
Fly strains
Stocks from the Bloomington Drosophila Stock Center (NIH P40OD018537) were used
in this study. *Tdc2-GAL4* (COLE et al. 2005) RRID:BDSC_9313, *GluRIIA-GAL4*
RRID:BDSC_84637, *GluRIID-GAL4* RRID:BDSC_84638, and *TRH-GAL4*
RRID:BDSC_84694 (DENG et al. 2019)
vGlut-GAL4 RRID:BDSC_60312, *vGlut-GAL4-DBD* (DIAO et al. 2015)
Tdc2-AD (DIONNE et al. 2018) RRID:BDSC_601902
TRH-GAL4-DBD RRID:BDSC_70371
TRH-AD RRID:BDSC_70975

GluRIIB-GAL4 RRID:BDSC_60333 and *GluRIIE-GAL4* RRID:BDSC_60332
(Diao et al. 2015)
Oct-TyrR-GAL4 RRID:BDSC_77735 and *GluRIIC-GAL4* RRID_83268 (Lee et al. 2018)
GluRIIA-GFP RRID:BDSC_99517 (Beckers et al. 2024)
5-HT1A-GAL4 RRID:BDSC_84588, *5-HT1B-GAL4* RRID:BDSC_84589 , *5-HT2A-GAL4*
RRID:BDSC_84590, *5-HT2B-GAL4* RRID:BDSC_84591, and *5-HT7-GAL4*
RRID:BDSC_84592 (Deng et al. 2019)
TβH-GFP RRID:BDSC_80113 (Li-Kroeger et al. 2018)
fru-GAL4 RRID:BDSC_66696
UAS-CD8-mCherry RRID:BDSC_27392
ProtB-GFP (Manier et al. 2010) RRID:BDSC_58406
AbdB-AD (Diao et al. 2024)
vGlut-AD (Gao et al. 2008)
IA2-GFP (Yu et al. 2025)
UAS-BONT-C (Han et al. 2022)
dsx-GAL4-DBD (Shirangi et al. 2016)
vGlut-40XV5 and *vGlut-40XMYC* (Stowers 2025)
B2RT-7XMYC-vAChT (Tison et al. 2020)
RSRT-9XV5-vGAT (Certel et al. 2022)
OAMB-GAL4, *OAα2R-GAL4*, *OAβ1R-GAL4*, *OAβ2R-GAL4*, *OAβ3R-GAL4* (McKinney
et al. 2020)
FRT-F3 OAMB-10XV5 (Bakshinska et al. 2025)
B3RT-Tdc2-LexA, *B3RT-vGlut-LexA*, *RSRT-STOP-RSRT-6XV5-vMAT*, *UAS-B3* and
20XUAS-R (Sherer et al. 2020)
20XUAS-B2 (Williams et al. 2019)
13LexAop-6XmCherry (Shearin et al. 2014)
998
Stocks original to this report
Tdc2-GAL4-DBD (*JK65C*)
B3RT-Tdc2-LexA germline excision
B3RT-vGlut-LexA germline excision
RSRT-6XV5-vMAT germline excision
FRT-F3-OAα2R-20XV5
FRT-F3-OAα2R-20XV5 germline inversion
B2RT-STOP-B2RT-OAβ2R-40XV5
B2RT-OAβ2R-40XV5 germline excision
5-HT7-20XV5
MHC-GCAMP8m (*VK27*)
1010
Entry clones
The *R4-GAL4-DBD-R3* entry clone contains a 640bp insert encoding the GAL4 DNA
binding domain. The *L1-MHC-5'Reg-R5* entry clone contains 5254bp of upstream
regulatory sequence from the Drosophila Myosin Heavy Chain (MHC) gene. The *L5-*
GCAMP8m-L2 entry clone contains a 1276bp insert encoding GCAMP8m (Zhang et al.
2023). Entry clones were generated as previously described (Petersen and Stowers
2011).

1018

Expression clones
Tdc2-GAL4-DBD was constructed by combining entry clones *L1-Tdc2-5'Reg-L4* and *L3-*
Tdc2-3'Reg-L2 (SHEARIN et al. 2013) with the *R4-GAL4-DBD-R3* entry clone and the
destination vector *pDESTP10* (SHEARIN et al. 2013). *MHC-GCAMP8m* was assembled
by combining entry clones *L1-MHC-5'Reg-R5* and *L5-GCAMP8m-L2* with *pDESTP10-*
SPEC. *pDESTP10-Spec* is a derivative of *pDESTP10* in which AmpR was replaced
with SpecR.
1026

Donor plasmid assembly
Donor plasmids were assembled starting with gene synthesis of the homology arms
(Synbio). The gene synthesis also included recombinase target sites and mutations in
the PAM sites associated with the guide RNAs. Homology arms typically extended
~750bp beyond the cut sites of the guide RNAs. STOP cassettes were added via
Gibson cloning. Multimerized epitope tags (STOWERS 2025) were added via *Asc I/Not I*
restriction enzyme cloning. The *OAα2R-FRT-F3-20XV5* donor plasmid contained a pair
of *FRT-F3* sites appropriately arranged for conditional expression of OAα2R-20XV5
after inversion/excision by FLP recombinase (FISHER *et al.* 2017) with the 20XV5 fused
to the carboxy-terminus. The *OAβ2R-B2RT-STOP-B2RT-40XV5* donor plasmid
contained B2 recombinase target sites flanking a STOP cassette such that OAβ2R-
40XV5 was expressed after excision of the STOP cassette with 40XV5 fused to the
carboxy-terminus. The 5-HT7-20XV5 donor plasmid contained a 20XV5 multimerized
epitope tag fused to the carboxy-terminus but was not designed for conditional
expression.
1042

Guide RNA plasmid assembly
Guide RNAs were assembled in the *pCFD4* double-guide RNA plasmid as previously
described (PORT *et al.* 2014). Two double guide RNA plasmids were used with each
donor construct for genome editing. The guide RNA plasmids and the guide RNAs they
contain are as follows: *pCFD4-OAα2RGd12*-TCTAAATCAATATCAGTG and
AACATAAACATCCTAAAA; *pCFD4-OAα2R-Gd34*-ACCTGTAACAGCACACAT and
GGCATGTATCAGCACTAC; *pCFD4-OAβ2R-Gd56*-ATAAATGAGATTATACTC and
CATGCATCTGGAATTTAT; *pCFD4-OAβ2R-Gd78*-CGGCGGGCGGATCCCGCC and
AGGATGGCAGGGAGTCGA; *pCFD4-5-HT7-Gd56*-TGGGCGATGAGAGGCACG and
TGGGCGATGAGAGGCACG; *pCFD4-5-HT7-Gd78*-CCACGAGGACCTCCATTC and
GGGTCTTCCCTAGGTTTC.
1054

Genome editing
For each genome edit, the donor plasmid was injected together with both *pCFD4* guide
RNA plasmids into the *nos-Cas9 attP40* fly strain by Rainbow Transgenic Flies. Adult
flies that were injected as embryos were crossed to 3rd chromosome balancer stocks.
Candidate progeny males from this first cross containing the *TM6b* balancer
chromosome were crossed to yw; n-syb-GAL4, UAS-FLP/TM6c (*OAα2R-FRT-F3-*
20XV5, yw; 20XUAS-B2; n-syb-GAL4 (*OAβ2R-B2RT-STOP-B2RT-40XV5*) or *yw;*
Ly/TM6c (*5-HT7-20XV5*) fly strain and larval progeny were screened by

immunostaining. Males whose progeny exhibited positive immunostaining were
subsequently crossed to a third chromosome balancer stock to establish stable lines.
1065

Germline excision/inversion
Germline inversion of *OAα2R-FRT-F3-20XV5* was accomplished by crossing to *yw*;
nos-GAL4; *UAS-FLP*. Males containing all three genetic components were crossed to a
third chromosome balancer stock. Individual progeny males from this cross were
crossed a second time to the third chromosome balancer stock and non-Tb larva were
screened by anti-V5 immunostaining to identify germline excisions. A similar strategy
was used with *OAβ2R-B2RT-STOP-B2RT-40XV5* for a germline excision except the
initial cross was to *yw*; *nos-GAL4*; *UAS-B2*. *vGlut-LexA* was generated by crossing
B3RT-vGlut-LexA to *yw*: *nos-GAL4*; *UAS-B3*. Progeny from this cross containing all
three genetic components were crossed to a second chromosome balancer stock.
Single progeny males were then crossed to a *13XLexAop-6XmCherry* reporter and
larva were screened for expression in the ventral nerve cord to identify germline
excision positives. A similar strategy was used to create *B3RT-Tdc2-LexA* starting with
B3RT-Tdc2-LexA (MCKINNEY et al. 2020). *RSRT-6XV5-vMAT* was generated by
crossing *RSRT-STOP-RSRT-6XV5-vMAT* with *yw*; *nos-GAL4*; *UAS-R*. Progeny from
this cross containing all three genetic components were crossed to a second
chromosome balancer stock. Single progeny males were crossed to the second
balancer a second time and larval progeny were subjected to anti-V5 immunostaining to
identify germline excisions.
1085

Immunostaining
Male reproductive systems were dissected in PBS and fixed in 4% paraformaldehyde
for 30 minutes with rotation. After 2X brief washes with PBS, they were incubated in
PBS containing 2% Triton-X 100 and 2% BSA (PBSTB) for at least two hours with
rotation. Subsequently, they were incubated in primary antibodies PBSTB for one hour
with rotation, followed by overnight incubation at 4$^\text{o}$C without rotation to prevent tissue
damage. After 3X 5-minute washes in PBSTB, male reproductive systems were
incubated for 2-3 hours in PBSTB with secondary antibodies with rotation. After three 5-
minute washes in PBS, they were incubated in PBS containing Phalloidin 405 for one
hour. After two 5-minute washes in PBSTB and one 5-minute wash in PBS, male
reproductive systems were mounted on glass slides in Vectashield Plus (Vector
Laboratories, Cat. #H-1900) using a coverslip that was glued on the edges with clear
nail polish.
1099

Primary antibodies and dilution factors: The SYN (3C11) mAb 1:20 (KLAGGES et al.
1996). Brp nc82 mAb 1:40 (deposited at the DSHB by Eric Buchner), Dlg 4F3 mAb 1:40
(PARNAS *et al.* 2001), were obtained from the Developmental Studies Hybridoma Bank,
created by the NICHD of the NIH and maintained at The University of Iowa, Department
of Biology, Iowa City, IA 52242. Mouse anti-V5 SV5-P-K (Novus NBP2-52703) 1:200;
rat anti-V5 SV5-P-K (Novus NBP2-81037) 1:200, rabbit anti-V5 SV5-P-K (Novus NBP2-
52653) 1:200; Human anti-V5 (Novus NBP2-81035) 1:200; rat anti-MYC 9E10 (Novus
NBP2-81020) 1:200, rabbit anti-MYC 9E10 (Novus NBP2-52636), mouse anti-MYC
4A6; (Millipore Sigma) 1:200, mouse anti-vGlut 1:10 (BANERJEE et al. 2021), mouse anti-

GFP 3E6 (Thermo-Fisher A11120) 1:200; Rabbit anti-GFP (Thermo-Fisher G10362); rat
anti-mCherry 16D7 (ThermoFisher-M11217) 1:200, rabbit anti-Tdc2 pab0822-P,
Covalab) 1:500; rat anti-5-HT YC5 (Abcam ab6336) 1:300.
1112
Secondary antibodies and dilution factors: Goat anti-Mouse Alexa 405 (Thermo-Fisher
A31553) 1:400; Goat anti-Mouse Alexa 488 (Jackson Immunoresearch 115-545-166)
1:400; Donkey anti-Rat Alexa 488 (Jackson Immunoresearch 712-546-153) 1:400, Goat
anti-Rabbit Alexa 488 (Thermo-Fisher A32731) 1:400, Goat anti-Mouse Alexa 568
(Thermo Fisher A11031) 1:400, Goat anti-Rabbit Alexa 568 (Thermo Fisher
A11036)1:400, Donkey anti-Rat Alexa 568 (Thermo-Fisher A78946) 1:400; Goat anti-
Mouse Alexa 647 (Thermo-Fisher A32728) 1:400; Goat anti-Rabbit Alexa 647 (Thermo-
Fisher A32733) 1:400; Goat anti-Rat Alexa 647 (Thermo-Fisher A48265 1:400); Donkey
anti-Human Alexa 790 (Jackson Immunoresearch 709-655-149) 1:400. Muscles were
stained with Phalloidin iFluor405 (Abcam AB176752) 1:800.
1123
1124 Confocal microscopy was performed using a Leica Stellaris DMI8 inverted confocal
1125 scanning light microscope in the microscopy facility at the Montana State University
1126 Center for Biofilm Engineering.
1127
Expansion microscopy
The Ten-fold Robust Expansion Microscopy (TREx) protocol was used to expand the
male ED (ED) (DAMSTRA et al. 2023). Briefly, Drosophila male reproductive systems
were dissected and fixed for at least thirty minutes in 4% paraformaldehyde. The ED's
were then dissected out and immunostained as described above. Tissues were then
incubated in Acryloyl-X solution, 1:100 dilution (Invitrogen, catalog #2604348),
overnight, at room temperature and protected from light. The next day, the ED's were
rinsed in PBS and incubated in TRex solution at 4 C, for 45-60 minutes. 0.15%
ammonium persulfate and 0.15% TEMED was added and the TREx mixture was quickly
transferred to chamber slides. TREx mixture with ED's were placed in wells (1 ED per
well), coverslipped and allowed to polymerize at 37 C for at least 45 minutes. After gel
polymerization, each gel-ED was trimmed to eliminate excess gel. Gel-ED's were
incubated in proteinase K (Biolabs, catalog #P81075, 800U/ml) 1:100 solution,
overnight at room temperature. Expansion of gel-ED's was done the following day.
Three washes in water were performed, at least 20 minutes per wash. DAPI
(Thermofisher Scientific, catalog# D1306), 2ug/ml, was added in the first wash to aid in
tissue visualization. After the third wash, hydrogel-ED's were trimmed again and placed
in chamber slides. 1 hydrogel-ED per well and coverslipped. Imaged expanded ED's in
upright confocal (Leica Stellaris 8 DIVE) equipped with a ×40 NA = 0.8 water objective. z
Stack- images were deconvoluted using the Lightning confocal function and 3D images
were generated for each image. ED's typically expanded approximately tenfold.
1149
Colocalization Analysis
Digitally deconvolved- 3D images of expanded EDs were analyzed using ImageJ
software (SCHINDELIN et al. 2012). To exclude background in an unbiased manner, we
thresholded all images by using the Otsu threshold and calculated Manders' coefficients
by using the BIOP JACop.plugin. A scatter plot of green and red pixel intensities

(associated with each of the two markers, Vmat and Vglut) was generated by using the
colocalization function. Manders' coefficient ranges from 0 to 1, indicating no or
complete colocalization. Prior to doing the colocalization analysis of each image, a
region of interest (ROI) was traced using the stain from the CD8 marker. This ROI
corresponded to an axon terminal.
1160
Mating assays
Mating assays involved placing single 3–7-day old virgin males with single 3-7-day old
virgin females in the same well of a 12 well tissue culture plate without anesthesia. The
wells of the 12 well plates were half-filled with agarose. Flies were videotaped for three
hours. Mating assays were conducted at 25C and 50% humidity in a room dedicated to
behavioral assays. Mating was scored by viewing the videos and scorers were blind to
the genotypes.
1168
Fertility assays
The fertility assays involved placing single 5 day old or 30-day old virgin males with four
3-7-day old virgin females in fresh vials without anesthesia for six hours. After six hours
the males were removed and the four females returned to the vials for 72hrs at which
point the females were removed. Six days after removal of the females the pupae on
the walls of the vials were counted.   The counting was done blind to the genotypes.
Fly culturing for all stages of the fertility assays were conducted at 25C and 50% in an
incubator.
1177
1178 **Animal use**
1179 All experiments involving *Drosophila melanogaster* were conducted in accordance with
1180 approved protocols and standard laboratory practices. *Drosophila melanogaster* is not
1181 considered a vertebrate animal and accordingly does not require Montana State
1182 University Institutional Animal Care and Use Committee (IACUC) approval.
1183
1184 **Data availability**
1185 All plasmids and their complete sequences are available upon request. Fly strains
1186 original to this publication will be deposited at the Bloomington Drosophila Stock Center
1187 or made available upon request.
1188
Acknowledgements

Stocks obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537)
were used in this study. Antibodies used in this study were obtained from the
Developmental Studies Hybridoma Bank created by the NICHD of the NIH and
maintained at The University of Iowa, Department of Biology. This work used a Leica
Stellaris DMI8 inverted confocal scanning light microscope in the Center for Biofilm
Engineering-Bioimaging and Analytical Cores, RRID:SCR_026519, which is supported
by Montana State University. This work was supported by National Institutes of Health
(NIH) grant 2R01GM115510 to SJC and RSS.

1198
1199
1200 **Figure Legends**

1201

1202
1203   **Figure 1.** The Drosophila male reproductive system. A) schematic diagram showing
1204   paired testes, SVs (green), and AGs (purple), and a singular sphincter (red), ED (gray),
1205   and ejaculatory bulb. B) actual male reproductive system. Te-testes, SV-seminal
1206   vesicle, AG-accessory gland, Sph-sphincter, ED-ejaculatory duct, EB-ejaculatory bulb.
1207   Scale bar: 200μm.

1208
1209   **Figure 2.** Gal4 and Split-GAL4 expression patterns in the Drosophila male reproductive
1210   system. A) *Tdc2-GAL4*; B) *TRH-GAL4*; C) vGlut-GAL4; D) Tdc2-AD ∩ *vGlut-GAL4-*

1211   *DBD*; E) *vGlut-AD ∩ Tdc2-GAL4-DBD*; F) *TRH-AD ∩ vGlut-GAL4-DBD*; G) *vGlut-AD ∩*

1212   *TRH-GAL4-DBD*; H) *Tdc2-AD ∩ TRH-GAL4-DBD*. Scale bar: 200μm.

1213

1214   **Figure 3.** Expression of *TRH-GAL4* and *Tdc2-LexA* in the Drosophila male

1215   reproductive system. A, D, G, J, M, P) *TRH-GAL4, UAS-6XGFP*; B, E, H, K, N, Q)

1216   *Tdc2-LexA, LexAop-6XmCherry*. C, F, I, L, O, R) overlay. Scale bars: O-50μm; R-

1217   10μm.

1218

1219 **Figure 3S1.** Expression of *vGlut-LexA* and *TRH-GAL4* in the Drosophila male
1220 reproductive system. A, D, G, J, M, P) *vGlut-LexA*, *LexAop-6XmCherry*; B, E, H, K, N,
1221 Q) *TRH-GAL4, UAS-6XGFP.* C, F, I, L, O, R) overlay. Scale bars: O-50μm; R-10μm.

1222

1223 **Figure 3S2.** Expression of *vGlut-LexA* and *Tdc2-GAL4* in the Drosophila male
1224 reproductive system. A, D, G, J, M, P) *vGlut-LexA*; B, E, H, K, N, Q) *Tdc2-GAL4*,
1225 *LexAop-6XmCherry*. C, F, I, L, O, R) overlay. Scale bars: O-50μm; R-10μm.

1226
1227 **Figure 3S3.** Fruitless and Doublesex expression in the Drosophila male reproductive
1228 system. A) *fru-GAL4*, *UAS-6XGFP*; B) *vGlut-LexA*, *LexAop-6XmCherry*; C) overlay; D)
1229 Tdc2-AD ∩ dsx-GAL4; E) TRH-AD ∩ dsx-GAL4. Scale bar: 200μm.

1230

1231 **Figure 4.** Expression of vGlut, TβH-GFP, Tdc2, and 5-HT in the Drosophila male
1232 reproductive system. A) vGlut-40XV5; B) TβH-GFP; C) 5-HT; D) vGlut-40XV5, TβH-
1233 GFP overlay; E) vGlut-40XV5, 5-HT overlay; F) TβH-GFP, 5-HT overlay. G) vGlut-
1234 40XV5; H) Tdc2; I) 5-HT; J) vGlut-40XV5, Tdc2 overlay; K) vGlut-40XV5, 5-HT overlay;
1235 L) Tdc2, 5-HT overlay. Scale bar: 200μm.

Seminal Vesicle

1236

Figure 4S1. Expression of vGlut, TβH-GFP, Tdc2, and 5-HT in the SV of the Drosophila
male reproductive system. A, G) vGlut-40XV5; B, H) TβH-GFP; C, I) 5-HT; D, J) vGlut-
40XV5, TβH-GFP overlay; E, K) vGlut-40XV5, 5-HT overlay; F, L) TβH-GFP, 5-HT
overlay;   M, S) vGlut-40XV5; N, T) Tdc2; O, U) 5-HT; P, V) vGlut-40XV5, Tdc2 overlay;
Q, W) vGlut-40XV5, 5-HT overlay; R, X) Tdc2, 5-HT overlay. Scale bars: R-50μm; X-
10μm.

Ejaculatory Duct

1243

Figure 4S2. Expression of vGlut, TβH-GFP, Tdc2, and 5-HT in the ED of the Drosophila
male reproductive system. A, G) vGlut-40XV5; B, H) TβH-GFP; C, I) 5-HT; D, J) vGlut-
40XV5, TβH-GFP overlay; E, K) vGlut-40XV5, 5-HT overlay; F, L) TβH-GFP, 5-HT
overlay;   M, S) vGlut-40XV5; N, T) Tdc2; O, U) 5-HT; P, V) vGlut-40XV5, Tdc2 overlay;
Q, W) vGlut-40XV5, 5-HT overlay; R, X) Tdc2, 5-HT overlay. Scale bars: R-50μm; X-
10μm.

Accessory Gland

Accessory Gland

1250

Figure 4S3. Expression of vGlut, TβH-GFP, Tdc2, and 5-HT in the AG of the Drosophila
male reproductive system. A, G) vGlut-40XV5; B, H) TβH-GFP; C, I) 5-HT; D, J) vGlut-
40XV5, TβH-GFP overlay; E, K) vGlut-40XV5, 5-HT overlay; F, L) TβH-GFP, 5-HT
overlay; M, S) vGlut-40XV5; N, T) Tdc2; O, U) 5-HT; P, V) vGlut-40XV5, Tdc2 overlay;
Q, W) vGlut-40XV5, 5-HT overlay; R, X) Tdc2, 5-HT overlay. Scale bars: R-50μm; X-
10μm.

1257
1258 **Figure 4S4.** Expression of vGlut, TβH-GFP, and 5-HT at the junction of the SV and
1259 AGs with the ED of the Drosophila male reproductive system. A) vGlut-40XV5; B) TβH-
1260 GFP; C) 5-HT; D) vGlut-40XV5, TβH-GFP overlay; E) vGlut-40XV5, 5-HT overlay; F)
1261 TβH-GFP, 5-HT overlay. A) vGlut-40XV5; B) Tdc2; C) 5-HT; D) vGlut-40XV5, Tdc2
1262 overlay; E) vGlut-40XV5, 5-HT overlay; F) Tdc2, 5-HT overlay. Scale bar: 50μm.

1263

Figure 5. Co-conditional expression of vGlut-40XMYC and 6XV5-vMAT in the SV of serotonergic (TRH) or octopaminergic (Tdc2) neurons of the Drosophila male reproductive system. A-D) 63X. A) vGlut-40XMYC; B) 6XV5-vMAT; C) Tdc2; D) TRH>CD8-mCherry. E-L) 63X zoom 4X. E) vGlut-40XMYC; F) 6XV5-vMAT; G) Tdc2; H) TRH>CD8-mCherry; I) vGlut-40XMYC, 6XV5-vMAT overlay; J) vGlut-40XMYC, 6XV5-vMAT, Tdc2 overlay; K) 6XV5-vMAT, Tdc2 overlay; L) Tdc2, TRH>CD8-mCherry overlay. M-P) 63X. M) vGlut-40XMYC; N) 6XV5-vMAT; O) 5-HT; P) Tdc2>CD8-mCherry. Q-X) 63X zoom 4X. Q) vGlut-40XMYC; R) 6XV5-vMAT; S) 5-HT; T) Tdc2>CD8-mCherry; U) vGlut-40XMYC, 6XV5-vMAT overlay; V) vGlut-40XMYC, 6XV5-vMAT, 5-HT overlay; W) 6XV5-vMAT, 5-HT overlay; X) Tdc2, Tdc2>CD8-mCherry overlay. Scale bars: P-50μm; X-10μm.

Ejaculatory Duct

1275

Figure 5S1. Co-conditional expression of vGlut-40XMYC and 6XV5-vMAT in the ED of
serotonergic (TRH) or octopaminergic (Tdc2) neurons of the Drosophila male
reproductive system. A-D) 63X. A) vGlut-40XMYC; B) 6XV5-vMAT; C) Tdc2; D)
TRH>CD8-mCherry. E-L) 63X zoom 4X. E) vGlut-40XMYC; F) 6XV5-vMAT; G) Tdc2;
H) TRH>CD8-mCherry; I) vGlut-40XMYC, 6XV5-vMAT overlay; J) vGlut-40XMYC,
6XV5-vMAT, Tdc2 overlay; K) 6XV5-vMAT, Tdc2 overlay; L) Tdc2, TRH>CD8-mCherry
overlay. M-P) 63X. M) vGlut-40XMYC; N) 6XV5-vMAT; O) 5-HT; P) Tdc2>CD8-
mCherry. Q-X) 63X zoom 4X. Q) vGlut-40XMYC; R) 6XV5-vMAT; S) 5-HT; T)
Tdc2>CD8-mCherry; U) vGlut-40XMYC, 6XV5-vMAT overlay; V) vGlut-40XMYC, 6XV5-
vMAT, 5-HT overlay; W) 6XV5-vMAT, 5-HT overlay; X) Tdc2, Tdc2>CD8-mCherry
overlay. Scale bars: P-50μm; X-10μm.

Accessory Gland

A vGlut-40XMYC
B 6XV5-vMAT
C Tdc2
D TRH>CD8-mCherry

E vGlut-40XMYC
F 6XV5-vMAT
G Tdc2
H TRH>CD8-mCherry

I vGlut-40XMYC
6XV5-vMAT
J vGlut-40XMYC
6XV5-vMAT
Tdc2
K 6XV5-vMAT
Tdc2
L Tdc2
TRH>CD8-mCherry

M vGlut-40XMYC
N 6XV5-vMAT
O 5-HT
P Tdc2>CD8-mCherry

Q vGlut-40XMYC
R 6XV5-vMAT
S 5-HT
T Tdc2>CD8-mCherry

U vGlut-40XMYC
6XV5-vMAT
V vGlut-40XMYC
6XV5-vMAT
5-HT
W 6XV5-vMAT
5-HT
X 5-HT
Tdc2>CD8-mCherry

1287

Figure 5S2. Co-conditional expression of vGlut-40XMYC and 6XV5-vMAT in the AG of
serotonergic (TRH) or octopaminergic (Tdc2) neurons of the Drosophila male
reproductive system. A-D) 63X. A) vGlut-40XMYC; B) 6XV5-vMAT; C) Tdc2; D)
TRH>CD8-mCherry. E-L) 63X zoom 4X. E) vGlut-40XMYC; F) 6XV5-vMAT; G) Tdc2;
H) TRH>CD8-mCherry; I) vGlut-40XMYC, 6XV5-vMAT overlay; J) vGlut-40XMYC,
6XV5-vMAT, Tdc2 overlay; K) 6XV5-vMAT, Tdc2 overlay; L) Tdc2, TRH>CD8-mCherry
overlay. M-P) 63X. M) vGlut-40XMYC; N) 6XV5-vMAT; O) 5-HT; P) Tdc2>CD8-
mCherry. Q-X) 63X zoom 4X. Q) vGlut-40XMYC; R) 6XV5-vMAT; S) 5-HT; T)
Tdc2>CD8-mCherry; U) vGlut-40XMYC, 6XV5-vMAT overlay; V) vGlut-40XMYC, 6XV5-
vMAT, 5-HT overlay; W) 6XV5-vMAT, 5-HT overlay; X) Tdc2, Tdc2>CD8-mCherry
overlay. Scale bars: P-50μm; X-10μm.

1299
Figure 5S3. Co-conditional expression of vGlut-40XMYC and 6XV5-vMAT in
serotonergic neurons of the Drosophila ED using expansion microscopy. A, D) 6XV5-
vMAT; B, E) vGlut-40XMYC; C, F) overlay. A-C) 6XV5-vMAT and vGlut-40XMYC
expression above the threshold intensity values. D-F) Complete 6XV5-vMAT and vGlut-
40XMYC expression. G) Mander's plot of 6XV5-vMAT and vGlut-40XMYC. Vertical line
indicates intensity threshold value for 6XV5-vMAT. Horizontal line indicates intensity
threshold value for vGlut-40XMYC. H) Mander's coefficients. Grayscale signal in A-C)
denotes the CD8-mCherry plasma membrane marker. Scale bars: 500nm. Scale bars
have been corrected by 10X to reflect the empirically determined 10.5X expansion
factor.

Octopamine/Glutamate neurons

1310

Figure 5S4. Co-conditional expression of vGlut-40XMYC and 6XV5-vMAT in
octopaminergic neurons of the Drosophila ED using expansion microscopy. A, D)
6XV5-vMAT; B, E) vGlut-40XMYC; C, F) overlay. A-C) 6XV5-vMAT and vGlut-40XMYC
expression above the threshold intensity values. D-F) Complete 6XV5-vMAT and vGlut-
40XMYC expression. G) Mander's plot of 6XV5-vMAT and vGlut-40XMYC. Vertical line
indicates intensity threshold value for 6XV5-vMAT. Horizontal line indicates intensity
threshold value for vGlut-40XMYC. H) Mander's coefficients. Grayscale signal in A-C)
denotes the CD8-mCherry plasma membrane marker. Scale bars: 500nm. Scale bars
have been corrected by 10X to reflect the empirically determined 10.5X expansion
factor.

1321

Figure 5S5. Nuclei of unexpanded and expanded EDs. A) DAPI-stained nuclei from an
unexpanded ED. B) DAPI-stained nuclei of an expanded ED. C) Bar graph plot of
nuclei area (mean ± SE-black bars). For the experiment shown the calculated
expansion factor (post-ExM nuclear area/pre-ExM nuclear area) was 10.47, p-value <
0.00001. Fij/ImageJi open-source software was used to segment images and calculate
nuclear area. Scale bars: 25um.

1328

Figure 6. Expression of the synaptic vesicle marker Synapsin in combination with
vGlut-40XMYC and 6XV5-vMAT in the SV, ED, and AG of the Drosophila male
reproductive system. A-I) SV. A) vGlut-40XMYC; B) 6XV5-vMAT; C) Synapsin; D)
vGlut-40XMYC; E) 6XV5-vMAT; F) Synapsin; G) vGlut-40XMYC, 6XV5-vMAT overlay;
H) vGlut-40XMYC, Synapsin overlay; I) 6XV5-vMAT, Synapsin overlay. A'-I') ED. A')
vGlut-40XMYC; B') 6XV5-vMAT; C') Synapsin; D') vGlut-40XMYC; E') 6XV5-vMAT; F')
Synapsin; G') vGlut-40XMYC, 6XV5-vMAT overlay; H') vGlut-40XMYC, Synapsin
overlay; I') 6XV5-vMAT, Synapsin overlay. A''-I'') AG. A'') vGlut-40XMYC; B'') 6XV5-
vMAT; C'') Synapsin; D'') vGlut-40XMYC; E'') 6XV5-vMAT; F'') Synapsin; G'') vGlut-
40XMYC, 6XV5-vMAT overlay; H'') vGlut-40XMYC, Synapsin overlay; I'') 6XV5-vMAT,
Synapsin overlay. Scale bars: C''-50μm; I''-10μm.

1340

Figure 6S1. Expression of the active zone marker Brp in combination with vGlut-
40XMYC and 6XV5-vMAT in the SV, ED, and AG of the Drosophila male reproductive
system. A-H) SV. A) vGlut-40XMYC; B) 6XV5-vMAT; C) Brp; D) 6XV5-vMAT, Brp
overlay; E) vGlut-40XMYC; F) 6XV5-vMAT; G) Brp; H) 6XV5-vMAT, Brp overlay. I-P)
ED. I) vGlut-40XMYC; J) 6XV5-vMAT; K) Brp; L) 6XV5-vMAT, Brp; M) vGlut-40XMYC;
N) 6XV5-vMAT; O) Brp; P) 6XV5-vMAT, Brp overlay.   I-P) AG. Q) vGlut-40XMYC; R)
6XV5-vMAT; S) Brp; T) 6XV5-vMAT, Brp; U) vGlut-40XMYC; V) 6XV5-vMAT; W) Brp; X)
6XV5-vMAT, Brp overlay. Scale bars: P-50µm; X-10µm.
1349

1350
Figure 6S2. Expression of the post-synaptic density marker Dlg in combination with
vGlut-40XMYC and 6XV5-vMAT in the SV, ED, and AG of the Drosophila male
reproductive system. A-D) Complete reproductive system. A) vGlut-40XMYC; B) 6XV5-

vMAT; C) Dlg; D) 6XV5-vMAT, Dlg overlay; E-L) SV. E) vGlut-40XMYC; F) 6XV5-vMAT;
G) Dlg; H) 6XV5-vMAT, Dlg overlay; I) vGlut-40XMYC; J) 6XV5-vMAT; G) Dlg; H) 6XV5-
vMAT, Dlg overlay. I-P) ED. I) vGlut-40XMYC; K) 6XV5-vMAT; L) Dlg; M-P) ED muscle
surface. M) vGlut-40XMYC; N) 6XV5-vMAT; O) Dlg; P) 6XV5-vMAT, Dlg overlay Q) Dlg
in the ED epithelial cells. R-U) AG muscle surface. R) vGlut-40XMYC; S) 6XV5-vMAT;
T) Dlg; U) 6XV5-vMAT, Dlg overlay; V) Dlg in AG epithelial cells. Scale bars: D-200μm;
H-50μm; V-10μm.

Seminal vesicle

1361

Figure 6S3. Expression of the large core dense vesicle marker IA2-GFP in combination
with vGlut-40XMYC and 6XV5-vMAT in the SV of the Drosophila male reproductive
system. A-C) Complete reproductive system. A) vGlut-40XMYC; B) 6XV5-vMAT; C)
IA2-GFP; D-O) SV. D, J) vGlut-40XMYC; E, K) 6XV5-vMAT; F, L) IA2-GFP; G, M)
vGlut-40XMYC, 6XV5-vMAT overlay; H, N) vGlut-40XMYC, IA2-GFP overlay; I, O)
6XV5-vMAT, IA2 overlay. Scale bars: C-200μm; F-50μm; O-10μm.

1368

Figure 6S4. Expression of the large core dense vesicle marker IA2-GFP in combination
with vGlut-40XMYC and 6XV5-vMAT in the ED and AGs of the Drosophila male
reproductive system. A-L) ED. A, G) vGlut-40XMYC; B, H) 6XV5-vMAT; C, I) IA2-GFP;
D, J) vGlut-40XMYC, 6XV5-vMAT overlay; E, K) vGlut-40XMYC, IA2-GFP overlay; F, L)
6XV5-vMAT, IA2 overlay. M-X) AG. M, S) vGlut-40XMYC; N, T) 6XV5-vMAT; O, U) IA2-
GFP; P, V) vGlut-40XMYC, 6XV5-vMAT overlay; Q, W) vGlut-40XMYC, IA2-GFP
overlay; R, X) 6XV5-vMAT, IA2 overlay. Scale bars: R-50μm; X-10μm.

1376
1377 **Figure 6S5.** Expression of vGlut-40XMYC, 6XV5-vMAT, 7XMYC-vAChT, and 9XV5-
1378 vGAT in the SV, ED, and AG of the Drosophila male reproductive system. A-D) SV. A)
1379 vGlut-40XV5; B) 6XV5-vMAT; C) 7XMYC-vAChT; D) 9XV5-vGAT. E-H) ED. E) vGlut-
1380 40XV5; F) 6XV5-vMAT; G) 7XMYC-vAChT; H) 9XV5-vGAT. I-L) AG. I) vGlut-40XV5; J)
1381 6XV5-vMAT; K) 7XMYC-vAChT; L) 9XV5-vGAT. Scale bar: 50μm.

1382
1383   **Figure 7.** Glutamate receptor GAL4 expression patterns in Drosophila male
1384   reproductive system. A) GluRIIB; B) GluRIIC. Scale bar: 200μm.

1385
1386   **Figure 7S1.** OA receptor GAL4 expression patterns in Drosophila male reproductive
1387   system. A) OAMB; B) OAα2R; C) Oct-TyrR; D) OAβ1R; E) OAβ2R; F) OAβ3R. Scale
1388   bar: 200μm.

1389
1390 **Figure 7S2.** 5-HT receptor GAL4 expression patterns in Drosophila male reproductive
1391 system. A) 5-HT1A; B) 5-HT2A; C) 5-HT1B; D) 5-HT2B; E) 5-HT7. Scale bar: 200μm.

1392

1393   **Figure 8.** GluRIIA-GFP expression in the SV, ED, and AG of the Drosophila male
1394   reproductive system. A-F) SV. A, D) vGlut-40XV5; B,E) GluRIIA-GFP; C, F) vGlut-
1395   40XV5, GluRIIA-GFP overlay. G-L) ED. G, J) vGlut-40XV5; H, K) GluRIIA-GFP; I, L)
1396   vGlut-40XV5, GluRIIA-GFP overlay. M-R) AG. M, P) vGlut-40XV5; N, Q) GluRIIA-GFP;
1397   O, R) vGlut-40XV5, GluRIIA-GFP overlay. Scale bars: O-50μm; R-10μm.

1398

Figure 8S1. OAMB expression in the Drosophila male reproductive system. A-C)
Complete male reproductive system. A) OAMB-10XV5; B) Syn; C) OAMB-10XV5, Syn
overlay. D-I) SV. D, G) OAMB-10XV5; E, H) Syn; F, I) OAMB-10XV5, Syn overlay. J-R)
ED. J-L) muscle surface. J) OAMB-10XV5; K) Syn; L) OAMB-10XV5, Syn overlay. M-
O) muscles and epithelial layers. M) OAMB-10XV5; N) Syn; O) OAMB-10XV5, Syn
overlay. P-R) epithelial layer. P) OAMB-10XV5; Q) Syn; R) OAMB-10XV5, Syn overlay.
Scale bars: C-200μm; F, O-50μm; I, L, R-10μm.

Accessory Gland

Ejaculatory Bulb

1406

Figure 8S2. OAMB expression in the Drosophila male reproductive system. A-C) AG.
A) OAMB-10XV5; B) Syn; C) OAMB-10XV5, Syn overlay. D-F) Epithelial layer of AG.
D) OAMB-10XV5; E) Syn; F) OAMB-10XV5, Syn overlay. G-I) Muscle layer of AG. G)
OAMB-10XV5; H) Syn; I) OAMB-10XV5, Syn overlay. J-L) Ejaculatory bulb. J) OAMB-
10XV5; K) Syn; L) OAMB-10XV5, Syn overlay. M-O) muscle layer of ejaculatory bulb.
M) OAMB-10XV5; N) Syn; O) OAMB-10XV5, Syn overlay. P-Q) Epithelial layer of
ejaculatory bulb. P) OAMB-10XV5; Q) Syn; R) OAMB-10XV5, Syn overlay. Scale bars:
C, L-50μm; F, I, R-10μm.

1415

Figure 8S3. OAα2R-20XV5 expression in the Drosophila male reproductive system. A-
C) Complete male reproductive system. A) OAα2R-20XV5; B) Syn; C) OAα2R-20XV5,
Syn overlay. D-I) SV. D, G) OAα2R-20XV5; E, H) Syn; F, I) OAα2R-20XV5, Syn
overlay. Cross-section of SV. J) OAα2R-20XV5: K) Syn; L) OAα2R-20XV5, Syn
overlay. M-R) AG. M, P) OAα2R-20XV5; N, Q) Syn; O, R) OAα2R-20XV5, Syn overlay.
Scale bars: C-200µm; O-50µm; R-10µm.

1422

1423   **Figure 8S4.** OAα2R-20XV5 expression in the Drosophila male reproductive system. A-
1424   C) SV/ED junction. A, D) OAα2R-20XV5; B, E) Syn; C, F) OAα2R-20XV5, Syn overlay.
1425   G-L) Proximal anterior ED. G, J) OAα2R-20XV5; H, K) Syn; I, L) OAα2R-20XV5, Syn
1426   overlay. M-U) Distal anterior ED. M, P, S) OAα2R-20XV5; N, Q, T) Syn; O, R, U)
1427   OAα2R-20XV5, Syn overlay. Scale bars: C, I, O-50μm; F-25μm; L, U-10μm.

Seminal Vesicle

Ejaculatory Duct

1428

Figure 8S5. OAβ2R-40XV5 expression in the Drosophila male reproductive system. A-
C) Complete male reproductive system. A) OAβ2R-40XV5; B) Syn; C) OAβ2R-40XV5,
Syn overlay. D-I) SV. D, G,) OAβ2R; E, H) Syn; F, I) OAβ2R-40XV5, Syn overlay. J-L)
Cross section of SV. J) OAβ2R-40XV5; K) Syn; L) OAβ2R-40XV5, Syn overlay. M-R)
ED. M, P) OAβ2R; N, Q) Syn; O, R) OAβ2R-40XV5, Syn overlay. Scale bars: C-
200μm; O-50μm; R-10μm.

1435

Figure 8S6. OAβ2R-40XV5 expression in the Drosophila male reproductive system.

A-C) Junction of SV and ED. A) OAβ2R-40XV5; B) Syn; C) OAβ2R-40XV5, Syn

overlay. D-F) Cross section of ED. D) OAβ2R-40XV5; E) Syn; F) OAβ2R-40XV5, Syn

overlay. G-I) Junction of SV and ED. G) OAβ2R-40XV5; H) Syn; I) OAβ2R-40XV5, Syn

overlay. J-L) Distal SV. J) OAβ2R-40XV5; E) Syn; F) OAβ2R-40XV5, Syn overlay. M-

O) Cross section of distal SV. M) OAβ2R-40XV5; N) Syn; O) OAβ2R-40XV5, Syn

overlay. Scale bars: C, I-50μm; O-10μm.

Ejaculatory Duct

A — OAß2R-40XV5
B — Syn
C — OAß2R-40XV5 / Syn
D — OAß2R-40XV5
E — Syn
F — OAß2R-40XV5 / Syn

Accessory Gland

G — OAß2R-40XV5
H — Syn
I — OAß2R-40XV5 / Syn
J — OAß2R-40XV5
K — Syn
L — OAß2R-40XV5 / Syn

Ejaculatory Bulb

M — OAß2R-40XV5
N — Syn
O — OAß2R-40XV5 / Syn
P — OAß2R-40XV5
Q — Syn
R — OAß2R-40XV5 / Syn

1443

Figure 8S7. OAβ2R-40XV5 expression in the Drosophila male reproductive system. A-
C) Distal anterior ED. A) OAβ2R-40XV5; B) Syn; C) OAβ2R-40XV5, Syn overlay. D-F)
Cross section of distal anterior ED. D) OAβ2R-40XV5; E) Syn; F) OAβ2R-40XV5, Syn
overlay. G-L) AG. G, J) OAβ2R-40XV5; H, K) Syn; I, L) OAβ2R-40XV5, Syn overlay. M-
O) Ejaculatory bulb. M, O) OAβ2R-40XV5; N, Q) Syn; O, R) OAβ2R-40XV5, Syn
overlay. Scale bars: O-50μm; R-10μm.

1450

1451   **Figure 8S8.** 5-HT7-20XV5 expression in the Drosophila male reproductive system. A-
1452   C) Complete male reproductive system. A) 5-HT7-20XV5.; B) Syn; C) 5-HT7-20XV5,
1453   Syn overlay. D-I) SV. D, G) 5-HT7-20XV5.; E, H) Syn; F, I) 5-HT7-20XV5, Syn overlay.
1454   J-O) ED.   J, M) 5-HT7-20XV5.; K, N) Syn; L, O) 5-HT7-20XV5, Syn overlay. Scale bars:
1455   C-200μm; L-50μm; O-10μm.

Seminal vesicle

Seminal vesicle/ejaculatory duct junction

1456

Figure 8S9. 5-HT7-20XV5 expression in the Drosophila male reproductive system. A-I)
SV. A, D, G,) 5-HT7-20XV5.; B, E, H,) Syn; C, F, I,) 5-HT7-20XV5, Syn overlay. J-L)
Cross section of SV. J) 5-HT7-20XV5.; K) Syn; L) 5-HT7-20XV5, Syn overlay. M-O) SV
terminus. M) 5-HT7-20XV5.; N) Syn; O) 5-HT7-20XV5, Syn overlay. P-R) SV/ED
junction. P) 5-HT7-20XV5.; Q) Syn; R) 5-HT7-20XV5, Syn overlay. Scale bars: C, G-
50μm; F, O-10μm; R-25μm.

Accessory gland

Ejaculatory bulb

1463

Figure 8S10. 5-HT7-20XV5 expression in the Drosophila male reproductive system. A-
C) AG. A) 5-HT7-20XV5.; B) Syn; C) 5-HT7-20XV5, Syn overlay. D-F) Epithelial layer
of AG. D) 5-HT7-20XV5.; E) Syn; F) 5-HT7-20XV5, Syn overlay. G-I) Cross-section of
epithelial layer of AG. G) 5-HT7-20XV5.; H) Syn; I) 5-HT7-20XV5, Syn overlay. J-L)
Muscle layer of AG. J) 5-HT7-20XV5.; K) Syn; L) 5-HT7-20XV5, Syn overlay. M-O)
Ejaculatory bulb. M) 5-HT7-20XV5.; N) Syn; O) 5-HT7-20XV5, Syn overlay. Scale bars:
C, O-50μm; L-10μm.

1472
Figure 9. Low resolution images of nuclear expression of *VT019028-GAL4-DBD* in
combination with *TRH-AD* and *AbdB-AD* in Drosophila male and female adult nervous
system relative to 5-HT and Tdc2. A-D) nuclear expression using HIS2A-GFP marker.
A) male *VT019028-GAL4-DBD/TRH-AD*; B) female *VT019028-GAL4-DBD/TRH-AD*; C)
male *VT019028-GAL4-DBD/AbdB-AD*; B) female *VT019028-GAL4-DBD/AbdB-AD*. E-
H) 5-HT expression. E) male *VT019028-GAL4-DBD/*TRH-AD; F) female *VT019028-*

1479   *GAL4-DBD/TRH-AD*; G) male *VT019028-GAL4-DBD/AbdB-AD*; H) female *VT019028-*
1480   *GAL4-DBD/AbdB-AD*. Tdc2 expression. I) male *VT019028-GAL4-DBD/TRH-AD*; J)
1481   female *VT019028-GAL4-DBD/TRH-AD*; K) male *VT019028-GAL4-DBD/AbdB-AD*; L)
1482   female *VT019028-GAL4-DBD/AbdB-AD*. Scale bar: 200μm.

1483
Figure 9S1. High resolution images of nuclear expression of *VT019028-GAL4-DBD* in
combination with *TRH-AD* in posterior ventral nerve cord of Drosophila male adult
nervous system relative to 5-HT and Tdc2. A-E) Dorsal view of complete stack of
confocal images. A) His2A-GFP; B) 5-HT; C) Tdc2; D) His2A-GFP, 5-HT overlay; E)
His2A-GFP, Tdc2 overlay. F) Ventral view of His2A-GFP/5-HT overlay of subset of
slices containing the dorsal cluster of 5-HT neurons; G-K) Dorsal view of subset of
slices containing the dorsal cluster of 5-HT neurons. G) His2A-GFP; H) 5-HT; I) Tdc2;
J) His2A-GFP, 5-HT overlay; K) His2A-GFP, Tdc2 overlay. L-P) Dorsal view of subset of
slices containing the ventral cluster of 5-HT neurons. L) His2A-GFP; M) 5-HT; N) Tdc2;
O) His2A-GFP, 5-HT overlay; P) His2A-GFP, Tdc2 overlay. Scale bar: 50μm.

Male VT019028-DBD/AbdB-AD

Figure 9S2. High resolution images of nuclear expression of *VT019028-GAL4-DBD* in combination with *AbdB-AD* in posterior ventral nerve cord of Drosophila male adult nervous system relative to 5-HT and Tdc2. A-E) Dorsal view of complete stack of confocal images. A) His2A-GFP; B) 5-HT; C) Tdc2; D) His2A-GFP, 5-HT overlay; E) His2A-GFP, Tdc2 overlay. F-J) Dorsal view of subset of slices containing the dorsal cluster of 5-HT neurons. F) His2A-GFP; G) 5-HT; H) Tdc2; I) His2A-GFP, 5-HT overlay; J) His2A-GFP, Tdc2 overlay. K-O) Dorsal view of subset of slices containing the ventral cluster of 5-HT neurons. K) His2A-GFP; L) 5-HT; M) Tdc2; N) His2A-GFP, 5-HT overlay; O) His2A-GFP, Tdc2 overlay. Scale bar: 50μm.

1505
Figure 10. Comparison of mating duration between genotypes. Bar plots show the
mean ± SD for each group with individual data points overlaid on each bar. Statistical
comparisons were made using unpaired two-tailed t tests where appropriate. Asterisks
above indicate significance levels: p < 0.001 (**), p < 0.0001 (***), ns = not significant.
(A) Columns from left to right: TRH ∩ AbdB > BONT-C: n= 10, mean= 19.11, SD=4.251.
Tdc2 ∩ AbdB > BONT-C: n=14, mean=21.80, SD= 3.155. AbdB ∩ VT019028 > BONT-
C: n=12, mean=24.75, SD=4.247. TRH ∩ VT019028 > BONT-C: n=11, mean=28.74,
SD=2.155. AbdB ; BONT-C: n=8, mean=27.07, SD=4.907. BONT-C: n=7, mean=28.75,
SD=4.809. WT Canton-S: n=9, mean=26.91, SD=3.473.

1515
1516 **Figure 10S1.** Expression of ProtB-GFP fluorescent marker of sperm nuclei in male
1517 reproductive system of control and *TRH-AD*/*VT019028-GAL4-DBD* neuron-silenced
1518 flies interrupted mid-mating. A) control. B) *TRH-AD*/*VT019028-GAL4-DBD* neurons
1519 silenced with BONT-C. Prot-GFP-Protamine GFP. SV-SV, ED-ED. Scale bar: 75μm.

1520
Figure 10S2. Comparison of male drosophila fecundity between genotypes. Bar plots
show the mean ± SD for each group with individual data points overlaid on each bar.
Figure 10S1A males were aged 5 days before copulation. Figure 10S1B males were
aged 30 days before copulation. (A) Columns from left to right: TRH ∩ VT019028 →
vGlut-: n=34, mean=173.2, SD=24.22. AbdB ∩ TRH, Tdc2 → vGlut-: n=20,
mean=175.3, SD=22.92. TRH ∩ VT019028: n=32, mean=149.0, SD=26.10. TRH: n=28,
mean=172.8, SD=21.11. VT019028: n=25, mean=148.6, SD=20.81. (B) TRH
∩ VT019028 → vGlut-: n=14, mean=133.9, SD=37.64. TRH ∩ VT019028: n=19,
mean=123.7, SD, 25.02. TRH: n=13, mean=106.3, SD=41.93. VT019028: n=9,
mean=131.0, SD=38.88.

1531
1532 **Figure 10S3.** vGlut expression in AGs of control and vGlut conditional mutant. A-C)
1533 control male. A) vGlut; B) 6XmCherry; C) overlay. D-F) experimental male conditionally
1534 silenced for vGlut in *TRH-AD/VT019028-GAL4* neurons. Scale bar: 50μm.

1535
1536 **Video 1.** Male reproductive system of a ProtB-GFP control. Sperm with green
1537 fluorescent nuclei are visible in the ejaculatory duct of a male subjected to a mid-mating
1538 interruption.

1539
1540 **Video 2.** Male reproductive system of a ProtB-GFP experimental male in which a subset
1541 of the *TRH-AD/VT019028-GAL4-DBD* subset of serotonergic neurons innervating the
1542 male reproductive system have been silenced. Sperm with green fluorescent nuclei are
1543 restricted to the seminal vesicle and never enter the ejaculatory duct from a male
1544 subjected to a mid-mating interruption.

1545
1546 **Video 3.** GCAMP8m Ca++ imaging of an ejaculatory duct experiencing spontaneous
1547 peristaltic waves of muscle contractions.

1548
1549 **Video 4.** GCAMP8m Ca++ imaging of the SV/ED junction showing the initiation site of
1550 the spontaneous peristaltic waves.

1551
1552 **Video 5.** GCAMP8m Ca++ imaging of the SV/ED junction showing the subtle variation
1553 in the initiation site of spontaneous peristaltic waves as compared to Video 4.

1554
1555   **Video 6.** GCAMP8m Ca++ imaging of the SV showing spontaneous uncoordinated
1556   muscle activity.
1557
1558   **Video 7.** GCAMP8m Ca++ imaging of an AG showing spontaneous uncoordinated
1559   muscle activity.
1560
1561   **Video 8.** GCAMP8m Ca++ imaging of a testes showing spontaneous uncoordinated
1562   muscle activity.
1563
1564   **Video 9.** GCAMP8m Ca++ imaging of the AG showing strong coordinated muscle
1565   activity.
1566
1567
1568

Genotype	Male Fertile	Female Fertile
yw, *TRH-GAL4DBD/AbdB-AD, UAS-BONT-C*	No	No
yw, *Tdc2-GAL4DBD/AbdB-AD, UAS-BONT-C*	Yes	No
yw, *VT019028-GAL4-DBD/AbdB-AD, UAS-BONT-C*	No	Yes
yw, *TRH-AD*; *UAS-BONT-C/VT019028-GAL4-DBD*	No	Yes
yw, *AbdB-AD/UAS-BONT-C*	Yes	Yes
yw, *UAS-BONT-C*	Yes	Yes
Canton-S wildtype	Yes	Yes

1569
1570   **Table 1.** Male and female fertility phenotypes.
1571
1572
1573   **Complete Genotypes**
1574
1575   **Figure 1.** B) *yw*.
1576
Figure 2. A) *yw*: *Tdc2-GAL4/+*: *UAS-CD8-mCherry/+*; B) *yw*: *TRH-GAL4/+*: *UAS-CD8-*
mCherry/+; C) *yw*: *vGlut-GAL4/+*: *UAS-CD8-mCherry/+*; D) *yw*: *vGlut-GAL4-DBD/+*;
Tdc2-AD/UAS-CD8-mCherry/+; E) *yw*; *vGlut-AD/+*; *Tdc2-GAL4-DBD, UAS-CD8-*
mCherry; F) *yw*; *TRH-AD/vGlut-GAL4-DBD*; *UAS-CD8-mCherry/+*; G) *yw*; *vGlut-AD/+*;
TRH-GAL4-DBD/UAS-CD8-mCherry; H) *yw*; *Tdc2-AD/TRH-GAL4-DBD, UAS-CD8-*
mCherry.
1583
1584   **Figure 3.** *yw*; *B3RT-Tdc2-LexA/20XUAS-6XGFP*; *TRH-GAL4/13XLexAop-6XmCherry*.
1585
1586   **Figure 3S1.** *yw*; *B3RT-vGlut-LexA/20XUAS-6XGFP*; TRH-GAL4/*13XLexAop-*
1587   *6XmCherry*.

1588

Figure 3S2. *yw*; *B3RT-vGlut-LexA/Tdc2-GAL4*; TRH-GAL4/*20XUAS-6XGFP/13XLexAop-6XmCherry*.

Figure 3S3. A-C) *yw*; *B3RT-vGlut-LexA/20XUAS-6XGFP*; fru-GAL4/*13XLexAop-6XmCherry*. D) *yw*; *Tdc2-AD/dsx-GAL4-DBD, UAS-CD8-mCherry*. E) *yw*; *TRH-AD/+*; *dsx-GAL4-DBD, UAS-CD8-mCherry/+*.

Figure 4. A-F) *w, TBH-GFP; vGlut-40XV5/vGlut-40XV5*. J-L) *yw; vGlut-40XV5/vGlut-40XV5*.

Figure 4S1. A-L) *w, TBH-GFP; vGlut-40XV5/vGlut-40XV5*. M-X) yw; *vGlut-40XV5/vGlut-40XV5*.

Figure 4S2. A-L) *w, TBH-GFP; vGlut-40XV5/vGlut-40XV5*. M-X) yw; *vGlut-40XV5/vGlut-40XV5*.

Figure 4S3. A-L) *w, TBH-GFP; vGlut-40XV5/vGlut-40XV5*. M-X) yw; *vGlut-40XV5/vGlut-40XV5*.

Figure 4S4. A-F) *w, TBH-GFP; vGlut-40XV5/vGlut-40XV5*.

Figure 5. A-L) *yw; B2RT-STOP-B2RT-40XMYC, RSRT-STOP-RSRT-6XV5-vMAT/+; TRH-GAL4, UAS-CD8-mCherry, UAS-B2, UAS-R*. M-O) *yw; B2RT-STOP-B2RT-40XMYC, RSRT-STOP-RSRT-6XV5-vMAT/Tdc2-GAL4; UAS-CD8-mCherry, UAS-B2, UAS-R/+*.

Figure 5S1. A-L) *yw; B2RT-STOP-B2RT-40XMYC, RSRT-STOP-RSRT-6XV5-vMAT/+; TRH-GAL4, UAS-CD8-mCherry, UAS-B2, UAS-R*. M-O) *yw; B2RT-STOP-B2RT-40XMYC, RSRT-STOP-RSRT-6XV5-vMAT/Tdc2-GAL4; UAS-CD8-mCherry, UAS-B2, UAS-R/+*.

Figure 5S2. A-L) *yw; B2RT-STOP-B2RT-40XMYC, RSRT-STOP-RSRT-6XV5-vMAT/+; TRH-GAL4, UAS-CD8-mCherry, UAS-B2, UAS-R*. M-O) *yw; B2RT-STOP-B2RT-40XMYC, RSRT-STOP-RSRT-6XV5-vMAT/Tdc2-GAL4; UAS-CD8-mCherry, UAS-B2, UAS-R/+*.

Figure 5S3. A-I) *yw; B2RT-STOP-B2RT-40XMYC, RSRT-STOP-RSRT-6XV5-vMAT/+; TRH-GAL4, UAS-CD8-mCherry, UAS-B2, UAS-R*.

Figure 5S4. A-I) *yw; B2RT-STOP-B2RT-40XMYC, RSRT-STOP-RSRT-6XV5-vMAT/Tdc2-GAL4; UAS-CD8-mCherry, UAS-B2, UAS-R/+*.

Figure 5S5. A-B) *yw*.

Figure 6. A-I) *yw; vGlut-40XMYC, 6XV5-vMAT/vGlut-40XMYC, 6XV5-vMAT*. A'-I') *yw; vGlut-40XMYC, 6XV5-vMAT/vGlut-40XMYC, 6XV5-vMAT*. A''-I'') *yw; vGlut-40XMYC, 6XV5-vMAT/vGlut-40XMYC, 6XV5-vMAT*.

Figure 6S1. A-X) *yw; vGlut-40XMYC, 6XV5-vMAT/vGlut-40XMYC, 6XV5-vMAT*.

Figure 6S2. A-V) *yw; vGlut-40XMYC, 6XV5-vMAT/vGlut-40XMYC, 6XV5-vMAT*.

Figure 6S3. A-O) *yw; vGlut-40XMYC, 6XV5-vMAT/IA2-GFP*.
Figure 6S4. A-O) *yw; vGlut-40XMYC, 6XV5-vMAT/IA2-GFP*.

Figure 6S5. A, E, I) *yw; vGlut-40X5/vGlut-40XV5*. B, F, J) *yw; 6XV5-vMAT/6XV5-vMAT*. C, G, K) *yw; 7XMYC-vAChT/7XMYC-vAChT*. D, H, L) *yw; 9XV5-vGAT/9XV5-vGAT*.

Figure 7. A) *yw; GluRIIB-GAL4/+; UAS-CD8-mCherry*. B) *yw; GluRIIC-GAL4/+; UAS-CD8-mCherry*.

Figure 7S1. A) *yw; OAMB-GAL4/UAS-CD8-mCherry*. B) *yw; OAα2R-GAL4/UAS-CD8-mCherry*. C) *yw; Oct-TyrR-GAL4/UAS-CD8-mCherry*. D) *yw; OAβ1R-GAL4/UAS-CD8-mCherry*. E) *yw; OAβ2R-GAL4/UAS-CD8-mCherry*. F) *yw; OAβ3R-GAL4/UAS-CD8-mCherry*.

Figure 7S2. A) *yw; 5-HT71A-GAL4/+; UAS-CD8-mCherry/+*. B) *yw; 5-HT71B-GAL4/+; UAS-CD8-mCherry/+*. C) *yw; 5-HT2A-GAL4/UAS-CD8-mCherry*. D) *yw; 5-HT2B-GAL4/UAS-CD8-mCherry*. E) *yw; 5-HT7-GAL4/UAS-CD8-mCherry*.

Figure 8. A-R) *yw; vGlut-40XV5/GluRIIA-GFP*.

Figure 8S1. A-R) *yw; OAMB-10XV5/OAMB-10XV5*.

Figure 8S2. A-R) *yw; OAMB-10XV5/OAMB-10XV5*.

Figure 8S3. A-R) *yw; OAα2R-20XV5/OAα2R-20XV5*.

Figure 8S4. A-U) *yw; OAα2R-20XV5/OAα2R-20XV5*.

Figure 8S5. A-R) *yw; OAβ2R-40XV5/OAβ2R-40XV5*.

Figure 8S6. A-O) *yw; OAβ2R-40XV5/OAβ2R-40XV5*.

Figure 8S7. A-R) *yw; OAβ2R-40XV5/OAβ2R-40XV5*.

Figure 8S8. A-O) *yw; 5-HT7-20XV5/5-HT7-20XV5*.

Figure 8S9. A-R) *yw; 5-HT7-20XV5/5-HT7-20XV5*.

Figure 8S10. A-O) *yw*; *5-HT7-20XV5/5-HT7-20XV5*.

Figure 9. A, E, I) male *yw*; *UAS-H2A-GFP/TRH-AD*; *VT019028-GAL4-DBD/+*. B, F, J) female *yw*; *UAS-H2A-GFP/TRH-AD*; *VT019028-GAL4-DBD/+*. C, G, K) male *yw*; *UAS-H2A-GFP/+*; *VT019028-GAL4-DBD/AbdB-AD*. D, H, L) female *yw*; *UAS-H2A-GFP/+*; *VT019028-GAL4-DBD/AbdB-AD*.

Figure 9S1. A-P) male *yw*; *UAS-H2A-GFP/TRH-AD*; *VT019028-GAL4-DBD/+*.

Figure 9S2. A-O) male *yw*; *UAS-H2A-GFP/+*; *VT019028-GAL4-DBD/AbdB-AD*.

Figure 9S3. A) male *yw*; *UAS-H2A-GFP/TRH-AD*; *VT019028-GAL4-DBD/+*. B) male *yw*; *UAS-H2A-GFP/+*; *VT019028-GAL4-DBD/AbdB-AD*.

Figure 10. A) column 1-*yw*; *TRH-GAL4-DBD/AbdB-AD, UAS-BONT-C/+*; column 2-*yw*; *Tdc2-GAL4-DBD/ AbdB-AD, UAS-BONT-C*. column 3-*yw*; *AbdB-AD, UAS-BONT-C/VT019028-GAL4-DBD*. column 4-*yw*; *TRH-AD/+*; *VT019028-GAL4-DBD, UAS-BONT-C*. column 5-*yw*; *AbdB-AD, UAS-BONT-C*. column 6-*UAS-BONT-C*. column 7-wildtype *Canton-S*. B) column 1-*yw*; *AbdB-AD, UAS-BONT-C/VT019028-GAL4-DBD*. column 2-*yw*; *AbdB-AD, UAS-BONT-C*. column 3-*UAS-BONT-C*. column 1-wildtype *Canton-S*. C) column 2-*yw*; *TRH-AD/+*; *VT019028-GAL4-DBD, UAS-BONT-C*. column 3-*yw*; *AbdB-AD, UAS-BONT-C*. column 4-*UAS-BONT-C*. column 7-wildtype *Canton-S*. D) column 1-*yw*; *TRH-GAL4-DBD/AbdB-AD, UAS-BONT-C/+*; column 2-*yw*; *AbdB-AD, UAS-BONT-C*. column 3-*UAS-BONT-C*. column 4-wildtype *Canton-S*. E) column 1-*yw*; *Tdc2-GAL4-DBD/ AbdB-AD, UAS-BONT-C*. column 2-*yw*; *AbdB-AD, UAS-BONT-C*. column 3-*UAS-BONT-C*. column 4-wildtype *Canton-S*.

Figure 10S1. A) *yw*; *ProtB-GFP/+*; B) *yw*; *ProtB-GFP/+*; *VT019028-GAL4-DBD, UAS-BONT-C*.

Figure 10S2. A) column 1-*yw*; *B3RT-vGlut-LexA/TRH-AD*, vGlut^{SS1}; *VT019028-GAL4-DBD/UAS-B3*. column 2-*yw*; *B3RT-vGlut-LexA/vGlut^{SS1}*; *TRH-GAL4-DBD, Tdc2-GAL4-DBD/AbdB-AD, UAS-B3*. column 3-*yw*; *B3RT-vGlut-LexA/TRH-AD*; *VT019028-GAL4-DBD/UAS-B3*. column 4-*yw*; *B3RT-vGlut-LexA/TRH-AD*, vGlut^{SS1}; *UAS-B3/+*. column 5-*yw*; *B3RT-vGlut-LexA/vGlut^{SS1}*; *VT019028-GAL4-DBD/UAS-B3*. B) column 1-*yw*; *B3RT-vGlut-LexA/TRH-AD*, vGlut^{SS1}; *VT019028-GAL4-DBD/UAS-B3*. column 2- *yw*; *B3RT-vGlut-LexA/TRH-AD*; *VT019028-GAL4-DBD/UAS-B3*. column 3- *yw*; *B3RT-vGlut-LexA/TRH-AD*, vGlut^{SS1}; *UAS-B3/+*. column 4-*yw*; *B3RT-vGlut-LexA/vGlut^{SS1}*; *VT019028-GAL4-DBD/UAS-B3*.

Figure 10S3. A-F) *yw*; *B3RT-vGlut-LexA/TRH-AD*, vGlut^{SS1}; *VT019028-GAL4-DBD/UAS-B3, LexAop-6XmCherry*.

Video 1. *yw*; *ProtB-GFP/+*.

Video 2. *yw*; *ProtB-GFP/+*; *VT019028-GAL4-DBD, UAS-BONT-C*.

1725
1726 **Videos 3-9.** *yw*: *MHC-GCAMP8m*.
1727
1728
1729 **References**
1730

Abbe, E., 1873 Beitrage zur Theorie des microskops und der mikroskopischen
Wahrnehmung. Anatomie IX**:** 413-468.
Acebes, A., Y. Grosjean, C. Everaerts and J. F. Ferveur, 2004 Cholinergic control of
synchronized seminal emissions in Drosophila. Curr Biol 14**:** 704-710.
Allen, J. M., T. E. Adrian, K. Tatemoto, J. M. Polak, J. Hughes *et al.*, 1982 Two novel
related peptides, neuropeptide Y (NPY) and peptide YY (PYY) inhibit the
contraction of the electrically stimulated mouse vas deferens. Neuropeptides 3**:**
71-77.
Alwaal, A., B. N. Breyer and T. F. Lue, 2015 Normal male sexual function: emphasis on
orgasm and ejaculation. Fertil Steril 104**:** 1051-1060.
Bairati, A., 1968 [The structure and ultrastructure of the male genital apparatus of the
Drosophila melanogaster Meig. 2. The genital duct and accessory glands]. Italian
Journal of Zoology 2**:** 105-182.
Baker, C. A., X. J. Guan, M. Choi and M. Murthy, 2024 The role of fruitless in specifying
courtship behaviors across divergent Drosophila species. Sci Adv 10**:** eadk1273.
Bakshinska, D., W. Y. Liu, R. Schultz, R. S. Stowers, A. Hoagland *et al.*, 2025 Synapse-
specific catecholaminergic modulation of neuronal glutamate release. Proc Natl
Acad Sci U S A 122**:** e2420496121.
Banerjee, S., S. Vernon, W. Jiao, B. J. Choi, E. Ruchti *et al.*, 2021 Miniature
neurotransmission is required to maintain Drosophila synaptic structures during
ageing. Nat Commun 12**:** 4399.
Beckers, C. J., A. Mrestani, F. Komma and S. Dannhauser, 2024 Versatile Endogenous
Editing of GluRIIA in Drosophila melanogaster. Cells 13.
Beguelini, M. R., C. C. I. Puga, F. F. Martins, A. H. S. Betoli, S. R. Taboga *et al.*, 2013
Morphological Variation of Primary Reproductive Structures in Males of Five
Families of Neotropical Bats. Anatomical Record-Advances in Integrative
Anatomy and Evolutionary Biology 296**:** 156-167.
Billeter, J. C., and S. F. Goodwin, 2004 Characterization of Drosophila fruitless-gal4
transgenes reveals expression in male-specific fruitless neurons and innervation
of male reproductive structures. J Comp Neurol 475**:** 270-287.
Billeter, J. C., E. J. Rideout, A. J. Dornan and S. F. Goodwin, 2006a Control of male
sexual behavior in Drosophila by the sex determination pathway. Curr Biol 16**:**
R766-776.
Billeter, J. C., A. Villella, J. B. Allendorfer, A. J. Dornan, M. Richardson *et al.*, 2006b
Isoform-specific control of male neuronal differentiation and behavior in
Drosophila by the fruitless gene. Curr Biol 16**:** 1063-1076.
Bretman, A., C. Fricke and T. Chapman, 2009 Plastic responses of male Drosophila
melanogaster to the level of sperm competition increase male reproductive
fitness. Proc Biol Sci 276**:** 1705-1711.

Brown, G. L., and J. S. Gillespie, 1957 The output of sympathetic transmitter from the spleen of the cat. J Physiol 138**:** 81-102.

Buck, S. A., S. A. Rubin, T. Kunkhyen, C. D. Treiber, X. Xue *et al.*, 2023 Sexually dimorphic mechanisms of VGLUT-mediated protection from dopaminergic neurodegeneration. bioRxiv.

Burnstock, G., 2009 Purinergic cotransmission. Exp Physiol 94**:** 20-24.

Burnstock, G., 2014 Purinergic signalling in the reproductive system in health and disease. Purinergic Signal 10**:** 157-187.

Carro-Juarez, M., S. L. Cruz and G. Rodriguez-Manzo, 2003 Evidence for the involvement of a spinal pattern generator in the control of the genital motor pattern of ejaculation. Brain Res 975**:** 222-228.

Carter, D., M. Champney, B. Hwang and R. M. Eglen, 1995 Characterization of a postjunctional 5-HT receptor mediating relaxation of guinea-pig isolated ileum. Eur J Pharmacol 280**:** 243-250.

Castellanos, M. C., J. C. Tang and D. W. Allan, 2013 Female-biased dimorphism underlies a female-specific role for post-embryonic Ilp7 neurons in Drosophila fertility. Development 140**:** 3915-3926.

Certel, S. J., B. D. McCabe and R. S. Stowers, 2022 A conditional GABAergic synaptic vesicle marker for Drosophila. J Neurosci Methods 372**:** 109540.

Chan, A. K., and P. Y. von der Weid, 2003 5-HT decreases contractile and electrical activities in lymphatic vessels of the guinea-pig mesentery: role of 5-HT 7-receptors. Br J Pharmacol 139**:** 243-254.

Chang Chien, C. C., L. W. Hsin and M. J. Su, 2015 Activation of serotonin 5-HT(7) receptor induces coronary flow increase in isolated rat heart. Eur J Pharmacol 748**:** 68-75.

Chehensse, C., P. Facchinetti, S. Bahrami, P. Andrey, J. M. Soler *et al.*, 2017 Human spinal ejaculation generator. Ann Neurol 81**:** 35-45.

Cizmar, D., L. Roller, M. Pillerova, K. Slama and D. Zitnan, 2019 Multiple neuropeptides produced by sex-specific neurons control activity of the male accessory glands and gonoducts in the silkworm Bombyx mori. Sci Rep 9**:** 2253.

Cole, S. H., G. E. Carney, C. A. McClung, S. S. Willard, B. J. Taylor *et al.*, 2005 Two functional but noncomplementing Drosophila tyrosine decarboxylase genes: distinct roles for neural tyramine and octopamine in female fertility. J Biol Chem 280**:** 14948-14955.

Crickmore, M. A., and L. B. Vosshall, 2013 Opposing dopaminergic and GABAergic neurons control the duration and persistence of copulation in Drosophila. Cell 155**:** 881-893.

Damstra, H. G. J., B. Mohar, M. Eddison, A. Akhmanova, L. C. Kapitein *et al.*, 2023 Ten-fold Robust Expansion Microscopy. Bio Protoc 13**:** e4698.

DeLecce, T., G. S. Vance, V. Zeigler-Hill, L. L. M. Welling and T. K. Shackelford, 2025 Ejaculate Adjustment in Response to Sperm Competition Risk in Humans. Arch Sex Behav 54**:** 277-287.

Demerec, M., 1950 *Biology of Drosophila*. Wiley, New York,.

Deng, B., Q. Li, X. Liu, Y. Cao, B. Li *et al.*, 2019 Chemoconnectomics: Mapping Chemical Transmission in Drosophila. Neuron 101**:** 876-893 e874.

Deshpande, S. A., Z. Freyberg, H. O. Lawal and D. E. Krantz, 2020 Vesicular
neurotransmitter transporters in Drosophila melanogaster. Biochim Biophys Acta
Biomembr 1862**:** 183308.
Deshpande, S. A., E. W. Rohrbach, J. D. Asuncion, J. Harrigan, A. Eamani *et al.*, 2022
Regulation of Drosophila oviduct muscle contractility by octopamine. iScience 25**:**
104697.
DiAntonio, A., 2006 Glutamate receptors at the Drosophila neuromuscular junction. Int
Rev Neurobiol 75**:** 165-179.
DiAntonio, A., S. A. Petersen, M. Heckmann and C. S. Goodman, 1999 Glutamate
receptor expression regulates quantal size and quantal content at the Drosophila
neuromuscular junction. J Neurosci 19**:** 3023-3032.
Diao, F., H. Ironfield, H. Luan, F. Diao, W. C. Shropshire *et al.*, 2015 Plug-and-play
genetic access to drosophila cell types using exchangeable exon cassettes. Cell
Rep 10**:** 1410-1421.
Diao, F., D. Vasudevan, E. S. Heckscher and B. H. White, 2024 Hox gene-specific
cellular targeting using split intein Trojan exons. Proc Natl Acad Sci U S A 121**:**
e2317083121.
Dionne, H., K. L. Hibbard, A. Cavallaro, J. C. Kao and G. M. Rubin, 2018 Genetic
Reagents for Making Split-GAL4 Lines in Drosophila. Genetics 209**:** 31-35.
Drobnis, E. Z., and A. K. Nangia, 2017 Cardiovascular/Pulmonary Medications and Male
Reproduction. Adv Exp Med Biol 1034**:** 103-130.
Dwivedi, S., M. Bautista, S. Shrestha, H. Elhasasna, T. Chaphekar *et al.*, 2021
Sympathetic signaling facilitates progression of neuroendocrine prostate cancer.
Cell Death Discov 7**:** 364.
Eberhard, W. G., 1985 *Sexual selection and animal genitalia.* Harvard University Press,
Cambridge, Mass.
Fisher, Y. E., H. H. Yang, J. Isaacman-Beck, M. Xie, D. M. Gohl *et al.*, 2017 FlpStop, a
tool for conditional gene control in Drosophila. Elife 6.
Fromme, L., D. R. Yogui, M. H. Alves, A. L. J. Desbiez, M. Langeheine *et al.*, 2021
Morphology of the genital organs of male and female giant anteaters (
). Peerj 9.
Gage, M. J. G., and R. R. Baker, 1991 Ejaculate Size Varies with Sociosexual Situation
in an Insect. Ecological Entomology 16**:** 331-337.
Gao, S., S. Y. Takemura, C. Y. Ting, S. Huang, Z. Lu *et al.*, 2008 The neural substrate of
spectral preference in Drosophila. Neuron 60**:** 328-342.
Garbaczewska, M., J. C. Billeter and J. D. Levine, 2013 Drosophila melanogaster males
increase the number of sperm in their ejaculate when perceiving rival males. J
Insect Physiol 59**:** 306-310.
Gilbert, D. G., 1981 Ejaculate Esterast 6 and Initial Sperm Use by Female Drosophila
melanogaster. Journal of Insect Physiology 27**:** 641-650.
Gomes, L. F., J. P. Badke, U. Zama, H. Dolder and J. Lino-Neto, 2012 Morphology of
the male reproductive system and spermatozoa in Centris Fabricius, 1804
(Hymenoptera: Apidae, Centridini). Micron 43**:** 695-704.
Goulding, M., S. Bourane, L. Garcia-Campmany, A. Dalet and S. Koch, 2014 Inhibition
downunder: an update from the spinal cord. Curr Opin Neurobiol 26**:** 161-166.

Gowda, S. B. M., P. D. Paranjpe, O. V. Reddy, D. Thiagarajan, S. Palliyil *et al.*, 2018 GABAergic inhibition of leg motoneurons is required for normal walking behavior in freely moving Drosophila. Proc Natl Acad Sci U S A 115**:** E2115-E2124.

Guerrero, S. M., M. L. Calderón, G. R. de Pérez and M. P. R. Pinilla, 2004 Morphology of the male reproductive duct system of (Crocodylia, Alligatoridae). Annals of Anatomy-Anatomischer Anzeiger 186**:** 235-245.

Han, Y., C. Chien, P. Goel, K. He, C. Pinales *et al.*, 2022 Botulinum neurotoxin accurately separates tonic vs. phasic transmission and reveals heterosynaptic plasticity rules in Drosophila. Elife 11.

He, K., and D. Dickman, 2025 Building and modifying diverse synaptic properties: Insights from Drosophila. Curr Opin Neurobiol 92**:** 102995.

Hiramoto, A., J. Jonaitis, S. Niki, H. Kohsaka, R. D. Fetter *et al.*, 2021 Regulation of coordinated muscular relaxation in Drosophila larvae by a pattern-regulating intersegmental circuit. Nat Commun 12**:** 2943.

Honda, K., A. Miyata-Osawa and T. Takenaka, 1985 alpha 1-Adrenoceptor subtype mediating contraction of the smooth muscle in the lower urinary tract and prostate of rabbits. Naunyn Schmiedebergs Arch Pharmacol 330**:** 16-21.

Hopkins, B. R., I. Sepil, M. L. Thezenas, J. F. Craig, T. Miller *et al.*, 2019 Divergent allocation of sperm and the seminal proteome along a competition gradient in Drosophila melanogaster. Proc Natl Acad Sci U S A 116**:** 17925-17933.

Inoue, M., T. Kitazawa, J. Cao and T. Taneike, 2003 5-HT7 receptor-mediated relaxation of the oviduct in nonpregnant proestrus pigs. Eur J Pharmacol 461**:** 207-218.

Jois, S., Y. B. Chan, M. P. Fernandez and A. K. Leung, 2018 Characterization of the Sexually Dimorphic fruitless Neurons That Regulate Copulation Duration. Front Physiol 9**:** 780.

Kaufman, B. P. a. D., M., 1942 Utilization of sperm by Female Drosophila Melanogaster. The American Naturalist 76**:** 445-469.

Klagges, B. R., G. Heimbeck, T. A. Godenschwege, A. Hofbauer, G. O. Pflugfelder *et al.*, 1996 Invertebrate synapsins: a single gene codes for several isoforms in Drosophila. J Neurosci 16**:** 3154-3165.

Kreiner, T., W. Sossin and R. H. Scheller, 1986 Localization of Aplysia neurosecretory peptides to multiple populations of dense core vesicles. J Cell Biol 102**:** 769-782.

Lange, A. B., A. Kisana, J. Leyria and I. Orchard, 2023 The Male Reproductive System of the Kissing Bug, Rhodnius prolixus Stal, 1859 (Hemiptera: Reduviidae: Triatominae): Arrangements of the Muscles and the Myoactivity of the Selected Neuropeptides. Insects 14.

Lee, G., and J. C. Hall, 2001 Abnormalities of male-specific FRU protein and serotonin expression in the CNS of fruitless mutants in Drosophila. J Neurosci 21**:** 513-526.

Lee, P. T., J. Zirin, O. Kanca, W. W. Lin, K. L. Schulze *et al.*, 2018 A gene-specific T2A-GAL4 library for Drosophila. Elife 7.

Leung, E., L. K. Walsh, M. T. Pulido-Rios and R. M. Eglen, 1996 Characterization of putative 5-ht7 receptors mediating direct relaxation in Cynomolgus monkey isolated jugular vein. Br J Pharmacol 117**:** 926-930.

Li-Kroeger, D., O. Kanca, P. T. Lee, S. Cowan, M. T. Lee *et al.*, 2018 An expanded toolkit for gene tagging based on MiMIC and scarless CRISPR tagging in Drosophila. Elife 7.

1906 Liao, A., F. Cavigliasso, L. Savary and T. J. Kawecki, 2024 Effects of an
1907   entomopathogenic fungus on the reproductive potential of Drosophila males.
1908   Ecol Evol 14**:** e11242.
1909 Macartney, E. L., V. Zeender, A. Meena, A. N. De Nardo, R. Bonduriansky *et al.*, 2021
1910   Sperm depletion in relation to developmental nutrition and genotype in
1911   Drosophila melanogaster. Evolution 75**:** 2830-2841.
1912 Maciel, E. D. S., R. Zieri and S. M. de Almeida-Santos, 2024 Male genital system of
1913   Ameiva ameiva (Squamata: Teiidae). Anat Rec (Hoboken) 307**:** 3596-3605.
1914 Magnon, C., S. J. Hall, J. Lin, X. Xue, L. Gerber *et al.*, 2013 Autonomic nerve
1915   development contributes to prostate cancer progression. Science 341**:** 1236361.
1916 Majane, A. C., J. M. Cridland and D. J. Begun, 2022 Single-nucleus transcriptomes
1917   reveal evolutionary and functional properties of cell types in the Drosophila
1918   accessory gland. Genetics 220.
1919 Manier, M. K., J. M. Belote, K. S. Berben, D. Novikov, W. T. Stuart *et al.*, 2010 Resolving
1920   mechanisms of competitive fertilization success in Drosophila melanogaster.
1921   Science 328**:** 354-357.
1922 Marciniak, P., A. Urbanski, M. Kudlewska, M. Szymczak and G. Rosinski, 2017 Peptide
1923   hormones regulate the physiological functions of reproductive organs in Tenebrio
1924   molitor males. Peptides 98**:** 35-42.
1925 Matsumoto-Miyai, K., M. Yoshizumi and M. Kawatani, 2015 Regulatory Effects of 5-
1926   Hydroxytryptamine Receptors on Voiding Function. Adv Ther 32 Suppl 1**:** 3-15.
1927 McCullough, E. L., E. Whittington, A. Singh, S. Pitnick, M. F. Wolfner *et al.*, 2022 The life
1928   history of Drosophila sperm involves molecular continuity between male and
1929   female reproductive tracts. Proc Natl Acad Sci U S A 119**:** e2119899119.
1930 McKinney, H. M., L. M. Sherer, J. L. Williams, S. J. Certel and R. S. Stowers, 2020
1931   Characterization of Drosophila octopamine receptor neuronal expression using
1932   MiMIC-converted Gal4 lines. J Comp Neurol 528**:** 2174-2194.
1933 Michel, M. C., and W. Vrydag, 2006 Alpha1-, alpha2- and beta-adrenoceptors in the
1934   urinary bladder, urethra and prostate. Br J Pharmacol 147 Suppl 2**:** S88-119.
1935 Miyazaki, T., H. Kobayashi and T. Tosaka, 1998 Presynaptic inhibition by noradrenaline
1936   of the EPSC evoked in neonatal rat sympathetic preganglionic neurons. Brain
1937   Res 790**:** 170-177.
1938 Moatt, J. P., C. Dytham and M. D. Thom, 2014 Sperm production responds to perceived
1939   sperm competition risk in male Drosophila melanogaster. Physiol Behav 131**:**
1940   111-114.
1941 Monastirioti, M., C. E. Linn, Jr. and K. White, 1996 Characterization of Drosophila
1942   tyramine beta-hydroxylase gene and isolation of mutant flies lacking octopamine.
1943   J Neurosci 16**:** 3900-3911.
1944 Norville, K., S. T. Sweeney and C. J. Elliott, 2010 Postmating change in physiology of
1945   male Drosophila mediated by serotonin (5-HT). J Neurogenet 24**:** 27-32.
1946 O'Connor, S. C., K. L. Brain and M. R. Bennett, 1999 Individual sympathetic varicosities
1947   possess different sensitivities to alpha 2 and P2 receptor agonists and
1948   antagonists in mouse vas deferens. Br J Pharmacol 128**:** 1739-1753.
1949 Oliveira, M. S., J. E. Serrao, G. Dias, L. C. B. Martins and V. A. Araújo, 2021 Anatomy
1950   and histology of the male reproductive tractof Machtima crucigera (Fabricius,
1951   1775) (Heteroptera: Coreidae). Zoologischer Anzeiger 293**:** 156-162.

Parnas, D., A. P. Haghighi, R. D. Fetter, S. W. Kim and C. S. Goodman, 2001
Regulation of postsynaptic structure and protein localization by the Rho-type
guanine nucleotide exchange factor dPix. Neuron 32**:** 415-424.
Pauls, D., C. Blechschmidt, F. Frantzmann, B. El Jundi and M. Selcho, 2018 A
comprehensive anatomical map of the peripheral octopaminergic/tyraminergic
system of Drosophila melanogaster. Sci Rep 8**:** 15314.
Pavlou, H. J., A. C. Lin, M. C. Neville, T. Nojima, F. Diao *et al.*, 2016 Neural circuitry
coordinating male copulation. Elife 5.
Pelletier, G., and R. Leclerc, 1979 Localization of Leu-enkephalin in dense core vesicles
of axon terminals. Neurosci Lett 12**:** 159-163.
Petersen, L. K., and R. S. Stowers, 2011 A Gateway MultiSite recombination cloning
toolkit. PLoS One 6**:** e24531.
Pizzari, T., C. K. Cornwallis, H. Lovlie, S. Jakobsson and T. R. Birkhead, 2003
Sophisticated sperm allocation in male fowl. Nature 426**:** 70-74.
Port, F., H. M. Chen, T. Lee and S. L. Bullock, 2014 Optimized CRISPR/Cas tools for
efficient germline and somatic genome engineering in Drosophila. Proc Natl Acad
Sci U S A 111**:** E2967-2976.
Pound, N., and M. J. G. Gage, 2004 Prudent sperm allocation in Norway rats,
:: a mammalian model of adaptive ejaculate adjustment. Animal Behaviour 68**:** 819-823.
Prins, N. H., M. R. Briejer, P. J. Van Bergen, L. M. Akkermans and J. A. Schuurkes,
1999 Evidence for 5-HT7 receptors mediating relaxation of human colonic
circular smooth muscle. Br J Pharmacol 128**:** 849-852.
Qi, Y. X., G. Xu, G. X. Gu, F. Mao, G. Y. Ye *et al.*, 2017 A new Drosophila octopamine
receptor responds to serotonin. Insect Biochem Mol Biol 90**:** 61-70.
Recio, P., M. V. Barahona, L. M. Orensanz, S. Bustamante, A. C. Martinez *et al.*, 2009
5-hydroxytryptamine induced relaxation in the pig urinary bladder neck. Br J
Pharmacol 157**:** 271-280.
Rohrbach, E. W., J. D. Asuncion, P. Meera, M. Kralovec, S. A. Deshpande *et al.*, 2024a
Heterogeneity in the projections and excitability of tyraminergic/octopaminergic
neurons that innervate the Drosophila reproductive tract. Front Mol Neurosci 17**:**
1374896.
Rohrbach, E. W., E. M. Knapp, S. A. Deshpande and D. E. Krantz, 2024b Expression
and potential regulatory functions of Drosophila octopamine receptors in the
female reproductive tract. G3 (Bethesda) 14.
Scheibner, J., A. U. Trendelenburg, L. Hein and K. Starke, 2001 Stimulation frequency-
noradrenaline release relationships examined in alpha2A-, alpha2B- and
alpha2C-adrenoceptor-deficient mice. Naunyn Schmiedebergs Arch Pharmacol
364**:** 321-328.
Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair *et al.*, 2012 Fiji: an
open-source platform for biological-image analysis. Nat Methods 9**:** 676-682.
Segawa, T., H. Murakami, H. Ogawa and H. Yajima, 1978 Effect of enkephalin and
substance P on sympathetic nerve transmission in mouse vas deferens. Jpn J
Pharmacol 28**:** 13-19.
Sepil, I., B. R. Hopkins, R. Dean, M. L. Thezenas, P. D. Charles *et al.*, 2019 Quantitative
Proteomics Identification of Seminal Fluid Proteins in Male Drosophila
melanogaster. Mol Cell Proteomics 18**:** S46-S58.

Shearin, H. K., A. R. Dvarishkis, C. D. Kozeluh and R. S. Stowers, 2013 Expansion of
the gateway multisite recombination cloning toolkit. PLoS One 8**:** e77724.
Shearin, H. K., I. S. Macdonald, L. P. Spector and R. S. Stowers, 2014 Hexameric GFP
and mCherry reporters for the Drosophila GAL4, Q, and LexA transcription
systems. Genetics 196**:** 951-960.
Shen, H., R. A. M. Marino, R. A. McDevitt, G. H. Bi, K. Chen *et al.*, 2018 Genetic
deletion of vesicular glutamate transporter in dopamine neurons increases
vulnerability to MPTP-induced neurotoxicity in mice. Proc Natl Acad Sci U S A
115**:** E11532-E11541.
Sherer, L. M., E. Catudio Garrett, H. R. Morgan, E. D. Brewer, L. A. Sirrs *et al.*, 2020
Octopamine neuron dependent aggression requires dVGLUT from dual-
transmitting neurons. PLoS Genet 16**:** e1008609.
Shirangi, T. R., A. M. Wong, J. W. Truman and D. L. Stern, 2016 Doublesex Regulates
the Connectivity of a Neural Circuit Controlling Drosophila Male Courtship Song.
Dev Cell 37**:** 533-544.
Siwicki, K. K., and E. A. Kravitz, 2009 Fruitless, doublesex and the genetics of social
behavior in Drosophila melanogaster. Curr Opin Neurobiol 19**:** 200-206.
Somogyi, G. T., and W. C. de Groat, 1990 Modulation of the release of
[3H]norepinephrine from the base and body of the rat urinary bladder by
endogenous adrenergic and cholinergic mechanisms. J Pharmacol Exp Ther
255**:** 204-210.
Soni, K. K., H. S. Jeong and S. Jang, 2022 Neurons for Ejaculation and Factors
Affecting Ejaculation. Biology (Basel) 11.
Starke, K., 1972 Alpha sympathomimetic inhibition of adrenergic and cholinergic
transmission in the rabbit heart. Naunyn Schmiedebergs Arch Pharmacol 274**:**
18-45.
Stowers, R. S., 2025 Multimerized Epitope Tags for High Sensitivity Protein Detection.
G3 (Bethesda).
Susic-Jung, L., C. Hornbruch-Freitag, J. Kuckwa, K. H. Rexer, U. Lammel *et al.*, 2012
Multinucleated smooth muscles and mononucleated as well as multinucleated
striated muscles develop during establishment of the male reproductive organs of
Drosophila melanogaster. Dev Biol 370**:** 86-97.
Tainio, H., 1995 Peptidergic innervation of the human prostate, seminal vesicle and vas
deferens. Acta Histochem 97**:** 113-119.
Tayler, T. D., D. A. Pacheco, A. C. Hergarden, M. Murthy and D. J. Anderson, 2012 A
neuropeptide circuit that coordinates sperm transfer and copulation duration in
Drosophila. Proc Natl Acad Sci U S A 109**:** 20697-20702.
Tison, K. V., H. M. McKinney and R. S. Stowers, 2020 Demonstration of a Simple
Epitope Tag Multimerization Strategy for Enhancing the Sensitivity of Protein
Detection Using Drosophila vAChT. G3 (Bethesda) 10**:** 495-504.
Tomas, A. L., M. A. G. Bento, L. D. Mutti, F. J. Zara and L. S. L. Greco, 2019 New
insights in the male anatomy, spermatophore formation, and sperm structure in
Atyidae: The red cherry shrimp. Invertebrate Biology 138**:** 17-28.
Trendelenburg, A. U., A. Meyer, W. Klebroff, S. Guimaraes and K. Starke, 2003
Crosstalk between presynaptic angiotensin receptors, bradykinin receptors and

alpha 2-autoreceptors in sympathetic neurons: a study in alpha 2-adrenoceptor-
deficient mice. Br J Pharmacol 138**:** 1389-1402.

Trendelenburg, A. U., K. Starke and N. Limberger, 1994 Presynaptic alpha 2A-
adrenoceptors inhibit the release of endogenous dopamine in rabbit caudate
nucleus slices. Naunyn Schmiedebergs Arch Pharmacol 350**:** 473-481.

Truitt, W. A., and L. M. Coolen, 2002 Identification of a potential ejaculation generator in
the spinal cord. Science 297**:** 1566-1569.

Verhulst, E. C., and L. van de Zande, 2015 Double nexus--Doublesex is the connecting
element in sex determination. Brief Funct Genomics 14**:** 396-406.

Villella, A., and J. C. Hall, 2008 Neurogenetics of courtship and mating in Drosophila.
Adv Genet 62**:** 67-184.

Wedell, N., and P. A. Cook, 1999 Butterflies tailor their ejaculate in response to sperm
competition risk and intensity. Proceedings of the Royal Society B-Biological
Sciences 266**:** 1033-1039.

Wigby, S., N. C. Brown, S. E. Allen, S. Misra, J. L. Sitnik *et al.*, 2020 The Drosophila
seminal proteome and its role in postcopulatory sexual selection. Philos Trans R
Soc Lond B Biol Sci 375**:** 20200072.

Williams, J. L., H. K. Shearin and R. S. Stowers, 2019 Conditional Synaptic Vesicle
Markers for Drosophila. G3 (Bethesda) 9**:** 737-748.

Wolfner, M. F., 1997 Tokens of love: functions and regulation of Drosophila male
accessory gland products. Insect Biochem Mol Biol 27**:** 179-192.

Wong, R., and A. B. Lange, 2014 Octopamine modulates a central pattern generator
associated with egg-laying in the locust, Locusta migratoria. J Insect Physiol 63**:**
1-8.

Wuest, M., B. Eichhorn, M. O. Grimm, M. P. Wirth, U. Ravens *et al.*, 2009
Catecholamines relax detrusor through beta 2-adrenoceptors in mouse and beta
3-adrenoceptors in man. J Pharmacol Exp Ther 328**:** 213-222.

Yamamoto, D., 2007 The neural and genetic substrates of sexual behavior in
Drosophila. Adv Genet 59**:** 39-66.

Yartsev, V. V., and S. S. Evseeva, 2021 The Male Urogenital System of a Salamander
2073
2074 (Amphibia, Caudata). Current Herpetology 40**:** 10-21.

2075 Yu, J., Y. Zhang, K. Clements, N. Chen and L. C. Griffith, 2025 Genetically-encoded
2076   markers for confocal visualization of single dense core vesicles. Commun Biol 8**:**
2077   383.

2078 Zhang, Y., M. Rozsa, Y. Liang, D. Bushey, Z. Wei *et al.*, 2023 Fast and sensitive GCaMP
2079   calcium indicators for imaging neural populations. Nature 615**:** 884-891.

2080