1   The Illusion of Polygenicity in Pool-seq Genetic Mapping studies: Insufficient Power Can Mask
2   Simple Genetic Architectures
3
4

5   Anthony D. Long[1], Katherine M. Hanson[2], Stuart J Macdonald[2]
6
7   [1] Ecology and Evolutionary Biology, 321 Steinhaus Hall, University of California, Irvine, CA
8   92697, USA
9
10   [2] Department of Molecular Biosciences, University of Kansas, 1200 Sunnyside Avenue,
11   Lawrence, KS 66045, USA
12
13   Corresponding Author:
14
15   Anthony D. Long
16   Ecology and Evolutionary Biology
17   321 Steinhaus Hall
18   University of California
19   Irvine, CA 92697
20
21   Email: tdlong@uci.edu
22

Abstract:

Pool-seq (pooled sequencing) combines DNA from multiple individuals prior to sequencing,
enabling population-level allele frequency estimation without individual genotyping. When
employed in Genome Wide Association Studies (GWAS) pool-seq faces a fundamental power
limitation in that errors on allele frequency estimates are proportional to sequence coverage.
Although this power limitation is widely appreciated, pool-seq GWAS lacking unambiguous hits
are often interpreted as showing a highly polygenic genetic architecture. We illustrate the
limitation of inferring architecture from Manhattan plots using empirical data from a *Drosophila*
zinc resistance mapping study. Despite achieving an average of >700× sequencing coverage in
case and control pools, a directly ascertained SNP-based GWAS failed to reveal clear evidence
for major-effect loci. A unique feature of the dataset is that an advanced intercross multiparent
population, with known founders, was employed as the base population for the GWAS. We
leverage this unique population structure to carry out a second GWAS using imputed haplotype
frequency estimates, which in contrast revealed localized regions of major effect. A third
reanalysis of the same data using imputed SNP genotypes derived from the founder haplotype
frequency estimates uncovered a similar major gene architecture. The key difference between
approaches lies in statistical power: directly ascertained SNP counts have errors proportional to
sequencing coverage whereas known founder imputation-based approaches can be
considerably more accurate. This work highlights that insufficiently powered GWAS studies can
mask simple genetic architectures and create the illusion of polygenicity through statistical noise
alone.

Pool-seq (pooled sequencing) is a cost-effective genomic approach where DNA from multiple
individuals is combined prior to short-read sequencing, allowing researchers to estimate the
allele frequency in a population without individually genotyping each individual (Futschik and
Schlötterer 2010; Schlötterer et al. 2014). Pool-seq has been widely employed across diverse
research contexts including population genomics (Kofler et al. 2011; Czech et al. 2024),
experimental evolution studies (Burke et al. 2010; Turner et al. 2011; Orozco-terWengel et al.
2012; Burke et al. 2014), agricultural breeding programs (Michelmore et al. 1991), and genetic
mapping in model organisms (Ehrenreich et al. 2010; Macdonald et al. 2022). A fundamental
principle of pool-seq is that, for autosomal loci, when the number of chromosomes in the pool,
$2N$, substantially exceeds the depth of sequence coverage, C (*i.e., $2N >> C$*), directly
ascertained SNP frequency estimates provide unbiased estimates of allele frequencies in the
population from which the sample is drawn (Futschik and Schlötterer 2010). Furthermore, the
precision of these frequency estimates follows binomial sampling theory, with standard errors
proportional to $\sqrt{(pq/C)}$, where p and q are the reference and alternative allele frequencies.
59
Pool-seq has been increasingly applied to case-control genome-wide association studies
(GWAS), where researchers compare allele frequencies between phenotypically distinct pools
of individuals to identify genetic loci contributing to complex trait variation (Huang et al. 2012;
Bastide et al. 2013; Morozova et al. 2015; Fochler et al. 2017; Zhou et al. 2017; Macdonald et
al. 2022; Macdonald and Long 2022). This approach is analogous to the human case-control
GWAS design (Risch and Merikangas 1996; Wellcome Trust Case Control Consortium 2007), in
which a phenotypically extreme group of N_1 affected case individuals are compared to N_2
control individuals, except that rather than genotyping individuals via SNP arrays, pools of DNA
from case or control individuals are subjected to short read sequencing to coverages C_1 and C_2.
The standard GWAS tests for differences in allele frequencies between the $2N_1$ case versus $2N_2$
control autosomal alleles, whereas under the pool-seq design the tests are for differences
between the C_1 case versus C_2 control reads per locus. Several studies have successfully
implemented this approach across model organisms demonstrating its feasibility for trait
dissection (Craig et al. 2009; Bastide et al. 2013; Lirakis et al. 2022; Davies and Myles 2023).
However, much like a standard GWAS, pool-seq GWAS faces the fundamental trade-off
between controlling false positive rates through stringent significance thresholds and
maintaining sufficient statistical power to detect true associations. For example, The Wellcome
Trust Case Control Consortium established that human GWAS requires significance thresholds
of at least -log10(p-value) > 6, paired with on the order of three thousand fully genotyped case
versus control individuals, to reliably detect subtle effect risk alleles while controlling for multiple
tests (Wellcome Trust Case Control Consortium 2007).
81
The power limitations of case-control comparisons using pool-seq are well-established
(Baldwin-Brown et al. 2014; Kofler and Schlötterer 2014): to reliably detect allele frequency
differences <10%, pools of thousands of individuals sequenced to >1000X average coverage
are likely required. Despite this constraint, Manhattan plots of -log10(p-value) against genomic
location from experiments with much lower sequence coverage, and that show little evidence of
"hits", are often interpreted as representing a highly polygenic architecture (Huang et al. 2012;
Orozco-terWengel et al. 2012; Morozova et al. 2015; Fochler et al. 2017; Zhou et al. 2017;

Barghi et al. 2019; Lirakis et al. 2022). A polygenic architecture may indeed underlie the trait,
but Manhattan plots from modestly powered studies are a poor proxy for genetic architecture.
To quantify this limitation, we conducted simulations of pooled sequencing case-control
experiments with pools of 10,000 individuals, varying levels of sequencing coverage (400X,
1000X and 5000X), and a single SNP with a true allele frequency difference of 4% or 8%
between pools (Figure 1). Such allele frequency differences would be considered large by the
standards of human disease GWAS studies, yet the sample sizes and coverages required to
detect these loci are never achieved in published pool-seq studies. Our simulations suggest that
only at 5000X coverage for an allele frequency difference of 8% is the allele of large effect
reliably detected. It would be easy to conclude from the Manhattan plots for the more modestly
powered experiments shown in Figure 1 that the simulated trait has a polygenic basis, and that
many of the sites with highest LOD scores are true, but very modest-effect causative loci. The
combination of limited power and genome-wide testing can generate apparent polygenic signals
through statistical noise alone. This phenomenon, where insufficient statistical power masks a
relatively simple genetic architecture, represents a critical interpretive challenge in pool-seq
studies.
105
A recently published dataset in *Drosophila melanogaster* allows us to uniquely illustrate the
problem in an empirical as opposed to a simulated dataset. Hanson et al. (Hanson et al. 2025)
utilized a base population derived by combining 663 Drosophila Synthetic Population Resource
(DSPR) 8-way, known founder advanced intercross recombinant inbred lines. They collected 0-
24 hour old embryos and exposed them to either 25 mM zinc chloride or water. Case samples
were generated from the ~7% most zinc-resistant adult females that survived the zinc treatment,
while controls represent samples of individuals that developed successfully in normal media
conditions. Across 12 replicate selection experiments, a total of 4831 control and 3448 case
individuals were sampled and pooled for sequencing. A total of 739× and 983× sequence
coverage for the zinc-treated "case" and water-treated control pools was obtained, respectively
(Figure 2). Ignoring the unique population structure of the base population employed, the
experiment is a large case-control GWAS by Drosophila standards. Indeed, it is conceivable
that the power of the experiment exceeds that of a typical 200-500 inbred-line-based mapping
study (Macdonald et al. 2022).
120
We carried out a GWAS of these data by contrasting directly ascertained allele frequencies
between cases and controls, summarizing the experiment with a Manhattan plot (Figure 3A).
Although 642 SNPs exceed a Bonferroni significance threshold, these significant associations
show little apparent localization to specific genomic regions and instead appear dispersed
throughout the genome. Based on these data alone one could easily conclude that zinc
resistance in this population has a highly polygenic genetic architecture.
127
The dataset of Hanson et al. (Hanson et al. 2025) is unique in that the base population
employed is derived from an advanced intercross synthetic population originating from eight
known founder strains. This experimental design allows the frequency of those eight founder
alleles to be estimated from the pool-seq data throughout the genome using a sliding window
haplotype estimator (Long et al. 2011; Kessner et al. 2013; Burke et al. 2014; Tilk et al. 2019;

Linder et al. 2020; Linder et al. 2022; Macdonald et al. 2022). Specifically, Hanson et al.
estimated founder haplotype frequencies in overlapping 1.5-cM windows across the genome for
each sample. These eight founder haplotype frequencies were then tested for differentiation as
a function of genomic location. This known-founder haplotype-based testing strategy underlies
the X-QTL (or bulked segregant) analysis approach to mapping QTL (Ehrenreich et al. 2012;
Macdonald et al. 2022; Macdonald and Long 2022). Based on this X-QTL statistical approach,
employing the same raw data as the directly ascertained SNP GWAS described above, Hanson
et al. concluded that resistance is associated with regions of major effect localized at
centimorgan scales. They further identified and validated several major genes in these intervals
(including *MTF-1*, a gene central to zinc homeostasis). Figure 3B reproduces the scan of
Hanson et al and clearly shows highly significant peaks of association localized to specific
genomic regions. The results of this zinc-resistance mapping are consistent with other studies
where multiparent population (MPP)-based mapping have identified regions of large effect (King
et al. 2012; King et al. 2014; Najarro et al. 2015; Everman et al. 2021). They present a stark
contrast to the dispersed association pattern observed with directly ascertained SNP-based
analysis.
149
We claim that the fundamental reason the haplotype-based test succeeded in detecting major-
effect loci, while the direct SNP analysis failed, is simply due to poor power of the latter
approach. Despite achieving greater than 700× sequencing coverage per treatment, the directly
ascertained SNP approach still suffers from relatively large binomial sampling errors on the SNP
frequency estimates. In turn, this noise in allele frequency estimation obscures genuine allele
frequency differences between pools. In contrast, the haplotype-based approach leverages the
critical statistical advantage that haplotype frequency estimates have much smaller errors than
directly ascertained SNP frequencies. The error on imputed haplotype frequencies are
proportional to ~75-100% of the binomial sampling errors on twice the number of individuals per
treatment, as opposed to the average short-read coverage (Tilk et al. 2019; Linder et al. 2020).
In the zinc chloride experiment the average number of autosomal alleles per treatment (2×4140
= 8280) is roughly ten-fold greater than the average sequence coverage per treatment (700×),
and as a result, haplotype frequencies are estimated with much greater certainty than directly
ascertained SNP frequencies. The haplotype approach effectively circumvents the coverage
limitation of directly ascertaining SNP frequencies. At sequencing coverages much less than
2N, as is typically the case with pool-seq datasets, imputing founder haplotype frequencies is a
"trick" that can be used to achieve frequency estimation errors approaching genotyping every
individual.
168
To illustrate that the difference we observe between the haplotype-based approach (Figure 3B)
and the directly ascertained SNP-based approach (Figure 3A) is primarily due to sampling error
on frequency estimates, and not somehow associated with the decision to employ haplotypes or
some other detail of X-QTL mapping procedure, we used the haplotype frequency estimates to
impute SNP frequencies. That is, we imputed SNP frequencies throughout the genome as the
sum of the Hadamard product of the founder frequency estimates and known SNP states in the
founders at any given location. Although counter-intuitive, just like the haplotype frequencies,
these imputed SNP frequencies are estimated much more accurately than the directly

ascertained SNP counts (Tilk et al. 2019; Linder et al. 2020). We then converted imputed SNP
frequencies to pseudo-counts based on the number of individuals in each pool and carried out a
third GWAS on the imputed SNP counts (Figure 3C). The plot demonstrates that when we use
the more accurate imputed, as opposed to directly ascertained, SNP allele frequencies we
recover the clear peaks of association seen in the haplotype-based analysis. This confirms that
the failure of direct SNP analysis stems from poor power. Looking closely at the tests on
imputed SNP frequencies (Supplementary Figure 1) it further appears as though one could fit
several lines to that data. This pattern represents sets of SNPs that are "in phase" with one
another with respect to different founder chromosomes and is expected in multiparent panels
derived from a modest number of founders.
187
The power limitations we demonstrate are not specific to synthetic populations or X-QTL
methodology but represent a fundamental challenge for pooled sequencing approaches. While
our analysis employed a synthetic population derived from eight known founders, the underlying
statistical problem, that errors on directly ascertained SNP frequency estimates are large
relative to subtle allele frequency differences between pools, applies broadly to any pooled
sequencing design. In outbred populations lacking known founder structure, researchers cannot
employ the imputation strategy that rescued power in the illustrative example we provide here.
The apparent success of haplotype-based approaches in synthetic populations stems not from
any inherent superiority of haplotypes over SNPs, but simply because haplotype frequencies
are estimated more accurately, effectively circumventing the coverage limitations that plague
direct SNP ascertainment.
199
A criticism that can be leveled at the illustrative example presented in this work is that the
statistical tests for directly ascertained SNPs use total counts derived from sequence coverage
(see Methods), whereas the two imputation-based tests use total counts associated with twice
the number of individuals in the pool from which DNA is made. Counts associated with the 2N
individuals in the DNA pools result in higher power, since 2N is more than ten times the
sequence coverage per pool. However, the total sequence coverage is the true count
associated with directly ascertained SNPs, whereas the 2N associated with imputation-based
tests incorrectly assumes imputation is equivalent to perfectly genotyping the N individuals in
each pool. Imputation is simply not that effective. But the decision to multiply frequencies by 2N
to obtain counts *only* impacts the statistical test as a multiplicative constant throughout the
genome. As a result, the imputation-based Manhattan plots of Figure 3B&C simply have an
"inflated" Y-axis scale. But this inflation constant does not contribute to the noise of the directly
ascertained GWAS.
213
A second criticism is that an MPP is not appropriate for a GWAS-type analysis, since a synthetic
population derived from an 8-way cross is not the type of population typically employed in a
GWAS study (generally a sample of natural chromosomes lacking a known set of founders).
This is reflected in Figure 3 where for the two imputation-based scans the "hits" extend over
centimorgan sized intervals, a much larger region than the localization signal expected of a
GWAS. But this difference doesn't detract from the value of our illustration. We observe that
directly ascertained SNP frequencies at coverages approaching 700X per sample are

insufficient to detect the real, but subtle frequency shifts apparent in this dataset, a result
independent of the genetic details of the population employed. If anything, the problem is more
dire in a population harboring lower levels of linkage disequilibrium as a function of distance. In
such a case one is trying to detect the signal associated with only a handful of SNPs in LD with
a causative factor, and a lower genome-wide *p*-value threshold is likely required to control for
false positives.
227
In *Drosophila melanogaster*, we are aware of four traits have been dissected using both an
MPP-panel based mapping approach using the *Drosophila* Synthetic Population Resource
(DSPR; King et al. 2012) and a GWAS-based analysis using collection of inbred lines called the
Drosophila Genetic Reference panel (DGRP; Mackay et al. 2012; Huang et al. 2014) where the
genetic architecture appears complex: resistance to caffeine (Najarro et al. 2015), copper
toxicity (Everman et al. 2021; Everman et al. 2023), boric acid resistance (Najarro et al. 2017),
and starvation stress (Everman et al. 2019). In all four cases DSPR mapping-based approaches
have successfully identified loci of large effect, while in contrast DGRP association-based
approaches have concluded that trait variation is highly polygenic and due to dozens to
hundreds of loci of subtle effect spread throughout the genome. Notably the association hits are
subtle enough that they rely on a suggestive threshold for significance, often a -log10(p-value) >
5. Our observations with respect to pool-seq case-control designs suggest that power deficits
can lead to incorrect assumptions about the polygenicity of complex traits, and we speculate
that a similar pattern may explain the contradictory results observed in these inbred line studies.
The highly polygenic signal in these four association studies is consistent with DGRP-based
GWAS outcomes more generally (Mackay and Huang 2018), as well as those in other
Drosophila GWAS-type studies employing individual-based genotyping (Pallares et al. 2023).
Perhaps the observed difference between DGRP and DSPR outcomes is similarly driven by
differences in power. Overall, in Drosophila the genetic basis of trait variation dissected using
GWAS-type approaches appears highly polygenic, while in contrast traits dissected using QTL-
mapping type approaches appear more oligogenic, and this remains true even when holding the
trait constant. It is possible that the apparent contradiction between polygenicity versus handfuls
of more major effect loci is driven by power. Many traits currently characterized as highly
polygenic based on dispersed GWAS signals are consistent with the traits truly being polygenic,
but equally consistent with a more oligogenic architecture that studies simply are poorly
powered to detect.
254
255   **Methods:**
256
The zinc chloride mapping dataset is described in detail in Hanson et al. (Hanson et al. 2025).
Briefly, the study utilized a population derived by intercrossing 663 Drosophila Synthetic
Population Resource (DSPR) "A" founder population recombinant inbred lines. After 31-36
generations of synthetic population maintenance, 12 replicate zinc selection assays were
performed. To perform the assays, 0–24-hour old embryos were collected from the base
population and transferred to control (water) or treatment (25-mM zinc chloride) bottles.
Emerged adult females were collected, and pooled DNA samples were prepared for both control
and zinc-selected groups, DNA libraries were prepared using the Illumina DNA Prep protocol

and sequenced with PE150 reads on an Illumina NovaSeq 6000. Reads were mapped to the *D.*
melanogaster Release 6 reference genome, REF/ALT counts were obtained for every SNP, and
founder haplotype frequencies estimated in overlapping 1.5-cM windows across the genome for
each sample.
269
Here, we employ the raw SNP tables and haplotype frequency estimates from Hanson et al.
(2025). But unlike that earlier work all the statistical tests in the present study employ Cochran–
Mantel–Haenszel (CHM) tests on counts as opposed to ANOVA on square root transformed
frequency estimates. The zinc chloride experiment consisted of twelve replicate control versus
zinc-selected comparisons, so CHM is an appropriate test. Additionally, the use of the CHM test
for all three analytical approaches allows for more intuitive comparisons between the study
designs. For directly ascertained SNP scans the CHM tests employed raw REF vs ALT counts
tables, and as a result the total counts in the analysis at any given SNP is the total coverage at
the SNP. In contrast the two imputation-based designs use the raw counts and founder
genotypes to impute haplotype or SNP frequencies at 10kb steps, or at each SNP in the
genome, respectively. SNP frequencies are imputed by multiplying the closest set of haplotype
frequency estimates by the founder states (generally 0 or 1) for each haplotype. To convert per-
sample imputed haplotype or SNP frequencies to counts, those frequencies are multiplied by
twice the number of individuals contributing to that pooled DNA sample. The number of
individuals in a pool is a genome-wide constant per pool, as a result it will not generate variation
between sites, instead it will result in -log10(p-values)'s inflated by a constant factor.
286
MSPrime (Kelleher and Lohse 2020; Baumdicker et al. 2022) was used to simulate 500
haplotypes from a neutrally evolving 1Mb region under *Drosophila melanogaster*-like
parameters (N_e=1e6, u=5e-9, r=2e-8). These haplotypes were sampled with replacement to
create a ten thousand haplotype control base population. A similar 10,000 haplotype case
population was created by sampling haplotypes with replacement from the control population
conditional on a single focal SNP having an allele frequency difference of 4% or 8% from the
controls. Frequencies were then estimated at all common SNPs for both case and control
populations, and then pool-seq allele counts obtained via random negative binomial draws (size
= mu = expected coverage) conditional on the known frequencies and expected sequencing
coverages of 400X, 1000X, or 5000X. The negative binomial was employed so that realized
coverage varied among SNP positions according to an over-dispersed poisson with a variance
twice the mean (equal to expected coverage). Chi-square tests were carried out at each SNP
and -log10(p-values) obtained.
300
301 **Data and Methods Availability:**
302
303 The raw data, methods for calling SNPs, and haplotyping calling are described in the Hanson
304 2025. Briefly, DSPR RILs are available from the BDSC, raw FASTQ files are in the NCBI SRA
305 under accession number PRJNA1127662, and all code to replicate the analyses is available on
306 GitHub (https://github.com/Hanson19/Zinc-X-QTL). The analysis code, and code to reproduce
307 the figures of this work are hosted on Github (https://github.com/tdlong/Compare_3_GWAS.git).
308

309 **Acknowledgements:**

311 This work was supported by NIH award R01-OD034064 (to SJM and ADL).

312 **Figures**

314
315

Figure 1: Simulated pool-seq case/control experiments. We simulated 500 haplotypes from a
1Mb region with typical *D. melanogaster* population genetic parameters. These haplotypes were
instantaneously expanded to create 10,000 allele Case versus Control pools, subsequently
pool-sequenced to expected coverages of 400X, 1000X, or 5000X. The case haplotype pools
were sampled conditioning on a single intermediate frequency focal SNP (labelled in red) having
a true allele frequency difference between cases or controls of 4% or 8%. We simulate a 1Mb
region here and note a -log10(p) threshold of 5 is commonly used in *D. melanogaster* GWAS
studies for statistical significance.

324
325
Figure 2: Sampling properties of an X-QTL experiment challenging a *Drosophila melanogaster*
synthetic population with zinc chloride (Hanson et al. 2025). The experiment consisted of 12
paired replicates of Control versus Zinc-selected treatments (Trt). For each treatment and
replicate N individuals were sampled to create DNA pools (left panel), and these were
subsequently pool-sequenced to the coverage depicted in the right panel. Replicates 5, 6, 8, 11
and 12 had two separate, equally sized control pools.

Figure 3: Zinc chloride developmental resistance Manhattan plots associated with three

different approaches to statistical testing: (A) directly ascertained REF and ALT SNP counts (we

do not display a statistical threshold but note a -log10(p) threshold of 5 is widely used in

Drosophila GWAS), (B) Imputed founder haplotype counts, and (C) imputed SNP counts. X-

axes are scaled in genetic as opposed to physical distances. *p*-values are obtained from

Cochran–Mantel–Haenszel tests on counts over the 12 replicates of the experiment. For directly

ascertained SNPs, the counts used are observed REF vs ALT counts. For the imputation-based

tests, *pseudo*-counts are the product of haplotype (or SNP) frequency estimates and twice the

number of individuals per DNA pool. These pseudo-counts will inflate -log10(p-values) by a

constant multiplicative factor.

Supplementary Figure 1: Zinc chloride developmental resistance Manhattan plots associated

with three different approaches to statistical testing for chromosome 3R: (A) directly ascertained

REF and ALT SNP counts, (B) Imputed founder haplotype counts, and (C) imputed SNP counts.

These panels are otherwise the same as Figure 3, except for panel A where we limit the Y-axis

to 15.

References

Baldwin-Brown JG, Long AD, Thornton KR. 2014. The power to detect quantitative trait loci using resequenced, experimentally evolved populations of diploid, sexual organisms. Mol Biol Evol. 31(4):1040–1055.

Barghi N, Tobler R, Nolte V, Jakšić AM, Mallard F, Otte KA, Dolezal M, Taus T, Kofler R, Schlötterer C. 2019. Genetic redundancy fuels polygenic adaptation in Drosophila. PLoS Biol. 17(2):e3000128.

Bastide H, Betancourt A, Nolte V, Tobler R, Stöbe P, Futschik A, Schlötterer C. 2013. A genome-wide, fine-scale map of natural pigmentation variation in Drosophila melanogaster. PLoS Genet. 9(6):e1003534.

Baumdicker F, Bisschop G, Goldstein D, Gower G, Ragsdale AP, Tsambos G, Zhu S, Eldon B, Ellerman EC, Galloway JG, et al. 2022. Efficient ancestry and mutation simulation with msprime 1.0. Genetics. 220(3). doi:10.1093/genetics/iyab229. http://dx.doi.org/10.1093/genetics/iyab229.

Burke MK, Dunham JP, Shahrestani P, Thornton KR, Rose MR, Long AD. 2010. Genome-wide analysis of a long-term evolution experiment with Drosophila. Nature. 467(7315):587–590.

Burke MK, Liti G, Long AD. 2014. Standing genetic variation drives repeatable experimental evolution in outcrossing populations of Saccharomyces cerevisiae. Mol Biol Evol. 31(12):3228–3239.

Craig JE, Hewitt AW, McMellon AE, Henders AK, Ma L, Wallace L, Sharma S, Burdon KP, Visscher PM, Montgomery GW, et al. 2009. Rapid inexpensive genome-wide association using pooled whole blood. Genome Res. 19(11):2075–2080.

Czech L, Spence JP, Expósito-Alonso M. 2024. Grenedalf: Population genetic statistics for the next generation of pool sequencing. Bioinformatics. 40(8):btae508.

Davies T, Myles S. 2023. Pool-seq of diverse apple germplasm reveals candidate loci underlying ripening time, phenolic content, and softening. Fruit Research. 3(1):0–0.

Ehrenreich IM, Bloom J, Torabi N, Wang X, Jia Y, Kruglyak L. 2012. Genetic architecture of highly complex chemical resistance traits across four yeast strains. PLoS Genet. 8(3):e1002570.

Ehrenreich IM, Torabi N, Jia Y, Kent J, Martis S, Shapiro JA, Gresham D, Caudy AA, Kruglyak L. 2010. Dissection of genetically complex traits with extremely large pools of yeast segregants. Nature. 464(7291):1039–1042.

Everman ER, Cloud-Richardson KM, Macdonald SJ. 2021. Characterizing the genetic basis of copper toxicity in Drosophila reveals a complex pattern of allelic, regulatory, and behavioral variation. Genetics. 217(1):1–20.

Everman ER, Macdonald SJ, Kelly JK. 2023. The genetic basis of adaptation to copper pollution in Drosophila melanogaster. Front Genet. 14:1144221.

Everman ER, McNeil CL, Hackett JL, Bain CL, Macdonald SJ. 2019. Dissection of Complex, Fitness-Related Traits in Multiple Drosophila Mapping Populations Offers Insight into the

Genetic Control of Stress Resistance. Genetics. 211(4):1449–1467.

Fochler S, Morozova TV, Davis MR, Gearhart AW, Huang W, Mackay TFC, Anholt RRH. 2017.
Genetics of alcohol consumption in Drosophila melanogaster. Genes Brain Behav. 16(7):675–
685.

Futschik A, Schlötterer C. 2010. The next generation of molecular markers from massively
parallel sequencing of pooled DNA samples. Genetics. 186(1):207–218.

Hanson KM, Long AD, Macdonald SJ. 2025. Extreme QTL mapping enables the identification of
novel zinc toxicity response loci in Drosophila melanogaster. Genetics.:iyaf173.

Huang W, Massouras A, Inoue Y, Peiffer J,Ràmia M, Tarone AM, Turlapati L, Zichner T, Zhu D,
Lyman RF, et al. 2014. Natural variation in genome architecture among 205 Drosophila
melanogaster Genetic Reference Panel lines. Genome Res. 24(7):1193–1208.

Huang W, Richards S, Carbone MA, Zhu D, Anholt RRH, Ayroles JF, Duncan L, Jordan KW,
Lawrence F, Magwire MM, et al. 2012. Epistasis dominates the genetic architecture of
Drosophila quantitative traits. Proc Natl Acad Sci U S A. 109(39):15553–15559.

Kelleher J, Lohse K. 2020. Coalescent simulation with msprime. Methods Mol Biol. 2090:191–
230.

Kessner D, Turner TL, Novembre J. 2013. Maximum likelihood estimation of frequencies of
known haplotypes from pooled sequence data. Mol Biol Evol. 30(5):1145–1158.

King EG, Kislukhin G, Walters KN, Long AD. 2014. Using Drosophila melanogaster to identify
chemotherapy toxicity genes. Genetics. 198(1):31–43.

King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, Long AD, Macdonald SJ.
2012. Genetic dissection of a model complex trait using the Drosophila Synthetic Population
Resource. Genome Res. 22(8):1558–1566.

Kofler R, Orozco-terWengel P, De Maio N, Pandey RV, Nolte V, Futschik A, Kosiol C,
Schlötterer C. 2011. PoPoolation: a toolbox for population genetic analysis of next generation
sequencing data from pooled individuals. PLoS One. 6(1):e15925.

Kofler R, Schlötterer C. 2014. A guide for the design of evolve and resequencing studies. Mol
Biol Evol. 31(2):474–483.

Linder RA, Majumder A, Chakraborty M, Long A. 2020. Two Synthetic 18-Way Outcrossed
Populations of Diploid Budding Yeast with Utility for Complex Trait Dissection. Genetics.
215(2):323–342.

Linder RA, Zabanavar B, Majumder A, Hoang HC-S, Delgado VG, Tran R, La VT, Leemans
SW, Long AD. 2022. Adaptation in Outbred Sexual Yeast is Repeatable, Polygenic and Favors
Rare Haplotypes. Mol Biol Evol. 39(12). doi:10.1093/molbev/msac248.
http://dx.doi.org/10.1093/molbev/msac248.

Lirakis M, Nolte V, Schlötterer C. 2022. Pool-GWAS on reproductive dormancy in Drosophila
simulans suggests a polygenic architecture. G3 (Bethesda). 12(3):jkac027.

Long Q, Jeffares DC, Zhang Q, Ye K, Nizhynska V, Ning Z, Tyler-Smith C, Nordborg M. 2011.

PoolHap: inferring haplotype frequencies from pooled samples by next generation sequencing. PLoS One. 6(1):e15292.

Macdonald SJ, Cloud-Richardson KM, Sims-West DJ, Long AD. 2022. Powerful, efficient QTL mapping in Drosophila melanogaster using bulked phenotyping and pooled sequencing. Genetics. 220(3). doi:10.1093/genetics/iyab238. http://dx.doi.org/10.1093/genetics/iyab238.

Macdonald SJ, Long AD. 2022. Discovery of malathion resistance QTL in Drosophila melanogaster using a bulked phenotyping approach. G3 . 12(12). doi:10.1093/g3journal/jkac279. http://dx.doi.org/10.1093/g3journal/jkac279.

Mackay TFC, Huang W. 2018. Charting the genotype-phenotype map: lessons from the Drosophila melanogaster Genetic Reference Panel. Wiley Interdiscip Rev Dev Biol. 7(1). doi:10.1002/wdev.289. http://dx.doi.org/10.1002/wdev.289.

Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y, Magwire MM, Cridland JM, et al. 2012. The Drosophila melanogaster Genetic Reference Panel. Nature. 482(7384):173–178.

Michelmore RW, Paran I, Kesseli RV. 1991. Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. Proc Natl Acad Sci U S A. 88(21):9828–9832.

Morozova TV, Huang W, Pray VA, Whitham T, Anholt RRH, Mackay TFC. 2015. Polymorphisms in early neurodevelopmental genes affect natural variation in alcohol sensitivity in adult drosophila. BMC Genomics. 16:865.

Najarro MA, Hackett JL, Macdonald SJ. 2017. Loci Contributing to Boric Acid Toxicity in Two Reference Populations of Drosophila melanogaster. G3 . 7(6):1631–1641.

Najarro MA, Hackett JL, Smith BR, Highfill CA, King EG, Long AD, Macdonald SJ. 2015. Identifying Loci Contributing to Natural Variation in Xenobiotic Resistance in Drosophila. PLoS Genet. 11(11):e1005663.

Orozco-terWengel P, Kapun M, Nolte V, Kofler R, Flatt T, Schlötterer C. 2012. Adaptation of Drosophila to a novel laboratory environment reveals temporally heterogeneous trajectories of selected alleles. Mol Ecol. 21(20):4931–4941.

Pallares LF, Lea AJ, Han C, Filippova EV, Andolfatto P, Ayroles JF. 2023. Dietary stress remodels the genetic architecture of lifespan variation in outbred Drosophila. Nat Genet. 55(1):123–129.

Risch N, Merikangas K. 1996. The future of genetic studies of complex human diseases. Science. 273(5281):1516–1517.

Schlötterer C, Tobler R, Kofler R, Nolte V. 2014. Sequencing pools of individuals - mining genome-wide polymorphism data without big funding. Nat Rev Genet. 15(11):749–763.

Tilk S, Bergland A, Goodman A, Schmidt P, Petrov D, Greenblum S. 2019. Accurate Allele Frequencies from Ultra-low Coverage Pool-Seq Samples in Evolve-and-Resequence Experiments. G3 . 9(12):4159–4168.

Turner TL, Stewart AD, Fields AT, Rice WR, Tarone AM. 2011. Population-based resequencing
of experimentally evolved populations reveals the genetic basis of body size variation in
Drosophila melanogaster. PLoS Genet. 7(3):e1001336.

Wellcome Trust Case Control Consortium. 2007. Genome-wide association study of 14,000
cases of seven common diseases and 3,000 shared controls. Nature. 447(7145):661–678.

Zhou S, Luoma SE, St Armour GE, Thakkar E, Mackay TFC, Anholt RRH. 2017. A Drosophila
model for toxicogenomics: Genetic variation in susceptibility to heavy metal exposure. PLoS
Genet. 13(7):e1006907.