Intersecting experimental evolution and CRISPR screens to identify novel toxin resistance loci

Michele Marconcini[1], Steeve Cruchet[1], Srishti Goswami[2], Raghuvir Viswanatha[2], Matthew Butnaru[2,3], Joydeep De[4], Camilla Roselli[4], Dafni Hadjieconomou[4], Norbert Perrimon[2,3], Stephanie E. Mohr[2], Richard Benton[1,*]

[1]Center for Integrative Genomics
Faculty of Biology and Medicine
University of Lausanne
CH-1015
Lausanne
Switzerland

[2]Department of Genetics
Blavatnik Institute
Harvard Medical School
Boston
MA 02115
USA

[3]Howard Hughes Medical Institute
Boston
MA 02115
USA

[4]Institut du Cerveau-Paris Brain Institute
Sorbonne Université
Inserm
CNRS
Hôpital Pitié-Salpêtrière
75013
Paris
France

*Corresponding author:
E: richard.benton@unil.ch
T: ++41(0)21 692 3932

Short title: Orthogonal selection screens for toxin resistance genes

Abstract

Understanding toxin resistance in insects is key to appreciate niche adaptations, but remains challenging due to its often complex genetic basis. A well-known example is the specialized association of *Drosophila sechellia* with noni fruit (*Morinda citrifolia*), which is toxic to most other insects, including the closely-related drosophilids *D. simulans* and *D. melanogaster*. Noni toxicity is due to its high concentration of octanoic acid (OA), but the mechanisms that determine sensitivity or resistance to OA remain poorly understood. Here, we experimentally-evolved *D. simulans* with increased OA resistance, identifying multiple loci under selection. Cross-referencing these with a genome-wide, OA-resistance CRISPR screen in a *D. melanogaster* cell line highlighted two proteins: Kraken, a putative detoxification enzyme expressed in digestive and renal tissues, and Alkbh7, a mitochondrial protein linked to fatty acid metabolism. Both genes show elevated expression in *D. sechellia* and OA-resistant *D. simulans*. In *D. melanogaster*, *kraken* mutants are more OA-sensitive, while *Alkbh7* overexpression increased OA resistance. Importantly, mutation of these genes in *D. sechellia* reduced OA tolerance. Our identification of genes underlying OA resistance in laboratory and natural contexts demonstrates how complementary, cross-species selection approaches can provide insights into toxin susceptibility and adaptation mechanisms, with practical applications in the characterization of novel insecticides.

Introduction

How insects resist toxic chemicals in the environment is of both basic scientific interest – reflecting insects' arms race with plant food sources [1, 2] or venomous predators [3] – and applied interest, because humans have long used chemicals to combat insect disease vectors and agricultural pests [4]. The best-understood insecticides are those that bind to and modulate neuronal ion channels, where resistance arises through changes in these target proteins [5, 6]. For example, dichlorodiphenyltrichloroethane (DDT) and pyrethroids prevent closure of voltage-gated sodium channels, resulting in neuronal excitotoxicity and organismal paralysis; diverse insect species displaying resistance for these chemicals have independently-arising mutations in genes encoding these channels [7, 8]. Similarly, neonicotinoids bind to and hyperactivate nicotinic acetylcholine receptors, leading to neuronal death; mutations in various receptor subunits can confer resistance [9]. Even in such well-studied examples, however, it is thought that other neuronal targets exist [10] and/or that the chemicals impact receptors in non-neuronal tissues [11]. For many insecticides – as well as other environmental pollutants [12] – the mode of action is incompletely, or not at all, characterized [6].

Resistance observed in nature is only partially understood, reflecting the contribution of many genes beyond those encoding target proteins, which can underlie distinct defense mechanisms [13-15]. For example, thickening of the cuticle to reduce entry of insecticides into the body is widely considered an effective resistance mechanism [16, 17]. Cuticular changes have been linked to overexpression of proteins implicated in exoskeleton development (e.g., cuticle proteins, ABC transporters) but their precise roles, and how they are upregulated, remain unclear [17]. Increased expression of metabolic enzymes that degrade insecticides is another commonly observed adaptation in resistant strains, sometimes resulting from copy-number increases in the corresponding genes [18]. Duplicated metabolic genes can also undergo neofunctionalization to produce enzymes with altered specificity, as

illustrated by Cytochrome P450s in strains of the brown planthopper that recently evolved resistance to the neonicotinoid imidacloprid [19], as well as by more ancient evolution of glutathione S-transferases during transition of the *Scaptomyza* genus of Drosophilidae to herbivory to counter toxic isothiocyanates [20]. However, the lack of experimental resources in many insect species renders it difficult to characterize the causal contribution of specific genetic changes.

A compelling natural model clade study the evolution of toxin resistance is the trio of closely-related drosophilid species, *Drosophila melanogaster*, *Drosophila simulans* and *Drosophila sechellia* (Fig. 1A). *D. sechellia* is an extreme specialist on the toxic noni fruit of the *Morinda citrifolia* shrub [21, 22]. Amongst *D. sechellia*'s numerous adaptations [22-27], this species has evolved high resistance to octanoic acid (OA), an abundant noni chemical that is toxic to other drosophilids and more divergent insects [22, 28-33]. Previous pioneering studies of *D. sechellia*'s resistance to OA has been studied through genomic mapping-based approaches in *D. sechellia*/*D. simulans* hybrids, which revealed this trait to be polygenic, comprising at least five genomic regions [28, 34-36]. One of the main effect loci encompasses a cluster of 18 genes including members of the *Osiris* (*Osi*) family, which are thought to function in cuticle patterning [37, 38]. RNA interference (RNAi) knockdown of some *Osi* genes in *D. melanogaster* led to reduced tolerance of OA [39, 40]. However, genetic evidence for the contribution of these genes to OA resistance in *D. sechellia* is lacking.

Given the challenges of identifying candidate genes through genomic mapping-based strategies, in this work we took two complementary approaches to investigate the evolution of OA resistance in this drosophilid trio: experimental evolution in *D. sechellia*'s closest relative, *D. simulans*, through an Evolve-and-Resequence strategy [41], and CRISPR/Cas9-based genome-wide genetic screening in cultured *D. melanogaster* cells [42]. We show how the intersection of these approaches can identify novel genes contributing to OA susceptibility/resistance in these drosophilids, establishing these methods and this species clade as a powerful paradigm to investigate ecotoxicology.

Results

Experimental evolution of OA resistance in *D. simulans*

A previous study demonstrated that *D. simulans* strains can be experimentally evolved for higher OA resistance, enabling the identification of genomic regions (through microsatellite markers) responding to selection [43]. This success inspired us to take an Evolve-and-Resequence approach, which uses high-throughput sequencing to analyze the entire genome of populations before and after exposure to specific selective pressures, thereby facilitating the identification of adaptive loci from standing genetic variation in controlled and replicable settings [41, 44-47]. For studies of insecticide resistance, it has been predominantly used to examine changes in frequency of alleles of known genes rather than as a means to identify novel loci [48, 49].

The susceptibility/resistance of drosophilids to OA can be easily assessed by exposing flies to the toxin in a culture tube and counting the survivors after 24 h (Fig. 1B). In line with previous observations [34, 43], we found that a dose of 3 μl of pure OA (mixed with 1 ml glucose solution) leads to death of ~60% of *D. simulans* but has no effect on *D. sechellia* (Fig. 1C). By collecting the *D. simulans* survivors, we

reasoned that we could allow these to mate and establish the next generation (Fig.
1B), thus enriching genetic variants conferring higher resistance, similarly to [43].
Repeating the process over multiple generations and multiple replicates would allow
us to track phenotypic changes in resistance and investigate the associated
trajectories in allele frequencies.

Standing genetic variation should be present in the evolving populations for
natural selection to act upon [50, 51]. As the spontaneous mutation rate in drosophilids
is too low for artificial selection [52], we generated ten independent polymorphic base
populations of *D. simulans* from 93 isogenic strains [53] by round-robin crossing (Fig.
1D). These lines contain millions of SNPs, but we note that they were sampled from a
single wild population in California, therefore reflecting the genetic variation present in
that locality. The base populations therefore might not include all variants found across
the species' range, including those in regions where *D. sechellia* co-occurs. To
minimize bottleneck effects, each population was kept at a size of ~1000 flies. At each
generation, ~500 flies from each population were randomly collected and selected for
resistance to 3 μl OA. Over the course of 25 generations, all the populations showed
increased levels of survivability, suggesting parallel adaptive evolution (Fig. 1E). Given
the consistent responses across all populations, we also analyzed the phenotypic
changes grouping all replicates, finding that the mean survivability significantly
increased from ~43% at generation 0 to ~78% at generation 25 (Fig. 1F).

As we kept the same amount of OA throughout the 25 generations of selection,
the selective pressure lowered as the populations' survivability increased, evidenced
as a larger fraction of flies surviving and reproducing. Viewing the phenotype of
populations both individually (Fig. 1E) and collectively (Fig. 1F), it was apparent that
the change in phenotype plateaued at around generation 15. For continued evolution,
we therefore gradually increased the amount of OA during generations 26-31 (Fig.
1G), reaching a selection pressure comparable to the initial phase of the experiment
with 9 μl OA (~38% mean survivability across all populations) (Fig. 1G). We then
performed a second round of selection at this dose, observing an increase in
survivability across all populations, reaching ~70% by generation 50 (Fig. 1H). As the
populations adapted to the new condition, a second plateau started around generation
40 (Fig. 1H-I). Over 50 generations we were therefore able to evolve *D. simulans* to
have approximately 5-fold higher survivability to 9 μl OA compared to the unevolved
populations (Fig. 1J), though still falling short of the survival exhibited by *D. sechellia*
(Fig. 1J). These plateaus can be explained by the decreasing selective pressure
exerted by fixed OA volumes as an ever greater proportion of flies adapt and survive
the treatment. Determining whether an upper limit exists that would constrain the
evolution of a *D. sechellia*-like level of phenotypic resistance would require additional
selection rounds using progressively higher OA concentrations over an extended (and
unknown) number of generations. OA is only one component of noni fruit, although
probably the major toxic chemical [31]. We therefore tested three of our evolved
populations for resistance to noni fruit pulp, finding that they exhibited much higher
survival on this substrate compared to the same original populations maintained under
neutral conditions (i.e., no OA selection) (Fig. 1K).

Specialization of *D. sechellia* on toxic noni is associated with a reduced egg-
laying capacity [54, 55] and a shorter (albeit strain-specific) lifespan [26, 56, 57]. Our
success in evolving *D. simulans* with a partial *D. sechellia*-like resistance trait begs the
question as to whether these selected strains exhibit other phenotypic changes as a
trade-off for higher toxin resistance. We tested whether the evolved resistant
populations of *D. simulans* exhibit similar phenotypic changes in control conditions.

However, none of the three focal populations tested exhibited significant decreases in
the number of eggs laid (Fig. 1L), developmental viability (Fig. 1M) or lifespan (Fig.
1N). Thus, at least within the timeframe of our experimental evolution experiment, *D.*
simulans did not display obvious phenotypic trade-offs for the gain in OA resistance.
204
205　**Dynamic directional selection of allele frequencies**
206
We hypothesized that the observed phenotypic changes in *D. simulans* depended on
the directional selection of alleles of consistent loci across the ten populations, while
most of the genome evolves under neutrality. Allele frequencies at the selected loci
should therefore consistently change across all populations, allowing us to statistically
distinguish them from randomly evolving loci (Fig. 2A). We sequenced the genomes
of the populations of evolved (generation 25 and generation 50) and unevolved
(generation 0) flies, identifying 5,127,840 high-quality variants distributed across all
chromosomes. Pairwise identity-by-state analysis indicated that as the number of
generations increased, the proportions of shared alleles decreased, both between time
points and between pairs of populations (left-to-right and top-to-bottom, respectively,
in Fig. 2B).
To identify the selected genetic variants underlying the phenotypic adaptation,
we analyzed the genome-wide allele frequency changes both between generations 0
and 25, and between generations 0 and 50, across all populations. The identification
of significant changes in frequency is non-trivial, and many methods have been
developed [58-60]. We therefore assessed the significance of frequency changes
using two complementary approaches: the Cochran–Mantel–Haenszel (CMH) test
[61, 62], a widely-used method that tests for allele frequency changes between two
time points, and Bait-ER [60], a Bayesian method that tests for selection while explicitly
modelling genetic drift. Both tests identified large portions of the genome under
positive selection at generation 25 and 50 (Fig. 2C,D). The majority of prominent peaks
were found by both methods and across time points (Fig. 2E).
Based on the analysis of genome-wide linkage disequilibrium decay (Fig. S1A),
we grouped the significant SNPs identified by each of the two methods into genomic
blocks according to their proximity (i.e., <50 kb). We retrieved all the candidate genes
that fall within the block intervals, obtaining a total of 439 and 401 genes for the CMH
and Bait-ER, respectively, at generation 25, and 1238 and 174 genes at generation
50 (Data S3). These genes likely include true targets of selection but also neighboring
genes affected by hitchhiking. The increase in CMH candidate genes over time – in
contrast to the decrease observed with Bait-ER – might reflect the CMH test's
tendency to overestimate significance and to be insensitive to divergence among
replicate populations.
The overlap of all significant genes between the two time points is <30%. This
might reflect that a large fraction of the candidate genes results from noise caused by
genetic drift and draft (hitchhiking) and/or reflect a shift in the phenotypic contribution
of genes under the stronger selection (i.e., the higher dose of OA) during generations
27-50. Moreover, previous work on *Drosophila* using temporal sampling showed that
adaptation involving complex traits can lead to non-overlapping sets of selected SNPs,
in contrast with the simple expectation of continuous increases in allele frequency until
fixation [41, 63]. Gene ontology analysis is complicated at this stage as most hits are
likely hitchhikers of the true causal genes. It is also hard to compare currently to
previous mapping studies in *D. simulans/D. sechellia* hybrids [34-36] or in the earlier
D. simulans experimental evolution experiment [43], which only identified wide

genomic regions. The sole exception, a 170-kb window on 3R containing only 18 genes (including several *Osiris* genes) [64] did not emerge in our analysis.

A genome-wide, pooled CRISPR screen for OA susceptibility and resistance genes in *D. melanogaster* S2R+ cells

Given that these previous studies also contain extensive hitchhiking signals and substantial noise across large genomic regions, comparing their results with our analyses would likely yield numerous false positives and still leave a large pool of putative loci, offering limited benefit for fine-mapping. We therefore sought an orthogonal screening approach. Motivated by previous observations of OA toxicity for insect Sf9 cultured cells [33], we found that OA also leads to reduced viability of *D. melanogaster* S2R+ cells (Fig. 3A). This phenotype prompted us to perform a selection experiment in this cell line through exploitation of a genome-wide, pooled CRISPR/Cas9 mutagenesis approach (Fig. 3B) [42]. In brief, a library of sgRNA-encoding plasmids (6-8 sgRNAs/gene) was transfected into Cas9-expressing cells where site-specific plasmid integration predominantly limits targeting to a single gene per cell. Two replicate cultures of this pool of cells were either left untreated, or exposed to OA in the medium, at two different concentrations (Fig. 3B). After 26 days in culture, DNA was extracted from the cell pool, then sgRNA sequences were PCR amplified and sequenced to determine those that were enriched after OA selection (reflecting genes whose loss enhances cell viability, referred to hereafter as "susceptibility genes") or depleted (reflecting genes whose loss leads to lower OA resistance, hereafter "resistance genes").

To identify candidate OA susceptibility/resistance genes, we first filtered out genes essential for general cell survival [42] and genes that were not detectably expressed in S2R+ cells (using DGET [65]. (Removal of such genes are inevitable limitations of this cell-based screening approach; see Discussion). After comparing results from each condition and replicate, we identified the top 5% susceptibility genes (Fig. 3C) and top 5% resistance genes (Fig. 3D). Amongst the 87 susceptibility genes, there was no single stand-out hit, suggesting that – at least in S2R+ cells – OA does not have a single major cellular target (unlike, for example, insecticidal bacterial toxins that bind a specific cell surface receptor [66]). Gene-ontology analysis of these susceptibility genes revealed a significant enrichment of genes involved membrane trafficking (Fig. 3C). As OA has previously been hypothesized to disrupt cellular membranes [33, 67], the recovery of these hits suggests that general reduction in membrane transport reduces OA uptake into cells and/or renders membranes less susceptible to OA treatment. 22 OA resistance genes were identified, encoding diverse types of proteins (Fig. 3D), suggesting there are multiple mechanisms by which cells can resist the toxic effects of OA.

We next examined which of the genes identified in the S2R+ cell selection experiments might be relevant for evolution of OA resistance in whole animals by determining those also present within significant blocks of the experimental evolution analyses (Fig. 3E,F). Four susceptibility genes were identified: CG1620 (a predicted SANT-MYB domain transcription factor), CG4291 (a putative component of the splicing machinery), *Nph* (a chromatin remodeling factor [68]) and *Debcl* (a pro-apoptotic member of the Bcl-2 family [69]); these are all likely to have indirect roles in conferring susceptibility to OA given their broad functions. By contrast, two resistance genes have potentially direct roles in conferring tolerance to OA, *kraken* and CG14130. *kraken* encodes a putative serine hydrolase [70] and stood out both

because it was the top hit in the resistance gene screen (Fig. 3D), and because of its location in one of the most prominent peaks in the experimental evolution at generations 25 and 50 (Fig. 3F and Fig. 2C-D). CG14130 encodes the orthologue of mammalian *Alkbh7*, which is involved in regulating fatty acid metabolism [71, 72], and is located within a significant peak at generation 25 (Fig. 3F and Fig. 2C). Therefore, intersecting the experimental evolution and CRISPR screening results allowed us to narrow down tens or hundreds of potential candidates to two focal genes for downstream investigations.

Sequence, expression and evolutionary analyses of candidate OA resistance genes

To assess the candidacy of *kraken* and CG14130 (hereafter *Alkbh7*) in contributing to OA resistance in animals, we first examined their sequence and expression in *D. melanogaster*. Kraken belongs to a large family of serine hydrolases, which encompass lipases, esterases and proteases that use a nucleophilic serine residue in the active site for hydrolysis of substrates [70]. Consistent with the conservation of this serine in Kraken (Fig. 4A and Fig. S2A), an activity-based proteomic screen of the serine hydrolase superfamily in *D. melanogaster* provided evidence for *in vivo* catalytic activity of Kraken [73]. *kraken* is broadly expressed in adults, with particular enrichment in the gut and Malpighian tubules (Fig. 4B), similar to previous observations of *kraken* expression in the embryo by RNA *in situ* hybridization [70]. Using the Fly Cell Atlas [74], we examined sub-tissue adult expression of *kraken*: in the Malpighian tubules, *kraken* is expressed at highest levels in the principal cells (Fig. 4C), which are involved in ion transport and detoxification. In the gut, *kraken* transcripts are most robustly detected in various types of enterocytes, which secrete digestive enzymes [75] (Fig. 4C). However, Kraken lacks an N-terminal signal sequence [76], suggesting that the protein remains in the cytoplasm of these cells, unlike canonical digestive enzymes [75], and more similar to detoxification enzymes of the Cytochrome P450 and Glutathione S-transferase families [77]. To examine *kraken* expression *in situ*, we generated a *kraken* promoter-Gal4 driver line, which induced expression of a fluorescent reporter in cells of the proventriculus, the crop duct and the hindgut (Fig. 4D). No GFP expression was detected in Malpighian tubules (data not shown), which we suspect reflects the lack of necessary regulatory sequences in the driver transgene.

Alkbh7 is a member of the alpha-ketoglutarate-dependent hydroxylase (Alkbh) family (Fig. 4A), which are enzymes that act on diverse substrates in processes such as biosynthesis, metabolism and epigenetic regulation [78]. As in mammals [72], *D. melanogaster Alkbh7* displays broad tissue expression (Fig. 4B). Murine Alkbh7 is localized to the mitochondrial matrix [72]. This property appears to be conserved in the *Drosophila* protein, which has a mitochondrial targeting sequence (Fig. 4A and Fig. S2B) [79]; consistently, we identified Alkbh7 within *D. melanogaster* proteomes of the mitochondrial matrix [80, 81].

Although the sequences of both Kraken and Alkbh7 are overall highly similar across *D. melanogaster*, *D. simulans* and *D. sechellia* (Fig. S2A,B), we wondered whether we could detect signs of selection at these loci in *D. sechellia* employing population sequence datasets of *D. sechellia* and *D. simulans* [82]. Using the McDonald-Kreitman test [83] on the coding sequences of both genes, we found evidence for gene-level positive selection on *kraken* but not *Alkbh7* (Fig. S2C). Because selection might act pervasively at individual sites rather than along the entire

protein coding region, we also implemented a codon-based analysis using FUBAR
(Fast, Unconstrained Bayesian AppRoximation) [84], which identified 1 and 4 codons
under diversifying positive selection in *kraken* and *Alkbh7*, respectively (Fig. S2D).
Although their functional relevance remains unclear, such mutations might reflect
underlying evolutionary dynamics at the protein level.
We next compared the expression levels of these genes both in laboratory
strains of *D. sechellia* and *D. simulans* (Fig. 4E), as well as in RNA-seq datasets of
various strains of *D. sechellia*, *D. simulans* and *D. melanogaster* [85-87] (Fig. 4F, Fig.
S3A,B). While there were some exceptions, overall both *kraken* and *Alkbh7* show
higher expression levels in *D. sechellia* adults and larvae compared to the other
species. Importantly, in our experimentally-evolved *D. simulans* populations, we also
found that both *kraken* and *Alkbh7* were more highly expressed compared to
unselected populations (Fig. 4G). The expression differences between species, and
between unevolved and evolved *D. simulans*, reflect variation in basal levels of
expression, as prior exposure to OA did not lead to changes in expression in either *D.*
sechellia or *D. simulans* (Fig. S3C).
366
367 **Functional contributions of *kraken* and *Alkbh7* to OA resistance**
368
Given the expression differences observed in the naturally and experimentally evolved
OA-resistant drosophilids, we asked whether artificially modulating gene expression
would impact OA tolerance. We first performed ubiquitous RNAi of these genes in *D.*
melanogaster (Fig. S4A) and assessed the effects on resistance to OA using a short-
term, plate-based assay [28] (see Methods). *kraken*[RNAi] led to significantly diminished
OA resistance compared to genetic controls, while *Alkbh7*[RNAi] did not have a
discernable effect (Fig. 5A). We next generated deletion mutants for these genes in
D. melanogaster (Fig. S4B). Similar to the RNAi experiments, *kraken* mutants, but not
Alkbh7 mutants, displayed reduced resistance to OA (Fig. 5B). All genotypes remained
fully viable when no OA was added (Fig. S5).
Loss-of-function analyses might fail to uncover a role for *Alkbh7* in whole
animals if there are redundant resistance mechanisms absent from S2R+ cells. We
therefore took a complementary approach by overexpressing these genes in *D.*
melanogaster. Strikingly, overexpression of *Alkbh7* resulted in survivability of *D.*
melanogaster to levels close to those observed for *D. sechellia* under the same
experimental conditions (Fig. 5C, Fig. S4C), indicating that this gene is sufficient to
confer high-level OA resistance. By contrast, *kraken* overexpression did not have any
detectable impact on OA resistance (Fig. 5C); a similar lack of effect was seen when
overexpressing *D. sechellia kraken* (Fig. S2E). The absence of a phenotypic effect
upon *kraken* overexpression is surprising, given its elevated expression in *D. sechellia*
and in evolved *D. simulans* but this could reflect either insufficient overexpression
levels (Extended Data Fig. 4c) and/or genotype-by-genotype interactions that are not
recapitulated when manipulating a single locus in a *D. melanogaster* background.
A key question is whether these genes contribute to OA resistance of *D.*
sechellia. Implementing CRISPR/Cas9-based genome-engineering in this species
[22, 27], we generated and validated null mutants in these genes in *D. sechellia* (Fig.
S4B) and tested their resistance to OA in the plate assay. Although we did not detect
any significant difference to genetically-matched wild-type flies (Fig. 5D), this was
anticipated given the polygenic nature of OA resistance in *D. sechellia* [28, 34-36], and
the modest or absent phenotypes of equivalent mutants in *D. melanogaster* (Fig. 5B).
We reasoned that more subtle differences might be detected by monitoring *D.*

sechellia's survivability over longer periods of time, when repeated contact with OA was enforced by blending it with the food. Indeed, we observed higher mortality rates of both *kraken* and *Alkbh7* mutants compared to genetically-matched wild-type flies over a period of 6 days (Fig. 5E). By contrast, both mutants survived equally well as controls in the absence of OA over this time period (Fig. 5E). These data provide the first evidence for the contribution of specific genes to OA resistance in *D. sechellia*. However, future allelic swap studies with *D. simulans* or *D. melanogaster* would be required to assess if and how they contribute to species-specific differences in OA sensitivity (e.g., through structural and/or regulatory differences in the genes).

Discussion

Identifying specific loci underlying phenotypic evolution is challenging due to the polygenicity of the vast majority of adaptive traits. Investigating the evolution of chemical toxicity is an attractive model trait because of the relatively easy phenotypic read-out. To gain insight into the genetic basis of susceptibility and resistance to a natural toxin, OA, we have intersected the results of selection over vastly different timescales: weeks in cell lines, years in experimentally-evolved laboratory strains, and millennia during speciation.

Unlike insecticides that activate or antagonize broadly-expressed neuronal ion channels to cause organismal death, the mode of action of OA appears to be mechanistically very different, capable of inducing death at the level both of whole organisms and of cultured cells. Beyond insect cell lines (our work and [33]), OA is also toxic to budding yeast [88] and bacteria [89, 90], suggesting that this toxin acts via a more fundamental cellular mechanism. Indeed, a genome-wide, knock-out screen in yeast for genes required for tolerance of OA highlighted contributions of several core cellular processes, such as vesicle trafficking, mitochondrial function or chromatin regulation [88], reminiscent of our CRISPR screen results. Consistent with this idea, the results of our experimental evolution and CRISPR screening, as well as earlier mapping efforts in *D. sechellia*/*D. simulans* hybrids [28, 34-36] revealed multiple genomic regions/genes contributing to the resistance to OA.

Considered separately, the experimental evolution and cell-based CRISPR screens identified many candidates. That there was very limited overlap between the screen results is unsurprising given their very different designs. Notably, the loss-of-function, cell-based screen cannot identify genes that are not expressed in the cell line, nor those essential for cell viability in the absence of OA. Nevertheless, both lists are likely to contain multiple genes relevant to understanding OA susceptibility and resistance, and hence mode(s) of action of OA. Here, we focused on two candidates, *kraken* and *Alkbh7*, which were common hits of both screens. We note, however, that neither of these genes were identified in previous studies of OA resistance [28, 34-36], which could reflect several factors, such as differences in OA toxicity assay conditions, developmental stages, species and molecular readouts (DNA or RNA), and the evident multilayered, polygenic nature of this trait (discussed further in [91]).

The (partial) necessity, but insufficiency, of *kraken*, and lack of necessity but sufficiency of *Alkbh7* to confer OA resistance implies fundamentally different roles of these proteins. The expression of *kraken* in the gut and renal system supports a hypothesis of this protein as a detoxification enzyme, although future work will be necessary to determine whether Kraken has a direct role in OA degradation. The mild loss-of-function phenotype of *kraken* in both *D. melanogaster* and *D. sechellia* likely reflects substantial redundancy of defense mechanisms, including other potential

detoxification mechanisms, especially given the plethora of enzymes expressed in the fly gut and Malpighian tubules [75, 92]. Indeed, extremely few associations between such enzymes and substrates and/or physiological roles have previously been defined. One exception is the Cytochrome P450 gene *Cyp6g1*, which has a key role in Malpighian tubules in conferring resistance to DDT [77]. Our data on Kraken therefore provide one of the first links between a natural toxic chemical and a putative detoxification enzyme.

The mitochondrial Alkbh7 enzyme likely has a distinct role in OA resistance. Insights from mammals are informative: mice lacking *Alkbh7* are obese, with higher body fat levels, proposed to result from a defect in metabolism of fatty acids [72]. Molecularly, one study suggests that Alkbh7's metabolic role stems from its regulation of RNA editing to control mitochondrial protein translation [71], although it is unclear whether this is its only function. While the *Alkbh7* loss-of-function phenotype is mild in drosophilids, it is striking that artificially elevated expression in *D. melanogaster* is sufficient to produce *D. sechellia*-like levels of OA resistance. This result implies a potent effect of Alkbh7 on the mitochondrial metabolic network to modulate toxin resistance, perhaps through control of fatty acid metabolism; such ideas can be tested in future studies. Intriguingly, our implication of Alkbh7 in the niche adaptation of *D. sechellia* has a potential parallel in bats, where this gene displays evidence of positive selection during dietary diversification of different species [93].

Taken together, our work has demonstrated the power of orthogonal selection approaches in the laboratory to understand the evolution of toxin resistance, which might be relevant to understand a historic paradigm of this phenomenon [23], as well as offering clues as to how the natural toxin OA exerts its lethal effects. Moreover, the methodology – notably cell-based CRISPR screening, which can be extended to both gain-of-function approaches [94] and mosquito cell lines [95] – is applicable to explore the molecular mechanisms of susceptibility and resistance of insects to the wide-range of artificial toxic chemicals used in the environment [12]. Such knowledge will have application in the future characterization of novel types of insecticides.

Author contributions

M.M. and R.B. conceived the project. M.M. designed and performed the experimental evolution experiment, generated and analyzed all transgenic and mutant flies, and performed the comparative gene expression and molecular evolutionary analyses. S.C. generated the base populations and provided technical support for the experimental evolution experiment. M.M., R.B., R.V. and S.E.M. designed the cell screen. S.G. and R.V. tested and defined cell screen assay conditions. S.G. and M.B. performed the CRISPR/Cas9 screen and prepared DNA for sequencing. R.V. analyzed the screen data, and M.M., R.B., R.V., S.E.M and N.P. contributed to interpretation of screen results. J.D. and C.R. performed histological experiments, supervised by D.H. M.M. and R.B. wrote the paper with input from other authors. All authors approved the final version of the manuscript.

Acknowledgements

We thank Tadeusz Kawecki and Ana Marija Jakšić for advice on experimental evolution experiments, Tina Lenče and Jean-Yves Roignant for the gift of the *D. melanogaster Alkbh7* sgRNA transgenic line, the Bloomington *Drosophila* Stock Center (NIH P40OD018537) and the Vienna *Drosophila* Resource Center for *D.*

melanogaster stocks, Christian Schlötterer and Neda Barghi for the *D. simulans* lines,
Blaise Tissot-Dit-Sanfin for maintenance of *M. citrifolia* plants and Ambra Masuzzo for
providing noni fruits. Figure icons were created with Biorender.com. We are grateful
to Tadeusz Kawecki, Philippe Reymond, Michael Shahandeh and members of the
Benton laboratory for discussions throughout the project and comments on the
manuscript. D.H. is funded by an ERC Starting Grant (101117267). N.P. is an
Investigator of the Howard Hughes Medical Institute. Research in R.B.'s laboratory
was supported by the University of Lausanne, an ERC Advanced Grant (833548) and
the Swiss National Science Foundation (310030_219185).

509
510 **Declaration of interests**

511
512 The authors declare no competing interests.

513
514 **Data availability**

515
516 Source data is provided in the supplementary data files. All other data, new materials
517 and code are available from the corresponding author upon reasonable request.
518

Materials and Methods

Drosophila strains and husbandry

Drosophila stocks were maintained on standard wheat flour–yeast–fruit–juice medium under a 12 h light:12 h dark cycle at 25°C. Wild-type strains, as well as mutant and transgenic lines used in this study are listed in Table S1. Experimental evolution was performed using a subset of published *D. simulans* isogenic strains [53] (Table S2).

Generation of mutant and transgenic flies

Loss-of-function mutants: we generated *D. melanogaster* mutants for *kraken* and *Alkbh7* using in vivo CRISPR/Cas9 mutagenesis [96] by crossing flies expressing Cas9 in the germline (*nanos-Cas9*) to those expressing sgRNAs targeting the desired gene. For *kraken*, we used P{hsFLP}1,y^1,w^{1118};P{HD_CFD02326}attP40 as a source of sgRNAs driven by *Act5C-GAL4*; for *Alkbh7*, we used the CG14130 sgRNA line described in [97]. Mutants were validated by genomic PCR and sequencing (Fig. S4B).

D. sechellia mutant flies were generated by WellGenetics Inc. using modified methods of [98]. For *kraken* (GM16772/LOC6617307) the upstream sequence CTGTAATGCTGGACGCGGAT[TGG] (PAM in square brackets) and the downstream sequence CCTGGTCACCCCGGATCGAG[TGG] were cloned separately into a U6 promoter plasmid. The 3xP3-RFP cassette, which contains a floxed 3xP3-RFP flanked by a 5' homology arm (999 bp: from -1043 nt to -45 nt relative to the *kraken* ATG) and a 3' homology arm (1063 bp: +1388 nt to +2450 nt) were cloned into *pUC57-Kan* as a donor template for repair. For *Alkbh7* (GM25300/LOC6605344) the upstream sequences ACTAATAAGAGTCGACCGGG[TGG], AACAGTCAAACAGCTGTTCC[TGG] and downstream sequences GTCCCAGTTTCAGGGTACGC[TGG], ATGGAAATGCGACGTGTCCT[GGG] were cloned separately into a U6 promoter plasmid. The 3xP3-RFP cassette, which contains a floxed 3xP3-RFP, 5' homology arm (1037 bp: -1121 nt to -85 nt relative to the *Alkbh7* ATG) and 3' homology arm (964 bp: +940 nt to +1903 nt) were cloned into *pUC57-Kan*. DNA plasmids encoding the sgRNAs and the corresponding homology-directed repair template were microinjected into embryos of *Dsec.07 pBAC(nos:Cas9,3P3-YFP)* [27]. In both cases, a single F1 line expressing the 3xP3-RFP selection marker was obtained, which was validated by genomic PCR and sequencing (Fig. S4B).

D. melanogaster and D. sechellia mutant flies were backcrossed for six generations to Canton-S and *Dsec*07, respectively.

Transgenic flies: to make *kraken-Gal4*, we amplified putative 5' regulatory regions for *D. melanogaster kraken* (encompassing the entire intergenic region between the last exon of the upstream gene and the first exon of *kraken*) from w^{1118} genomic DNA (all oligonucleotides are listed in Table S6), and subcloned the PCR product into *pGal4-attB* [99]. To make the *UAS-Dmel\kraken* and *UAS-Dsec\kraken* transgenes, we amplified the regions from the start to the stop codons from w^{1118} and *Dsec07* genomic DNA and subcloned them into a *pUASTattB* vector [100]. Transgenic flies were obtained by phiC31 integrase-mediated transgenesis into attP2 performed by BestGene Inc. and verified by sequencing.

Toxin resistance assays

Tube assay: resistance to OA for the experimental evolution setup was scored adapting a previous protocol [34, 43]. In brief, flies were lightly anesthetized with CO_2 and ~50 animals were placed in standard cultured tubes (25 mm diameter × 95 mm height, Milian SA) and provided with 1 ml of a 3% glucose solution on a tissue paper (Kimtech, 7552). 3 μl of pure OA (Sigma-Aldrich, CAS 124-07-2, C2875) were directly pipetted onto the tissue while the flies remained anesthetized. Direct contact with the acid was scrupulously avoided until all flies recovered. The fraction of alive flies was scored after 24 h. For the experimental evolution experiment, the surviving flies were collected (see below).

Plate assay: to measure OA resistance in the genetically-manipulated *D. melanogaster* and *D. sechellia* strains, we used a plate assay [28]. Compared to the experiments in tubes, this assay offers greater temporal resolution and higher dynamic range (in contrast to the experimental evolution experiment, we did not strive to maintain a certain fraction of surviving flies), as well as greater statistical power from tests of fewer animals. Ten female flies were lightly anesthetized with CO_2 and placed in a Petri dish (60 mm diameter × 15 mm height) in which 2 μl of pure OA (or 30 μl for *D. sechellia*) were spotted in the center of the underside of the lid. The flies were allowed to recover (typically <5 min), and immobile/dead were scored every 5 min for 1 h. The survival curves were calculated and plotted using the R package survminer (10.32614/CRAN.package.survminer). The statistical analysis was conducted fitting a mixed effect Cox regression model as implemented in the R package coxme [101]. Genotype was included as a fixed effect, and replicate dishes were modelled as a random effect.

Noni resistance assay: noni fruit was collected from *M. citrifolia* plants grown in the University of Lausanne greenhouses. Five ripe (pale yellow) fruits were homogenized together, and the pulp puree frozen at -20°C. The noni assay was performed with the same setup as for tube assay described above, using 2 g of thawed noni puree placed on the bottom of the tube. Noni contains 3.06 g OA/kg fruit [102]; assuming an equivalent quantity in our fruits, 2 g noni pulp should contain 6.73 μl OA, approximately double the amount used for the experimental evolution between generations 1 to 25, and two thirds of the concentration during generations 30-50.

Long-term assay: To assess the long-term effects of OA (i.e., beyond 24 h), two-day-old mated female flies were individually housed in vials containing food supplemented with OA. OA was thoroughly mixed into Formula 4-24® Instant Drosophila Medium, Blue (Carolina Biological Supply) to a final concentration of 0.9%. Fresh OA-containing food was provided every two days, and mortality was monitored daily until all flies in the experimental group died. Statistical analysis was run as for the plate assay.

Fecundity and *developmental viability*: 10 mated adult female flies (4-7 days old) were allowed to lay eggs for 24 h on agar plates (2% agar, 2.5% sucrose in a 1:3 grape juice:water mix). The total number of eggs on each plate was counted. From each plate, 100 eggs were collected and transferred to standard fly food. Emerging adults were counted and removed daily until no further flies emerged.

Lifespan: Two-day-old mated female flies were individually housed in vials containing standard fly food and monitored daily until death. Fresh food was provided every 2–3 days.

Experimental evolution

Selection procedure: to generate the base populations for experimental evolution, we used 93 isogenic lines (Table S2) of the *D. simulans* panel established by [53]. We first crossed each of the lines together following a round-robin design to ensure equal contribution of each line to the starting genetic pool; 5 males and 5 females from each cross were then combined in ten independent replicates (Fig. S1D). These ten populations were maintained in bottles (57 mm diameter ×103 mm height, Milian SA) of standard fly food at high population size (~1000 individuals per population) to minimize the effect of genetic drift. Each population was subdivided into smaller groups of ~250 individuals per bottle (i.e., four bottles per population) to reduce competition. At each generation, individuals from the four bottles were mixed to prevent bottle-specific genetic drift. The experimental evolution experiment was initiated 10 generations after the creation of these populations.

From each population, 500 individuals (2-7 day old) were randomly collected and further split into 10 groups of 50 mixed-sex flies (i.e., ~5000 flies total). Each group of 50 flies was then placed under selection with 3 µl of OA for 24 h (as described above); the survivors were collected and allowed to breed in food vials for 6-8 days, transferring them to fresh vials every 2-3 days. In the next generation, 500 flies per population were collected only from the last seeded vial, to maximize the probability of these flies being parented by the OA-resistant males and females and not offspring generated by females that mated before the selection procedure. These flies were then subjected to the same selection procedure. This process was repeated for 25 generations, and survivability was scored at each generation.

A second round of experimental evolution was performed by increasing the amount of OA to 9 µl between generations 31 and 50; this amount was determined empirically during generation 26-31 to re-establish the survivability close to the initial level (Fig. 1G). All successive generations were maintained following the same procedure.

As a control, the ten original populations were maintained in parallel in the same conditions as previously described (i.e. ~1000 individuals per population subdivided into smaller groups of ~250 individuals per bottle).

DNA sequencing and analysis: we pooled ~200 individual flies per population per time point (generations 0, 25 and 50). Genomic DNA (and total RNA for qPCR, as described below) extraction was performed using the Quick DNA/RNA Miniprep plus kit (Zymo Research, D7003) following the manufacturer's instructions. Samples were stored at -80°C.

We performed paired-end whole genome sequencing on an Illumina NovaSeq 6000 following Nextera DNA flex preparation, at an average coverage of >200×. Read quality was assessed with Fastqc [103] and adapters were trimmed with Trimmomatic [104] providing the Nextera adapters sequences. We aligned the trimmed reads to the *D. simulans* reference genome (NCBI GCF_016746395.2) using bwa-mem [105] and the results were sorted and compressed using samtools [106]. Duplicate reads were then removed with Picard MarkDuplicates (https://broadinstitute.github.io/picard/).

Unplaced scaffolds and alignments with mapping quality <20 were removed with samtools. We then combined all the aligned sequences into a samtools pileup file and split this into individual chromosomes. Regions with zero coverage were removed with custom scripts. Sync files for each chromosome were generated with the mpileup2sync.pl script implemented in Popoolation2 [107] and used as input files for all the Evolve-and-Resequence analyses.

To explore the genetic variation of our samples we generated vcf files with vcftools 0.1.17 [108] and binary files with plink1.9 [109, 110]. We calculated the identity-by-state (IBS) matrix as implemented in plink1.9 (function -distance 'ibs').

Evolve-and-Resequence analysis: allele frequency changes between generation 0 and generations 25 or 50 were investigated via Popoolation2 and Bait-ER [60].

We performed a Cochran–Mantel–Haenszel (CMH) test via the cmh-test.pl script implemented in Popoolation2 comparing base and evolved populations with parameters --min-count 12 and --min-coverage 50. We estimated a plausible significance threshold approximating the top 0.000005% of the simulated p-values based on the effect of genetic drift alone by running genome-wide forward simulations under neutrality in mimicree2 [111]. We used the same genomic information from the original 93 lines as input and the average real survivability ratios observed throughout the selection experiment as –selection-regime parameter and 500 as –population-size parameter (Fig. S1B).

For Bait-ER, the input sync files were filtered to match --min-count 12 and --min-coverage 50 and scaled to a uniform coverage of 100 as in [58] to avoid numerical problems due to high coverage. We ran the algorithm on each chromosome separately. Bait-ER models the effect of genetic drift directly using estimates of the effective population size (N_e). We calculated N_e estimates for each chromosome in each replicate population with the R package poolSeq [112] and used the median value of each chromosome as "Population_size" parameter (Table S3). Other parameters included "Sampling correction" = 0 (i.e., binomial) and "Prior parameters" =0.001 (shape), 0.001 (rate). Data points were considered significant when the absolute natural logarithm of the Bayes Factor exceeded the conservative threshold of log(0.99/0.01) as in [60].

The linkage disequilibrium decay in the 10 evolved populations was analysed across the whole genome using PopLDdecay [113]. Linkage disequilibrium maps of focal regions were generated with LDBlockShow [114] using the Solid Spine method based on D' statistics originally introduced in Haploview [115].

For the genome-wide screen, all the genes within selection blocks were considered potential candidates. Based on the results on linkage disequilibrium decay, a block was arbitrarily defined as a genomic interval where at least two significant SNPs were found at a distance less than 50 kb. Consecutive SNPs respecting this definition were grouped into larger blocks and visualized as circus plots using the R package circlize [116]. The genes overlapping such blocks were extracted using BEDtools intersect [117]. Orthologous genes in *D. melanogaster* were retrieved using a combination of the OMA browser [118], DAVID [119, 120], BLAST [121], NCBI [122] and FlyBase [123], and subsequent manual curation.

Genome-wide CRISPR/Cas9 screen in *D. melanogaster* S2R+ cells

Library design, construction, and transfection: design, synthesis and cloning of the genome-wide sgRNA library are previously described [124]. To determine sub-lethal doses of OA appropriate for screening, the effect of a gradient of OA supplementation into media on attP+ Cas9+ cells was determined using CellTiter-Glo (Promega) following the manufacturer's protocol; luminescence measurements were made using a Spectramax Paradigm (Molecular Devices).

For the screen, attP+ Cas9+ *D. melanogaster* S2R+ cells (RRID:CVCL_UD30) were transfected using Effectene and a 1:1 molar ratio of sgRNA library and pIntAC, as described [124]. Briefly, transfected cells were distributed in 5 ml aliquots across ten 100 mm culture dishes and incubated overnight, then 5 ml of fresh media was added to each plate. After 4 days, cells were subjected to puromycin selection for 12 days, subculturing every 4 days, to establish a stable sgRNA-expressing cell library. For the selection assay, 2.5×10^7 cells were plated onto four 15 cm diameter cell culture dishes to ensure sufficient sgRNA coverage, with each sgRNA represented approximately 1000 times [124].

OA selection assay: library-containing cells were left untreated or treated with either 0.075 µg/µl or 0.05 µg/µl OA (Sigma-Aldrich, CAS 124-07-2, C2875). Each replicate consisted of four 15 cm diameter cell culture dishes seeded with cells, achieving ~1000× coverage. Two replicate sets were included for each of the two OA treatment conditions and one replicate for the untreated control. During the selection period, cells were split 1-2 times per week to avoid overgrowth. An aliquot of cells from each replicate was collected and stored for genomic prep following each split and at the end of the assay. The final samples were collected after 26 days of OA or control treatment.

Genomic DNA extraction and sequencing: genomic DNA was extracted from samples collected during the screen and at the final endpoint using a Zymo Quick-gDNA Miniprep kit (Zymo Research, D3025). DNA fragments containing the sgRNA sequences were amplified by PCR using what we calculated to be at least 1000 genomes per sgRNA as templates for each sample. PCR fragments were in-line barcoded using a previously described approach [42, 124, 125]. Next-generation sequencing (Illumina NextSeq) was performed at the Biopolymers Facility at Harvard Medical School (RRID:SCR_007175). FASTQ sequence files were demultiplexed using in-line barcodes as sequence tags as described previously [42, 124]. All further processing was done using MAGeCK (v.0.5.4 and 0.5.9.4) [126] to generate readcount files and perform robust rank aggregation (RRA) analysis and Maximum likelihood estimation directly on the generated readcount files using default parameters. Plasmid readcounts were used to normalize RRA values.

After removing essential genes, we considered as candidates the top 5% genes overlapping across the two conditions that were over-represented or under-represented in the OA-exposed cell populations compared to the control cells (i.e., representing "susceptibility" and "resistance" genes, respectively). The gene-set enrichment analysis was performed with PANGEA 1.1 Beta using the *Drosophila* GO subsets (GO slim) and a p-value cut-off of 0.05 [127].

qPCR analysis

Reverse transcription of total RNA (extracted as described above) was performed with the PrimeScript first-strand cDNA synthesis kit (Takara, 6110A) following the manufacturer's instructions. Samples were stored at -80°C.

Quantitative PCR experiments were conducted on a QuantStudio 6 Flex Real-Time PCR System using the PowerUp™ SYBR™ Green Master Mix following the manufacturer's instructions. Each sample consisted of at least three biological replicates. We performed three technical replicates per reaction and removed replicates where the standard deviation was equal or greater than 0.5. We determined target specific amplification efficiencies for each gene (Table S4, Table S6). Relative quantification and statistical analysis were performed with the software qbase+ v3.4 [128] using Calibrated Normalized Relative Quantities (CNRQs). Normalization was performed on the geometric mean of the relative quantities for the three most commonly-used reference targets: *Act5C*, *αTub84B* and *RpL32* [129]. We assessed reference target stability by the geNorm expression stability value (M) and the coefficient of variation of the normalized reference gene relative quantities (CV) (Table S5) [128, 130].

Population genetics analysis

Publicly available WGS data for 41 *D. sechellia* and 20 *D. simulans* genomes were downloaded from the Sequence Read Archive (PRJNA215932 and PRJNA395473) [82, 131] with fasterq-dump [132]. FASTQ files were processed and aligned as described in the "Experimental evolution" section. Aligned files were phased with samtools phase [106] and the genes of interest were extracted with pilon [133] to find variation among populations. CDS sequences were aligned based on amino acid information using translatorX [134]. Gene sequences were aligned using Clustal Omega [135]. McDonald-Kreitman test, Tajima's D and Fu and Li's statistics were calculated in DNAsp 6 [136]. FUBAR analyses (Fast, Unconstrained Bayesian AppRoximation) were performed in HyPhy 2.5 [137]. The consensus sequences of *D. simulans* and *D. sechellia* were computed by SnapGene (www.snapgene.com) "export consensus" with a >50% threshold. The sequence alignment and secondary structures were depicted using ESPript 3 [138].

Histology

Intestinal tissues and Malpighian tubules were dissected in phosphate-buffered saline (PBS) and fixed at room temperature for 45 min in PBS, 4% paraformaldehyde. Staining of muscle was performed using a mushroom phalloidin AlexFluor647 probe (Life Technologies, A22287, diluted 1:200 in PBS + 0.2% Triton X-100) for 45 min. Samples were mounted in Vectashield with DAPI (Vector Labs, H-1200-10).

A Leica STELLARIS-DIVE confocal microscope was used to generate all confocal images, using a water-immersive 40× objective or an oil-immersive 63× objective. The images were acquired using both HyDs as well as PMTs tailored for the fluorophores of each sample.

Statistics and reproducibility

Data were analysed and plotted in R 4.0.3 (R Core Team 2021).

Figure legends

Fig. 1. Experimental evolution of *D. simulans* with increased OA resistance.
A Phylogeny of the drosophilid trio studied in this work. Mya: million years ago.
B Schematic of octanoic acid (OA) resistance tube assay (see Materials and Methods) and selection procedure.
C Species differences in resistance of *D. simulans* 04 and *D. sechellia* 07 strains to 3 µl OA.
D Schematic illustrating the generation of the base populations for Evolve-and-Resequence (see Materials and Methods).
E Changes in survivability to 3 µl OA of the ten *D. simulans* populations over generations 0-25. Box plots represent the interquartile ranges of the survivability for all the replicates; blue lines shows the generalized additive model (GAM) regression; grey shading represents the 95% confidence interval for the mean responses at each time point.
F As in **E** for the 10 populations combined.
G OA resistance for the combined 10 populations with increasing OA volumes during generation 25-31.
H As in **E**, with 9 µl OA during generations 31-50.
I As in **H** for the 10 populations combined.
J Comparison of OA resistance of unevolved *D. simulans* (i.e., the original 10 outcrossed populations maintained on standard fly food; see Materials and Methods), evolved (generation 50 (G50)) *D. simulans*, and *D. sechellia*.
K-N Phenotypic comparison of unevolved and evolved (G50+) *D. simulans* populations 1-3 for **K** noni resistance (G54), **L** fecundity (G54), **M** developmental viability (G57), and **N** lifespan (G57). For **K-M**, significance was assessed using the unpaired t-test correcting for multiple testing. NS p>0.05, ***p<0.001. For **L-N**, tests were run on standard media. No significant differences between unevolved and evolved flies were detected using the Cox regression model. See Materials and Methods for details and Data S1 for raw data.

Fig. 2. Dynamic directional selection of allele frequencies during experimental evolution.
A Schematic of the analysis.
B Identity-by-state analysis. The heatmap depicts the proportion of shared alleles between populations, with values ranging from 0.8 (blue, lowest shared proportion) to 1 (red, identical alleles).
C,D Manhattan plot showing the probability of allele frequency changes between **C** generations 0 and 25, and **D** generations 0 and 50. Top panel shows the results of the CMH test; orange dots indicate significant variants at a threshold of $-\log_{10}(p) \geq 40$, a conservative approximation of the top 0.000005% simulated SNPs under neutrality (Fig. S1B). Bottom panel shows the absolute log-Bayes factors estimated by Bait-ER; orange dots indicate significant variants at the conservative threshold log(0.99/0.01). Raw data in Data S2.
E Circular plots illustrating overlap of predicted genomic regions under directional selection between the two statistical approaches at generation 25 (left) and generation 50 (middle), or between generation 25 and 50 (right). Raw data in Data S3.

Fig. 3. A genome-wide pooled CRISPR screen for OA susceptibility and resistance genes in *D. melanogaster* cultured cells.

A Left: images of S2R+ cells cultured in the absence or presence of OA. Scale bar, 50 μm. Right: dose-dependent effects of OA on S2R+ cell viability, through measurement of ATP levels with a CellTiter-Glo assay (see Materials and Methods).

B Schematic of CRISPR screen design.

C Top 5% "susceptibility" genes (see Materials and Methods and Data S4). Box plots represent the interquartile ranges for all the replicates of the Z-scores obtained through maximum likelihood estimation to test for selection. The inset (bottom-right) displays the gene-ontology analysis of these genes.

D As in **C**, for the top 5% "resistance" genes.

E Venn diagram showing the overlap between all the genes identified in the experimental evolution at G25 and G50, and genes associated with susceptibility and resistance in the CRISPR screen.

F Subportion of the top panel (CMH test) of the Manhattan plot from Fig. 2C for chromosome arms 2L and 3L, with the red arrows highlighting the regions magnified on the right showing the significant variants and linkage disequilibrium blocks (triangles) for *kraken* and *Alkbh7* (CG14130).

Fig. 4. Sequence and expression analyses of *kraken* and *Alkbh7*.

A AlphaFold protein models for *D. melanogaster* Kraken (AF-O18391-F1-model_v4) and Alkbh7 (AF-Q9VTP1-F1-model_v4) [139, 140], highlighting a conserved putative catalytic serine in the active site in Kraken, and the mitochondrial targeting sequence in Alkbh7 (predicted by MitoFates [79]). See also Fig. S2A,B.

B Tissue-specific expression of *D. melanogaster kraken* and *Alkbh7* from bulk RNA-sequencing data from the Fly Atlas 2.0 [141].

C tSNE plots illustrating *D. melanogaster kraken* expression in single-cell transcriptomes of the Malpighian tubules and gut from the Fly Cell Atlas (Stringent 10x datasets) [74].

D Left: whole gut of an adult female expressing GFP (green, endogenous fluorescence signal) under the control of a *kraken* promoter Gal4 driver (genotype: *kraken-Gal4/UAS-FB1.1*, aged 4-5 days); muscles are labelled by phalloidin (red) to reveal tissue morphology. Imaging was performed with a 40× objective; scale bars, 200 μm. Right: higher magnification images of the regions indicated with dashed boxes. Imaging was performed with a 63× objective; scale bars, 100 μm.

E Expression levels of the *kraken* and *Alkbh7* in the indicate strains of adult female *D. sechellia* and *D. simulans* measured by qPCR (Data S6). Expression is represented as calibrated and normalized relative quantities (CNRQ). Significance was assessed using the unpaired t-test correcting for multiple testing and represented using letter codes.

F Expression levels of *kraken* and *Alkbh7* in the indicated strains and species at adult or larval stages as indicated from published RNA-seq data [85-87]. Significance was assessed using the unpaired t-test correcting for multiple testing and represented using letter codes when multiple replicates were present in the original data.

G Expression levels of the candidate genes at generation 0, 25 and 50 of the experimentally-evolved *D. simulans* populations measured by qPCR. Significance was assessed using the paired t-test correcting for multiple testing and represented using letter codes.

Fig. 5. Functional validation of the contribution of *kraken* and *Alkbh7* to OA resistance.

A-C Survival curves of adult flies of the indicated genotypes tested in the plate assay with 2 µl OA for *kraken* and *Alkbh7* RNAi (**A**), mutants (**B**) and overexpression (**C**). Each replicate consisted of 10 2-7 day-old female flies. N replicates ≥6, n flies ≥60. Shading indicates 95% confidence intervals. Genotypes: **A** *act-Gal4/+;UAS-kraken^{RNAi}/+, UAS-kraken^{RNAi}/+, act-Gal4/+, act-Gal4/+;UAS-Alkbh7^{RNAi}/+, UAS-Alkbh7^{RNAi}/+, act-Gal4/+* **B** *kraken^1, Alkbh7^1, Canton-S* (wild-type genetic background control for both mutants), **C** *act-Gal4/+;UAS-kraken$^{o/e}$/+, UAS-kraken$^{o/e}$/+, act-Gal4/+, act-Gal4/+;UAS-Alkbh7$^{o/e}$/+, UAS-Alkbh7$^{o/e}$/+, act-Gal4/+.*

Significance is based on the resulting mixed effect Cox regression model. NS $p>0.05$, *$p<0.05$, **$p<0.01$, ***$p<0.001$. The same data for *D. sechellia* are shown in both plots in **C**.

D Survival curves of wild-type (*Dsec07*), and *kraken* (*Dsec\kraken^{RFP}*) and *Alkbh7* (*Dsec\Alkbh7^{RFP}*) mutant *D. sechellia* in the plate assay with 30 µl OA. Statistics as in **A**.

E Long-term survivability of the same genotypes as in **D** (n = 60 per condition) in control conditions (no OA) (left) and food supplemented with 0.9% OA (right). Statistics as in **A**.

Raw data in Data S7.

Supplementary Information

Supplementary Tables

Table S1. *Drosophila* strains.

Genotype	Reference/Source
D. sechellia 07 (*Dsec07*)	*Drosophila* Species Stock Center (DSSC):14021-0248.07
D. simulans 04 (*Dsim04*)	DSSC:14021-0251.004
D. melanogaster Canton-S (*DmelCS*)	
y[1] w[*]; P{Act5C-GAL4-w}E1/CyO (act-Gal4)	RRID:BDSC_25374
y[1] w[1118] (used as background for act-Gal4)	RRID:BDSC_6598
y[1] w[67c23] (background of Alkbh7$^{o/e}$)	RRID:BDSC_6599
y[1] w[67c23]; P{y[+mDint2] w[+mC]=EPgy2}EY05855 (Alkbh7$^{o/e}$)	RRID:BDSC_16679
y,w[1118];P{attP,y[+],w[3']]} (background of UAS-kraken^{RNAi} and UAS-Alkbh7^{RNAi})	VDRC:v60100
P{KK100356}VIE-260B (UAS-kraken^{RNAi})	VDRC:v105604
P{KK102727}VIE-260B (UAS-Alkbh7^{RNAi})	VDRC:v109669
y[1] M{w[+mC]=nanos-Cas9.P}ZH-2A w[*] (nanos-cas9)	RRID:BDSC_54591
P{hsFLP}1, y1 w1118; P{HD_CFD02326}attP40/CyO-GFP (sgRNA line for Dmel\kraken CRISPR mutant)	VDRC:v342579
CG14130 cris (gRNA 1+2)/CyO (*sgRNA line for Dmel\Alkbh7 CRISPR mutant*)	[97]
y[1] w[67c23];;UAS-kraken (kraken$^{o/e}$)	This study
y[1] w[67c23];;UAS-Dsec\kraken (Dsec\kraken$^{o/e}$)	This study
y[1] w[67c23]; P{y[+t7.7]=CaryP}attP2 (background of kraken$^{o/e}$)	This study
Dmel\kraken-Gal4	This study
UAS-FB1.1 (GFP reporter)	[142]
Dmel\kraken^1	This study
Dmel\Alkbh7^1	This study
Dsec\kraken^{RFP}	This study
Dsec\Alkbh7^{RFP}	This study

Table S2. *D. simulans* lines used to generate the base populations.
- see separate Excel file.

Table S3. Effective population size estimates from the experimentally-evolved populations.

	2L	2R	3L	3R	4	X
pop1	80.65681	101.5796	107.7145	100.5748	369.3642	107.3616
pop2	135.7243	56.29977	76.40593	91.41878	333.638	53.66758
pop3	74.08474	119.4491	95.20849	128.4216	193.5982	99.40552
pop4	83.05221	89.62312	112.3929	116.7979	171.2875	112.1581
pop5	123.0386	106.8179	107.448	81.31624	364.4287	83.55315
pop6	98.8018	77.03804	60.71772	147.1788	124.9405	75.48775
pop7	119.4038	101.4515	96.22689	114.8849	431.7489	114.6792
pop8	136.6513	176.4308	70.8519	130.9822	639.7962	78.70108
pop9	95.58733	101.7351	122.8352	78.80166	109.3757	69.50865
pop10	141.862	101.6789	107.512	83.0109	376.8733	71.34583
mean	108.8863	103.2104	95.73134	107.3388	311.5051	86.58684
median	109.1028	101.6293	101.8374	107.7299	349.0333	81.12711

Table S4. qPCR primer efficiency.

Gene	Efficiency	SE
kraken	1.998	0.041
Alkbh7	1.977	0.016
Act5C	1.976	0.022
αTub84B	2.027	0.023
RpL32	2.053	0.031

Table S5. geNorm analysis of reference target stability.
- see separate Excel file.

Table S6. Oligonucleotides.
- see separate Excel file.

Supplementary Figure legends

Fig. S1. Linkage disequilibrium decay, and comparison of simulated and observed p-value distributions in the experimentally evolved populations.
A Linkage disequilibrium decay at generations 25 (left) and 50 (right). Black curves represent the trajectory of r2 values (y-axis left) as genomic distances increase (x-axis). Red curves represent the same but for D' values (y-axis right).
B Violin plots showing the distribution of p-values from the CMH test, comparing allele frequency changes between generation 25 and 0 (left), and generation 50 and 0 (right). Blue violins represent p-values derived from forward evolution simulations, while orange violins correspond to p-values from experimentally evolved populations. Red dashed lines mark the top 0.000005% of simulated p-values, serving as the significance threshold for the experimental data.

Fig. S2. Sequence and molecular evolutionary analyses of *kraken* and *Alkbh7*.
A,B Protein sequence alignment and secondary structure of Kraken (**A**) and Alkbh7 (**B**) from the reference sequence of *D. melanogaster* (NP_477265.1 and NP_648511.2) and the consensus sequences of *D. simulans* and *D. sechellia*. Sequence data from PRJNA215932 and PRJNA395473 (see Materials and Methods). Black triangles indicate the residues where signs of selection were detected by the FUBAR analysis (see **D**).
C Results from the McDonald-Kreitman test. NI = neutrality index; alpha = proportion of substitutions driven by positive selection; p-value = p-value resulting from a two-tailed Fisher's exact test.
D FUBAR analysis to detect pervasive selection as implemented in HyPhy. position = codon position; alpha = mean posterior synonymous substitution rate; beta = mean posterior nonsynonymous substitution rate; posterior prob = posterior probability of positive selection; codons = alternative codons, Freq Dsec = frequency of the alternative codons in *D. sechellia* in the same order as codons; Freq Dsim = frequency of the alternative codons in *D. simulans* in the same order as codons.
E Survival curves of *D. melanogaster* overexpressing *Dmel\kraken* and *Dsec\kraken* exposed to 2 µl OA. Each replicate consisted of 10 2-7 day-old female flies. Genotypes: *act-Gal4/+;UAS-Dmel\kraken/+*, *act-Gal4/+;UAS-Dsec\kraken/+*. N replicates ≥6, n flies ≥60. Shading indicates 95% confidence intervals. Significance is based on the resulting mixed effect Cox regression model. NS p>0.05. Raw data in Data S5.

Fig. S3. Additional expression analyses of *kraken* and *Alkbh7*.
A Bar plots showing the expression levels of *kraken* and *Alkbh7* across various adult *Drosophila* tissues and species. Data from [143].
B Bar plots showing the expression levels of *kraken* and *Alkbh7* across various larval *Drosophila* tissues and species. Data from [85] ("medium diet" condition).
C Bar plots showing the effect of exposure to 9 µl OA (in the tube assay) on *kraken* and *Alkbh7* expression in *D. sechellia* and three evolved (G50) *D. simulans* populations.

Fig. S4. Validation of RNAi and overexpression lines and generation of mutants for *kraken* and *Alkbh7*.

A Quantitative RT-PCR analysis of RNAi-mediated knockdown of *kraken* and *Alkbh7* in *D. melanogaster*. Genotypes as in Fig. 5A.

B Schematics of the CRISPR/Cas9 design to generate *D. melanogaster* and *D. sechellia* mutants for *kraken* and *Alkbh7*. Validation was performed by PCR and Sanger sequencing. The *Dsec\kraken^{RFP}* mutant comprised a 1431 bp deletion (-44 to +1,387 nt relative to the ATG), replaced with the 3xP3-RFP cassette. The *Dsec\Alkbh7^{RFP}* mutant was a 1023 bp deletion (-84 nt to +939 nt relative to the ATG), replaced with the 3xP3-RFP cassette.

C Quantitative RT-PCR analysis of Gal4/UAS-mediated overexpression of *D. melanogaster kraken* and *Alkbh7*. Genotypes as in Fig. 5C.

Fig. S5. Control survival curves in the absence of OA.

Survival curves of adult flies of the indicated genotypes tested in the plate assay without OA. Each replicate consisted of 10 2-7 day-old female flies. N replicates ≥5, n flies ≥50. All genotypes display full viability, rendering datapoints invisible beneath the topmost displayed genotype (*UAS-kraken^{RNAi}*). Genotypes as in Fig. 5A-B.

Data S1. Source data for Fig. 1

Data S2. Experimental evolution data available at:
https://drive.google.com/file/d/1Gy3k9eYvQ8xQMjlsTCqWevRaO_b1__JY/view?usp=sharing

Data S3. Source data for Fig. 2

Data S4. Source data for Fig. 3

Data S5. Source data for Fig. S2

Data S6. Source data for Fig. 4

Data S7. Source data for Fig. 5

References

1. Erb, M., and Reymond, P. (2019). Molecular Interactions Between Plants and Insect Herbivores. Annu Rev Plant Biol *70*, 527-557.
2. Jones, A.C., Felton, G.W., and Tumlinson, J.H. (2022). The dual function of elicitors and effectors from insects: reviewing the 'arms race' against plant defenses. Plant Mol Biol *109*, 427-445.
3. Smith, J.J., Herzig, V., King, G.F., and Alewood, P.F. (2013). The insecticidal potential of venom peptides. Cell Mol Life Sci *70*, 3665-3693.
4. Umetsu, N., and Shirai, Y. (2020). Development of novel pesticides in the 21st century. J Pestic Sci *45*, 54-74.
5. Raisch, T., and Raunser, S. (2023). The modes of action of ion-channel-targeting neurotoxic insecticides: lessons from structural biology. Nat. Struct. Mol. Biol. *30*, 1411-1427.
6. Sparks, T.C., Crossthwaite, A.J., Nauen, R., Banba, S., Cordova, D., Earley, F., Ebbinghaus-Kintscher, U., Fujioka, S., Hirao, A., Karmon, D., et al. (2020). Insecticides, biologics and nematicides: Updates to IRAC's mode of action classification - a tool for resistance management. Pesticide biochemistry and physiology *167*, 104587.
7. Busvine, J.R. (1951). Mechanism of resistance to insecticide in houseflies. Nature *168*, 193-195.
8. Dong, K. (2007). Insect sodium channels and insecticide resistance. Invert Neurosci *7*, 17-30.
9. Matsuda, K., Ihara, M., and Sattelle, D.B. (2020). Neonicotinoid Insecticides: Molecular Targets, Resistance, and Toxicity. Annu Rev Pharmacol Toxicol *60*, 241-255.
10. Soderlund, D.M., and Bloomquist, J.R. (1989). Neurotoxic actions of pyrethroid insecticides. Annu. Rev. Entomol. *34*, 77-96.
11. Di Prisco, G., Cavaliere, V., Annoscia, D., Varricchio, P., Caprio, E., Nazzi, F., Gargiulo, G., and Pennacchio, F. (2013). Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees. Proceedings of the National Academy of Sciences of the United States of America *110*, 18466-18471.
12. Gandara, L., Jacoby, R., Laurent, F., Spatuzzi, M., Vlachopoulos, N., Borst, N.O., Ekmen, G., Potel, C.M., Garrido-Rodriguez, M., Bohmert, A.L., et al. (2024). Pervasive sublethal effects of agrochemicals on insects at environmentally relevant concentrations. Science *386*, 446-453.
13. Clarkson, C.S., Temple, H.J., and Miles, A. (2018). The genomics of insecticide resistance: insights from recent studies in African malaria vectors. Curr Opin Insect Sci *27*, 111-115.
14. Lucas, E.R., Nagi, S.C., Egyir-Yawson, A., Essandoh, J., Dadzie, S., Chabi, J., Djogbenou, L.S., Medjigbodo, A.A., Edi, C.V., Ketoh, G.K., et al. (2023). Genome-wide association studies reveal novel loci associated with pyrethroid and organophosphate resistance in *Anopheles gambiae* and *Anopheles coluzzii*. Nat Commun *14*, 4946.
15. Catteruccia, F. (2020). Malaria-carrying mosquitoes get a leg up on insecticides. Nature *577*, 319-320.
16. Balabanidou, V., Kefi, M., Aivaliotis, M., Koidou, V., Girotti, J.R., Mijailovsky, S.J., Juarez, M.P., Papadogiorgaki, E., Chalepakis, G., Kampouraki, A., et al.

1089   (2019). Mosquitoes cloak their legs to resist insecticides. Proc Biol Sci *286*,
1090   20191091.

17. Balabanidou, V., Grigoraki, L., and Vontas, J. (2018). Insect cuticle: a critical determinant of insecticide resistance. Curr Opin Insect Sci *27*, 68-74.

18. Weetman, D., Djogbenou, L.S., and Lucas, E. (2018). Copy number variation (CNV) and insecticide resistance in mosquitoes: evolving knowledge or an evolving problem? Curr Opin Insect Sci *27*, 82-88.

19. Zimmer, C.T., Garrood, W.T., Singh, K.S., Randall, E., Lueke, B., Gutbrod, O., Matthiesen, S., Kohler, M., Nauen, R., Davies, T.G.E., et al. (2018). Neofunctionalization of Duplicated P450 Genes Drives the Evolution of Insecticide Resistance in the Brown Planthopper. Curr Biol *28*, 268-274 e265.

20. Gloss, A.D., Vassao, D.G., Hailey, A.L., Nelson Dittrich, A.C., Schramm, K., Reichelt, M., Rast, T.J., Weichsel, A., Cravens, M.G., Gershenzon, J., et al. (2014). Evolution in an ancient detoxification pathway is coupled with a transition to herbivory in the drosophilidae. Mol. Biol. Evol. *31*, 2441-2456.

21. Jones, C.D. (2005). The genetics of adaptation in *Drosophila sechellia*. Genetica *123*, 137-145.

22. Auer, T.O., Shahandeh, M.P., and Benton, R. (2021). *Drosophila sechellia*: A Genetic Model for Behavioral Evolution and Neuroecology. Annu Rev Genet *55*, 527-554.

23. R'Kha, S., Capy, P., and David, J.R. (1991). Host-plant specialization in the *Drosophila melanogaster* species complex: a physiological, behavioral, and genetical analysis. Proceedings of the National Academy of Sciences of the United States of America *88*, 1835-1839.

24. Dekker, T., Ibba, I., Siju, K.P., Stensmyr, M.C., and Hansson, B.S. (2006). Olfactory shifts parallel superspecialism for toxic fruit in *Drosophila melanogaster* sibling, *D. sechellia*. Curr Biol *16*, 101-109.

25. Alvarez-Ocana, R., Shahandeh, M.P., Ray, V., Auer, T.O., Gompel, N., and Benton, R. (2023). Odor-regulated oviposition behavior in an ecological specialist. Nat Commun *14*, 3041.

26. Shahandeh, M.P., Abuin, L., Decker, L.L., Cergneux, J., Koch, R., Nagoshi, E., and Benton, R. (2024). Circadian plasticity evolves via regulatory changes in a neuropeptide gene. Nature *635*, 951-959.

27. Auer, T.O., Khallaf, M.A., Silbering, A.F., Zappia, G., Ellis, K., Alvarez-Ocana, R., Arguello, J.R., Hansson, B.S., Jefferis, G.S.X.E., Caron, S.J.C., et al. (2020). Olfactory receptor and circuit evolution promote host specialization. Nature *579*, 402-408.

28. Jones, C.D. (1998). The genetic basis of *Drosophila sechellia*'s resistance to a host plant toxin. Genetics *149*, 1899-1908.

29. Legal, L., Moulin, B., and Jallon, J.M. (1999). The Relation between Structures and Toxicity of Oxygenated Aliphatic Compounds Homologous to the Insecticide Octanoic Acid and the Chemotaxis of Two Species of *Drosophila*. Pesticide biochemistry and physiology *65*, 90-101.

30. Markow, T.A. (2019). Host use and host shifts in *Drosophila*. Curr Opin Insect Sci *31*, 139-145.

31. Legal, L., Chappe, B., and Jallon, J.M. (1994). Molecular basis of *Morinda citrifolia* (L.): Toxicity on *Drosophila*. J Chem Ecol *20*, 1931-1943.

32. Kaczmarek, A., Katarzyna Wronska, A., and Irena Bogus, M. (2024). Octanoic acid kills Lucilia sericata (Diptera: Calliphoridae) by affecting two major

defence systems: cuticular free fatty acids and immunocompetent cells. J
Invertebr Pathol *206*, 108165.

33. Kaczmarek, A., Wronska, A.K., Kazek, M., and Bogus, M.I. (2022). Octanoic
Acid-An Insecticidal Metabolite of *Conidiobolus coronatus* (Entomopthorales)
That Affects Two Majors Antifungal Protection Systems in *Galleria mellonella*
(Lepidoptera): Cuticular Lipids and Hemocytes. International journal of
molecular sciences *23*.

34. Amlou, M., Pla, E., Moreteau, B., and David, J.R. (1997). Genetic analysis by
interspecific crosses of the tolerance of *Drosophila sechellia* to major aliphatic
acids of its host plant. Genet Sel Evol *29*, 511-522.

35. Amlou, M., Moreteau, B., and David, J.R. (1998). Larval tolerance in the
Drosophila melanogaster species complex toward the two toxic acids of the *D.*
sechellia host plant. Hereditas *129*, 7-14.

36. Huang, Y., and Erezyilmaz, D. (2015). The Genetics of Resistance to *Morinda*
Fruit Toxin During the Postembryonic Stages in *Drosophila sechellia*. G3
(Bethesda) *5*, 1973-1981.

37. Sun, Z., Inagaki, S., Miyoshi, K., Saito, K., and Hayashi, S. (2024). Osiris
gene family defines the cuticle nanopatterns of *Drosophila*. Genetics *227*.

38. Ando, T., Sekine, S., Inagaki, S., Misaki, K., Badel, L., Moriya, H., Sami, M.M.,
Itakura, Y., Chihara, T., Kazama, H., et al. (2019). Nanopore Formation in the
Cuticle of an Insect Olfactory Sensillum. Curr Biol *29*, 1512-1520 e1516.

39. Andrade Lopez, J.M., Lanno, S.M., Auerbach, J.M., Moskowitz, E.C., Sligar,
L.A., Wittkopp, P.J., and Coolon, J.D. (2017). Genetic basis of octanoic acid
resistance in *Drosophila sechellia*: functional analysis of a fine-mapped
region. Mol. Ecol. *26*, 1148-1160.

40. Lanno, S.M., Shimshak, S.J., Peyser, R.D., Linde, S.C., and Coolon, J.D.
(2019). Investigating the role of *Osiris* genes in *Drosophila sechellia* larval
resistance to a host plant toxin. Ecology and evolution *9*, 1922-1933.

41. Schlotterer, C., Kofler, R., Versace, E., Tobler, R., and Franssen, S.U. (2016).
Combining experimental evolution with next-generation sequencing: a
powerful tool to study adaptation from standing genetic variation. Heredity
(Edinb) *116*, 248.

42. Viswanatha, R., Li, Z., Hu, Y., and Perrimon, N. (2018). Pooled genome-wide
CRISPR screening for basal and context-specific fitness gene essentiality in
Drosophila cells. Elife *7*.

43. Colson, I. (2004). *Drosophila simulans*' response to laboratory selection for
tolerance to a toxic food source used by its sister species *D. sechellia*.
Evolutionary Ecology *18*, 15-28.

44. Barghi, N., Tobler, R., Nolte, V., Jaksic, A.M., Mallard, F., Otte, K.A., Dolezal,
M., Taus, T., Kofler, R., and Schlotter, C. (2019). Genetic redundancy fuels
polygenic adaptation in *Drosophila*. PLOS Biol. *17*, e3000128.

45. Turner, T.L., and Miller, P.M. (2012). Investigating natural variation in
Drosophila courtship song by the evolve and resequence approach. Genetics
191, 633-642.

46. Kelly, J.K., and Hughes, K.A. (2019). Pervasive Linked Selection and
Intermediate-Frequency Alleles Are Implicated in an Evolve-and-
Resequencing Experiment of *Drosophila simulans*. Genetics *211*, 943-961.

47. Shahrestani, P., King, E., Ramezan, R., Phillips, M., Riddle, M., Thornburg,
M., Greenspan, Z., Estrella, Y., Garcia, K., Chowdhury, P., et al. (2021). The
molecular architecture of *Drosophila melanogaster* defense against *Beauveria*

1188		*bassiana* explored through evolve and resequence and quantitative trait locus
1189		mapping. G3 (Bethesda) *11*.
1190	48.	Zoh, M.G., Bonneville, J.M., Tutagata, J., Laporte, F., Fodjo, B.K.,
1191		Mouhamadou, C.S., Sadia, C.G., McBeath, J., Schmitt, F., Horstmann, S., et
1192		al. (2021). Experimental evolution supports the potential of neonicotinoid-
1193		pyrethroid combination for managing insecticide resistance in malaria vectors.
1194		Sci Rep *11*, 19501.
1195	49.	Sadia, C.G., Bonneville, J.M., Zoh, M.G., Fodjo, B.K., Kouadio, F.A., Oyou,
1196		S.K., Koudou, B.G., Adepo-Gourene, B.A., Reynaud, S., David, J.P., et al.
1197		(2024). The impact of agrochemical pollutant mixtures on the selection of
1198		insecticide resistance in the malaria vector *Anopheles gambiae*: insights from
1199		experimental evolution and transcriptomics. Malar J *23*, 69.
1200	50.	Long, A., Liti, G., Luptak, A., and Tenaillon, O. (2015). Elucidating the
1201		molecular architecture of adaptation via evolve and resequence experiments.
1202		Nat. Rev. Genet. *16*, 567-582.
1203	51.	Teotonio, H., Chelo, I.M., Bradic, M., Rose, M.R., and Long, A.D. (2009).
1204		Experimental evolution reveals natural selection on standing genetic variation.
1205		Nat Genet *41*, 251-257.
1206	52.	Keightley, P.D., Ness, R.W., Halligan, D.L., and Haddrill, P.R. (2014).
1207		Estimation of the spontaneous mutation rate per nucleotide site in a
1208		*Drosophila melanogaster* full-sib family. Genetics *196*, 313-320.
1209	53.	Signor, S.A., New, F.N., and Nuzhdin, S. (2018). A Large Panel of *Drosophila*
1210		*simulans* Reveals an Abundance of Common Variants. Genome Biol. Evol.
1211		*10*, 189-206.
1212	54.	R'Kha, S., Moreteau, B., Coyne, J.A., and David, J.R. (1997). Evolution of a
1213		lesser fitness trait: egg production in the specialist *Drosophila sechellia*. Genet
1214		Res *69*, 17-23.
1215	55.	Markow, T.A., Beall, S., and Matzkin, L.M. (2009). Egg size, embryonic
1216		development time and ovoviviparity in *Drosophila* species. J Evol Biol *22*, 430-
1217		434.
1218	56.	Abe, M., Kamiyama, T., Izumi, Y., Qian, Q., Yoshihashi, Y., Degawa, Y.,
1219		Watanabe, K., Hattori, Y., Uemura, T., and Niwa, R. (2022). Shortened
1220		lifespan induced by a high-glucose diet is associated with intestinal immune
1221		dysfunction in *Drosophila sechellia*. J Exp Biol *225*.
1222	57.	Watada, M., Hayashi, Y., Watanabe, K., Mizutani, S., Mure, A., Hattori, Y.,
1223		and Uemura, T. (2020). Divergence of *Drosophila* species: Longevity and
1224		reproduction under different nutrient balances. Genes Cells *25*, 626-636.
1225	58.	Vlachos, C., Burny, C., Pelizzola, M., Borges, R., Futschik, A., Kofler, R., and
1226		Schlotterer, C. (2019). Benchmarking software tools for detecting and
1227		quantifying selection in evolve and resequencing studies. Genome Biol *20*,
1228		169.
1229	59.	Kojima, Y., Matsumoto, H., and Kiryu, H. (2020). Estimation of population
1230		genetic parameters using an EM algorithm and sequence data from
1231		experimental evolution populations. Bioinformatics *36*, 221-231.
1232	60.	Barata, C., Borges, R., and Kosiol, C. (2023). Bait-ER: A Bayesian method to
1233		detect targets of selection in Evolve-and-Resequence experiments. J Evol
1234		Biol *36*, 29-44.
1235	61.	Cochran, W.G. (1954). Some Methods for Strengthening the Common Chi-
1236		Squared Tests. Biometrics *10*, 417-451.

1237   62.   Mantel, N., and Haenszel, W. (1959). Statistical aspects of the analysis of
1238   data from retrospective studies of disease. J Natl Cancer Inst *22*, 719-748.
1239   63.   Orozco-terWengel, P., Kapun, M., Nolte, V., Kofler, R., Flatt, T., and
1240   Schlotterer, C. (2012). Adaptation of *Drosophila* to a novel laboratory
1241   environment reveals temporally heterogeneous trajectories of selected alleles.
1242   Mol. Ecol. *21*, 4931-4941.
1243   64.   Hungate, E.A., Earley, E.J., Boussy, I.A., Turissini, D.A., Ting, C.T., Moran,
1244   J.R., Wu, M.L., Wu, C.I., and Jones, C.D. (2013). A locus in *Drosophila*
1245   *sechellia* affecting tolerance of a host plant toxin. Genetics *195*, 1063-1075.
1246   65.   Hu, Y., Comjean, A., Perrimon, N., and Mohr, S.E. (2017). The *Drosophila*
1247   Gene Expression Tool (DGET) for expression analyses. BMC Bioinf. *18*, 98.
1248   66.   Xu, Y., Viswanatha, R., Sitsel, O., Roderer, D., Zhao, H., Ashwood, C.,
1249   Voelcker, C., Tian, S., Raunser, S., Perrimon, N., et al. (2022). CRISPR
1250   screens in *Drosophila* cells identify Vsg as a Tc toxin receptor. Nature *610*,
1251   349-355.
1252   67.   Borrull, A., Lopez-Martinez, G., Poblet, M., Cordero-Otero, R., and Rozes, N.
1253   (2015). New insights into the toxicity mechanism of octanoic and decanoic
1254   acids on *Saccharomyces cerevisiae*. Yeast *32*, 451-460.
1255   68.   Emelyanov, A.V., Rabbani, J., Mehta, M., Vershilova, E., Keogh, M.C., and
1256   Fyodorov, D.V. (2014). *Drosophila* TAP/p32 is a core histone chaperone that
1257   cooperates with NAP-1, NLP, and nucleophosmin in sperm chromatin
1258   remodeling during fertilization. Genes Dev *28*, 2027-2040.
1259   69.   Galindo, K.A., Lu, W.J., Park, J.H., and Abrams, J.M. (2009). The Bax/Bak
1260   ortholog in *Drosophila*, Debcl, exerts limited control over programmed cell
1261   death. Development *136*, 275-283.
1262   70.   Edwin Chan, H.Y., Harris, S.J., and O'Kane, C.J. (1998). Identification and
1263   characterization of kraken, a gene encoding a putative hydrolytic enzyme in
1264   *Drosophila melanogaster*. Gene *222*, 195-201.
1265   71.   Zhang, L.S., Xiong, Q.P., Pena Perez, S., Liu, C., Wei, J., Le, C., Zhang, L.,
1266   Harada, B.T., Dai, Q., Feng, X., et al. (2021). ALKBH7-mediated
1267   demethylation regulates mitochondrial polycistronic RNA processing. Nat Cell
1268   Biol *23*, 684-691.
1269   72.   Solberg, A., Robertson, A.B., Aronsen, J.M., Rognmo, O., Sjaastad, I.,
1270   Wisloff, U., and Klungland, A. (2013). Deletion of mouse Alkbh7 leads to
1271   obesity. J Mol Cell Biol *5*, 194-203.
1272   73.   Kumar, K., Mhetre, A., Ratnaparkhi, G.S., and Kamat, S.S. (2021). A
1273   Superfamily-wide Activity Atlas of Serine Hydrolases in *Drosophila*
1274   *melanogaster*. Biochemistry *60*, 1312-1324.
1275   74.   Li, H., Janssens, J., De Waegeneer, M., Kolluru, S.S., Davie, K., Gardeux, V.,
1276   Saelens, W., David, F.P.A., Brbic, M., Spanier, K., et al. (2022). Fly Cell Atlas:
1277   A single-nucleus transcriptomic atlas of the adult fruit fly. Science *375*,
1278   eabk2432.
1279   75.   Lemaitre, B., and Miguel-Aliaga, I. (2013). The digestive tract of *Drosophila*
1280   *melanogaster*. Annu Rev Genet *47*, 377-404.
1281   76.   Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gislason, M.H., Pihl,
1282   S.I., Tsirigos, K.D., Winther, O., Brunak, S., von Heijne, G., and Nielsen, H.
1283   (2022). SignalP 6.0 predicts all five types of signal peptides using protein
1284   language models. Nat. Biotechnol. *40*, 1023-1025.

1285   77.   Yang, J., McCart, C., Woods, D.J., Terhzaz, S., Greenwood, K.G., ffrench-
1286   Constant, R.H., and Dow, J.A. (2007). A *Drosophila* systems approach to
1287   xenobiotic metabolism. Physiol Genomics *30*, 223-231.
1288   78.   Xu, B., Liu, D., Wang, Z., Tian, R., and Zuo, Y. (2021). Multi-substrate
1289   selectivity based on key loops and non-homologous domains: new insight into
1290   ALKBH family. Cell Mol Life Sci *78*, 129-141.
1291   79.   Fukasawa, Y., Tsuji, J., Fu, S.C., Tomii, K., Horton, P., and Imai, K. (2015).
1292   MitoFates: improved prediction of mitochondrial targeting sequences and their
1293   cleavage sites. Molecular & cellular proteomics : MCP *14*, 1113-1126.
1294   80.   Chen, C.L., Hu, Y., Udeshi, N.D., Lau, T.Y., Wirtz-Peitz, F., He, L., Ting, A.Y.,
1295   Carr, S.A., and Perrimon, N. (2015). Proteomic mapping in live *Drosophila*
1296   tissues using an engineered ascorbate peroxidase. Proceedings of the
1297   National Academy of Sciences of the United States of America *112*, 12093-
1298   12098.
1299   81.   Sen, A., and Cox, R.T. (2022). Loss of *Drosophila* Clueless differentially
1300   affects the mitochondrial proteome compared to loss of Sod2 and Pink1.
1301   Frontiers in physiology *13*, 1004099.
1302   82.   Schrider, D.R., Ayroles, J., Matute, D.R., and Kern, A.D. (2018). Supervised
1303   machine learning reveals introgressed loci in the genomes of *Drosophila*
1304   *simulans* and *D. sechellia*. PLOS Genet. *14*, e1007341.
1305   83.   McDonald, J.H., and Kreitman, M. (1991). Adaptive protein evolution at the
1306   Adh locus in *Drosophila*. Nature *351*, 652-654.
1307   84.   Murrell, B., Moola, S., Mabona, A., Weighill, T., Sheward, D., Kosakovsky
1308   Pond, S.L., and Scheffler, K. (2013). FUBAR: a fast, unconstrained bayesian
1309   approximation for inferring selection. Mol. Biol. Evol. *30*, 1196-1205.
1310   85.   Watanabe, K., Kanaoka, Y., Mizutani, S., Uchiyama, H., Yajima, S., Watada,
1311   M., Uemura, T., and Hattori, Y. (2019). Interspecies Comparative Analyses
1312   Reveal Distinct Carbohydrate-Responsive Systems among *Drosophila*
1313   Species. Cell Rep *28*, 2594-2607 e2597.
1314   86.   Ma, S., Avanesov, A.S., Porter, E., Lee, B.C., Mariotti, M., Zemskaya, N.,
1315   Guigo, R., Moskalev, A.A., and Gladyshev, V.N. (2018). Comparative
1316   transcriptomics across 14 *Drosophila* species reveals signatures of longevity.
1317   Aging Cell *17*, e12740.
1318   87.   Kalra, S., Lanno, S., Sanchez, G., and Coolon, J.D. (2024). cis- and trans-
1319   regulatory contributions to a hierarchy of factors influencing gene expression
1320   variation. Communications biology *7*, 1563.
1321   88.   Mota, M.N., Matos, M., Bahri, N., and Sa-Correia, I. (2024). Shared and more
1322   specific genetic determinants and pathways underlying yeast tolerance to
1323   acetic, butyric, and octanoic acids. Microb Cell Fact *23*, 71.
1324   89.   Liu, P., Chernyshov, A., Najdi, T., Fu, Y., Dickerson, J., Sandmeyer, S., and
1325   Jarboe, L. (2013). Membrane stress caused by octanoic acid in
1326   *Saccharomyces cerevisiae*. Appl Microbiol Biotechnol *97*, 3239-3251.
1327   90.   National Center for Biotechnology Information PubChem Compound
1328   Summary for CID 379, Octanoic Acid. Retrieved January 19, 2024.
1329   91.   Marconcini, M., Fragnière, C., Masuzzo, A., and Benton, R. (2025). Genome-
1330   wide association studies identify new candidate genes and tissues underlying
1331   resistance to a natural toxin in drosophilids. bioRxiv.
1332   92.   Dow, J.A.T., Simons, M., and Romero, M.F. (2022). *Drosophila melanogaster*:
1333   a simple genetic model of kidney structure, function and disease. Nat Rev
1334   Nephrol *18*, 417-434.

93. Gutierrez-Guerrero, Y.T., Ibarra-Laclette, E., Martinez Del Rio, C., Barrera-Redondo, J., Rebollar, E.A., Ortega, J., Leon-Paniagua, L., Urrutia, A., Aguirre-Planter, E., and Eguiarte, L.E. (2020). Genomic consequences of dietary diversification and parallel evolution due to nectarivory in leaf-nosed bats. Gigascience *9*.

94. Xia, B., Viswanatha, R., Hu, Y., Mohr, S.E., and Perrimon, N. (2023). Pooled genome-wide CRISPR activation screening for rapamycin resistance genes in *Drosophila* cells. Elife *12*.

95. Viswanatha, R., Mameli, E., Rodiger, J., Merckaert, P., Feitosa-Suntheimer, F., Colpitts, T.M., Mohr, S.E., Hu, Y., and Perrimon, N. (2021). Bioinformatic and cell-based tools for pooled CRISPR knockout screening in mosquitos. Nat Commun *12*, 6825.

96. Port, F., Strein, C., Stricker, M., Rauscher, B., Heigwer, F., Zhou, J., Beyersdorffer, C., Frei, J., Hess, A., Kern, K., et al. (2020). A large-scale resource for tissue-specific CRISPR mutagenesis in *Drosophila*. Elife *9*.

97. Lenče, T. (2021). PhD Thesis: The role of m6A modification on mRNA processing in *Drosophila melanogaster*. Volume PhD. (Johannes Gutenberg University Mainz).

98. Kondo, S., and Ueda, R. (2013). Highly Improved Gene Targeting by Germline-Specific Cas9 Expression in *Drosophila*. Genetics.

99. Croset, V., Rytz, R., Cummins, S.F., Budd, A., Brawand, D., Kaessmann, H., Gibson, T.J., and Benton, R. (2010). Ancient protostome origin of chemosensory ionotropic glutamate receptors and the evolution of insect taste and olfaction. PLOS Genet. *6*, e1001064.

100. Bischof, J., Maeda, R.K., Hediger, M., Karch, F., and Basler, K. (2007). An optimized transgenesis system for *Drosophila* using germ-line-specific phiC31 integrases. Proceedings of the National Academy of Sciences of the United States of America *104*, 3312-3317.

101. Therneau, T.M. (2024). coxme: Mixed Effects Cox Models. R package version 2.2-22.

102. Pino, J.A., Marquez, E., and Castro, D. (2009). Volatile and non-volatile acids of noni (*Morinda citrifolia* L.) fruit. J Sci Food Agric *89*, 1247-1249.

103. Wood, S.N. (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. Journal of the Royal Statistical Society: Series B (Statistical Methodology) *73*, 3-36.

104. Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics *30*, 2114-2120.

105. Li, H. (2013). Aligning Sequence Reads, Clone Sequences and Assembly Contigs with BWA-MEM. arXiv.

106. Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane, T., McCarthy, S.A., Davies, R.M., et al. (2021). Twelve years of SAMtools and BCFtools. Gigascience *10*.

107. Kofler, R., Pandey, R.V., and Schlotterer, C. (2011). PoPoolation2: identifying differentiation between populations using sequencing of pooled DNA samples (Pool-Seq). Bioinformatics *27*, 3435-3436.

108. Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., et al. (2011). The variant call format and VCFtools. Bioinformatics *27*, 2156-2158.

109. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller, J., Sklar, P., de Bakker, P.I., Daly, M.J., et al. (2007). PLINK: a tool set

for whole-genome association and population-based linkage analyses. Am J
Hum Genet *81*, 559-575.

110. Purcell, S. (2020). PLINK v1.90b6.21.

111. Vlachos, C., and Kofler, R. (2018). MimicrEE2: Genome-wide forward
simulations of Evolve and Resequencing studies. PLoS Comput Biol *14*,
e1006413.

112. Taus, T., Futschik, A., and Schlotterer, C. (2017). Quantifying Selection with
Pool-Seq Time Series Data. Mol. Biol. Evol. *34*, 3023-3034.

113. Zhang, C., Dong, S.S., Xu, J.Y., He, W.M., and Yang, T.L. (2019).
PopLDdecay: a fast and effective tool for linkage disequilibrium decay
analysis based on variant call format files. Bioinformatics *35*, 1786-1788.

114. Dong, S.S., He, W.M., Ji, J.J., Zhang, C., Guo, Y., and Yang, T.L. (2021).
LDBlockShow: a fast and convenient tool for visualizing linkage disequilibrium
and haplotype blocks based on variant call format files. Briefings in
bioinformatics *22*.

115. Barrett, J.C., Fry, B., Maller, J., and Daly, M.J. (2005). Haploview: analysis
and visualization of LD and haplotype maps. Bioinformatics *21*, 263-265.

116. Gu, Z., Gu, L., Eils, R., Schlesner, M., and Brors, B. (2014). circlize
Implements and enhances circular visualization in R. Bioinformatics *30*, 2811-
2812.

117. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for
comparing genomic features. Bioinformatics *26*, 841-842.

118. Altenhoff, A.M., Train, C.M., Gilbert, K.J., Mediratta, I., Mendes de Farias, T.,
Moi, D., Nevers, Y., Radoykova, H.S., Rossier, V., Warwick Vesztrocy, A., et
al. (2021). OMA orthology in 2021: website overhaul, conserved isoforms,
ancestral gene order and more. Nucleic Acids Res. *49*, D373-D379.

119. Sherman, B.T., Hao, M., Qiu, J., Jiao, X., Baseler, M.W., Lane, H.C.,
Imamichi, T., and Chang, W. (2022). DAVID: a web server for functional
enrichment analysis and functional annotation of gene lists (2021 update).
Nucleic Acids Res. *50*, W216-W221.

120. Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and
integrative analysis of large gene lists using DAVID bioinformatics resources.
Nat Protoc *4*, 44-57.

121. Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer,
K., and Madden, T.L. (2009). BLAST+: architecture and applications. BMC
Bioinf. *10*, 421.

122. Sayers, E.W., Beck, J., Bolton, E.E., Brister, J.R., Chan, J., Connor, R.,
Feldgarden, M., Fine, A.M., Funk, K., Hoffman, J., et al. (2025). Database
resources of the National Center for Biotechnology Information in 2025.
Nucleic Acids Res. *53*, D20-D29.

123. Ozturk-Colak, A., Marygold, S.J., Antonazzo, G., Attrill, H., Goutte-Gattat, D.,
Jenkins, V.K., Matthews, B.B., Millburn, G., Dos Santos, G., Tabone, C.J., et
al. (2024). FlyBase: updates to the *Drosophila* genes and genomes database.
Genetics *227*.

124. Viswanatha, R., Entwisle, S., Hu, Y., Kim, A.R., Reap, K., Butnaru, M., Qadiri,
M., Mohr, S.E., and Perrimon, N. (2025). Higher resolution pooled genome-
wide CRISPR knockout screening in *Drosophila* cells using integration and
anti-CRISPR (IntAC). Nat Commun *16*, 6498.

125. Viswanatha, R., Brathwaite, R., Hu, Y., Li, Z., Rodiger, J., Merckaert, P.,
Chung, V., Mohr, S.E., and Perrimon, N. (2019). Pooled CRISPR Screens in
Drosophila Cells. Current protocols in molecular biology *129*, e111.

126. Li, W., Xu, H., Xiao, T., Cong, L., Love, M.I., Zhang, F., Irizarry, R.A., Liu, J.S.,
Brown, M., and Liu, X.S. (2014). MAGeCK enables robust identification of
essential genes from genome-scale CRISPR/Cas9 knockout screens.
Genome Biol *15*, 554.

127. Hu, Y., Comjean, A., Attrill, H., Antonazzo, G., Thurmond, J., Chen, W., Li, F.,
Chao, T., Mohr, S.E., Brown, N.H., et al. (2023). PANGEA: a new gene set
enrichment tool for *Drosophila* and common research organisms. Nucleic
Acids Res. *51*, W419-W426.

128. Hellemans, J., Mortier, G., De Paepe, A., Speleman, F., and Vandesompele,
J. (2007). qBase relative quantification framework and software for
management and automated analysis of real-time quantitative PCR data.
Genome Biol *8*, R19.

129. Lu, J., Yang, C., Zhang, Y., and Pan, H. (2018). Selection of Reference
Genes for the Normalization of RT-qPCR Data in Gene Expression Studies in
Insects: A Systematic Review. Frontiers in physiology *9*, 1560.

130. Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De
Paepe, A., and Speleman, F. (2002). Accurate normalization of real-time
quantitative RT-PCR data by geometric averaging of multiple internal control
genes. Genome Biol *3*, RESEARCH0034.

131. Rogers, R.L., Cridland, J.M., Shao, L., Hu, T.T., Andolfatto, P., and Thornton,
K.R. (2014). Landscape of standing variation for tandem duplications in
Drosophila yakuba and *Drosophila simulans*. Mol. Biol. Evol. *31*, 1750-1766.

132. Leinonen, R., Akhtar, R., Birney, E., Bonfield, J., Bower, L., Corbett, M.,
Cheng, Y., Demiralp, F., Faruque, N., Goodgame, N., et al. (2010).
Improvements to services at the European Nucleotide Archive. Nucleic Acids
Res. *38*, D39-45.

133. Walker, B.J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S.,
Cuomo, C.A., Zeng, Q., Wortman, J., Young, S.K., et al. (2014). Pilon: an
integrated tool for comprehensive microbial variant detection and genome
assembly improvement. PLOS ONE *9*, e112963.

134. Abascal, F., Zardoya, R., and Telford, M.J. (2010). TranslatorX: multiple
alignment of nucleotide sequences guided by amino acid translations. Nucleic
Acids Res. *38*, W7-13.

135. Madeira, F., Madhusoodanan, N., Lee, J., Eusebi, A., Niewielska, A., Tivey,
A.R.N., Lopez, R., and Butcher, S. (2024). The EMBL-EBI Job Dispatcher
sequence analysis tools framework in 2024. Nucleic Acids Res. *52*, W521-
W525.

136. Rozas, J., Ferrer-Mata, A., Sanchez-DelBarrio, J.C., Guirao-Rico, S., Librado,
P., Ramos-Onsins, S.E., and Sanchez-Gracia, A. (2017). DnaSP 6: DNA
Sequence Polymorphism Analysis of Large Data Sets. Mol. Biol. Evol. *34*,
3299-3302.

137. Kosakovsky Pond, S.L., Poon, A.F.Y., Velazquez, R., Weaver, S., Hepler,
N.L., Murrell, B., Shank, S.D., Magalis, B.R., Bouvier, D., Nekrutenko, A., et
al. (2020). HyPhy 2.5-A Customizable Platform for Evolutionary Hypothesis
Testing Using Phylogenies. Mol. Biol. Evol. *37*, 295-299.

138. Robert, X., and Gouet, P. (2014). Deciphering key features in protein
structures with the new ENDscript server. Nucleic Acids Res. *42*, W320-324.

139. Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova,
G., Yuan, D., Stroe, O., Wood, G., Laydon, A., et al. (2022). AlphaFold Protein
Structure Database: massively expanding the structural coverage of protein-
sequence space with high-accuracy models. Nucleic Acids Res. *50*, D439-
D444.
140. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly
accurate protein structure prediction with AlphaFold. Nature *596*, 583-589.
141. Krause, S.A., Overend, G., Dow, J.A.T., and Leader, D.P. (2022). FlyAtlas 2
in 2022: enhancements to the *Drosophila melanogaster* expression atlas.
Nucleic Acids Res. *50*, D1010-D1015.
142. Hadjieconomou, D., Rotkopf, S., Alexandre, C., Bell, D.M., Dickson, B.J., and
Salecker, I. (2011). Flybow: genetic multicolor cell labeling for neural circuit
analysis in *Drosophila melanogaster*. Nat Methods *8*, 260-266.
143. Bontonou, G., Saint-Leandre, B., Kafle, T., Baticle, T., Hassan, A., Sanchez-
Alcaniz, J.A., and Arguello, J.R. (2024). Evolution of chemosensory tissues
and cells across ecologically diverse Drosophilids. Nat Commun *15*, 1047.
1500

A

Dmel\kraken
Dsim\kraken
Dsec\kraken

AB hydrolase-1

catalytic serine

B

Dmel\Alkbh7
Dsim\Alkbh7
Dsec\Alkbh7

mitochondrial targeting

C

McDonald & Kreitman Test

	NI	alpha	p-value
kraken	0.083	0.917	0.018*
Alkbh7	0.544	0.456	0.477

D

HyPhy FUBAR

	position	alpha	beta	post. prob.	codons	Freq *Dsec*	Freq *Dsim*
kraken	27	2.503	40.230	0.979	S / A / T	1 / 0 / 0	0.658 / 0.237 / 0.105
Alkbh7	66	0.768	9.440	0.977	H / Y	1 / 0	0.806 / 0.194
	105	0.665	7.158	0.957	V / I / L	1 / 0 / 0	0.839 / 0.097 / 0.064
	126	0.783	13.505	0.989	D / E	1 / 0	0.774 / 0.226
	141	0.723	9.841	0.982	R / K	1 / 0	0.871 / 0.129

E

Dmel\kraken vs Dsec\kraken

Survival probability

Time (min)

2 µl OA
■ *act-Gal4>UAS-Dmel\kraken* | NS
■ *act-Gal4>UAS-Dsec\kraken* |