Interannual variation in *Protocalliphora* blowfly (Diptera: Calliphoridae) parasitism of

Tree Swallows (*Tachycineta bicolor*) breeding in southern Québec's farmlands: 16 years of

data on species composition, morphometry and hyperparasitism by *Nasonia* wasps

(Hymenoptera: Pteromalidae)

Authors :

Coroller-Chouraki, Simon[1,2,4] ; Savage, Jade[3] ; Pelletier, Fanie[1] ; Garant, Dany[1] ; Bélisle, Marc[1,2]

Authors Affiliations :

[1]Département de biologie, Université de Sherbrooke, 2500 Boulevard de l'Université,

Sherbrooke (Québec), Canada

[2]Centre d'étude de la forêt, Pavillon des Sciences biologiques, Montréal (Québec), Canada

[3]Department of Biology and Biochemistry, Bishop's University, Sherbrooke (Québec), Canada

[4]Simon.Coroller@USherbrooke.ca

ABSTRACT

Protocalliphora are among the most prevalent ectoparasites of altricial birds across the Holarctic. Yet, their spatial and temporal dynamics of infestations, species composition, and interactions with parasitoids remain poorly understood. We present a 16-year (2004-2019) multi-site study of *Protocalliphora* infestations based on 2,673 Tree Swallow (*Tachycineta bicolor*) nests collected across a 10,200-km² gradient of agricultural intensity in Québec, Canada. *Protocalliphora* nest infestation prevalence and load varied markedly across space and time, and showed cyclical patterns at ~75% of sites, suggesting the influence of regional and local processes. Yearly rates of hyperparasitism of *Protocalliphora* puparia by *Nasonia* parasitoid wasps were also high but variable (48-90%), likely contributing to the temporal oscillations in *Protocalliphora* prevalence and load. Substantial interannual shifts in the relative abundance of three *Protocalliphora* species (*bennetti*, *metallica*, and *sialia*) emphasized the importance of species-level resolution in blowfly ecological studies. Considerable overlap in puparia size among species challenged the utility of traditional diagnostic traits for species identification. Finally, dormancy or mortality of *Nasonia* occurred in 3-16% of *Protocalliphora* puparia depending on year. These findings highlight the importance of long-term, multi-trophic, and spatially explicit monitoring to unravel the complex ecological drivers of host-parasite-parasitoid dynamics in increasingly human-modified environments such as agroecosystems.

INTRODUCTION

Avian ectoparasites encompass an exceptionally diverse array of invertebrate taxa, ranging from leeches (Hirudinea) and ticks (Ixodida) to insects from various orders, including lice (Phthiraptera), true bugs (Hemiptera), fleas (Siphonaptera), and flies (Diptera; Rothschild & Clay 1952; Clayton et al. 2010). Many of these ectoparasites are hematophagous, a feeding mode that has evolved independently multiple times within Diptera and which occurs in many families of both the Nematocera and Brachycera sub-orders (e.g., Simuliidae, Ceratopogonidae, Hippoboscidae, Nycteribiidae). However, significant differences exist among dipteran ectoparasites in terms of host selection and specificity, whether they are obligatory blood-feeders or not, and regarding diet differentiation across life stages. For instance, whereas the larvae of *Protocalliphora* blowflies (Calliphoridae) are obligatory blood-feeding parasites of nestling belonging to a wide diversity of altricial bird species, the adults appear to feed on the pollen of herbaceous plants (Sabrosky et al. 1989; Bennett & Whithworth 1991). While the biology and ecology of hematophagous flies parasitizing humans and domestic animals are well known (e.g., Marquardt et al. 2004), flies parasitizing wild animals have received much less attention or have mainly been studied through the costs they can inflict on their hosts. *Protocalliphora* blowflies exemplify this case as most previous studies focused on measuring the impacts that their larvae may have on the nestlings and reproductive success of altricial birds (Simon et al. 2004; Maziarz et al. 2022). In recent years, however, the environmental factors that drive the prevalence and intensity of bird nest infestations by *Protocalliphora* have been increasingly addressed (Mennerat et al. 2021; Merino et al. 2024). Yet only a few studies have so far formally investigated the determinants of bird nest infestations by *Protocalliphora* in multiple years and sites (Mennerat et al. 2021; Maziarz et al. 2022; Merino et al. 2024). Moreover, these studies, which mainly focused

on the effects of weather or climatic factors, were conducted in a small number of forested sites,

and thus, on a limited range of habitat heterogeneity. Thus, we have limited information on the

spatial and temporal variability of infestations, and thereby on the influence of abiotic and biotic

factors on the host-parasite dynamics involving *Protocalliphora*.

The lack of long-term monitoring data on bird nest infestations by *Protocalliphora* also concerns

the species of blowflies involved in these interactions. At least 40 species of *Protocalliphora*

blowflies have been described in the Holarctic region (Sabrosky et al. 1989), many of which have

overlapping distribution ranges, resulting in occasional mixed infestations within a bird nest

(Daoust et al. 2012). Moreover, the morphological identification of *Protocalliphora* species is

challenging, at least to the non-specialists, and has been considered to be reliable only in adults

(Heeb et al. 2000). Yet, locating and sampling adult blowflies in the field is no easy task (Stiner

1969; Bennett & Whitworth 1991), thus larvae and puparia often remain the primary evidence of

blowfly parasitism and parasitic load (Jánošková et al. 2010). Despite significant progress

regarding the taxonomy and identification of *Protocalliphora*, notably through the work of

Sabrosky et al. (1989) and Whitworth (2002, 2003, 2019), identifying *Protocalliphora* species

based on puparia morphology is complicated by the fact that key criteria, such as distance between

specific spines, prothoracic fringe diameter, and distance between posterior spiracles can vary

widely within species and overlap between species (Jánošková et al. 2010; Jamriška & Modrý

2013). Furthermore, identification based on DNA barcoding using the Folmer region of the

common COI gene fails to discriminate between most described *Protocalliphora* species

(Whitworth et al. 2007). These identification challenges, especially considering the high parasite

load found in some nests, have understandably led ecologists to neglect *Protocalliphora* species

composition and typically lump all individuals under *"Protocalliphora spp."* (e.g., Heeb et al.

2000; Bańbura et al. 2004; Hannam 2006; Musgrave et al. 2019). Using *Protocalliphora* as a

homogeneous entity can, however, bias species abundance estimates, and hinder the study of host-

parasite dynamics by neglecting the potential competition that may occur among *Protocalliphora*

species as well as their interactions with their hyperparasites within host nests.

Other overlooked aspects that may impact host-parasite dynamics involving *Protocalliphora*

include within and between-species variation in body size that may partly reflect the quality of the

growth conditions they experienced as larvae and ultimately their metamorphosis and emergence

success, notably through their susceptibility to hyperparasitism by gregarious *Nasonia* "jewel

wasps" (Hymenoptera: Pteromalidae; Werren & Loehlin 2009). These generalist wasps, which

parasitize *Protocalliphora* at the pupal stage, are known for their effective control of insect host

populations, including agricultural pests, and play a substantial role in the top-down regulation of

calyptrate flies. Most research on *Nasonia*, including the well-known *N. vitripennis*, and *N.

giraulti*, which attack *Protocalliphora* (Daoust et al. 2012; Garrido-Bautista et al. 2019), has

however largely focused on their genetics and behavioral ecology under laboratory conditions

(e.g., Werren & Loehlin 2009; Deveux et al. 2023). Past studies hence provide limited insights into

the ecological interactions between *Nasonia* and *Protocalliphora* in natural habitats, and how their

top-down regulation potential against blowflies may fluctuate under different conditions in either

space or time. These gaps in our understanding of *Protocalliphora* extend beyond taxonomy,

morphometry, and distribution. Long-term studies focusing on the prevalence and abundance of

blowfly larvae within nests have noted important annual fluctuations, after accounting for some

environmental conditions (Wesolowski, 2001; Daoust et al. 2012; Musgrave et al. 2019; Merino

et al. 2024). However, only Merino et al. (2024) have described cyclical trends in abundance and

to our knowledge, no details on temporal variation in species composition and hyperparasitic

pressures have been documented.

In this study, we quantify the spatial and temporal variability in the prevalence, intensity, and

species composition of Tree Swallow (*Tachycineta bicolor*) nest infestations by *Protocalliphora*

over a 16-year period within a 10,200-km² gradient of agricultural intensity in southern Québec,

Canada. We also report species-specific size distributions for the puparia of three species of

Protocalliphora, temporal variation in hyperparasitism rates by *Nasonia* wasps as well as the

proportion of this hyperparasite dormancy within its host's puparia.

METHODS

Study system and field sampling

METHODS

Study system and field sampling

We collected *Protocalliphora* puparia between 2004 and 2019 within a network of 400 nest boxes

distributed equally among 40 farms located along a 10,200-km² gradient of agricultural intensity

in Southern Québec, Canada (Ghilain & Bélisle 2008). Nest boxes were placed ~1.5 m above

ground and spaced by ~50 m along a linear transect that followed field edges. While the primary

objective of the above network initiated in 2004 was to quantify the impact of land use on Tree

Swallow breeding ecology, it has evolved into a comprehensive platform for diverse

investigations, including the effects of *Protocalliphora* parasitic load on the swallows' breeding

success (Daoust et al. 2012; Pigeon et al. 2012; Sigouin et al. 2021). At the end of each breeding

season, we collected and bagged the entire content of nest boxes exclusively occupied by Tree

Swallows, a single-brooded species. Collected nests were stored at 4°C prior to processing, which

included 2 days at -80°C to kill any remaining living organisms before sorting under a ventilated

hood. *Protocalliphora* puparia found within nest material were counted and stored in vials filled

with 75% ethanol and referenced according to farm and nestbox IDs and year of sampling. Note

that *Protocalliphora* puparia from 2008 to 2009 were collected under two protocols by Daoust et

al. (2012) to raise some pupae in the lab and collect emerging flies or *Nasonia* wasps. As a result,

some specimens were no longer available for further examination in this study (i.e., for

hyperparasitism, species composition, and morphometry). Nevertheless, our dataset allowed us to

investigate nest infestation and parasitic load using nests from all nest boxes occupied by Tree

Swallows with a single clutch event leading to ≥ 1 nestling on a given year between, 2004 and

2019 (including 2008 and 2009), for a total of 2,673 nests (Table 1).

Specimen examination

Due to the considerable number of *Protocalliphora* puparia and the significant time commitment

required for individual specimen examination, a subsample of 263 infested nests was selected for

hyperparasitism assessment, species identification, and morphometric measurements of puparia

found therein (Table 1). Infested nests were subsampled as follows: two infested nests (when

available) were randomly sampled from each of 10 farms randomly selected to represent the

gradient of agricultural intensity on each year (excluding 2008 and 2009). When selected farms

had nests with no or very low levels of infestation by *Protocalliphora* (i.e., < 100 puparia in total

for all 10 nest boxes of a farm in a given year), additional nests, or farms if needed, were randomly

selected until a minimum of 100 puparia could be processed for a given year. All *Protocalliphora*

specimens from the resulting subsample of 263 nests (n = 3,994 puparia) were examined and only

those with adequate structural integrity for the determination or measurement of a given attribute

were considered in data analyses regarding hyperparasitism, species composition, and

morphometry (Table 1).

Hyperparasitism of Protocalliphora by Nasonia

Species-level identification of *Nasonia* wasps was not conducted in the present work due to fall

sample collection when most hyperparasites had already emerged and those that did not were killed

before completing their development. However, early subsampling and rearing of *Protocalliphora*

pupae by Daoust et al. (2012) within the same study system allowed the post-emergence

identification of two *Nasonia* species, *N. vitripennis* and *N. giraulti*, the first accounting for > 90%

of *Protocalliphora* infestation events.

The occurrence of *Nasonia* hyperparasitism on a *Protocalliphora* puparium is typically diagnosed

by the presence of one or more exit holes of 0.5-1 mm in diameter. However, *Nasonia* can also die

or overwinter as pupae inside their host instead of emerging at the end of the active season

(Garrido-Bautista 2019). Because puparia containing a dead *Protocalliphora* pupa and those

containing unemerged *Nasonia* will look identical, we systematically dissected *Protocalliphora*

puparia showing no sign of either fly or wasp emergence to verify the occurrence of *Nasonia*

hyperparasites to discriminate pupation failures linked to hyperparasitism from those resulting

from other causes. All 3,608 *Protocalliphora* puparia assessed for *Nasonia* infestation were hence

classified into one of the following categories: no evidence of *Nasonia* hyperparasitism, presence

of *Nasonia* emergence holes, and presence of unemerged *Nasonia* pupae without exit holes.

Protocalliphora species composition

We identified 3,009 well-preserved *Protocalliphora* puparia based on morphology following

Sabrosky et al. (1989), and Whitworth (2002, 2003, 2019), with additional feedback from T.L.

Whitworth (pers. comm.). Consistently with Daoust et al. (2012), who conducted research within

the same study system, only three species were identified: *Protocalliphora sialia* Shannon *&*

Dobroscky, *Protocalliphora bennetti* (Whitworth)*, and *Protocalliphora metallica* (Townsend)*.

Protocalliphora morphometry

To investigate inter and intra-specific variability in maximum length and width of *Protocalliphora*

puparia, each puparium was photographed using a Leica M165FC stereomicroscope equipped with

a 0.63X planapo objective, a Leica MC170HD color camera, and the Leica LAS X image-capture

software (v 3.0.3). All measurements were performed with Fiji image analysis software

(Schindelin et al. 2012). This allowed us to model any photographed puparium as an ellipsoidal-

shaped object and to calculate its theoretical maximum length and width. This approach minimized

measurement biases as puparia from which a *Protocalliphora* had successfully emerged would

otherwise appear shorter.

Data analysis

Interannual variability in Tree Swallow nest infestation by *Protocalliphora* between 2004 and

2019 was first evaluated by calculating, for each sampling year, the proportion of infested nests as

well as the average number of *Protocalliphora* puparia per infested nest (±SD). We then assessed

how the annual mean nest infestation rate across farms varied in time between 2004 and 2019, and

to what level each farm followed that global trend, if any, using a Generalized Linear Additive

Mixed Model (GAMM) fitted with the mgcv package (v.1.9-1; Wood 2023) in R (v.4.3.1, R Core

Team, 2023). More specifically, we modeled the proportion of infested nests using a global smooth

term for Year and group-level smooth terms allowing for farm-specific temporal trends that may

differ in shape and wiggliness from the global trend (i.e., as per Model *GI* in Pedersen et al. 2019).

The GAMM was fitted using a binomial distribution and logit link function. Temporal

autocorrelation in the residuals was assessed using the acf function in R (Simpson 2018). We then

performed the same exploratory analyses for the average number of *Protocalliphora* puparia per

infested nest with a GAMM that used a tweedie distribution and a log link function due to

substantial overdispersion with respect to a (truncated) Poisson distribution. Detailed codes, model

diagnostics and results for theses analyses are provided in Supplementary Material 1.

Interannual variability in *Nasonia* hyperparasitism of *Protocalliphora* puparia was evaluated by

contrasting the yearly proportion of non-infested to infested puparia and the proportion of infested

puparia from which *Nasonia* wasps had emerged against those still containing *Nasonia* pupae or

adults. Interannual variability in *Protocalliphora* species composition was assessed by calculating

the yearly relative proportion of each species for all specimens that could be unambiguously

identified. Hyperparasitism rates and species composition from 2008-2009 were obtained from

Daoust et al. (2012).

Intra- and between-species variability in the maximum length and width of puparia of

Protocalliphora sialia, *Protocalliphora bennetti*, and *Protocalliphora metallica* were first

assessed using density plots and boxplots based on the raw data. We then assessed inter-specific

differences in mean puparium maximum length and width using Generalized Linear Mixed Models

(GLMMs) fitted with the glmmTMB R package (Brooks et al. 2023). In this analysis

Protocalliphora bennetti, the intermediate-size species (Sabrosky et al. 1989; Whitworth 2002),

was designated as the reference category. Farm ID and Year were included as random effects to

account for environmental variation and to control for potential confounding effects. Nestbox ID

has been included as a random effect, but this led to convergence issues due to the low number of

repeated measures within each nest. Therefore, this variable was omitted from the analysis. The

proportion of variance explained by Farm ID and Year random effects were estimated following

Nakagawa and Schielzeth (2013). Species-specific maximum length and width being

approximately normally distributed, GLMMs were fitted using a Gaussian distribution and identity

link.

RESULTS

Spatiotemporal variation in Protocalliphora prevalence and load

Protocalliphora nest infestation occurred widely throughout the study area. All 40 monitored

farms harbored ≥ 1 infested Tree Swallow nest at some point between 2004 and 2019 and between

67.65% and 92.31% of farms had ≥ 1 infested nest depending on year. Overall, 45.90% (1,227) of

the 2,673 Tree Swallow nests with a single clutch event leading to ≥ 1 nestling were infested by

≥ 1 *Protocalliphora* larva based on counts of puparia. The proportion of infested nests varied

substantially both among years (range: 28.76% in 2012 to 69.33% in 2008; Fig.1A) and farms

(range: 13.37% to 79.49%). While most farms showed yearly fluctuations in nest infestation rate

that followed the global, nonlinear trend observed across the study area, about 30% of farms

showed mild to strong departures from that global trend (n = 587; GAMM R^2= 48.8%;

Supplementary Material 1). Temporal autocorrelation in the GAMM residuals at time lags of 1, 2,

3 and 5 years supported the presence of a cyclical nest infestation dynamics with a ~3-year period

at the scale of the study area (Supplementary Material 1).

A total of 22,630 puparia were counted among infested nests, resulting in an overall estimated

average load (± SD) of 18.44 ± 19.16 *Protocalliphora* per infested nest. The average number of

Protocalliphora per infested nest exhibited variability both within and across years, fluctuating

between 10.89 ± 8.08 (2012) and 27.40 ± 23.90 (2008) with a yearly coefficient of variation

ranging between 74.21 and 97.11% (Fig. 1A). The overall frequency distribution of the number of

Protocalliphora per infested nest was, however, highly skewed to the right with a median of 12

puparia (range: 1-141 puparia; Fig. 1B). Interestingly, most farms showed yearly fluctuations in

average *Protocalliphora* loads per infested nest that followed a global, nonlinear trend, which

peaks and troughs coincided with those observed for the proportion of nests infested by

Protocalliphora (n = 1,227; GAMM R^2= 12.2%; Supplementary Material 1). However, the farms

that deviated from the global load trend (i.e., 22.5% of farms) were not the same as for the

proportion of infested nests and the GAMM residuals showed no temporal autocorrelation pattern.

Hyperparasitism of Protocalliphora by Nasonia

Nasonia likely occurred throughout the study area as they were detected at least once in 30 of the

31 farms from which *Protocalliphora* puparia were subsampled to estimate hyperparasitism rate.

Overall, *Nasonia* hyperparasitized ≥ 1 puparium in 85.08% of the *Protocalliphora* infested nests

and 68.82% of the 3,608 subsampled puparia that could be examined (Table.1). Importantly,

12.81% of these hyperparasitized *Protocalliphora* puparia contained non-emerged *Nasonia*

individuals. These non-emerged individuals occurred nearly exclusively at the pupal stage,

suggesting that they either failed to produce adults or entered diapause for the coming winter.

Hyperparasitism by *Nasonia* remained high across years, ranging between 48.25% (2006) and

90.48% (2012) of *Protocalliphora* puparia (Fig.2). Note that Daoust et al. (2012) estimated

hyperparasitism rates in 2008 and 2009 based only on the presence of exit holes made by *Nasonia*

to exit puparia and reported among the lowest hyperparasitism rates of the study period (i.e.,

50.35% and 39.29%, respectively). The proportions of hyperparasitized *Protocalliphora* puparia

from which *Nasonia* did not emerge before winter varied substantially, ranging from 3.07% in

2006 to 16.36% in 2019 (Fig.2).

Protocalliphora species composition

Of the 3,009 *Protocalliphora* puparia suitable for species identification, 82.62% belonged to *P.*

sialia, 15.75% to *P. bennetti* (15.75%) and 3.62% to *P. metallica* (Table 1). *Protocalliphora sialia*

was consistently the most abundant species in each year, ranging from of 61.72% in 2011 to

98.41% the following year (Fig. 3). In contrast, *P. bennetti* exhibited more pronounced fluctuations

in relative abundance, ranging from quasi absence (0.79%) in 2012 to 32.01% in 2011.

Protocalliphora metallica, the least abundant species in our Tree Swallow study system, also

showed important variation in its relative abundance. It was either completely absent from our

subsamples in 2005 and 2016 or equally to more abundant than *P. bennetti* in 2004, 2012 and 2017,

but never exceeded 9.27% (2017) of *Protocalliphora* puparia in any given year (Fig.3). Data

extrapolated from Daoust et al. (2012) for 2008 and 2009 also showed a strong dominance of *P.*

sialia, with proportions of 96.59% and 90.74%, respectively. Overall, all three species were found

over most of the study area (Supplementary Material 2).

Protocalliphora morphometry

The maximum puparium length and width of the three *Protocalliphora* species were obtained from

a subsample of 1,982 individuals (Table 1). The distribution of maximum length and width of the

three species were unimodal and relatively symmetrical, though slightly skewed to the left in *P.*

metallica (Fig. 4). Despite substantial intraspecific variation leading to distributional overlap, the

GLMMs showed species differed in their average morphometry. *P. bennetti* exhibited a mean

puparium length ($\pm$ SD) of 8.18 $\pm$ 0.67 mm (range: 5.24-9.93 mm), while *P. metallica* showed a

significantly shorter mean length (7.34 $\pm$ 0.71 mm; range: 5.82-9.29 mm; *P* < 0.001), and *P. sialia*

a longer one (8.74 $\pm$ 0.58 mm; range: 5.70-10.27 mm; *P* < 0.001). Regarding width, *P. metallica*

had a narrower puparium (3.36 ± 0.41 mm, range: 2.25-4.16 mm; $P < 0.001$) and *P. sialia* a wider

one (3.89 ± 0.32 mm, range: 1.89-5.50 mm; $P < 0.001$) compared to *P. bennetti* (3.50 ± 0.39 mm,

range: 1.84-4.41 mm), respectively. The proportion of variance in length and width explained by

both the random (i.e., Farm ID and Year) and fixed (i.e., species) effects of the linear mixed models

was 36.45% and 30.57% for the maximum puparium length and width, respectively. Random

effects alone accounted for 51.58% and 44.16% of these proportions, respectively. These results

suggest that morphometric variation of *Protocalliphora* puparia is strongly caused by

agroenvironmental variables that have not been taken into account in the analysis.

DISCUSSION

Our study presents a rare time series on the prevalence of bird nest infestation by *Protocalliphora* blowflies and their abundance within infested nests. We found that infestation rates and loads of *Protocalliphora* within Tree Swallow nests were high over a 16-year period (2004-2019), yet widely variable in both space and time along a 10,200-km^2 gradient of agricultural intensity in southern Québec, Canada. Moreover, we found that these large temporal fluctuations were indicative of a cyclical dynamics that was shared by ~75% of the sampled locations (i.e., farms) and thus likely driven by both large-scale and local conditions or processes. Our study also shows that although *Protocalliphora* pupae were subjected to hyperparasitic pressures by *Nasonia* wasps that can vary as much as double among years, these were high enough to kill more than half of the *Protocalliphora* pupae produced in most years. Such an impact suggests that *Protocalliphora* and *Nasonia* may be part of a host-parasitoid dynamics responsible for the oscillations in *Protocalliphora* infestation rates and loads that we report. Consistent with the previous 2-year investigation in our study system by Daoust et al. (2012), we found both mono- and multispecific infestations in nests, involving three species of *Protocalliphora*, with *P. sialia* clearly dominating (>80% of puparia).. However, we found higher proportions of *P. bennetti* (~2x) and *P. metallica* (~4x) than reported by Daoust et al. (2012), emphasizing the value of multi-year studies in revealing the complete range of species composition variability in these flies. Lastly, we report highly variable puparium length and width with overlapping frequency distributions among *Protocalliphora* species. This finding suggests that these morphometric traits are not reliable as discrete diagnostic characters and not to be used in isolation for identifying *Protocalliphora* species (Whitworth 2002). It also raises important questions about the ecological factors driving

this variability and the potential consequences of size variation for emergence success and host

selection by parasitoids.

Spatiotemporal variation in Protocalliphora prevalence and load

The prevalence and intensity of bird nest infestations by *Protocalliphora* have generally been

found to vary widely in both space and time. For example, Bennett and Whitworth's (1992)

comprehensive examination across 73 bird species and 4,668 sampled nests revealed infestation

rates of 52% within the Rocky Mountain region (Utah, USA) versus 24% in the Great Lakes region

(Ontario, Canada). The mechanisms underlying such variation likely involve multiple dependent

and interactive factors and few studies have attempted to quantify the importance and

spatiotemporal scales at which they operate. While large-scale geographic variation may result

from differences in climate, landscape habitat composition, bird host community (species

composition and abundance), and hyperparasitic pressure (Sabrosky et al. 1989; Bennett and

Whitworth 1992; Daoust et al., 2012a, 2012b; Garrido-Bautista et al. 2019), local variation may

origin from factors as specific as bird host brood size and hatching date (Williams 2017; Wolfe-

Merritt et al. 2022), and nest type and material affecting the nest microclimate (Dawson et al. 2011;

Griebel et al. 2020). With respect to Tree Swallows breeding within nest boxes, previous research

reported *Protocalliphora* infestation rates ranging from 89% to 100% in most study sites across

Canada (Rendell & Verbeek 1996; Dawson et al. 2005; Gentes et al. 2007; Griebel et al. 2020).

Infestation rates are generally much higher than those we found either across years or farms within

our study area (overall rate = 45.90%). In contrast to other study sites, many of which were located

near wetlands, ours were located on farms distributed along a gradient of agricultural intensity

with very low, if any, wetland cover (Garrett et al. 2022). Hence, we hypothesize that the much

lower infestation rates we observed is linked to intensive agricultural practices that may impact *Protocalliphora* through multiple pathways, including directly through wetland drainage or pesticide use, or indirectly by reducing the density of their bird hosts (e.g., Robert et al. 2019; Wagner et al. 2021; Powell et al. 2024). Evidence for such negative impacts has also been reported in our study area by Daoust et al. (2012b) who found that lower loads of *Protocalliphora* in Tree Swallow nests located occurred within agro-intensive landscape during a 2-year period (2008-2009).

Annual infestation rates of Tree Swallow nests by *Protocalliphora* varied between 29% and 63% across the 16 years that spanned our study. This level of interannual variation, which was observed at the scale of the entire study area (i.e., 10,200-km^2), was comparable to what we observed among nestboxes from a single farm (i.e., < 1 km^2 ; Supplementary Material 3). Wesolowski (2001) also reported highly variable annual nest infestation rates by *Protocalliphora* in Marsh Tits (*Parus palustris*) that ranged from 27% to 88% in four 0.5-km^2 plots separated by ~1-2 km during an 8-year period in eastern Poland. Similarly, Musgrave et al. (2019), found annual infestation rates between 30% and 91% in Western Bluebird (*Sialia mexicana*) nests monitored over 17 years in a ~100-km^2 area of northern New Mexico (USA). Despite potential confounding effects, the above comparisons suggest that nest infestation rates by *Protocalliphora* typically vary substantially among years across a range of spatial scales. Given the lack of reported time series of infestation rates (and loads; but see Merino et al. 2024) in the literature, the level of temporal autocorrelation among years and how it varies across spatial scales are largely unknown and prevent the investigation of the ecological mechanisms underlying such fluctuations. Our observation that yearly fluctuations in both *Protocalliphora* nest infestation rates and loads were to some degree cyclical and synchronous among ~75% of the farms sampled over 10,200 km^2 suggest that both

regional (e.g., climatic) and local (e.g., crop rotations), as well as some density-dependent (e.g.,

hyperparasitic pressure by *Nasonia* wasps) processes may be at play.

The observed loads of *Protocalliphora* within infested nests exhibited a right-skewed distribution,

a phenomenon commonly observed in the field in parasitology (Wilson and Grenfell 1997).

Furthermore, a high degree of variability between nests, between sites, and across years were

reported (Griebel and Dawson 2019; Remeš and Krist, 2005). Part of the interannual variation is

often associated with weather (Merino & Potti 1996; Dawson et al. 2005; Mennerat et al. 2021;

Castaño-Vazquez et al. 2021; Merino et al. 2024) and spatial variation is typically linked to micro-

habitat and nest-specific factors, such as the amount of feather lining or the species and number of

nestling hosts (Eeva et al. 1994; Merino & Potti 2001; Tomás et al. 2007; Dawson et al. 2011;

Williams 2017; Wolfe-Merritt et al. 2022). In our study system, part of the between-farm variation

likely results from the diversified agricultural contexts in which Tree Swallows breed. Daoust et

al. (2012b) found fewer *Protocalliphora* in Tree Swallow nests located in more agrointensive

areas, particularly in maize and soybean monocultures. This pattern could result from several direct

and indirect effects associated with intensive agriculture, such as reduced shelter and food

resources for adult flies, lower avian host availability at both the landscape and brood scale and

increased toxicological effects from pesticide exposure either directly or through their avian hosts

(Daoust et al. 2012). Given the rarity of studies that examined the effects of agricultural intensity

on ectoparasite of wild birds, more extensive and long-term research is needed to determine the

mechanisms by which agriculture generates spatiotemporal variation and ultimately shapes host-

parasite dynamics.

Hyperparasitism of Protocalliphora by Nasonia

Interannual variation in the proportions of *Protocalliphora* puparia hyperparasitized by *Nasonia* wasps showed no clear patterns while ranging between 48.25% and 90.48% among years, with an overall rate of 68.82%. These proportions are generally higher from those of Jamriška & Lučeničová (2019), who observed an average hyperparasitism rate of 37.91% over four years in Slovakia, and from those of Bennett & Whitworth (1991), who reported an overall rate of ~17% from two regions of North America, namely Utah and Ontario. In contrast, Grillenberger et al. (2009) reported a rate of ~44% in New York state (USA), while Garrido-Bautista et al. (2020) reported rates from 47% to 56% for southern Spain, and from 61% to 79% for central Spain, figures much closer to our own. While more extensive longitudinal and ecological studies are needed, current findings suggest that geographical and environmental contexts may significantly influence the likelihood of *Nasonia* hyperparasitism, a primary determinant of *Protocalliphora* emergence success. However, it is crucial to recognize that, in addition to the substantial control exerted by *Nasonia* wasps on *Protocalliphora* communities, uninfested *Protocalliphora* individuals may also die during pupation due to other ecological factors, such as environmental contamination. The specific impacts of those factors on *Protocalliphora* emergence success have yet to be thoroughly investigated and quantified.

Nasonia parasitoid wasps have mostly served as model species in numerous laboratory studies addressing themes in genetics, developmental biology, and behavioral ecology (Werren and Loehlin 2009). The ecological factors driving host selection and hyperparasitism by *Nasonia* under natural conditions are thus poorly understood (Garrido-Bautista 2019; Peters 2010). This is particularly true given the wide diversity of hosts in which they can lay eggs and the potential for regional variations due to their widespread distribution. This leaves the causes of yearly

fluctuations in *Protocalliphora* puparia hyperparasitic events or of dormancy within the host

puparia unresolved. Future research on *Nasonia* ecology would likely benefit from considering

factors such as vegetation type and density, as these features may influence host hyperparasitism

by affecting the availability of food and shelter (Wylies 1958). Avian nest availability associated

with these environmental factors could also potentially induce host-dependent effects, contributing

to an increased incidence of hyperparasitism on *Protocalliphora*.

This study reports the first quantitative exploration of dormancy and/or mortality of *Nasonia*

within their host in the wild. We noted consistently low yearly proportions of *Protocalliphora*

puparia containing dormant or dead *Nasonia* (range: 3.07-16.36%), with no discernible temporal

trend. Several biotic and abiotic factors, such as weather, luminosity or host and parasitoid body

condition, may affect the production of diapausing *Nasonia* offspring (Saunders et al. 1969; Yoder

et al. 1994; Bertossa et al. 2010; Li et al. 2014; Paolucci et al. 2013). Mechanisms linked to

phenology could thereby impact the dormancy and survival of *Nasonia* larvae. For instance,

Protocalliphora pupation phenology, which in turn determines host availability and development

conditions for *Nasonia*, must partly depend on avian host breeding phenology. Moreover,

agricultural intensity may affect *Nasonia* development and emergence success as pesticide use can

alter hyperparasitoid wasps cognition and fitness (Schöfer et al. 2023, 2024; Sulg et al. 2023), as

well as the availability and quality of their hosts (Li et al. 2014; Daoust et al. 2012).

Protocalliphora species composition and morphometry

Bird nestlings from the Nearctic region are hosts to at least 28 *Protocalliphora* species (Sabrosky

et al. 1989; updated by Whitworth 2002). While some bird blowfly species appear to specialize on

specific hosts through behavioral adaptations, some parasitize a wide array of bird species (Bennett

& Whitworth 1992), and the occurrence of multiple *Protocalliphora* species within a single host

nest is not a rare event (Daoust et al. 2012). Our longitudinal study, one of the very few that

describe *Protocalliphora* species composition, reports that such compositions can vary

substantially across years. Indeed, our results show that certain species (in this case, *P. metallica*)

may be practically absent in some years, only to reappear in relatively high numbers the following

year. Even the highly prevalent *P. sialia* fluctuated widely, representing 58.93% of the

Protocalliphora specimens in 2011 to being almost the only species found in infested nests the

following year (98.32%). It thus seems essential to conduct longitudinal studies considering

variability in species composition before drawing any conclusions about host specialization to

avian parasites and mixed infestations.

While descriptive statistics of some morphometric measurements of *Protocalliphora* puparia can

be found in the literature, few, if any, studies report the actual frequency distributions of these

features that may help discriminating among species of this cryptic group (Heeb et al. 2000;

Whitworth 2002, 2003; Jamriška & Modry 2013). The mean, standard deviation, and range

estimates of puparium length and width for *P. bennetti, P. metallica, and P. sialia* that we obtained

via photogrammetry align well with published estimates (Whitworth et al. 1991; Whitworth 2002).

Yet, the fact that density plots highlighted substantial intraspecific variation in both puparium

length and width, as well as significant overlap among the three studied species, indicates that

these metrics should not be used on their own for *Protocalliphora* identification, even within a

small regional species pool. Nevertheless, the ecological factors contributing to this high

intraspecific variation, with certain puparia being more than twice as large and long as others, is

unclear. Sexual dimorphism (Sabrosky et al. 1989) or factors like host availability and quality, as

well as developmental period and temperature, which are often interconnected, have been

hypothesized as causes of size variation in *Protocalliphora*. For instance, underfed larvae may

conclude their development as 'runts' (Sabrosky et al. 1989; Bennett & Whitworth 1991).

Alternatively, intensive agricultural practices, including pesticide use, have been reported as either

direct or indirect drivers of phenotypic variation (Wisniewska & Prokopy 1997; Verheyen & Stoks

2019; Gérard et al. 2022; Ponce-Méndez et al. 2022) and pupation success (Heneberg et al. 2020;

Mahdjoub et al. 2020) in non-targeted arthropod species.

All the ecological parameters we studied showed substantial variation in space, time, or among

individuals, warranting that future studies addressing the mechanisms driving host-parasite

dynamics involving *Protocalliphora* adopt a long-term monitoring and multi-trophic approach in

contrasting environments. Investigating the tri-trophic relationships involving birds,

Protocalliphora, and *Nasonia* across diverse agroecosystems presents a promising avenue for such

research. Given the complex interplay among environmental conditions and multi-trophic

interactions, we further suggest that such endeavor be tackled using a causal inference framework

to limit biases, especially given that each trophic level may respond to different factors, at different

spatiotemporal scales, and potentially nonlinearly or in different directions, which has the potential

to amplify, attenuate, and even create or mask effects (Arif and MacNeil 2023; Grace 2024).

ACKNOWLEDGMENTS

We are deeply grateful to the many farm owners who have collaborated to our long-term

monitoring program. We also thank U de Sherbrooke for its institutional support and the funding

agencies that made this research possible, including the Natural Sciences and Engineering

Research Council of Canada (NSERC), the Fonds de recherche du Québec – Nature et technologies

(FRQ-NT), and the Canada Research Chairs program. We warmly acknowledge the contribution

of the many students, field assistants, and technicians, and especially L. Glaude and Y. Sageau,

who participated to the collection and sorting of nests over the years. This study was conducted

with the approval of U de Sherbrooke animal care committee, and we gratefully recognize all those

whose efforts, on the ground and behind the scenes, contributed to making this work possible.

REFERENCES

Arif, S., MacNeil, M. A., 2023. Applying the structural causal model framework for observational causal inference in ecology. *Ecological Monographs* 93, e1554. https://doi.org/10.1002/ecm.1554

Bańbura, J., Perret, P., Blondel, J., Thomas, D. W., Cartan-Son, M., Lambrechts, M. M., 2004. Effects of Protocalliphora parasites on nestling food composition in Corsican Blue Tits *Parus caeruleus*: consequences for nestling performance. *Acta Ornithologica* 39(2), 93–103.

Bennett, G. F., Whitworth, T. L., 1991. Studies on the life history of some species of Protocalliphora (Diptera: Calliphoridae). *Canadian Journal of Zoology* 69(8), 2048–2058.

Bennett, G. F., Whitworth, T. L., 1992. Host, nest, and ecological relationships of species of Protocalliphora (Diptera: Calliphoridae). *Canadian Journal of Zoology* 70(1), 51–61.

Bertossa, R. C., van Dijk, J., Beersma, D. G., Beukeboom, L. W., 2010. Circadian rhythms of adult emergence and activity but not eclosion in males of the parasitic wasp *Nasonia vitripennis*. *Journal of Insect Physiology* 56(7), 805–812.

Brooks, M., Bolker, B., Kristensen, K., Maechler, M., Magnusson, A., McGillycuddy, M., 2023. Package 'glmmtmb'. *R Package Version* 1(1), 7.

Castaño-Vázquez, F., Schumm, Y. R., Bentele, A., Quillfeldt, P., Merino, S., 2021. Experimental manipulation of cavity temperature produces differential effects on parasite abundances in blue tit nests at two different latitudes. *International Journal for Parasitology: Parasites and Wildlife* 14, 287–297.

Clayton, D. H., Koop, J. A., Harbison, C. W., Moyer, B. R., Bush, S. E., 2010. How birds combat ectoparasites.=

Daoust, S. P., Savage, J., Whitworth, T. L., Bélisle, M., Brodeur, J., 2012a. Diversity and abundance of ectoparasitic blow flies Protocalliphora (Diptera: Calliphoridae) and their Nasonia (Hymenoptera: Pteromalidae) parasitoids in tree swallow nests within agricultural lands of southern Québec, Canada. *Annals of the Entomological Society of America* 105(3), 471–478.

Daoust, S. P., Bélisle, M., Savage, J., Robillard, A., Baeta, R., Brodeur, J., 2012b. Direct and indirect effects of landscape structure on a tri-trophic system within agricultural lands. *Ecosphere* 3(11), 1–19.

Dawson, R. D., Hillen, K. K., Whitworth, T. L., 2005. Effects of experimental variation in temperature on larval densities of parasitic Protocalliphora (Diptera: Calliphoridae) in nests of tree swallows (Passeriformes: Hirundinidae). *Environmental Entomology* 34(3), 563–568.

Dawson, R. D., O'Brien, E. L., Mlynowski, T. J., 2011. The price of insulation: costs and benefits of feather delivery to nests for male tree swallows *Tachycineta bicolor*. *Journal of Avian Biology* 42(2), 93–102.

Deveux, M., Christiaens, O., Smagghe, G., Taning, N. T. C., 2023. *Nasonia vitripennis* as a parasitoid wasp model for pest-specific dsRNA safety assessment. *Entomologia Generalis* 43(1), 139–146.

Eeva, T., Lehikoinen, E., Nurmi, J., 1994. Effects of ectoparasites on breeding success of great tits (*Parus major*) and pied flycatchers (*Ficedula hypoleuca*) in an air pollution gradient. *Canadian Journal of Zoology* 72(4), 624–635.

Garrido-Bautista, J., Moreno-Rueda, G., Baz, A., Canal, D., Camacho, C., Cifrián, B., … Potti, J., 2020. Variation in parasitoidism of Protocalliphora azurea (Diptera: Calliphoridae) by *Nasonia vitripennis* (Hymenoptera: Pteromalidae) in Spain. *Parasitology Research* 119(2), 559–566.

Gentes, M. L., Whitworth, T. L., Waldner, C., Fenton, H., Smits, J. E., 2007. Tree swallows (*Tachycineta bicolor*) nesting on wetlands impacted by oil sands mining are highly parasitized by the bird blow fly Protocalliphora spp. *Journal of Wildlife Diseases* 43(2), 167–178.

Gérard, M., Baird, E., Breeze, T., Dominik, C., Michez, D., 2022. Impact of crop exposure and agricultural intensification on the phenotypic variation of bees. *Agriculture, Ecosystems & Environment* 338, 108107.

Ghilain, A., Bélisle, M., 2008. Breeding success of tree swallows along a gradient of agricultural intensification. *Ecological Applications* 18(5), 1140–1154.

Grace, J. B., 2024. An integrative paradigm for building causal knowledge. *Ecological Monographs* 94, e1628. https://doi.org/10.1002/ecm.1628

Grillenberger, B. K., Van De Zande, L., Bijlsma, R., Gadau, J., Beukeboom, L. W., 2009. Reproductive strategies under multiparasitism in natural populations of the parasitoid wasp *Nasonia* (Hymenoptera). *Journal of Evolutionary Biology* 22(3), 460–470.

Griebel, I. A., Dawson, R. D., 2019. Predictors of nestling survival during harsh weather events in an aerial insectivore, the Tree Swallow (*Tachycineta bicolor*). *Canadian Journal of Zoology* 97, 81–90.

Hannam, K., 2006. Ectoparasitic blow flies (*Protocalliphora spp.*) and nestling Eastern Bluebirds (*Sialia sialis*): direct effects and compensatory strategies. *Canadian Journal of Zoology* 84(7), 921–930.

Heeb, P., Kölliker, M., Richner, H., 2000. Bird-ectoparasite interactions, nest humidity, and ectoparasite community structure. *Ecology* 81(4), 958–968.

Heneberg, P., Bogusch, P., Astapenková, A., Řezáč, M., 2020. Neonicotinoid insecticides hinder
the pupation and metamorphosis into adults in a crabronid wasp. *Scientific Reports* 10(1), 7077.

Jánošková, V., Országh, I., Jamriška, J., Kopáni, M., 2010. Pre-imaginal stages of the blowfly
Protocalliphora falcozi in nests of the tree sparrow (*Passer montanus*). *Entomologica Fennica*
21(2), 107–116.

Jamriška, J, Lucenicova, T. Impact of the parasitic wasp *Nasonia* spp. (Walker, 1836)
Hymenoptera: Pteromalidae) on populations of bird blowflies *Protocalliphora* spp. Hough, 1899
(Diptera: Calliphoridae). J Microbiol Exp. 2019;7(1):6-7. DOI: 10.15406/jmen.2019.07.00230

Jamriška, J., Modrý, D., 2013. Description of the Puparium of Protocalliphora nourtevai (Insecta:
Diptera: Calliphoridae). *The Journal of Parasitology* 99(5), 896–898.

Li, Y., Zhang, L., Zhang, Q., Chen, H., Denlinger, D. L., 2014. Host diapause status and host
diets augmented with cryoprotectants enhance cold hardiness in the parasitoid Nasonia
vitripennis. *Journal of Insect Physiology* 70, 8–14.

Lučeničová, T., Nordt, A., Klenke, R., Jamriška, J., 2013. Prevalence of bird blow fly infestation
in nests of the European Blackbird (*Turdus merula*). *The Wilson Journal of Ornithology* 125(4),
848–851.

Mahdjoub, H., Blanckenhorn, W. U., Lüpold, S., Roy, J., Gourgoulianni, N., Khelifa, R., 2020.
Fitness consequences of the combined effects of veterinary and agricultural pesticides on a non-
target insect. *Chemosphere* 250, 126271.

Mennerat, A., Charmantier, A., Perret, P., Hurtrez-Boussès, S., Lambrechts, M. M., 2021.
Parasite intensity is driven by temperature in a wild bird. *Peer Community Journal* 1.

Maziarz, M., Broughton, R. K., Chylarecki, P., Hebda, G., 2022. Weather impacts on interactions
between nesting birds, nest-dwelling ectoparasites and ants. *Scientific Reports* 12(1), 17845.

Marquardt, W. H. (Ed.), 2004. *Biology of Disease Vectors.* Elsevier.

Merino, S., Potti, J., 1996. Weather dependent effects of nest ectoparasites on their bird hosts.
Ecography 19(2), 107–113.

Merino, M., García-Del Río, M., Castaño-Vázquez, F., Merino, S., 2024. A long-term study on
the impact of climatic variables on two common nest-dwelling ectoparasites of the Eurasian blue
tit (*Cyanistes caeruleus*). *Integrative Zoology*.

Musgrave, K., Bartlow, A. W., Fair, J. M., 2019. Long-term variation in environmental
conditions influences host–parasite fitness. *Ecology and Evolution* 9(13), 7688–7703.

Paolucci, S., Van De Zande, L., Beukeboom, L. W., 2013. Adaptive latitudinal cline of
photoperiodic diapause induction in the parasitoid Nasonia vitripennis in Europe. *Journal of*
Evolutionary Biology 26(4), 705–718.

Peters, R. S., 2010. Host range and offspring quantities in natural populations of Nasonia vitripennis (Walker, 1836) (Hymenoptera: Chalcidoidea: Pteromalidae). *Journal of Hymenoptera Research* 19(1), 128–138.

Pigeon, G., Baeta, R., Bélisle, M., Garant, D., Pelletier, F., 2013. Effects of agricultural intensification and temperature on immune response to phytohemagglutinin in Tree Swallows (*Tachycineta bicolor*). *Canadian Journal of Zoology* 91(2), 56–63.

Ponce-Méndez, M., García-Martínez, M. A., Serna-Lagunes, R., Lasa-Covarrubias, R., Presa-Parra, E., Murguía-González, J., Llarena-Hernández, C., 2022. Local Agricultural Management Filters Morphological Traits of the South American Palm Weevil (*Rhynchophorus palmarum* L.; Coleoptera: Curculionidae) in Ornamental Palm Plantations. *Agronomy* 12(10), 2371.

Puchala, P., 2004. Detrimental effects of larval blow flies (*Protocalliphora azurea*) on nestlings and breeding success of Tree Sparrows (*Passer montanus*). *Canadian Journal of Zoology* 82(8), 1285–1290.

R Core Team, 2023. *R: A Language and Environment for Statistical Computing.* R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Remeš, V., Krist, M., 2005. Nest design and the abundance of parasitic Protocalliphora blow flies in two hole-nesting passerines. *Écoscience* 12, 549–553. https://doi.org/10.2980/i1195-6860-12-4-549.1

Rendell, W. B., Verbeek, N. A., 1996. Are avian ectoparasites more numerous in nest boxes with old nest material? *Canadian Journal of Zoology* 74(10), 1819–1825.

Robert, M., Hachey, M. H., Lepage, D., Couturier, A. R. (éd.), 2019. *Second Atlas of the Breeding Birds of Southern Québec.* Regroupement Québec Oiseaux; Service canadien de la faune (Environnement et Changement climatique Canada); Bird Studies Canada, Montréal, Québec, Canada.

Rothschild, M., Clay, T., 1957. *Fleas, flukes and cuckoos: A study of bird parasites.*

Sabrosky, C. W., Bennett, G. F., Whitworth, T. L., 1989. Bird blow flies (Protocalliphora) in North America (Diptera: Calliphoridae) with notes on Palearctic species. *Smithsonian Institution Press*.

Saunders, D. S., 1969. Diapause and photoperiodism in the parasitic wasp *Nasonia vitripennis*, with special reference to the nature of the photoperiodic clock. In *Symposia of the Society for Experimental Biology* 23, 301–329.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., … Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. *Nature Methods* 9, 676–682.

Schöfer, N., Ackermann, J., Hoheneder, J., Hofferberth, J., Ruther, J., 2023. Sublethal effects of four insecticides targeting cholinergic neurons on partner and host finding in the parasitic wasp *Nasonia vitripennis*. *Environmental Toxicology and Chemistry* 42(11), 2400–2411.

Schöfer, N., Ratschmann, G., Ruther, J. 2024. Effects of sub-nanogram doses of acetamiprid, dimethoate, flupyradifurone, and sulfoxaflor on courtship, mating, and olfactory host finding of the parasitic wasp *Lariophagus distinguendus*. *Entomologia Experimentalis et Applicata*, *172*(8), 666-678.

Sigouin, A., Bélisle, M., Garant, D., Pelletier, F., 2021. Agricultural pesticides and ectoparasites: potential combined effects on the physiology of a declining aerial insectivore. *Conservation Physiology* 9(1), coab025.

Simon, A., Thomas, D., Blondel, J., Perret, P., Lambrechts, M. M., 2004. Physiological ecology of Mediterranean blue tits (*Parus caeruleus L.*): effects of ectoparasites (*Protocalliphora spp.*) and food abundance on metabolic capacity of nestlings. *Physiological and Biochemical Zoology* 77(3), 492–501.

Stiner, F. M., 1969. Overwintering by *Protocalliphora metallica* (Diptera: Calliphoridae). *Annals of the Entomological Society of America* 62(5), 1205–1206. https://doi.org/10.1093/aesa/62.5.1205a

Sulg, S., Kaasik, R., Kallavus, T., Veromann, E., 2023. Toxicity of essential oils on cabbage seedpod weevil (Ceutorhynchus obstrictus) and a model parasitoid (*Nasonia vitripennis*). *Frontiers in Agronomy* 5, 1107201.

Tomás, G., Merino, S., Moreno, J., Morales, J., Martínez-de La Puente, J., 2007. Impact of blood parasites on immunoglobulin level and parental effort: a medication field experiment on a wild passerine. *Functional Ecology*, 125–133.

Verheyen, J., Stoks, R., 2019. Shrinking body size and physiology contribute to geographic variation and the higher toxicity of pesticides in a warming world. *Environmental Science & Technology* 53(19), 11515–11523.

Wagner, D. L., Grames, E. M., Forister, M. L., Berenbaum, M. R., Stopak, D., 2021. Insect decline in the Anthropocene: Death by a thousand cuts. *Proceedings of the National Academy of Sciences of the United States of America* 118(2), e2023989118.

Werren, J. H., Loehlin, D. W., 2009. The parasitoid wasp Nasonia: an emerging model system with haploid male genetics. *Cold Spring Harbor Protocols* 2009(10), pdb-emo134.

Wesołowski, T., 2001. Host–parasite interactions in natural holes: Marsh Tits (*Parus palustris*) and blow flies (Protocalliphora falcozi). *Journal of Zoology* 255(4), 495–503.

Whitworth, T. L., Bennett, G. F., 1992. Pathogenicity of larval Protocalliphora (Diptera: Calliphoridae) parasitizing nestling birds. *Canadian Journal of Zoology* 70(11), 2184–2191.

Whitworth, T. L., 2002. Two new species of North American Protocalliphora Hough (Diptera:
Calliphoridae) from bird nests. *Proceedings of the Entomological Society of Washington* 104,
801–811.

Whitworth, T. L., 2003. A key to the puparia of 27 species of North American Protocalliphora
Hough (Diptera: Calliphoridae) from bird nests and two new puparial descriptions. *Proceedings*
of the Entomological Society of Washington 105, 995–1033.

Whitworth, T. L., Dawson, R. D., Magalon, H., Baudry, E., 2007. DNA barcoding cannot reliably
identify species of the blowfly genus Protocalliphora (Diptera: Calliphoridae). *Proceedings of*
the Royal Society B: Biological Sciences 274(1619), 1731–1739.

Whitworth, T., 2019. Keys to the genera and species of blow flies (Diptera: Calliphoridae) of
America, North of Mexico. In *Forensic Entomology*, 413–443. CRC Press.

Williams, D., 2017. *Parasite-host interactions between Protocalliphora spp. (Diptera:*
Calliphoridae) and tree swallows Tachycineta bicolor. Doctoral dissertation, University of
Northern British Columbia.

Wilson, K., Grenfell, B. T., 1997. Generalized linear modelling for parasitologists. *Parasitology*
Today 13, 33–38. https://doi.org/10.1016/S0169-4758(96)40009-6

Wisniewska, J., Prokopy, R. J., 1997. Pesticide effect on faunal composition, abundance, and
body length of spiders (Araneae) in apple orchards. *Environmental Entomology* 26(4), 763–776.

Wolfe-Merritt, C. J., Hartman, L. E., Barragan, E. M., Hellman, B. M., Pigott, S., Rodríguez-
Ferraro, A., Tori, W. P., 2022. Identity and characteristics of feathers used as lining in Tree
Swallow (*Tachycineta bicolor*) nests in Indiana and Ohio. *The Wilson Journal of Ornithology*
134(3), 473–484.

Wylie, H. G., 1958. Factors That Affect Host Finding by *Nasonia vitripennis* (Walker)
(Hymenoptera: Pteromalidae). *The Canadian Entomologist* 90(10), 597–608.

Yoder, J. A., Rivers, D. B., Denlinger, D. L., 1994. Water relationships in the ectoparasitoid
Nasonia vitripennis during larval diapause. *Physiological Entomology* 19(4), 373–378.

Table 1. Number of Tree Swallow nests and *Protocalliphora* puparia collected and analysed

between 2004 and 2019 in southern Québec, Canada. No puparia from 2008-2009 could be

included in the hyperparasitism, species composition, and morphometry datasets as they were

used in a previous by Daoust et al. (2012) and were no longer suitable for further processing.

Samples	Datasets			
	All data	Hyperparasitism	Species composition	Morphometry
# of nests examined	2,673	263	263	263
# of nests infested by *Protocalliphora* with > 1 puparium suitable for processing	1,227	259	216	174
# of nests infested by *Nasonia*	-	218	-	-
# of puparia examined	22,630	3,994	3,994	3,994
# of puparia excluded (damaged)		386	985	2,012
# of puparia without *Nasonia*	-	1,125	-	-
# of puparia with *Nasonia*	-	2,483	-	-
# of puparia with non-emerged *Nasonia*	-	318	-	-
# of puparia of *P. sialia*	-	-	2,426	1,569
# of puparia of *P. bennetti*	-	-	474	347
# of puparia of *P. metallica*	-	-	109	66

Figure 1. A) Patterns of Tree Swallow nest infestation by *Protocalliphora* observed between

2004 and 2019 in southern Québec, Canada. Left y-axis and corresponding red dots (± error

bars) show the yearly average (± SD) number of *Protocalliphora* puparia per infested nest. Right

y-axis and corresponding bars show the yearly number of infested nests (grey) over the yearly

total number of examined nests (i.e., nests that harboured a single clutch that experienced at least

one hatching event; white). **B**) Frequency distribution of the number of *Protocalliphora* puparia

found within infested nest ($n = 1,227$).

Figure 2. Interannual rates of *Protocalliphora* puparia hyperparasitism by *Nasonia* within Tree

Swallow nests collected in southern Québec, Canada, between 2004 and 2019. *Protocalliphora*

puparia were classified as uninfested when they showed no signs of hyperparasitism, and as

infested, if *Nasonia* exit holes were observed on the puparia or *Nasonia* first instars were found

when *Protocalliphora* puparia were dissected. (*) Proportions for 2008 and 2009 were derived

from Daoust et al. (2012; Table 3) based only on the presence of holes made by *Nasonia* to exit

puparia.

Figure 3. Interannual *Protocalliphora* species composition within Tree Swallow nests collected

in southern Québec, Canada, between 2004 and 2019. Three species were identified (*P. bennetti*,

light green; *P. metallica*, grey; *P. sialia*, dark green). (*) Proportions for 2008 and 2009 were

derived from Daoust et al. (2012; Table 1).

Figure 4. Density distribution of morphometric measurements (length and width) of

Protocalliphora puparia for *P. bennetti* (light green, n = 347), *P. metallica* (grey, n = 66), and *P.*

sialia (dark green, n = 1,569) collected in southern Québec, Canada, between 2004 and 2019.

Mean (dotted line) and quartiles (via boxplot) are shown.