C-Nucleosides Stabilize RNA by Reducing Nucleophilicity at 2'-OH

Dipanwita Banerjee,[1] Lu Xiao,[1] Pavitra S. Thacker,[1] Jayanta Kundu,[2] Muthiah Manoharan,[2] and Eric T. Kool[1*]

[1]Department of Chemistry, Stanford University, Stanford, CA, USA; [2]Alnylam Pharmaceuticals, Cambridge, MA, USA

*Corresponding author: kool@stanford.edu

ABSTRACT

Nucleotides with carbon substitution for heteroatoms are common in biological and therapeutic RNAs. Important examples include the C-nucleosides pseudouridine and N1-methyl-pseudouridine; these modifications were reported to slow degradation of large RNAs, but the mechanism is unknown. We measured kinetics of thermal and enzymatic cleavage at a single bond of synthetically modified RNAs, and find that carbon substitution markedly reduces strand cleavage rates in RNA by both mechanisms. Studies of nucleophilic acylation reactions of RNAs and of small alcohols of varied pK_a suggest that reduced inductive effects resulting from carbon substitution for electronegative atoms results in both higher pK_a and lower nucleophilicity. The results provide insight into native transcriptome modifications as well as RNA therapies.

INTRODUCTION

Modified RNAs are under intensive development as drugs, genetic therapies, and vaccines. These bioactive species invariably incorporate numerous nucleobase and sugar modifications to provide enhanced stability to degradation and reduced innate immune responses.[1–3] Many of these modifications are inspired and adopted from modifications first identified in naturally occurring RNAs.[4-6] Over 150 chemical modifications have been characterized in native RNAs, adding a regulatory layer to RNA biology by affecting cellular processes such as gene expression,[4] splicing,[5] and translation.[6,7]

One of the chief motivations for incorporating ribonucleotide modifications in RNA therapies is the fact that the biopolymer is highly susceptible to degradation, resulting in strand cleavage due to thermal and enzymatic processes. This strand cleavage directly involves the 2'-hydroxyl (2'-OH) group of the ribose sugar, which undergoes intramolecular nucleophilic attack at the neighboring phosphate, breaking the 3'-5' phosphodiester bond in an in-line fashion and causing cleavage of the RNA (Figure 1).[8,9] This nucleophilic attack occurs spontaneously under thermal conditions, especially at elevated pH,[10] or is facilitated by nuclease enzymes that provide general acid/base catalysis and promote the required conformation changes for this attack.[11,12] Chemical modification or replacement of the 2'-OH can be beneficial, extending the lifetime of bioactive RNAs in the cell.[13,14] Studies of modified nucleotides have provided important outcomes in improving RNA therapies and vaccines for increasing target specificity and reducing immunogenicity.[2] However, mechanistic understanding of the effects of nucleotide modifications on RNA degradation is less well explored. Such understanding can provide both basic insights into existing naturally occurring modifications in RNA, as well as inspire improved designs for future RNA therapies.

Figure 1. The effects of pyrimidine nucleotide modifications on RNA stability. (A) Modifications that alter the nucleophilic reactivity of the 2′-OH group can affect RNA strand cleavage rates and acylation rates. (B) We hypothesize that modifications that reduce the number of electronegative N/O atoms (highlighted in blue) within two bonds of 2′-OH lower its acidity (increase its pK_a), as is known for small-molecule alcohols. This inductive electron-withdrawing effect of N/O stabilizes the oxyanion form of the hydroxyl; conversely, carbon substitution for these electronegative atoms decreases the population of the anionic form. (C) Strongly-stacking uridine analogs test the effects of base stacking on 2′-OH reactivity; increased stacking stabilizes the helical conformation, which has lower nucleophilic reactivity. Analogs with altered acidity will influence the fraction of the highly reactive oxyanion form. (D) Structures of uridine and analogs employed in this study, with differences highlighted in green.

Modified nucleotides can potentially protect RNA from degradation by multiple mechanisms. One clear mechanism is direct modification or replacement of the 2'-OH group to provide a chemical block to intramolecular nucleophilic attack. Examples of this strategy

include 2'-O-methyl groups, 2'-O-acyl groups, and 2'-fluoro groups, all of which completely prevent cleavage. However, other nucleotide modifications more remote from the 2'-OH group (Figure 1) also have the potential to enhance stability against strand cleavage. Kariko *et al.* reported that replacement of uridine by pseudouridine (Ψ) in mRNAs provides a measure of stabilization against degradation in cells; however, the mechanism by which this occurred was not determined.[15] This modified nucleoside is an isomer of uridine in which the base is attached to ribose through carbon rather than nitrogen, resulting in a C-nucleoside (Figure 1B). A more recent study showed that Ψ replacement of U enhances stability of an mRNA to thermal degradation.[16] In general, studies of thermal degradation of unmodified mRNAs have shown that single-stranded, flexible regions are hotspots for cleavage.[16] Thus it is plausible that base modifications that thermodynamically enhance stacking and helicity have the potential to indirectly stabilize RNAs against strand cleavage. Interestingly, thermodynamics studies have shown that Ψ stabilizes RNA helices moderately relative to uridine,[17] providing one possible mechanism for suppressing strand cleavage.

Here we also consider a different and previously unrecognized mechanism for RNA strand stabilization by carbon-substituted nucleotides, involving differences in the inherent nucleophilicity of the 2'-OH group. Studies of the nucleophilicity of the 2'-OH group toward acylating agents at physiological pH suggest that the main reactive form of this hydroxyl is the oxyanion (Figure 1C),[18] which is rare at pH 7.4 but rapidly equilibrates. The pK_a of the 2'-OH group in mononucleotides is approximately 12.5,[19] *ca.* 3 pK units lower that of than reference alcohols such as ethanol or isopropanol. We have suggested that the relative acidity of the 2'-hydroxyl group in canonical RNA is at least in part due to inductive effects of three highly electronegative atoms in proximity: the 3'-oxygen, the 4'-oxygen, and the N1 nitrogen of the base (Figure 1B).[20] In this light, we note that C-nucleosides commonly employed in therapeutic RNAs (Ψ and N1-methyl Ψ (m1Ψ)) have one of these three electronegative atoms replaced by carbon. Thus, it seems possible that the nucleophilic reactivity of the 2'-OH group might be

reduced in such C-nucleosides as a result of lower acidity; this might be expected to reduce rates of reactivity with electrophiles including both external acylating agents as well as the neighboring phosphodiester group (Figures 1B,C). To date, no kinetics data for these nucleosides have been available for either of these forms of reactivity.

In sum, multiple physical and chemical factors of nucleotide modifications have the potential to affect RNA strand stability against thermal and enzymatic cleavage. In this study, we analyze these factors in short RNAs by use of specifically placed carbon modifications (Figure 1D). We compare reactivity of C-5 methylated nucleotides, as such methylation is documented to enhance stacking,[21] to unmethylated analogs, and to test the effects of the absence or presence of electronegative atoms on nucleophilicity and cleavage, we study RNAs with C-nucleosides Ψ and m1Ψ, as well as cU, a carbacyclic uridine analog.[22] Nucleophilic reactivity is assessed with thermal and enzymatic bond cleavage as well as with acylation reactions.

Overall, our results show that methyl modifications that alter base stacking have mixed and moderate effects on nucleophilicity and strand cleavage. However, we find that the 2'-OH group of the canonical N-nucleoside uridine is significantly more reactive as a nucleophile than the analogous hydroxyl of C-nucleosides, resulting in considerably higher rates of acylation and of thermal strand cleavage. Enzymatic cleavage by ribonuclease A is also strongly suppressed at Ψ and m1Ψ as compared with U. Studies of the reactivity of small alcohols of varied pK_a suggest that a major contributor to this difference in nucleophilicity is inductive effects on 2'-OH acidity as a result of C versus N substitution. Taken together, our studies provide new mechanistic insight into the effects of native RNA modifications on their biological properties, and provide information that may aid in future improvements in the stability of therapeutic RNAs.

RESULTS AND DISCUSSION

Shielding of RNA thermal degradation by nucleotide modifications. While our primary studies were carried out with short synthetic oligoribonucleotides, we first confirmed that a pseudonucleotide replacement for uridine can stabilize mRNA to thermal degradation under our assay conditions (Figure 2A). A previous study reported this for an mRNA;[15] here we sought to confirm the effects of m1Ψ under our conditions using the enhanced green fluorescence protein (eGFP) mRNA context. Although this replacement occurs at less than 20% of the nucleotides of this RNA, our results confirmed stabilization of the strand, increasing the half-life of the full-length RNA by approximately 2-fold (Figure 2A). This is consistent with the previous measurements of mRNAs containing Ψ and m1Ψ,[16] confirming that they provide a degree of stabilization in a large RNA even as minority substitutions among canonical nucleotides.

For RNA polymers, all internucleotide linkages can simultaneously undergo cleavage, making quantitative measurements of rates challenging. To determine the effects of nucleotide modification on individual phosphodiester bond cleavage, we designed short (11 nt) RNA oligonucleotides (Table S1) containing only one scissile phosphodiester bond at the 3′ side of uridine and analogs (Figure 2B), with the adjacent nucleotides 2′-O-methylated to block cleavage at all other positions. Importantly, 2′-O-methylated RNAs retain a helical conformation nearly identical to that of unmethylated RNA oligonucleotides.[23] To investigate the effect of base stacking context for the central uridine analogues, we studied contexts of polyU and polyA sequence (Table S1). Adenine is the most strongly stacking base in RNA, while uracil is the least.[24] UV-monitored thermal melting studies in buffer (pH 7.0) confirmed that no significant higher-order intramolecular or intermolecular structure was formed by the RNA oligomers (Figure S1).

Figure 2. Shielding of RNA thermal degradation by base modifications. (A) mRNA stabilities assessed over time (water, 37 °C) for unmodified eGFP RNA and m1Ψ-modified eGFP RNA measured by capillary electrophoresis. (B) Molecular mechanism for thermal in-line cleavage of scissile phosphodiester bond in RNA oligonucleotides. Nucleotides with unstable and stable 3′-phosphodiester bonds are colored red and blue, respectively. (C, D) Plots of the cleavage of 11 nt RNAs after the central U or modified nucleotide by thermal cleavage at 37 °C in CHES buffer solution (pH 10.0) in U and A contexts, respectively. Data are averages from 3 replicates with error bars showing standard deviations. (E) Comparison of the rate constants (k_{clv}) derived from the thermal degradation shown in C and D. (F, G) Circular dichroism (CD) spectra for 11 nt RNAs with U and A contexts, respectively, with a central U or analog.

Kinetics of single bond cleavage events were assessed from base-catalyzed cleavage reactions of the oligonucleotides, monitored with gel electrophoresis. The mechanistic pathway of cleavage of RNA polymers involves in-line phosphoester transfer (Figure 2B).[8,25] The RNAs were incubated at 37 °C in a solution of 50 mM CHES (pH 10) containing 10 mM $MgCl_2$, similar to prior studies.[26] After quenching the reactions, the fraction of strand cleavage was quantified by denaturing polyacrylamide gel electrophoresis (PAGE) (Figure S2). Plots of intact RNA fraction *vs* time were fit with first-order exponential decay curves (Figures 2C,D and S3).

Strand cleavage rate constants (k_{clv}) are given in Table 1, and are compared graphically in Figure 2E. Data were obtained for uridine and four uridine analogs in two contexts. Comparing

Table 1. Rate constants for thermal cleavage at 3′ of uridine and analogues in oligoribonucleotides.[a]

U context	k_{clv} (min^{-1})	A context	k_{clv} (min^{-1})
<u>UUUUU</u> **U** <u>UUUUU</u>	3.7 (0.10) x 10^{-4}	<u>AAAAA</u> **U** <u>AAAAA</u>	4.9 (0.10) x 10^{-4}
<u>UUUUU</u> **m5U** <u>UUUUU</u>	0.99 (0.04) x 10^{-4}	<u>AAAAA</u> **m5U** <u>AAAAA</u>	1.3 (0.05) x 10^{-4}
<u>UUUUU</u> **Ψ** <u>UUUUU</u>	0.36 (0.02) x 10^{-4}	<u>AAAAA</u> **Ψ** <u>AAAAA</u>	0.15 (0.01) x 10^{-4}
<u>UUUUU</u> **m1Ψ** <u>UUUUU</u>	0.38 (0.02) x 10^{-4}	<u>AAAAA</u> **m1Ψ** <u>AAAAA</u>	0.26 (0.02) x 10^{-4}
<u>UUUUU</u> **cU** <u>UUUUU</u>	0.83 (0.03) x 10^{-4}	<u>AAAAA</u> **cU** <u>AAAAA</u>	0.85 (0.03) x 10^{-4}

[a]Rate constants derived from 20% PAGE analysis. Underlined nucleotides possess 2′-*O*-methyl groups to block cleavage. Conditions: CHES buffer (50 mM, pH 10.0) containing 10 mM Mg^{2+} at 37 °C. Data are averages of 3-4 replicates; error as standard deviations are given in parentheses.

the adjacent sequence contexts of polyU versus polyA, we find moderate and mixed effects; for the N-nucleotides U and m5U, cleavage was slower in the U context relative to the A context by 1.3-fold in rate constants. However, for the C-nucleotides Ψ and m1Ψ, the reverse was true, with slightly slower cleavage occurring in the A context. Previous studies have shown that sequence context can affect RNA cleavage rates significantly.[27]

Adjacent (nearest-neighbor) bases affect local RNA structure and stability by differential stacking free energies.[17] Methylation of uridine (in m5U) enhances base stacking and stabilizes helical conformations of RNA,[21] and *N*1-methylation of Ψ similarly stabilizes base stacking.[6,28] m5U and m1Ψ are both known in therapeutic RNAs, as is the structurally related methylation of cytosine in m5C. Here we find that, as was seen for the varied adjacent base context, methylation also has moderate and conflicting effects. C-5 methylation of uracil results in nearly identical 3.6-3.8-fold stabilization of the internucleotide linkage against cleavage in both sequence contexts. In contrast, analogous methylation of Ψ results in either no effect or a small enhancement of cleavage rate.

Importantly, the data also allow a comparison of the effects of C-nucleoside to N-nucleoside substitution on the cleavage rates of the short RNAs. Results were consistent across context and methylation status, and were considerably greater in magnitude; all C-nucleoside

substitutions result in strong stabilization against strand cleavage. Substituting Ψ for U stabilizes the RNA phosphodiester linkage against cleavage by substantial kinetic factors of 10-fold (U context) and 32-fold (A context). Similarly, substitution of m1Ψ for m5U slows strand cleavage by 2.7-fold to 5.3-fold. Thus, the C-nucleoside substitution results in strong suppression of in-line attack on the phosphodiester by the 2′-OH group. While a previous study showed stabilization by Ψ of partially-substituted RNAs from thermal degradation,[16] the current data provide quantitative measures of the effect on a specific bond.

Structural studies of the nucleosides indicate that the pseudouridine modification alters sugar conformation slightly; Ψ has a 48:52 North/South conformational preference, slightly shifted from that of uridine (53:47 North/South).[29] The Ψ modification also moderately enhances base stacking relative to U depending on sequence context.[17,30] CD spectra of the oligonucleotides (Figures 2F,G) showed evidence for moderate reduction of helicity in the U context by the base modifications. However, no changes in helicity were seen for the A context. Our data show that C-nucleoside substitution has a stronger suppressive effect on strand cleavage in the A context, where no change in overall helicity is seen, suggesting that global changes in oligomer conformation are unlikely to be the source of the protective effect. Thus,the data suggest a more localized explanation for most of the lower reactivity of 2′-OH groups in Ψ and m1Ψ.

Base modifications protect RNA from enzymatic degradation. To test whether enzymatic bond cleavage is also affected by uridine nucleotide modifications, we carried out quantitative measurements of cleavage with ribonuclease A, one of the most well-studied nuclease enzymes.[31] RNase A is a metal ion-independent ribonuclease,[32] and its catalytic activity involves two histidine residues for general base/acid catalysis (Figure 3A).[33] Although RNase A-catalyzed degradation follows a similar mechanistic pathway as in-line thermal cleavage, it provides ~10^{11}-fold rate acceleration.[34] RNase A prefers to cleave the phosphodiester bond to

Figure 3. Modifications protect RNA from ribonuclease cleavage. (A) Scheme of substrates employed in RNase A catalyzed cleavage of a specific internucleotide linkage in RNA oligonucleotides. (B) Plots showing decay of RNA phosphodiester by enzymatic cleavage at 37 °C with RNase A for oligonucleotides containing U, m5U, Ψ, and m1Ψ with time. (C) Linear plots of ln(remaining RNA fraction) vs time for unmodified and modified uridine-containing RNAs. Negative slopes indicate pseudo-first-order rate constants, from which second-order rate constants were obtained. (D) Comparison of apparent 2nd-order rate constants of RNase A-catalyzed degradation ($k_{clv(app)}$) for A-context oligonucleotides (see Table 1). Data were obtained from 3 replicates with error bars showing standard deviations.

the 3′ side of pyrimidine nucleotides and 5′ to purines,[31] enabling our 11 nt oligomers in the polyA context to act as appropriate enzyme substrates. We are aware of no prior studies testing the effects of C-nucleoside structure or methylation on RNase A nuclease activity.

We measured bond cleavage for the 11 nt sequences in the polyA context with comparison of U, m5U, Ψ, and m1Ψ. Due to very high enzymatic activity (Figure S4A), we used dilute RNase A solutions (10 ng mL⁻¹) with a large excess of RNA substrate to follow initial rates of cleavage at 37 °C in 50 mM Tris-HCl buffer (pH 7.4). The relative fractions of uncleaved oligonucleotide were quantified with denaturing PAGE (Figure S4) and plotted against time (Figure 3B). The K_M of RNase A has been reported as 1100 μM;[35] as our concentrations were

far below this, the data were fit well with pseudo-first-order kinetics (Figure 3B,C). Apparent second-order rate constants of RNase cleavage ($k_{clv(app)}$) were derived from the linear pseudo-first-order plots (Figure 3C), and data are plotted in Figure 3D. The results show that both base methylation and C1-substitution decrease ribonuclease cleavage rates markedly. Methylation of U slowed cleavage by 23-fold, and also hindered cleavage at Ψ (compared to m1Ψ) by a smaller factor of 2.1-fold. Most notably, C-nucleotide substitution had an even larger effect on cleavage, slowing strand scission by factors of 52-fold in the unmethylated case and 4.7-fold in the methylated case. Thus, the data show that both structural carbon substitutions have a strongly suppressing effect on a human ribonuclease-mediated RNA degradation.

Structures of RNase A with nucleotide substrates show close contacts of the enzyme with the pyrimidine base, explaining its preference for cleavage after a pyrimidine nucleotide in RNA.[36] It is possible that methylation sterically inhibits interactions with the preferred substrate, although more detailed kinetics studies will be needed to shed light on this possibility. Regardless of mechanism, the results suggest that naturally occurring C-5 methylation of RNA pyrimidines may have a similar suppressive effect on cleavage of the 3′-phosphodiester bond. Although 5-methylC is known in mRNAs, and has been shown to prolong cellular lifetimes of the RNA,[37] its effect on a specific nuclease enzyme has yet to be studied.

We find that Ψ strongly suppresses RNase A cleavage to the 3′ side of the nucleotide. Given the frequent occurrence of Ψ in natural RNAs, it is clear that one biological effect is likely to be suppression of enzymatic degradation rates at the bond 3′ to the nucleotide, although we add caution that this is one of multiple human ribonucleases. For RNase A, it seems difficult to attribute structural effects in the enzyme/RNA complex to explain this relatively large effect, since Ψ is no larger than U and adopts a very similar conformation. Thus, we hypothesize a

Figure 4. Shielding of RNA thermal degradation by carbacyclic uridine modifications. (A) Molecular mechanism for thermal in-line cleavage of scissile phosphodiester bond in carbacyclic uridine modified 11 nt RNA oligonucleotides. Nucleotide with cleavable 3′-phosphodiester bond is colored red. (B,C) Representative images of 20% denaturing PAGE analysis for thermal cleavage of cU RNAs in U and A contexts, respectively, after incubating substrate oligonucleotides in CHES buffer (pH 10.0) solution with Mg^{2+} at 37 °C. (D, E) Decay curves from band intensities; inset plots depict the first-order decay behavior, with slopes yielding rate constants k_{clv} (min^{-1}). Data were obtained from 3 replicates with error bars showing standard deviations.

local chemical effect of pseudouridine C-nucleoside substitution on reducing the inherent nucleophilicity of the attacking 2′-OH group (see below).

Carbon substitution in the ribose sugar also slows strand cleavage. The above data show that carbon substitution for electronegative N at the C1′ carbon of ribose can strongly reduce strand cleavage by thermal or enzymatic processes. We hypothesize that this replacement lowers nucleophilic reactivity of the nearby 2′-OH group by reducing the inductive effect of nearby electronegative atoms that stabilize the anionic form of the hydroxyl (Figures 1B,C). Although this was tested above with two C-nucleosides, we sought a further test of this hypothesis by replacing one of the other electronegative atoms in the vicinity of 2′-OH. To this end, we employed the recently described carbacyclic uridine analog cU (Figure 1),[22] which has a chemical structure identical to U but replaces the C4′ ring oxygen with carbon. Importantly, this substitution resides the same number of chemical bonds from the 2′-OH as the carbon

Figure 5. Effect of 2′-OH nucleophilicities on acylation. (A) Molecular mechanism and experimental workflow of RNA 2′-OH acylation with 10 different unmodified and modified oligonucleotides (Table S1). MALDI-TOF quantifies the rate constants of the acylation using the initial rates method. (B) Comparison of normalized rate constants for the acylation of modified nucleotides in the RNA oligonucleotides with corresponding cleavage rate constants. Normalization was done with respect to the value obtained for unmodified U in the U context. (C) Experimental workflow to study kinetics of acylation for alcohols with NAI at 37 °C or meNAI at 37 °C (pH 7.4). (D) Molecular structures of alcohols with pK_a range (12-15) at 25 °C to study the correlation of nucleophilicity with pK_a. (E) Plot of the logarithms of rate constants for primary alcohols (black circle) against reported pK_a, bracketing likely RNA pK_a values. Previously reported log(k_{Acyl}) and pK_a values for nucleotides are also shown (red triangles).

substitution in the pseudonucleosides (Figure 1), which is relevant because inductive effects depend strongly on the number of intervening chemical bonds.[38] We prepared

oligoribonucleotides having the same polyU and polyA contexts as the other modifications, and again measured thermal cleavage rates (Figure 4). Comparison of strand scission rates again revealed a clear shielding effect of the substitution on the strand cleavage with cU. Carbacyclic uridine decreased k_{clv} by 4.5-fold and 5.8-fold compared to unmodified uridine in the pyrimidine and purine contexts. This effect is similar in magnitude to the suppressive effect of C-for-N replacement in the pseudonucleotides (Figure 2E). CD spectra revealed that the RNAs containing cU exhibit moderately or slightly reduced global helicity as the analogous RNAs with U, respectively (Figure S5), suggesting that the stabilization in this case was not due to a large change in conformation, but is instead localized near the modification.

Carbon substitutions reduce nucleophilicity as measured by 2′-OH acylation. The above data show that carbon substitutions near the 2′-OH group of a ribonucleotide in RNA lower the nucleophilicity of this hydroxyl for the neighboring phosphate group. To evaluate nucleophilicity of this group in a different context and at neutral pH, we employed a different electrophile to test. The 2′-OH group of RNA can be modified by acylating agents in water, a fact that has been widely used in mapping RNA structure in SHAPE methods[39] as well as in strategies for high-yield labelling and stabilization of RNA.[40] Thus, we used an acylation reaction here to further evaluate the nucleophilicity of 2′-OH (Figure 5A). Notably, no prior study has to our knowledge quantified absolute acylation kinetics in full RNA strands, the results of which can be relevant both to RNA structure mapping and to development of new methods and reagents for RNA labelling.

As described above, these RNA sequences (Table S1) contain only one free 2′-OH available for acylation, simplifying the measurement of acylation yields and rates. Reactions were quantified from relative peak intensities measured by MALDI-TOF mass spectrometry over time (Figure 5A). Rate constants of acylation reaction (k_{acyl}) were determined with an

established RNA acylimidazole reagent (NAI), employed previously to study kinetics of acylation with mono- and di-nucleotides.[18] Data are shown in Table 2 and Figure S6.

Table 2. Rate constants of acylation reactions[a] at the central nucleotide of RNA oligonucleotides without or with modifications.

U oligonucleotide	k_{acyl} (M^{-1}min^{-1})	A oligonucleotide	k_{acyl} (M^{-1}min^{-1})
U	9.7 (0.2) x10^4	U	3.7 (0.9) x10^4
m5U	8.7 (0.2) x10^4	m5U	3.6 (0.9) x10^4
Ψ	2.7 (0.7) x10^4	Ψ	0.92 (0.20) x10^4
m1Ψ	2.7 (0.5) x10^4	m1Ψ	1.7 (0.4) x10^4
cU	1.5 (0.3) x10^4	cU	0.55 (0.16) x10^4

[a]Rate constants of RNA 2′-OH acylation using NAI in pH 7.4 buffer at 37 °C were determined from relative peak intensities measured by mass spectrometry. Standard deviation (SD) values are given in parentheses. See full sequences in Table 1.

Comparison of rates (Figure 5B) clearly distinguishes the effects of carbon substitution and base stacking. The k_{acyl} decreased by 3.6-fold comparing U to Ψ and 3.2-fold from m5U vs. m1Ψ modifications, whereas no comparable change was observed due to the methylation of U and Ψ in the polyU context. This suggests that there is a negligible effect of base stacking on acylation rates due to methylation, but there is significant reduction of k_{acyl} by C-1 carbon substitution. A similar and somewhat larger effect was seen for the cU modification, which decreased k_{acyl} by over 6-fold as a result of replacing electronegative O with C in the sugar ring. Interestingly, the A-context RNAs showed somewhat lower k_{acyl} values compared to the U context, reduced by 1.9 to 2.6-fold depending on modification. This is consistent with the prior observation of lower acylation reactivity of polyA as compared with polyU.[41]

Overall, although the correlation is not quantitative, each pairwise comparison of N/O substitution by carbon shows the same qualitative effect on acylation rate as it does in strand cleavage rate (Figure 5B). For example, in the polyU context, acylation is decreased for Ψ as compared to U, and strand cleavage rate is also decreased. The same is true for Ψ versus U in the polyA context, and indeed for the pairwise comparisons of U to cU and m5U to m1Ψ. In

all cases, substitution of nitrogen or oxygen by carbon decreases acylation rate and decreases strand scission rate as well.

Correlations of nucleophilicity and pK_a of alcohols. Taken together, the above data show clearly that carbon replacement of electronegative atoms near the 2′-OH group strongly reduces nucleophilicity as measured both by acylation (with an external electrophile) and strand self-cleavage (an internal electrophile). This is consistent with the notion that the replacement of the electronegative atoms reduces the acidity of the 2′-OH group via inductive effects. It is difficult to measure the pK_a of this group directly, as it is already high (*ca*. 12.5-13.5) in native RNA,[19] and titrations at high pH are likely to cause rapid strand cleavage. Thus we sought to connect nucleophilicity of the 2′-OH with its pK_a by studying a series of small alcohols with pK_a values between 12-15 (Figures 5C,D), spanning the range of reported pK_as of the 2′-OH of ribonucleotides.[19] The pK_a of ethanol is 15.9 at 25 °C, and substitution by electronegative atoms at the 2-carbon is well documented to inductively reduce the pK_a of the OH group.[42]

We measured acylation kinetics for the alcohols in pH 7.4 buffer at 37 °C, and initial reaction rates (Table 3) were determined by [1]H-NMR (Figures 5C and S7). The data confirmed an inverse relationship of pK_a with nucleophilicity as measured by acylation rate; a linear plot of log(k_{acyl}) against pK_a showed a slope of −0.91 (Figure S8), documenting an average 9-fold increase of kinetic rate per unit of pK_a for the alcohols. Thus, the data confirm that these alcohols react with NAI at neutral pH at rates governed by their pK_a values, and the less acidic they are, the lower their nucleophilicity as measured by acylation rate. The data are entirely consistent with a mechanism of reaction involving the anion form of the alcohol reacting with the electrophile, which we hypothesize is the case for RNA as well.

Table 3. Rate constants of acylation reaction for alcohols[a]

Alcohol	k_{acyl} (NAI) (M^{-1}min^{-1})	k_{acyl} (meNAI) (M^{-1}min^{-1})
2,2,2-Trifluoroethanol	17.0 ± 6.1	15.0 ± 2.5
2,2,2-Trichloroethanol	7.9 ± 1.5	2.4 ± 1.8
2,2-Dichloroethanol	3.6 ± 1.2	1.2 ± 0.3
2-Chloroethanol	0.13 ± 0.06	0.049 ± 0.015

[a]Rate constants of acylation in presence of NAI (37 °C) and meNAI (25 °C) in buffer (pH 7.4) solution were derived from ^{1}H NMR analysis of initial rates.

Previously determined pK_a[19] and corresponding k_{acyl} values[18] for mononucleotides were fit well in our plot of the smaller alcohols (log(k_{acyl}) *vs* pK_a, Figures S9, S10, and 5E) using a closely analogous reagent (meNAI) and temperature (25 °C), suggesting that the reactivities of ribonucleotides toward NAI are also well explained by this mechanism. If we assume that pK_a shifts due to carbon substitution for electronegative atoms were entirely responsible for the relative acylation rates seen for the oligonucleotides (Figures 5E and S8), then the 3.6-fold reduction of 2′-OH acylation rate from U to Ψ modification suggests that the carbon substitution causes a *ca.* 0.6 pK_a unit increase for the 2′-OH of Ψ relative to U. The larger 6-fold reduction in acylation rates caused by cU substitution relative to U suggests a somewhat greater increase in pK_a of approximately 0.9, consistent with the greater electronegativity of oxygen relative to nitrogen.

After this work was completed, a preprint describing relevant studies of dinucleotides containing Ψ and m1Ψ was posted.[43] Comparing U to Ψ, the authors observed a 2-fold decrease in thermal cleavage rate and a 10-fold decrease in enzymatic cleavage rate by RNase A, consistent with the current findings albeit considerably smaller in magnitude, possibly due to the small size of the substrates in that study. Molecular dynamics simulations suggested a 2.9 kcal/mol free energy difference for proton transfer from 2′-OH for U vs. Ψ, attributed to pK_a difference. That finding is also generally consistent with the current results, although a 2.9 kcal/mol difference suggests a pK_a difference of *ca.* 2.1 pK units, which is significantly larger

than our experimental findings and greater than the differences in documented pK_as of relevant small alcohols such as dichloroethanol and trichloroethanol. Unlike the current study, the authors observed no methylation-associated difference in enzymatic cleavage rates for m1Ψ versus Ψ, whereas we observe substantial slowing as a result of methylation in two nucleotide contexts.

It should be noted that our data show that acylation rates in the RNA oligomers are far higher (by a factor of *ca*. 10^4) than for the current small alcohols as well as for mononucleotides measured previously; however, comparisons of trends within the context of the polymer (such as U versus Ψ) remain consistent. For example, trichloroethanol (pK_a 12.8), having three electronegative atoms, is more reactive to acylation than dichloroethanol (pK_a 13.5), and the 2′-OH group of U in RNA (also having three electronegative atoms) is more reactive than that of Ψ (with two electronegative atoms) by a similar factor. Work is ongoing to understand why the 2′-OH group in RNA is much more reactive than small-molecule alcohols of similar pK_a.

CONCLUSIONS

Taken as a whole, our data show that carbon substitution in ribonucleotides, particularly for the atoms attached at C1′ of ribose, results in strongly reduced nucleophilicity of the 2′-OH group in RNAs, causing slower rates of attack on external electrophiles such as acylating agents, and reduced rates of attack on the neighboring phosphodiester group as well. This lower nucleophilicity in carbon-substituted analogs results in markedly slower RNA strand cleavage rates at the phosphodiester bond 3′ to the nucleotide where substitution occurs. This has relevance both to native RNAs and to therapeutic RNAs containing Ψ and m1Ψ modifications, providing quantitative data for this protective effect as well as basic understanding of its physicochemical origins. The biological roles of Ψ in cellular RNAs are complex[44] and are under active study;[45–47] however, the current data provide clear and quantitative documentation of the protective effect of this modification against both thermal and enzymatic cleavage. The

data also suggest that broader carbon substitution for O4′, N1 and other atoms may be worthy of consideration in the design of future modification of RNAs for therapeutic use.

We have further shown that alcohols with similar pK_a as ribonucleotides have similar nucleophilic reactivity, and that alcohols with reduced number of electronegative atoms have reduced reactivity. In RNA, we show that this also the case with Ψ and m1Ψ relative to their N-nucleoside counterparts U and m5U. Thus, although direct pK_a measurements of 2′-OH in RNA are not yet available, the data suggest that the C-nucleosides Ψ and m1Ψ stabilize RNA by reducing the acidity of their associated 2′-OH groups as compared with canonical uridine, due to reduced inductive stabilization of the anionic form of the 2′-OH group.

In addition to effects on 2′-OH acidity and nucleophilicity, our data also show that methyl substitution of bases at C-5 of U and analogously at N1 of Ψ can also have protective effects on stabilizing phosphodiester bonds in RNA, depending on the context. Both of these substitutions enhance base stacking, which seems likely to be relevant to explaining their protective effect, as the substitution is too remote from the 2′-OH to cause a direct effect on reactivity. We observe that the effect for thermal cleavage is mixed (depending on context, as is the case for stacking generally[17]) and requires more study before its origins and scope can be fully understood. However, the effect has not been reported previously and is worthy of consideration in the biological context. The analogous C-5 methylation of cytosine in RNA is known[37] and the current results suggest that, like m5U, it has the potential to affect the stability of adjacent phosphodiester bonds. Interestingly, the adenine modification m6A, also widely observed in cellular RNA, is also known to enhance base stacking,[48] and recent experiments have shown that it does indeed affect the reactivity of adjacent nucleotides substantially, enabling the use of acylating agents to detect positions of m6A in cellular RNA.[49] Thus, the current data suggest that these other methyl substitutions in native RNAs bear further study in light of possible alterations of nearby 2′-OH reactivity.

ACKNOWLEDGEMENTS

We thank the U.S. National Institutes of Health (GM145357) for support.

REFERENCES

(1)   Corey, D. R.; Damha, M. J.; Manoharan, M. Challenges and Opportunities for Nucleic Acid Therapeutics. *Nucleic Acid Ther.* **2022**, *32* (1), 8–13.

(2)   Ho, L. L. Y.; Schiess, G. H. A.; Miranda, P.; Weber, G.; Astakhova, K. Pseudouridine and N1-Methylpseudouridine as Potent Nucleotide Analogues for RNA Therapy and Vaccine Development. *RSC Chem. Biol.* **2024**, *5* (5), 418–425.

(3)   Wang, S.; Weissman, D.; Dong, Y. RNA Chemistry and Therapeutics. *Nat. Rev. Drug Discov.* **2025**. https://doi.org/10.1038/s41573-025-01237-x.

(4)   Roundtree, I. A.; Evans, M. E.; Pan, T.; He, C. Dynamic RNA Modifications in Gene Expression Regulation. *Cell* **2017**, *169* (7), 1187–1200.

(5)   Shi, H.; Chai, P.; Jia, R.; Fan, X. Novel Insight into the Regulatory Roles of Diverse RNA Modifications: Re-Defining the Bridge between Transcription and Translation. *Mol. Cancer* **2020**, *19* (1), 1–17.

(6)   Monroe, J.; Eyler, D. E.; Mitchell, L.; Deb, I.; Bojanowski, A.; Srinivas, P.; Dunham, C. M.; Roy, B.; Frank, A. T.; Koutmou, K. S. N1-Methylpseudouridine and Pseudouridine Modifications Modulate MRNA Decoding during Translation. *Nat. Commun. 2024 151* **2024**, *15* (1), 1–11.

(7)   Lin, S.; Kuang, M. RNA Modification-Mediated MRNA Translation Regulation in Liver Cancer: Mechanisms and Clinical Perspectives. *Nat. Rev. Gastroenterol. Hepatol.* **2024**, *21* (4), 267–281.

(8)   Oivanen, M.; Kuusela, S.; Lönnberg, H. Kinetics and Mechanisms for the Cleavage and Isomerization of the Phosphodiester Bonds of RNA by Brønsted Acids and Bases. *Chem. Rev.* **1998**, *98* (3), 961–990.

(9)   Greis, K.; Kirschbaum, C.; Taccone, M. I.; Götze, M.; Gewinner, S.; Schöllkopf, W.; Meijer, G.; von Helden, G.; Pagel, K. Studying the Key Intermediate of RNA Autohydrolysis by Cryogenic Gas-Phase Infrared Spectroscopy. *Angew. Chemie - Int. Ed.* **2022**, *61* (19), 1–7.

(10)   Chheda, U.; Pradeepan, S.; Esposito, E.; Strezsak, S.; Fernandez-Delgado, O.; Kranz, J. Factors Affecting Stability of RNA – Temperature, Length, Concentration, PH, and Buffering Species. *J. Pharm. Sci.* **2024**, *113* (2), 377–385.

(11)   Usher, D. A.; Richardson, D. I. J.; Eckstein, F. Absolute Stereochemistry of the Second Step of Ribonuclease Action. *Nature* **1970**, *228* (5272), 663–665.

(12)   Roberts, G. C. K.; Dennis, E. A.; Meadows, D. H.; Cohen, J. S.; Jardetzky, O. The Mechanism of Action of Ribonuclease. *Proc. Natl. Acad. Sci.* **1969**, *62* (4), 1151–1158.

(13)   Manoharan, M.; Akinc, A.; Pandey, R. K.; Qin, J.; Hadwiger, P.; John, M.; Mills, K.; Charisse, K.; Maier, M. A.; Nechev, L.; Greene, E. M.; Pallan, P. S.; Rozners, E.; Rajeev, K. G.; Egli, M. Unique Gene-Silencing and Structural Properties of 2′-Fluoro-Modified SiRNAs. *Angew. Chemie Int. Ed.* **2011**, *50* (10), 2284–2288.

(14)   Fang, L.; Xiao, L.; Jun, Y. W.; Onishi, Y.; Kool, E. T. Reversible 2′-OH Acylation Enhances RNA Stability. *Nat. Chem.* **2023**, *15* (9), 1296–1305.

(15)   Karikó, K.; Muramatsu, H.; Welsh, F. A.; Ludwig, J.; Kato, H.; Akira, S.; Weissman, D. Incorporation of Pseudouridine into MRNA Yields Superior Nonimmunogenic Vector with Increased Translational Capacity and Biological Stability. *Mol. Ther.* **2008**, *16* (11), 1833–1840.

(16)   Leppek, K.; Byeon, G. W.; Kladwang, W.; Wayment-Steele, H. K.; Kerr, C. H.; Xu, A. F.; Kim, D. S.; Topkar, V. V.; Choe, C.; Rothschild, D.; Tiu, G. C.; Wellington-Oguri, R.; Fujii, K.; Sharma, E.; Watkins, A. M.; Nicol, J. J.; Romano, J.; Tunguz, B.; Diaz, F.; Cai, H.; Guo, P.; Wu, J.; Meng, F.; Shi, S.; Participants, E.; Dormitzer, P. R.; Solórzano, A.; Barna, M.; Das, R. Combinatorial Optimization of MRNA Structure, Stability, and Translation for RNA-Based Therapeutics. *Nat. Commun.* **2022**, *13* (1),1536.

(17)   Kierzek, E.; Malgowska, M.; Lisowiec, J.; Turner, D. H.; Gdaniec, Z.; Kierzek, R. The Contribution of Pseudouridine to Stabilities and Structure of RNAs. *Nucleic Acids Res.* **2014**, *42* (5), 3492–3501.

(18)   Jash, B.; Kool, E. T. Conjugation of RNA via 2′-OH Acylation: Mechanisms Determining Nucleotide Reactivity. *Chem. Commun.* **2022**, *58* (22), 3693–3696.

(19)   Velikyan, I.; Acharya, S.; Trifonova, A.; Földesi, A.; Chattopadhyaya, J. The p K a 's of 2'-Hydroxyl Group in Nucleosides and Nucleotides. *J. Am. Chem. Soc.* **2001**, *123* (12), 2893–2894.

(20)   Velema, W. A.; Kool, E. T. The Chemistry and Applications of RNA 2′-OH Acylation. *Nat. Rev. Chem.* **2020**, *4* (1), 22–37.

(21)   Wang, S.; Kool, E. T. Origins of the Large Differences in Stability of DNA and RNA Helixes: C-5 Methyl and 2′-Hydroxyl Effects. *Biochemistry* **1995**, *34* (12), 4125–4132.

(22)   Akabane-Nakata, M.; Chickering, T.; Harp, J. M.; Schlegel, M. K.; Matsuda, S.; Egli, M.; Manoharan, M. RNAs Containing Carbocyclic Ribonucleotides. *Org. Lett.* **2022**, *24* (2), 525–530.

(23)   Popenda, M.; Biala, E.; Milecki, J.; Adamiak, R. W. Solution Structure of RNA Duplexes Containing Alternating CG Base Pairs: NMR Study of r(CGCGCG)2 and 2'-O-Me(CGCGCG)2 under Low Salt Conditions. *Nucleic Acids Res.* **1997**, *25* (22), 4589–4598.

(24)   Sugimoto, N.; Turner, D. H.; Kierzek, R. Sequence Dependence for the Energetics of Dangling Ends and Terminal Base Pairs in Ribonucleic Acid. *Biochemistry* **1987**, *26* (14), 4554–4558.

(25)   Pia Járvinen, Mikko Oivanen, and H. L. Interconversion and Phosphoester Hydrolysis of 2',5'- and 3',5'-Di Núcleo Side Monophosphates: Kinetics and Mechanisms. *J. Org. Chem.* **1991**, *56*, 5396–5401.

(26)   Li, Y.; Breaker, R. R. Kinetics of RNA Degradation by Specific Base Catalysis of Transesterification Involving the 2γ-Hydroxyl Group. *J. Am. Chem. Soc.* **1999**, *121* (23), 5364–5372.

(27)   Kaukinen, U.; Lyytikäinen, S.; Mikkola, S.; Lönnberg, H. The Reactivity of Phosphodiester Bonds within Linear Single-Stranded Oligoribonucleotides in Strongly Dependent on the Base Sequence. *Nucleic Acids Res.* **2002**, *30* (2), 468–474.

(28)   Dutta, N.; Deb, I.; Sarzynska, J.; Lahiri, A. Structural and Thermodynamic Consequences of Base Pairs Containing Pseudouridine and N1-methylpseudouridine in RNA Duplexes. *ChemistrySelect* **2025**, *10* (4). https://doi.org/10.1002/slct.202400006.

(29)   Chang, Y. C.; Herath, J.; Wang, T. H. H.; Chow, C. S. Synthesis and Solution Conformation Studies of 3-Substituted Uridine and Pseudouridine Derivatives. *Bioorganic Med. Chem.* **2008**, *16* (5), 2676–2686.

(30)   Davis, D. R. Stabilization of RNA Stacking by Pseudouridine. *Nucleic Acids Res.* **1995**, *23* (24), 5020–5026.

(31)   Raines, R. T. Ribonuclease A. *Chem. Rev.* **1998**, *98* (3), 1045–1065.

(32)   Cuchillo, C. M.; Nogués, M. V.; Raines, R. T. Bovine Pancreatic Ribonuclease: Fifty Years of the First Enzymatic Reaction Mechanism. *Biochemistry* **2011**, *50* (37), 7835–7841.

(33)   Messmore, J. M.; Fuchs, D. N.; Raines, R. T. Ribonuclease A: Revealing Structure-Function Relationships with Semisynthesis. *J. Am. Chem. Soc.* **1995**, *117* (31), 8057–8060.

(34)   Thompson, J. E.; Kutateladze, T. G.; Schuster, M. C.; Venegas, F. D.; Messmore, J. M.; Raines, R. T. Limits to Catalysis by Ribonuclease A. *Bioorg. Chem.* **1995**, *23* (4), 471–481.

(35)   Moiseyev, G. P.; Yakovlev, G. I.; Lysov, Y. P.; Chernyi, A. A.; Polyakov, K. M.; Oivanen, M.; Lönnberg, H.; Beigelman, L. N.; Efimtseva, E. V.; Mikhailov, S. N. Determination of the Nucleotide Conformation in the Productive Enzyme-Substrate Complexes of RNA-Depolymerases. *FEBS Lett.* **1997**, *404* (2–3), 169–172.

(36)   Kelemen, B. R.; Schultz, L. W.; Sweeney, R. Y.; Raines, R. T. Excavating an Active Site: The Nucleobase Specificity of Ribonuclease A. *Biochemistry* **2000**, *39* (47), 14487–14494.

(37)   Yang, Y.; Wang, L.; Han, X.; Yang, W.-L.; Zhang, M.; Ma, H.-L.; Sun, B.-F.; Li, A.; Xia, J.; Chen, J.; Heng, J.; Wu, B.; Chen, Y.-S.; Xu, J.-W.; Yang, X.; Yao, H.; Sun, J.; Lyu, C.; Wang, H.-L.; Huang, Y.; Sun, Y.-P.; Zhao, Y.-L.; Meng, A.; Ma, J.; Liu, F.; Yang, Y.-G. RNA 5-Methylcytosine Facilitates the Maternal-to-Zygotic Transition by Preventing Maternal MRNA Decay. *Mol. Cell* **2019**, *75* (6), 1188-1202.e11.

(38)   Bianchi, G.; Howarth, O. W.; Samuel, C. J.; Vlahov, G. Long-Range σ-Inductive Interactions through Saturated C–C Bonds in Polymethylene Chains. *J. Chem. Soc., Perkin Trans. 2* **1995**, No. 7, 1427–1432.

(39)   Wilkinson, K. A.; Merino, E. J.; Weeks, K. M. Selective 2′-Hydroxyl Acylation Analyzed by Primer Extension (SHAPE): Quantitative RNA Structure Analysis at Single Nucleotide Resolution. *Nat. Protoc.* **2006**, *1* (3), 1610–1616.

(40)   Shioi, R.; Kool, E. T. Chemical Diversity of Reagents That Modify RNA 2′-OH in Water: A Review. *Chem. Sci.* **2024**, *15* (39), 15968–15982.

(41)   Xiao, L.; Fang, L.; Kool, E. T. Acylation Probing of "Generic" RNA Libraries Reveals Critical Influence of Loop Constraints on Reactivity. *Cell Chem. Biol.* **2022**, *29* (8), 1341-1352.e8.

(42)   Ballinger, P.; Long, F. A. Acid Ionization Constants of Alcohols. II. Acidities of Some Substituted Methanols and Related Compounds. *J. Am. Chem. Soc.* **1960**, *82* (4), 795–798.

(43)   Gutierrez, C. S.; Silkenath, B.; Kojasoy, V.; Pich, J. A.; Lim, D. C.; Raines, R. T.

Pseudouridine Residues as Substrates for Serum Ribonucleases. *bioRxiv* **2025**, 1–22. https://doi.org/10.1101/2025.06.18.660232.

(44)   Borchardt, E. K.; Martinez, N. M.; Gilbert, W. V. Regulation and Function of RNA Pseudouridylation in Human Cells. *Annu. Rev. Genet.* **2020**, *54*, 309–336.

(45)   Bérouti, M.; Wagner, M.; Greulich, W.; Piseddu, I.; Gärtig, J.; Hansbauer, L.; Müller-Hermes, C.; Heiss, M.; Pichler, A.; Tölke, A. J.; Witte, G.; Hopfner, K.-P.; Anz, D.; Sattler, M.; Carell, T.; Hornung, V. Pseudouridine RNA Avoids Immune Detection through Impaired Endolysosomal Processing and TLR Engagement. *Cell* **2025**, 1–16.

(46)   Liu, J.; Yan, X.; Wu, H.; Ji, Z.; Shan, Y.; Wang, X.; Ran, Y.; Ma, Y.; Li, C.; Zhu, Y.; Gu, R.; Wen, H.; Yi, C.; Chen, P. R. RNA Codon Expansion via Programmable Pseudouridine Editing and Decoding. *Nature* **2025**, No. April 2024. https://doi.org/10.1038/s41586-025-09165-x.

(47)   Luo, N.; Huang, Q.; Zhang, M.; Yi, C. Functions and Therapeutic Applications of Pseudouridylation. *Nat. Rev. Mol. Cell Biol.* **2025**. https://doi.org/10.1038/s41580-025-00852-1.

(48)   Liu, B.; Merriman, D. K.; Choi, S. H.; Schumacher, M. A.; Plangger, R.; Kreutz, C.; Horner, S. M.; Meyer, K. D.; Al-Hashimi, H. M. A Potentially Abundant Junctional RNA Motif Stabilized by M6A and Mg2+. *Nat. Commun.* **2018**, *9* (1), 1–10.

(49)   Xiao, L.; Fang, L.; Zhong, W.; Kool, E. T. RNA infrastructure profiling illuminates transcriptome structure in crowded spaces. *Cell Chem. Biol.* **2024**, *31*, 2156-2167.