Title

Kinematics and morphological correlates of descent strategies in arboreal mammals suggest early upright postures in euprimates

Keywords

Locomotion, kinematics, gait, body mass, limb proportions, head size

Authors

Séverine L.D. Toussaint[1,2]*, Dionisios Youlatos[3], John A. Nyakatura[2]

*Corresponding author

Affiliations

[1] Center for Research on Paleontology-Paris, UMR 7207, CNRS/MNHN/SU, 8 rue Buffon CP38, 75005 Paris, France.

[2] Laboratory of Comparative Zoology, Institute of Biology, Humboldt University of Berlin, Philippstrasse 13, Haus 2, 10115 Berlin, Germany.

[3] Department of Zoology, School of Biology, Aristotle University of Thessaloniki, Panepistimioupoli, 54124 Thessaloniki, Greece.

Abstract

Ascending and descending sloping and vertical branches is critical for arboreal locomotion and likely played a major role in early primate evolution. While most studies have focused on ascent, descending behaviors also provide insight into the functional significance of arboreal adaptations. To test how descending vertical supports of varying diameters affects locomotor abilities, we quantified postural and kinematic features during descents and ascents on vertical supports in 21 eutherian and metatherian mammals, and examined their relation to morphology. Primates showed greater variability in descent behaviors, using tail-first and side postures more often than other mammals, which predominantly descended head-first. Overall, animals adopted several kinematic adjustments to enhance stability during descent compared to ascent, including slower speeds, higher duty factors, and greater use of asymmetrical gaits. Additionally, vertical descent strategies reflected trade-offs among body mass, limb proportions, and head mass. Using a morphology-based model, we then inferred possible descent behaviors in 13 extinct euarchontoglires. Our results suggest that ancestral adaptations

for vertical locomotion may have promoted frequent upright (head-up) postures in early primates.

Introduction

Arboreal environments impose strong physical constraints on vertebrate morphology and locomotor adaptations, due to the diversity of supports varying in orientation, diameter, and compliance. For arboreal mammals, the ability to navigate sloping and vertical trunks, branches, and lianas is essential, as these constitute the most available supports in trees. Climbing on vertical supports is a key aspect of arboreal locomotion in both fully arboreal and scansorial species, enabling access to the canopy from the ground, diverse food sources, enhanced vigilance, predator avoidance, and secure resting and nesting sites [1]. Climbing abilities in mammals date back to the Jurassic [2–4] and have played a central role in the evolution of euarchontoglires, especially primates [5–13]. Adaptations for climbing vertical supports have been identified in early Paleogene primates through postcranial fossil analyses [14–17]. Frequent vertical climbing may have been a prerequisite for the development of hallucal and pollical grasping in early primates [18,19], and has been proposed as an alternative scenario to the hypothesis of cautious displacements on narrow branches of angiosperm tree peripheries [20–22].

Several extant mammal species have been proposed as models for early primate arboreal adaptations, including small strepsirrhines [23,24], platyrrhines [13,19,25], treeshrews [26], rodents [27–29], carnivorans [30], and marsupials [19,31–35]. While not all arboreal mammals travel on narrow terminal branches, all rely on vertical supports to navigate canopy strata. However, few comparative studies have examined vertical climbing in non-primate arboreal mammals, and most have focused on ascents [11,36–38]. Although behaviors such as dropping, jumping, or gliding allow arboreal animals to move to lower tree layers, the ability to safely descend sloping and vertical supports remains critical, yet largely understudied [39].

Non-primate arboreal mammals often adopt a head-first posture when descending vertical supports, as reported in some rodents [12,40,41] and procyonid carnivorans [30,42]. This strategy relies on reversed feet and functional claws that enable secure gripping of tree bark during descent. Head-first descent is also common in prehensile-tailed mammals, such as certain platyrrhine primates [43–45] and the kinkajou [30]. In species lacking a prehensile tail, and in primates that lack claws and use their digits to wrap around supports, alternative descent strategies are required. In a study of nine small to medium-sized strepsirrhines, Perchalski [46] found that lemurs under 1 kg predominantly used head-first descents regardless of support properties, whereas larger species (>1 kg) increasingly adopted tail-first postures on supports angled at 45° or more,

depending on species. Both head-first and tail-first strategies involve orthograde postures with

the body aligned parallel to the support [7]. Perchalski also recorded other descent behaviors,

including leaping, dropping, and asymmetrical descents that did not fit into strict head-first or

tail-first categories. These findings suggest that in strepsirrhines, body mass and locomotor

adaptations influence the choice of descent posture. However, such behavioral variation also

highlights the need for broader comparative data across arboreal mammals to fully understand

the functional and ecological significance of vertical descent strategies in an evolutionary

context.

From a locomotor perspective, speed and duty factor (DF, the percentage of stride duration

a limb remains in contact with the support) are key indicators of arboreal locomotor

performance [47]. Studies across tetrapods have shown that animals generally move more slowly

on narrow supports than on wider ones [48,49]. Small primates [24,50,51], rodents [12,52,53], and

marsupials [34,54] typically decrease speed and increase duty factor, especially for the forelimbs,

during descents compared to ascents on inclined supports and compared to locomotion on

horizontal supports. These mechanisms supposedly help enhance stability by allowing more

cautious displacements [39,48,53].

Gait patterns also reflect arboreal adaptations, particularly in primates [36,47,55,56]. Vertical

climbing employs both symmetrical (walk, trot, run) and asymmetrical (bound, half-bound,

gallop) gaits, similar to above-branch locomotion. Symmetrical gaits are defined as locomotor

patterns in which the footfalls of a girdle (a pair of fore- or hindlimbs) are evenly spaced in

time, with the right and left limbs of a pair of limbs being approximately 50% out of phase with

each other [56,57]. Symmetrical gaits can be further divided into two types: diagonal-sequence

gaits, in which a hindlimb footfall is followed by that of the contralateral forelimb, and lateral-

sequence gaits, in which a hindlimb footfall is followed by that of the ipsilateral forelimb [56,58,59].

In contrast, asymmetrical gaits are characterized by unevenly spaced footfalls within a girdle,

with the right and left limbs moving in near synchrony [60]. Asymmetrical gaits, often used by

small mammals at high speeds [48,61], allow to increase the distance traveled, but often include an

aerial phase [60], raising support reaction forces and potentially causing oscillations or breakage

on narrow horizontal or sloping branches, complicating locomotor control and safety. Benefits

of asymmetrical gaits have thus been proposed to be related to both body mass and support

diameter [62]. On vertical supports, such as trunks or lianas, these risks are reduced due to

different force distributions and gravity orientation. Symmetrical gaits are widespread in

primates and have been linked to the use of the fine branch niche, as enhanced grasping and

hindlimb dominance supposedly increase stability during cautious locomotion on narrow

horizontal or sloping supports [56,59,63,64]. Notably, primates adjust their symmetrical gaits with support inclination, favoring diagonal-sequence gaits during ascents and lateral-sequence gaits during descents [51,65–69]. Symmetrical gaits are also common in rodents [12,52], scandentians [70], carnivorans [30], and marsupials [71–74]. However, most kinematic studies have focused on horizontal or moderately sloped supports (~45°), rarely addressing vertical (~90°) supports or analyzing both symmetrical and asymmetrical gaits comprehensively. As a result, comparative data on how vertical supports of varying diameters affect descent and ascent kinematics remain limited, hindering our understanding of this common arboreal behavior.

In addition to body mass, limb proportions influence locomotor and postural strategies [46,65]. The relative length of autopodials, particularly manual and pedal digits, is closely linked to grasping ability and thus affects locomotor performance in primates [13,19,21,25,75] and small marsupials [19,32]. Species with relatively short autopods may struggle to maintain a secure grip during vertical descent. Fore- and hindlimb length ratios have also been proposed as predictors of postural adaptation in vertical locomotion, with shorter forelimbs generating greater hindlimb traction during head-first descents [36,46]. Consequently, a lower intermembral index (forelimb-to-hindlimb length ratio) increases the risk of forward pitching when descending vertically head-first compared to tail-first [46].

Variation of weight distribution along the body, notably caused by head mass differences, may also influence postural strategies during vertical descent (i.e., head-first vs. tail-first). Primates exhibit the highest brain-to-body mass ratio among mammals [76]. This relative brain expansion, traced back to early primate evolution, is a key synapomorphy alongside a divergent hallux and pollex, nails replacing claws, elongated hindlimbs and autopods, and large convergent eyes [21,77,78]. However, the evolutionary sequence and interplay between locomotor and cognitive specializations remain unresolved, despite consensus on the small body size [13,79] and arboreality [80–82] of primate ancestors. While brain enlargement likely increases relative head mass, its effects on arboreal locomotion have yet to be investigated in primates or other mammals. Given that head mass influences both the center of mass and moment of inertia, greater head mass may increase toppling risk during vertical head-first descent.

In this study, we aim to test (a) whether postural and kinematic adjustments during vertical descents versus ascents vary consistently across mammalian groups; (b) whether these strategies correlate with specific morphological parameters; and (c) whether such correlations can be used to infer descent postures in fossils, shedding light on the evolutionary history of vertical arboreal locomotion, particularly in primates. To address these questions, we analyzed locomotor sequences of descents and ascents on vertical supports in 21 small- to medium-sized

arboreal mammal species, including taxa proposed as modern analogues of early primates. We first examined the effect of support diameter (small, medium, large) on postural strategies and kinematics (speed, duty factor, gait) during descents relative to ascents. We then assessed the relationships between descent strategies and morphological parameters (body mass, limb proportions, and relative head mass) for each species. These data allowed us to build a model incorporating seven significant morphological predictors, which we applied to infer vertical descent behavior in 13 extinct euarchontoglires, including adapiforms (early strepsirrhines), omomyiforms (early haplorhines), a sciuromorph (early rodent), and plesiadapiforms (stem primates).

We predict that:

H1) Postural strategies: species under 1 kg and/or possessing a prehensile tail will exhibit head-first descent postures more frequently than larger species, which will favor tail-first postures [46].

H2) Kinematic adjustments: animals will adopt more cautious locomotor patterns during vertical descents compared to ascents, characterized by reduced speed, increased duty factors, and a higher proportion of symmetrical gaits, particularly on narrow supports [48,52,53,65,70].

H3) Morphological correlates: the intermembral index will positively correlate with the proportion of head-first descents, whereas relative autopod length and head size will correlate negatively with head-first descent frequency [21,36,46].

H4) Evolutionary implications: early euarchontoglires, due to their small size and generalized locomotor abilities, likely employed a high proportion of head-first descents [12,53], whereas early euprimates may have increasingly favored tail-first descents associated with larger body and brain, and elongated hindlimbs and autopods [21,77,78].

Results

Descent strategies on vertical supports among arboreal mammals

We recorded the locomotion of 57 adult individuals from 21 species: 6 strepsirrhine primates, 5 platyrrhine primates, 1 scandentian, 3 rodents, 3 carnivorans, and 3 marsupials (Fig. 1, Table S1). We categorized supports in three diameters: small (i.e., twigs), medium (i.e., branches), and large (i.e., boughs and very large trunks) relative to the size of the animals' hands and feet (see section Materials and Methods for more details). Whenever possible, we

analyzed up to 10 ascents and 10 descents per individual on each support type. In total, we

analyzed 1,400 ascents and 1,390 descents.

We identified three distinct strategies of descent: head-first descent, side descent, and tail-

first descent (Fig. 1A) which partly overlap with Perchalski's ethogram [46]. During a head-first

descent the body is positioned head-down, parallel to the support, hands and feet are anchored

to the support, and the face is directed downward (see *Trichosurus vulpecula* in Fig. 1A).

During a side descent the body is rotated perpendicularly to the support and the face is also

directed downward (see *Plecturocebus cupreus* in Fig. 1A). During a tail-first descent the body

is positioned head-up and parallel to the support similarly to ascents, and the face is either

directed horizontally or downward (see *Eulemur mongoz* in Fig. 1A). We found a clear

difference between primates and non-primates in the occurrence of each descent strategy on

each support type (Fig. 1B). Most non-primate species, although very diverse in body mass and

morphology, showed only head-first descents independently of the orientation and diameter of

the supports. The racoons, the largest species of this study (~ 6 kg), were an exception and

mainly exhibited tail-first descents. The other studied carnivorans, i.e., the coatis and the

kinkajous (the latter being the only prehensile-tailed of this study along with the brushtail

possum *Trichosurus vulpecula*), as well as all marsupials, rodents, and treeshrews exhibited

only head-first descents independently of the support diameter (Fig. 1B, D). In sharp contrast,

primates rarely used head-first descents on vertical supports except for lorises

(*Xanthonycticebus pygmaeus,* strepsirrhine) and tamarins (*Saguinus imperator* and *S. oedipus*,

platyrrhines), the smallest primate species of this study (~ 360 g to ~ 470 g respectively). The

other strepsirrhines (i.e., lemurs) descended vertical supports mainly using tail-first descents

with higher proportions on small supports, and the other platyrrhines generally used side

descents with also increased proportions on small supports. The difference between

phylogenetic groups is substantiated by a significant phylogenetic signal contained in the

overall proportion of head-first descent strategy on vertical supports (k=0.58, $p = 6.6.10^{-4}$, effect

size= 2.57).

Figure 1. Strategies of descent on vertical supports of various diameters by species. A. Photographs illustrating the three different strategies of descent identified in this study. Black arrows represent the axis of the body regarding the support and the direction of the movement, and grey arrows represent the direction of the gaze. **B.** Proportions (in %) of each descent type (yellow = head-first descent, light green = side descent, and dark green = tail-first descent) occurring on each support category (vertical branches of large, medium, and small diameters represented by brown icons at the top of the graph), by species. Proportions of all individuals of the same species (n) were averaged. See Table S1 for details on the individuals studied, and Table S2 for the mean body masses (BM) references. NA = no data for the given condition. Results of statistical tests between support diameters (ANOVAs or MANOVAs on bootstrapped samples, confirmed by post hoc tests) are represented with annotations defined as NS (non-significant): $p > 0.05$, *: $p < 0.05$, **: $p < 0.01$, and ***: $p < 0.001$. See SuppTables A for associated p-values. **C.** Associated phylogeny of the species studied with branch length obtained from http://timetree.org [130]. **D.** Photographs of individuals of various species descending vertical supports of different diameters.

Kinematic adaptations during ascents and descents on vertical supports

For each complete ascent and descent sequence of locomotion, characterized by successive lift-off, touchdown, and following lift-off of all four limbs, we quantified: the absolute speed, the relative speed (in relation to the individual' body length), the forelimbs and hindlimbs' duty factors (Fig. 2), and the gait type (categorized as symmetrical or asymmetrical, Fig. 3).

We found a significant general tendency for all species to reduce their absolute speed by approximately 30 to 43 % when descending head-first on vertical supports compared to ascents, regardless of support diameter (Fig. 2A, SuppTables B). In contrast, primates did not significantly reduce their speed compared to ascents when descending sideways or tail-first (Fig. 2A, SuppTables B) . On the contrary, in primates the relative speeds decreased during side and tail-first descent on all supports by between 17 and 46 %, indicating that animals tended to decrease the distance travelled compared to ascents (Fig 2B, SuppTables C). But during head-first descents the relative speeds tended to increase by ∼ 10 % on medium supports, while it decreased on small and large supports. It is interesting to note that the smallest species of our dataset, i.e. rodents and treeshrews (< 200 g), exhibited the highest relative speeds compared to other taxa, thus travelling larger distances in proportion to their size (Fig. 2B). On the other hand, all studied animals tended to increase both forelimbs and hindlimbs duty factors during descents compared to ascents, particularly during head-first descents on all supports by ∼ 4%, and during side descents on medium and small vertical supports, but not when descending tail-first (Fig. 2C, SuppTables D).

Figure 2. Kinematics of locomotion during ascents and descents on vertical supports of varying diameters. A. Mean absolute speeds by species during ascents and each descent strategy on the vertical supports of large, medium, and small diameters (represented by brown icons on the top left of the plots), with box plot representations of the medians for all species combined in ascents and each descent strategy. **A.** Mean relative speeds (based on individuals' body length) by species during ascents and each descent strategy on each support type, with box plot representations of the medians for all species combined in ascents and each descent strategy. **C.** Mean duty factors by species, differentiating forelimbs (left box plots) and hindlimbs (right box plots), during ascents and each descent strategy on each support type, with box plot representations of the medians for all species combined in ascents and each descent strategy. Results of statistical tests between ascents and each descent strategy (two-sided Mann Whitney U tests) are represented with annotations defined as NS: $p > 0.05$, *: $p < 0.05$, **: $p < 0.01$, and ***: $p < 0.001$. See SuppTables B, C and D for associated p-values. Percentages correspond to the relative variation of each kinematic variable mean in each descent condition compared to ascents.

We found that gait contains a low, yet significant phylogenetic signal ($K_{mult} = 0.238$; $p = 4.31.10^{-2}$; Effect Size = 1.72), suggesting the impact of shared evolutionary history of related

species on locomotor characteristics. We additionally found that there is a significant similarity between phylogenetically related species and dissimilarity across phylogenetically distant groups, allowing us to merge the gait data at the phylogenetic group level (as defined in Fig. 1C) to obtain a more general picture of the large dataset (MANOVA: p = $2.09.10^{-2}$, Wilks' λ = $3.36.10^{-3}$, df = 5, SuppTables E). Considering the difference in the proportions of descent strategies obtained between strepsirrhines and platyrrhines, we decided to keep these two subgroups separate rather than grouping all primates together. When analyzing the effect of vertical descent strategies on gait types by phylogenetic groups, we found that animals exhibited a mosaic of gait types depending on the species, support diameter and direction of movement. Interestingly, the proportion of asymmetrical gaits significantly increased during descents on all supports for all groups studied (Fig. 3, SuppTables F). In effect, during vertical descents, the animals exhibited more bounds, gallop, and related asymmetrical gaits with relatively high duty factors, not only during side and tail-first descents, but also during head-first descents (Fig. S1). Primates and marsupials overall exhibited a higher proportion of symmetrical gaits compared to the other studied mammals and particularly diagonal and lateral sequence - diagonal couplets (DSDC and LSDC respectively), especially during vertical ascents (Fig. 3, see Fig. S1 for detailed gait types by species and support diameters). We also documented symmetrical gaits in rodents and carnivorans. Primates, marsupials, and rodents exhibited more symmetrical gaits during ascents on vertical supports of small and medium diameter compared to large vertical supports. Carnivorans, especially raccoons, exhibited a relatively small proportion of symmetrical gaits on large and medium supports during ascents, and treeshrews used only asymmetrical gaits during both ascents and descents on vertical supports of medium and large diameter. Furthermore, strepsirrhines exhibited some lateral sequence – lateral couplets (LSLC) only during vertical descents, and rodents and carnivorans exhibited more DSDC during descents and more LSDC during ascents.

Figure 3. Proportions (in %) of gait types by support diameters for ascents and each descent strategies, for all phylogenetic groups. Results of statistical tests on the fraction of symmetrical versus asymmetrical gaits by group between ascents and descents on each support (two-sided Wilcoxon signed rank tests) are represented with annotations defined as NS: p > 0.05, *: p < 0.05, **: p < 0.01, and ***: p < 0.001. NA = not applicable. DSDC: Diagonal sequence - diagonal couplet, LSDC: Lateral sequence – diagonal couplet, LSLC: Lateral sequence – lateral couplet. See SuppTables F for associated post hoc corrected p-values. See Fig. S1 for detailed proportions of gait types by species and support diameters.

Correlations between vertical descent strategies and morphological features

To investigate links between vertical descent strategies and morphological parameters, we collected ten morphological variables including body mass (Table S1), eight intrinsic body proportions calculated from measurements taken on each studied animal (Table 1), and relative head mass of each species, approximated by its encephalization quotient (EQ) (Table S2). We used the EQ [83] as it quantifies whether the brain mass of a given species is lower (<1) or higher (>1) than expected from a general brain-to-body allometric relationship. We calculated the mean EQ for each species from their mean endocranial volumes, collected either from the literature or from our own measurements for previously documented species (Tables S2, S3). To ensure that the EQs of the different species studied are comparable and meaningful, we tested the allometry between the brain and body masses in our dataset following [84] and found a significant and positive slope for all species (major axis orthogonal regression on log

transformed values: slope = 0.36, r^2 = 0.92, p = $5.0.10^{-12}$), indicating a negative allometry (r = -0.97, df = 19, p = $2.0.10^{-13}$), and similar allometric coefficients when restricting the analysis to phylogenetic groups (Fig. S2).

Table 1. Definitions of body measurements performed on extant and extinct individuals and calculation of limbs proportions.

Length measured	Description
Body length	Total body length without tail (from nose to rump)
Tail length	Total tail length
Forelimb length	Total length of the forelimb (stylopod + zeugopod + autopod)
Hindlimb length	Total length of the hindlimb (stylopod + zeugopod + autopod)
Hand length	Maximal length from the tip of the 3rd digit to the proximal carpus
Foot length	Maximal length from the tip of the 3rd digit to the proximal tarsus
Pollex length	Maximal length from the tip of the distal phalanx to the proximal part of the metacarpus
Hallux length	Maximal length from the tip of the distal phalanx to the proximal part of the metatarsus
Proportion calculated	**Description**
% Tail to body length	$\dfrac{Tail\ length \times 100}{Body\ length}$
% Forelimb to body length	$\dfrac{Forelimb\ length \times 100}{Body\ length}$
% Hindlimb to body length	$\dfrac{Hindlimb\ length \times 100}{Body\ length}$
Intermembral index	$\dfrac{Forelimb\ length\ without\ hand \times 100}{Hindlimb\ length\ without\ foot}$
% Hand to forelimb length	$\dfrac{Hand\ length \times 100}{Forelimb\ length\ without\ hand}$
% Foot to hindlimb length	$\dfrac{Foot\ length \times 100}{Hindlimb\ length\ wihout\ foot}$
% Pollex to hand length	$\dfrac{Pollex\ length \times 100}{Hand\ length}$
% Hallux to foot length	$\dfrac{Hallux\ length \times 100}{Foot\ length}$

Among the ten morphological variables analyzed, we found that the forelimb and hindlimb relative lengths correlate with each other and with the mean EQ, and the hindlimb and tail relative lengths correlate with each other (SuppTables G). Moreover, we found that body mass, EQ, relative forelimb and hindlimb lengths, and relative tail length significantly and positively correlate with the proportion of head-first descent strategy on vertical supports, while the intermembral index significantly and negatively correlates with the proportion of head-first descent strategy (all diameters combined) (Fig. 4 A, B, C, Fig. S3 A, B, C, SuppTables H). Also, in primates, only the relative length of the hallux correlates significantly and negatively

with the proportion of side descents (Fig. 4 D, SuppTables I). Primates stand out from the other studied groups in having relatively longer hindlimbs (> 70% of body length, Fig. 4C), forelimbs (> 70% of body length, Fig S3A), tails (> 90% of body length, except for the lorises who have a very reduced tail, Fig. S3C), and high EQs (> 1.5, Fig. 4B). In comparison, carnivorans, albeit being the largest studied species (> 3kg, Fig. 4A), possess relatively shorter limbs (~ 60% of body length, Fig. 4C, Fig. S3A), and medium EQs (> 1 and < 1.2, Fig. 4B). The studied rodents, scandentians, and marsupials also all possess relatively shorter limbs (< 70% of body length, Fig. 4C, Fig. S3A), and tails (< 90% of body length, Fig. S3C), and show small to medium body masses (< 3 kg), and low EQs (≤ 1).

Figure 4. Morphological correlates with vertical descents and prediction of descent strategy in extinct species. Spearman correlations and associated corrected p-values between the proportion of head descents by extant species and **A.** the logarithm of mean body mass, **B.** the mean encephalization quotient (EQ), **C.** the hindlimb to body length ratio (in %), and **D.** between the proportion of side descent in extant primate species only and the hallux to foot length ratio (in %). Each plot includes also the head

descent proportion (**A.**, **B.**, **C.**) and side descent proportion (**D.**) of fossils predicted with the Schafer's multiple imputation procedure. See Table 1 for body measurements definitions and Tables S2 and S3 for mean body masses, ECV and calculated EQ by species. See Fig. S3 for the plots of the percentage of head descent against the six other morphological variables analyzed by species, and SuppTables H and I for Spearman rhos and associated corrected p-values between each morphological variable and the proportion of respectively head (SuppTables H) and side descents (SuppTables I) on vertical supports. See Table S4 for calculated morphological variables of extinct species and SuppTable J for the resulting predictions of their descent behavior.

Integrating extant behavioral and morphological data to hypothesize vertical descent strategy in early euarchontoglires

Based on these results, we attempted to infer the likely descent strategies of extinct species, using their known body proportions. We selected 13 fossils relevant to the understanding of early primate evolution including 7 euprimates, 5 plesiadapiformes and 1 rodent with relatively complete postcranial and/or cranial remains (Table S2, S4). We used a Multiple Imputation (MI) method to predict descent behaviors given partial fossil morphological data. Following the benchmark of Clavel and colleagues [85] which compared several MI methods for the precise problem of morphometric fossil estimation, we used Schafer's MI procedure from the R package *norm*, with 5000 runs of 50 steps Markov-Chains. We calculated the prediction of head-first descent proportion on vertical supports (all diameters combined) for all fossils, using all extant and extinct species, and retaining all six morphological variables that significantly correlate with head-first descent (Fig. 4A, B, C, Fig. S3A, B, C). We also computed the prediction of side descents only for primate and plesiadapiform fossils, retaining all extant primate species and the respective correlated variable, i.e., the relative length of the hallux (Fig. 4D).

Fossil species were mostly reconstructed as employing a relatively high proportion of vertical head-first descent strategy (> 75%), except for *Darwinius masillae* and *Europolemur kelleri,* two medium-sized (660g and ~1.5kg respectively) adapiformes (early strepsirrhines primates), that were estimated at around 30% and 60% of vertical head-first descent strategy respectively (Fig. 4A, C, Fig. S3A, B, C, SuppTables J). *Darwinius* remains are very well preserved and we were able to calculate all the morphological variables for this species. For *Europolemur*, however, we could only include the estimated body mass and foot proportions in the analysis. The other adapiforms of this study, *Adapis parisiensis* and *Notharctus tenebrosus*, were reconstructed with high head-first descent proportions (~ 83% and ~ 86% respectively) despite being larger (~ 1.5kg and ~ 9kg respectively). However, we could not find suitable illustrations of the known postcranial elements of these species in the literature that could be

reliably incorporated into this study. Thus, we only included their reconstructed body mass and EQ, the latter being relatively low (0.56 and 0.26 respectively) compared with extant strepsirrhines. Interestingly, the omomyiforms (early haplorrhines) studied here, *Archicebus achilles*, *Microchoerus erinaceus*, and *Tetonius homunculus* were reconstructed with very high head-first vertical descent proportions (> 90%, Fig 4A, B, SuppTables J), but they are also known mainly from their cranial remains and only body mass and EQ were used in the head-first descent model. Yet, their EQs are also very low (< 1) compared with those of extant haplorrhines (Fig 4B, Table S2, SuppTables J). Concerning the plesiadapiformes and the fossil rodent studied, which display variable body masses ranging from the small *Carpolestes simpsoni* (~ 100 g) to the medium-sized plesiadapids (~ 2 kg, Fig. 4A, Table S2), they were all reconstructed as exhibiting a relatively high proportion of head-first descent strategy (> 75%, SuppTables J). Those species possess relatively elongated limbs whose proportions fall between extant non-primates and primates (Fig. 4C, Fig S3A, B, Table S4), but retain a low EQ (Fig. 4B, Table S2). Concerning the reconstruction of side descent strategy based on hallucal proportions, our model reconstructed the omomyiform *Archicebus achilles* with a low side descent proportion (23%, Fig 4D, Table S4, SuppTables J), despite its small size (~ 25g) but relatively very long hallux (50% of foot length), similar to extant strepsirrhines. The same holds for the adapiforms *Darwinius masillae* and *Europolemur kelleri* that were estimated with moderate side descent proportions (33% and 55% respectively, Fig. 4D, SuppTable J). Finally, the plesiadapiforms *Carpolestes simpsoni* and *Plesiadapis insignis,* with hallucal proportions similar to the studied platyrrhines, were reconstructed with relatively high side descent proportions (respectively 58% and 75%, Fig. 4D, SuppTable J).

Finally, we investigated if the proportion of head-first descent on vertical supports follows a particular evolutionary pattern. Based on the literature, we included extinct taxa in the phylogeny of extant species (Fig. 5) and compared several quantitative models of evolution using the Akaike Information Criterion (AIC). We considered simple models commonly included in studies of morphological evolutionary dynamics: a single-rate Brownian Motion model (BM), the Early Burst model (EB), and a single-peak Ornstein-Uhlenbeck model (OU). We also considered models with two optima: a multiple-rate BM model (BMM) and a multiple-peak OU model (OUM). We considered various relevant phylogenetic nodes where peak shifts or rate changes could have occurred. We found that OUM models with a peak-shift for 'euprimates + plesiadapiforms' or 'euprimates only' yielded the best fits (AIC= 9.46 and 9.48 respectively) both clearly favored from the other evolutionary models tested (SuppTables K).

These models have a head-first descent proportion theta peak of ~44% and ~45% respectively,

for a theta peak of ~95% and ~93% for other mammals groups respectively. After rescaling the

phylogenetic tree height to 1, the strength of return to the optimum is quite high (alpha = 7.49

and 7.51 for the 'euprimates + plesiadapiforms' and 'euprimates only' models, respectively)

and the phylogenetic half-life is small (0.093 and 0.092, respectively) confirming that these

OUM models fit our dataset best [86].

Figure 5. Phylogeny of studied species, including extant and extinct taxa, and representation of
the best evolutionary model of head-first descent proportion. We calculated a consensus tree for
extant species based on 1000 trees from vertlife.org, to which we added extinct taxa using age and
phylogenetic topology from literature. See Table S2. for estimated age of extinct species and associated
references. The best evolutionary model of head-first descent proportion is an Ornstein-Uhlenbeck
evolutionary model with two optimums (OUM) with a peak shift occurring at the node 'plesiadapiformes
+ euprimates'. The associated theta peaks of head-first descent proportions of the two subgroups
'plesidapiforms + euprimates' vs 'others' are represented in red and black respectively.

Discussion

The variability of vertical descent strategies in primates compared to other mammals

To date, this study is the first comparative quantification of ascent and descent strategies on

vertical supports across a broad sample of small- to medium-sized arboreal mammals. Overall,

our results reveal that vertical descent strategies are strongly influenced by phylogenetic

relationships and not solely by body mass, contrary to our initial hypothesis (H1). Specifically,

not all species >1 kg systematically descended tail-first. Primates exhibited greater variability

in descent strategies—including head-first, side, and tail-first postures—whereas scandentians,

rodents, carnivorans, and marsupials primarily employed head-first descents, with the exception

of raccoons. Both head-first and tail-first postures involve the body remaining parallel to the

support, as previously documented in rodents [12,40,41], procyonids [30,42], prehensile-tailed

platyrrhines [43–45], and non-prehensile tailed strepsirrhines and marsupials [46,73]. However, we

also identified a distinct "side descent" posture, observed exclusively in primates, in which the

body is rotated perpendicular to the support while the face remains directed downward. This

posture likely partly corresponds to the "other descent strategies" described by Perchalski

(2021). We quantified side descents in both strepsirrhines and platyrrhines, with non-callitrichid

platyrrhines using it most frequently (>70%) across all support diameters. In contrast, the

smallest primates (i.e., pygmy slow loris *N. pygmaeus*, ~360 g; and tamarins *S. imperator* ~475

g, *S. oedipus* ~410 g) primarily employed head-first descents, while larger strepsirrhines,

particularly lemurs, used tail-first descents in >80% of trials across all support diameters.

Primates generally increased the use of tail-first and side descents on small vertical supports,

suggesting that narrow supports impose additional constraints favoring more upright postures.

This marked behavioral divergence among phylogenetic groups underscores the role of

evolutionary history and supports recent findings emphasizing the influence of shared

ecological niches and postural adaptations in shaping primate locomotor performance [65,69,87–89].

The kinematic adjustments of arboreal mammals during vertical descents

In terms of locomotion, we found that all species reduced their speed and increased both

forelimb and hindlimb duty factors during head-first descents compared to ascents on vertical

supports, partially supporting our second hypothesis (H2). This pattern, previously observed in

small primates [24,50,51], rodents [12,52,53], and marsupials [34,54,73], suggests that head-first descent

imposes greater demands on maintaining secure grips, leading to more cautious movements

regardless of support diameter, characterized by increased limb contact times and reduced

velocity [48,53]. However, when employing side or tail-first descent strategies, primates did not

significantly reduce absolute speed nor markedly increase duty factors, but instead decreased

relative distance traveled per cycle. This unexpected result suggests that tail-first and side

descent strategies are highly effective, allowing primates to maintain comparable speeds while

adopting flexible manual and pedal postures [19,90]. During vertical ascent, locomotion primarily

relies on hindlimb propulsion [8,91], whereas head-first descent requires increased forelimb

braking and frictional control at the extremities [8,50,51,66,92,93]. Our results suggest that the upright

descent strategies employed by primates mitigate these braking demands on the forelimbs while

preserving static stability and high descent velocity.

Moreover, we found that gait patterns are strongly constrained by phylogenetic history,

consistent with previous findings in primates [65,69,87–89]. Primates, marsupials, rodents, and

carnivorans exhibited both symmetrical and asymmetrical gaits, though primates and

marsupials employed the highest proportions of symmetrical gaits, particularly DSDC and

LSDC gaits. This aligns with earlier reports of habitual DSDC gait use in primates, some

marsupials, and carnivorans, specifically the kinkajou *Potos flavus* [49,56,59,63,73,74,94]. Interestingly,

only strepsirrhines increased their use of LSLC gaits during descents, as previously observed

in platyrrhines on sloped supports [51,65–69], whereas rodents and carnivorans favored more DSDC

gaits during descents compared to more LSDC gaits in ascents. Both DSDC and LSLC gaits

have been proposed to enhance stability by facilitating precise fore- and hindfoot placement,

especially during horizontal walking on narrow branches [49,63,95]. On the other hand, LSDC gaits

promote stability by keeping the location of the center of mass relative to the support polygon

of the limbs [58,59,63,68,95,96]. Most previous studies have focused on symmetrical gaits, particularly

DS, due to their proposed adaptive significance for fine-branch locomotion in primates [56,63,65].

However, several studies have also reported increased use of asymmetrical and LS gaits on

sloped and vertical supports in primates and other arboreal mammals, emphasizing the

importance of behavioral flexibility rather than strict kinematic specialization in arboreal

environments to cope with complex 3D environments [12,53,65,66,94,97].

We found that the proportion of symmetrical gaits increased on small vertical supports,

partially validating our second hypothesis (H2), but only during ascents. During descents, the

proportion of asymmetrical gaits significantly increased across all supports and species,

regardless of descent strategy. This key finding suggests that vertical locomotion requires

distinct kinematic adjustments: symmetrical gaits enhance climbing efficiency, whereas

asymmetrical gaits facilitate stability during descents. Symmetrical gaits, characterized by

regular stance and swing phases, allow for controlled force application, lower reaction forces,

control and transfer of the moments and torques imposed on the body axes, and minimized

mediolateral deviations, keeping the center of mass within the support polygon throughout the

stride [56,59,63,64]. Conversely, asymmetrical gaits permit higher speeds, reduce metabolic cost,

provide an energetic input to locomotion through the sagittal movement of the spine, and

stabilize locomotion by reducing peak forces and center of mass oscillations through shorter,

more frequent strides [24,52,53,60–62]. Overall, while symmetrical gaits enhance stability on narrow

supports during horizontal or oblique locomotion and during vertical climbing [11,52,64,66,71,74],

asymmetrical gaits may offer superior control and security during vertical descents by enabling
simultaneous grasping with both fore- and hindlimbs [51–53,60–62,70].

The correlation between morphological adaptations and vertical descent strategies

We identified a clear influence of body proportions on head-first descent behavior in our
dataset, with body mass, relative forelimb, hindlimb, and tail lengths, and relative head size all
significantly affecting descent strategies. Heavier animals with relatively larger heads as well
as elongated limbs and tails were more likely to avoid head-first descents, partially supporting
our third hypothesis (H3). As predicted, the intermembral index (IMI) was negatively correlated
with the proportion of head-first descents [36,46]. This relationship was largely driven by
strepsirrhines, which exhibited particularly low IMI values compared to other taxa.
Strepsirrhines also performed the fewest head-first descents and the highest proportion of tail-
first descents, highlighting the impact of hindlimb elongation on vertical locomotor
performance by shifting the center of mass posteriorly. In contrast, platyrrhines displayed higher
IMI values (> 80), and all non-primate species exhibited relatively shorter limbs (forelimbs <
55% and hindlimbs < 70% of body length) compared to primates. These findings support
previous hypotheses that elongated hindlimbs may increase the risk of imbalance during head-
first descent, by simultaneously increasing the forelimbs' braking demand and the hindlimbs'
traction requirements [36,46].

We also detected a significant relationship between tail length and descent strategy: primates
generally possessed proportionally longer tails (> 90% of body length) than the non-primate
species. Although tails primarily function in balance during horizontal arboreal locomotion [98–
100] and serve to stabilize the body during leaping or aerial maneuvers [101], increased tail length
may also contribute to a posterior shift of the center of mass that discourages head-first descent.

Interestingly, we also found a negative correlation between relative head size (approximated
by the EQ) and the proportion of head-first descents, suggesting that larger head mass may
further bias animals toward more upright postures during vertical descent. To our knowledge,
this study is the first to explicitly assess the influence of body mass distribution—and
particularly relative head mass—in the context of arboreal locomotion. Within our dataset,
primates, especially platyrrhines, exhibited higher encephalization quotients (EQ > 1.5) and
engaged in head-first descent postures less frequently than other mammals, which possessed
lower EQ values (< 1.2) and relied more on head-first strategies. A larger brain not only
demands increased cerebral blood flow, which may impose physiological limitations during
acrobatic arboreal behaviors, but also increases head mass, thereby raising the moment of

inertia and shifting the center of mass anteriorly. These factors support our hypothesis that an
enlarged head may mechanically constrain animals to avoid head-first vertical descent by
increasing the risk of forward toppling during head-first postures, particularly in primates.
However, it is noteworthy that certain larger arboreal primates (> 5 kg), such as howler monkeys
[43] and orangutans [102], both characterized by higher encephalization and, in the case of
orangutans, markedly elongated forelimbs and elevated intermembral indices [103], have been
observed performing head-first descents in their natural environments. Although these
behaviors appear relatively infrequent, they underscore the need for further investigation into
how brain expansion and associated morphological changes interact to influence arboreal
locomotor strategies across taxa.

Contrary to our expectations (H3), we did not find a significant correlation between the
relative size of the autopods and the proportion of head-first descents. However, within
primates, we found a negative correlation between the hallux length and the proportion of side
descents, with the longer the hallux the less side descents employed. Digital elongation of the
hand and foot in primates has long been interpreted as enhancing grasping efficiency on
arboreal supports [13,19,21,25,75,104]. During vertical ascent the superior point of attachment (i.e., the
hands) ensures a critical hold, whereas in head-first descents both hands and/or feet may ensure
holding. In contrast, tail-first or side descents may allow primates to rely more extensively on
the foot for secure hold, supporting most of the weight. Strepsirrhines possess a highly
specialized foot with a very divergent elongated hallux, while some platyrrhines possess a less
elongated hallux but bear claw-like nails on their distal phalanges which they use to cling to
tree bark similarly to other mammals. These differences in hand and foot morphology among
primates are reflected in their vertical descent behavior: strepsirrhines more frequently adopt
tail-first descents, relying more heavily on the foot as a secure point of support, whereas
platyrrhines more often employ side descents, using both hands and feet to grasp the support
efficiently. Taken together, our results highlight that efficient vertical descent is a complex
phenomenon governed by the interplay of kinematic adjustments, locomotor flexibility, body
mass distribution, limb proportions, grasping ability, and shared evolutionary history among
species. Primates exhibit particular adaptations for vertical descent, including upright postures
and a remarkable capacity for locomotor flexibility, which may derive from their ancestral
adaptation to navigating supports of varying size and orientation [65]. In contrast, other arboreal
mammals appear to have been more restricted to the use of less diversified supports during their
evolutionary histories.

Implications for the reconstruction of early primates locomotion

Our inference model reconstructed fossil species as employing relatively high proportions of head-first descent (> 75%), except for the two medium-sized adapiforms *Darwinius masillae* and *Europolemur kelleri* (estimated at ~30% and 60%, respectively), which already displayed relatively long hindlimbs and tail. This validates our fourth hypothesis (H4) and suggests that postural adaptations for vertical descent in primates evolved progressively. Although some species already reached relatively high body mass, such as the adapiform *Notharctus tenebrosus* (~9 kg, [105], the plesiadapiform *Plesiadapis sp.* (~2 kg, [106]), or the rodent *Ischyromys typus* (~1.3 kg, [107]), they retained low EQs and moderate limb elongation. Interestingly, the plesiadapoids *Carpolestes simpsoni* and *Plesiadapis insignis*, whose phylogenetic positions within Euarchonta remain debated [80,108–110], have been reconstructed with a high proportion of side descent, associated with their slightly elongated hallux similar to extant platyrrhines. Plesiadapiforms are central to the debate on primate origins as the family Plesiadapidae is often regarded as a stem or sister group to euprimates [109,111,112]. However, plesiadapiforms are mainly reconstructed as generalized quadrupedal walkers and climbers [20,111,113,114]. In contrast, early primates (including both omomyiforms and adapiforms) are all reconstructed as vertical climbers and leapers at variable degrees [115–117]. Nonetheless, our evolutionary analysis corroborates current phylogenetic hypotheses suggesting that euprimates and plesiadapiforms share an adaptive regime distinct from the other groups studied [best OUM model: euprimates + plesiadapiforms vs. other], highlighting that descent strategy on vertical supports was tightly constrained throughout evolution.

These results lead us to suggest a refinement of the existing scenario on the order of acquisition of early primate specializations. Considering that early euarchontoglires were probably small to very small (≤ 100g [110]) with shorter hindlimbs, autopods, and reduced brain size, it is plausible that they primarily used mostly head-first descents and asymmetrical gaits on vertical supports [12,53]. As euprimates evolved enhanced grasping abilities, elongated hindlimbs and tail, and eventually larger brains, they likely began to adopt side and tail-first vertical descent postures. The evolutionary success of early primates likely resided in their ability to colonize environments which are mechanically more challenging and difficult to access by other animals, such as horizontal terminal branches and small vertical supports. The frequent use of vertical supports, such as trunks, lianas, climbing plants, or even tall grass thickets, favored the early acquisition of a divergent hallux and pollex, improving grasping ability [15,18,19]. These supports were abundant in the tropical forests of the northern hemisphere during the Paleogene, the likely origin place and time of primates [80,118,119]. The use of these

supports may have facilitated rapid access to the canopy, provided platforms for scanning for food or predators, efficient escape routes and rapid changes of height, and possibly privileged access to a greater diversity of food sources [120]. Autopodial innovations were followed hindlimb elongation, likely enhancing leaping abilities as inferred in various early euprimate fossils [16,121]. Altogether, these hindlimb specializations, followed by a progressive increase of the body and tail lengths, and brain size, probably altered body mass distribution. This may have further promoted side- and tail-first vertical descent strategies, allowing efficient vertical navigation while maintaining environmental awareness. Such adaptations may have thus favored primates to frequently adopt upright postures when negotiating vertical supports in an early stage of their evolutionary history, while increasing the use of symmetrical gaits to negotiate other types of supports such as narrow horizontal branches. Building on previous hypotheses on autopodial grasping evolution of primates [21,23,122,123], upright orthograde postures would have also freed the hand from locomotion, while the foot remained specialized for efficient support holding. Functional differences between hand and foot have emerged early in primates [5,92,124,125], with the anchoring foot playing a fundamental role in body support, and the hand redirected toward other activities, such as food grasping and manipulation, substantiated by the great diversity of manual postures in extant species [19,126,127].

Limitations of experimental conditions and future directions

Every study that includes experiments with animals housed in zoos displays some limitations. Some enclosures required enrichment with new supports and in some cases the animals did not readily use the support provided (see the Materials and Methods section). It is therefore possible that this experimental bias may have had an impact on the proportion of descent behaviors and the kinematics quantified for some species. For instance, although raccoons and coatis have been observed to descend head-first on small and medium supports in the wild (personal observations) they did not use them in their enclosures. Thus, quantifying vertical descent strategies as well as the relative frequency of vertical support use in the wild would provide more conclusive evidence regarding the vertical descent repertoire in those species. Moreover, comparing kinematics of vertical ascents and descents in more species of larger body mass (> 5 kg) with varying morphological proportions, such as apes and large carnivorans, would help to better understand the complex parameters at play regarding mechanical adaptation to arboreal locomotion in larger mammals. Such studies will also allow to develop reconstruction models better adapted to large fossils, particularly those primates with

609 relatively lower EQ than extant species, thereby refining our understanding of the evolution of
610 those adaptations within this group.

Materials and Methods

Experimental design

<u>Animals studied</u>

We collected data on animals housed in zoological parks except for the forest dormice
Dryomys nitedula which were wild caught (Table S1). In zoos, individuals were video recorded
directly in their enclosures. All the enclosures were large enough to allow unconstrained
locomotion of the animals. We placed non-treated wooden branches of variable diameters and
orientations in each enclosure before filming to ensure equal access to a variety of support
types. All studied individuals were adults, in good shape, and did not display any stereotypical
behavior before or during the experiments. The two individuals of *Dryomys nitedula* were
caught in their natural habitat in Vasilika, Greece, with the use of specialized wooden nest boxes
positioned in trees (permit by the Hellenic Ministry of Environment and Energy 8817/242).
They were immediately transferred and housed to glass terrariums (L= 100; W= 40; H= 50 cm)
enriched with tree branches at room temperature and with normal photoperiod (complying to
the regulations of the Ethics committee of the School of Biology, Aristotle University of
Thessaloniki, Greece). After observations and filming directly in these terrariums, the animals
were released at the same location they were originally captured. Animal observation was
performed in compliance with the International Primatological Society (IPS) Ethical Guidelines
for the Use of Nonhuman Primates in Research and the Association for the Study of Animal
Behavior (ASAB) and the Animal Behavior Society (ABS) ethical guidelines for the use of
animals in research [128].

<u>Behavioral data collection</u>

We focused on vertical supports (90°± 5°). We categorized support diameters into three
categories: small (support circumference < hand and foot lengths), medium (support
circumference ≈ hand and foot lengths), and large (support circumference > hand and foot
lengths) following previously described methods [19], resulting in three support types tested for
each individual. Sequences were recorded using a portable video camera (Panasonic HC-V770
camcorder 120fps, full HD 1080p). To ensure both a complete view and close ups of the animals

during locomotion, we also used three small action video cameras (Mobius ActionCam 60fps,

HD 720p) installed in three different view angles (frontal, lateral and ventral) and in relatively

close distance (between 20 and 50cm depending on the size of the animals) from each support

type. After a habituation period, the individuals were recorded using an alternation of scan

sampling and focal-animal sampling methods [129] in continuous recording sessions of 10 to 30

min. Animals were filmed as they moved freely over supports, and we stimulated them by

providing small pieces of food placed randomly along the supports when necessary. For

nocturnal species (Table S1), observations were conducted either in artificial or real nocturnal

conditions with addition of red lighting to enable recording without disturbing the animals.

<u>Video analysis</u>

We analyzed videos using Adobe Premiere Elements 12. All video analysis was performed

by the same person (SLDT) to avoid inter-observer biases. We focused on locomotion only and

excluded other behaviors. We analyzed a minimum of three and up to ten passages in ascent

and ten passages in descent for each individual on each support type (i.e. small, medium, and

large), resulting in up to thirty ascents and thirty descents analyzed for each individual, with the

following exceptions: we could not obtain data for *Tupaia belangeri* on small supports, for

Graphiurus murinus on small supports; for *Typhlomys chapensis* on medium supports, for

Procyon lotor on small supports, and for *Nasua nasua* on the medium and small supports. For

Tupaia belangeri, *Graphiurus murinus*, and *Typhlomys chapensis*, these supports were not

available during the data acquisition process. However, for *Procyon lotor* and *Nasua nasua*,

these supports had been added and were available in the enclosures, but were not used by the

animals.

<u>Kinematic analysis of locomotion</u>

To quantify the absolute speed of the individuals for each sequence of locomotion we

extracted two snapshots and their associated frame number from the beginning and the end of

the movement, using videos with a lateral view relative to the support. We merged both images

using the product layer functionality in Adobe Photoshop CS6. We then measured in ImageJ

the distance travelled by the individuals during each stride using markers of known length

positioned in each support as references. We calculated the associated speed as the ratio of this

distance to the duration separating the snapshots.

We also calculated the relative speed for each sequence of locomotion as the ratio of the

absolute speed to the body length of each animal studied, defined as the total length between

the top of the nose to the base of the tail. We used snapshots of each individual to measure body

lengths in ImageJ, using a scale marker as a metric reference positioned on top of the branches.

We analyzed the strides for each sequence of locomotion by recording the frame number of

the first touchdown, the following lift-off and the subsequent touchdown of each limb on the

support. We calculated the associated stride duration of each limb, using the known frame rate

of the associated videos. We then calculated the duty factor (DF) of each limb as the percentage

of time spent in contact with the support relatively to the total stride duration. Forelimbs and

hindlimbs duty factors were averaged to obtain the mean DF.

We classified the gait type following previously described methodology [61]. We used the

midpoints of each limb contact intervals (i.e., the temporal midpoint between first touchdown

and following lift-off). We categorized gaits as symmetrical when the temporal lag between left

and right forelimbs and hindlimbs midpoints were both $\geq 40\%$ and $\leq 60\%$ of the total stride

duration, and asymmetrical otherwise. Further categorization among symmetrical gaits were

made using the limb phase, which corresponds to the percentage of total stride duration

separating a hindlimb touchdown from its ipsilateral forelimb touchdown. We thus calculated

the left and right limb phases using the midpoints of forelimbs and hindlimbs contact intervals.

Left and right limb phases were then averaged to obtain the mean limb phase for each stride.

We categorized as lateral sequence – lateral couplet (LSLC) when the mean limb phase was $\geq$

0% and $< 25\%$, as lateral sequence – diagonal couplet (LSDC) when $\geq 25\%$ and $< 50\%$, as

diagonal sequence – diagonal couplet (DSDC) when ≥ 50 and $< 75\%$, and as diagonal sequence

– lateral couplet (DSLC) when $\geq 75\%$ and $\leq 100\%$. Classification of asymmetrical gaits used

the stance period, which corresponds to the percentage of limb pair total contact duration

separating left and right limbs of the same pair. We classified as bounds the gaits when both

forelimb and hindlimb stance periods were $\leq 10\%$, as half-bounds those when either (i) the

forelimbs stance period was $\geq 10\%$ and the hindlimbs stance period was $\leq 10\%$, or (ii) when

the forelimbs stance period was $\leq 10\%$ and the hindlimbs stance period was $\geq 10\%$, and as

gallop when both forelimb and hindlimb stance periods values were $\geq 10\%$.

<u>Morphological data collection</u>

For each studied individual, we extracted three to five lateral view screenshots from the

videos. On these images we measured lengths in ImageJ allowing to calculate eight intrinsic

body proportions (Table 1). We calculated the average proportions over screenshots for each

individual, and ultimately the average proportions by species.

The mean body mass of each species studied was gathered from the literature (Table S1). We could have used the actual body masses of certain individuals studied in zoos, but we could not have access to this information for most species, so we decided to use the standard reference values from literature. Additionally, we calculated the mean encephalization quotient (EQ) of these species as a proxy for their relative head to body size ratio and to allow comparison with fossil specimens for which the EQ is commonly studied and provided in the literature (Table S2). Prior to the calculation of the EQ in extant species, we collected the mean endocranial volumes (ECV) either from the literature or from own measurements in the case of species that were not previously documented (Table S3). For these species we quantified the ECV from preserved skulls of specimens housed in the anatomical collections of the Museum für Naturkunde of Berlin, by filling their endocranial cavity with chia seeds of known volumic mass to infer their volume. We thus provide hereby the mean endocranial volumes for the treeshrew *Tupaia belangeri*, the rodents *Dryomys nitedula*, *Graphiurus murinus* and *Typhlomys cinereus*, and the marsupial *Petaurus breviceps* (Table S3).

For comparative purposes, we collected from the literature the reconstructed body mass and EQ of key fossil specimens of euarchontoglires with relatively complete postcranial and/or cranial remains (Table S2). Whenever possible, we also measured the same postcranial lengths as for extant species, using published photographs (see Table S2 for associated references, and Table S4 for body proportions calculated in fossils).

Construction of phylogenetic tree including fossil taxa

For illustration purposes in Fig. 1C, we used a phylogeny of extant species from timetree.org [130] which derives from an aggregation of molecular phylogenetic data.

For phylogenetic comparative analysis (phylogenetic signals and evolutionary model comparisons) we generated 1000 trees from the tip-date DNA-only version of mammalian trees of the studied extant species from vertlife.org [80]. We computed a consensus tree using the *consensus* function of the R package *ape* and made it strictly ultrametric by extending the tips with the *force.ultrametric* function from the R package *phytools*. To build and compare evolutionary models of the fossils' descent behaviors, we supplemented this tree of extant species with the extinct taxa studied (Fig. 5) using estimated age and phylogenetic topology from literature (Table S2).

Statistical analysis

Statistical analysis was performed with RStudio version 1.3.1093 (R version 3.6.3).

<u>Proportions of descent strategies</u>

To assess the effect of support diameter (e.g., small vs. medium vs. large) on the proportion

of descent types between individuals of the same species, we performed either analysis of

variance (ANOVA) or multivariate analysis of variance (MANOVA) tests (respectively *aov*

and *manova* functions in R) depending on the number of descent strategies exhibited (Fig. 1B,

SuppTables A). To even out the testing power among species for which we had a different

number of individuals, we bootstrapped 10 individuals and 10 descent events on each support

category to perform the tests. P-values were then Benjamini-Hochberg corrected.

To assess the extent of phylogenetic information contained in the proportion of head-first

descents on vertical supports, we computed Blomberg's K phylogenetic signal [131] using the

physignal function from the *geomorph* package in R with 50000 iterations.

<u>Locomotion</u>

To test for the global differences of speed, relative speed, and duty factor between ascents

and each descent type, we performed two-sided Mann Whitney U tests on 40 bootstrapped

events (20 ascents and 20 descents) for all species on each support diameter. P-values were then

Benjamini-Hochberg corrected. We also calculated the mean percentage of variation between

ascents and each descent category (Fig. 2, SuppTables B, C, D). For tests over duty factor, we

considered both forelimbs and hindlimbs combined.

We calculated the frequency of all possible gait type values occurring for each individual on

each support category differentiating ascents and the different descent strategies when relevant.

These frequencies were PCA-transformed to obtain linearly independent values which serve as

low-dimensional representation of the overall gait data. Using the same *physignal* function as

described above for phylogenetic signal computation of head-first proportion on vertical

supports, we computed the K_{mult} statistic [132] with 50000 iterations on the PCA-transformed data

of gaits, keeping only the first principal components that retain 95% of variance, to assess the

extent of phylogenetic information contained in gaits.

To check for enough similarities between species of the same phylogenetic group as defined

in Fig. 1C, and dissimilarities between groups to justify pooling data at the group level, we

performed MANOVAs with Benjamini-Hochberg corrected post hoc pairwise comparisons on

PCA-transformed frequencies of gaits, keeping principal components explaining at least 95%

of variance, using the *manova* function from the *stats* package in R and Wilks's test

(SuppTables E).

To test for changes of the fraction of symmetrical versus asymmetrical gaits between ascents and descents for each group of species and on each support diameter separately, we performed two-sided Wilcoxon ranked signed tests on bootstrapped samples. For each group and each support, we took 100 bootstrapped species and sampled 20 events (10 ascents and 10 descents) at random. The resulting p-values were Benjamini-Hochberg corrected (Fig. 3, SuppTables F).

<u>Morphological correlates</u>

To control that brain allometry is homogeneous among all phylogenetic groups, to be able to compare EQ between species, we computed major axis orthogonal regressions, following the recommendation of Martin and Barbour [84], between the Log transformed brain and body masses, over all species and by phylogenetic group using the *sma* package in R (Fig. S2).

To assess the level of correlation among the 10 morphological variables, we computed Spearman correlations on all pairs. P-values were then Benjamini-Hochberg corrected (SuppTables G).

We computed Spearman correlations to quantify the relation between morphological variables and the proportions of head-first descent (for all species) or side descent (for primates) on vertical supports (all diameters combined) and the resulting p-values were Benjamini-Hochberg corrected (Fig. 4, Fig. S3, SuppTables H, I).

<u>Estimation of extinct species vertical descent strategy</u>

We isolated the six morphological variables for which we obtained a significant correlation with head-first descent proportions on vertical supports (body mass, EQ, relative forelimb length, relative hindlimb length, intermembral index and relative tail length). We used the Schafer's Multiple Imputation procedure from the R package *norm* to compute estimations of the fossils' proportions of head-first descent given their respective morphological features (Fig. 4 A, B, C, Fig. S3 A, B, C, SuppTables J). We simulated 5000 runs of Markov Chains with 50 steps to infer the head-descent proportions of these 13 extinct species. We conducted the same procedure to then infer the proportion of side descent in euprimates and plesiadapiforms fossils by restricting the set of extant species to primates, and using the only morphological variable which correlated with side descents (relative hallux to foot length, Fig. 4 D, SuppTables J).

<u>Evolutionary model comparison for head-first descent strategy</u>

To evaluate if the proportion of head-first descent strategy on vertical supports follows a specific evolutionary model, we compared different quantitative models of trait evolution in all

extant and extinct taxa studied. We first fitted three simple models: Early Burst (EB), Brownian Motion (BM) and Ornstein-Ulhenbeck (OU) using the *mvEB, mvBM and mvOU* functions from the R package *mvMorph* respectively [133]. Then, we considered models with two distinct optima within the phylogeny. We fitted the Brownian Motion and Ornstein-Ulhenbeck models with multiple optima (respectively BMM and OUM) using the *mvBM* and *mvOU* functions. We tested a change of the evolutionary optimum at different phylogenetic nodes: I) euprimates, II) euprimates + plesiadapiforms, III) euarchontans (euprimates + plesiadapiforms + scandentians), and IV) euarchontoglires (euarchontans + rodents). We compared models using the Akaike Information Criterion (AIC) and estimated the phylogenetic half-life in the OUM models that best fitted head-first descent strategies in our sample, using the *halflife* function from *mvMorph*, after rescaling the phylogenetic tree height to 1 (SuppTables K).

Acknowledgements

We greatly thank the staff of the Zoological and Botanical Parc of Mulhouse, the Zoological Park of Paris, and the Spaycific'Zoo (France) for access to their animals and for their assistance during material installation and data collection. We thank Dr. A. Panyutina and Dr. A. Kuznetsov (Lomonossov Moscow State University and Russian Academy of Sciences) and the Moscow Zoo staff (Russian Federation) for video recordings of *Graphiurus murinus*, *Typhlomys chapensis*, and *Tupaia belangeri*. We thank Dr. N.E. Karantanis and the staff of the Nowe Zoo Poznan (Poland) for the video recordings of *Xanthonycticebus pygmaeus*. We thank the staff of the Laboratoire d'Ecologie Generale in Brunoy (France) for the video recording of *Caluromys philander*. We thank C. Funk from the Museum für Naturkunde of Berlin for access to the collection of mammal specimens. We thank Dr. A. Llamosi for his help with statistical analysis.

References

1. Hildebrand, M. *Analysis of vertebrate structure, 4th edn.* (John Wiley & Sons, Inc., New York, 1995).

2. Zheng, X., Bi, S., Wang, X. & Meng, J. A new arboreal haramiyid shows the diversity of crown mammals in the Jurassic period. *Nature* **500**, 199–202 (2013).

3. Bi, S., Wang, Y., Guan, J., Sheng, X. & Meng, J. Three new Jurassic euharamiyidan species reinforce early divergence of mammals. *Nature* **514**, 579–584 (2014).

4. Meng, Q.-J. *et al.* An arboreal docodont from the Jurassic and mammaliaform ecological diversification. *Science* **347**, 764–768 (2015).

5. Wood Jones, F. *Arboreal Man*. (Edward Arnold, 1916).

6. Cartmill, M. Climbing. in *Functional Vertebrate Morphology* (eds. Hildebrand, M., Bramble, D. M., Liem, K. F. & Wake, D. B.) 73–88 (Harvard University Press, 1985). doi:10.4159/harvard.9780674184404.c5.

7.   Hunt, K. D. *et al.* Standardized descriptions of primate locomotor and postural modes. *Primates* **37**, 363–387 (1996).

8.   Hirasaki, E., Kumakura, H. & Matano, S. Biomechanical analysis of vertical climbing in the spider monkey and the
Japanese macaque. *Am. J. Phys. Anthropol.* **113**, 455–472 (2000).

9.   Isler, K. 3D-kinematics of vertical climbing in hominoids. *Am. J. Phys. Anthropol.* **126**, 66–81 (2005).

10.   Hanna, J. B., Schmitt, D. & Griffin, T. M. The energetic cost of climbing in primates. *Science* **320**, 898 (2008).

11.   Hanna, J. B., Granatosky, M. C., Rana, P. & Schmitt, D. The evolution of vertical climbing in primates: evidence
from reaction forces. *J. Exp. Biol.* **220**, 3039–3052 (2017).

12.   Karantanis, N.-E., Rychlik, L., Herrel, A. & Youlatos, D. Vertical Locomotion in Micromys minutus (Rodentia:
Muridae): Insights into the Evolution of Eutherian Climbing. *J. Mamm. Evol.* **25**, 277–289 (2018).

13.   Nyakatura, J. A. Early primate evolution: insights into the functional significance of grasping from motion analyses
of extant mammals. *Biol. J. Linn. Soc.* **127**, 611–631 (2019).

14.   Dagosto, M. The postcranial morphotype of primates. *PRIMATE Orig. Adapt. Evol.* 489–534 (2007)
doi:10.1007/978-0-387-33507-0_15.

15.   Gebo, D. L. Vertical clinging and leaping revisited: Vertical support use as the ancestral condition of strepsirrhine
primates. *Am. J. Phys. Anthropol.* **146**, 323–335 (2011).

16.   Boyer, D. M., Toussaint, S. & Godinot, M. Postcrania of the most primitive euprimate and implications for primate
origins. *J. Hum. Evol.* **111**, 202–215 (2017).

17.   Yapuncich, G. S., Feng, H. J., Dunn, R. H., Seiffert, E. R. & Boyer, D. M. Vertical support use and primate origins.
Sci. Rep. 1–10 (2019) doi:10.1038/s41598-019-48651-x.

18.   Szalay, F. S. & Dagosto, M. Evolution of hallucial grasping in the primates. *J. Hum. Evol.* **17**, 1–33 (1988).

19.   Toussaint, S., Llamosi, A., Morino, L. & Youlatos, D. The Central Role of Small Vertical Substrates for the Origin
of Grasping in Early Primates. *Curr. Biol.* **30**, 1–14 (2020).

20.   Bloch, J. I. & Boyer, D. M. Grasping primate origins. *Science* **298**, 1606–1610 (2002).

21.   Cartmill, M. New views on primate origins. *Evolutionary Anthropology: Issues, News, and Reviews* vol. 1 105–111
(1992).

22.   Sussman, R. W., Rasmussen, D. & Raven, P. H. Rethinking Primate Origins Again. *Am. J. Primatol.* **75**, 95–106
(2013).

23.   Toussaint, S., Herrel, A., Ross, C. F., Aujard, F. & Pouydebat, E. Substrate Diameter and Orientation in the Context
of Food Type in the Gray Mouse Lemur, Microcebus murinus: Implications for the Origins of Grasping in Primates.
Int. J. Primatol. **36**, 583–604 (2015).

24.   Shapiro, L. J., Kemp, A. D. & Young, J. W. Effects of Substrate Size and Orientation on Quadrupedal Gait
Kinematics in Mouse Lemurs (Microcebus murinus). *J. Exp. Zool. Part A Ecol. Genet. Physiol.* **325**, 329–343
(2016).

25.   Kirk, E. C., Lemelin, P., Hamrick, M. W., Boyer, D. M. & Bloch, J. I. Intrinsic hand proportions of euarchontans
and other mammals: Implications for the locomotor behavior of plesiadapiforms. *J. Hum. Evol.* **55**, 278–299 (2008).

26.   Sargis, E. J. The postcranial morphology of *Ptilocercus lowii* (Scandentia, Tupaiidae) and its implications for
primate supraordinal relationships. *PRIMATE Orig. Adapt. Evol.* 51–82 (2007) doi:10.1007/978-0-387-33507-0_3.

27.   Urbani, B. & Youlatos, D. Positional behavior and substrate use of Micromys minutus (Rodentia: Muridae): Insights
for understanding primate origins. *J. Hum. Evol.* **64**, 130–136 (2013).

28.   Orkin, J. D. & Pontzer, H. The Narrow Niche hypothesis: Gray squirrels shed new light on primate origins. *Am. J.
Phys. Anthropol.* **144**, 617–624 (2011).

29.   Byron, C. D., Kunz, H., Matuszek, H., Lewis, S. & Van Valkinburgh, D. Rudimentary pedal grasping in mice and
implications for terminal branch arboreal quadrupedalism. *J. Morphol.* **272**, 230–240 (2011).

30.   McClearn, D. Locomotion, Posture, and Feeding Behavior of Kinkajous, Coatis, and Raccoons. *J. Mammal.* **73**,
245–261 (1992).

31. Youlatos, D. Hallucal grasping behavior in Caluromys (Didelphimorphia: Didelphidae): Implications for primate
pedal grasping. *J. Hum. Evol.* **55**, 1096–1101 (2008).

32. Lemelin, P. & Schmitt, D. Origins of grasping and locomotor adaptations in primates: Comparative and
experimental approaches using an opossum model. *PRIMATE Orig. Adapt. Evol.* 329–380 (2007) doi:10.1007/978-
0-387-33507-0_10.

33. Shapiro, L. J. & Young, J. W. Is primate-like quadrupedalism necessary for fine-branch locomotion? A test using
sugar gliders (Petaurus breviceps). *J. Hum. Evol.* **58**, 309–319 (2010).

34. Shapiro, L. J., Young, J. W. & VandeBerg, J. L. Body size and the small branch niche: Using marsupial ontogeny to
model primate locomotor evolution. *J. Hum. Evol.* **68**, 14–31 (2014).

35. Youlatos, D., Moussa, D., Karantanis, N.-E. & Rychlik, L. Locomotion, postures, substrate use, and foot grasping in
the marsupial feathertail glider Acrobates pygmaeus (Diprotodontia: Acrobatidae): Insights into early euprimate
evolution. *J. Hum. Evol.* **123**, 148–159 (2018).

36. Preuschoft, H. What does " arboreal locomotion " mean exactly and what are the relationships between " climbing ",
environment and morphology ? *Z. Morphol. Anthropol.* 171–188 (2002) doi:10.1098/rstb.2011.0050.

37. Antunes, V. Z., Delciellos, A. C. & Vieira, M. V. Postural climbing behaviour of didelphid marsupials: Parallels
with primates. *Oecologia Aust.* **20**, 54–69 (2016).

38. Clemente, C. J. *et al.* Moving in complex environments: A biomechanical analysis of locomotion on inclined and
narrow substrates. *J. Exp. Biol.* **222**, (2019).

39. Birn-Jeffery, A. V. & Higham, T. E. The scaling of uphill and downhill locomotion in legged animals. *Integr. Comp.*
Biol. **54**, 1159–1172 (2014).

40. Youlatos, D. Substrate use and locomotor modes of the Neotropical pygmy squirrel Sciurillus pusillus (E. Geoffroy,
1803) in French Guyana. *Zool. Stud.* **50**, 745–750 (2011).

41. Youlatos, D. & Panyutina, A. A. Habitual bark gleaning by cambodian striped squirrels tamiops rodolphii (rodentia:
Sciuridae) in Cat Tien National Park, South Vietnam. *Mammal Study* **39**, 73–81 (2014).

42. Jenkins, F. A. & McClearn, D. Mechanisms of hind foot reversal in climbing mammals. *J. Morphol.* **182**, 197–219
(1984).

43. Youlatos, D. & Gasc, J. Critical foraging locomotor patterns: Head-first vertical descent in the red howler monkey
(Alouatta seniculus). *Z. Morphol. Anthropol.* **80**, 65–77 (1994).

44. Garber, P. A. & Rehg, J. A. The ecological role of the prehensile tail in white-faced capuchin (Cebus capucinus).
Am. J. Phys. Anthropol. **110**, 325–339 (1999).

45. Lawler, R. R. & Stamps, C. The relationship between tail use and positional behavior in Alouatta palliata. *Primates*
43, 147–152 (2002).

46. Perchalski, B. Headfirst descent behaviors in a comparative sample of strepsirrhine primates. *Am. J. Primatol.* **83**, 1–
16 (2021).

47. Cartmill, M., Lemelin, P. & Schmitt, D. Primate gaits and primate origins. *PRIMATE Orig. Adapt. Evol.* 403–435
(2007) doi:10.1007/978-0-387-33507-0_12.

48. Lammers, A. R. & Stakes, S. A. Kinetics of Symmetrical Versus Asymmetrical In-Phase Gaits During Arboreal
Locomotion. *J. Exp. Zool. Part A Ecol. Integr. Physiol.* 159–171 (2024) doi:10.1002/jez.2878.

49. Schmitt, D. & Lemelin, P. Origins of primate locomotion: Gait mechanics of the woolly opossum. *Am. J. Phys.*
Anthropol. **118**, 231–238 (2002).

50. Hesse, B., Nyakatura, J. A., Fischer, M. S. & Schmidt, M. Adjustments of Limb Mechanics in Cotton-top Tamarins
to Moderate and Steep Support Orientations: Significance for the Understanding of Early Primate Evolution. *J.*
Mamm. Evol. **22**, 435–450 (2015).

51. Nyakatura, J. A., Fischer, M. S. & Schmidt, M. Gait parameter adjustments of cotton-top tamarins (Saguinus
oedipus, Callitrichidae) to locomotion on inclined arboreal substrates. *Am. J. Phys. Anthropol.* **135**, 13–26 (2008).

52. Karantanis, N. E., Rychlik, L., Herrel, A. & Youlatos, D. Comparing the Arboreal Gaits of Muscardinus avellanarius
and Glis glis (Gliridae, Rodentia): A First Quantitative Analysis. *Mammal Study* **42**, 161–172 (2017).

53. Wölfer, J., Aschenbach, T., Michel, J. & Nyakatura, J. A. Mechanics of Arboreal Locomotion in Swinhoe's Striped
Squirrels: A Potential Model for Early Euarchontoglires. *Front. Ecol. Evol.* **9**, (2021).

54. Lammers, A. R., Earls, K. D. & Biknevicius, A. R. Locomotor kinetics and kinematics on inclines and declines in
the gray short-tailed opossum Monodelphis domestica. *J. Exp. Biol.* **209**, 4154–4166 (2006).

55. Vilensky, J. A. & Larson, S. G. Primate Locomotion: Utilization and Control of Symmetrical Gaits. *Annu. Rev.*
Anthropol. **18**, 17–35 (1989).

56. Hildebrand, M. Symmetrical gaits of primates. *Am. J. Phys. Anthropol.* **26**, 119–130 (1967).

57. Hildebrand, M. Analysis of the symmetrical gaits of tetrapods. *Folia Biotheor.* **6**, 9–22 (1966).

58. Shapiro, L. J. & Raichlen, D. A. Lateral sequence walking in infant Papio cynocephalus: Implications for the
evolution of diagonal sequence walking in primates. *Am. J. Phys. Anthropol.* **126**, 205–213 (2005).

59. Cartmill, M., Lemelin, P. & Schmitt, D. Understanding the adaptive value of diagonal-sequence gaits in primates: A
comment on Shapiro and Raichlen, 2005. *Am. J. Phys. Anthropol.* **133**, 822–825 (2007).

60. Hildebrand, M. Analysis of Asymmetrical Gaits. *J. Mammal.* **58**, 131–156 (1977).

61. Dunham, N. T., McNamara, A., Shapiro, L. J., Phelps, T. & Young, J. W. Asymmetrical gait kinematics of free-
ranging callitrichine primates in response to changes in substrate diameter and orientation. *J. Exp. Biol.* **223**, (2020).

62. Chadwell, B. A. & Young, J. W. Angular Momentum and Arboreal Stability in Common Marmosets (Callithrix
jacchus). **576**, 565–576 (2015).

63. Cartmill, M., Lemelin, P. & Schmitt, D. Support polygons and symmetrical gaits in mammals. *Zool. J. Linn. Soc.*
136, 401–420 (2002).

64. Lemelin, P. & Cartmill, M. The effect of substrate size on the locomotion and gait patterns of the kinkajou (Potos
flavus). *J. Exp. Zool. Part A Ecol. Genet. Physiol.* **313 A**, 157–168 (2010).

65. Granatosky, M. C., Schmitt, D. & Hanna, J. Comparison of spatiotemporal gait characteristics between vertical
climbing and horizontal walking in primates. *J. Exp. Biol.* **222**, (2019).

66. Nyakatura, J. A. & Heymann, E. W. Effects of support size and orientation on symmetric gaits in free-ranging
tamarins of Amazonian Peru: implications for the functional significance of primate gait sequence patterns. *J. Hum.*
Evol. **58**, 242–251 (2010).

67. Prost, J. H. & Sussman, R. W. Monkey locomotion on inclined surfaces. *Am. J. Phys. Anthropol.* **31**, 53–58 (1969).

68. Rollinson, J. & Martin, R. D. Comparative aspects of primate locomotion, with special reference to arboreal
Cercopithecines. *Symp. Zool. Soc. London* **48**, 377–427 (1981).

69. Shapiro, L. J., Dunham, N. T., McNamara, A., Young, J. W. & Hieronymus, T. L. Assessing the Determinants of
Platyrrhine Quadrupedal Gait Kinematics in an Ecological and Phylogenetic Framework. *Am. J. Primatol.* **87**,
(2025).

70. Granatosky, M. C., Toussaint, S. L. D., Young, M. W., Panyutina, A. & Youlatos, D. The northern treeshrew
(Scandentia: Tupaiidae: Tupaia belangeri) in the context of primate locomotor evolution: A comprehensive analysis
of gait, positional, and grasping behavior. *J. Exp. Zool. Part A Ecol. Integr. Physiol.* **337**, 645–665 (2022).

71. Cartmill, M. *et al.* The gaits of marsupials and the evolution of diagonal-sequence walking in primates. *Am. J. Phys.*
Anthropol. 1–16 (2019) doi:10.1002/ajpa.23959.

72. Lemelin, P., Schmitt, D. & Cartmill, M. Footfall patterns and interlimb co-ordination in opossums (Family
Didelphidae): Evidence for the evolution of diagonal-sequence walking gaits in primates. *J. Zool.* **260**, 423–429
(2003).

73. Gaschk, J. L., Frère, C. H. & Clemente, C. J. Quantifying koala locomotion strategies: Implications for the evolution
of arborealism in marsupials. *J. Exp. Biol.* **222**, (2019).

74. Karantanis, N. E., Youlatos, D. & Rychlik, L. Diagonal gaits in the feathertail glider Acrobates pygmaeus

(Acrobatidae, Diprotodontia): Insights for the evolution of primate quadrupedalism. *J. Hum. Evol.* **86**, 43–54 (2015).

75.   Lemelin, P. & Jungers, W. L. Body size and scaling of the hands and feet of prosimian primates. *Am. J. Phys.*

Anthropol. **133**, 828–840 (2007).

76.   Boddy, A. M. *et al.* Comparative analysis of encephalization in mammals reveals relaxed constraints on anthropoid

primate and cetacean brain scaling. *J. Evol. Biol.* **25**, 981–994 (2012).

77.   Szalay, F. S. The Beginnings of Primates. *Evolution (N. Y).* **22**, 19 (1968).

78.   Martin, R. D. *Primate origins and evolution: A phylogenetic reconstruction.* (Princeton, NJ: Princeton University

Press., 1990).

79.   Dagosto, M., Gebo, D., Ni, X. & Smith, T. Estimating body size in early primates: The case of Archicebus and

Teilhardina. *J. Hum. Evol.* **115**, 8–19 (2018).

80.   Upham, N. S., Esselstyn, J. A. & Jetz, W. *Inferring the mammal tree: Species-level sets of phylogenies for questions*

in ecology, evolution, and conservation. *PLoS Biology* vol. 17 (2019).

81.   Hughes, J. J., Berv, J. S., Chester, S. G. B., Sargis, E. J. & Field, D. J. Ecological selectivity and the evolution of

mammalian substrate preference across the K–Pg boundary. *Ecol. Evol.* **11**, 14540–14554 (2021).

82.   Dunn, R. H. *et al.* New euprimate postcrania from the early Eocene of Gujarat, India, and the strepsirrhine–

haplorhine divergence. *J. Hum. Evol.* **99**, 25–51 (2016).

83.   Jerison, H. J. Evolution of the brain and intelligence. *Acad. Press. New York* (1973).

84.   Martin, R. D. & Barbour, A. D. Aspects of line-fitting in bivariate allometric analyses. *Folia Primatol.* **53**, 65–81

(1989).

85.   Clavel, J., Merceron, G. & Escarguel, G. Missing data estimation in morphometrics: How much is too much? *Syst.*

Biol. **63**, 203–218 (2014).

86.   Cooper, N., Thomas, G. H., Venditti, C., Meade, A. & Freckleton, R. P. A cautionary note on the use of Ornstein

Uhlenbeck models in macroevolutionary studies. *Biol. J. Linn. Soc.* **118**, 64–77 (2016).

87.   Dunham, N. T. *et al.* Effects of substrate and phylogeny on quadrupedal gait in free-ranging platyrrhines. *Am. J.*

Phys. Anthropol. **170**, 565–578 (2019).

88.   Janisch, J. *et al.* Ecological and phylogenetic influences on limb joint kinematics in wild primates. *Zool. J. Linn.*

Soc. **202**, (2024).

89.   Wimberly, A. N., Slater, G. J. & Granatosky, M. C. Evolutionary history of quadrupedal walking gaits shows

mammalian release from locomotor constraint. *Proc. R. Soc. B Biol. Sci.* **288**, (2021).

90.   Toussaint, S. Towards primate origins: hands and feet in interdisciplinary perspective. (2018).

91.   Hanna, J. B. & Schmitt, D. Locomotor energetics in primates: Gait mechanics and their relationship to the energetics

of vertical and horizontal locomotion. *Am. J. Phys. Anthropol.* **145**, 43–54 (2011).

92.   Preuschoft, H. What does 'arboreal locomotion' mean exactly and what are the relationships between 'climbing',

environment and morphology? *Z. Morphol. Anthropol.* **83**, 171–188 (2002).

93.   Preuschoft, H., Witte, H. & Fischer, M. Locomotion in nocturnal prosimians. in *Creatures of the Dark* (eds.

Alterman, L., Doyle, G. A. & Izard, M. K.) 453–472 (Springer US, 1995). doi:10.1007/978-1-4757-2405-9_27.

94.   Granatosky, M. C., Tripp, C. H., Fabre, A. C. & Schmitt, D. Patterns of quadrupedal locomotion in a vertical

clinging and leaping primate (Propithecus coquereli) with implications for understanding the functional demands of

primate quadrupedal locomotion. *Am. J. Phys. Anthropol.* **160**, 644–652 (2016).

95.   Shapiro, L. J. & Raichlen, D. A. A response to Cartmill et al.: Primate gaits and arboreal stability. *Am. J. Phys.*

Anthropol. **133**, 825–827 (2007).

96.   Lammers, A. R. & Zurcher, U. Stability During Arboreal Locomotion. *Theor. Biomech.* 319–334 (2011)

doi:10.5772/23751.

97.   Granatosky, M. C. A Review of locomotor diversity in mammals with analyses exploring the influence of substrate

use, body mass and intermembral index in primates. *J. Zool.* **306**, 207–216 (2018).

98.   Young, J. W., Russo, G. A., Fellmann, C. D., Thatikunta, M. A. & Chadwell, B. A. Tail function during arboreal
1028   quadrupedalism in squirrel monkeys (Saimiri boliviensis) and tamarins (Saguinus oedipus). *J. Exp. Zool. Part A*
1029   *Ecol. Genet. Physiol.* **323**, 556–566 (2015).

1030   99.   Stevens, N. J. The effect of branch diameter on primate gait sequence pattern. *Am. J. Primatol.* **70**, 356–362 (2008).

1031   100.   Larson, S. G. & Stern, J. T. Maintenance of above-branch balance during primate arboreal quadrupedalism:
1032   Coordinated use of forearm rotators and tail motion. *Am. J. Phys. Anthropol.* **129**, 71–81 (2006).

1033   101.   Fukushima, T. *et al.* Inertial Tail Effects during Righting of Squirrels in Unexpected Falls: From Behavior to
1034   Robotics. *Integr. Comp. Biol.* **61**, 589–602 (2021).

1035   102.   Thorpe, S. K. S. & Crompton, R. H. Orangutan positional behavior and the nature of arboreal locomotion in
1036   Hominoidea. *Am. J. Phys. Anthropol.* **131**, 384–401 (2006).

1037   103.   Fleagle, J. G. *Primate Adaptation and Evolution*. (Elsevier, 2013). doi:10.1016/C2009-0-01979-5.

1038   104.   Young, J. W. & Chadwell, B. A. Not all fine-branch locomotion is equal: Grasping morphology determines
1039   locomotor performance on narrow supports. *J. Hum. Evol.* **142**, 102767 (2020).

1040   105.   Harrington, A. R., Silcox, M. T., Yapuncich, G. S., Boyer, D. M. & Bloch, J. I. First virtual endocasts of adapiform
1041   primates. *J. Hum. Evol.* **99**, 52–78 (2016).

1042   106.   Orliac, M. J., Ladeveze, S., Gingerich, P. D., Lebrun, R. & Smith, T. Endocranial morphology of Palaeocene
1043   Plesiadapis tricuspidens and evolution of the early primate brain. *Proc. R. Soc. B Biol. Sci.* **281**, 20132792–
1044   20132792 (2014).

1045   107.   Bertrand, O. C. & Silcox, M. T. First virtual endocasts of a fossil rodent: Ischyromys typus (Ischyromyidae,
1046   Oligocene) and brain evolution in rodents. *J. Vertebr. Paleontol.* **36**, (2016).

1047   108.   Janecka, J. E. *et al.* Molecular and Genomic Data Identify the Closest Living Relative of Primates. *Science* **318**,
1048   792–794 (2007).

1049   109.   Chester, S. G. B., Williamson, T. E., Bloch, J. I., Silcox, M. T. & Sargis, E. J. Oldest skeleton of a plesiadapiform
1050   provides additional evidence for an exclusively arboreal radiation of stem primates in the Palaeocene. *R. Soc. Open*
1051   *Sci.* **4**, 170329 (2017).

1052   110.   Silcox, M. T. & López-Torres, S. Major Questions in the Study of Primate Origins. *Annu. Rev. Earth Planet. Sci.* **45**,
1053   113–137 (2017).

1054   111.   Silcox, M. T. & Gunnell, G. F. *Plesiadapiformes. Evolution of Tertiary Mammals of North America* (2008).
1055   doi:10.1017/CBO9780511541438.015.

1056   112.   Silcox, M. T., Bloch, J. I., Boyer, D. M., Chester, S. G. B. B. & López-Torres, S. The evolutionary radiation of
1057   plesiadapiforms. *Evol. Anthropol.* **26**, 74–94 (2017).

1058   113.   Boyer, D. M. *New cranial and postcranial remains of late Paleocene Plesiadapidae ("Plesiadapiformes,"*
1059   *Mammalia) from North America and Europe: description and evolutionary implications*. Stony Brook, NY: Ph. D.
1060   Dissertation, Stony Brook University (2009).

1061   114.   Youlatos, D. & Godinot, M. Locomotor adaptations of Plesiadapis tricuspidens and Plesiadapis n. sp. (Mammalia,
1062   Plesiadapiformes) as reflected in selected parts of the postcranium. *Anthropologica* **82**, 103–118 (2004).

1063   115.   Ni, X. *et al.* The oldest known primate skeleton and early haplorhine evolution. *Nature* **498**, 60–64 (2013).

1064   116.   Marigó, J., Verrière, N. & Godinot, M. Systematic and locomotor diversification of the Adapis group (Primates,
1065   Adapiformes) in the late Eocene of the Quercy (Southwest France), revealed by humeral remains. *J. Hum. Evol.* **126**,
1066   71–90 (2019).

1067   117.   Franzen, J. L. *et al.* Complete primate skeleton from the middle Eocene of Messel in Germany: Morphology and
1068   paleobiology. *PLoS One* **4**, (2009).

1069   118.   Williams, B. A., Kay, R. F. & Kirk, E. C. New perspectives on anthropoid origins. *Proc. Natl. Acad. Sci.* **107**, 4797–
1070   4804 (2010).

119.   Pozzi, L. *et al.* Primate phylogenetic relationships and divergence dates inferred from complete mitochondrial

genomes. *Mol. Phylogenet. Evol.* **75**, 165–183 (2014).

120. Del Rio, C., Haevermans, T. & De Franceschi, D. First record of an Icacinaceae Miers fossil flower from le Quesnoy
(Ypresian, France) amber. *Sci. Rep.* **7**, 1–8 (2017).

121. Ravosa, M. & Dagosto, M. *PRIMATE ORIGINS: Adaptations and Evolution*. (Springer US, 2007). doi:10.1007/978-
0-387-33507-0.

122. Sussman, R. W. Primate origins and the evolution of angiosperm. *Am. J. Primatol.* **23**, 209–223 (1991).

123. Peckre, L. R. *et al.* Food mobility and the evolution of grasping behaviour: A case study in strepsirrhine primates. *J.*
Exp. Biol. **222**, (2019).

124. Cartmill, M. Arboreal adaptations and the origin of the order Primates. in *The Functional and Evolutionary Biology*
of Primates (ed. Tuttle, R.) 97–122 (1972).

125. Patel, B. A. *et al.* Distinct functional roles of primate grasping hands and feet during arboreal quadrupedal
locomotion. *J. Hum. Evol.* **88**, 79–84 (2015).

126. Pouydebat, E., Fragaszy, D. & Kivell, T. L. GRASPING IN PRIMATES: TO FEED, TO MOVE AND HUMAN
SPECIFICITIES. *Growth (Lakeland)* **26**, 1–24 (2006).

127. Boyer, D. M., Yapuncich, G. S., Chester, S. G. B., Bloch, J. I. & Godinot, M. Hands of early primates. *Am. J. Phys.*
Anthropol. **152**, 33–78 (2013).

128. ASAB & ABS. Guidelines for the treatment of animals in behavioural research and teaching. *Anim. Behav.* **159**, I–
XI (2020).

129. Altmann, J. Observational Study of Behavior : Sampling Methods. *Behaviour* **49**, 227–267 (1974).

130. Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A Resource for Timelines, Timetrees, and
Divergence Times. *Mol. Biol. Evol.* **34**, 1812–1819 (2017).

131. Blomberg, S. P., Garland, T. & Ives, A. R. Testing for phylogenetic signal in comparative data: behavioral traits are
more labile. *Evolution (N. Y).* **57**, 717–745 (2003).

132. Adams, D. C. A generalized K statistic for estimating phylogenetic signal from shape and other high-dimensional
multivariate data. *Syst. Biol.* **63**, 685–97 (2014).

133. Clavel, J., Escarguel, G. & Merceron, G. mvMORPH: An R package for fitting multivariate evolutionary models to
morphometric data. *Methods Ecol. Evol.* **6**, 1311–1319 (2015).

1099

1100 ## Fundings

1101 Research funding was from the Fyssen Foundation and the Alexander von Humboldt
1102 Foundation postdoctoral grants (to SLDT) and from the German Research Foundation grant
1103 DFG NY63 2/1 (to JAN). Travel funding for behavioral data collection in Poland and The
1104 Russian Federation was from Erasmus International Mobility funds (to DY).

1105

Author contributions

Conceptualization: SLDT

Methodology: SLDT, DY, JAN

Investigation: SLDT, DY, JAN

Visualization: SLDT

Funding acquisition: SLDT, DY, JAN

Writing—original draft: SLDT

Writing—review & editing: SLDT, DY, JAN

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

All aggregated data needed to evaluate the conclusions of this study are included in this paper and its supplementary materials. Raw video data are available upon reasonable request which should be addressed to SLDT and DY. The R scripts used in this study are also available upon request and should be addressed to SLDT.