Hyper-dominant species drive unexpected functional shifts in North American birds

Heng-Xing Zou[1,2*], Adam C. Smith[3,4], Yiluan Song[1,2,5], Kai Zhu[1,2], and Brian C. Weeks[2]

1. Institute for Global Change Biology, University of Michigan, Ann Arbor, MI 48109

2. School for Environment and Sustainability, University of Michigan, Ann Arbor, MI 48109

3. Department of Biology, Carleton University, Ottawa K1S 5B6, ON, Canada

4. Canadian Wildlife Service, National Wildlife Research Centre, Environment and Climate

Change Canada, Ottawa K1S 5R2, ON, Canada

5. Michigan Institute for Data and AI in Society, University of Michigan, Ann Arbor, MI 48109

* Corresponding author. Email: hxzou.ecology@gmail.com

Number of words in Abstract: 189

Number of words in the main text: 3303

Number of words in Methods: 3608

Number of references: 70

Number of figures in the main text: 3

Abstract

Global environmental changes have led to widespread intraspecific shifts in functional

traits across large numbers of widely distributed taxa. However, how within-species

functional shifts predict community-level changes is unknown. We summarized seven key

traits that are hypothesized to shift adaptively under environmental changes using local

community-weighted means across North American bird communities. We find that

communities shifted toward larger body mass and smaller appendage sizes over time, the

opposite of widely observed within-species temporal changes in North American birds.

These unexpected patterns arise from changes in the relative abundance of a very small

number of hyper-dominant species, with 23 out of 470 (5%) species explaining the

community-level trajectories in functional traits. These species are known to be heavily

impacted—both positively and negatively—by human activities, and the idiosyncratic

drivers of population sizes result in weak associations between community-level

functional shifts and major bioclimatic variables but strong associations with human-

induced land use changes. Our results illustrate the complexity in scaling from within-

species responses to global change to community-level responses and highlight the

importance of species-specific responses to the environment beyond climate when

predicting functional and taxonomic diversity under global change.

Main Text

Rapid climate change has driven widespread shifts in species' abundances and

distributions (1–3), as well as changes in key functional or life history traits (4, 5). Both

shifts in species abundances across space and functional traits through time are expected

to alter the taxonomic and functional diversity of ecological communities (6, 7). Global

change drivers are often general, such as broadly consistent increases in temperature and

widespread human-mediated habitat degradation (8, 9). Many of these changes could lead

to similarly consistent responses among taxa; for example, as the world warms,

thermoregulatory demands are likely to shift similarly across broad groups of species,

driving parallel adaptations to facilitate heat dissipation (10). However, impacts of other

global change factors, such as land use change, could be more idiosyncratic among

species, potentially resulting in inconsistent responses (e.g., variable shifts in range (11),

phenology (12), and morphology (13) across species). These inconsistencies may pose

remarkable challenges for predicting how ecological communities will respond to global

change from within-species responses.

Birds are a model system for relating species- and community-level responses to

global change as they have comprehensive distributional (14), phylogenetic (15),

behavioral (16), and functional trait (17) data, as well as long-term time series data on

shifts in range, phenology, and morphology. There is reason to expect that responses to

rapid environmental changes are general and directional among avian communities.

Within species, shifts in key functional traits have been relatively uniform: broadly, species

have become smaller (4, 18–20) and their appendages, such as beaks, tarsi, and wings,

have become longer (13). These shifts are often interpreted within the context of

thermoregulation or adaptation to more fragmented habitats and urbanization (10, 21, 22).

There is growing evidence that some within-species responses to climate change (e.g.,

range shifts) have resulted in community-level changes. Bird species have generally

shifted poleward and up in elevation, likely in response to the warming climate (2, 12, 23),

and these distributional changes are thought to have contributed to consistent local

changes in functional richness and evenness across large scales (7, 24, 25).

Should within-species changes in morphology represent consistent selection favoring

particular phenotypes (e.g., smaller body size) under changing climate and environments,

we may expect that species with that phenotype should be increasing in relative

abundance, shifting the community-weighted mean (CWM) of that trait in parallel with the

intra-specific trends. Alternatively, scaling up within-species trends to the community level

may be more complex. Idiosyncratic responses of individual species to the broad array of

environmental changes that are occurring contemporaneously may further lead to

inconsistencies between shared within-species responses to a single driver (e.g., climate

change) and community-level functional shifts (20). Relatedly, within-species changes

may be adaptive, but this micro-evolutionary response may not predict variation in fitness

among species along the same trait axis. Finally, while many within-species changes have

been associated with environmental change, it is unclear if these changes are adaptive or

the result of plasticity, in which case they may be less likely to predict shifts in fitness

among species. Evaluating the generality of these functional responses across the scales

is important in predicting shifts in bird community structure and ecosystem function under

rapid environmental changes.

Unexpected community-level shifts in functional traits

To test if community-level functional shifts align with previous hypotheses based on

within-species changes, we characterized changes in functional traits of local

communities based on abundance data for 470 species spanning half a century of

sampling in North America (North American Breeding Bird Survey, NABBS; Fig. 1A; (26)). To

obtain robust estimates of local community composition, we aggregated abundance data

within one-degree grids, aligning with the design of NABBS to have at least one route per

grid with suitable access roads (see Methods). We calculated community-weighted means

(CWMs) of seven key morphological and demographic traits: beak size, wing size, body

mass, relative beak length, relative wing length, clutch size, and generation length. We

calculated corrected beak and wing lengths (Methods); like relative beak/wing lengths,

these measurements correct for the associations between beak/wing length and body

mass (27). Similarly, we also corrected clutch size and generation lengths by body mass

(Methods). CWMs are the average of functional trait values among species within a

community, weighted by species abundances (28) and provide more mechanistic

interpretations of functional shifts at the community level, more direct comparisons with

within-species shifts, and are less biased by the presence of rare species than ordination-

based approaches (29). To better compare with within-species expectations of functional

shifts, we examined how shifts in CWMs are associated with time and environmental

variables. We considered major climatic attributes and human-induced land use changes,

which have greatly affected ecological communities (9, 30, 31).

We find that community-level shifts in key morphological traits are often in opposite

directions from within-species expectations based on adaptations to changing climates.

CWM of body mass increases with temperature of the warmest quarter (mean = 0.168,

standard error = 0.00893, p < 0.0001; Fig. 1C), contrary to within-species studies that have

found reductions in body size as the climate has warmed over time (4, 19). The CWMs of

beak and wing sizes (the first PC axes from phylogenetic PCA of four beak measurements

and three wing measurements, respectively; see Methods) also positively associate with

temperature (beak: 0.0354 ± 0.00177; wing: 0.0295 ± 0.00205; both p < 0.0001), but

relative beak and wing lengths both show strongly negative associations with temperature

(beak: -0.177 ± 0.00868; wing: -0.326 ± 0.0117; both p < 0.0001). These results contrast

with within-species trends toward longer relative appendage size (4, 19, 20) that are often

attributed to selection towards increased effective surface area and heat dissipation

efficiency under warming (10). The CWM of clutch size positively associates with

temperature (0.290 ± 0.00980, p < 0.0001) but negatively associates with temperature

seasonality (-0.112 ± 0.00480, p < 0.0001; Fig. 1C). Both are contrary to within-species

patterns: higher ambient temperature is associated with smaller clutch sizes often

attributed to increased potential for hatching failure, and higher intra-annual fluctuations

in temperature may select for larger clutch sizes as a bet-hedging strategy (32–34).

Generation length represents the average age of parents in any population of a species (35,

36) and higher intra-annual fluctuations in temperature should select for earlier

reproduction (35, 37). In contrast, we observed a positive association with temperature

seasonality (0.0339 ± 0.00393, $p < 0.0001$). Because temperature and precipitation

seasonality only directly affect year-round residents (35), we re-analyzed the associations

between climate seasonality and demographic traits separately for species that undertake

long-distance seasonal migrations and those that do not according to (38). The signs of all

associations are consistent among migrants and residents, but the positive association

between the CWM of generation length and temperature seasonality is non-significant (fig.

S1-S2).

Many studies have also suggested human-induced land use changes should favor

species with certain traits, especially in urban areas where habitats are most modified (21,

22). We found that these land use changes often have stronger associations with

community-level functional traits, but some of them are again contrary to expectations.

The CWM of body mass is positively associated with human-modified forests and drylands

(cultured land), and villages and urban areas (settlements; Fig. 1C; table S1). These results

are different from the expectation that smaller body size should be favored in more human-

dominated landscapes such as urban areas due to lower resource and habitat

requirements (21, 22). The CWM of relative wing length is positively associated with

cultured land (0.103 ± 0.00272, $p = 0.000196$) but not agricultural land and dense

settlements, again different from the expectation that urban-adapted species should have

larger wing-to-body ratios to assist with dispersal among fragmented habitats (21, 22). All

three human-induced land use changes are negatively associated with the CWM of clutch

size (agriculture: -0.272 ± 0.0277, cultured: -0.244 ± 0.0259, settlements: -0.128 ± 0.0184;

all $p < 0.0001$), contrary to the expectations that anthropogenic environmental stress

should favor species with larger clutch sizes (21, 22). On the other hand, the CWM of

generation length is positively associated with all three human-induced land use changes

(agriculture: 0.373 ± 0.0239, cultured: 0.331 ± 0.0223, settlements: 0.395 ± 0.0157; all $p <$

0.0001). Because maximum longevity has a strongly positive correlation with generation

length (35, 36), our result is therefore consistent to the expectation from previous studies

that longer-lived species are favored in urban areas (21, 22).

We found that across North America, community-level shifts in morphological and

demographic traits often contrast with species- and population-level expectations. We did

not consider how species traits may change across space or time, which is unlikely to alter

results because the magnitudes of within-species trait changes are much smaller than

typical among-species variation in these traits (39). In addition to advancing our

understanding of how communities have shifted through time, our findings shed new light

on the interpretation of well-documented functional shifts within species. If these changes

in morphological and demographic traits are adaptive to environmental changes, then it is

intuitive to expect that species with traits in these more adaptive directions would increase

in abundance and lead to shifts in the CWMs of the trait that are aligned with intra-specific

trends. However, the shifts we find at the community level are often contrary to the shifts

that have been broadly observed within species, which we interpret in three possible ways.

First, observed within-species functional shifts may not be adaptive, or may not have

meaningful effects on fitness (39–41). Indeed, although some morphological responses

are consistent over time, whether they are indeed adaptive to environmental changes

remains unclear (39, 40). With more studies finding evidence both supporting and

opposing longstanding hypotheses based on the adaptation of climate and environmental

change (e.g., warming and urbanization (10, 22)), the lack of consistency on a community

level in our results may further indicate that within-species morphological changes may

not be responding to the same single driver in the environment. Second, if observed

intraspecific functional shifts are indeed adaptive, then our results indicate that

microevolutionary processes may not necessarily predict relative fitness differences

among species (42). Third, factors that drive adaptive intraspecific changes in functional

traits (e.g., climate change) may have weaker effects on relative abundance than other

drivers that do not drive intraspecific trait evolution but do determine differential sensitivity

to more significant population-level pressures (e.g., human-induced land use change;

(43)). As such, inconsistencies between intraspecific trait changes and shifts at the

community level demonstrate the complexities involved in scaling up from species to

community level dynamics in the context of multifaceted global environmental change.

Nearly universal functional shifts across space

Continental-scale analyses of functional shifts may mask local and regional

environmental factors. We explored the generality of functional shifts across local

communities in North America by analyzing the temporal trajectory of each local

community within a functional space constructed by a PCA that reduced the CWMs of the

seven traits into two major axes (Fig. 2A). The first PC axis (which explains 43.9% of the

variance) mostly represents variation in traits related to absolute body size (beak and wing

sizes and body mass); the second PC axis (which explains 16.9% of the variance) mostly

reflects variation in clutch size, generation length, and relative wing length (Fig. 2A). We

focus on the locations of the community in functional trait space at the start and end years

(1970 and 2021), summarized by their positions in quadrants, and the overall direction

(angle) of the shift for each community, summarized by the quadrant it points to (Fig. 2A).

In 1970, most local communities in central North America were characterized by larger

body mass, larger absolute appendage sizes, larger clutch size, and shorter generation

lengths (Fig. 2B; fig. S3). In 2021, many local communities, especially at lower latitudes,

were characterized by smaller clutch sizes and longer generation lengths. Examining the

directions of these shifts from 1970 to 2021, we observe an east-west divide, with local

communities in the Eastern Forests and eastern parts of the Prairie regions shifting

towards smaller clutch sizes and longer relative beak and wing lengths, whereas many

other major avifaunal biomes in the west and north have shifted towards larger body mass

and clutch sizes (Fig. 2B; fig. S4; see fig. S5 for a map of avifaunal biomes). Overall, these

local-level functional shifts are broadly consistent with the results from the continent-level

trends, that CWM of body mass have generally increased with time and temperature, and

that clutch size and generation lengths have generally decreased with human-induced

land use changes (Fig. 1C). The highly consistent community-level functional shifts that

contrast with within-species expectations highlight that even when there are nearly

universal changes in the environment over time that can drive consistent intraspecific

functional trends among species (e.g. consistent size declines associated with increasing

temperatures; (4)), these conserved trends may not scale simply to analogous functional

shifts of local communities.

Functional shifts driven by a few hyper-dominant species

As a consequence of global environmental changes, shifts in the composition of a

community may emerge due to consistent changes across many of the species in a

community (e.g. consistent declines in the abundances of smaller species, resulting in an

increase in the CWM of mass through time) or extreme shifts in the abundance of a

relatively small number of species that either become hyper-dominant while others are

extirpated or decline rapidly transitioning away from being a hyper-dominant species (44).

To better understand the mechanisms underlying the unexpected functional shifts on the

community level, we quantified the contributions of each species to the temporal trends of

CWMs within each local community, allowing us to determine the magnitude of the

contributions of individual species to community-level functional shifts. For each trait, we

ranked the species by the number of local communities in which they contribute greatly to

the temporal trend (i.e., they are within the top 10% of species driving the community trend

in that local community). We found 4.89% of species (23 out of 470) had overwhelmingly

important contributions to functional shifts across North America (Fig. 3; fig. S6-S12). To

illustrate the importance of these top contributors to community-level functional shifts, we

calculated CWMs for two subcommunities: (1) only the "top contributors", and (2) the rest

of the species (hereafter the "standard contributors"). We then repeated all the analyses of

the temporal trends and environmental associations with CWMs and compared the results

of each subcommunity with the results derived using the full set of species (the "complete

communities").

In general, many unexpected associations between shifts in CWMs of the complete

communities and bioclimatic variables can be explained by the idiosyncratic responses of

top contributors to human-induced land use changes. Top contributors drive the

unexpected positive association between body mass and temperature (0.269 ± 0.0111; $p <$

0.0001) and negative associations between relative beak and wing lengths and

temperature (beak: -0.243 ± 0.00827; wing: -0.387 ± 0.0106; both $p < 0.0001$; Fig. 1C).

These unexpected morphological shifts can be explained by a few large-bodied species

that are increasingly abundant in urban, suburban, and inhabited wildlands despite

warming temperature, such as Canada Goose (*Branta canadensis*) and Wild Turkey

(*Meleagris gallopavo*; Fig. 3), and species with relatively longer beaks and wings that are

declining over time, such as Western Meadowlark (*Sturnella neglecta*), Horned Lark

(*Eremophila alpestris*), and Barn Swallow (*Hirundo rustica*). On the other hand, relative

beak and wing lengths of the top contributors are strongly positively associated with

proportions of settlements (beak: 0.287 ± 0.0165; wing: 0.0836 ± 0.0187; both $p < 0.0001$),

whereas these associations are strongly negative for the standard contributors (beak: -

0.278 ± 0.0174, $p < 0.0001$; wing: -0.0644 ± 0.0184, $p = 0.000593$; Fig. 1C). The positive

association for relative wing sizes may arise from the increase of long-winged songbirds,

such as American Robin (*Turdus migratorius*) and Red-eyed Vireo (*Vireo olivaceus*) in urban

areas, while the positive association for relative beak sizes may arise from the decrease of

small-beaked species such as House Sparrow (*Passer domesticus*; Fig. 3).

The unexpected associations between CWMs of demographic traits and climatic

variables are again driven by the top contributors, as indicated by their responses to land

use changes. One of the most important contributors to the CWM of clutch size is Northern

Bobwhite (*Colinus virginianus*; Fig. 3), which has a large clutch size and has been declining

precipitously due to habitat degradation (45), corresponding to stronger negative

associations with agriculture and settlements of top contributors (agriculture: -0.305 $\pm$

0.0297; settlements: -0.257 $\pm$ 0.0192; both p < 0.0001; Fig. 1C). The strongly positive

associations between CWMs of generation length and agriculture and cultured lands are

also driven by the top contributors (agriculture: 0.394 $\pm$ 0.0240; cultured land: 0.267 $\pm$

0.0225; both p < 0.0001; Fig.1C), such as increasing abundances of Canada Goose and

Northern Cardinal (*Cardinalis cardinalis*), both of which have above-average generation

lengths (Fig. 3). Overall, our analyses provide strong evidence that changes in the

abundance of a few hyper-dominant species have been responsible for driving the

functional shifts in North American avian communities. Contrary to expectations that

climatic conditions drive consistent morphological changes among species, these

unexpected functional shifts on the community level are driven by the highly idiosyncratic

responses of individual species as they benefit or perish because of human-induced

changes in land cover and habitats (45–48).

At the continental scale, most of the unexpected functional shifts and their strong

associations with human-induced land use changes can be explained by the top

contributors, which are often stronger than, or even opposite to, the associations of CWMs

calculated with the standard contributors. To test the generality of this result among local

communities, we repeated the analyses on the temporal trajectories for the top

contributors and standard contributors. In 1970, subcommunities composed of the top

contributors were mostly characterized by having larger body mass, appendage sizes, and

longer generation lengths (Fig. 2A-B). In 2021, subcommunities with top contributors have

all shifted to smaller clutch sizes and longer generation lengths (Fig. 2B; fig. S3). Strikingly,

spatial patterns of overall directions of functional shifts from 1970 to 2021 are nearly

identical between complete communities and top contributors, indicating that the top

contributors have driven most of the observed functional shifts of complete communities

(Fig. 2B). In contrast, there were no widespread shifts in functional positions within the

standard contributors between 1970 and 2021 (Fig. 2B; fig. S3). The uniform functional

shifts of top contributors are further illustrated by the temporal decrease in spatial

heterogeneity of functional positions (-0.000906 ± 0.000333; $p = 0.00894$), whereas spatial

heterogeneity has increased for complete communities and standard contributors

(complete: 0.00289 ± 0.000515, $p < 0.0001$; standard: 0.000605 ± 0.000267, $p = 0.0279$;

Fig. 2C; table S2). Overall, our observations emphasize the hyper-dominance of a few

species in driving nearly universal changes in CWMs of functional traits (49, 50), indicating

that environmental drivers may not induce shared responses across all species within

communities. Together, our results highlight the importance of considering the role of a

few key contributors in designing conservation and management strategies. Instead of the

entire community, future studies should also consider analyzing species contributions in

subcommunities with specific ecological roles, such as predators, herbivores, or

scavengers, to better understand the effect of global change on biotic interactions and

ecosystem functioning.

Conclusion

Rapid changes in global environments are driving within-species responses and are

reshaping ecological communities, but are these community-level shifts emergent

outcomes of general within-species processes? In North American avian communities, we

find significant shifts in the functional traits of communities that are not driven by

consistent responses among species, or by obvious climatic drivers, but instead are the

outcome of the idiosyncratic rise and fall of a few highly impactful species in response to

land use changes. These widespread, abundant, and often functionally distinctive species

have undergone strong changes in relative abundance, leading to local and continental

functional shifts that are not aligned with expectations derived from within-species trends.

Our evidence highlights the challenges in predicting both within-species and community-

level changes in functional traits from single environmental factors. On the other hand, our

approach provides a direction in predicting community composition by focusing on the

abundance and fitness of key species that contribute greatly to community structure. By

analyzing summary metrics such as community-weighted means of traits essential to

ecosystem functioning, we can further project the broader impact of species redistribution

and identify their key contributors. Integration of functional ecology across scales will be

essential to more accurately predict biotic responses to rapid global change.

Acknowledgements

We thank Casey Youngflesh for comments on the manuscript, and members of the Weeks and Zhu labs and IGCB Postdoc Fellows for their feedback on the project. We thank Olivia Stein for creating the illustrations used in Figure 1 and 3. We thank the numerous volunteers who collected data for the North American Breeding Bird Survey. H.-X.Z was supported by the Postdoctoral Fellowship from the Institute for Global Change Biology at the University of Michigan. Y.S. was supported by the Eric and Wendy Schmidt AI in Science Postdoctoral Fellowship, a Schmidt Sciences program. K.Z. was supported by the National Science Foundation award 2306198 and USDA McIntire-Stennis Capacity Grant award 25-PAF01509. B.C.W. was supported by the David and Lucile Packard Foundation. This research was supported in part through computational resources and services provided by the Advanced Research Computing at the University of Michigan, Ann Arbor.

Authors contributions

H.X.-Z. conceived the idea and designed the study with K.Z. and B.C.W.; H.X.-Z. curated and analyzed data with supervision from K.Z. and B.C.W. A.C.S. assisted with estimating and adjusting bird abundance; Y.S. assisted with analyzing species contributions. H.X.-Z. wrote the first draft, and all authors contributed to subsequent review and editing.

Data and materials availability

340   All primary data are publicly available, and all generated data will be permanently

341   archived upon acceptance. Code is available on GitHub:

342   https://github.com/hengxingzou/Zouetal2025bioRXiv and will be archived upon

343   acceptance.

Methods

Species abundances and trends

We estimated species relative abundance based on data from the North American

Breeding Bird Survey (NABBS), a long-term survey program started in 1966 (26). We then

transformed estimates of relative abundance (relative number of individuals of a given

species, compared to other times and locations; not to be confused with relative

proportions of species abundance in a community) into estimates of abundance (total

number of individual birds, comparable across species, locations, and times) following the

methods used in previous broad-scale studies of North American bird populations (38).

NABBS is a structured monitoring program that has tracked changes in bird communities

across much of North America, since the late 1960s. The NABBS supplies annual

estimates of population change across most of Canada and the United States for almost

all terrestrial bird species (with exceptions that include some waterbirds and nocturnal

species; (26)). NABBS surveys are conducted each breeding season (May, June, or July)

along fixed routes of 40 km at 50 evenly spaced, independent stops. At each stop, the

observer counts all identifiable birds for 3 minutes within a 400 m radius. The routes are

established following secondary roads, in a random design, stratified within grid-cells of

one-degree latitude and longitude. NABBS routes cover a wide range of the United States

and Canada, with at least one route in most grids with suitable roads.

To estimate species relative abundances, we downloaded the 2022 release that

contains survey data from 1966 to 2021 from the U.S. Geological Survey website. We

consider grid cells of one degree to be "local" communities; this scale represents the

design stratification of the NABBS and is small enough that we do not expect significant

variation of bioclimatic conditions within cells and large enough to contain enough survey

data to estimate bird relative abundances (at least one NABBS route). To estimate relative

abundances, we fit hierarchical Bayesian models with a spatial intrinsic conditional

autoregressive structure using the R package *bbsBayes2* (51). To summarize, the model

estimates relative abundance indices from NABBS counts on each route and year with a

negative binomial distribution and weakly informative priors, and the model fitting of a

species from shared information among all sites ("spatial strata"; in our case, one-by-one-

degree grids). In addition, the model accounts for the spatial connection among strata by

constructing a network graph, in which estimates in each stratum draw information from

neighbors to estimate spatial variance and to assist in the model fitting (a "spatial

smoothing" among neighbors). This spatial model not only allows for the estimation of

relative abundances on a finer spatial scale but also outperforms the nonspatial

counterpart in predicting population trajectories over time in poorly sampled locations

(52). After accounting for spatial and temporal autocorrelation and variability in

observation effort, the output of the model is a continuous year series of relative

abundance estimates within the entire historical and current spatial range of each species.

See (52) for the model structure and an expanded technical description of the model

structure.

Prior to model fitting, we consolidated all records to the species level, pooling all

subspecies or morphotypes and removing observations of unidentified species. The Great

White Heron was combined with the Great Blue Heron (*Ardea herodias*), and the former

Cordilleran, Pacific-slope, and unidentified Western Flycatchers were all pooled into

Western Flycatcher (*Empidonax difficilis*). We excluded species with fewer than 50 records

in the whole NABBS dataset (across all routes and years). We also excluded the White-

crowned Pigeon (*Patagioenas leucocephala*), which was only recorded in one grid-cell. To

constrain our analyses to high-quality data, we limited the analysis to data from 1970 to

2021, removed regions outside of the continental United States and Canada (south of

25°N, north of 55°N, and west of 135°W), and excluded local communities with fewer

than 10 species in all years. Finally, we excluded nocturnal and pelagic species, as

identified by the EltonTraits 1.0 dataset (16), because the NABBS protocol makes it poorly

suited for estimating relative abundances for these groups. In total, we conducted

analyses on 470 species in 1261 grids (Fig. 1A). We matched species taxonomy in NABBS

and functional trait datasets (see below) to the 2023 version of the Clements taxonomy

(15).

We used a spatially explicit, "first difference" model to estimate time series of annual

relative abundance (i.e., the species' local population trajectory) from the NABBS data for

each of these 470 species. This model has been commonly used to estimate population

status and trends from the NABBS data (52, 53). It estimates a species' population

trajectory in each grid as a series of annual differences from the previous year, sharing

information on the estimated annual differences in a spatially explicit way (i.e., among

neighboring grids using an intrinsic conditional autoregressive structure). However, we do

not formally compare model performance to determine the best model structure for each

species because: (1) model comparison via leave-one-out cross-validation requires strong

assumptions that individual observations are conditionally independent, which is unlikely

with NABBS data; (2) different models may fit better for different species, making the

pooling of all species relative abundances within a local community difficult; (3) previous

work comparing alternative time series models with the NABBS data has shown that the

estimated population trajectories are very similar among models (51, 52).

To estimate species relative abundances, we first fit the Bayesian models of all

selected species with 1000 warm-up and 1000 sampling iterations, then obtained the

posterior median of relative abundance indices in each year using the *bbsBayes2* R-

package. The large spatial models for each species contain thousands to tens of

thousands of parameters; we therefore used a less conservative convergence criterion by

considering models as converged if no parameters had Rhats > 1.1 as converged. For 31

out of 470 species for which not all the parameters converged, we reran the models, using

2000 iterations as warm up and retaining the next 2000 sampling iterations, then we

closely examined the model diagnostics for the few parameters that still had not

converged in 10 species' models. The spatial model estimates parameters for each local

community, enabling us to plot the maximum Rhat for each local community in the

species' range. We found that for American Robin (*Turdus migratorius*), grid cells that

included a parameter with Rhat > 1.1 were from data-sparse locations outside of our study

region (north of 25˚N, south of 55˚N, and east of 135˚W), so we refit the model with only

the data in our study region for 4000 sampling iterations. The rest of the species have

patchy geographic distributions in some or all parts of their ranges: local communities in

which species are detected are not spatially adjacent, leading to fewer neighboring

communities and, therefore, poor performance of the spatial model. For these nine

species, we reran the model with an alternative approach to identifying spatial neighbors

(Voronoi tessellation of the grid-centers within the species' range); this method increases

the total number of neighbor relationships (i.e., broadens the definition of neighbor beyond

the queen's-rule adjacency used in the default approach; (51)), increases the sharing of

information among grids, and thereby improved the model fit, especially for species with

more patchy distributions (51, 52).

Finally, we transformed the relative abundances from the Bayesian models for each of

the 470 species into estimates of abundance (i.e., total number of birds for each species,

year, and grid-cell), based on methods used in earlier analyses (38, 54) that account for

variation in each species' detection probabilities during NABBS sampling protocols. The

relative abundance of each species in each year and local community is multiplied by a

series of adjustment factors that account for variations in detectability of different species.

Briefly, distance adjustments account for the distances at which different species can be

detected by visual or auditory signals, pair adjustments account for the probability that

one individual of the breeding pair may be more easily detected (e.g., singing male vs.

skulking female), and time-of-day adjustments account for the variation of detectability by

the time of the survey among different species (54). NABBS was designed to compare

abundance within species, not between species; these adjustments are therefore

essential to our community-level analyses for not only improving accuracy but also making

estimates between species comparable. We obtained the adjustment coefficients of 402

out of 470 species from (38). For the rest of the species, which are mostly waterbirds (34

species) and waterfowl (28 species), we followed previous methodologies to calculate the

time-of-day adjustments (54). We then filled in missing distance and pair adjustments

using the means for species within the same genus (if available), family, or order. Finally,

we estimated distance and pair adjustments for Wood Stork (*Mycteria americana*), the only

species without all the above information, by its behavioral patterns (no difference in

detectability between sexes, and detectable from a long distance; (55)). In general, these

methods remove the biases in observed counts among species by adjusting for the higher

detectability of large, conspicuous, and soaring species such as pelicans, herons, and

crows and the lower detectability of smaller songbirds and inconspicuous species such as

warblers, wrens, and hummingbirds. We did not rerun the analyses by removing species

with newly calculated adjustments because it would fundamentally change the

community compositions. Rather, we checked the sensitivity by rerunning all analyses

without adjustments for all species. Although these adjustments are necessary to

accurately compare abundances among species, we have provided supplemental results

for the full analysis without these adjustments (fig. S13-S15). Some of our conclusions are

not qualitatively affected by these adjustments. In fact, our main conclusion that a small

number of species greatly contribute to functional shifts across North America remained,

because many common, large-bodied species that often have extreme values in functional

traits are down-weighted by the adjustments. A complete species list with the adjustment

factors used to estimate corrected abundances can be found in **Supplementary Data S1**.

Functional traits, community-weighted means, and species contributions

We obtained the following morphological traits from AVONET (17): beak length (both

the tip of the beak to the culmen and the tip of the beak to the nares), beak width, beak

depth, the length of the relaxed wing chord, Kipp's distance, the length from the carpal

joint to the tip of the longest secondary, and body mass. We obtained data on clutch size

from (56) and generation length from (36). Generation length is a composite measurement

that roughly represents the average parental age (35). To obtain a holistic size metric for

beak and wing while accounting for their phylogenetic correlations among species, we

conducted phylogenetic principal component analyses (pPCA) separately on the four beak

traits and three wing traits of the 470 species in our analyses using the R package *phytools*

(57) and the dated tree from the 2023 version of Clements taxonomy (15). The first PC axes

(PC1) explained 78.04% of and 86.94% of the total variance for the beak and wing traits,

respectively (fig. S16). To correct for the tight correlation between beak and wing lengths,

clutch size, and generation length with body mass (27, 33, 35), we conducted phylogenetic

linear models for log-transformed values of each of these traits, with log-transformed body

mass as the predictor variable, then extracted the residuals from each model (58). For

beak and wing lengths, these residuals are considered as separate traits from beak and

wing sizes, and we named them relative beak/wing lengths for the consistency with

previous studies (27). Together, we calculated community-weighted means for seven

functional traits: beak PC1, wing PC1, relative beak length, relative wing length, body

mass, corrected clutch size, and corrected generation length.

We calculated community-weighted means (CWMs) at each year t from the relative

abundance of species present in each grid (28):

$$CWM_t = \sum_{i=1}^{n} A_{i,t} x_i \ (1),$$

where $A_{i,t}$ is the local relative abundance of species i at year t, calculated as the

proportion of species i's relative abundance at year t over the sum of relative abundances

of all species in that community and year. n is the number of species in the local

community (i.e., $\sum_{i=1}^{n} A_{i,t} = 1$), and x_i is the value of the trait Q for species i. We

characterized the temporal trend of CWM for each trait Q by a simple linear model:

$$CWM_{Q,t} = \beta_{0,Q} + \beta_Q t + \epsilon_{Q,t},$$

$$t = t_{original} - 1969, \text{ and}$$

$$\epsilon_{Q,t} \sim Normal(0, \sigma_Q{}^2) \ (2),$$

where the estimated $\widehat{\beta_Q}$ is the temporal slope, and the transformed year is the difference

between the original year and 1969, such that the first year in the data (1970) was set to 1.

All t in the following models were transformed as such.

To further partition the contributions of each species to the temporal slope of CWMs,

we decomposed $\widehat{\beta_Q}$ by first presenting it as the Ordinary Least Squares (OLS) solution

using the variance-covariance form:

$$\widehat{\beta_Q} = \frac{Cov(CWM_{Q,t}, t)}{Var(t)} \ (3).$$

Combining Eqns. 1 and 3, $\widehat{\beta_Q}$ can be decomposed as

$$\widehat{\beta_Q} = \frac{Cov(CWM_{Q,t}, t)}{Var(t)} = \frac{Cov(\sum_{i=1}^{n} A_{i,t} x_i)}{Var(t)} = \sum_{i=1}^{n} \frac{Cov(A_{i,t}, t)}{Var(t)} x_i \ (4).$$

On the other hand, we can estimate the temporal slope of species i's relative

abundance changes as:

$$A_{i,t} = \beta_{0,i} + \beta_{A_i} t + \epsilon_{i,t}, \text{ and}$$

519
$$\epsilon_{i,t} \sim Normal(0, \sigma_i^2) \ (5).$$

520   Eqn. 5 can be written with the OLS solution using the variance-covariance form as well:

521
$$\widehat{\beta_{A_i}} = \frac{Cov(A_{i,t}, t)}{Var(t)} \ (6).$$

522   Combining Eqns. 4 and 6, the temporal slope of the CWM of trait x is then

523
$$\widehat{\beta_Q} = \sum_{i=1}^{n} \widehat{\beta_{A_i}} x_i \ (7),$$

i.e., the contribution of species i is its temporal slope times its trait value. Note that this

decomposition (Eqn. 7) is mathematically exact—the temporal trend of CWM change is

exactly the summation of individual species trends (see fig. S17-S18 for an example using

real data from a local community).

Using this decomposition, we calculated species contributions to temporal changes in

CWMs in each local community. For each trait, we first ranked the species in each local

community by absolute values of their contributions, then counted the number of local

communities in which a species ranked in the top 10% in that local community. We then

ranked all species by this count and pooled the top 10 species for each of the seven traits,

defined as the "top contributors" (fig. S6-S12). For each local community, we conducted

the following analyses using three sets of species: the full set (complete community), a

subcommunity containing only the species among the top contributors, and a

subcommunity containing the rest of the species (standard contributors). We also

recalculated CWMs for the top contributors and the standard contributors separately for

further analyses. Note that no variables (CWMs, and relative abundance) were

transformed or scaled in the calculation, but time was subtracted from the starting year

(see Eqn. 2). However, for better visualization, we scaled all trait values to their averages

among all 470 species and used these scaled values as the x-axis of Fig. 3. We did not fit

actual linear models to test for statistical significance; therefore, adjustment for multiple

comparisons were not necessary. To test the sensitivity of our analyses, we conducted all

analyses with the top five instead of top 10 contributor for each trait and found no

qualitative changes in our main results (fig. S19-20).

Temporal shifts in the functional space

To further characterize the temporal change of local communities, we mapped the

community in a grid each year in a functional space shared among all communities over

years, constructed with a PCA of CWMs of the five key traits (beak PC1, wing PC1, body

mass, generation length, and clutch size). We then tracked the location of each local

community over time within the functional space, creating a trajectory, and characterized

the community trajectory using the position of their start and end points (1970 and 2021)

and the overall direction of change from 1970 to 2021. The overall direction (effective

angle) was measured as the angle relative to the positive x-axis of the vector pointing from

the beginning (1970) to the end (2021) of the trajectory (Fig. 3A). Finally, we constructed

and analyzed trajectories from the CWMs of the two subcommunities: top contributors

and standard contributors. To ensure all trajectories were comparable, we projected the

subcommunities onto the PC axes of the PCA with the CWMs of the complete community

so that all trajectories share the same axes. After obtaining the position of each local

community in the functional space, measured in quadrants, we spatially mapped the local

community each year and calculated the spatial heterogeneity by Moran's *I* statistic using

the *spdep* package (59). We compared the time series of spatial heterogeneity calculated

from the CWMs of the base community and the two subcommunities to further illustrate

the contribution of the top 23 species to the spatial patterns of functional shift over time.

To compare patterns of functional shifts across major biogeographical regions, we

categorized each local community into its bird conservation regions (BCRs; (60)). For

spatial grids that fall into multiple BCRs, we calculated the percentage area in each BCR

and only considered the one with the highest percentage. We further generalized the BCRs

in the continental U.S. and Canada into seven major "avifaunal biomes" (61): Arctic,

Northern Forest, Pacific, Intermountain West, Southwest, Prairie, and Eastern (fig. S5).

Environmental data

We quantified the association between shifts in CWMs and changes in key

environmental variables to explore the potential mechanisms of community-level

functional shifts and to facilitate their comparisons with within-species expectations. We

obtained a time series of the mean temperature and precipitation in the warmest quarter

of the year ($bio10$ and $bio18$) from the WorldClim2 dataset (62). The warmest quarter

coincides with both the breeding season of most North American birds and the period

when NABBS is conducted. Both $bio10$ and $bio18$ are strongly correlated with mean

annual temperature (MAT; linear model, slope = 0.650, 95% CI = [0.649, 0.651], $p < 0.0001$)

and mean annual precipitation (MAP; slope = 0.173, 95% CI = [0.172, 0.174], $p < 0.0001$);

therefore, we do not expect results to be qualitatively different if using MAT and MAP

instead. To evaluate the effect of seasonality on key demographic traits, we calculated the

seasonality of temperature and precipitation as the difference between the warmest and

coldest months ($bio5 - bio6$) and the wettest and driest months ($bio13 - bio14$; (63)). To

explore the relationship between human-induced land use change and functional shifts,

we obtained land cover data from the HYDE 3.3 database, which estimates historical

anthropogenic land use changes (64). The HYDE data contain 20 categories of

anthropogenic biomes, or "anthromes" (9, 30); we generalized land use types into four

main categories: settlements, which contains urban areas and villages; agriculture, which

contains croplands and rangelands; cultured land, which contains woodland and drylands

with human inhabitants, and wild land. Settlements and agriculture are intensely impacted

by human activities, cultured land receives less human impact, and wildland contains

woodland and drylands with minimum human impacts. To match the spatial resolution of

estimated species abundances, we calculated the proportions of areas in each land use

category in a one-degree latitude and longitude grid cell. We obtained yearly time series of

all the above environmental data from 1970 to 2021 and aggregated them to the same one-

degree grid cells as our abundance data.

Model fitting

Before fitting the full model, we first fit the following linear mixed-effect models with

the R package *lme4* (65):

$$CWM_Q = \beta_{0,Q} + \beta_{1,Q} t + b_{g,Q} + \epsilon_Q,$$

$$b_{g,Q} \sim Normal\left(0, \sigma_g^2\right), \text{and}$$

$$\epsilon_Q \sim Normal(0, \sigma^2) \text{ (8)}.$$

Specifically, we separately modeled the CWM of each trait Q as a linear response of

transformed year t with slope $\beta_{1,Q}$, intercept $\beta_{0,Q}$, grid-level random effect $b_{g,Q}$, and the

overall residual ϵ_Q. We used a simple grid-level random effect because there was no

spatial autocorrelation in the residuals of the full model (see below). Both $b_{g,Q}$ and ϵ_Q are

normally distributed. We weighted the fitted residual $\widehat{\epsilon_Q}$ by the inverse of the overall

abundance of the local community within each grid. This weighting corrects for the

potentially higher measurement errors in communities with lower overall abundance. In all

linear mixed effects models, we log-transformed the CWM of body mass. Because we fit

the CWM of each trait separately, we adjusted the p-value of the models using the

Benjamini-Hochberg method (66). Fitting the year-only model yielded significant slopes for

CWMs of all traits but generation length (table S3).

Then, to explore whether the year effect still accounted for variances unexplained by

all environmental variables (bioclimatic variables and human-induced land use

proportions), we fit a year-environment model:

$$CWM_Q = \beta_{0,Q} + \beta_{1,Q}t + E\beta_{2,Q} + b_{g,Q} + \epsilon_Q \text{ (9)},$$

where E is a vector of environmental variables, $\beta_{2,Q}$ is a vector of slopes corresponding to

each environmental variable, and $b_{g,Q}$ and ϵ_Q follow definitions in Eqn. 8. Because the land

use type is proportional data (summing to 1), we removed wild land when fitting the model.

We then scaled all the environmental variables to ensure the slopes are comparable in

magnitude. In this model, year slopes for CWMs of all traits were significant and had the

same signs comparing to the year-only model (**Supplementary Data S3**), indicating that

year explained variations in CWMs of most traits even after accounting for all

environmental variables, which changed temporally, and should be considered as a fixed

effect rather than a temporal random effect.

Finally, we fit the full model with year, environmental variables, year-environment

interactions, and grid-level random effects:

$$CWM_Q = \beta_{0,Q} + \beta_{1,Q}t + E\beta_{2,Q} + (t \times E)\beta_{3,Q} + b_{g,Q} + \epsilon_Q \text{ (10)},$$

where $t \times E$ denotes the interactive effects between year and environmental variables. We

analyzed the models of each CWM using variograms in the R package *gstat* (67) and found

no evidence of strong spatial autocorrelation in the residuals (fig. S21). We then fit the full

model (Eqn. 10) with CWMs of the two subcommunities, consisting of the top contributors

and the standard contributors, then compared their slopes of the same predictor

variables. We fit separate models rather than using group-level slopes within a single

model because CWMs do not contain species identities and therefore can only be

calculated for different groups of species. See **Supplementary Data S4** for full diagnostics

of the models.

Temperature and precipitation seasonality may only affect the life history strategies of

residents who are exposed to climatic extremes throughout the whole year. Therefore, we

analyzed the associations between temperature/precipitation seasonality and the two

demographic traits, clutch size and generation length, separately for residents and

migrants. To do so, we filtered species by their migration status according to (38), then

separately calculated the CWMs for each group in local communities.

Software

All analyses were conducted by R version 4.4.1. (68). In addition to specific packages

mentioned in methods, all data wrangling and visualization were conducted using the

package *tidyverse* (69), and all geospatial analyses were conducted using the package *sf*

(70).

Reference

1. C. Parmesan, *et al.*, Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature* **399**, 579–583 (1999).

2. M. W. Tingley, W. B. Monahan, S. R. Beissinger, C. Moritz, Birds track their Grinnellian niche through a century of climate change. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 19637–19643 (2009).

3. I.-C. Chen, J. K. Hill, R. Ohlemüller, D. B. Roy, C. D. Thomas, Rapid Range Shifts of Species Associated with High Levels of Climate Warming. *Science* **333**, 1024–1026 (2011).

4. B. C. Weeks, *et al.*, Shared morphological consequences of global warming in North American migratory birds. *Ecology Letters* **23**, 316–325 (2020).

5. S. A. MacLean, S. R. Beissinger, Species' traits as predictors of range shifts under contemporary climate change: A review and meta-analysis. *Global Change Biology* **23**, 4094–4105 (2017).

6. K. Zhu, *et al.*, Rapid shifts in grassland communities driven by climate change. *Nat Ecol Evol* **8**, 2252–2264 (2024).

7. M. A. Jarzyna, W. Jetz, A near half-century of temporal change in different facets of avian diversity. *Global Change Biology* **23**, 2999–3011 (2017).

8. J. Rogelj, M. Meinshausen, R. Knutti, Global warming under old and new scenarios using IPCC climate sensitivity range estimates. *Nature Clim Change* **2**, 248–253 (2012).

9. E. C. Ellis, K. Klein Goldewijk, S. Siebert, D. Lightman, N. Ramankutty, Anthropogenic transformation of the biomes, 1700 to 2000. *Global Ecology and Biogeography* **19**, 589–606 (2010).

10. S. Ryding, M. Klaassen, G. J. Tattersall, J. L. Gardner, M. R. E. Symonds, Shape-shifting: changing animal morphologies as a response to climatic warming. *Trends in Ecology & Evolution* **36**, 1036–1048 (2021).

11. C. Moritz, *et al.*, Impact of a century of climate change on small-mammal communities in Yosemite National Park, USA. *Science* **322**, 261–264 (2008).

12. M. H. C. Neate-Clegg, B. A. Tonelli, M. W. Tingley, Advances in breeding phenology outpace latitudinal and elevational shifts for North American birds tracking temperature. *Nat Ecol Evol* 1–10 (2024). https://doi.org/10.1038/s41559-024-02536-z.

13. S. Ryding, A. McQueen, M. Klaassen, G. J. Tattersall, M. R. E. Symonds, Long- and short-term responses to climate change in body and appendage size of diverse Australian birds. *Global Change Biology* **30**, e17517 (2024).

690   14. D. W. Gibbons, P. F. Donald, H.-G. Bauer, L. Fornasari, I. K. Dawson, Mapping avian
691   distributions: the evolution of bird atlases. *Bird Study* **54**, 324–334 (2007).

692   15. E. J. McTavish, *et al.*, A complete and dynamic tree of birds. *Proceedings of the National*
693   *Academy of Sciences* **122**, e2409658122 (2025).

694   16. H. Wilman, *et al.*, EltonTraits 1.0: Species-level foraging attributes of the world's birds and
695   mammals. *Ecology* **95**, 2027–2027 (2014).

696   17. J. A. Tobias, *et al.*, AVONET: morphological, ecological and geographical data for all birds.
697   *Ecology Letters* **25**, 581–597 (2022).

698   18. J. A. Sheridan, D. Bickford, Shrinking body size as an ecological response to climate change.
699   *Nature Clim Change* **1**, 401–406 (2011).

700   19. V. Jirinec, *et al.*, Morphological consequences of climate change for resident birds in intact
701   Amazonian rainforest. *Science Advances* **7**, eabk1743 (2021).

702   20. C. Youngflesh, J. F. Saracco, R. B. Siegel, M. W. Tingley, Abiotic conditions shape spatial and
703   temporal morphological variation in North American birds. *Nat Ecol Evol* **6**, 1860–1870
704   (2022).

705   21. A. P. Møller, Successful city dwellers: a comparative study of the ecological characteristics
706   of urban birds in the Western Palearctic. *Oecologia* **159**, 849–858 (2009).

707   22. M. H. C. Neate-Clegg, *et al.*, Traits shaping urban tolerance in birds differ around the world.
708   *Current Biology* **33**, 1677-1688.e6 (2023).

709   23. C. S. Rushing, J. A. Royle, D. J. Ziolkowski, K. L. Pardieck, Migratory behavior and winter
710   geography drive differential range shifts of eastern birds in response to recent climate
711   change. *Proceedings of the National Academy of Sciences* **117**, 12897–12903 (2020).

712   24. A. M. Schipper, *et al.*, Contrasting changes in the abundance and diversity of North
713   American bird assemblages from 1971 to 2010. *Global Change Biology* **22**, 3948–3959
714   (2016).

715   25. J.-Y. Barnagaud, P. Gaüzère, B. Zuckerberg, K. Princé, J.-C. Svenning, Temporal changes in
716   bird functional diversity across the United States. *Oecologia* **185**, 737–748 (2017).

717   26. J. R. Sauer, *et al.*, The first 50 years of the North American Breeding Bird Survey. *The Condor*
718   **119**, 576–593 (2017).

719   27. S. Ryding, A. McQueen, Multiple regression, not ratios, for analyzing relative appendage
720   size. *Evol Ecol* (2025). https://doi.org/10.1007/s10682-025-10346-3.

28. S. Lavorel, *et al.*, Assessing functional diversity in the field – methodology matters!
Functional Ecology **22**, 134–147 (2008).

29. S. Villéger, N. W. H. Mason, D. Mouillot, New Multidimensional Functional Diversity Indices
for a Multifaceted Framework in Functional Ecology. *Ecology* **89**, 2290–2301 (2008).

30. E. C. Ellis, *et al.*, People have shaped most of terrestrial nature for at least 12,000 years.
Proceedings of the National Academy of Sciences **118**, e2023483118 (2021).

31. K. L. Thompson, J. M. Chase, R. Remelgado, C. Meyer, The interacting effects of climate and
land-cover change on bird communities in the United States. *Ecography* **2025**, e07949
(2025).

32. R. E. Ricklefs, Geographical Variation in Clutch Size among Passerine Birds: Ashmole's
Hypothesis. *The Auk* **97**, 38–49 (1980).

33. W. Jetz, C. H. Sekercioglu, K. Böhning-Gaese, The Worldwide Variation in Avian Clutch Size
across Species and Space. *PLOS Biology* **6**, e303 (2008).

34. C. B. Cooper, W. M. Hochachka, G. Butcher, A. A. Dhondt, Seasonal and Latitudinal Trends in
Clutch Size: Thermal Constraints During Laying and Incubation. *Ecology* **86**, 2018–2031
(2005).

35. C. Youngflesh, *et al.*, Environmental Variability Shapes Life History of the World's Birds.
Ecology Letters **28**, e70077 (2025).

36. J. P. Bird, *et al.*, Generation lengths of the world's birds and their implications for extinction
risk. *Conservation Biology* **34**, 1252–1261 (2020).

37. W. M. Schaffer, Optimal Reproductive Effort in Fluctuating Environments. *The American*
Naturalist **108**, 783–790 (1974).

38. K. V. Rosenberg, *et al.*, Decline of the North American avifauna. *Science* **366**, 120–124
(2019).

39. A. Nord, E. Persson, J. K. R. Tabh, E. Thoral, Shrinking body size may not provide meaningful
thermoregulatory benefits in a warmer world. *Nat Ecol Evol* **8**, 387–389 (2024).

40. J. K. R. Tabh, *et al.*, Limited evidence that body size shrinking and shape-shifting alleviate
thermoregulatory pressures in a warmer world. *Commun Biol* **8**, 1–13 (2025).

41. C. Teplitsky, J. A. Mills, J. S. Alho, J. W. Yarrall, J. Merilä, Bergmann's rule and climate change
revisited: Disentangling environmental and genetic responses in a wild bird population.
Proceedings of the National Academy of Sciences **105**, 13492–13496 (2008).

42. C. P. terHorst, *et al.*, Evolution in a Community Context: Trait Responses to Multiple Species
Interactions. *The American Naturalist* **191**, 368–380 (2018).

43. T. Dias, *et al.*, Long-term increases in wing length occur independently of changes in climate
and climate-driven shifts in body size. *Proceedings of the Royal Society B: Biological*
Sciences **292**, 20242556 (2025).

44. C. H. Trisos, C. Merow, A. L. Pigot, The projected timing of abrupt ecological disruption from
climate change. *Nature* **580**, 496–501 (2020).

45. F. Hernández, L. A. Brennan, S. J. DeMaso, J. P. Sands, D. B. Wester, On reversing the
northern bobwhite population decline: 20 years later. *Wildlife Society Bulletin* **37**, 177–188
(2013).

46. J. R. Allan, J. S. Kirby, C. J. Feare, The biology of canada geese Branta canadensis in relation
to the management of feral populations. *Wildlife Biology* **1**, 129–143 (1995).

47. J. L. Tapley, R. K. Abernethy, J. E. Kennamer, N. Carolina, Status and distribution of the wild
turkey in 2004 in *Proceedings of the National Wild Turkey Symposium*, (2005), pp. 21–32.

48. I. Benmazouz, *et al.*, Corvids in Urban Environments: A Systematic Global Literature Review.
Animals **11**, 3226 (2021).

49. K. Princé, B. Zuckerberg, Climate change in our backyards: the reshuffling of North
America's winter bird communities. *Global Change Biology* **21**, 572–585 (2015).

50. G. J. Di Cecco, A. H. Hurlbert, Anthropogenic drivers of avian community turnover from local
to regional scales. *Global Change Biology* **28**, 770–781 (2022).

51. B. P. M. Edwards, A. C. Smith, S. LaZerte, bbsBayes2: Hierarchical Bayesian Analysis of North
American BBS Data. (2023). Deposited 2023.

52. A. C. Smith, *et al.*, Spatially explicit Bayesian hierarchical models improve estimates of avian
population status and trends. *Ornithological Applications* **126**, duad056 (2024).

53. W. A. Link, J. R. Sauer, D. K. Niven, Model selection for the North American Breeding Bird
Survey: A comparison of methods. *The Condor* **119**, 546–556 (2017).

54. J. Stanton, P. Blancher, K. Rosenberg, A. Panjabi, W. Thogmartin, Estimating uncertainty of
North American landbird population sizes. *Avian Conservation and Ecology* **14** (2019).

55. M. C. Coulter, J. A. Rodgers Jr., J. C. Ogden, F. C. Depkin, Wood Stork (Mycteria americana),
version 1.0. *Birds of the World* (2020).
https://doi.org/10.2173/bow.woosto.01species_shared.bow.project_name.

56. N. P. Myhrvold, *et al.*, An amniote life-history database to perform comparative analyses
784   with birds, mammals, and reptiles. *Ecology* **96**, 3109–3109 (2015).

785   57. L. J. Revell, phytools 2.0: an updated R ecosystem for phylogenetic comparative methods
786   (and other things). *PeerJ* **12**, e16505 (2024).

787   58. Y. Xu, *et al.*, Ecological predictors of interspecific variation in bird bill and leg lengths on a
788   global scale. *Proceedings of the Royal Society B: Biological Sciences* **290**, 20231387 (2023).

789   59. R. Bivand, R Packages for Analyzing Spatial Data: A Comparative Case Study with Areal Data.
790   *Geographical Analysis* **54**, 488–518 (2022).

791   60. J. R. Sauer, J. E. Fallon, R. Johnson, Use of North American Breeding Bird Survey Data to
792   Estimate Population Change for Bird Conservation Regions. *The Journal of Wildlife*
793   *Management* **67**, 372–389 (2003).

794   61. T. L. Crewe, J. D. McCracken, P. D. Taylor, D. Lepage, A. E. Heagy, The Canadian Migration
795   Monitoring Network-Réseau canadien de surveillance des migrations: Ten-year report on
796   monitoring landbird population change. *CMMN-RCSM Scientific Technical Report* **1** (2008).

797   62. S. E. Fick, R. J. Hijmans, WorldClim 2: new 1-km spatial resolution climate surfaces for global
798   land areas. *International Journal of Climatology* **37**, 4302–4315 (2017).

799   63. A. J. Boyce, B. G. Freeman, A. E. Mitchell, T. E. Martin, Clutch size declines with elevation in
800   tropical birds. *The Auk* **132**, 424–432 (2015).

801   64. C. G. M. K. Goldewijk, Anthropogenic land-use estimates for the Holocene; HYDE 3.2. DANS
802   Data Station Archaeology. https://doi.org/10.17026/dans-25g-gez3. Deposited 29
803   September 2017.

804   65. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using lme4.
805   *Journal of Statistical Software* **67**, 1–48 (2015).

806   66. Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate: A Practical and Powerful
807   Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*
808   *(Methodological)* **57**, 289–300 (1995).

809   67. E. J. Pebesma, Multivariable geostatistics in S: the gstat package. *Computers & Geosciences*
810   **30**, 683–691 (2004).

811   68. R Core Team, R: A language and environment for statistical computing. (2024).

812   69. H. Wickham, *et al.*, Welcome to the Tidyverse. *Journal of Open Source Software* **4**, 1686
813   (2019).

814   70. E. Pebesma, Simple Features for R: Standardized Support for Spatial Vector Data. *The R*
815   *Journal* (2018).

Figure Legends

Fig. 1: Overview of the analyses and associations between functional shifts and environmental variables. **A.** Map of the study region, showing the 1261 local communities (one by one degree grids) included in the analyses. **B.** Overview of the analyses. **C.** Associations between community-weighted means (CWMs) of seven key functional traits and main environmental variables, including climate, human-induced land use, and year. CWMs are calculated from either all species in the communities (complete communities), only the top contributors, or only the rest of the species (standard contributors). All statistics are shown in table S1. Illustration by Olivia Stein.

Fig. 2: Analyzing the temporal trajectory of local communities in the functional space. **A.** The functional space constructed by PCA of all communities by their community-weighted mean measurements over time, and a temporal trajectory of one local community (40°N, 110°W) moving through the functional space over time, showing the start and end points and the overall direction (effective angle). In the trajectory, lighter shades represent earlier years, and darker shades represent later years. The two panels represent the same functional space, but the scales are adjusted for better visualization. **B.** Maps of each local community's start and end positions in the functional space, measured by the quadrants in **A**, and the overall direction of the functional shifts, measured by the quadrants to which the trajectory travels. Columns show results for complete communities, subcommunities with top contributors, and subcommunities with the standard contributors. **C.** Spatial heterogeneity of functional positions of the base communities, top contributors, and the rest of the species, calculated as Moran's *I* from the maps. A Moran's *I* closer to 0 means lower spatial heterogeneity.

Fig. 3: Species contributions to the community-weighted means (CWMs) of seven key functional traits. Panels only include species that contribute substantially (ranked top 10% of all species in a local community) in at least one local community; point size denotes the number of grids in which a species contributes greatly. The sign of the contribution depends on the species' temporal trends of relative abundance and its trait value, e.g., if a species with below-average trait value increases in relative abundance, it will contribute negatively by lowering the CWM of the trait. For each trait, the identity of the top 10 species that contribute to the CWMs of the most local communities is labeled; pooling all these species results in 25 top contributors of the observed functional shifts over time, shown on the right. For each panel, the x-axis is the scaled trait values, and the y-axis is the median of the temporal slopes of relative abundances across each species' range. Shades denote the absolute magnitudes of species contributions: colder colors represent negative contributions, and warmer colors represent positive contributions. Axes and scales are different for each panel for better visualization. Illustrations by Olivia Stein.