Historical shifts, geographic biases, and biological constraints shape mammal species discovery

Matheus de T. Moroti[1,†], Jhonny J. M. Guedes[1,2,†,*], Guilherme M. Missio[1],

Giovana L. Diegues[1], Alexandra M. R. Bezerra[3], and Mario R. Moura[1,4]

[1]*Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, SP, 13083-862, Brazil*

[2]*Departamento de Ecologia, Instituto de Ciências Biológicas, Universidade Federal de Goiás – Campus Samambaia, Goiânia, GO, 74690-900, Brazil*

[3]*Mastozoologia, Coordenação de Zoologia, Museu Paraense Emílio Goeldi, Belém, PA ,66077-830, Brazil*

[4]*Departamento de Sistemática e Ecologia, Universidade Federal da Paraíba, João Pessoa, PB, 58051-900, Brazil*

[*]*Corresponding author: Jhonny J. M. Guedes. Programa de Pós-Graduação em Ecologia e Evolução, Departamento de Ecologia, Universidade Federal de Goiás, Goiânia, GO, 74690-900, Brazil. E-mail: jhonnyguds@gmail.com*

† These authors contributed equally to this work.

Abstract

Species descriptions in taxonomy have become increasingly comprehensive, yet disparities persist across taxa and regions. We assess temporal trends in mammal species descriptions (1990–2025) using four proxies of comprehensiveness—counts of examined specimens and compared taxa, number of pages (only from the Methods/Results sections), and number of evidence lines (i.e., analytical tools and techniques). Using generalized linear models, we assessed how these proxies are explained by factors associated with species' biology, geography, and taxonomic practice. Most new species derive from tropical regions, particularly rodents and bats, reflecting global discovery hotspots. Descriptions have grown more rigorous over time, with expanded specimen sampling, broader taxonomic comparisons, and integrative methods. However, disparities

emerge along geographic and biological axes: descriptions from temperate regions

incorporate more evidence lines, while small-bodied and tropical species (especially bats)

remain understudied due to sampling biases and resource limitations. Body size inversely

correlates with description length, as smaller species often require advanced diagnostics.

Species-rich genera show greater comprehensiveness, likely due to heightened diagnostic

scrutiny. Our findings highlight progress in taxonomic rigor but underscore persistent

gaps tied to geography, body size, and accessibility of analytical tools. Addressing these

disparities requires targeted investments in local capacity, equitable collaboration, and

accessible methodologies to strengthen global taxonomic infrastructure and support

conservation priorities.

Key words: Linnean shortfall, Mammalia, species descriptions, taxonomic practice,

taxonomy.

1. Introduction

Species represent fundamental units in biodiversity science (Braby et al. 2024), where their names constitute testable hypotheses subject to revision in light of new data (Bremer et al. 1990; Raposo et al. 2017; Streicher et al. 2023; Gippoliti et al. 2024). Although taxonomic science is inherently dynamic and absolute stability is undesirable (Dominguez and Wheeler 1997; Dubois 2007), minimizing unnecessary changes in species identities can increase usability of biodiversity databases and improve scientific communication, data sharing, and conservation planning (Bertrand and Härlin 2006; Sullivan et al. 2023; Hamdan et al. 2024; Mahony et al. 2024). Following the development and popularization of integrative taxonomy, contemporary species descriptions tend to be more robust, incorporating more comprehensive diagnostic criteria (Sangster and Luksenburg 2015; Poulin and Presswell 2016; Guedes et al. 2024b) and contrasting with older descriptions, which generally relied on simpler diagnostics (Sangster and Luksenburg 2015; Poulin and Presswell 2016; Guedes et al. 2020). This helps explain the observed decline in synonymization rates over time (Solow et al. 1995; Alroy 2002; Baselga et al. 2008; Guedes et al. 2025), although this pattern may reflect a temporal bias (Alroy 2002) or changes in taxonomic practice (Lumpers vs. Splitters debate, Isaac et al. 2004). Understanding how temporal trends in species description comprehensiveness are related to improved sampling and analytical frameworks helps to inform best practices that can strengthen taxonomic science.

Comprehensive descriptions can help mitigate common pitfalls in taxonomic practices (Dayrat 2005; Wüster et al. 2021; Braby et al. 2024), such as oversplitting due to insufficient diagnostic criteria and failure to accurately distinguish natural variation within species (Isaac et al. 2004), or undersplitting resulting from overlooked phenotypic

or genetic divergence (Chek et al. 2003; Hendrixson and Bond 2005; Mutanen et al.

2016). While the accumulated evidence highlights temporal shifts in taxonomic practices

(Sangster and Luksenburg 2015; Padial and De la Riva 2021; Guedes et al. 2024b), recent

studies also reveal pronounced spatial variation (De Queiroz 2007; Freeman and Pennell

2021; Padial and De la Riva 2021). A well-documented latitudinal taxonomic bias exists,

wherein biodiversity-rich but resource-limited regions (e.g., the Tropics) often exhibit

lower taxonomic resolution compared to temperate, high-income regions (Freeman and

Pennell 2021; Diniz Filho et al. 2023; Guedes et al. 2025). This spatial variation likely

emerges from differences in access to resources and tools, though international

collaborations and open-access data initiatives have the potential to mitigate these

imbalances (Nakamura et al. 2023; Carneiro et al. 2025; Weksler et al. 2025). Addressing

these gaps is critical to ensuring robust taxonomic frameworks that support downstream

research on fields dependent on taxonomic data (Diniz Filho et al. 2023; Lessa et al.

2024).

Because citation-based metrics are poor proxies of the comprehensiveness of

taxonomic publications (Valdecasas et al. 2000; Valdecasas 2011; Pyke 2014), more

direct metrics are needed to investigate trends in the robustness of species descriptions

and to inform whether many taxonomic changes are likely to occur in the future across

organismal groups. Using metrics like page counts and the number of evidence lines or

number of techniques adopted (e.g., morphometrics, cytogenetics, internal anatomy,

molecular analysis) in descriptions, studies have generally observed an increase in

comprehensiveness of descriptions across birds (Sangster and Luksenburg 2015), reptiles

(Guedes et al. 2024a), mollusks (Abreu et al. 2025), helminths (Poulin and Presswell

2016), and other taxa (Benton 2008; Pante et al. 2015). Here, we investigate how several

proxies of comprehensiveness in species description have changed over time and assess

their potential determinants.

We focus our assessment on recent mammals, a group with a relatively stable

alpha taxonomy when compared to other vertebrates (e.g., amphibians and reptiles; see

Moura and Jetz 2021; Guedes et al. 2024b), but yet still dynamic (Parsons et al. 2022).

Specifically, we explore whether temporal variation is evident across four proxies of

comprehensiveness in species descriptions, namely: (1) the number of specimens used to

describe the new taxa, (2) count of taxa to which the new species is compared, (3) page

counts, and (4) number of evidence lines used in the description. We examine whether

variability in these proxies is related to species' biology, geography, and taxonomic

practice. Additionally, we inspect how accessibility to specific resources and tools is

associated with the internationalization of species descriptions. Given that many

mammalian taxa still lack comprehensive taxonomic studies (Feijó and Brandão 2022),

and that most formal descriptions represent the splitting of one species into many (Isaac

et al. 2004; Burgin et al. 2018), it is likely that several taxonomic changes lie ahead.

Despite its focus on mammals, our study uses taxon-independent proxies for description

comprehensiveness, which combined with its global scope, help yield broader

implications for developing protocols and practices that can increase data quality and

usage in biodiversity research (Mahony et al. 2024).

2. Material and methods

2.1. Data collection

We examined the descriptions of 1,116 mammal species (including terrestrial, freshwater,

and marine) published between 1990 and 2025, covering 36 years of mammal discovery.

Focusing on post-1990 descriptions minimize inconsistencies from historical variations

in taxonomic standards, such as the cladistics and molecular systematics, ensuring

comparable scientific and socioeconomic trends. We only considered "original"

descriptions, disregarding redescriptions, revalidations, and other taxonomic works.

Taxonomic nomenclature and species account follows the Mammal Diversity Database

Version 2.1 (MDD 2025). For biogeographic regionalization, each species was assigned

to a realm based on the coordinates of its type-locality, using the spatial data from

Dinerstein et al. (2017).

For each species in our database, we extracted four proxies for the species

description comprehensiveness: (1) number of specimens - type-specimens plus all

referred specimens of the new species used to obtain diagnostic traits (type-series size

and all referred specimens were highly correlated, Pearson's $r = 0.60$, $p < 0.001$); (2)

number of taxa compared - the count of close-related species directly compared with the

new species; (3) number of pages - used as a proxy for the depth of taxonomic

information. The presence of taxonomic keys, figures, or expanded results, for instance,

typically represents a more detailed description and increases page and word count. We

quantified this by dividing each page into quadrants and counting only those containing

Methods and Results sections. For studies describing multiple species, the total count was

divided by the number of species. Our fourth proxy was represented by the (4) number of

evidence lines present in the formal description, scored from 1 to 9 lines of evidence

based on the presence or absence of the following analyzed characters: (i) dentition (e.g.

dental formula, tooth shape, etc), (ii) internal soft anatomy (e.g. internal morphology of

the stomach, glans, etc.), (iii) external qualitative morphology (e.g. comparison of head

and tail shapes), (iv) microscopic-based trichology, (v) coloration patterns, (vi)

karyotype, (vii) external morphometry (e.g., head-body length, hind foot length, forearm

length), (viii) osteological morphometry (from teeth, skull, mandible and other skeleton

parts), and (ix) DNA (e.g. allozyme, mtDNA, nucDNA, multiLoci). These proxies

showed low multicollinearity (Variance Inflation Factor, VIF < 1.5; Table S1) and limited

pairwise correlations (Spearman's r correlation range = 0.00-0.11; Fig. S1), indicating

they capture relatively distinct aspects of description comprehensiveness.

To represent potential determinants of species description comprehensiveness, we

used the seven variables: (1) year of species description; (2) number of authors; (3)

average number of countries per author, the country counts derived from authors

affiliations were divided by author counts to avoid sampling effects associated with

multiauthored works (see Fig. S2 for pairwise correlation between authors and countries

counts); (4) maximum body mass, obtained from original descriptions and from Moura

et al. (2024), (5) per-genus species richness based on the year of species descriptions (i.e.,

for each species, we tallied the total number of currently accepted species described in

the same genus until their year of description); (6) absolute latitude of type-locality; and

whether description was based on (7) taxonomic revision, which was considered as such

only if the work explicitly mentioned the term "revision" (or similar) in its title, abstract,

keywords, or main text. To allow cross-group comparisons, we also recorded data on

taxonomic ranks.

2.2. Statistical analyses

Temporal trends in our proxies of the species description comprehensiveness (number of

lines of evidence, number of pages, number of specimens, and number of taxa compared)

were investigated using Spearman's (r_s) correlation between the year of publication and

annual means for the metrics analyzed across species described in each year. We assessed

multicollinearity among continuous predictors using the Variation Inflation Factors (VIF;

Mansfield and Helms 1982), where strong multicollinearity is often attributed to variables

holding VIF > 10 (Kutner et al. 2005). All variables in our analysis had VIF values below

2 (Table S2), so none were excluded.

We assessed potential determinants of species description comprehensiveness,

using generalized linear models (GLMs; Zuur et al. 2009). We used a negative binomial

error distribution to analyze the number of specimens and number of taxa compared, both

of which are count data. The number of pages was calculated as a continuous variable, so

we applied a $\log_{10}$-transformation and modelled it using a GLM with a Gaussian error

distribution. Although the number of evidence can be considered a count variable, given

its hump-shaped distribution (Fig. S1), we used both negative binomial and Gaussian

distributions, comparing model fit and selecting the one with the lowest AIC values

(Burnham and Anderson 2002). Each response variable was modelled against year of

species description, number of authors ($\log_{10}$-transformed), average number of countries

per author ($\log_{10}$-transformed), body mass ($\log_{10}$-transformed), per-genus species

richness ($\log_{10}$-transformed), and occurrence of taxonomic revision. Continuous

predictors were z-transformed before modelling, allowing comparisons of their effect

sizes. Because of varying amounts of missing data among response and predictor

variables, GLMs were performed with datasets differing slightly in the number of

modelled species: number of lines of evidence ($n = 937$ species); number of pages ($n =$

950), number of specimens ($n = 933$); and number of taxa compared ($n = 942$). All

analyses were performed in the R software version 4.3.1 (R Core Team 2023).

All models were fitted using the package *MASS* (Venables and Ripley 2002). We

assessed explained variation through Nagelkerke's pseudo-R^2 for negative binomial

models (response variables 'number of taxa compared' and 'number of specimens') and

the coefficient of determination (R^2) for the Gaussian model (response variables 'number

of evidence' and 'number of pages'), both computed with the R package *performance*

(Lüdecke et al. 2021). We assessed the necessity of phylogenetic regression models by

investigating the phylogenetic autocorrelation of model residuals through Moran's *I*

correlograms, computed across 14 distance classes (Revell 2010). Correlograms were

derived from the average results of 100 fully sampled phylogenies for mammals (Upham

et al. 2019), with trees trimmed to include only modelled species. Phylogenetic

autocorrelation analyses were conducted with the R packages *phylobase* (Bolker et al.

2020) and *phylosignal* (Keck et al. 2016).

To further investigate if accessibility of costly analytical tools is reflected across

international collaboration, we applied a Kruskal-Wallis analysis to test whether the

average number of countries per authors differed among descriptions using (i) molecular

biology, (ii) taxonomic review, or (iii) other evidence lines. *Post-hoc* comparisons were

conducted using Dunn's test, with pairwise contrasts adjusted for multiple testing using

the Bonferroni correction method (Bonferroni 1936). All analyses were conducted using

the *rstatix* R package (Kassambara 2023). See Data Availability section for raw data and

R code (Moroti et al. 2025).

3. Results

Most mammal species described between 1990 and 2025 were from Africa, South

America, Southeast Asia, and islands near Australia (Fig. 1). The Afrotropical realm

stands out in terms of primate descriptions, whereas the few descriptions from the Nearctic were mostly from rodents and Eulipotyphla (e.g., hedgehogs, moles, true shrews). The majority of the 1,116 species described comprised rodents (41%, n = 458) and bats (26.2%, n = 293), a trend observed globally as well as across most biogeographic realms. Descriptive statistics for comprehensiveness metrics across all mammals, rodents, bats, and when considering mammals excluding bats and rodents, are summarized in Table 1.

Figure 1. Global distribution of mammal species described between 1990 and 2025. Each point on the map represents a species' type locality, while donut charts indicate the proportion of species described within each taxonomic order. Points are coloured according to their respective taxonomic order, with rodents in blue, bats in green, and the different shades of red representing other mammals. To improve visualization, taxonomic orders holding less than 5% of species descriptions were grouped under 'Other taxa' label in the donut charts. The total sample size (N) is lower than the 1,116 species described during this period due to missing type-locality data for some species. Two species occurring in the Oceania realm were suppressed due to low representation.

Table 1. Descriptive statistics for metrics of species descriptions' comprehensiveness. For each group, we report the median and the range (minimum–

233 maximum) for the number of evidence lines, number of specimens, number of pages, and
234 number of taxa compared.

	All mammals	Non-bats & non-rodents	Bats	Rodents
N. of evidence lines	5 (1–8)	5 (1–7)	5 (1–8)	5 (1–8)
N. of specimens	8 (1–409)	7 (1–345)	7 (1–327)	9 (1–409)
N. of pages	7.5 (0.23–179)	6.5 (0.23–48.2)	8.1 (0.8–179)	8.1 (1–130)
N. of taxa compared	5 (0–77)	4 (0–46)	5 (1–34)	4 (0–77)

3.1. Temporal trends in mammal description comprehensiveness

The annual mean number of evidence lines used in species descriptions increased when

analyzing all mammals ($r_s = 0.75$; $p < 0.001$), non-bat & non-rodent mammals ($r_s = 0.51$;

$p = 0.006$), bats ($r_s = 0.63$; $p < 0.001$) and rodents ($r_s = 0.64$; $p < 0.001$; Fig 2a). There

was no change in the number of pages over time (Fig. 2b), but an increase in the number

of specimens when considering all mammals together ($r_s = 0.49$; $p = 0.02$, Fig. 2c) and

non-bat & non-rodent mammals ($r_s = 0.44$; $p = 0.03$). The number of taxa compared also

increased for all mammals combined ($r_s = 0.61$; $p < 0.001$, Fig. 2d), non-bat & non-rodent

mammals ($r_s = 0.56$; $p < 0.001$) and for bats separately ($r_s = 0.7$; $p < 0.001$), but not for

rodents ($r_s = 0.25$; $p = 0.71$). In addition, the incorporation of molecular analyses in

species descriptions has grown over time, with a noticeable increase in the proportion of

taxa described annually using molecular data (Figs S3-S4).

Figure 2. Variation in proxies of mammal species descriptions comprehensiveness over time (1990–2025). Panels show overall trends for all mammals (gray), non-rodents and non-bats (red), bats (green), and rodents (blue). Proxies include the (a) number of evidence used in species descriptions, (b) number of pages in taxonomic description papers, (c) number of specimens examined in species descriptions, and (d) number of taxa compared to distinguish new species. rs denotes the Spearman's correlation between the year of description and annual means for each variable. We applied Bonferroni corrections to p-values to account for multiple comparisons. Vertical lines represent standard errors, and shaded areas around the regression lines represent 95% confidence intervals.

3.2. Internationalization of mammal descriptions

We found a temporal decrease in the average number of countries per author for all taxonomic groups (Fig. 3). In addition, this 'internationalization indicator' differed across descriptions using analytical tools, with studies using molecular biology and taxonomic

review showing lower values of average number of countries per author. In bats and

rodents, only molecular-based descriptions involved fewer countries relative to other

lines of evidence.

Figure 3. Average number of countries per author involved in mammal species
descriptions across different lines of evidence and over time. Patterns are shown for
(a) all mammals, (b) non-bats and non-rodents, (c) bats, and (d) rodents. 'Other
evidences' refers to descriptions that do not include molecular diagnoses and were not
derived from taxonomic revisions. Molecular approaches exhibit different levels of
internationalization compared to taxonomic reviews and other evidence types across
mammals, bats, and rodents. In the violin plot, post-hoc comparisons were performed
using estimated marginal means, with pairwise contrasts adjusted for multiple testing
using the Bonferroni method. Groups that do not share letters are significantly different
from each other.

The use of molecular data in descriptions varies across mammalian orders, with

Pilosa (4 spp.), Carnivora (5 spp.), Artiodactyla (19 spp.), Peramelemorphia (5 spp.),

Macroscelidea (3 spp.), Didelphimorphia (25 spp.), Rodentia (n = 458 spp.), and

Eulipotyphla (146 spp.) frequently incorporating genetic evidence (more than 50%

described species). In comparison, Dasyuromorphia (13 spp.) lacked such data entirely.

International collaborations were common in molecular-based descriptions, except in

Lagomorpha, with nine descriptions associated with a single country (Fig. 4a).

Taxonomic revisions contributed substantially to new species descriptions in Pilosa,

Monotremata, and Carnivora (> 50%), but were rare in species-rich groups like Rodentia

and Chiroptera and absent in depauperate orders (e.g., Peramelemorphia,

Paucituberculata, Microbiotheria, Macroscelidea, Lagomorpha, and Hyracoidea). While

international collaboration in revision-based descriptions was idiosyncratic, primates and

afrosoricids exhibited broader multinational involvement—i.e., >75% of descriptions

involving two or more countries (Fig. 4b).

Figure 4. Proportion of species descriptions using (a) molecular analysis or (b) taxonomic review for each mammal order from 1990–2025. The bars indicate the proportion of the total number of species described per order over this period. The different shades of gray indicate the proportion of these descriptions involving one, two, and three or more countries.

3.3. Determinants of comprehensiveness in mammal descriptions

GLMs explained 2–30% of the variation in taxonomic comprehensiveness metrics (Fig. 5). Across all four proxies tested, the number of taxa compared emerged as the best-explained one for descriptions in all mammals and rodents, whereas the number of specimens was the most explained metric in non-bat & non-rodent mammals and bats. We did not find phylogenetic structure in GLM residuals, as indicated by phylogenetic correlograms (Fig. S5).

We found the year of description, number of species/genus, and absolute latitude among the most important predictors of species description comprehensiveness. Recent descriptions used more evidence, referred specimens, compared taxa, and (to some extent) more pages across all taxonomic mammalian orders and non-bat & non-rodent mammals, although such effect was less pronounced in bats (Fig. 5). Species-rich genera

correlated with more specimens and compared taxa overall, except for bats specimens.

Our results revealed a latitudinal gradient, with the number of evidence lines decreasing

in mammal descriptions from tropical regions. We also observed fewer specimens in

descriptions of non-rodent and non-bat species towards the Tropics. However, more taxa

are compared in the Tropics for rodents.

Figure 5. Influence of putative drivers on proxies of mammal species description comprehensiveness. Standardized coefficients from the generalized linear models fit to all species and for bats and rodents separately are shown. Horizontal lines denote the 95% confidence intervals around each coefficient. Predictors with significant effect sizes (p-value < 0.05) are those whose horizontal lines do not cross 0. Explained variation is presented (Model R^2) as Nagelkerke's pseudo-R^2 for negative binomial models (Number of taxa compared and Number of specimens) and as the coefficient of determination (R^2) for the Gaussian model (Number of evidence and Number of pages).

Other predictors affected only some taxonomic groups. For instance, we found

fewer evidence lines and pages in descriptions of small-sized mammals that are neither

rodents nor bats, and all mammals when considering them together. In bats, fewer

specimens were used to describe small-sized species, but descriptions produced via

taxonomic review generally included more specimens. For non-bat and non-rodent

mammals, taxonomic reviews tended to promote descriptions with fewer pages.

The two remaining predictors, number of authors and average number of countries

per author (an indicator of internationalization), showed more contrasting trends across

taxonomic groups and proxies of description comprehensiveness. Non-rodent and non-

bat species had fewer specimens in multi-authored descriptions. In contrast, bat

descriptions with more authors included more evidence, and international collaborations

had limited influence on description comprehensiveness, as descriptions of non-rodent

and non-bat mammals with multinational authorship consistently included fewer

specimens.

4. Discussion

Most new mammal species described between 1990 and 2025 come from Tropical

regions, particularly rodents and bats. This pattern aligns with mammal discovery

hotspots (Moura and Jetz 2021; Parsons et al. 2022) and corresponds to the highest

diversity of both orders in these regions (Abreu et al. 2024; MDD 2025). Although the

rate of new mammal descriptions remains slower than in ectothermic tetrapods (Moura

et al. 2018), hundreds of species await formal descriptions (Moura and Jetz 2021; Parsons

et al. 2022), with 68.8% increase in recognized species since the 1980s (~65 new

species/year; Burgin et al. 2025). Many additions result from taxonomic splits, reflecting

taxonomic uncertainties often rooted in species descriptions of limited

comprehensiveness, likely due to the historical unavailability or accessibility of tools that

are now commonly used in taxonomic research (Sangster and Luksenburg 2015; Poulin

and Presswell 2016; Guedes et al. 2024b). We find a clear temporal progression toward

more rigorous practices in mammal descriptions, including an increased use of multiple

evidence lines, expanded sampling of referred specimens, and broader taxonomic

comparisons (Fig. 2). Our results reveal substantial heterogeneity in determinants of

species description comprehensiveness, likely associated with taxon-specific sampling

challenges.

The increase over time in comprehensiveness of mammal species descriptions has

mirrored trends observed across various taxa (Sangster and Luksenburg 2015; Poulin and

Presswell 2016; Guedes et al. 2024b). While this pattern aligns with the principles of

integrative taxonomy (Dayrat 2005; Will et al. 2005), it also shows that many taxonomists

were already applying integrative approaches before the formalization of the concept

(Valdecasas et al. 2008). The higher comprehensiveness of recent descriptions can be

attributed to technological advancements and reduced costs, which have democratized

access to analytical tools (Nakamura et al. 2023; Guedes et al. 2024b), like micro-CT

scanning, DNA sequencing, and automated platforms for taxonomic analysis (Thomer et

al. 2018; Grenié et al. 2023; e.g., Blackburn et al. 2024). Taken together, these

technological advances have accelerated species descriptions and expanded the scope and

depth of taxonomic knowledge, contributing to a more comprehensive understanding of

biodiversity. Despite improvements in access to analytical tools and resources, disparities

still persist (e.g., Abreu et al. 2025), as evidenced by a greater number of evidence lines

in mammal descriptions from temperate regions (Fig. 5), where wealthy economies

predominate relative to low latitudes.

While descriptions of new mammal species now involve more authors and

countries overall (Fig. S6), the growth in authorship size has outpaced international

collaboration (Fig. 3). Consequently, we find a lower average number of countries per

author in recent descriptions, which is also noticeable across molecular-based

descriptions. These findings may relate to novel standards in authorship criteria of recent

descriptions (Allen et al. 2019; Lin 2024). For instance, more collectors of mammal type-

specimens have authored recent descriptions (Fig. S7). This suggests that recent species

descriptions may be based on newly collected specimens obtained through projects led

by these collectors, or that there is an increasing recognition of the contributions made by

fieldwork personnel. In addition, the somewhat limited internationalization of molecular-

based descriptions corroborates DNA sequencing in taxonomic research, which is

nowadays more affordable by researchers in biodiversity-rich regions in the Global

South. For instance, the Human Genome Project (1990-2003) required 3 US$ billions,

whereas advances like next−generation sequencing reduced costs to less than $1,000 per

genome (Schmidt and Hildebrandt 2017; Wetterstrand 2025). Although this shift is

evident in the rising prevalence of molecular-based diagnoses in mammal descriptions

(Figs S4-S5), the advent of next-generation sequencing has introduced greater

methodological heterogeneity and costly approaches that are not necessarily accessible to

Global South researchers.

Among the most consistent findings of this study is the higher comprehensiveness

of descriptions made within speciose genera. On the one hand, this finding may reflect a

sampling effect, as species-rich genera necessitate more extensive within-group

comparisons as additional species accumulate over time (Fig. 2). On the other hand, alpha

taxonomy is inherently comparative, and distinguishing closely related species becomes

particularly challenging in species-rich genera (Gaston et al. 1995), where greater

phenotypic and genetic variation may require more specimens to accurately delineate

species. Indeed, we found that more speciose genera from all groups analyzed together,

or separately, involve more taxa compared. Similarly, more evidence is used for diagnosis

in rodents and non-bats & non-rodents. On the other hand, only in mammals analyzed

together, and non-bats & non-rodents have more specimens included. This pattern may

be related to the difficulty of collecting specimens from small and nocturnal, difficult-to-

sample animals, such as rodents and bats (Meiri 2018; Moura and Jetz 2021).

Species descriptions of larger mammals contain fewer evidence lines and are

shorter than those of smaller species, especially in non-flying mammals. While larger

species' morphologies are typically diagnosable without advanced imaging, smaller

species often require it for accurate identification (Ziegler and Sagorny 2021). Thus, body

size shows a negative relationship with evidence lines, contrasting to its positive

associations with species description probability and data availability (Meiri 2018; Moura

and Jetz 2021). The inverse association between body mass and description length

appears to be mediated through the number of evidence lines. That is, small species

demand more pages to document their additional evidence lines, a pattern supported by

the moderate but significant correlation between page count and evidence lines (Fig. S1).

Notably, body mass influenced specimen availability only in bats, once again reflecting

taxon-specific sampling challenges. Unlike bats, which face size-biased mist-netting

limitations, non-flying mammals (especially rodents) are more efficiently collected

through passive methods like pitfalls and baited live traps (Tobler et al. 2008), with

minimal size-related bias (Williams and Braun 1983; Bovendorp et al. 2017). These

sampling limitations result in fewer specimens of bats relative to non-flying mammals,

as evidenced by GBIF's global holdings of preserved specimens (1.16 million bats versus

3.6 million rodents and 1.89 million other non-flying mammals; GBIF 2025), although

these figures may also reflect the global distribution of known species by order (MDD

2025).

Our study highlights the significant progress in the comprehensiveness of

mammal species descriptions since the 1990s, driven by technological advancements and

the broader adoption of integrative taxonomic practices. Additionally, our findings

confirm that taxonomic practices align with the well-documented latitudinal gaps in

taxonomic resolution (Freeman and Pennell 2021; Guedes et al. 2025), with implications

that extend beyond mammalogy. The core question of how taxonomic effort shapes our

understanding of biodiversity is universal. The pattern we uncover in mammals, which

are driven by socioeconomic inequalities, taxon-specific sampling biases, and the broader

taxonomic impediment (i.e., systemic barriers limiting both taxonomic output and end-

user reliability; Ebach et al. 2011; Moura et al. 2025), exemplifies the systemic issues

faced in biodiversity systematics. Bats, rodents, and potentially other small-sized Tropical

organisms face compounded challenges, from methodological and sampling constraints

to underrepresentation in either molecular diagnoses or revisionary studies (Guedes et al.

2024a; Abreu et al. 2025). Addressing these challenges requires targeted investments in

building local capacity and inclusive collaboration frameworks (Nakamura et al. 2023;

Carneiro et al. 2025; Moura et al. 2025). In the absence of such interventions, the

interoperability of taxonomic data risks being delayed (Sandall et al. 2023), ultimately

undermining research on conservation, systematics, and evolution.

Acknowledgments

We thank Pedro C. Estrela for valuable comments on a previous version of our

manuscript. JJMG thanks Conselho Nacional de Desenvolvimento Científico e

Tecnológico (CNPq, proc. 381394/2024-7) and Coordenação de Aperfeiçoamento de

Pessoal de Nível Superior (CAPES, proc. 88887.478942/2020-00) for research grants.

We also acknowledge support from Rio de Janeiro Foundation (FAPERJ) for grant to

ARB (E-26/203.454/2023), and São Paulo Research Foundation (FAPESP) for grants to

MTM (#2023/14506-5), GMM (#2023/16169-6), GLD (#2022/14674-2), and MRM

(#2021/11840-6 and #2022/12231-6). This work is also a contribution of the National

Institute of Science and Technology (INCT) in Ecology, Evolution, and Biodiversity

Conservation funded by CNPq (grant 465610/2014-5/ 409197/2024-6).

Data Sharing

The R-script and raw dataset used in the analyses of this study is available at Zenodo

Digital Repository (Moroti et al. 2025) as well as on GitHub (https://github.com/jhonny-

guedes/mammal_desc_trends)

Conflict of Interest Statement

The authors declare they have no conflict of interest.

References

Abreu, E.C.T., da Silva, E.L. and Moura, M.R. 2025. Geopolitical impacts on the

description of new terrestrial mollusc species. *Proceedings of the Royal Society B:*

Biological Sciences 292(2055). Available at:

https://royalsocietypublishing.org/doi/10.1098/rspb.2025.1428.

Abreu, E.F. et al. 2024. Lista de Mamíferos do Brasil (2024-1) [Data set]. doi:

https://doi.org/10.5281/zenodo.14536925.

Allen, L., O'Connell, A. and Kiermer, V. 2019. How can we ensure visibility and

diversity in research contributions? How the Contributor Role Taxonomy (CRediT) is

helping the shift from authorship to contributorship. *Learned Publishing* 32(1), pp. 71–

74. Available at: https://onlinelibrary.wiley.com/doi/10.1002/leap.1210.

Alroy, J. 2002. How many named species are valid? *Proceedings of the National*

Academy of Sciences of the United States of America 99(6), pp. 3706–3711. doi:

10.1073/pnas.062691099.

Baselga, A., Lobo, J.M., Hortal, J., Jiménez-Valverde, A. and Gómez, J.F. 2008.

Assessing alpha and beta taxonomy in eupelmid wasps: Determinants of the probability

of describing good species and synonyms. *Journal of Zoological Systematics and*

Evolutionary Research 48(1), pp. 40–49. doi: 10.1111/j.1439-0469.2009.00523.x.

Benton, M.J. 2008. Fossil quality and naming dinosaurs. *Biology Letters* 4(6), pp. 729–

732. doi: 10.1098/rsbl.2008.0402.

Bertrand, Y. and Härlin, M. 2006. Stability and Universality in the Application of

Taxon Names in Phylogenetic Nomenclature. Lee, M. ed. *Systematic Biology* 55(5), pp.

848–858. Available at: https://academic.oup.com/sysbio/article/55/5/848/1668137.

Blackburn, D.C. et al. 2024. Increasing the impact of vertebrate scientific collections

through 3D imaging: The openVertebrate (oVert) Thematic Collections Network.

BioScience 74(3), pp. 169–186. Available at:

https://academic.oup.com/bioscience/article/74/3/169/7615104.

Bolker, B. et al. 2020. phylobase: Base Package for Phylogenetic Structures and

Comparative Data. Available at: https://cran.r-project.org/package=phylobase.

Bonferroni, C. 1936. Teoria statistica delle classi e calcolo delle probabilita.

Pubblicazioni del R istituto superiore di scienze economiche e commericiali di firenze

8, pp. 3–62.

Bovendorp, R.S., MCCleery, R.A. and Galetti, M. 2017. Optimising sampling methods

for small mammal communities in Neotropical rainforests. *Mammal Review* 47(2), pp.

148–158. Available at: https://onlinelibrary.wiley.com/doi/10.1111/mam.12088.

Braby, M.F., Hsu, Y.-F. and Lamas, G. 2024. How to describe a new species in zoology

and avoid mistakes. *Zoological Journal of the Linnean Society*. doi:

10.1093/zoolinnean/zlae043.

Bremer, K., Bremer, B., Karis, P. and Källersjö, M. 1990. Time for change in

taxonomy. *Nature* 434, pp. 202–202.

Burgin, C.J. et al. 2025. How many mammal species are there now? Updates and trends

in taxonomic, nomenclatural, and geographic knowledge. Percequillo, A. ed. *Journal of*

Mammalogy. Available at: https://academic.oup.com/jmammal/advance-

article/doi/10.1093/jmammal/gyaf047/8253815.

Burgin, C.J., Colella, J.P., Kahn, P.L. and Upham, N.S. 2018. How many species of

mammals are there? *Journal of Mammalogy* 99(1), pp. 1–14. doi:

10.1093/jmammal/gyx147.

Burnham, K.P. and Anderson, D.A. 2002. *Model Selection and Multimodel Inference: A*

Practical Information-Theoretic Approach. 2nd ed. Burnham, K. P. and Anderson, D.

A. eds. Fort Collins, CO: Springer.

Carneiro, C.M. et al. 2025. Toward a Global Science of Conservation Genomics:

Coldspots in Genomic Resources Highlight a Need for Equitable Collaborations and

Capacity Building. *Molecular Ecology*. Available at:

https://onlinelibrary.wiley.com/doi/10.1111/mec.17729.

Chek, A.A., Austin, J.D. and Lougheed, S.C. 2003. Why is there a tropical–temperate

disparity in the genetic diversity and taxonomy of species? *Evolutionary Ecology*

Research 5, pp. 69–77.

Dayrat, B. 2005. Towards integrative taxonomy. *Biological Journal of the Linnean*

Society 85(3), pp. 407–415. doi: 10.1111/j.1095-8312.2005.00503.x.

Dinerstein, E. et al. 2017. An Ecoregion-Based Approach to Protecting Half the

Terrestrial Realm. *BioScience* 67(6), pp. 534–545. doi: 10.1093/biosci/bix014.

Diniz Filho, J.A.F. et al. 2023. Macroecological links between the Linnean, Wallacean,

and Darwinian shortfalls. *Frontiers of Biogeography* 15(2), pp. 1–11. doi:

10.21425/F5FBG59566.

Dominguez, E. and Wheeler, Q.D. 1997. Forum – Taxonomic Stability is Ignorance.

Cladistics 13(4), pp. 367–372. Available at:

https://onlinelibrary.wiley.com/doi/10.1111/j.1096-0031.1997.tb00325.x.

Dubois, A. 2007. Naming taxa from cladograms: some confusions, misleading

statements, and necessary clarifications. *Cladistics* 23(4), pp. 390–402. Available at:

https://onlinelibrary.wiley.com/doi/10.1111/j.1096-0031.2007.00151.x.

Ebach, M.C., Valdecasas, A.G. and Wheeler, Q.D. 2011. Impediments to taxonomy and

users of taxonomy: accessibility and impact evaluation. *Cladistics* 27(5), pp. 550–557.

Available at: https://onlinelibrary.wiley.com/doi/10.1111/j.1096-0031.2011.00348.x.

Feijó, A. and Brandão, M.V. 2022. Taxonomy as the first step towards conservation: an

appraisal on the taxonomy of medium- and large-sized Neotropical mammals in the 21st

century. *Zoologia (Curitiba)* 39, p. e22007. doi: 10.1590/s1984-4689.v39.e22007.

Freeman, B.G. and Pennell, M.W. 2021. The latitudinal taxonomy gradient. *Trends in*

Ecology and Evolution 36(9), pp. 778–786. doi: 10.1016/j.tree.2021.05.003.

Gaston, K.J., Blackburn, T.M. and Loder, N. 1995. Which species are described first?:

the case of North American butterflies. *Biodiversity and Conservation* 4, pp. 119–127.

GBIF. 2025. *GBIF Home Page*. Available at: https://www.gbif.org [Accessed: 30 June

2025].

Gippoliti, S., Farina, S. and Andreone, F. 2024. Lost species, neglected taxonomy, and

the role of natural history collections and synonymization in the identification of the

World's forgotten biodiversity. *Nature Conservation* 56, pp. 119–126. Available at:

https://natureconservation.pensoft.net/article/132036/.

Grenié, M., Berti, E., Carvajal-Quintero, J., Dädlow, G.M.L., Sagouis, A. and Winter,

M. 2023. Harmonizing taxon names in biodiversity data: A review of tools, databases

and best practices. *Methods in Ecology and Evolution* 14(1), pp. 12–25. Available at:

https://besjournals.onlinelibrary.wiley.com/doi/10.1111/2041-210X.13802.

Guedes, J.J.M., Feio, R.N., Meiri, S. and Moura, M.R. 2020. Identifying factors that

boost species discoveries of global reptiles. *Zoological Journal of the Linnean Society*

190(4), pp. 1274–1284. doi: 10.1093/zoolinnean/zlaa029.

Guedes, J.J.M., Gomes De Lima, H.V., Mendonça, L.R., Chen-Zhao, R., Diniz-Filho,

J.A.F. and Moura, M.R. 2024a. Temporal trends in global reptile species descriptions

over three decades. *Systematics and Biodiversity* 22(1), p. 2419832. doi:

10.1080/14772000.2024.2419832.

Guedes, J.J.M., Moura, M.R., Jardim, L. and Diniz-Filho, J.A.F. 2025. Global Patterns

of Taxonomic Uncertainty and its Impacts on Biodiversity Research. *Systematic Biology*

In Press. Available at: https://academic.oup.com/sysbio/advance-

article/doi/10.1093/sysbio/syaf010/8016068.

Guedes, J.J.M., Valentim, H., Mendonça, L.R., Chen-Zhao, R., Diniz-Filho, J.A.F. and

Moura, M.R. 2024b. Temporal trends in global reptile descriptions over three decades

[Dataset]. *Zenodo*. doi: https://doi.org/10.5281/zenodo.13485016.

Hamdan, B. et al. 2024. When a name changes everything: taxonomy and conservation

of the Atlantic bushmaster (Lachesis Daudin, 1803) (Serpentes: Viperidae: Crotalinae).

Systematics and Biodiversity 22(1). doi: 10.1080/14772000.2024.2366215.

Hendrixson, B.E. and Bond, J.E. 2005. Testing species boundaries in the Antrodiaetus

unicolor complex (Araneae: Mygalomorphae: Antrodiaetidae): "Paraphyly" and cryptic

diversity. *Molecular Phylogenetics and Evolution* 36(2), pp. 405–416. Available at:

https://linkinghub.elsevier.com/retrieve/pii/S1055790305000382.

Isaac, N.J.B., Mallet, J. and Mace, G.M. 2004. Taxonomic inflation: its influence on

macroecology and conservation. *Trends in Ecology & Evolution* 19(9), pp. 464–469.

doi: 10.1016/j.tree.2004.06.004.

Kassambara, A. 2023. rstatix: Pipe-Friendly Framework for Basic Statistical Tests, R

package version 0.7.2.

Keck, F., Rimet, F., Bouchez, A. and Franc, A. 2016. Phylosignal: An R package to

measure, test, and explore the phylogenetic signal. *Ecology and Evolution* 6(9), pp.

2774–2780. doi: 10.1002/ece3.2051.

Kutner, M.H., Nachtsheim, C.J., Neter, J. and Li, W. 2005. *Applied Linear Statistical*

Models. 5th ed. New York: McGraw-Hill Irwin.

Lessa, T., Stropp, J., Hortal, J. and Ladle, R.J. 2024. How taxonomic change influences

forecasts of the Linnean shortfall (and what we can do about it)? *Journal of*

Biogeography. Available at: https://onlinelibrary.wiley.com/doi/10.1111/jbi.14829.

Lin, Z. 2024. Modernizing authorship criteria and transparency practices to facilitate

open and equitable team science. *Accountability in Research*, pp. 1–24. Available at:

https://www.tandfonline.com/doi/full/10.1080/08989621.2024.2405041.

Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P. and Makowski, D. 2021.

performance: An R Package for Assessment, Comparison and Testing of Statistical

Models. *Journal of Open Source Software* 6(60), p. 3139. doi: 10.21105/joss.03139.

Mahony, S., Kamei, R.G., Brown, R.M. and Chan, K.O. 2024. Unnecessary splitting of

genus-level clades reduces taxonomic stability in amphibians. *Vertebrate Zoology* 74,

pp. 249–277. doi: 10.3897/vz.74.e114285.

Mansfield, E.R. and Helms, B.P. 1982. Detecting Multicollinearity. *The American*

Statistician 36(3), pp. 158–160. doi: 10.1080/00031305.1982.10482818.

MDD. 2025. *Mammal Diversity Database, version 2.1*. Available at:

https://www.mammaldiversity.org/.

Meiri, S. 2018. Traits of lizards of the world: Variation around a successful

evolutionary design. *Global Ecology and Biogeography* 27(10), pp. 1168–1172. doi:

10.1111/geb.12773.

Moroti, M. de T., Guedes., J.J.M., Missio, G.M., Diegues, G.L., Bezerra, A.M.R. and

Moura, M.R. 2025. Historical shifts, geographic biases, and biological constraints shape

mammal species discovery [Dataset]. *Zenodo*. doi: 10.5281/zenodo.17279492.

Moura, M., Carvalho, R., Ceron, K., Guedes, J., Moroti, M. and Nakamura, G. 2025.

Local expertise anchors biodiversity documentation, but geopolitical power drives

parachute discovery. Available at: https://www.researchsquare.com/article/rs-

7724270/v1.

Moura, M.R. et al. 2024. A phylogeny-informed characterisation of global tetrapod

traits addresses data gaps and biases. *PLoS Biology* 22(7), p. e3002658. doi:

10.1371/journal.pbio.3002658.

Moura, M.R., Costa, H.C., Peixoto, M.A., Carvalho, A.L.G., Santana, D.J. and

Vasconcelos, H.L. 2018. Geographical and socioeconomic determinants of species

discovery trends in a biodiversity hotspot. *Biological Conservation* 220, pp. 237–244.

doi: 10.1016/j.biocon.2018.01.024.

Moura, M.R. and Jetz, W. 2021. Shortfalls and opportunities in terrestrial vertebrate

species discovery. *Nature Ecology & Evolution* 5(5), pp. 631–639. doi:

10.1038/s41559-021-01411-5.

Mutanen, M. et al. 2016. Species-Level Para- and Polyphyly in DNA Barcode Gene

Trees: Strong Operational Bias in European Lepidoptera. *Systematic Biology* 65(6), pp.

1024–1040. Available at: https://academic.oup.com/sysbio/article-

lookup/doi/10.1093/sysbio/syw044.

Nakamura, G., Soares, B.E., Pillar, V.D., Diniz-Filho, J.A.F. and Duarte, L. 2023. Three

pathways to better recognize the expertise of Global South researchers. *npj Biodiversity*

2(1), p. 17. doi: 10.1038/s44185-023-00021-7.

Padial, J.M. and De la Riva, I. 2021. A paradigm shift in our view of species drives

current trends in biological classification. *Biological Reviews* 96(2), pp. 731–751. doi:

10.1111/brv.12676.

Pante, E., Schoelinck, C. and Puillandre, N. 2015. From integrative taxonomy to species

description: One step beyond. *Systematic Biology* 64(1), pp. 152–160. doi:

10.1093/sysbio/syu083.

Parsons, D.J., Pelletier, T.A., Wieringa, J.G., Duckett, D.J. and Carstens, B.C. 2022.

Analysis of biodiversity data suggests that mammal species are hidden in predictable

places. *Proceedings of the National Academy of Sciences* 119(14). doi:

10.1073/pnas.2103400119.

Poulin, R. and Presswell, B. 2016. Taxonomic Quality of Species Descriptions Varies

over Time and with the Number of Authors, but Unevenly among Parasitic Taxa.

Systematic Biology 65(6), pp. 1107–1116. doi: 10.1093/sysbio/syw053.

Pyke, G.H. 2014. Evaluating the quality of taxonomic publications: A simple alternative

to citations and effort. *BioScience* 64(11), pp. 961–962. doi: 10.1093/biosci/biu154.

De Queiroz, K. 2007. Species Concepts and Species Delimitation. *Systematic Biology*

56(6), pp. 879–886. Available at:

https://academic.oup.com/sysbio/article/56/6/879/1653163.

R Core Team. 2023. R: A Language and Environment for Statistical Computing. *R*

Foundation for Statistical Computing. Available at: https://www.r-project.org/.

Raposo, M.A., Stopiglia, R., Brito, G.R.R., Bockmann, F.A., Kirwan, G.M., Gayon, J.

and Dubois, A. 2017. What really hampers taxonomy and conservation? A riposte to

Garnett and Christidis (2017). *Zootaxa* 4317(1), pp. 179–184.

Revell, L.J. 2010. Phylogenetic signal and linear regression on species data. *Methods in*

Ecology and Evolution 1, pp. 319–329. doi: 10.1111/j.2041-210X.2010.00044.x.

Sandall, E.L. et al. 2023. A globally integrated structure of taxonomy to support

biodiversity science and conservation. *Trends in Ecology & Evolution* 38(12), pp.

1143–1153. Available at:

https://linkinghub.elsevier.com/retrieve/pii/S016953472300215X.

Sangster, G. and Luksenburg, J.A. 2015. Declining rates of species described per

taxonomist: Slowdown of progress or a side-effect of improved quality in taxonomy?

Systematic Biology 64(1), pp. 144–151. doi: 10.1093/sysbio/syu069.

Schmidt, B. and Hildebrandt, A. 2017. Next-generation sequencing: big data meets high

performance computing. *Drug Discovery Today* 22(4), pp. 712–717. Available at:

https://linkinghub.elsevier.com/retrieve/pii/S1359644617300582.

Solow, A.R., Mound, L.A. and Gaston, K.J. 1995. Estimating the Rate of Synonymy.

Systematic Biology 44(1), pp. 93–96.

Streicher, J.W., Wiens, J.J., Jocqué, M., García-Vázquez, U.O. and Smith, E.N. 2023.

Molecular phylogenetic analyses reveal both underestimation and overestimation of

species diversity in northern rain frogs (Craugastor). *Journal of Vertebrate Biology*

72(23072). doi: 10.25225/jvb.23072.

Sullivan, K.A. et al. 2023. Building a community-based taxonomic resource for

digitization of parasites and their hosts. Lozier, J. ed. *Insect Systematics and Diversity*

7(6). Available at:

https://academic.oup.com/isd/article/doi/10.1093/isd/ixad023/7439483.

Thomer, A.K., Twidale, M.B. and Yoder, M.J. 2018. Transforming Taxonomic

Interfaces. *Proceedings of the ACM on Human-Computer Interaction* 2(CSCW), pp. 1–

23. Available at: https://dl.acm.org/doi/10.1145/3274442.

Tobler, M.W., Carrillo-Percastegui, S.E., Leite Pitman, R., Mares, R. and Powell, G.

2008. An evaluation of camera traps for inventorying large- and medium-sized

terrestrial rainforest mammals. *Animal Conservation* 11(3), pp. 169–178. doi:

10.1111/j.1469-1795.2008.00169.x.

Upham, N.S., Esselstyn, J.A. and Jetz, W. 2019. Inferring the mammal tree: Species-

level sets of phylogenies for questions in ecology, evolution, and conservation. *PLoS*

Biology 17(12), pp. 1–44. doi: 10.1371/journal.pbio.3000494.

Valdecasas, A.G. 2011. An index to evaluate the quality of taxonomic publications.

Zootaxa 62(2925), pp. 57–62. doi: 10.11646/zootaxa.2925.1.6.

Valdecasas, A.G., Castroviejo, S. and Marcus, L.F. 2000. Reliance on the citation index

undermines the study of biodiversity. *Nature* 403(6771), p. 698. doi:

10.1038/35001751.

Valdecasas, A.G., Williams, D. and Wheeler, Q.D. 2008. "Integrative taxonomy" then

and now: A response to Dayrat (2005). *Biological Journal of the Linnean Society* 93(1),

pp. 211–216. doi: 10.1111/j.1095-8312.2007.00919.x.

Venables, W.N. and Ripley, B.D. 2002. *Modern Applied Statistics with S.* Fourth Edi.

New York: Springer.

Weksler, M. et al. 2025. A comprehensive survey of mammal collections and genetic

resources in South America: challenges and directions. *Biological Journal of the*

Linnean Society 146(1). Available at:

https://academic.oup.com/biolinnean/article/doi/10.1093/biolinnean/blaf069/8253696.

Wetterstrand, K.A. 2025. *DNA Sequencing Costs: Data from the NHGRI Genome*

Sequencing Program (GSP). Available at: www.genome.gov/sequencingcostsdata.

Will, K.W., Mishler, B.D. and Wheeler, Q.D. 2005. The perils of DNA barcoding and

the need for integrative taxonomy. *Systematic Biology* 54(5), pp. 844–851. doi:

10.1080/10635150500354878.

Williams, D.F. and Braun, S.E. 1983. Comparison of Pitfall and Conventional Traps for

Sampling Small Mammal Populations. *The Journal of Wildlife Management* 47(3), p.

841. Available at: https://www.jstor.org/stable/3808622?origin=crossref.

Wüster, W., Thomson, S.A., O'shea, M. and Kaiser, H. 2021. Confronting taxonomic

vandalism in biology: conscientious community self-organization can preserve

nomenclatural stability. *Biological Journal of the Linnean Society* 133(3), pp. 645–670.

Available at: https://academic.oup.com/biolinnean/article/133/3/645/6240088.

Ziegler, A. and Sagorny, C. 2021. Holistic description of new deep sea megafauna

(Cephalopoda: Cirrata) using a minimally invasive approach. *BMC Biology* 19(1), p. 81.

Available at: https://bmcbiol.biomedcentral.com/articles/10.1186/s12915-021-01000-9.

Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A. and Smith, G.M. 2009. *Statistics*

for Biology and Health. Springer.

722