From cells to organism – how natural selection causes metabolic scaling

Peter F. Pelz[1]

[1]Chair of Fluid Systems, Technische Universität Darmstadt, Otto-Bernd Straße 2, Darmstadt, 64287, Hessen, Germany.

Contributing authors: peter.pelz@tu-darmstadt.de;

Abstract

The cell is the power station of life. Surprisingly, to date there is still no metabolic scaling theory that links cellular respiration to the organismal metabolism and predicts the 'mouse-to-elephant curve', also known as Kleiber's law, in an approach that is consistent with physicochemical principles. This paper shows that for a consistent model, the novel concept of the optimised Metabolic Module (MM) is the missing link between cell and organism. It is shown how evolutionary selection under resource scarcity optimises the MM towards (a) light-weight design and (b) resource efficiency. Thus, Darwin's evolution by natural selection is simulated by model-based optimisation. The final model presented is complete (for the entire mass range of the organism of different taxonomic classes), concise (it uses only five scale-invariant physicochemical constants), clear (it predicts all metabolic rates within the uncertainty range of a scale model observed in measurements) and consistent with Murray's law of capillary blood flow and cell metabolism. The model features observed asymptotes for both 'small' protists and 'large' endotherms. It predicts the mass-dependent metabolic rate of protists, planarians, ectotherms and endotherms with the usual uncertainty of any scaling theory.

1 Introduction

The cellular architecture of life

The idea of a cellular architecture of life is common. In this image the growth of the organism is achieved by the multiplication of scale-invariant cells, and the metabolic rate of the organism results from the multiplication of cells that are regarded as

metabolic power stations. To account for the chemical nature of cellular respiration, the chemical reaction rate k is the most relevant physicochemical parameter. In 'small' organisms, the metabolic rate is proportional to the reaction rate k and the body mass m. This is to be expected in a complete cellular architecture of life. However, the situation is different for 'large' animals: their metabolism must be proportional to $k^{3/8}$ and $m^{3/4}$, as will be shown in this paper.

'Small' organisms are protists such as plankton. 'Large' organisms include ectotherms (cold-blooded organisms) such as reptiles and amphibians and endotherms (warm-blooded organisms) such as birds and mammals and most planarians (flatworms) as their typical size is above the diffusion length. As will be shown, the diffusion length is the relevant measure for distinguishing between 'large' and 'small', a physiological parameter determined by the diffusion coefficient and the reaction rate of cell metabolism.

The metabolic rate of 'small' organisms

In the following, we will first look at the metabolism of the 'small' organism. The resulting scaling equation is well known. Nevertheless, the derivation is valuable because some of the arguments are used below to derive a general metabolic scaling theory that is valid for 'small' and 'large' organisms. As a reminder, oxygen molecules are transported through the cell tissue by diffusion, resulting from a concentration field c limited by the available oxygen concentration: $c \leq c_{\text{avail}}$. In compliance with Fick's first law of diffusion, the transport speed of the molecules is proportional to their molecular diffusion coefficient $\mathcal{D}$ in the cell tissue multiplied by the gradient of the concentration field $|\nabla c| < c_{\text{avail}}/\delta$. The longest transport path for molecules is equal to the thickness δ of the cell tissue. Simple organisms such as plankton have a tissue thickness that must be much smaller than the diffusion length, $\delta_{\text{diff}} := (\mathcal{D}/k)^{1/2} \gg \delta$, as only in these 'small' organisms does the diffusive transport of oxygen always meet the cell's need per unit of time $\dot{c}$. This rate is limited by the reaction rate and the available oxygen concentration: $-\dot{c} = k\,c \leq k\,c_{\text{avail}}$. Therefore, the molar flow rate for 'small' organism with the volume V is given by the mass balance $J = -\int_V \dot{c}\,\mathrm{d}V \sim k\,c_{\text{avail}}V$, i.e. the supply, the left-hand side, corresponds to the need, the right-hand side. With the abbreviation $Q := J/c_{\text{avail}}$ the kinematic character of this mass balance becomes obvious, i.e. $Q = k\,V$. With the energy density $e = c_{\text{avail}}\,|\Delta G^\circ|$ and the mass density $\varrho = m/V$ of the cell tissue, the metabolic rate, measured as heat production rate, can be derived:

$$\dot{E} = k\,\frac{e}{\varrho}\,m. \tag{1}$$

Here $|\Delta G^\circ|$ is the molar Gibbs free energy for the exothermic oxidation of nutrients within the cells.

The given derivation of the scaling equation $\dot{E} \propto m$ is consistent with the idea of a cellular architecture of life: the cell is the power station of life. Equation (1) is therefore nothing more than a summation of the power contributions of all the cells of a small organism.

Diffusion limit on growth for 'small' organisms

The idea of a cellular architecture of life has a limitation: for an organism with a mass m far above the diffusion limit on growth, i.e. $m \gg \varrho\, \delta_{\mathrm{diff}}^3 = \varrho\, v_{\mathrm{diff}} = m_{\mathrm{diff}}$, the diffusive supply with oxygen is insufficient to prevent cell death. As shown in this paper, with increasing mass of the organism, this leads to a change from the 'diffusion asymptote' $\dot{E} \propto m$ for $m \ll m_{\mathrm{diff}}$ to the 'convection asymptote' $\dot{E} \propto m^{3/4}$ for $m \gg m_{\mathrm{diff}}$. The latter asymptote is known as Kleiber's law [1] or as the 'mouse-to-elephant curve', see figs. 2 and 3.

To date, there is (i) no general metabolic theory that applies to all organisms, from a plankton with a mass of only 0.01 picogram to a blue whale with a mass of more than 100 tonnes. This general theory must include both asymptotes mentioned above; there is (ii) no consistent theory linking cell metabolism to organismal metabolism; and there is (iii) no theory that is consistent with the physiology of blood flow in capillaries [2–4].

The theory presented here fills this gap. It links cell metabolism and the metabolism of the organism via the Metabolic Module (MM), which has been optimised to the size of the animal through natural selection. The concept of a scalable MM is presented here for the first time. The results might be useful for model-based drugs and tissues design.

Darwin's evolution by natural selection is simulated as model-based optimisation. The solution of the optimisation contributes to the understanding of the metabolism of organisms; the scaling of the metabolic rate resulting from evolution under resource scarcity is explained axiomatically. This new approach is discussed in the following.

2 From cellular to optimised modular architecture of life

The idea of an optimised modular architecture of life goes further than the purely cellular picture, as the theory, its asymptotes and their validation will show below.

Every organ and every muscle is supplied with nutrients and oxygen by a capillary network consisting of a large number of modules, see fig. 1. Others [2, 3] consider the capillary of these modules to be scale-invariant like cells, which is in contradiction to the measurements of capillary bundles [5, 6] and conflicts with Murray's law of capillary flow [7]. In contrast, the concept of an optimised Metabolic Module (MM) is presented here for the first time. The MM is optimised to the size of the organism through natural selection. Understanding the scaling of the size and the shaping of the slenderness of the MM by Darwin's evolution by selection is key to linking cell metabolism and organismal metabolism.

Sequential growth strategies

In the new perspective presented here, nature applies two sequential evolutionary growth strategies, which are sketched as vertical and horizontal directions in fig. 1. The horizontal direction shows volume growth by scaling and shaping the MM. The vertical direction shows consecutive growth by multiplication. For cell tissues and organisms

Fig. 1 Growth of the organism through first scaling and shaping of the Metabolic Module (MM) and second growth through multiplication of the MMs

that are smaller than the diffusion size limit, growth is achieved by cell multiplication. For organisms above the size limit, the MMs are the optimised modules of the organism, which are scaled and shaped in the first evolutionary step and multiplied by a factor $n \geq 1$ in the second evolutionary step. This means that volume v of the module results in the volume $V = n\,v$ of the organism and the blood volume flow Q of the organism is given by the volume flow q through the capillary of the module as $Q = n\,q$.

Figure 1 shows that the MM is a cylindrical tissue structure of length l, which is supplied with nutrients and oxygen by a concentric capillary of diameter d. The tissue thickness is δ and the effective outer diameter is correspondingly $d + 2\delta$. The nutrients and oxygen are first transported by convection in the capillary and finally by diffusive transport within the cell tissue in order to meet the needs of cell metabolism.

Selection pressures caused by resource scarcity

Scarcity of resources and the struggle for survival leads to optimisation pressure. In fact, there are two optimisation pressures that scale and shape the MM, namely (a) minimum mass of the MM for a given power (lightweight design) and (b) optimal use of available nutrients and oxygen (resource efficiency). Both are achieved through natural selection, which is simulated here using model-based optimisation.

The principle of lightweight design is also known as the 'physiological principle of minimum work' [7]. Lightweight design is essential for high acceleration in animal locomotion. But it is also crucial for two other reasons: firstly, the mass-to-power ratio should be minimised when energy reserves are limited, e.g. during the breeding season. Secondly, a minimum power-related mass is required to limit the energy expenditure for cell renewal.

(a) lightweight design: With minimal blood mass and minimal dissipation, the optimal ratio between the blood volume flow q through the capillary and its diameter d is given by Murray's law [5, 7] $q = d_{\text{opt}}^3 / (32\,\tau)$. It results from a constrained optimisation problem. The design variable is the capillary diameter d, the constraint is the given volume flow q, and the objective function to be minimised is the sum of the blood mass and the energy loss due to capillary flow.

The given form of Murray's law with the physiological constant τ in the denominator given here is new; it shows how nature realises Murray's law: Adaptive growth is often triggered by the prevailing shear stress. If Murray's law applies, the blood cells in capillary flow experience the same wall shear stress and wall shear velocity τ^{-1} regardless of the capillary diameter, the flow velocity, the size of the organism and its taxonomic classification.

Murray's law therefore provides an equilibrium state of growth in a cybernetic loop. Like any other closed control loop, it contains a measuring element, a controller and a controlled variable. The latter is the capillary diameter. The measuring elements are the cells forming the capillary wall and measuring wall shear rate $\dot{\gamma}$. If $\dot{\gamma}$ deviates from τ^{-1} in a way that is unacceptable for the organism, the organism responds with a change in diameter. This adaptation happens at the cellular level until the wall shear rate is approximately equal to τ^{-1} [8–11].

As shown in the section on model validation, τ^{-1} is a physiological parameter for blood flow that is known in connection with primary haemostasis. For this reason, too, it can be summarised that Murray's law forms a solid basis for the theory presented here.

(b) resource efficiency: In the following, the convective-diffusive transport in capillary and tissue as well as the exothermic reaction in the cell, the cell metabolism is considered. Let us call the oxygen concentration at the inlet of the MM $c_1 = c_{\text{basal}} + c_{\text{avail}}$ and c_2 at the outlet of the capillary. The capillary is too short for $c_2 \approx c_1$, as the allocated and hence available oxygen concentration c_{avail} is obviously not utilised. Conversely, the capillary is too long for $c_2 \approx 0$, as there is then a risk of cell death in the exit area of the MM. The resources are only utilised efficiently if c_2 assumes the basal concentration c_{basal}: $(c_1 - c_2)_{\text{opt}} = c_{\text{avail}}$.

Mass balance for the Metabolic Module

With the two optimisation results for (a) lightweight construction, i.e. $q = d_{\text{opt}}^3 / (32\,\tau)$, and (b) resource efficiency, i.e. $c_{\text{avail}} = (c_1 - c_2)_{\text{opt}}$, the mass balances for the capillary, for the blood-tissue interface and for the tissue itself can be specified. The integral forms are used in the scaling theory presented here. At the end of this article, we provide an outlook on how the corresponding conservation equations can be solved in differential form as a boundary value problem. However, this has no influence on the model concept, which is why the concise integral model is completely sufficient here.

The mass balance for the capillary for resource efficient operation gives the convective molar flow rate:

$$J = q\,(c_1 - c_2)_{\text{opt}} = q\,c_{\text{avail}}.$$

For a resource efficient organism, the convective molar flow rate equals the diffusive molar flow rate into the cell tissue across the blood-tissue interface $S = \pi\,d\,l$ with the radial normal vector $\vec{n} = \vec{e}_r$:

$$J = \int\limits_{\text{interface}} \vec{j} \cdot \vec{n}\,\mathrm{d}S = \int\limits_{S=\pi dl} -\mathcal{D}\,\nabla c \cdot \vec{e}_r\,\mathrm{d}S \approx D\,c_{\text{avail}}\,\pi\,l\,\frac{d}{\delta}.$$

Fick's first law of diffusion is applied here, in which the vector of molar diffusion flux vector is proportional to the concentration gradient: $\vec{j} = -\mathcal{D}\,\nabla c$. The radial component of the concentration gradient $\nabla c \cdot \vec{e}_r$ is inhomogeneous. It has a maximum at the capillary inlet and a minimum at the outlet. The right-hand side of the upper equation is therefore an average value for the MM. This is captured by the dispersion coefficient of the cell tissue D. This is a parameter that must be much smaller than the diffusion coefficient $D/\mathcal{D} \ll 1$ [12, 13]. The factor $D/\mathcal{D}$ can be determined either by solving the reaction-diffusion equation for the cell tissue, as explained in the outlook section of this article, or by using measured metabolic rates. The latter is done in the following section.

The model is closed with the cell metabolism. At each point in the cell tissue, the temporal change in concentration $\dot{c}$ due to the exothermic reaction is balanced by the diffusive supply of nutrients and oxygen. The integral form of the reaction-diffusion equation for the volume of the cell tissue v_{tiss} is the same as for 'small' animals of volume v, eq. (1), i.e.

$$J = -\int\limits_{\text{tissue volume}} \dot{c}\,\mathrm{d}V = \int\limits_{v_{\text{tiss}}} k\,c\,\mathrm{d}V \approx k\,c_{\text{avail}}\,v_{\text{tiss}}.$$

However, the volume of the cell tissue in a 'large' organism is only a fraction $v_{\text{tiss}}/v < 1$ of the total volume of the MM. Darwin's principle of selection for (a) lightweight design and (b) resource efficient solutions was explained above by first modelling the physiology of the MM axiomatically and then optimising the MM based on the model. The central idea is that nature scales and shapes MMs under scarcity using selection heuristics, while model-based optimisation simulates this. If the concept is viable, and this is demonstrated in the following (i) by model validation using measured metabolic rates, (ii) by independent derivation via dimensional analysis, and (ii) by successful model use for predicting the heart rate of mammals, then both strategies lead to the same functional relationship:

$$q = q(v,\,k,\,D,\,\tau), \tag{2}$$

which results from the solution of the non-linear system of the five equations:

$$v = \frac{\pi}{4}\,l\,d^2 \left(1 + 2\frac{\delta}{d}\right)^2, \quad v_{\text{tiss}} = \pi\,l\,d^2\,\frac{\delta}{d}\left(1 + \frac{\delta}{d}\right), \quad q = \frac{d^3}{32\,\tau} = D\,\pi\,l\,\frac{d}{\delta} = k\,v_{\text{tiss}}$$

For a given volume flow q, the five unknowns v, v_{tiss}, d, l, δ are derived from the system of equations, and therefore also $q = q(v,\,k,\,D,\,\tau)$.

3 Model validation

The only difference between the taxonomy classes protists, planarians, ectotherms and endotherms in the presented idea, see fig. 1, is the number n of MMs. The order of magnitude of MMs in mammals is one billion [2], which gives an upper limit for the number of MMs: $n < 10^9$. Figure 2 therefore shows the solution of the system of equations as black solid lines for $\log n = 0, 3, 8$ as parameter to distinct the set of curves. The number of MMs for the various taxonomic classes was determined based on measurement data, see fig. 3. The data obtained is consistent with the upper limit provided [2].

All other physicochemical properties, i.e. ϱ, e, k, D, τ, are determined by the tissue and blood physiology and are therefore the same for all organism, i.e. they are scale-invariant. The following typical parameters are identified on the basis of measurements and experience and used as constant model parameters in the following: The reaction rate for glucose oxidation at normal condition is $k \approx 0.1\,\mathrm{s}^{-1}$ [14]. The change in molar Gibbs free energy for glucose oxidation is $\Delta G^\circ = -2.87\,\mathrm{MJ\,mol^{-1}}$ [15] and the assumed available oxygen concentration is $c_{\mathrm{avail}} = 0.02\,\mathrm{mol\,m^{-3}}$, resulting in the energy density $e = c_{\mathrm{avail}}\,|\Delta G^\circ| = 57.4\,\mathrm{kJ\,m^{-3}}$. To prevent primary haemostasis, the required shear rate within the capillaries is $\tau^{-1} \approx 10^3\,\mathrm{s}^{-1}$ [8, 9]. The molecular coefficient of diffusion in dilute solutions is $\mathcal{D} = 10^{-9}\,\mathrm{m^2\,s^{-1}}$ [13]. The dispersion coefficient is three orders of magnitude smaller, $D = 10^{-3}\,\mathcal{D}$. As already mentioned, this takes into account the inhomogeneous concentration gradient field in the cell tissue of the MM. D is therefore an effective value for the MM.

Given the conciseness of the model, it is remarkable that the metabolic rate of all taxonomic classes and all organism sizes can be predicted (completeness) not only relatively but also absolutely within the usual limits of accuracy of a scaling model. In addition, three further arguments for the validity of the theory are presented below, namely its asymptotes, independent confirmation by dimensional analysis, and the scaling of heart rate.

4 Kleiber's law is an asymptote of Equation (2)

Figure 2 shows the metabolic rate as a function of body mass. This curve is also known as 'mouse-to-elephant curve'. Two expected asymptotes $\dot{E} \propto m$ and $\dot{E} \propto m^{3/4}$ are clearly visible.

Both special solutions are asymptotes of the given system of five non-linear equations: for $\delta/d \gg 1$ the asymptotic solution is $q = k\,v$. For $\delta/d \ll 1$, the diffusion limit on growth is reached, and the tissue thickness is limited by the diffusion length $\delta \to \delta_{\mathrm{diff}} = (D/k)^{1/2}$. The explicit form of the asymptotic solution is then

$$q = \frac{\pi}{2}\,\frac{(D\,k)^{3/8}}{(2\,\tau)^{1/4}}\,v^{3/4}.$$

This result is confirmed by an independent derivation using dimensional analysis [16]. Based on Buckingham's Pi theorem, the functional relationship $q = q(v, k, D, \tau)$

Fig. 2 Metabolic rate as a function of organism size for three different numbers of metabolic modules represented by the given theory; all given physicochemical and physiological properties for cell tissue and capillary flow are held constant and assume typical values. The model proves to be a general theory for the entire mass range. The diffusion asymptote and the convection asymptote (Kleiber's law) as asymptotic limits of the general model are clearly visible.

is equivalent to $q = (D\,k)^{3/8}\,v^{3/4}\,\tau^{-1/4}\,\mathrm{fn}(v/v_{\mathrm{diff}},\,k\tau)$. When the argument of a dimensionless function approaches zero (or infinity), it assumes a constant function value. Therefore, $\mathrm{fn}(v_{\mathrm{diff}}/v \to 0) \to \mathrm{const.}$, where the numerical value of the constant, namely $\pi/2^{5/4}$, cannot be derived from the dimensional analysis. Nevertheless, dimensional analysis also provides the result $q \propto v^{3/4}$ valid for 'large' organisms. In addition to validating the model through experiments, see fig. 3, this provides strong support for the theory presented.

When growing by multiplying a number of $n \geq 1$ MMs with the mass and energy density ϱ and e respectively, the two asymptotes are given by eq. (1) for $m \ll m_{\mathrm{diff}}$ and

$$\dot{E} = e\,\frac{\pi}{2}\,\frac{n^{1/4}}{\varrho^{3/4}}\,\frac{(D\,k)^{3/8}}{(2\,\tau)^{1/4}}, \quad \text{for } m \gg m_{\mathrm{diff}}. \tag{3}$$

Equation (1) is the 'diffusion asymptote', eq. (3) the 'convection asymptote', known as Kleiber's law. To the author's knowledge this is the first link from cell metabolism to the metabolism of the organism. The theory is convincingly confirmed by available experiments, see fig. 3.

5 From passive to active

As organisms are at rest most of the time, it can be assumed that the MMs are optimised for this operating state. Therefore, the calculated volume flow $Q = n\,q$ is for the resting state. For mammals, the ratio of heart volume to body volume is geometrically similar, i.e. $\alpha := V_{\text{heart}}/V \approx 6\,\%$ independent of the body sizes V [17]. As a first approximation, the volume flow is proportional to the product of heart volume and heart rate: $Q = f\,V_{\text{heart}} = \alpha\,fV$. With $Q = n\,q$ and $V = n\,v$ the heart rate at rest is therefore

$$f = \frac{Q}{V_{\text{heart}}} = \frac{1}{\alpha}\frac{Q}{V} = \frac{1}{\alpha}\frac{q}{v} = \frac{1}{\alpha}\frac{\pi}{2}\,(\varrho n)^{1/4}\,\frac{(D\,k)^{3/8}}{(2\,\tau)^{1/4}}\,m^{-1/4}.$$

In fact, this axiomatically derived scaling law $f \propto m^{-1/4}$ agrees with the observations [18]. This is a further indication of the validity of the concept and theory presented here.

Fig. 3 Model vs. measurement: the solid lines show the calculated metabolic rate as a function of organism size for three different numbers of MMs represented by the given theory; all physicochemical and physiological properties for cell tissue and capillary flow are kept constant. The symbols show measured metabolic rates for protists, planarians, ectotherms and endotherms. The dashed line applies to $\log n = 8$ and an activity factor of $a = 4$, which corresponds to an increase of heart rate by approximately the same factor. Data are taken from [19–23].

A mammal adapts its metabolism to a required activity by increasing the heart rate by an activity factor $a := f_{\text{active}}/f$. For example, a rower's heart rate can increase from a resting rate of $f = 50\,\text{bpm}$ to $f_{\text{active}} = 200\,\text{bpm}$ during a race. The activity factor is therefore $a = 4$. This increases the volume flow to $Q_{\text{active}} = V_{\text{heart}}\,f_{\text{heart}} = V_{\text{heart}}\,af = a\,Q$. The metabolic rate for the active organism is hence $\dot{E}_{\text{active}} = a\dot{E}$ or $\log \dot{E}_{\text{active}} = \log a + \log \dot{E}$. This constant shift in the vertical direction can be seen in fig. 3 for $\log (n = 8)$: the metabolic rate of mammals at rest is the reference, see solid line for $a = 1$ in fig. 3. The dashed line is the same line, but shifted vertically by the amount $\log (a = 4)$.

6 Conclusion and outlook

The paper links the physiology of cell tissue with the metabolism of organisms concisely, clearly and consistently with physiology such as Murray's law. In this way, the theory presented opens up a new and useful perspective on the metabolism of organisms. It can also be used to provide an insight into evolution under resource scarcity. Darwin's evolution by selection is interpreted as an optimisation heuristic. This heuristic is rationalised by model-based optimisation of an axiomatic model. The optimisation pressures are directed towards (a) lightweight design and (b) resource efficiency. Together, these two pressures lead to Kleiber's law.

In this paper, the conservation laws are formulated and solved in integral form. For a detailed analysis of the concentration field in the MM, the convection-diffusion-reaction equation must be formulated and solved in differential form for both the capillary and the cell tissue. This associated boundary value problem can be approached numerically or analytically. The analytical solution leads to an eigenvalue problem in which the eigenvalues are determined by the boundary values. For this, the solution must be developed in a Bessel series. The optimisation with regard to (a) lightweight design and (b) resource efficiency is then carried out on the basis of the numerical or analytical solution, as in the present work. This would lead to more details, but the essence of the results remains unchanged.

Acknowledgements. I thank the anonymous reviewers for their curiosity and efforts. I would like to thank my co-workers for their supportive team spirit. Finally, I would like to thank my family from the core of my heart for their love, without which life is not possible.

Declarations

The author declares that he has no competing interests.

References

[1] Kleiber, M.: Body size and metabolism. Hilgardia **6**(11), 315–353 (1932) https://doi.org/10.3733/HILG.V06N11P315

[2] West, G.B., Brown, J.H., Enquist, B.J.: A general model for the origin of allometric scaling laws in biology. Science **276**(5309), 122–126 (1997) https://doi.org/10.1126/SCIENCE.276.5309.122,

[3] Banavar, J.R., Maritan, A., Rinaldo, A.: Size and form in efficient transportation networks. Nature **399**(6732), 130–132 (1999) https://doi.org/10.1038/20144;KWRD=SCIENCE

[4] Ballesteros, F.J., Martinez, V.J., Luque, B., Lacasa, L., Valor, E., Moya, A.: On the thermodynamic origin of metabolic scaling. Scientific Reports **8**(1), 1–10 (2018) https://doi.org/10.1038/S41598-018-19853-6;TECHMETA=129,141;SUBJMETA=1464,319,443,57,631;KWRD=BIOENERGETICS, METABOLISM

[5] Taber, L.A., Ng, S., Quesnel, A.M., Whatman, J., Carmen, C.J.: Investigating Murray's law in the chick embryo. Journal of Biomechanics **34**(1), 121–124 (2001) https://doi.org/10.1016/S0021-9290(00)00173-1

[6] Dawson, T.H.: Scaling laws for capillary vessels of mammals at rest and in exercise. Proceedings of the Royal Society B: Biological Sciences **270**(1516), 755–763 (2003) https://doi.org/10.1098/RSPB.2002.2304;REQUESTEDJOURNAL:JOURNAL:RSPB1990;WEBSITE:WEBSITE:RSJ-SITE;WGROUP:STRING:PUBLICATION

[7] Murray, C.D.: The Physiological Principle of Minimum Work. Proceedings of the National Academy of Sciences **12**(3), 207–214 (1926) https://doi.org/10.1073/PNAS.12.3.207

[8] Gogia, S., Neelamegham, S.: Role of fluid shear stress in regulating VWF structure, function and related blood disorders. Biorheology **52**(5-6), 319–335 (2015) https://doi.org/10.3233/BIR-15061/ASSET/6AAF6FEA-0D4B-4715-A9FC-1163E42CAC59/ASSETS/IMAGES/LARGE/10.3233{_}BIR-15061-FIG4.

[9] Khamassi, J., Bierwisch, C., Pelz, P.: Geometry optimization of branchings in vascular networks. Physical Review E **93**(6), 062408 (2016) https://doi.org/10.1103/PHYSREVE.93.062408/FIGURES/15/MEDIUM

[10] Mazzag, B.M., Tamaresis, J.S., Barakat, A.I.: A Model for Shear Stress Sensing and Transmission in Vascular Endothelial Cells. Biophysical Journal **84**(6), 4087 (2003) https://doi.org/10.1016/S0006-3495(03)75134-0

[11] Barakat, A.I., Lieu, D.K., Gojova, A.: Secrets of the code: Do vascular endothelial cells use ion channels to decipher complex flow signals? Biomaterials **27**(5), 671–678 (2006) https://doi.org/10.1016/J.BIOMATERIALS.2005.07.036

[12] Taylor, G.I.: Dispersion of soluble matter in solvent flowing slowly through a tube.

Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences **219**(1137), 186–203 (1953) https://doi.org/10.1098/RSPA.1953.0139

[13] Probstein, R.F.: Physicochemical Hydrodynamics. Butterworth-Heinemann, Boston (1989)

[14] Tao, Z., Raffel, R.A., Souid, A.K., Goodisman, J.: Kinetic Studies on Enzyme-Catalyzed Reactions: Oxidation of Glucose, Decomposition of Hydrogen Peroxide and Their Combination. Biophysical Journal **96**(7), 2977–2988 (2009) https://doi.org/10.1016/J.BPJ.2008.11.071

[15] Tran, Q.H., Unden, G.: Changes in the proton potential and the cellular energetics of Escherichia coli during growth by aerobic and anaerobic respiration or by fermentation. European Journal of Biochemistry **251**(1-2), 538–543 (1998) https://doi.org/10.1046/J.1432-1327.1998.2510538.X;PAGEGROUP:STRING:PUBLICATION

[16] Bridgman, P.W.: Dimensional Analysis. Yale University Press, New Haven (1922)

[17] Schmidt-Nielsen, K.: Animal Physiology: Adaptation and Environment, 5th edn. Cambridge University Press, ??? (1997). https://doi.org/10.1017/9780521570985

[18] Seymour, R.S., Blaylock, A.J.: The principle of laplace and scaling of ventricular wall stress and blood pressure in mammals and birds. Physiological and Biochemical Zoology **73**(4), 389–405 (2000) https://doi.org/10.1086/317741,

[19] DeLong, J.P., Okie, J.G., Moses, M.E., Sibly, R.M., Brown, J.H.: Shifts in metabolic scaling, production, and efficiency across major evolutionary transitions of life. Proceedings of the National Academy of Sciences of the United States of America **107**(29), 12941–12945 (2010) https://doi.org/10.1073/PNAS.1007783107/SUPPL{_}FILE/SD02.

[20] Hemmingsen, A.M.: Energy metabolism as related to body size and respiratory surface, and its evolution. Report of Steno Memorial Hospital . Rep Steno Mom Hosp Copenhagen **9**, 1–110 (1960)

[21] West, G.: Complexity, regularity, diversity and scale from the complexity of diversity Santa Fe Institute (2018). https://www.youtube.com/watch?time_continue=1971&v=WcwqcKkkkcg

[22] Thommen, A., Werner, S., Frank, O., Philipp, J., Knittelfelder, O., Quek, Y., Fahmy, K., Shevchenko, A., Friedrich, B.M., Jülicher, F., Rink, J.C.: Body size-dependent energy storage causes Kleiber's law scaling of the metabolic rate in planarians. eLife **8** (2019) https://doi.org/10.7554/ELIFE.38187

[23] Pelz, P.F., Vergé, A.: Validated biomechanical model for efficiency and speed of rowing. Journal of Biomechanics **47**(13), 3415–3422 (2014) https://doi.org/10.