Subunit-specific roles of LRRC8 proteins in determining glutamate permeability of astrocytic volume-regulated anion channels

Madison L. Chandler, Ariella D. Sprague, Julia W. Nalwalk, and Alexander A. Mongin*

Department of Neuroscience and Experimental Therapeutics, Albany Medical College, Albany, New York, United States

*Correspondence: A.A. Mongin (mongina@amc.edu)

Running Head: COMPOSITION OF ASTROCYTIC LRRC8/VRAC CHANNELS

ABSTRACT

Volume-regulated anion channels (VRACs) are ubiquitous chloride channels that play a crucial role in cell volume regulation but are also involved in many other physiological processes. VRACs are heteromers of proteins from the leucine-rich repeat-containing family 8 (LRRC8A–E), with LRRC8A being essential. Other LRRC8 subunits are expressed in a cell type-specific manner and modulate the biophysical properties of VRACs, including permeability to small signaling molecules. Here, we used primary astrocyte cultures from wild-type and genetically modified C57BL/6 mice to investigate (i) LRRC8 subunit composition of endogenous VRACs in the brain and (ii) the subunit determinants of VRAC permeability to the excitatory neurotransmitter glutamate. qPCR and RNA-seq revealed high expression of *Lrrc8a–d* in mouse forebrain and astrocytes. As expected, *Lrrc8a* deletion abolished VRAC activity, measured as swelling-activated release of the glutamate analogue D-[³H]aspartate. RNAi knockdown of individual subunits established that LRRC8A and LRRC8C are key components of astrocytic glutamate-permeable VRACs, with their siRNAs reducing radiotracer release by 85% and 56%, respectively. Downregulation of LRRC8D had a moderate effect, which depended on the severity of swelling. Combined silencing of LRRC8C and LRRC8D suggested that these subunits act in distinct channel populations. LRRC8B silencing alone was ineffective but partially rescued glutamate release in LRRC8C- or LRRC8D-knockdown cells. Overall, these findings indicate that astrocytic glutamate-permeable VRACs are primarily composed of LRRC8A and LRRC8C, with a possible structural role for LRRC8B. The refined understanding of VRAC subunit composition may inform the development of targeted VRAC inhibitors to mitigate glutamate excitotoxicity in neurological disorders.

NEW & NOTEWORTHY

We examined the subunit composition of volume-regulated anion channels (VRACs), which contribute to pathological glutamate release in various neurological disorders. VRACs are heteromers formed by LRRC8A–E proteins, but the subunit configuration and functional roles within native channels remain poorly understood. In mouse astrocytes, we determined that glutamate-permeable VRACs are composed primarily of LRRC8A and LRRC8C, with structural support from LRRC8B. These findings provide a foundation for developing subunit-specific VRAC inhibitors as potential neuroprotective therapies.

Keywords: Astrocytes, Glutamate release, LRRC8, LRRC8A, VRAC

INTRODUCTION

Volume-regulated anion channels (VRACs), also known as volume-sensitive outwardly rectifying anion channels (VSOR) or volume-sensitive organic osmolyte/anion channels (VSOAC), are swelling-activated chloride (Cl^-) channels found in virtually all vertebrate cells (1-3). VRAC activity is typically assessed by measuring swelling-induced Cl^- currents ($I_{Cl,vol}$) in cells exposed to hypoosmotic media [e.g., early studies (4-7)]. However, in addition to Cl^-, VRACs are also permeable to a range of small negatively charged or uncharged organic molecules, collectively referred to as organic osmolytes (1, 8-11). As their name implies, VRACs play a key role in cell volume control and drive regulatory volume decrease (RVD), a fundamental physiological process that restores cell volume after osmotic swelling. RVD is driven primarily by the electrically coupled efflux of cytosolic Cl^- (via VRACs) and K^+ (via separate K^+ channels), with additional contributions from bicarbonate and organic osmolytes (1-3). Beyond their role in RVD, VRACs have been implicated in numerous other physiological and pathological processes, including regulation of membrane potential, cell proliferation, apoptosis, immune responses, and paracrine signaling [reviewed in (12-16)].

In the context of brain physiology, VRACs are thought to exert a strong influence over the physiological and pathological regulation of brain excitability [reviewed in (13, 17-19)]. This unique role is due to VRACs' permeability to several amino acid-derived neurotransmitters and neuromodulators, including aspartate, glutamate, GABA, and taurine (8, 20, 21). Hence, once open, these channels have considerable potential to affect electrical activity in neural networks (16). VRACs' influence becomes even more pronounced in neural pathologies that promote pathological cell swelling, such as stroke, traumatic brain injury, and epilepsy. In these disorders, swelling-induced activation of VRACs leads to the release of cytosolic glutamate and contributes to excitotoxic neuronal damage (17, 22, 23). In mouse, rat, and canine models of stroke, the putative but structurally diverse VRAC blockers—tamoxifen, DCPIB, dicoumarol, and broma-diolone—potently reduce pathological glutamate release and ischemic brain infarction with a considerable therapeutic window of up to 3 hours after the interruption of blood flow (24-28). However, none of the currently available small molecule inhibitors possess sufficient specificity to confirm that their neuroprotective effects are mediated exclusively through inhibition of VRAC. For nearly three decades, the unknown molecular identity of VRACs has hindered progress in understanding their diverse roles in the brain as well as their potential as a therapeutic target [for a historical perspective, see (29-31)].

In 2014, the laboratories of Ardem Patapoutian and Thomas J. Jentsch independently solved the long-standing enigma of VRAC identity. Using genome-wide siRNA screens combined with a high-throughput YFP quenching assay to measure Cl^- permeability, they identified leucine-rich repeat-containing family 8 member A (LRRC8A, also known as Swell1) as an essential VRAC component (32, 33). LRRC8A belongs to a family of five LRRC8 proteins (LRRC8A-E), which share high sequence homology and a common evolutionary origin with vertebrate pannexins (34). Subsequent studies in CRISPR/Cas9-edited cells further established that LRRC8A must heteromerize with at least one other isoform, LRRC8C, LRRC8D, or LRRC8E, to form a fully functional channel (33, 35). Interestingly, LRRC8A/LRRC8B heteromers remain functionally silent for reasons that are not fully understood (33). Based on the structural homology with pannexins, LRRC8 proteins were predicted to assemble into hexamers with one central ion-conducting pore (34). The hexameric model of VRAC has been confirmed by multiple cryo-electron microscopy (cryo-EM) studies (36-39). However, more recent cryo-EM work indicates that LRRC8 channels can also adopt an alternative heptameric stoichiometry, such as LRRC8C homo-heptamers (40, 41).

The discovery of the indispensable role of LRRC8A has enabled the use of molecular
genetics strategies to study the physiological role of VRACs. The Geha laboratory produced the
first global LRRC8A knockout (KO) (42) to investigate a clinical case of congenital agammaglobu-
linemia, a severe immunoglobulin deficiency due to the loss of B lymphocytes, in a patient with a
C-terminal truncation of LRRC8A (43). Unexpectedly, LRRC8A KO mice developed a lethal
phenotype: ~80% of them died in utero, with the surviving animals rapidly dying due to multi-organ
failure (42). In contrast, tissue- and cell type-specific LRRC8A deletions are generally better
tolerated. For example, mice with conditional deletion of LRRC8A in CNS astrocytes show normal
brain development and remain viable and fertile (44). However, astrocytic LRRC8A KO resulted
in altered neuronal communication and deficits in learning and memory, likely due to reduced
astrocytic glutamate release and consequent decreased activation of neuronal glutamate
receptors (44). In another study, astrocytic VRACs were shown to mediate tonic release of the
inhibitory neurotransmitter GABA in the ventral tegmental area, influencing addictive drug reward
behavior (45). Together, these findings suggest that brain-specific LRRC8A deletion disrupts both
excitatory and inhibitory signaling, with a broad impact on brain physiology. The significance of
VRACs in the brain is underscored by the striking phenotype of mice carrying a brain-wide deletion
of LRRC8A in neurons, astrocytes, and oligodendrocytes, using Nestin[Cre]-driven recombination.
Nestin[Cre+]: LRRC8A$^{-/-}$ mice develop spontaneous seizures and all die between late adolescence
and early adulthood (46-48). Our group has linked this lethal phenotype to changes in GABAergic
signaling and reactive astrogliosis (47); however, other molecular deficits may also contribute.

The lethal phenotype resulting from LRRC8A deletion in the brain (47) has raised
concerns about the therapeutic potential of VRAC blockers and highlighted the need for further
investigation into the structure and function of these channels within the CNS. Unlike the essential
LRRC8A subunit, the partnering LRRC8 isoforms (LRRC8B–E) are expressed in a cell type-
specific manner and modulate the biophysical properties of VRACs, including their permeability
to small signaling molecules. The best-established example of this functional diversity is the
critical role of LRRC8A and LRRC8D in forming VRACs that preferentially conduct small,
uncharged molecules; these include myo-inositol, neutral zwitterions such as taurine, GABA, and
glutamine, as well as chemotherapeutic agents like cisplatin and carboplatin (21, 49, 50). Studies
in model cell lines suggest that glutamate-permeable VRACs can form through alternative
heteromeric assemblies, with LRRC8A pairing separately with LRRC8C, LRRC8D, or LRRC8E
or in combination with multiple subunits such as LRRC8C+LRRC8D [see (21, 50, 51) and
Discussion]. Despite this extensive body of work, the precise subunit composition and
stoichiometry of native VRAC heteromers remain unresolved. To address this knowledge gap,
the present study investigated the LRRC8 subunit determinants of endogenous glutamate-
permeable VRACs in primary mouse astrocytes, employing both an RNAi approach and
genetically modified animals.

MATERIALS AND METHODS

Ethical Approval

All animal procedures used in the present study were approved by the Institutional Animal Care and Use Committee of Albany Medical College (ACUP 24-03002), and strictly conformed to the Guide for the Care and Use of Laboratory Animals as adopted by the U.S. National Institutes of Health (https://grants.nih.gov/grants/olaw/guide-for-the-care-and-use-of-laboratory-animals.pdf).

Animals

This study utilized two types of mice to collect brain tissue and generate primary astrocyte cultures. Wild type (WT) animals were of the C57BL/6J background, originally obtained from the Jackson Laboratory (Farmington, CT, USA; strain #000664, RRID: IMSR_JAX:000664) and bred in-house. Additionally, we used mice with conditional knockout of LRRC8A in the brain (bLRRC8A KO). These were produced by breeding $Lrrc8a^{flox/flox}$ females (gift of Dr. Rajan Sah, Washington University School of Medicine, St. Louis, MO, USA) with commercially available $Nestin^{Cre/+}$ males (B6.Cg-Tg(Nes-cre)1Kln/J; Jackson Lab stock #003771, RRID:IMSR_JAX:003771). Heterozygous $Nestin^{Cre/+}$:$Lrrc8a^{flox/+}$ males were further crossed with $Lrrc8a^{flox/flox}$ female mice to produce $Nestin^{Cre/+}$:$Lrrc8a^{flox/flox}$ mice (bLRRC8A KO) and littermate controls. This breeding scheme produces four genotypes, as presented in Fig. 1A. The specificity and efficacy of this genetic strategy were thoroughly validated in our previous work (47, 48). To visualize Cre activity and monitor for unintentional germline recombination, mice were additionally inbred with a strain carrying the Ai9 tdTomato Cre reporter (B6.Cg-Gt(ROSA)26Sor$^{tm9(CAG-tdTomato)Hze}$/J; Jackson Lab stock #007909, RRID:IMSR_JAX:007909). For simplicity, the presence of the reporter allele is not shown in the breeding diagram. All mice were housed in a temperature, humidity, and light-controlled facility on a 12:12 h light/dark cycle and given free access to food and water.

Brain Tissue Collection and Processing

For brain tissue collection, 14-week-old WT mice were euthanized with a lethal injection of sodium pentobarbital and perfused transcardially, first with room temperature phosphate-buffered saline (PBS), followed by ice-cold PBS. For qRT-PCR analysis, brains were rapidly removed and immediately transferred to a large dissection dish containing ice cold PBS additionally treated with the RNase inhibitor diethyl pyrocarbonate (DEPC, 0.1%). The olfactory bulbs, cerebellum and brainstem were removed, and a midline sagittal cut was made to separate the hemispheres. Each hemisphere was snap-frozen in liquid nitrogen and stored at –80°C until further processing.

Primary Cultures of Mouse Astrocytes

Primary cultures of mouse astrocytes were prepared from brain cortices of newborn (P0-P2) pups of both sexes as most recently described in (48) with minor modifications. Briefly, neonates were anesthetized by cooling and rapidly decapitated. Brains were harvested and

transferred into ice-cold sterile Dulbecco's PBS without calcium and magnesium (ThermoFisher,
Scientific, Waltham, MA, USA, cat. #14190144). Cortical tissue was dissected from meninges,
minced using a sterile surgical blade, incubated with 500 μL of the recombinant protease TrypLE
Express (ThermoFisher, cat. #12605), and additionally triturated by pipetting. Triturated tissue
was digested with TrypLE for 5 min at room temperature (22°C). Enzymatic digestion was
terminated by adding Earl's minimal essential medium (MEM, ThermoFisher, cat. #11095)
supplemented with 10% heat-inactivated horse serum (HIHS, ThermoFisher, cat. #26050) and
penicillin/streptomycin (P/S, ThermoFisher, cat. #15140). The cell suspension was next filtered
through a 70-μm nylon mesh cell strainer (Corning, Corning, NY, USA, cat. # 431751). To obtain
an enriched astrocyte population, cells extracted from either half brain (for WT), or full brain
(bLRRC8A KO) were separately plated onto T75 flasks and grown in a humidified atmosphere of
5% CO_2/balance air at 37°C. With this approach, primary cell cultures typically reached confluency
after around two weeks and were maintained up to six weeks until replating for functional assays.
The purity of astrocyte populations was periodically verified with immunofluorescence by staining
for the astrocyte marker, glial fibrillary acidic protein (anti-GFAP, MilliporeSigma, St. Louis, MO,
USA, cat. #MAB360, RRID:AB_11212597; 1:400 dilution) and was ≥95%.

Quantitative Real-Time PCR (qRT-PCR)

Relative gene expression and efficiency of siRNA gene knockdowns were assessed using
qRT-PCR. Total RNA was isolated from brain tissue or primary astrocyte cultures using the Aurum
Total RNA Mini Kit (Bio-Rad, Hercules, CA, USA, cat. #732-6820) following the manufacturer's
instructions. Processing of tissue samples included on-column DNase digestion to eliminate
genomic DNA contamination. RNA concentration was measured with a NanoDrop 1000
spectrophotometer (ThermoFisher, RRID:SCR_016517). RNA was immediately converted to
cDNA using the iScript cDNA synthesis kit (Bio-Rad, cat. #1708891), with 1 μg of total RNA per
20 μL reaction. The resulting cDNA was analyzed for the relative abundance of *Lrrc8* transcripts
by qPCR using gene-specific primers and SYBR Green Supermix (Bio-Rad, cat. #1708882) in a
CFX96 Real-Time PCR Detection System (Bio-Rad, RRID:SCR_018064). Expression levels were
normalized to the housekeeping gene *Rpl13a*, and where applicable, further normalized to control
samples using the $2^{-\Delta\Delta C(T)}$ method (52). As an internal quality control, expression of an additional
housekeeping gene, *Rps20*, and the abundant transcript *Gapdh* were also quantified. All qPCR
primers were obtained as validated GeneGlobe assays from Qiagen (Germantown, MD, USA);
detailed catalog information is provided in Table 1.

Western Blotting

Relative expression of LRRC8A protein was assessed in lysates from primary astrocyte
cultures. Cells were lysed in a denaturing buffer containing 2% sodium dodecyl sulfate (SDS) and
8 mM EDTA, followed by dilution with 4× Laemmli sample buffer (Bio-Rad, cat. #4561033).
Lysates were boiled for 5 min, and 20 μg of total protein per sample was resolved on 10% Mini-
PROTEAN TGX gels (Bio-Rad, cat. #1610747) via SDS-PAGE. Proteins were then transferred to
polyvinylidene difluoride (PVDF) membranes (Bio-Rad, cat. #1620177). Membranes were
blocked for 20 min in Tris-buffered saline with 0.1% Tween-20 (TBS-T) containing 5% non-fat dry
milk and further incubated overnight at 4°C with mouse monoclonal anti-LRRC8A primary

antibody (Santa Cruz, cat. #sc-517113; RRID:AB_2928142; 1:1,000) diluted in the same blocking buffer. After washing in TBS-T, membranes were incubated for 2 h at room temperature with anti-mouse IgG HRP-conjugated secondary antibody (m-IgG Fc BP-HRP, Santa Cruz, cat. #sc-525416; 1:10,000) in 5% milk/TBS-T. Protein bands were visualized using enhanced chemiluminescence reagent (ECL; GE Healthcare, cat. #RPN2232) and a ChemiDoc Imaging System (Bio-Rad, RRID:SCR_019684). For loading controls, membranes were washed and re-probed for 30 min with HRP-conjugated anti-β-actin antibody (Millipore-Sigma, cat. #A3854; RRID:AB_262011; 1:50,000). Band intensities were quantified using ImageJ software (53) (NIH, RRID:SCR_003070). LRRC8A signals were normalized to β-actin immunoluminescence from the same blot.

RNAi-Mediated LRRC8 Gene Silencing

Targeted knockdown of LRRC8 family members was achieved using gene-specific siRNA constructs obtained from Qiagen (Valencia, CA, USA). AllStars Negative Control siRNA (Qiagen) was used as a non-targeting control. Primary astrocytes at ~70–80% confluency were transfected using Lipofectamine RNAiMAX (Thermo Fisher, cat. #13778075) according to the manufacturer's protocol. siRNA-Lipofectamine complexes were prepared in Opti-MEM (Thermo Fisher, cat. #31985), further diluted with Opti-MEM, and added to cells at final siRNA concentrations ranging from 10 to 50 nM (optimized for individual siRNAs). Details of the siRNA constructs, including catalog numbers and target sequences, are listed in Table 1. Knockdown efficiency for each construct was validated by qRT-PCR. After a 4 h incubation, cultures were supplemented with MEM containing 10% HIHS and penicillin plus streptomycin (MEM+HIHS+P/S). At 24 h post-transfection, the medium was fully replaced with fresh MEM+HIHS+P/S. mRNA expression levels were assessed 48 h post-transfection, and functional assays were performed 96 h post-transfection.

Radiotracer Assays of VRAC Activity

VRAC activity was evaluated by measuring the release of D-[^{3}H]aspartate, a non-metabolizable analog of glutamate, as extensively validated in our earlier studies [e.g., (50, 54, 55)]. Two complementary assay formats were used, employing astrocytes cultured either in multi-well plates or on glass coverslips. In all experiments, astrocytes were preloaded overnight in standard serum-containing medium supplemented with 2 µCi/mL D-[^{3}H]aspartate (Revvity, cat. #NET50100).

For medium-throughput screening of multiple siRNA constructs, cells were grown on poly-D-lysine-coated 12-well plates. After radiotracer loading, cultures were washed to remove extracellular isotope and equilibrated for 30 min at 37°C in isoosmotic Basal medium containing (in mM): 135 NaCl, 3.8 KCl, 1.2 $MgSO_4$, 1.3 $CaCl_2$, 1.2 KH_2PO_4, 10 D-glucose, and 10 HEPES (pH 7.4; osmolality 290 ±3 mOsm/kg). To inhibit radiotracer reuptake via the Na^+-dependent GLAST transporters, cells were exposed to LiCl-substituted media, in which NaCl was isoosmotically replaced with LiCl (56). Baseline release was measured during a 10-min incubation in isoosmotic Li^+-containing medium, followed by stimulation with hypoosmotic Li^+ medium (LiCl reduced by 50 mM; final osmolality ~200 mOsm/kg, ~30% reduction) to activate VRAC. Ten-min fractions of isoosmotic and hypoosmotic media were collected for analysis. Radiotracer release was quantified by liquid scintillation counting using a TRI-CARB 4910TR counter (Revvity), after

mixing samples with Ecoscint A scintillation fluid (National Diagnostics, cat. #LS-273). Integral
release values were normalized to the remaining intracellular D-[^{3}H]aspartate, determined after
cell lysis in 2% SDS containing 8 mM EDTA.

To resolve the kinetics of VRAC activation, D-[^{3}H]aspartate efflux assays were conducted
using a custom-fabricated Lucite superfusion chamber that has a depression on the bottom to
accommodate an 18×18 mm glass coverslip and a Teflon screw lid that leaves a ~200 μm space
above the cells. Coverslips with D-[^{3}H]aspartate-preloaded astrocytes were briefly washed with
Basal medium (for composition see above), transferred into the chamber, and superfused at a
constant rate of ~1.4 mL/min with either isoosmotic or hypoosmotic media. In this setup, we used
NaCl-containing media because released D-[^{3}H]aspartate was effectively removed with the flow.
One-min perfusate fractions were collected into scintillation vials using an automated CF-1
fraction collector (SpectraLab Scientific, Markham, ON, Canada). At the conclusion of each
experiment, cells were lysed in 2% SDS+8 mM EDTA, and D-[^{3}H]aspartate content in all samples
was quantified using a scintillation counter as described above. Data were expressed as fractional
release values, calculated at each time point as a ratio of released [^{3}H] to the total [^{3}H] content
(cumulative release + remaining tracer), using a custom Excel-based analysis program.

281
Statistics

All data are presented as mean values ± SD. n value corresponds to an individual animal (qPCR
and RNA-seq) or an independently transfected astrocyte culture. For cell culture experiments, all
results were reproduced in multiple cultures prepared from at least two different animals, unless
specified otherwise. Statistical differences between three or more groups were analyzed using
one-way ANOVA followed by Tukey's multiple comparisons test. For functional assays conducted
in a multi-well plate format, statistical differences were examined using repeated measures two-
way ANOVA assessing the effects of medium (cell swelling), treatment (siRNA), and their
interaction, followed by Tukey's multiple comparisons test. For clarity, we present only the effects
of treatment, but full analyses are included in the primary datasets deposited in the Open Science
Framework (osf.io). In RNAi expression assays, mRNA expression values were normalized to the
expression levels in siNC-treated cells and compared to a unity using one-sample t-test with
Bonferroni correction for multiple testing (adjusted p = p $\times$ number of comparisons). All statistical
analyses and graph preparation were performed using GraphPad Prism 10.0 (GraphPad
Software, San Diego, CA, USA; RRID:SCR_002798).

RESULTS

LRRC8A Deletion Abolishes Activity of Glutamate-Permeable VRACs in Astrocytes

To investigate the role of LRRC8 proteins in the formation of endogenous astrocytic
VRACs, we first employed a molecular genetics approach using *Nestin*[Cre]-driven recombination.
Nestin[Cre] targets neural precursor cells, leading to gene deletion in all cells of neuroectodermal
origin, including astrocytes (57, 58). Our conditional knockout breeding strategy produced four
genotypes: two types of controls (*Lrrc8a*[flox/+] and *Lrrc8a*[flox/flox] without Cre, referred to as fl/+ and
fl/fl), a heterozygous (HET) LRRC8A deletion, and a homozygous LRRC8A knockout (KO) (Fig.
1A1). Primary astrocyte cultures were prepared from neonatal brains of each genotype and
successful Cre recombination was confirmed using the Ai9 tdTomato reporter. Live-cell imaging
of tdTomato fluorescence showed complete Cre penetrance in both LRRC8A HET and KO

astrocytes (representative image in Fig. 1A2). Western blot analysis of astrocyte culture lysates revealed that heterozygous LRRC8A deletion reduced LRRC8A protein levels by approximately 60% (p <0.0001, Fig. 1B), while KO cultures showed a complete loss of LRRC8A immunoreactivity (p <0.0001, Fig. 1B, full western blot dataset in Supplemental Fig. 1).

Insert Fig. 1 near here

We next assessed functional VRAC activity using a D-[^{3}H]aspartate efflux assay. In control astrocytes, a 30% reduction in extracellular osmolality (to 200 mOsm) triggered a rapid increase in D-[^{3}H]aspartate release, reaching a peak rate of 2.6% within two minutes (an 8.7-fold rise over baseline) followed by a gradual decline consistent with the volume regulatory process. This kinetic profile closely matched our previous measurements of VRAC activation in both rat and mouse astrocytes (50, 54). Astrocytes with heterozygous LRRC8A deletion exhibited a 30% reduction in swelling-activated radiotracer efflux (p = 0.038, Fig. 1C, full analysis in Supplemental Fig. 2). As anticipated, LRRC8A KO astrocytes displayed a profound loss of swelling-induced glutamate release (p < 0.0001, Fig. 1C and Supplemental Fig. 2). However, a small, transient release component persisted during the initial three minutes of hypoosmotic exposure (Fig. 1C). This residual response was characterized by more rapid activation and inactivation kinetics as compared to typical VRAC-mediated release and is likely mediated by a distinct mechanism (see additional considerations in the following paragraph and extended Discussion). Notably, LRRC8A KO astrocytes also showed a substantial reduction in baseline glutamate release under isoosmotic conditions: 28% lower than that of littermate controls (p <0.0001 see inset in Fig. 1C). Together, these findings demonstrate that (i) LRRC8A containing VRACs are the primary conduit for swelling-induced glutamate release in astrocytes, and (ii) these channels also contribute to tonic glutamate release under resting, isoosmotic conditions.

To further investigate the apparent discrepancy between the 60% decrease in LRRC8A protein levels and the relatively modest 30% inhibition of swelling-activated glutamate release in HET cells, we performed additional functional assays using media with a more pronounced reduction in osmolality. When medium osmolality was decreased by ~50% (to 150 mOsm), VRAC activation was dramatically enhanced compared to the response observed with a 30% (to 200mOsm) osmotic reduction, resulting in an over 10-fold increase in maximal release rates between the two conditions (compare Y-axis scales in Fig. 1C and 1D). LRRC8A KO cells again showed a significant reduction in baseline glutamate release, a 40% decrease as compared to controls (p < 0.0001, Fig. 1D, consistent with Fig. 1C). Importantly, in severely swollen astrocytes, heterozygous LRRC8A deletion caused a much more pronounced 62% reduction in VRAC-mediated activity, which is close to what would be expected based on changes in protein expression in HET cells (p <0.0001, Fig. 1D, full analysis in Supplemental Fig. 3). These results may suggest that a reserve pool of LRRC8A-containing anion channels in the HET genotype can compensate under moderate swelling conditions, but this compensatory capacity is lost with more extreme osmotic stress/cellular swelling (see Discussion). This finding has implications for interpreting subsequent RNAi-based knockdown experiments.

Finally, an important observation was made while comparing the "residual" D-[^{3}H]aspartate release in LRRC8A KO astrocytes exposed to 30% and 50% reductions in medium osmolality. In LRRC8A KO cells, the magnitude of this residual release was indistinguishable between the two hypoosmotic conditions (the maximal release rates: 0.62 ± 0.25% vs. 0.51 ± 0.32%, p = 0.4975, insets in Fig. 1C and 1D). This was in striking contrast to control astrocytes, where VRAC activity increased by an order of magnitude under stronger osmotic stress (2.59 ± 0.55% vs. 26.38 ± 5.19%, p < 0.0001). This lack of sensitivity to the extent of cellular swelling further supports the conclusion that the residual release observed in LRRC8A KO astrocytes is not mediated by VRAC (see Discussion).

Relative Abundance of *Lrrc8* Transcripts in Mouse Brain and Primary Astrocyte Cultures

We next measured and compared the endogenous expression of all five *Lrrc8* isoforms in C57BL/6J mouse brains and primary astrocyte cultures derived from C57BL/6J neonates. qPCR analysis of samples prepared from forebrain tissue detected high levels of four *Lrrc8* transcripts, *Lrrc8a-Lrrc8d*, with the order of abundance *Lrrc8b>Lrrc8a≈Lrrc8d>Lrrc8c* (Fig. 2A). mRNA levels of the *Lrrc8* subunits were ~1–5% of the housekeeping gene ribosomal *Rpl13a*, or 0.5–2.5% of the highly prevalent *Gapdh* (Fig. 2A and Supplemental Fig. 4A). Consistent with literature reports, brain levels of *Lrrc8e* were very low-to-non-detectable (Fig. 2A). To independently validate our findings on *Lrrc8* expression, we additionally performed an in-silico analysis of publicly available RNAseq datasets. Figure 2B summarizes expression levels of the five *Lrrc8* isoforms in seven forebrain regions from five mice of both sexes from the Human Protein Atlas. The pattern of relative abundance of *Lrrc8* subunits in RNAseq datasets was nearly identical to those found in our qRT-PCR experiments: *Lrrc8b>Lrrc8d≈Lrrc8a>Lrrc8c*, and extremely low-to-nondetectable *Lrrc8e* (compare Fig. 2A and 2B). Transcripts Per Million (TPM) values of ≥10 confirmed that *Lrrc8a-d* transcripts are highly abundant in the mouse brain.

Insert Fig. 2 near here

In primary astrocyte cultures, the average levels of *Lrrc8* transcripts were comparable to those observed in forebrain tissue. Expression levels of the four predominant *Lrrc8* isoforms ranged from ~1–4% relative to *Rpl13a*, or 0.5–2% when normalized to *Gapdh* (Fig. 2C and Supplemental Fig. 4B). As in brain tissue, *Lrrc8e* expression was detected at trace levels (Fig. 2C). Notably, we observed cell type-specific differences in the relative expression of individual *Lrrc8* isoforms: cultured astrocytes exhibited significantly higher levels of *Lrrc8c* and markedly lower levels of *Lrrc8b* compared to brain tissue (compare Fig. 2A and Fig. 2C). These differences may reflect an intrinsic cell type-specific *Lrrc8* expression profile or could result from artifacts introduced by *in vitro* culture conditions.

Analysis of Subunit-Specific Contributions of LRRC8B-E to Glutamate-Permeable VRACs

To assess the individual contributions of the five LRRC8 isoforms to glutamate permeability, we employed an RNA interference (RNAi) approach using commercially available siRNA constructs. For each of the five *Lrrc8* mRNA species, we tested three distinct siRNA constructs targeting different regions of the mRNA (see Table 1 for target sequences). In cells treated with a non-targeting control siRNA (siNC) the relative expression profile remained unchanged (compare Fig. 2C and Fig. 3A1). All tested siRNA constructs achieved 70–90% knockdown of their respective targets (Fig. 3B1–F1). Therefore, all siRNA constructs were included in the initial functional screenings.

Insert Fig. 3 near here

To increase experimental throughput, initial VRAC activity assays were conducted in a multi-well plate format, measuring swelling-activated D-[^{3}H]aspartate release in response to a 30% reduction in medium osmolarity (see schematic Fig. 3A2). As detailed in the Methods section, these efflux assays were performed in Na$^+$-free medium to prevent reuptake of the tracer by Na$^+$-dependent GLAST transporters. For all experiments, repeated measures two-way ANOVA revealed a main effect of medium osmolarity (basal vs. hypoosmotic, p<0.0001), consistent with activation of VRACs (Fig. 3B2-F2). For simplicity, we present only the post-hoc analyses comparing the effects of individual siRNA constructs to siNC-treated controls under hypoosmotic conditions. We first tested the effect of siRNA-mediated knockdown of the essential LRRC8A

subunit. All three siRNAs targeting LRRC8A (designated siA1, siA3, and siA4) strongly reduced swelling-induced D-[^{3}H]aspartate release, decreasing it by 67–85% as compared to siNC controls (p < 0.0001 vs. siNC; Fig. 3B2). The degree of inhibition ranked siA1 > siA3 > siA4 (Fig. 3B2). Importantly, the high inhibitory potency of the LRRC8A knockdown resembled the near-complete loss of VRAC activity in LRRC8A KO astrocytes.

In contrast to the potent effect of LRRC8A knockdown, none of the siRNAs targeting LRRC8B (siB1, siB3, and siB4) inhibited VRAC activity (Fig. 3C2). However, repeated measures two-way ANOVA revealed a significant interaction between siRNA treatments and media osmolarity (p=0.02; Fig. 3C2), so we performed a post hoc analysis. Interestingly, one construct, siB1, produced a statistically significant 24% increase in swelling-activated D-[^{3}H]aspartate release (p = 0.002 vs. siNC; Fig. 3C2). For all subsequent experiments, we used siB4, which demonstrated high efficacy in reducing *Lrrc8b* mRNA levels, while not affecting functional VRAC activity.

All three siRNA constructs targeting LRRC8C (siC1, siC2, and siC4) significantly reduced VRAC activity (p <0.0001 vs. siNC; Fig. 3D2). The most effective, siC1, reduced swelling-activated D-[^{3}H]aspartate release by 56% (p <0.0001 vs. siNC; Fig. 3D2). The order of inhibition was siC1 > siC4 > siC2, which closely mirrored the efficacy of *Lrrc8c* mRNA knockdown (Fig. 3D1). Due to its superior efficacy, siC1 was used in all subsequent functional assays. The prominent contribution of LRRC8C to VRAC-mediated glutamate release in mouse astrocytes was surprising and contrasted with previous reports in other cell types, including our own findings in rat astrocytes, which showed that LRRC8C knockdown does not meaningfully impact swelling-activated D-[^{3}H]aspartate efflux (see (50) and Discussion).

LRRC8D knockdowns produced more nuanced effects. Two of the tested siRNA constructs, siD2 and siD4, reduced VRAC activity by 10% and 28%, respectively (p = 0.542 and 0.002 vs. siNC, Fig. 3E2). In contrast, siD1 produced a striking 60% increase in swelling-activated D-[^{3}H]aspartate release (p <0.0001 vs. siNC; Fig. 3E2). Follow-up qRT-PCR analysis revealed that siD1 induced off-target effects, leading to a paradoxical upregulation of *Lrrc8a* and *Lrrc8c* transcript levels by 2.5-fold and 1.7-fold, respectively (Supplemental Fig. 5). Based on its superior efficacy and lack of off target-effects, siD4 was used in all subsequent experiments. Overall, these findings suggest that LRRC8D plays a role in glutamate-permeable VRAC function, though its contribution is relatively limited.

Finally, we tested three siRNA constructs targeting LRRC8E (siE1, siE2, and siE4). Although the LRRC8E subunit is expressed in mouse astrocytes at only trace levels (see Fig. 2C), our prior work in rat astroglia has suggested that LRRC8E may be incorporated into astrocytic VRAC and modulate swelling-activated glutamate release (50). In the current study, LRRC8E knockdown had no biologically significant effect on VRAC activity in mouse astrocytes, albeit small increases in VRAC activity were observed with all siRNA constructs (p = 0.001 vs. siNC, Fig. 3F2).

Non-Redundant Functions of LRRC8C and LRRC8D in Glutamate-Permeable VRACs

While LRRC8C knockdown strongly suppressed swelling-activated D-[^{3}H]aspartate release (Fig. 3D2), the resulting inhibition was less pronounced than that observed with knock-down of the essential LRRC8A subunit (compared to Fig. 3B2). These findings suggest that other LRRC8 proteins may function redundantly with LRRC8C in forming heteromeric VRACs. This hypothesis aligns with the original discovery that individual LRRC8B–E subunits are dispensable for VRAC function in HEK293 and other cells (32, 33). It is also consistent with our earlier work showing that, in rat astrocytes, LRRC8C and LRRC8D serve as mutually replaceable partners of

LRRC8A in forming glutamate-permeable VRACs (50). To test if this redundancy extends to mouse astrocytes, we performed a double-knockdown of LRRC8C and LRRC8D.

We first verified that co-transfection with siRNAs targeting *Lrrc8c* (siC1) and *Lrrc8d* (siD4) effectively reduced target gene expression by ≥80%, both individually and in combination (p<0.01, p<0.05; Fig. 4A). Functional assays performed in a multi-well plate revealed that dual knockdown of LRRC8C and LRRC8D did not further decrease glutamate release as compared to LRRC8C knockdown alone (60% vs. 56% inhibition, p = 0.240, Fig. 4B). To assess VRAC activity with higher temporal resolution, we repeated the double-knockdown experiments using a Lucite superfusion system. In this setup, LRRC8C knockdown reduced the maximum rate of glutamate release by 33%, while LRRC8C+D knockdown lowered it by 44%; however, this difference was not statistically significant (p = 0.7591; Fig. 4C). Analysis of the integral VRAC-mediated release under hypoosmotic conditions showed a small difference between siC1 alone and the siC1+siD4 combination (p = 0.011, Supplemental Fig. 6A). While differences in the degree of VRAC inhibition between plate-based and superfusion assay formats were noted, the lack of additive effect between LRRC8C and LRRC8D knockdowns was consistent across both.

Insert Fig. 4 near here

Our LRRC8A KO data, particularly the results of the heterozygous LRRC8A deletion, suggest that in cells exposed to a 30% reduction in medium osmolality, VRAC channels may have a spared capacity obscuring the full effects of LRRC8 subunit knockdown. Therefore, we repeated the double-knockdown experiments in astrocytes exposed to a 50% reduction in medium osmolarity (Fig. 4D). Under these conditions, knockdowns of individual LRRC8C and LRRC8D subunits strongly inhibited VRAC activity (50% and 42% decrease in maximal release rates, respectively; p<0.0001 vs. siNC, Fig. 4D, additional analysis in Supplemental Fig. 6B). The double knockdown of LRRC8C plus LRRC8D produced partially additive inhibition as compared to single knockdowns (a 72% decrease, p=0.004, vs LRRC8C and p =0.0001 vs. LRRC8D alone, Fig. 4D and Supplemental Fig. 6B). Partial additivity of the double-knockdown (as opposed to synergistic effect) is consistent with the existence of separate LRRC8A+LRRC8C and LRRC8A+LRRC8D-containing heteromeric VRACs (see Discussion).

Functional Contribution of LRRC8B to LRRC8C- and LRRC8D-Containing VRACs

Given that the combined downregulation of LRRC8C and LRRC8D did not reach the full effect of LRRC8A knockdown, we hypothesized that another abundant LRRC8 isoform, namely LRRC8B, serves as a functional substitute for either LRRC8C and/or LRRC8D. To test this hypothesis, we explored the potential contribution of LRRC8B in double-knockdown siRNA assays. We predicted that knocking down either LRRC8C+LRRC8B or LRRC8D+LRRC8B would produce a more substantial functional effect than knocking down of LRRC8C or LRRC8D alone.

We first tested the efficacy of the double-knockdown of LRRC8B and LRRC8C. qRT-PCR analysis confirmed that siB4 and siC1 constructs were equally effective when transfected alone or in combination (p<0.01, p<0.05; Fig. 5A). Unexpectedly, in VRAC activity assays, the double knockdown of LRRC8B and LRRC8C resulted in a partial rescue of VRAC-mediated glutamate release as compared to the effect of LRRC8C knockdown on its own (p<0.0001 vs. siNC; Fig. 5B). Specifically, the siB4 construct had no effect (p= 0.98; Fig. 5B), siC1 suppressed VRAC activity by 51% (p<0.0001; Fig. 5B), and the combination of siB4 and siC1 paradoxically caused a smaller 25% inhibition (p < 0.0001, siB1+C1 vs. siC1 alone, Fig. 5B).

Insert Fig. 5 near here

In a similar fashion, we quantified the efficacy of the double-knockdown of LRRC8B and
LRRC8D. qRT-PCR analysis verified that siB4 and siD4 constructs were equally effective when
transfected alone or in combination (p<0.001; Fig. 5C). In functional assays, the double
knockdown of LRRC8B and LRRC8D resulted in a complete rescue of the VRAC-mediated
glutamate release as compared to the effect LRRC8D knockdown on its own (Fig. 5D).
Specifically, siB4 had no effect (p= 0.4145; Fig. 5D), siD4 reduced VRAC activity by 17%
(p<0.001; Fig. 5D), and the combination of siB4 and siD4 caused full recovery of glutamate
release in hypoosmotic medium (Fig. 5D).

It important to note that the rescue effects of co-transfection with siB4 were subunit
specific. In experiments involving the combined knockdown of LRRC8A+LRRC8B, the effect of
siA1+siB4 was no different from the effect of siA1alone (Supplemental Fig. 7A). Overall, these
data suggest that LRRC8B can act as a structural partner in LRRC8A+LRRC8C and
LRRC8A+LRRC8D containing channels.

DISCUSSION

The main objective of this study was to identify which LRRC8 isoforms contribute to the formation of glutamate-permeable VRACs in mouse astrocytes. VRAC activation in this cell type is considered a major source of pathological glutamate release and a possible therapeutic target in stroke and other CNS disorders [see ref. (44) and reviews (17, 18, 23)]. Our key findings can be summarized as follows: (1) Genetic knockout of the essential LRRC8A subunit confirms that LRRC8A-containing VRACs are the primary pathway for swelling-activated glutamate release in moderately swollen astrocytes and account for ~98% of the release under conditions of substantial cell swelling. (2) Results of siRNA-mediated knockdown of LRRC8A through LRRC8E, individually and in combination, support a model in which glutamate-permeable VRACs critically depend on LRRC8A and LRRC8C, with likely structural contribution from LRRC8B. (3) The LRRC8D subunit provides an additional contribution to swelling-activated glutamate release but is likely present in a distinct subpopulation of VRACs.

Despite extensive work in the field, there is ongoing debate over whether VRACs represent the primary pathway for swelling-activated glutamate release, particularly in astrocytes. Several studies have proposed alternative molecular candidates as major, or even primary, conduits for swelling- and/or Ca^{2+}-activated glutamate release from astrocytes, including SLCO2A1-containing Maxi Cl^- channels (59, 60), Ca^{2+}-activated BEST1 channels (61, 62), and channels formed by the Tweety homolog proteins TTYH1–3 (63). Furthermore, astroglial Cx43 hemichannels (64) and the $P2X_7$ receptor-activated pannexin complexes (65, 66) may also release amino acid transmitters in response to various stimuli. Our current findings in genetically modified astrocytes show that deletion of the essential LRRC8A subunit virtually abolishes swelling-activated glutamate release. Consistent with our earlier study (47), we observed a small residual release in LRRC8A KO astrocytes, which was insensitive to the extent of cell swelling. We speculate that this residual release in LRRC8A KO cells is mediated by one of the above-mentioned alternative permeability routes, which may be activated by Ca^{2+}. Although the molecular nature of this additional release component is not clear, it is cell volume independent, and its quantitative contribution is minimal. Interestingly, as compared to cells prepared from littermate controls, LRRC8A KO astrocytes display a significant reduction in tonic glutamate release under isoosmotic conditions. Overall, this part of our work suggests that (1) LRRC8A-containing VRACs dominate the swelling-activated release of cytosolic excitatory amino acids, and (2) VRACs exhibit limited tonic activity in non-swollen cells.

While the *Lrrc8a* gene product can be found in virtually all vertebrate cells, the mRNA and protein expression profiles of partnering *Lrrc8* subunits exhibit cell type-, tissue-, and species-specific differences (34, 67). For example, a recent comprehensive study in the mouse cochlea revealed that in this organ different cell types can express dramatically different assortments of LRRC8 proteins (68). Therefore, we first explored and compared the *Lrrc8* transcript profiles in mouse brain tissue and primary mouse astrocytes. qRT-PCR analysis of the mouse forebrain revealed abundant expression of *Lrrc8a* through *Lrrc8d* mRNA, while *Lrrc8e* levels were low to undetectable. These results were corroborated by an *in-silico* analysis of publicly available RNA-seq datasets. Interestingly, both approaches revealed particularly high brain expression of *Lrrc8b*. The relevance of this finding is not clear given that the LRRC8B protein is yet to be reliably associated with any established VRAC function. In mouse astrocyte cultures, qRT-PCR revealed generally similar *Lrrc8* expression patterns. However, we noted a shift in the relative abundance

of specific subunits: *Lrrc8c* was expressed at ~3-fold higher levels, while *Lrrc8b* expression was reduced ~3.5-fold compared to the forebrain. Whether these differences reflect a cell type-specific expression profile or adaptations to *in vitro* culture conditions remains to be determined. Notably, our analyses also revealed species-specific differences in *Lrrc8* transcript levels. In the human brain, *Lrrc8a* dominates over other *Lrrc8b-d* isoforms with *Lrrc8e* being largely absent (see RNA-seq data in a Supplemental Fig. 8). In rat brain and astrocytes, *Lrrc8a* through *Lrrc8d* are detected in similar quantities, whereas astrocytic *Lrrc8e* is expressed at low but functionally relevant levels (50). These species-dependent expression patterns are important to consider when interpreting functional data presented in this study.

Our understanding of the subunit composition of VRACs capable of conducting organic anions, including the neurotransmitters glutamate and aspartate, remains incomplete and somewhat contradictory. In CRISPR/Cas9-edited HEK293 cells, only LRRC8A+LRRC8D and LRRC8A+LRRC8E heteromers show significant conductance of the glutamate analog D-[^{3}H]as-partate, while LRRC8A+LRRC8C channels showed little activity (21). In comparison, the results of heterologous expression in *Xenopus* oocytes demonstrated that LRRC8A+LRRC8C and LRRC8A+LRRC8E channels exhibit measurable glutamate permeability (51). The latter *Xenopus* study also reported glutamate release through LRRC8A+LRRC8D heteromers, albeit at lower rates (51). Our own RNAi analysis in rat astrocytes indicated that endogenous glutamate-permeable VRACs are composed of LRRC8A, LRRC8C, and LRRC8D proteins, with a possible minor contribution from LRRC8E (50). In support of this complex heteromeric assembly, work in model cell lines, such as HEK and HeLa cells, has shown that individual deletion of LRRC8B through LRRC8E does not lead to an appreciable reduction of VRAC activity, suggesting that these subunits play interchangeable roles in channel function (33, 35). Given this background, we were surprised to find that in mouse astrocytes, swelling-activated glutamate release was strongly dependent on the expression of LRRC8C (56% inhibition observed following LRRC8C knock-down). To our knowledge, the only comparable recent finding is the complete loss of VRAC currents and swelling-activated cyclic dinucleotide transport in mouse T lymphocytes after genetic deletion of LRRC8C (69). Taken together, these data suggest that in mouse astrocytes, endogenous VRACs are predominantly formed by LRRC8A and LRRC8C, with limited functional input from other LRRC8 subunits.

In rat astrocytes, glutamate-permeable VRACs are formed with redundant contributions of LRRC8C and LRRC8D (50). In this species, knockdown of either LRRC8C or LRRC8D alone had no significant effect, whereas combined knockdown of both subunits caused strong inhibition of swelling-activated D-[^{3}H]aspartate release (50). Based on this observation, we tested the effect of combined LRRC8C and LRRC8D knockdown in mouse astroglia. Surprisingly, in mouse astrocytes subjected to moderate hypotonic stress (30% reduction in medium osmolality), a combination of siRNAs targeting LRRC8C and LRRC8D produced no additional inhibition beyond that seen with LRRC8C knockdown alone. This result was confirmed using two independent assay systems. However, when cells were exposed to more severe hypotonic stress (50% reduction in medium osmolality), LRRC8D knockdown alone exerted a stronger inhibitory effect on glutamate release, and the combined knockdown of LRRC8C and LRRC8D produced partially additive effects. The lack of synergy between LRRC8C and LRRC8D knockdown in mouse astrocytes is consistent with a model in which LRRC8C and LRRC8D contribute to distinct populations of VRACs (see Fig. 6). Previously, we and others have shown that LRRC8A+ LRRC8D heteromers form a specific VRAC subtype that preferentially conducts uncharged and net-neutral osmolytes, such as taurine, GABA, and myo-inositol, but also exhibits measurable

permeability to glutamate (21, 49, 50). The current data support the existence of two separate channel populations (LRRC8A+LRRC8C) and (LRRC8A+LRRC8D) with distinct permeability profiles.

<mark>*Insert Fig. 6 near here*</mark>

Finally, we would like to discuss potential contributions of LRRC8B. Although highly expressed in the brain [ref. (34) and current data], this subunit does not yet have an established physiological role. Even though LRRC8B is present in endogenous VRACs, heteroexpression of LRRC8A+LRRC8B in model cell lines or their reconstitution in lipid bilayers does not produce functional VRAC currents (21, 33, 35, 51). Among the five LRRC8 isoforms, only LRRC8A and LRRC8B contain charged arginines in the narrowest part of the channel (R103 and R99, respectively) (36-38). In contrast, LRRC8C-E contain neutral leucine or phenylalanine in the same position, and the presence of these subunits results in a wider VRAC pore (70). The only reported physiological role of LRRC8B is the increased Ca^{2+} leak from the endoplasmic reticulum in HEK293 cells overexpressing LRRC8B, and reciprocally, slower depletion of ER calcium stores caused by LRRC8B knockdown (71). Our current data show that in mouse astrocytes, individual LRRC8B knockdowns did not produce inhibition of glutamate-permeable VRACs (one of three effective siRNAs weakly *stimulated* VRAC activity). However, when LRRC8B was downregulated in combination with LRRC8C or LRRC8D, it partially rescued the inhibitory effects of targeting LRRC8C or LRRC8D subunits alone. Interestingly, in the CRISPR/ Cas9-edited HEK293 cells, LRRC8B deletion also caused a noticeable increase in swelling-activated D-[^{3}H]aspartate efflux (21). Together, these findings suggest that LRRC8B is present in glutamate-permeable VRACs but paradoxically restricts their permeability for this amino acid. We speculate that LRRC8B can structurally substitute for downregulated LRRC8C or LRRC8D in the heteromeric VRACs. However, the resulting LRRC8B "enrichment" may make the VRAC pore narrower, much like what has been reported for the homomeric LRRC8A assemblies (36, 37, 39).

In summary, this study provides important insights into the subunit composition of native glutamate-permeable VRACs in the mouse brain and may serve as a steppingstone for the development of subunit-specific VRAC blockers. Beyond their many physiological roles, VRACs are recognized as significant contributors to glutamate-driven tissue damage in stroke, traumatic brain injury, and possibly epilepsy (17, 22, 23). Adding to VRACs' clinical relevance, a recent study discovered that gain-of-function mutations in the human *LRRC8C* gene cause a multi-system disorder named TIMES [Telangiectasia, Impaired intellectual development, Microcephaly, Eye abnormalities, and Short stature (72)]. These mutations lead to constitutive VRAC activity and are associated with severe neurological deficits including an abnormally small brain, optic nerve atrophy, spasticity, and seizures (72). Targeting glutamate-permeable VRACs may mitigate hyperexcitability and neurotoxicity in CNS disorders, with more limited impact on broader VRAC functions in other organs and tissues.

DATA AVAILABILITY

All primary data included in this manuscript have been archived in the Open Science Framework depository (osf.io) and can be accessed at https://osf.io/cdxqf

SUPPLEMENTAL MATERIAL

The following supplemental figures for the manuscript are referred to in the text and can be found at https://osf.io/cdxqf or at https://doi.org/10.6084/m9.figshare.29636810

Supplemental Figure 1. Full-length western blots of LRRC8A and β-actin immunoreactivity in four sets of astrocyte culture lysates derived from LRRC8A fl/+, fl/fl, HET, and KO neonates.

Supplemental Figure 2. Extended statistical analysis of swelling-activated D-[^{3}H]aspartate release from genetically modified astrocytes in response to a 30% reduction in medium osmolarity.

Supplemental Figure 3. Extended statistical analysis of swelling-activated D-[^{3}H]aspartate release from genetically modified astrocytes in response to a 50% reduction in medium osmolarity.

Supplemental Figure 4. Additional qRT-PCR analysis and Ct values of *Lrrc8a-e* expression in mouse forebrain (A) and primary astrocyte cultures (B).

Supplemental Figure 5. qRT-PCR analysis of all five *Lrrc8* transcripts in primary mouse astrocytes treated with three independent *Lrrc8d* gene-specific siRNA constructs (siD1, siD2, siD4).

Supplemental Figure 6. Extended statistical analysis of VRAC activity in mouse astrocytes treated with siNC, siC1, siD4, or siC1+siD4 in response to a 30% or 50% reduction in medium osmolarity.

Supplemental Figure 7. Additivity experiments assessing the combinatory effect of double LRRC8A+B knockdown or LRRC8C+E knockdown on swelling-activated D-[^{3}H]aspartate release from mouse astrocytes.

Supplemental Figure 8. RNA-seq analysis of *LRRC8A-E* expression in four human brain regions (cortex, amygdala, caudate, and hippocampus)

ACKNOWLEDGMENTS

The authors thank Dr. Preeti Dohare for help with validating LRRC8D siRNA constructs and Antonio M. Fidaleo for his assistance in RNA-seq analyses.

GRANTS

This study was supported by the National Institutes of Health, grants R01 NS111943 and R21 NS140782 (to A.A.M.).

DISCLOSURES

The authors have no conflict of interest, financial or otherwise, to disclose.

AUTHOR CONTRIBUTIONS

Conceived and designed research: M.L.C. and A.A.M. Performed experiments: M.L.C., A.D.S., J.W.N. Analyzed data: M.L.C., A.D.S., A.A.M. Interpreted results: M.L.C. and A.A.M. Prepared figures: M.L.C. and A.A.M. Drafted manuscript: M.L.C. Edited and revised manuscript, M.L.C., A.D.S., J.W.N., and A.A.M. Approved final version of manuscript: M.L.C., A.D.S., J.W.N., and A.A.M.

REFERENCES

1. **Strange K, Emma F and Jackson PS**. Cellular and molecular physiology of volume-sensitive anion channels. *Am J Physiol* 270: C711-C730, 1996.

2. **Okada Y**. Volume expansion-sensing outward-rectifier Cl- channel: fresh start to the molecular identity and volume sensor. *Am J Physiol Cell Physiol* 273: C755-C789, 1997. doi: 10.1152/ajpcell.1997.273.3.C755

3. **Nilius B, Eggermont J, Voets T, Buyse G, Manolopoulos V and Droogmans G**. Properties of volume-regulated anion channels in mammalian cells. *Prog Biophys Mol Biol* 68: 69-119, 1997.

4. **Hazama A and Okada Y**. Ca2+ sensitivity of volume-regulatory K+ and Cl- channels in cultured human epithelial cells. *J Physiol* 402: 687-702, 1988.

5. **Cahalan MD and Lewis RS**. Role of potassium and chloride channels in volume regulation by T lymphocytes. *Soc Gen Physiol Ser* 43: 281-301, 1988. doi: n/a

6. **Worrell RT, Butt AG, Cliff WH and Frizzell RA**. A volume-sensitive chloride conductance in human colonic cell line T84. *Am J Physiol* 256: C1111-C1119, 1989.

7. **Solc CK and Wine JJ**. Swelling-induced and depolarization-induced C1-channels in normal and cystic fibrosis epithelial cells. *Am J Physiol* 261: C658-C674, 1991. doi: 10.1152/ajpcell.1991.261.4.C658

8. **Kimelberg HK, Goderie SK, Higman S, Pang S and Waniewski RA**. Swelling-induced release of glutamate, aspartate, and taurine from astrocyte cultures. *J Neurosci* 10: 1583-1591, 1990.

9. **Roy G and Malo C**. Activation of amino acid diffusion by a volume increase in cultured kidney (MDCK) cells. *J Membr Biol* 130: 83-90, 1992.

10. **Kirk K, Ellory JC and Young JD**. Transport of organic substrates via a volume-activated channel. *J Biol Chem* 267: 23475-23478, 1992.

11. **Jackson PS and Strange K**. Volume-sensitive anion channels mediate swelling-activated inositol and taurine efflux. *Am J Physiol* 265: C1489-C1500, 1993. doi: 10.1152/ajpcell.1993.265.6.C1489

12. **Eggermont J, Trouet D, Carton I and Nilius B**. Cellular function and control of volume-regulated anion channels. *Cell Biochem Biophys* 35: 263-274, 2001.

13. **Mongin AA**. Volume-regulated anion channel--a frenemy within the brain. *Pflugers Arch* 468: 421-441, 2016.

14. **Jentsch TJ**. VRACs and other ion channels and transporters in the regulation of cell volume and beyond. *Nat Rev Mol Cell Biol* 17: 293-307, 2016.

15. **Chen L, Konig B, Liu T, Pervaiz S, Razzaque YS and Stauber T**. More than just a
pressure relief valve: physiological roles of volume-regulated LRRC8 anion channels.
Biol Chem 400: 1481-1496, 2019. doi: 10.1515/hsz-2019-0189

16. **Okada Y**. Physiology of the volume-sensitive/regulatory anion channel VSOR/VRAC:
part 2: its activation mechanisms and essential roles in organic signal release. *J Physiol*
Sci 74: 34, 2024. doi: 10.1186/s12576-023-00897-x

17. **Kimelberg HK**. Astrocytic swelling in cerebral ischemia as a possible cause of injury
and target for therapy. *Glia* 50: 389-397, 2005.

18. **Mongin AA**. Disruption of ionic and cell volume homeostasis in cerebral ischemia: The
perfect storm. *Pathophysiology* 14: 183-193, 2007.

19. **Akita T and Okada Y**. Characteristics and roles of the volume-sensitive outwardly
rectifying (VSOR) anion channel in the central nervous system. *Neuroscience* 275: 211-
231, 2014.

20. **Hyzinski-Garcia MC, Rudkouskaya A and Mongin AA**. LRRC8A protein is
indispensable for swelling-activated and ATP-induced release of excitatory amino acids
in rat astrocytes. *J Physiol* 592: 4855-4862, 2014.

21. **Lutter D, Ullrich F, Lueck JC, Kempa S and Jentsch TJ**. Selective transport of
neurotransmitters and modulators by distinct volume-regulated LRRC8 anion channels.
J Cell Sci 130: 1122-1133, 2017.

22. **Mongin AA and Kimelberg HK**. Astrocytic swelling in neuropathology. In: *Neuroglia*,
edited by Kettenmann H and Ransom BR. Oxford/New York: Oxford University Press,
2005.

23. **Wilson CS and Mongin AA**. Cell volume control in healthy brain and neuropathologies.
Curr Top Membr 81: 385-455, 2018.

24. **Kimelberg HK, Feustel PJ, Jin Y, Paquette J, Boulos A, Keller RW, Jr. and Tranmer**
BI. Acute treatment with tamoxifen reduces ischemic damage following middle cerebral
artery occlusion. *Neuroreport* 11: 2675-2679, 2000.

25. **Kimelberg HK, Jin Y, Charniga C and Feustel PJ**. Neuroprotective activity of
tamoxifen in permanent focal ischemia. *J Neurosurg* 99: 138-142, 2003.

26. **Zhang Y, Zhang H, Feustel PJ and Kimelberg HK**. DCPIB, a specific inhibitor of
volume regulated anion channels (VRACs), reduces infarct size in MCAo and the
release of glutamate in the ischemic cortical penumbra. *Exp Neurol* 210: 514-520, 2008.

27. **Boulos AS, Deshaies EM, Dalfino JC, Feustel PJ, Popp AJ and Drazin D**. Tamoxifen
as an effective neuroprotectant in an endovascular canine model of stroke. *J Neurosurg*
114: 1117-1126, 2011.

28. **Chen J, Yang J, Chu J, Chen KH, Alt J, Rais R and Qiu Z**. The SWELL1 channel
promotes ischemic brain damage by mediating neuronal swelling and glutamate toxicity.
Adv Sci (Weinh) 11: e2401085, 2024. doi: 10.1002/advs.202401085

29. **Pedersen SF, Okada Y and Nilius B**. Biophysics and physiology of the volume-
regulated anion channel (VRAC)/volume-sensitive outwardly rectifying anion channel
(VSOR). *Pflugers Arch* 468: 371-383, 2016.

30. **Strange K, Yamada T and Denton JS**. A 30-year journey from volume-regulated anion
currents to molecular structure of the LRRC8 channel. *J Gen Physiol* 151: 100-117,
2019.

31. **Okada Y**. Physiology of the volume-sensitive/regulatory anion channel VSOR/VRAC.
Part 1: from its discovery and phenotype characterization to the molecular entity
identification. *J Physiol Sci* 74: 3, 2024. doi: 10.1186/s12576-023-00897-x

32. **Qiu Z, Dubin AE, Mathur J, Tu B, Reddy K, Miraglia LJ, Reinhardt J, Orth AP and
Patapoutian A**. SWELL1, a plasma membrane protein, is an essential component of
volume-regulated anion channel. *Cell* 157: 447-458, 2014.

33. **Voss FK, Ullrich F, Munch J, Lazarow K, Lutter D, Mah N, Andrade-Navarro MA,
von Kries JP, Stauber T and Jentsch TJ**. Identification of LRRC8 heteromers as an
essential component of the volume-regulated anion channel VRAC. *Science* 344: 634-
638, 2014. doi: 10.1126/science.1252826

34. **Abascal F and Zardoya R**. LRRC8 proteins share a common ancestor with pannexins,
and may form hexameric channels involved in cell-cell communication. *Bioessays* 34:
551-560, 2012.

35. **Syeda R, Qiu Z, Dubin AE, Murthy SE, Florendo MN, Mason DE, Mathur J, Cahalan
SM, Peters EC, Montal M and Patapoutian A**. LRRC8 proteins form volume-regulated
anion channels that sense ionic strength. *Cell* 164: 499-511, 2016. doi:
10.1016/j.cell.2015.12.031

36. **Deneka D, Sawicka M, Lam AKM, Paulino C and Dutzler R**. Structure of a volume-
regulated anion channel of the LRRC8 family. *Nature* 558: 254-259, 2018.

37. **Kasuya G, Nakane T, Yokoyama T, Jia Y, Inoue M, Watanabe K, Nakamura R,
Nishizawa T, Kusakizako T, Tsutsumi A, Yanagisawa H, Dohmae N, Hattori M,
Ichijo H, Yan Z, Kikkawa M, Shirouzu M, Ishitani R and Nureki O**. Cryo-EM structures
of the human volume-regulated anion channel LRRC8. *Nat Struct Mol Biol* 25: 797-804,
2018.

38. **Kefauver JM, Saotome K, Dubin AE, Pallesen J, Cottrell CA, Cahalan SM, Qiu Z,
Hong G, Crowley CS, Whitwam T, Lee WH, Ward AB and Patapoutian A**. Structure
of the human volume regulated anion channel. *eLife* 7: e38461, 2018.

39. **Kern DM, Oh S, Hite RK and Brohawn SG**. Cryo-EM structures of the DCPIB-inhibited
volume-regulated anion channel LRRC8A in lipid nanodiscs. *eLife* 8: e42636, 2019. doi:
10.7554/eLife.42636

40. **Rutz S, Deneka D, Dittmann A, Sawicka M and Dutzler R**. Structure of a volume-
regulated heteromeric LRRC8A/C channel. *Nat Struct Mol Biol* 30: 52-61, 2023.

41. **Takahashi H, Yamada T, Denton JS, Strange K and Karakas E**. Cryo-EM structures
of an LRRC8 chimera with native functional properties reveal heptameric assembly.
eLife 12: e82431, 2023. doi: 10.7554/eLife.82431

42. **Kumar L, Chou J, Yee CS, Borzutzky A, Vollmann EH, von Andrian UH, Park SY,
Hollander G, Manis JP, Poliani PL and Geha RS**. Leucine-rich repeat containing 8A
(LRRC8A) is essential for T lymphocyte development and function. *J Exp Med* 211: 929-
942, 2014.

43. **Sawada A, Takihara Y, Kim JY, Matsuda-Hashii Y, Tokimasa S, Fujisaki H, Kubota
K, Endo H, Onodera T, Ohta H, Ozono K and Hara J**. A congenital mutation of the
novel gene LRRC8 causes agammaglobulinemia in humans. *J Clin Invest* 112: 1707-
1713, 2003.

44. **Yang J, Vitery MDC, Chen J, Osei-Owusu J, Chu J and Qiu Z**. Glutamate-releasing
SWELL1 channel in astrocytes modulates synaptic transmission and promotes brain
damage in stroke. *Neuron* 102: 813-827, 2019.

45. **Yang J, Chen J, Liu Y, Chen KH, Baraban JM and Qiu Z**. Ventral tegmental area
astrocytes modulate cocaine reward by tonically releasing GABA. *Neuron* 2023.

46. **Zhou JJ, Luo Y, Chen SR, Shao JY, Sah R and Pan HL**. LRRC8A-dependent volume-
regulated anion channels contribute to ischemia-induced brain injury and glutamatergic
input to hippocampal neurons. *Exp Neurol* 332: 113391, 2020.

47. **Wilson CS, Dohare P, Orbeta S, Nalwalk JW, Huang Y, Ferland RJ, Sah R, Scimemi
A and Mongin AA**. Late adolescence mortality in mice with brain-specific deletion of the
volume-regulated anion channel subunit LRRC8A. *FASEB J* 35: e21869, 2021.

48. **Balkaya M, Dohare P, Chen S, Schober AL, Fidaleo AM, Nalwalk JW, Sah R and
Mongin AA**. Conditional deletion of LRRC8A in the brain reduces stroke damage
independently of swelling-activated glutamate release. *iScience* 26: 106669, 2023. doi:
10.1016/j.isci.2023.106669

49. **Planells-Cases R, Lutter D, Guyader C, Gerhards NM, Ullrich F, Elger DA,
Kucukosmanoglu A, Xu G, Voss FK, Reincke SM, Stauber T, Blomen VA, Vis DJ,
Wessels LF, Brummelkamp TR, Borst P, Rottenberg S and Jentsch TJ**. Subunit
composition of VRAC channels determines substrate specificity and cellular resistance
to Pt-based anti-cancer drugs. *EMBO J* 34: 2993-3008, 2015.

50. **Schober AL, Wilson CS and Mongin AA**. Molecular composition and heterogeneity of
the LRRC8-containing swelling-activated osmolyte channels in primary rat astrocytes. *J*
Physiol 595: 6939-6951, 2017.

51. **Gaitan-Penas H, Gradogna A, Laparra-Cuervo L, Solsona C, Fernandez-Duenas V,**
Barrallo-Gimeno A, Ciruela F, Lakadamyali M, Pusch M and Estevez R. Investigation
of LRRC8-mediated volume-regulated anion currents in Xenopus oocytes. *Biophys J*
111: 1429-1443, 2016.

52. **Livak KJ and Schmittgen TD**. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25: 402-408, 2001.

53. **Schneider CA, Rasband WS and Eliceiri KW**. NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* 9: 671-675, 2012.

54. **Abdullaev IF, Rudkouskaya A, Schools GP, Kimelberg HK and Mongin AA**.
Pharmacological comparison of swelling-activated excitatory amino acid release and Cl-
currents in rat cultured astrocytes. *J Physiol* 572: 677-689, 2006. doi:
10.1113/jphysiol.2005.103820

55. **Bowens NH, Dohare P, Kuo YH and Mongin AA**. DCPIB, the proposed selective
blocker of volume-regulated anion channels, inhibits several glutamate transport
pathways in glial cells. *Mol Pharmacol* 83: 22-32, 2013. doi: 10.1124/mol.112.080457

56. **Schober AL and Mongin AA**. Intracellular levels of glutamate in swollen astrocytes are
preserved via neurotransmitter reuptake and de novo synthesis: implications for
hyponatremia. *J Neurochem* 135: 176-185, 2015.

57. **Tronche F, Kellendonk C, Kretz O, Gass P, Anlag K, Orban PC, Bock R, Klein R**
and Schutz G. Disruption of the glucocorticoid receptor gene in the nervous system
results in reduced anxiety. *Nat Genet* 23: 99-103, 1999.

58. **Dubois NC, Hofmann D, Kaloulis K, Bishop JM and Trumpp A**. Nestin-Cre
transgenic mouse line Nes-Cre1 mediates highly efficient Cre/loxP mediated
recombination in the nervous system, kidney, and somite-derived tissues. *Genesis* 44:
355-360, 2006.

59. **Liu HT, Tashmukhamedov BA, Inoue H, Okada Y and Sabirov RZ**. Roles of two
types of anion channels in glutamate release from mouse astrocytes under ischemic or
osmotic stress. *Glia* 54: 343-357, 2006.

60. **Sabirov RZ, Merzlyak PG, Okada T, Islam MR, Uramoto H, Mori T, Makino Y,**
Matsuura H, Xie Y and Okada Y. The organic anion transporter SLCO2A1 constitutes
the core component of the Maxi-Cl channel. *EMBO J* 36: 3309-3324, 2017.

61. **Woo DH, Han KS, Shim JW, Yoon BE, Kim E, Bae JY, Oh SJ, Hwang EM,**
Marmorstein AD, Bae YC, Park JY and Lee CJ. TREK-1 and Best1 channels mediate

fast and slow glutamate release in astrocytes upon GPCR activation. *Cell* 151: 25-40,
2012.

62. **Oh SJ, Han KS, Park H, Woo DH, Kim HY, Traynelis SF and Lee CJ**. Protease
activated receptor 1-induced glutamate release in cultured astrocytes is mediated by
Bestrophin-1 channel but not by vesicular exocytosis. *Mol Brain* 5: 38, 2012.

63. **Han YE, Kwon J, Won J, An H, Jang MW, Woo J, Lee JS, Park MG, Yoon BE, Lee
SE, Hwang EM, Jung JY, Park H, Oh SJ and Lee CJ**. Tweety-homolog (Ttyh) family
encodes the pore-forming subunits of the swelling-dependent volume-regulated anion
channel (VRAC(swell)) in the brain. *Exp Neurobiol* 28: 183-215, 2019. doi:
10.5607/en.2019.28.2.183

64. **Ye ZC, Wyeth MS, Baltan-Tekkok S and Ransom BR**. Functional hemichannels in
astrocytes: a novel mechanism of glutamate release. *J Neurosci* 23: 3588-3596, 2003.

65. **Duan S, Anderson CM, Keung EC, Chen Y, Chen Y and Swanson RA**. P2X7
receptor-mediated release of excitatory amino acids from astrocytes. *J Neurosci* 23:
1320-1328, 2003.

66. **Iglesias R, Dahl G, Qiu F, Spray DC and Scemes E**. Pannexin 1: the molecular
substrate of astrocyte "hemichannels". *J Neurosci* 29: 7092-7097, 2009.

67. **Pervaiz S, Kopp A, von KL and Stauber T**. Absolute protein amounts and relative
abundance of volume-regulated anion channel (VRAC) LRRC8 subunits in cells and
tissues revealed by quantitative immunoblotting. *Int J Mol Sci* 20: 5879, 2019. doi:
10.3390/ijms20235879

68. **Knecht DA, Zeziulia M, Bhavsar MB, Puchkov D, Maier H and Jentsch TJ**.
LRRC8/VRAC volume-regulated anion channels are crucial for hearing. *J Biol Chem*
300: 107436, 2024. doi: 10.1016/j.jbc.2024.107436

69. **Concepcion AR, Wagner LE, Zhu J, Tao AY, Yang J, Khodadadi-Jamayran A,
Wang YH, Liu M, Rose RE, Jones DR, Coetzee WA, Yule DI and Feske S**. The
volume-regulated anion channel LRRC8C suppresses T cell function by regulating cyclic
dinucleotide transport and STING-p53 signaling. *Nat Immunol* 23: 287-302, 2022. doi:
10.1038/s41590-021-01105-x

70. **Yanushkevich S, Zieminska A, Gonzalez J, Anazco F, Song R, Arias-Cavieres A,
Granados ST, Zou J, Rao Y and Concepcion AR**. Recent advances in the structure,
function and regulation of the volume-regulated anion channels and their role in
immunity. *J Physiol* 2024. doi: 10.1113/JP285200

71. **Ghosh A, Khandelwal N, Kumar A and Bera AK**. Leucine-rich repeat-containing 8B
protein is associated with the endoplasmic reticulum Ca(2+) leak in HEK293 cells. *J Cell
Sci* 130: 3818-3828, 2017.

72. **Quinodoz M, Rutz S, Peter V, Garavelli L, Innes AM, Lehmann EF, Kellenberger S,**
901   **Peng Z, Barone A, Campos-Xavier B, Unger S, Rivolta C, Dutzler R and Superti-**
902   **Furga A**. De novo variants in LRRC8C resulting in constitutive channel activation cause
903   a human multisystem disorder. *EMBO J* 44: 413-436, 2025. doi: 10.1038/s44318-024-
904   00322-y

FIGURE LEGENDS

Figure 1: LRRC8A-containing VRACs are the main conduit for swelling-activated release of cytosolic glutamate.

A1, Breeding strategy for brain-specific deletion of LRRC8A using *Nestin*[Cre] recombination. The four genotypes generated are: *Lrrc8a*[flox/+] (fl/+), *Lrrc8a*[flox/flox] (fl/fl), *Lrrc8a*[flox/+]:*Nestin*[Cre/+] (HET), and *Lrrc8a*[flox/flox]:*Nestin*[Cre/+] (KO). **A2**, Representative live image of LRRC8A HET astrocytes expressing the Cre reporter Ai9. TdTomato fluorescence was captured at $320\times$ magnification. **B**, Western blot analysis of LRRC8A expression in primary astrocyte cultures from the four genotypes shown in (A1). **B1**, Representative full-length Western blot of LRRC8A immuno-reactivity and matching β-actin signal probed on the same membrane. **B2**, Quantification of LRRC8A expression normalized to β-actin and presented relative to the average signal from fl/+ samples. Data are the mean values ± SD from five cultures per group. ****p <0.0001, one-way ANOVA with Tukey's multiple comparisons test. **C**, VRAC activity in primary astrocytes assessed as swelling-activated release of D-[^{3}H]aspartate in response to a 30% reduction in medium osmolality (290 to 200 mOsm). Data are the means ± SD from 8–16 independent experiments in at least two different astrocyte cultures per genotype. *p < 0.05, ****p < 0.0001, HET (n=8) and KO (n=8) vs. controls (fl/+ and fl/fl, n=16), respectively. ##p < 0.01, HET vs. KO. One-way ANOVA of maximal release values with Tukey's multiple comparisons test. **Inset:** Hypoosmotic response in LRRC8A KO astrocytes shown on an expanded scale. **D**, VRAC activity in primary astrocytes exposed to a 50% reduction in medium osmolality (from 290 to 150 mOsm). Data are the means ± SD from 8–11 independent experiments in at least two different astrocyte cultures per genotype. ****p < 0.0001, HET (n=9) and KO (n=8) vs. controls (fl/+ and fl/fl, n=11); #p < 0.05, HET vs. KO. One-way ANOVA with Tukey's multiple comparisons test. **Inset:** Hypoosmotic response in LRRC8A KO astrocytes shown on an expanded scale.

Figure 2: Relative abundance of *Lrrc8a-e* transcripts in mouse brain and primary astrocyte cultures.

A, Quantification of mRNA expression levels for *Lrrc8a–e* in mouse forebrain lysates using qRT-PCR. Expression levels were normalized to ribosomal *Rpl13a* within the same sample. Data are the means ± SD of 7 mouse forebrains. **B**, RNA-seq analysis of *Lrrc8a-e* expression in 7 mouse forebrain regions from 5 mice of both sexes (data downloaded from the Human Protein Atlas). **C**, Quantification of mRNA expression levels for *Lrrc8a–e* in primary astrocyte cultures measured with qRT-PCR and normalized to ribosomal *Rpl13a*. Data are the means ± SD of 6 independent cultures.

Figure 3: Functional contribution of individual LRRC8A-E subunits to glutamate-permeable VRACs.

A1, Quantification of *Lrrc8a-e* mRNA expression levels in mouse astrocyte cultures treated with negative control siRNA (siNC). Expression levels were normalized to the housekeeping *Rpl13a*.

Data are the means ± SD of 4 independently prepared cell cultures. **A2**, Schematic of a radiotracer
assay of VRAC activity (created with BioRender.com, see Methods for additional details). **B-F**,
Effects of gene-specific siRNA constructs on *Lrrc8a-e* mRNA expression levels 48 h post-
transfection and swelling-activated D-[^{3}H]aspartate release 96 h post transfection. **B1-F1**, Relative
Lrrc8a-e expression was normalized to *Rpl13a* and compared to siNC. Values are the means ±
SD of 3-4 independent transfections in at least two different astrocyte cultures. *p<0.05, **p<0.01,
***p<0.001, one-population t-test with Bonferroni correction for multiple comparisons. **B2-F2**,
Integral D-[^{3}H]aspartate release values in cells exposed for 10 min to either isoosmotic (Basal) or
hypoosmotic (Hypo) conditions (see Methods). With the exception of *F2*, data are the means ±
SD of 6-9 independent transfections in two independently prepared cell cultures. For *Lrrc8e*
knockdowns in *F2*, data are the means ± SD of 3 independent transfections in one astrocyte
culture. ** p <0.01, *** p <0.001, ****p <0.0001, vs. siNC, repeated measures two-way ANOVA
with Tukey's multiple comparisons test (for clarity only the effect of siRNA under hypoosmotic
conditions are shown).

Figure 4: Non-redundant roles of LRRC8C and LRRC8D subunits in glutamate-permeable
VRACs.

A, Validation of siRNA efficacy in double knockdown experiments targeting *Lrrc8c* and *Lrrc8d*.
Astrocytes were treated with siRNA targeting *Lrrc8c* (siC1), *Lrrc8d* (siD4), or their combination
(siC1 + siD4). Efficacy of gene-specific knockdowns was verified by qRT-PCR. Data are the mean
normalized expression values ± SD in three independently transfected cell cultures. **p<0.01,
*p<0.05, one-population t-test with Bonferroni correction for multiple comparisons. **B**, Effect of
siC1, siD4, or siC1+siD4 on VRAC-mediated D-[^{3}H]aspartate release from cultured astrocytes
exposed to a 30% reduction in medium osmolality measured in a 12-well plate format. Data are
the means ± SD from 12 independent transfections in three different astrocyte cultures. ****p <
0.0001 vs. siNC, repeated measures two-way ANOVA with Tukey's multiple comparisons test
(only the effects of siRNAs under hypoosmotic conditions are shown). **C**, Kinetics of VRAC-
mediated D-[^{3}H]aspartate release from astrocyte cultures treated with siNC, siC1, or siC1+siD4
and exposed to a 30% reduction in medium osmolality. Data are the means ± SD in four-five
independent transfections. **Inset:** Statistical comparison of the maximal D-[^{3}H]aspartate release
values in the same experiments. *p<0.05, one-way ANOVA with Tukey's multiple comparisons
test. **D**, Kinetics of VRAC-mediated D-[^{3}H]aspartate release from astrocytes treated with siNC,
siC1, siD4, or siC1+siD4 and exposed to a 50% reduction in medium osmolality. Data are the
means ± SD in five-six independent transfections performed in two different astrocyte cultures.
Inset: Statistical comparison of the maximal D-[^{3}H]aspartate release values in the same
experiments. **p<0.01, ****p <0.0001, one-way ANOVA with Tukey's multiple comparisons test.

Figure 5: Functional contribution of LRRC8B to LRRC8C- and LRRC8D-containing
VRACs.

A, Validation of siRNA efficacy in double knockdown experiments targeting *Lrrc8b* and *Lrrc8c*.
Astrocytes were treated with siRNA targeting *Lrrc8b* (siB4), *Lrrc8c* (siC1), or their combination
(siB4+siC1). Efficacy of gene-specific knockdowns was verified by qRT-PCR. Data are the mean
normalized expression values ± SD in three independently transfected cell cultures. **p<0.01,
*p<0.05; one-population t-test with Bonferroni correction for multiple comparisons. **B**, Effects of

siNC, siB4, siC1, siB4+siC1 on VRAC-mediated D-[^{3}H]aspartate release from cultured astrocyte
exposed to a 30% reduction in medium osmolality, measured in a 12-well plate format. Data are
the means ± SD from 6 independent transfections in two different astrocyte cultures. ****p <
0.0001, repeated measures two-way ANOVA with Tukey's multiple comparisons test (only the
effects of siRNAs under hypoosmotic conditions are shown). **C**, qRT-PCR validation of double
siRNA knockdown experiments targeting *Lrrc8b* (siB4) and *Lrrc8d* (siD4). Astrocytes were treated
with siB4, siD4, or their combination (siB4+siD4). Data represent mean normalized expression
values ± SD in three-four independently transfected cell cultures. ***p<0.001, one-population t-
test with Bonferroni correction for multiple comparisons. **D**, Effects of siNC, siB4, siD4, siB4+siD4
on VRAC-mediated D-[^{3}H]aspartate release from cultured astrocyte exposed to a 30% reduction
in medium osmolality measured in a 12-well plate format. Data are the means ± SD from 6
independent transfections in two different astrocyte cultures. ***p<0.001, ****p<0.0001; repeated
measures two-way ANOVA with Tukey's multiple comparisons test (only the effects of siRNAs
under hypoosmotic conditions are shown).

1004

Figure 6: Proposed subunit composition of glutamate-permeable VRACs in astrocytes.

A and **B**, Theoretical predictions of the effects on siRNA knockdowns on VRAC activity based on
two models of VRAC composition. In *Model I*, VRACs are assembled from LRRC8A, LRRC8C,
and LRRC8D subunits, with LRRC8C and LRRC8D functioning as mutually redundant
components. In *Model II*, two independent VRAC populations are formed by either LRRC8A+
LRRC8C or LRRC8A+LRRC8D. In this case, LRRC8C and LRRC8D subunits are not functionally
redundant. **C**, *Experimental outcomes* showing the effects of LRRC8C (siC) and/or LRRC8D (siD)
knockdown on VRAC-mediated D-[^{3}H]aspartate release in mouse astrocytes subjected to
moderate (30% reduction in osmolarity; light bars) or severe (50% reduction in osmolarity; dark
bars) hypotonic stress. These results are consistent with the Model II in which two distinct VRAC
populations (labeled 1 and 2) mediate glutamate permeability. Additional data in the manuscript
indicate that LRRC8B likely plays a structural role in these channels. LRRC8A+LRRC8D-
containing VRACs provide a smaller contribution to swelling-activated glutamate release and their
activity is dependent on the degree of cellular swelling. ***p<0.001, *p=0.012; unpaired t-tests.

1019

TABLES

 Table 1. List of molecular biology tools used in qRT-PCR analyses and RNAi experiments in this
 study.

Target gene	Catalogue # qPCR primers	siRNA name (catalogue #)	siRNA target sequence
Negative Control (NC)	N/A	AllStars NC (SI03650318)	Scrambled (proprietary)
Lrrc8a	QT01167502	Mm_Lrrc8_1 (SI01296127)	CAGCCTCAAGAAGATGGTTAA
		Mm_Lrrc8_3 (SI01296141)	CAGAGTCATCTCAGGGTTAAA
		Mm_Lrrc8_4 (SI01296148)	AATGATGGTTTGGGTATTAAA
Lrrc8b	QT01541666	Mm_LOC433926_1 (SI01235311)	AAGAAGATGGTCAATCTGAAA
		Mm_LOC433926_3 (SI01235325)	ATCCTGTTTGTCCTTATTATA
		Mm_LOC433926_4 (SI01235332)	TAGGAGGATGAACTTTCCTAA
Lrrc8c	QT01042881	Mm_E430036I04Rik_1 (SI00988799)	ATCGGAGTTCTGCAAAGTTTA
		Mm_E430036I04Rik_2 (SI00988806)	GAGAAGTAACATATACATATA
		Mm_E430036I04Rik_4 (SI00988820)	CACCTTATACTGGCTGTTCTA
Lrrc8d	QT02255085	LRRC5_1 (SI01296043)	TTGCAAGTTATTACAAGATAA
		LRRC5_2 (SI01296050)	CAGCATGACAATCCTGGACAA
		LRRC5_3 (SI01296057)	CTGCTAAAGTATATGATTAAA
		LRRC5_4 (SI01296064)	AAGCCCTGTTTGAGAAAGTAA
Lrrc8e	QT01046269	Mm_1810049O03Rik_1 (SI00806043)	CACATTTAGCTATGTAATAAA
		Mm_1810049O03Rik_2 (SI00806050)	CTGAAGGTGATCTGTGTTAAA
		Mm_1810049O03Rik_4 (SI00806064)	CCCAACAAGGATCAGCATATA
GAPDH	QT01658692	N/A	N/A
Rpl13a	QT00267197	N/A	N/A
RPS20	QT00251433	N/A	N/A

LRRC8A-containing VRACs are the main conduit for swelling-activated release of cytosolic glutamate.

A1, Breeding strategy for brain-specific deletion of LRRC8A using *Nestin*[Cre] recombination. The four genotypes generated are: *Lrrc8a*[flox/+] (fl/+), *Lrrc8a*[flox/flox] (fl/fl), *Lrrc8a*[flox/+]:*Nestin*[Cre/+] (HET), and *Lrrc8a*[flox/flox]:*Nestin*[Cre/+] (KO). **A2**, Representative live image of LRRC8A HET astrocytes expressing the Cre reporter Ai9. TdTomato fluorescence was captured at 320× magnification. **B**, Western blot analysis of LRRC8A expression in primary astrocyte cultures from the four genotypes shown in (A1). **B1**, Representative full-length Western blot of LRRC8A immunoreactivity and matching β-actin signal probed on the same membrane. **B2**, Quantification of LRRC8A expression normalized to β-actin and presented relative to the average signal from fl/+ samples. Data are the mean values ± SD from five cultures per group. ****$p < 0.0001$, one-way ANOVA with Tukey's multiple comparisons test. **C**, VRAC activity in primary astrocytes assessed as swelling-activated release of D-[^{3}H]aspartate in response to a 30% reduction in medium osmolality (290 to 200 mOsm). Data are the means ± SD from 8–16 independent experiments in at least two different astrocyte cultures per genotype. *$p < 0.05$, ****$p < 0.0001$, HET (n=8) and KO (n=8) vs. controls (fl/+ and fl/fl, n=16), respectively. ##$p < 0.01$, HET vs. KO. One-way ANOVA of maximal release values with Tukey's multiple comparisons test. **Inset:** Hypoosmotic response in LRRC8A KO astrocytes shown on an expanded scale. **D**, VRAC activity in primary astrocytes exposed to a 50% reduction in medium osmolality (from 290 to 150 mOsm). Data are the means ± SD from 8–11 independent experiments in at least two different astrocyte cultures per genotype. ****$p < 0.0001$, HET (n=9) and KO (n=8) vs. controls (fl/+ and fl/fl, n=11); #$p < 0.05$, HET vs. KO. One-way ANOVA with Tukey's multiple comparisons test. **Inset:** Hypoosmotic response in LRRC8A KO astrocytes shown on an expanded scale.

Relative abundance of *Lrrc8a-e* transcripts in mouse brain and primary astrocyte cultures.

A, Quantification of mRNA expression levels for *Lrrc8a–e* in mouse forebrain lysates using qRT-PCR. Expression levels were normalized to ribosomal *Rpl13a* within the same sample. Data are the means ± SD of 7 mouse forebrains. **B**, RNA-seq analysis of *Lrrc8a-e* expression in 7 mouse forebrain regions from 5 mice of both sexes (data downloaded from the Human Protein Atlas). **C**, Quantification of mRNA expression levels for *Lrrc8a–e* in primary astrocyte cultures measured with qRT-PCR and normalized to ribosomal *Rpl13a*. Data are the means ± SD of 6 independent cultures.

Functional contribution of individual LRRC8A-E subunits to glutamate-permeable VRACs.

A1, Quantification of *Lrrc8a-e* mRNA expression levels in mouse astrocyte cultures treated with negative control siRNA (siNC). Expression levels were normalized to the housekeeping gene *Rpl13a*. Data are the means ± SD of 4 independently prepared cell cultures. **A2**, Schematic of a radiotracer assay of VRAC activity (created with BioRender.com, see Methods for additional details). **B-F**, Effects of gene-specific siRNA constructs on *Lrrc8a-e* mRNA expression levels 48 h post-transfection and swelling-activated D-[^{3}H]aspartate release 96 h post-transfection. **B1-F1**, Relative *Lrrc8a-e* expression was normalized to *Rpl13a* and compared to siNC. Values are the means ± SD of 3-4 independent transfections in at least two different astrocyte cultures. *p<0.05, **p<0.01, ***p<0.001, one-population t-test with Bonferroni correction for multiple comparisons. **B2-F2**, Integral D-[^{3}H]aspartate release values in cells exposed for 10 min to either isoosmotic (Basal) or hypoosmotic (Hypo) conditions (see Methods). With the exception of *F2*, data are the means ± SD of 6-9 independent transfections in two independently prepared cell cultures. For *Lrrc8e* knockdowns in *F2*, data are the means ± SD of 3 independent transfections in one astrocyte culture. ** p <0.01, *** p <0.001, ****p <0.0001, vs. siNC, repeated measures two-way ANOVA with Tukey's multiple comparisons test (for clarity only the effect of siRNA under hypoosmotic conditions are shown).

Non-redundant roles of LRRC8C and LRRC8D subunits in glutamate-permeable VRACs.

A, Validation of siRNA efficacy in double knockdown experiments targeting *Lrrc8c* and *Lrrc8d*. Astrocytes were treated with siRNA targeting *Lrrc8c* (siC1), *Lrrc8d* (siD4), or their combination (siC1 + siD4). Efficacy of gene-specific knockdowns was verified by qRT-PCR. Data are the mean normalized expression values ± SD in three independently transfected cell cultures. **p<0.01, *p<0.05, one-population t-test with Bonferroni correction for multiple comparisons. **B**, Effect of siC1, siD4, or siC1+siD4 on VRAC-mediated D-[³H]aspartate release from cultured astrocytes exposed to a 30% reduction in medium osmolality measured in a 12-well plate format. Data are the means ± SD from 12 independent transfections in three different astrocyte cultures. ****p < 0.0001 vs. siNC, repeated measures two-way ANOVA with Tukey's multiple comparisons test (only the effects of siRNAs under hypoosmotic conditions are shown). **C**, Kinetics of VRAC-mediated D-[³H]aspartate release from astrocyte cultures treated with siNC, siC1, or siC1+siD4 and exposed to a 30% reduction in medium osmolality. Data are the means ± SD in 4-5 independent transfections. **Inset:** Statistical comparison of the maximal D-[³H]aspartate release values in the same experiments. *p<0.05, one-way ANOVA with Tukey's multiple comparisons test. **D**, Kinetics of VRAC-mediated D-[³H]aspartate release from astrocytes treated with siNC, siC1, siD4, or siC1+siD4 and exposed to a 50% reduction in medium osmolality. Data are the means ± SD in 5-6 independent transfections performed in two different astrocyte cultures. **Inset:** Statistical comparison of the maximal D-[³H]aspartate release values in the same experiments. **p<0.01, ****p <0.0001, one-way ANOVA with Tukey's multiple comparisons test.

Functional contribution of LRRC8B to LRRC8C- and LRRC8D-containing VRACs.

A, Validation of siRNA efficacy in double knockdown experiments targeting *Lrrc8b* and *Lrrc8c*. Astrocytes were treated with siRNA targeting *Lrrc8b* (siB4), *Lrrc8c* (siC1), or their combination (siB4+siC1). Efficacy of gene-specific knockdowns was verified by qRT-PCR. Data are the mean normalized expression values ± SD in 3 independently transfected cell cultures. **p<0.01, *p<0.05; one-population t-test with Bonferroni correction for multiple comparisons. **B**, Effects of siNC, siB4, siC1, siB4+siC1 on VRAC-mediated D-[^{3}H]aspartate release from cultured astrocyte exposed to a 30% reduction in medium osmolality, measured in a 12-well plate format. Data are the means ± SD from 6 independent transfections in two different astrocyte cultures. ****p < 0.0001, repeated measures two-way ANOVA with Tukey's multiple comparisons test (only the effects of siRNAs under hypoosmotic conditions are shown). **C**, qRT-PCR validation of double siRNA knockdown experiments targeting *Lrrc8b* (siB4) and *Lrrc8d* (siD4). Astrocytes were treated with siB4, siD4, or their combination (siB4+siD4). Data represent mean normalized expression values ± SD in 3-4 independently transfected cell cultures. ***p<0.001, one-population t-test with Bonferroni correction for multiple comparisons. **D**, Effects of siNC, siB4, siD4, siB4+siD4 on VRAC-mediated D-[^{3}H]aspartate release from cultured astrocyte exposed to a 30% reduction in medium osmolality measured in a 12-well plate format. Data are the means ± SD from 6 independent transfections in two different astrocyte cultures. ***p < 0.001, ****p<0.0001; repeated measures two-way ANOVA with Tukey's multiple comparisons test (only the effects of siRNAs under hypoosmotic conditions are shown).

Figure 6 • M.L. Chandler et al.

Proposed subunit composition of glutamate-permeable VRACs in astrocytes

A and **B**, Theoretical predictions of the effects on siRNA knockdowns on VRAC activity based on two models of VRAC composition. In *Model I*, VRACs are assembled from LRRC8A, LRRC8C, and LRRC8D subunits, with LRRC8C and LRRC8D functioning as mutually redundant components. In *Model II*, two independent VRAC populations are formed by either LRRC8A+LRRC8C or LRRC8A+LRRC8D. In this case, LRRC8C and LRRC8D subunits are not functionally redundant. **C**, *Experimental outcomes* showing the effects of LRRC8C (siC) and/or LRRC8D (siD) knockdown on VRAC-mediated D-[³H]aspartate release in mouse astrocytes subjected to moderate (30% reduction in osmolarity; light bars) or severe (50% reduction in osmolarity; dark bars) hypotonic stress. These results are consistent with the Model II in which two distinct VRAC populations (labeled 1 and 2) mediate glutamate permeability. Additional data in the manuscript indicate that LRRC8B likely plays a structural role in these channels. LRRC8A+LRRC8D-containing VRACs provide a smaller contribution to swelling-activated glutamate release and their activity is dependent on the degree of cellular swelling. ***p<0.001, *p=0.012; unpaired t-tests.