1   **Resistance, heteroresistance and fitness costs drive colistin treatment failure during**

2   ***Acinetobacter baumannii* pneumonia**

4   **Running title:** Evolution of colistin resistance in mouse infection model

7   Juan Hernandez-Bird[1]; Bixi He[1]; Leah M. VanOtterloo[2]; Elizabeth B. Billings[1]; Wenwen Huo[1];

8   Gabriella I C Teodoro[1]; Haley Echlin[3]; Jason M. Rosch[3]; M. Stephen Trent[2]; Ralph R. Isberg[1*]

[1] Department of Molecular Biology and Microbiology, Tufts University School of Medicine,
MA, USA

[2] Department of Microbiology, College of Arts and Sciences, University of Georgia, GA, USA

[3] Department of Infectious Diseases, St Jude Children's Research Hospital, TN, USA

*Corresponding author contact: ralph.isberg@tufts.edu

Abstract:

Acinetobacter baumannii is an ESKAPE pathogen linked to healthcare-associated diseases. Due

to evolved resistance, last-resort antibiotics such as the lipooligosaccharide (LOS)-targeting

colistin are increasingly used to treat multidrug-resistant isolates. To track the evolution of

colistin resistance within a host, we performed sequential oropharyngeal infections in

immunocompetent or immune-depleted mice in the presence of inhaled colistin. Both resistant

and heteroresistant *A. baumannii* strains emerged with *pmrB* mutations that efficiently competed

with the susceptible parent in the presence of colistin. These *pmrB* mutants had a fitness cost in

the absence of colistin treatment but retained their ability to colonize the host. In contrast, LOS-

deficient *A. baumannii* mutants removed the target of colistin, but such mutants were unable to

colonize the lung. The two pathogenic *pmrB* mutants showed clear evidence of LOS

modification, which was linked to increased transcription of LOS modification enzymes,

including the product of the cryptic *eptA* gene. Spontaneous insertion mutations that caused

hyperexpression of *eptA* allowed the heteroresistant mutant to develop clinically-significant

colistin resistance. Insertion mutations upstream of the *eptA* gene or those disrupting *hns*, which

encodes a small histone-like protein, resulted in increased *eptA* transcript, linking expression of

this protein to clinically significant resistance. A resistant variant derived from the

heteroresistant parent was stable in the absence of drug, but continued passaging selected for

colistin-resensitized pseudorevertants that were largely due to disruption of the LOS

modification enzymes. Therefore, colistin heteroresistance is an early stage in the stepwise

acquisition of stable colistin resistance in *A. baumannii*.

Significance:

The mutational pathways leading to antibiotic-resistant infections and the role of the immune system in preventing them are poorly understood. Here we employed a colistin-treated mouse pneumonia model of *Acinetobacter baumannii* and observed the evolution of colistin-resistant and heteroresistant mutants in immune-depleted and immunocompetent hosts, respectively. We show that mutations that result in alteration of colistin's target drive evolutionary pathways to resistance, whereas removal of the target is unlikely to be clinically significant. We demonstrated that the heteroresistant mutant generates subpopulations with higher levels of resistance through insertion mutations in specific gene regions. This study furthers our understanding of how resistance emerges during infection and provides a genetic explanation for the transition from colistin heteroresistance to full resistance.

Introduction:

Antibiotic-resistant infections caused by the ESKAPE group of pathogens are an urgent global health concern, with lethality predicted to surpass cancer by 2050(1). Belonging to this group is *Acinetobacter baumannii*, a Gram-negative bacterium with a remarkable ability to evolve resistance to currently employed antibiotics(2). As an opportunistic pathogen, *A. baumannii* mainly affects wounded or immunocompromised patients in healthcare settings with respiratory, bloodstream and urinary tract infections being its most concerning clinical manifestations(3, 4). The World Health Organization currently classifies carbapenem-resistant *A. baumannii* in the highest priority category for research and development of treatment options(5). Therefore, understanding how this organism evolves to give rise to drug-resistant variants is of the upmost importance.

Resistance to the first-line antimicrobial carbapenems is increasingly common in *A.*

baumannii clinical isolates(6), so last-resort antibiotics are currently being employed against

extensively drug-resistant *A. baumannii*(7). Colistin is one such antimicrobial, but its usefulness

is limited due to concerns about its nephrotoxicity. New inhalation therapies are being pursued

for the treatment of respiratory infections in hopes of reducing toxic side effects(8, 9). The drug

is a positively charged peptide that binds to the negatively charged phosphate groups on the lipid

A component of lipopolysaccharide (LPS), or lipooligosaccharide (LOS) in the case of *A.*

baumannii, leading to envelope disruption and cell lysis(7).

Gram-negative bacteria have evolved multiple strategies to modify the lipid A component

of LPS/LOS to increase resistance to antimicrobial peptides(10). In *A. baumannii,* colistin

resistance is mainly caused by mutations within the *pmrCAB* operon(11). The PmrAB cassette is

a two-component system that responds to low pH and select divalent cations (Fe^{2+}, Zn^{2+}, Al^{3+}),

controlling the expression of multiple genes including the *pmrC* and the *naxD* operons(12).

pmrC, also known as *eptA*, encodes a phosphoethanolamine transferase that adds

phosphoethanolamine to the 1- and 4' phosphoryl groups in lipid A(13). *naxD* encodes a YdjC

family deacetylase that plays a part in the addition of galactosamine to the 1- phosphoryl group

in lipid A(14). These modifications reduce both the negative charge and LOS affinity for

polymyxins such as colistin(13). Mutations in the *pmrAB* genes can lead to the constitutive

activation of this system by the PmrA positive regulator, resulting in polymyxin resistance. In

addition, some strains of *Acinetobacter* possess an orphan homologue of the PmrC/EptA

phosphoethanolamine transferase, that is also under the control of PmrAB. For simplicity, we

will refer to the orphan copy as *eptA* and the *pmrAB*-linked copy as *pmrC.* Clinical isolates with

high colistin resistance have been characterized that overexpress *eptA* transcript as a

consequence of insertion sequence transposition into regions upstream of *eptA*(15, 16).

Mutations in *pmrAB* have also been linked to colistin heteroresistance in *A.*

baumannii(17). Antibiotic heteroresistant strains generate subpopulations resistant to the exposed

antibiotic at much higher frequencies than observed for sensitive strains(18). Heteroresistant

pathogens are troubling because current clinical microbiology methods frequently classify them

as susceptible, and their genetic background could provide an intermediate step towards the

evolution of traditional resistance(19–21). The genetic mechanisms facilitating this phenomenon

are poorly understood, but recent evidence points to transient gene amplification in the resistant

subpopulation, which occurs at a higher frequency than observed for spontaneous mutations(22).

Amplification and mutations in genes that regulate or mediate the modification of lipid A have

been reported to contribute to colistin heteroresistance in pathogens other than

Acinetobacter(23–26). The genetic basis for colistin heteroresistance in *A. baumannii*, however,

appears unrelated to amplification.

A. baumannii is one of the few known species with the ability to survive with an LOS-

deficient outer membrane(27). LOS-deficient mutants are readily isolated in a subset of clinical

isolates during *in-vitro* experimental evolution to polymyxins but are rarely reported in clinical

studies(28). Absence of LOS, usually due to mutations in the early LOS biosynthesis genes,

results in high-level colistin resistance but is accompanied by outer membrane instability and a

high fitness cost(29). Nevertheless, these LOS-deficient mutants may provide transition states to

more virulent, drug-resistant mutants(30, 31). Inactivation of PBP1A glycosyltransferase activity

allows viability in the presence of LOS knockout mutations(32). Additionally, mutations in the

pldA and *mla* genes, enzymes that act to reduce glycerophospholipid content in the outer

membrane, alleviate the severe growth defect in LOS-deficient strains(27). It is unknown if these

suppressor mutations allow LOS-deficient strains to retain their virulence.

Previously, we performed serial passaging of the low virulence *A. baumannii*

ATCC17978 strain in a murine pneumonia model during treatment with ciprofloxacin(33). In

immune-depleted animals, drug-tolerant/persistent mutants arose that resulted in antibiotic

treatment failure and the emergence of second-step drug-resistant mutants. Neither class of

mutants achieved high population levels in the lines passaged in immunocompetent hosts(33,

34). More recent multi-drug resistant *A. baumannii* clinical isolates have been proposed as more

relevant for infection modeling due to their ability to cause lethal pneumonia even in the absence

of immune depletion(35). One of these strains is *A. baumannii* LAC-4, a multi-drug resistant

specimen that was isolated from a Los Angeles County outbreak in 1997(36–38). In this study,

we performed experimental evolution in this highly mouse-virulent strain, identifying multiple

routes to colistin-resistant virulent isolates, as well as forbidden routes that result in avirulent

strains.

Results:

Sequential lung infections in the presence of nebulized colistin select for pmrB

mutations.

To better simulate the conditions in which *A. baumannii* evolves colistin resistance, we

performed 16 sequential lung infections with the LAC-4 strain in both immune-depleted and

immunocompetent mice treated with inhaled colistin (Fig. 1A). Immune-depleted animals

received two intraperitoneal injections of cyclophosphamide, a chemotherapeutic agent that

induces profound neutropenia and significantly reduces lymphocyte and monocyte counts(39,

40). Lung infections were established via oropharyngeal inoculation in both sets of animals.

Three separate lines were maintained per condition, and lung bacterial burden after 24-hour

infections was quantified (Figs. 1B, C). The inoculum CFU for the immune-depleted lines was

reduced in the second and fifth passage to ensure effective antibiotic selection throughout the

experiment (2×10^2 CFU below the expected bacterial burden; Fig. 1B). Line A showed

evidence of antibiotic treatment failure after 13 passages in the immune-depleted animals, with

bacterial burdens approaching that observed in untreated animals. In contrast, colistin treatment

consistently reduced bacterial burden in the other two immune-depleted lines and in the

immunocompetent lines throughout all passages (Figs. 1B, C).

Colistin-resistant *A. baumannii* isolates are defined by the Clinical Laboratory Standards

Institute (CLSI) as being able to grow at colistin concentrations $\geq 4\mu g/mL$(41). The parental

LAC-4 strain is colistin-susceptible, so saved pools were plated on LB agar supplemented with

0-4µg/mL colistin to detect the presence and abundance of colistin-resistant isolates (Figs. 1D, E

and SI Appendix, Fig. S1). During passaging in the immune-depleted animals, colistin-resistant

colonies emerged in all three lines, but only in line A did the proportion of resistant clones

increase and overgrow the susceptible ancestor (Fig. 1D). Whole genome sequencing (WGS) of

saved resistant isolates from line A revealed they all have the *pmrB(*T235I) mutation that has

been previously described in polymyxin-resistant *A. baumannii* strains(42).

Colistin-resistant colonies emerged in lines B and C in the last three passages through

immunocompetent animals (Fig. 1E). WGS of resistant isolates in line C revealed that they all

possessed the *pmrB*(A236E) mutation that had been associated previously with polymyxin

resistance(43). Due to their increase in proportion on solid medium containing 4µg/ml colistin on

successive passages through mice, the *pmrB*(T235I) and *pmrB*(A236E) clones were selected for

further characterization. Importantly, the *pmrB*(T235I) strain employed in this study contains

three additional IS*Aba* insertion mutations discussed later on. Additional mutants appeared

throughout the passaging that grew at low colistin concentrations. These were isolated, subjected

to MIC determination and WGS, but they were not studied further (Dataset S1 and SI Appendix,

Fig. S1). Thus, colistin treatment in both host immune states selected for *A. baumannii* isolates

with mutations in the region encoding the PmrB histidine kinase domain during passage in the

experimental pneumonia model.

The fitness cost of the LAC-4 pmrB mutants.

Mutations that promote antibiotic resistance are often associated with fitness costs(44,

45). The continuous spread of drug-resistant bacteria requires that resistant mutants compete

efficiently with fully virulent parental organisms in mammalian tissues(46). Therefore, we tested

the hypotheses that mutations selected during lung infections would confer resistance without

compromising the ability of the LAC-4 to colonize the host, and that these mutants would

overgrow the parental strain in the presence of colistin.

The two evolved *pmrB* mutants were compared to WT LAC-4 during growth in broth

culture. The *pmrB*(T235I) mutant, which had been selected in the immune-depleted animal, had

a significant increase in broth doubling time compared to WT (Fig. 2A; $p \leq 0.05$). In contrast, the

pmrB(A236E) mutant that was selected in the immunocompetent animal showed no significant

change in doubling time during broth growth (Fig. 2A). To determine the relative fitness of the

evolved *pmrB* mutants compared to WT during lung infection, strains were mixed 1:1

(mutant:WT) and competition experiments were performed during experimental lung infections

in the presence or absence of immune depletion (Figs. 2B, C). In the absence of colistin

treatment, the evolved *pmrB* mutants exhibited a competition deficiency against WT in the

immunocompetent host, with the *pmrB*(T235I) mutant exhibiting the larger defect (Fig. 2B). In

contrast, there was no significant competition defect against WT when the competition was

performed in immune-depleted hosts (Fig. 2C). The gentamycin-susceptible LAC-4 strain

(*ΔpABLAC2*) was included in this study since it exhibits no significant change in colistin

sensitivity and growth rate *in-vitro,* allowing us to perform WT vs WT mouse competitions as a

control (Figs. 2A-C). These results demonstrate that in the absence of colistin, *A. baumannii*

pmrB mutants experimentally evolved during lung infections can be negatively selected in the

presence of an intact immune response.

To reconstruct the fitness advantage of *pmrB* mutants relative to the parental control in

the presence of colistin, each mutant was mixed with WT at 5:95 ratio and passaged through

either immune-depleted or immunocompetent mice treated with colistin (Figs. 2D, E). After only

two successive passages in the lung, the *pmrB*(T235I) mutant lines reached 100% of the total

population in the immune-depleted host (Fig. 2E). Notably, during colistin treatment, this mutant

also reached a high proportion of the population (>85%) in all lines after three passages in the

immunocompetent host (Fig. 2D), even though this mutant competed poorly with wild type in

the absence of drug pressure (Fig. 2B). The *pmrB*(A236E) mutant also showed evidence for

effective competition during passaging in both host conditions, but constituted smaller

proportions of the populations than that observed for the *pmrB*(T235I) mutant, with competitions

not uniform among all the lines (compare Figs. 2D and 2E). Therefore, the *pmrB*(T235I) mutant

selected in the immune-depleted conditions established a colistin-resistant population more

efficiently than the *pmrB*(A236E) strain, even in the presence of an intact immune response.

Colistin-resistant LOS-deficient strains show poor fitness during murine pneumonia.

As the colistin-resistant mutations identified in the murine pneumonia model resulted in

altered regulation of the PmrAB control system but did not directly disrupt LOS biosynthesis

(Fig. 3A), we asked whether LOS biosynthesis dysfunction was an available mutational pathway

to achieve colistin resistance during LAC-4 infection. A subset of *A. baumannii* clinical isolates

can acquire resistance to polymyxins through mutations causing defective LOS biosynthesis

(LOS[(-)])(28). To determine whether the parental LAC-4 strain could tolerate LOS[(-)] mutations, we

first selected for colistin-resistant LAC-4 isolates and then screened for collateral vancomycin

sensitivity (Fig. 3B). LOS[(-)] mutants have a compromised outer membrane resulting in

susceptibility to antibiotics usually ineffective against Gram-negative bacteria(27). We identified

one such isolate and whole genome sequencing revealed a single *lpxA* G142R (*lpxA*[*]) mutation,

predicted to disrupt LOS synthesis (Dataset S2). SDS-polyacrylamide gel analysis of protease-

treated extracts showed that, in contrast to the *pmrB* mutants, the *lpxA*[*]-containing strain showed

no evidence of LOS comigrating with the parental WT LAC-4 strain (Fig. 3A).

Due to low fitness, LOS[(-)] strains revert to LOS[(+)] or acquire second-site mutations that

alter the glycerophospholipid composition of the outer membrane(27, 28, 32, 47). LOS[(-)]

suppressor mutants were selected by growing the LAC-4 *lpxA*[*] mutant on LB solid medium at

42°C, a temperature that exacerbates its growth defect and results in a fine lawn of growth (Fig

3B). After selection, 16 non-reverted colonies that grew above the lawn were saved and

subjected to WGS. Mutations identified in this fashion included a frameshift in *pldA,* as well as

missense mutations in the *ponA* gene, both of which had been previously identified as increasing

the growth rate of LOS[(-)] strains (Dataset S2) (27, 32). A quadruple mutant containing a mutation

in *msbA*, a gene involved in lipooligosaccharide transport across the membrane, was also

isolated(48). Additionally, *pldA* mutants with second step mutations in ABLAC_25350, which

encodes a hypothetical protein, were also isolated but were not studied further (Dataset S2).

Three non-identical LOS$^{(-)}$ suppressor mutants were chosen for further analysis, each of

which had either a *pldA* mutation or *pldA ponA* mutations (Fig. 3B). The three suppressor

mutations showed no evidence of the reversion to LOS production, as the LOS profile of

proteolyzed samples fractionated on SDS gels were indistinguishable from the *lpxA** mutant (Fig.

3A). Furthermore, analysis by thin layer chromatography (TLC) showed that these suppressor

mutants remained LOS deficient (Fig. 4C). Growth rates in broth cultures were determined for

each of the LOS$^{(-)}$ strains. The *lpxA** mutant had a drastic increase in doubling time (38mins)

compared to its WT parent (22mins) (compare Figs. 2A and 3C). All three LOS deficient

suppressor mutants exhibited significant improvements in doubling time (32-35mins; Fig. 3C).

Therefore LAC-4 tolerates LOS deficiency, and similarly to other *A. baumannii* strains,

mutations in the *pldA* and *ponA* genes enhance fitness.

Although readily selected *in-vitro*, LOS$^{(-)}$ clinical isolates have rarely, if ever, been

observed and none emerged during passaging(49). Few studies have tested the virulence of

LOS$^{(-)}$ *Acinetobacter baumannii* mutants(50–52), and none, to our knowledge, have tested the

impact of suppressor mutations on its pathogenesis. To determine their relative defects during

murine pneumonia, the LOS$^{(-)}$ strains were mixed at a 1:1 ratio with WT, inoculated into either

immune-depleted or immunocompetent mice, and competitive indices were determined (Figs.

3D, E). In both host states, all LOS$^{(-)}$ mutants had a devastating competition defect relative to

WT, varying from 10^{-4} – 10^{-5} compared to the initial inoculum (Figs. 3D, E). This colonization

efficiency was more than 1000X lower than observed for the *pmrB* mutants that had been

selected during lung infection (compare Figs. 3D, E vs Figs. 2B, C). The suppressor mutations

showed a fractional increase in the competitiveness relative to the parental *lpxA*[*] strain when

competitions were performed specifically in immunocompetent animals, with *lpxA*[*] *pldA*[*] *ponA*[*]

showing a significant increase in competitiveness (Fig. 3D). Even so, these suppressor mutations

fail to compensate for the drastic colonization defects exhibited by LOS[(-)] strains.

Paths to colistin resistance occur via LOS modification and elimination.

To both confirm and quantify the level of colistin resistance in our evolved strains,

colistin E-test strip analysis was performed (Fig. 4A). All LOS[(-)] strains surpassed the breakpoint

for clinical resistance ($\geq$4μg/mL) with the suppressor mutants showing colistin minimum

inhibitory concentration (MIC) values that were higher than the *lpxA*[*] parent (Fig. 4A). The WT

LAC-4 strain was confirmed to be colistin-susceptible and the *pmrB*(T235I) mutant was highly

colistin-resistant based on this MIC determination method (Fig. 4A). Surprisingly, the

pmrB(A236E) mutant only had a slight increase in colistin MIC compared to WT based on the E-

test determination and was clearly still susceptible based on the clinical drug resistance

breakpoint definition (Fig. 4A).

As mutations in *pmrB* appear to be preferentially selected during disease because of their

relatively high fitness compared to those lacking LOS(50–52), the genetic and biochemical basis

of the two *pmrB* alleles was pursued. Similar to our observations modeling disease during mouse

passage, colistin-resistant *Acinetobacter baumannii* clinical isolates primarily have mutations in

pmrB(53). PmrB is the sensor kinase of the PmrAB two-component system whose regulon

includes three genes involved in LOS modification(12). PmrC and its orphan homologue, EptA,

are transmembrane proteins able to add phosphoethanolamine to both the 1- and 4' phosphate

groups in lipid A (SI Appendix, Fig. S2) (42). In addition, PmrAB positively regulates an operon

including *naxD*, encoding a protein involved in the modification of the 1- phosphate group of

lipid A via the addition of galactosamine (14).

To confirm that resistance in the *pmrB* mutants was due to modifications associated with

upregulation of these genes, qRT-PCR analysis was performed. The transcript levels of *pmrC,*

eptA and *naxD* were all increased in both *pmrB* mutants, with *pmrB*(T235I) resulting in higher

upregulation than *pmrB*(A236E) relative to WT (Fig. 4B). When LOS preparations were

analyzed by TLC, both mutants showed high levels of modified lipid A compared to WT, but the

migration patterns of the modified forms were not identical in the two *pmrB* mutant strains (Fig.

4C). Serendipitously, the previously characterized colistin-resistant ATCC17978 R2 strain has a

pmrB(T235I) mutation that results in lipid A modification as a consequence of both PmrC and

NaxD activity(14, 42). The LAC-4 *pmrB*(T235I) mutant showed a similar lipid A profile to this

strain, providing further evidence that colistin resistance in these mutants results from these two

types of lipid A modification (Fig. 4C).

Mutations in *pmrB* in *A. baumannii* and other Gram-negative bacteria have also been

associated with colistin heteroresistance, which can be confirmed by performing population

analysis profiling (PAP)(54). This assay was performed with WT LAC-4 and the *pmrB* mutants,

which revealed that *pmrB*(A236E) is heteroresistant due to its ability to form colonies at low

frequencies at colistin concentrations 4-8X higher than the resistance breakpoint, even though

the MIC is considerably lower than the breakpoint(18, 54) (Fig. 4D). In contrast, the WT was

unable to grow at 1μg/mL colistin, which was 4-fold below the clinical breakpoint. The

pmrB(T235I) mutant was distinct from both these strains, as it had 100% colony-forming

efficiency up to 64μg/mL colistin (Fig. 4D). PAP of the LOS$^{(-)}$ strains provided further evidence

that the suppressor mutants have higher levels of colistin resistance compared to their *lpxA**

parent (Fig. 4D). Therefore, the *pmrB* mutant selected in the immune-depleted host (T235I) was

colistin-resistant while the one selected in the immunocompetent host (A236E) was colistin

heteroresistant. Notably, in contrast to the high resistance strain, the heteroresistant mutant had a

doubling time that was similar to WT during growth in broth culture (Fig. 2A).

Insertion sequences drive heteroresistance in the *pmrB*(A236E) strain.

Antibiotic heteroresistance is associated with strains that have subpopulations with higher

levels of drug resistance relative to the parental strain(18). The high resistance exhibited in these

subpopulations is often unstable and has been attributed to transient DNA amplification that

allows hyperproduction of proteins that drive increased drug resistance(22). Consistent with this

model, colistin heteroresistance in *Salmonella typhimurium* is driven by amplification of the

pmrD gene, which encodes a positive regulator of *pmrAB*(24), and is consequently unstable via

homologous recombination. *A. baumannii* does not encode *pmrD*, and little is known about what

causes colistin heteroresistance in this organism other than it being linked to activating *pmrAB*

mutations that drive upregulation of lipid A modifying enzymes(17). Therefore, starting with the

low resistance *pmrB*(A236E) strain, we selected for increased resistance followed by continued

growth in the absence of colistin selection to: 1) identify the genetic basis of variants that show

high colistin-resistance in this strain; 2) test the stability of the resistant variants in the absence of

colistin(18).

Three separate lines of the *pmrB*(A236E) strain were grown for 24 hours in LB broth

supplemented with either 2μg/mL (Fig. 5A) or 4μg/mL (Fig. 5B) colistin, which were then

passaged for 7 days in the absence of antibiotics. After drug exposure, colony forming efficiency

(CFE) at 16μg/mL colistin increased to close to 100% frequency from the initial ~5×10^{-4} CFE

before drug exposure, demonstrating enrichment for the colistin-resistant subpopulation. Once

colistin was removed, 5/6 tested lines retained their high CFE at 16μg/mL colistin after 7 days of

passaging (Figs. 5A, B). These results demonstrate that colistin treatment selects for the resistant

subpopulation derived from the *pmrB*(A236E) strain that largely remains stable in the absence of

antibiotic.

To detect genetic changes in the resistant *pmrB*(A236E) population, single colonies

grown at 16μg/mL colistin were collected from all lines after 24 hours colistin treatment (day 0),

and at 4 and 7 days after continuous passage in the absence of drug (Figs. 5A, B). E-test strips

revealed that all 18 isolates had increased their colistin MIC compared to their *pmrB*(A236E)

parent with 10/18 surpassing the breakpoint for clinical resistance (SI Appendix, Table S1, COL

MIC). Short-read whole genome sequencing was performed on a single colony from each

timepoint and confirmed they all retained the *pmrB*(A236E) mutation (SI Appendix, Table S1;

Materials and Methods). Surprisingly, only 5/18 isolates contained additional mutations detected

through short-read sequencing and only 1/18 exhibited evidence of genome amplification, which

was a 9 gene region not known to be associated with antibiotic resistance (Dataset S3 and SI

Appendix, Table S1).

Short-read sequencing analysis often fails to detect large insertions or other

rearrangements in the genome(55). Long read sequencing and de-novo chromosomal assembly

of the 18 isolates revealed that 17/18 isolates had acquired 1-4 insertion sequences (IS) at sites

not found in the *pmrB*(A236E) parent (Dataset S3 and SI Appendix, Table S1). Long-read

sequencing also revealed that the high resistance *pmrB*(T235I) strain contained the additional

insertion mutations (ABLAC_16720::IS*Aba25*, ABLAC_23900::IS*Aba1*, and

ABLAC_35150::IS*Aba13*) absent in the WT ancestor (Dataset S3 and SI Appendix, Fig. S3B).

The *A. baumannii* IS*Aba* insertion sequences are simple mobile genetic elements that can

cause mutations by gene inactivation or altering expression patterns of transcripts downstream

from transposition sites(56). The LAC-4 reference genome contains 6 different types of IS*Aba*

inserted at 81 locations, comprising approximately 2.75% of the chromosome(36). IS*Aba*

insertions into the region upstream of *eptA* and gene-inactivating insertions in *hns* have been

observed in colistin-resistant *A. baumannii* isolates, both resulting in increased *eptA*

transcription(15, 16). Consistent with insertions in these regions causing increased resistance,

16/18 of the colistin-selected *pmrB*(A236E) isolates contained IS*Aba* insertions upstream of *eptA*

or in the *hns* region (Figs. 5C, D; Dataset S3; SI Appendix, Figs. S4C, D and Table S1). qRT-PCR

analysis of *eptA* expression in representative isolates confirmed that IS*Aba* transposition at either

site increased *eptA* expression but not that of *pmrC* or *naxD* (Figs. 5E, F; SI Appendix, Figs. S4C,

D). Interestingly, several of the insertions upstream of *eptA* resulted in significantly lower

expression of *naxD,* consistent with galactosamine modification of lipid A not being a critical

determinant of increased colistin-resistance (SI Appendix, Fig. S4C)(12). TLC profiling of

selected colistin-resistant isolates showed similar lipid A profiles compared to the heteroresistant

parental *pmrB*(A236E) mutant (SI Appendix, Fig. S5). Thus, the stable colistin-resistant

pmrB(A236E) subpopulation is predominantly created through IS*Aba* transposition events

resulting in increased *eptA* expression.

Insertion mutations remain stable in the absence of colistin treatment.

The current models on heteroresistance postulate that the phenotypic stability of high

drug resistance variants depends on the severity of the fitness cost of second-step mutations(18).

The A-0-Hi isolate containing a single IS*Aba*25 insertion upstream of *hns* was chosen for further

studies because it is the only second-step mutation identified in the *pmrB*(A236E) background

that was associated with rapid loss of colistin resistance after subculture in the absence of drug

(Fig. 5D, Dataset S3 and SI Appendix, Fig. S6A). qRT-PCR confirmed that the upstream insertion

decreases *hns* gene expression in the A-0-Hi isolate compared to the *pmrB*(A236E) parent strain

(*SI Appendix*, Fig. S6B). To test the stability of colistin resistance in this mutant, three separate

lines of this A-0-Hi isolate were passaged for 14 days in the absence of antibiotic (Fig. 6A).

Colony forming efficiency at 16μg/mL colistin remained stable for 9 days (~120 generations),

then decreased to *pmrB*(A236E) levels between days 10-14 (Fig. 6A). Therefore, we were unable

to reproduce the instability seen in the *pmrB*(A236E) Col Hi, line A after passaging in antibiotic-

free media (Fig. 5B). However, *pmrB*(A236E) A-0-Hi was a single high-frequency isolate

collected from this line and may not accurately represent the potentially mixed population that

lost resistance.

A-0-Hi isolates from all three lines were collected at day 14 from LB plates

supplemented with 16 or 0 μg/mL colistin to identify the genetic changes that resulted in loss of

resistance (Fig. 6A). Population analysis profiling confirmed that A-0-Hi colonies collected at

day 14 from colistin-free plates, which we will refer to as pseudorevertants, had colistin

sensitivity near WT levels and lower than their heteroresistant *pmrB*(A236E) ancestor (Fig. 6B).

All A-0-Hi isolates collected at day 14 (either pseudorevertants or those remaining resistant)

retained the IS*Aba25* insertion near *hns*, indicating that simple excision of the element was not

the cause of the loss of drug resistance (*SI Appendix*, Fig. S7). Rather, linear amplicon

sequencing of the *pmrCAB* region revealed that all 18 colistin-sensitive pseudorevertants

acquired new mutations disrupting either the *pmrA* or *pmrB* genes (Fig. 6C and *SI Appendix*,

Table S2).

To test whether reversion to sensitivity was due to the fitness cost incurred by high

colistin resistance, growth rates in the absence of colistin were measured for the A-0-Hi strain

and representative pseudorevertants (Fig. 6D and *SI Appendix*, Fig. S8A). Only one

pseudorevertant exhibited a significant decrease in doubling time compared to its A-0-Hi parent,

with none reaching WT growth rates (Fig. 6D). During the continuous passaging experiment,

cultures were grown for hours in post-exponential phase prior to subculturing (Fig. 6A), so we

reasoned that the pseudorevertants may have a competitive advantage after exponential growth.

Therefore, we performed 1:1 competition experiments with all seven pseudorevertants against

their A-0-Hi parent in LB broth and measured the competitive indices at 3 hours (mid-

exponential) and 24 hours of culture (stationary). All pseudorevertants outcompeted their

colistin-resistant parent at 24 hours, while none did so after 3 hours of culture (*SI Appendix*, Figs.

S8B and C). Thus, colistin resistance in the *pmrB*(A236E) A-0-Hi insertion mutant is stable, but

continuous passaging in the absence of drug eventually selects for colistin-sensitive

pseudorevertants, with loss-of-function mutations in *pmrAB* being a common source of the

selective advantage (Fig. 6C).

Discussion:

In the murine pneumonia model, immune depletion results in a permissive environment

for *A. baumannii* replication, facilitating the emergence and selection of tolerant and drug-

resistant mutants that fail to reach high numbers in immunocompetent animals(33). We provide

further evidence that immune depletion accelerates the emergence of antibiotic resistance, with

mutants having altered colistin sensitivity arising more readily in the immune-depleted passages.

Furthermore, we found that bacterial fitness was another important determinant of resistance

evolution during infection, as highly colistin-resistant mutants lacking LOS(49) lost the ability to

colonize both immunocompetent and immune-depleted mouse lungs, even if they acquired

secondary mutations that suppressed their growth defect in broth (Figs. 3C, E). The inability of

LOS$^{(-)}$ mutants to colonize the host, even after they acquire suppressor mutations, is heartening

because LPS/LOS-targeting antibiotics that select for loss of LPS are likely to prevent

proliferation during disease(57). Altogether, these findings highlight the role of the host tissue

environment and the innate immune response in restricting the selection pool of antibiotic-

resistant variants(46).

Regardless of the host state, colistin treatment during pulmonary disease selected for

mutations that increased the expression of enzymes modifying LOS, with *pmrB* transcription-

activating mutants being enriched in our model, as previously observed in broth culture models

and in clinical isolates(58) (Figs. 1C, D). The evolved *pmrB* mutants were outcompeted by their

drug-sensitive parent during lung infections of immunocompetent mice but not in immune-

depleted hosts (Figs. 2B, C). Unlike the LOS$^{(-)}$ mutants, both *pmrB* mutants retained the ability

to grow in host tissues, allowing efficient competition with parental strains during colistin

treatment (Figs. 2D, E). A previous study provided evidence that modified LOS from *A.*

baumannii pmrB mutants induces higher TLR4 activation compared to unmodified LOS(59).

This may be related to reports that some *pmrB* mutants have increased virulence, perhaps as a

consequence of increased pattern recognition-driven inflammation(60, 61). Further work is

needed to determine how LOS modifications caused by *pmrB* mutations impact the pathogenesis

of *A. baumannii*.

We found that strains having mutations located one residue apart in the PmrB histidine

kinase domain resulted in very different phenotypes (Fig. 4). Both mutants accumulated

modified lipid A, with the doubly modified forms found at increased levels in the high resistance

*pmrB(*T235I) strain compared to the heteroresistant *pmrB*(A236E) strain (Fig. 4C). This

correlates well with the higher expression of *pmrC*, *eptA,* and *naxD* resulting from the

pmrB(T235I) mutation (Fig. 4B). The high-resistance *pmrB*(T235I) mutant, however, was found

to have three insertion element mutations that were absent both in the WT parent and the

pmrB(A236E) strain (Dataset S3; COL17 vs. COL23), possibly explaining these resistance

differences. When the *pmrB*(T235I) allele was backcrossed into WT LAC-4, the resulting strain

had a colistin MIC (4µg/mL) that reached the clinical breakpoint for resistance, which is 8X

higher than that of both the backcrossed and evolved *pmrB*(A236E) mutant strains (*SI Appendix*,

Fig. S3A). Population analysis profiling showed that both the evolved and backcrossed

pmrB(T235I) strains were able to form colonies all the way up to 64µg/mL colistin, with the

evolved strain having a higher colony forming efficiency at this concentration as predicted by its

higher E-test colistin MIC. In both backcrossed *pmrB* strains, an additional IS*Aba13* insertion

was detected in identical locations and orientation 61bp upstream ABLAC_20540 (Dataset S3).

Since this mutation does not confer resistance to the heteroresistant *pmrB*(A236E) backcrossed

strain (*SI Appendix*, Figure S3), we can reasonably conclude that the *pmrB*(T235I) allele is

responsible and sufficient for conferring LAC-4 with colistin resistance. Two additional IS*Aba*

insertions mutations were detected in the backcrossed *pmrB*(A236E) strain (Dataset S3);

underscoring the remarkable plasticity of the LAC-4 genome. These results indicate that *pmrB*

mutations located one residue apart can lead to either colistin resistance or heteroresistance.

Although the resistance level of the *pmrB*(T235I) backcrossed strain was above the

clinically defined breakpoint, it is notable that mouse passaging selected for a strain with even

higher levels of resistance, possibly due to the three additional IS insertions (*SI Appendix*, Fig.

S3A, COL17; MIC =16μg/mL). Evolved strains with the *pmrB*(T235I) mutation that harbored

two of these insertion mutations (strains COL29-31) exhibited only a mildly increased MIC

compared to the backcrossed mutant (*SI Appendix*, Fig. S3A) and did not outcompete the

population during mouse passage (Dataset S1; *SI Appendix*, Figs. S1 and S3). The third mutation,

found only in the highest resistance isolate, was in ABLAC_23900, encoding a putative TetR

family regulator (*SI Appendix*, Fig. S3B). This triple insertion strain showed a 3X increase in *hns*

expression compared to WT (*SI Appendix*, Fig. S6C), which could be attributable to any of these

insertions, including one (ABLAC_35150::IS*Aba*13) located 513bp upstream of *hns* (*SI*

Appendix, Fig. S6A). Coupled with our data showing that increased *eptA* expression is associated

with insertions having decreased *hns* expression (Fig. 5E and *SI Appendix*, Fig. S6B), it is

possible that misregulation of *hns* in any direction could increase colistin resistance via *eptA*

overexpression, which is clearly a key step in the progression to colistin resistance.

A confounding factor in the colistin resistance literature is that mutations in *pmrB* have

been associated with colistin susceptibility, resistance and heteroresistance[54]. Indeed, our

heteroresistant *pmrB*(A236E) mutation has been previously associated with colistin

resistance[43]. Based on the IS*Aba* insertions that we identified, the possibility emerges that

colistin-resistant *pmrB*(A236E) strains may have secondary mutations that activate *eptA*

expression. This is because the accumulation of added IS*Aba* insertions in the genome can be

easily missed, even when performing whole-genome sequencing[55]. This raises the possibility

that heteroresistance-conferring mutations like *pmrB*(A236E) present a double danger, because

their "susceptible" classification results in improper antibiotic treatment that is both toxic for the

patient and selects for higher resistance mutants. We do not believe that the selection of these

second-step insertion mutations is a phenomenon of growth in culture, as we were able to

identify similar insertion mutations in *pmrB*(A236E) isolates after continued passage in the

colistin-treated murine pneumonia model (Figs. 2D, E; *SI Appendix*, Fig. S9).

IS*Aba* transposition has been previously implicated in generating colistin-resistant

isolates(19, 51, 59)*.* It has been well-documented in several bacterial species that H-NS

preferentially binds to chromosomal regions that are AT-rich relative to the rest of the

genome(63). In *A. baumannii*, H-NS was shown to direct IS*Aba* transposition to these

regions(62). Notably, the region upstream of *eptA* in LAC-4 is highly AT rich (79%) compared

to the chromosome (61%), possibly explaining why 11/18 resistant isolates derived from

pmrB(A236E) had IS*Aba* insertions in this region (Fig. 5C and Dataset S3) (16). In contrast, the

hns gene itself, where 4/18 *pmrB*(A236E) high resistance isolates had insertions, is only 64%

TA, but the upstream intergenic region is 80.9% TA (Dataset S3).

Most of the isolates derived from the heteroresistant *pmrB*(A236E) strain had IS*Aba*

insertions in additional sites in the chromosome with unknown contributions to resistance

(Dataset S3). One hotspot was in an intergenic region (ABLAC_36890-ABLAC_36900) that

contained IS*Aba25* insertions in 6/18 *pmrB*(A236E)-derived isolates (*SI Appendix*, Fig. S10A).

This region was also TA-rich (79%) and was flanked by genes involved in capsule synthesis

(Dataset S3). Importantly, 5/6 of these isolates contained insertions in the vicinity of *eptA* and

hns with the remaining isolate containing *eptA*(R127H) and *clpS*(Q90*) mutations that could

contribute to colistin resistance (Figs. 5C, D; Dataset S3). The expression of the genes flanking

these insertion mutations was tested via qPCR in two representative *pmrB*(A236E)-derived

isolates, which contained either an *eptA*- or *hns*-disrupting insertion. In both isolates, the

downstream gene ABLAC_36890 (annotated as *weeH*) had a significant (2X) reduction in

expression while the upstream gene ABLAC_36900 only had increased expression when *hns*

was disrupted (*SI Appendix*, Figs. S10B and C). Mutations upstream of *weeH* are associated with

changes in the colony morphology, capsule, biofilm and motility of the mutant(62). The high

frequency of transposition events observed in this study highlights the potential role of insertion

elements in the adaptability of *A. baumannii*.

After ~120 generations of growth in broth, the colistin-resistant *pmrB*(A236) A-0-Hi

strain was outcompeted by colistin-sensitive pseudorevertants (Fig. 6A, B). The majority of these

pseudorevertants acquired loss-of-function mutations in *pmrB,* mainly in the region encoding the

histidine-kinase like ATPase domain (Fig. 6C and *SI Appendix*, Table S2)(64). Other studies

have observed colistin-resistant *A. baumannii* regain sensitivity after cessation of colistin

therapy(11, 65). Previously characterized colistin-sensitive pseudorevertants with mutations

disrupting the *pmrCAB* operon included isolates that may have been evolutionary dead ends,

because they produced colistin-resistant mutants at significantly lower frequencies than their WT

ancestors (65). There are several physiological changes in these revertants that could affect both

fitness in culture and colonization in the host. Deletion of *pmrA* and *pmrB* in *A. baumannii*

reduces motility, the production of biofilm, and the generation of outer membrane vesicles(66).

In mouse models, transposon insertion mutations in *pmrAB* are colonization-proficient,

indicating that loss-of-function of the regulator is compatible with maintenance of virulence (67,

68). It is not clear, however, if PmrAB inactivation provides an advantage during disease only

when such isolates compete against strains that constitutively activate the system.

In summary, we have presented evidence that growth of the heteroresistant *pmrB*(A236E)

parent in colistin selects for insertion mutations that drive acquisition of resistance (Figs. 5C, D;

Dataset S3; *SI Appendix*, Table S1). Similar insertion mutations can be selected after colistin

treatment of lungs infected with the *pmrB*(A236E) strain (Figs. 2D, E; *SI Appendix*, Fig. S9), or

by simply incubating cultures of this heteroresistant mutant on agar plates supplemented with

high colistin concentrations (*SI Appendix*, Fig. S11). Based on the results presented in this study,

we propose the following model. Strains having heteroresistance mutations, such as

pmrB(A236E), create a heterogenous population with varying degrees of lipid A modifications

(Fig. 7A). This enables a subpopulation to survive colistin challenge, potentially creating a stress

condition that activates transposition and results in resistance through insertions in the *eptA* or

hns regions (Fig. 7B). After cessation of antibiotic treatment, colistin-sensitive pseudorevertants

can emerge that acquire mutations disrupting *pmrAB* (Fig. 7C). Although this model predicts that

the *pmrB*(A236E) isolates with resistance-conferring secondary mutations should show an

increase in LOS modification relative to the *pmrB*(A236E) parent, this was not observed using

the TLC strategy described here (*SI Appendix*, Fig. S5). This technique does not quantify the

relative abundance of each type of single/modified lipid A form, such as double-modified

phosphoethanolamine versus singly modified galactosamine plus phosphoethanolamine, which

we suspect may underly increased colistin resistance (Fig. 7B).

This work demonstrates that evolution of bacterial pathogens is a fine interplay between

acquisition of antibiotic resistance and retention of fitness during growth and survival in tissues.

As multidrug-resistant organisms continue to proliferate in healthcare and agricultural settings,

accurately assessing the fitness costs of resistance mechanisms is crucial for identifying potential

Achilles' heels for novel treatments. Our work and others point to phosphoethanolamine

transferases (ex. PmrC/EptA), which can spread through horizontal gene transfer in *mcr*

plasmids(69), as attractive drug targets due to their involvement in colistin resistance and

heteroresistance(70). Furthermore, IS transposition should be considered when studying

antibiotic heteroresistance mechanisms that cannot be explained by gene amplification(22).

Lastly, our results support the model that *A. baumannii* LOS$^{(-)}$ mutants incur devastating fitness

costs that prevent effective colonization during disease, highlighting the crucial role of this

component in maintaining envelope integrity in Gram-negative bacteria(71).

Materials and Methods:

Unabridged Materials and Methods can be found in *SI Appendix*.

Bacterial strains. All strains of *Acinetobacter baumannii* (AB) are derivates of LAC-4

or ATCC17978UN(38, 72). Bacteria were grown in Lysogeny Broth (LB) or LB agar plates at

37^{o}C (*SI Appendix, SI Methods*).

Molecular cloning and mutant construction. The *pmrB*(T235I) and *pmrB*(A236E)

mutations were backcrossed from the COL17 and COL23 strains, respectively, into the parental

WT LAC-4 background (SI Appendix, SI Methods). This was accomplished by electroporation

and plasmid integration followed by homologous recombination, as described previously(73).

Measuring growth rate and competitive index in broth culture. Overnight cultures of

denoted strains were diluted 1:1000 in fresh LB broth in tubes. Cultures were grown at 37^{o}C in a

rotating roller set to 56rpm. OD$_{600}$ was measured at t=0, and at one hour time intervals through

6-8 hrs. The averaged log$_{10}$ values ± SEM over time were plotted. Mean doubling time and SEM

of 2-3 biological replicates was calculated (*SI Appendix, SI Methods*).

For competition experiments in broth, overnight cultures of denoted strains were adjusted

to the same OD, mixed at a 1:1 ratio and diluted into LB broth to a final OD of 0.002. Cultures

were grown at 37^{o}C for 3 and 24 hours. The number of colonies for the competing strains were

quantified at hours 0, 3 and 24. Competition Index (CI) = output(Strain 1/ Strain 2) / input(Strain

1/ Strain 2). Geometric Mean CI and SD was plotted (*SI Appendix, SI Methods*).

Selection of LOS deficient mutant and suppressors. Selection for a LOS deficient

(LOS[(-)]) LAC-4 mutant was performed as described (32, 74) by screening for colistin-resistant

colonies with collateral sensitivity to vancomycin (*SI Appendix, SI Methods*).

LOS[(-)] LAC-4 mutants with suppressor mutations that improved growth rate were

selected by plating the LOS[(-)] strain in LB plates incubated at 42°C for 3 days (*SI Appendix, SI*

Methods).

Animal protocols. All animal procedures were approved by the Institutional Animal

Care and Use Committee (IACUC) of Tufts University. All animals used in this study were 8-10

week-old female BALB/C mice obtained from Jackson Laboratories. All mice were housed in

ventilated caging systems (10–15 air changes per hour) at temperatures of 68–79 °F (~20–26 °C)

and 30–70% humidity, with 12 hr. light and dark cycle.

Bacterial passaging in pulmonary infection model. All passaging procedures were

performed in identical manner in both the immunocompetent and immune-depleted mice, with

inocula adjusted as described in Results(33). Immune depletion was induced by

cyclophosphamide pre-treatment. Lung infections were established by oropharyngeal

aspiration(33, 75).. At 4hrs. post infection, 8mg/kg colistin was directly administered into the

trachea using a liquid PenWu device (BioJane)(76). At 24-hours post infection, mice were

sacrificed using CO_2, and lungs were removed and processed. After outgrowing LAC-4 on LB

plates, the bacteria were collected and frozen stocks were prepared (*SI Appendix, SI Methods*).

For passage 2-16, the frozen bacteria from the previous passage were revived and grown

to mid-exponential phase at 37°C. These cultures were then adjusted to the appropriate OD_{600}

and used to create inocula. The infection, drug administration, and bacterial harvest were

performed in identical fashion at each passage (*SI Appendix, SI Methods*).

The same protocol was used for competitions between purified LAC-4 mutants and WT

in the presence of colistin treatment in either immunocompetent or immune-depleted mice. To

prepare the inocula, the mutants and WT were grown to mid-exponential phase, mixed at an

approximate 5:95 ratio (mutant:WT), and inoculated into mice. Mutant abundance was

calculated by quantifying number of colonies growing in drug plate (selective for *pmrB* mutants)

over number of colonies in no drug plate and then multiplied by 100 (*SI Appendix, SI Methods*).

Mouse competition assays. Mouse competitions were performed as described(33).

Immune depletion was induced with cyclophosphamide pre-treatment and infections were

initiated via oropharyngeal aspiration. At 24-hours post-infection, lungs were removed and

homogenized aseptically in PBS. The number of colonies for the competing strains were

quantified in the inocula and the lung homogenate. Geometric Mean CI and SD was plotted (*SI*

Appendix, SI Methods).

Isolation and quantification of mutants with altered colistin susceptibility. Pools

from each passage of the pulmonary infection model were thawed and CFU were determined on

LB plates in the absence of antibiotics and on graded concentrations of colistin (0 - 4μg/mL)(33).

Colony forming efficiency (CFE) = number of colonies on colistin-containing plate/number of

colonies in LB plate. For each colistin concentration tested, CFE was plotted as a function of

passage number. Colonies growing at colistin concentrations not viable for WT were saved for

further study (*SI Appendix, SI Methods*).

Population analysis profiling (PAP). Overnight cultures of WT LAC-4 and denoted

mutants were diluted into fresh LB and grown to exponential phase. CFE were then determined

on LB plates with increasing concentrations of colistin (0 - 64 μg/mL)(54). CFE were plotted as

a function of colistin concentration for each strain (*SI Appendix, SI Methods*).

613  **Antibiotic MIC determination.** Colistin MIC determinations were performed using the

614 E-test strip method(54). MIC was determined by identifying the site where the zone of inhibition

615 meets the strip (*SI Appendix, SI Methods*).

616  **Resistance stability assay.** The resistance stability assay was performed as

617 described(77). For the *pmrB*(A236E) passaging, three separate colonies were used to establish

618 three lines. Overnight cultures for each line were grown in LB at 37°C. These cultures (input)

619 were then diluted 1:100 in LB supplemented with either 2μg/mL (Lo) or 4μg/mL (Hi) colistin

620 and grown for 24 hours at 37°C. After 24 hrs. drug exposure (Day 0), each culture was diluted

621 1:100 into fresh LB broth without antibiotic and was grown for 24 hours. Cultures were

622 identically passaged in the absence of antibiotic for 7 days. At each day (input, day 0-7), colony

623 forming efficiency at 16μg/mL colistin relative to absence of drug was determined. Isolates able

624 to grow at 16μg/mL colistin were collected from day 0, 4, and 7 for further genome sequencing.

625 For the *pmrB*(A236E) A-0-Hi passaging, three lines were passaged in the antibiotic-free media

626 via 1:10,000 dilutions for 14 days. Isolates from day 14 growing at 0 and 16μg/mL colistin plates

627 were saved (*SI Appendix, SI Methods*).

628  **Quantification of transcription by qPCR**. Overnight cultures of denoted strains were

629 diluted into fresh LB and grown at 37°C until mid-exponential phase. RNA was extracted using

630 the Qiagen RNAeasy kit and subjected to cDNA synthesis using the Invitrogen SuperScript IV

631 VILO kit. The qPCR reactions were performed using the Applied Biosystems PowerUp SYBR

632 Green Master Mix and were run on a StepOnePlus Real-Time PCR system. Transcript level of

633 target genes were normalized to 16S ribosomal RNA and fold change compared to WT was

634 calculated using the ΔΔCt method (*SI Appendix, SI Methods*).

Whole genome sequencing. Genomic DNA (gDNA) was purified from denoted isolates

using the Qiagen DNeasy blood and tissue kit. Illumina Nextera reagents were used to prepare

the libraries for short-read sequencing as described(33). 100bp single-end reads were obtained

from pooled libraries using either a HiSeq2500 or a NovaSeq X Plus at the Tufts University Core

Facility (http://tucf-genomics.tufts.edu/). Mutations were identified by applying the BRESEQ

0.38.8 pipeline to align the reads to the LAC-4 reference genome, including chromosome and

two plasmids(78). (GenBank Accession: chromosome CP007712, pABLAC1 CP007713 and

pABLAC2 CP007714)(36). See Dataset S4 for a comprehensive list of mutations detected in the

evolved isolates.

For long-read sequencing, library preparation and sequencing was performed by the

Hartwell Center for Biotechnology at St. Jude Children's Hospital (SI Appendix). Alternatively,

gDNA was sent to Plasmidsaurus for Nanopore long-read sequencing. Sequencing reads were

de-novo assembled into genomes/contigs using the CANU pipeline(79). Mauve whole genome

alignments on Geneious Prime were performed to compare the chromosomes of the mutants

versus their parent. Insertions and deletions present in the mutants but not the parent were then

identified (*SI Appendix, SI Methods*). All sequencing information provided in this text has been

deposited in Genbank under the accession number PRJNA1273616.

LOS and lipid A visualization. For fractionation and staining of whole cell LOS, 10^9

CFUs of denoted strains were collected, washed and resuspended in 1X LDS sample buffer + 5%

β-mercaptoethanol, and boiled for 10 minutes. Samples were then treated with proteinase K and

run on a 4-12% Bis-Tris SDS-PAGE gel. Staining of LOS was performed using the ProQ

Emerald 300 kit, following manufacturer instructions (80).

Lipid A isolation for TLC analysis from denoted AB strain was performed as

described(80). Overnight cultures of denoted strains were diluted into fresh LB with 5

µCi/mL ^{32}P ortho-phosphoric acid (Perkin-Elmer) and grown to an $OD_{600} = 1$. Lipid A was

extracted using mild-acid hydrolysis followed by a Blight-Dryer extraction. Lipid A species were

separated by thin layer chromatography (TLC) in a chloroform/pyridine/88% formic acid/water

(50:50:16:5, vol/vol) solvent system. An Amersham Typhoon laser scanner with a

phosphorimaging screen was used to image the dry plates.

Statistical Analysis. GraphPad Prism was used for statistical analysis. For all statistical

analysis, One Way ANOVA followed by Dunnet's multiple comparison or unpaired two-tailed t-

tests were employed. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$; ns, not significant.

Figure Design. Illustrations in Figures 1A, 3B and 7 were created in BioRender.com.

Acknowledgements:

This work is supported by NIAID U01AI124302, U19AI158076, U19AI142780 and

R21AI128328. We would like to thank Albert Tai and Irina V. Grinvald for help with library

preparations and the next-generation sequencing support provided by S10OD032203 via Tufts

University Core Facility Genomics Core. We would like to thank the St. Jude Children's

Hospital Hartwell Center for Biotechnology for help with sequencing library preparation and

Pacbio sequencing. We would like to thank Eddie Geisinger, Tim Van Opijnen and all the

members of the Isberg lab for their feedback on the study.

Author Contributions:

J.H-B. and R.R.I devised the study. J.H-B., B.H., L.M.V., E.B.B. and G. I. C. T. performed the

wet lab experiments. H.E. and J.M.R. performed long-red sequencing and provided consultation

on library preparations. J.H-B and W.H. performed the sequencing analysis. L.M.V and M.S.T.

performed the LOS analysis and provided analysis of results.

Figure Legends:

Figure 1: Colistin treatment selects for *pmrB* mutations in *A. baumannii* during serial

oropharyngeal inoculations.

A. Passaging strategy. Immune depletion was induced by intraperitoneal (IP) cyclophosphamide

injections at 72- and 24- hours before infection. Three separate, co-housed lines derived from

different LAC-4 colonies were passaged per condition. *A. baumannii* LAC-4 was

oropharyngeally (OP) inoculated into mice and 4 hours post-infection (hpi), 8mg/kg aerosolized

colistin was administered via OP inhalation. At 24 hpi, the mice were euthanized, lungs were

removed, homogenized, and plated on LB agar plates. After overnight growth, the bacteria were

collected and saved for the next round of infections. **B-C.** Lung CFU after each passage in

colistin-treated mice. Dotted line: expected bacterial burden in the absence of colistin treatment

(Materials and Methods). At passage 2 in immune-depleted host, the inoculum size of $5X10^7$

CFU was reduced to $1X10^7$ CFU, followed by a reduction to $5X10^6$ CFU at passage 5 (arrows).

Inoculum size in immunocompetent host: $1X10^8$ CFU. **D-E.** Identification of colistin-resistant

mutants. Pools were serially diluted and plated on LB plates supplemented with 0 or 4 μg/mL

colistin. Colony forming efficiency (CFE) was calculated by the ratio of colonies growing in the

presence or absence of drug. Dotted line: limit of detection. Isolates that grew in colistin plates

were whole-genome sequenced to reveal the presence of the noted *pmrB* mutations.

Figure 2: Mutations in *pmrB* selected during pulmonary infection incur a fitness cost in the absence of colistin.

A. Kinetics of *pmrB* mutant growth in broth. Denoted LAC-4 strains were incubated in LB broth, and mass increase was measured at one-hour intervals. Mean doubling time $\pm$ SEM of three biological replicates shown. **B-C.** Competition of *pmrB* mutants vs. WT during pulmonary infections. Denoted strains were mixed at approximately 1:1 and inoculated via oropharyngeal aspiration into animals treated under noted conditions. 24hpi, mice were euthanized, and lungs were removed, homogenized in cold PBS, and plated on LB plates with or without colistin. Competition Index (CI) determined as described (Materials and Methods). Geometric Mean CI $\pm$ SD of 9-16 biological replicates was determined. **D-E.** The *pmrB*(T235I) strain efficiently competes with WT LAC-4 during pulmonary infections in the presence of colistin. WT LAC-4 and noted mutants were mixed at approximately 95:5 ratio followed by oropharyngeal inoculation into mice. 8mg/kg of colistin was administered 4hpi. At 24 hpi, the mice were euthanized, lungs were removed, homogenized, and plated on LB agar plates (Materials and Methods). Frozen pools from each passage were serially diluted and spot plated in LB plates supplemented with 0, 2 or 8μg/mL colistin. Mutant abundance was calculated as described (Materials and Methods). **A-C.** Statistical analysis of doubling time and CIs was performed using One-way ANOVA, followed by Dunnett's multiple comparisons (Materials and Methods). *$P < 0.05$, ****$P < 0.0001$; ns, not significant.

Figure 3: LOS-deficient *A. baumannii* isolates are defective for lung colonization.

A. LOS[(-)] strains lack carbohydrate sidechain. LOS isolated from denoted strains was fractionated on SDS-PAGE and carbohydrate was stained with ProQ Emerald 300. The denoted *E. coli* strain does not produce O-antigen and should migrate closely to *A. baumannii* LOS. The

bottom band present in all samples is an unknown glycolipid or intermediate species. **B.** Protocol

for selection of LOS-deficient LAC-4 and higher fitness variants. WT LAC-4 was plated on LB

plates supplemented with 10μg/mL colistin and colonies were restreaked onto identical medium.

An isolate ($lpxA^*$) that grew at 10μg/mL colistin but not 10μg/mL vancomycin was saved and

sequenced. 10^9 CFU of the isolate were plated on LB at 42°C and papillae growing above the

lawn after 72 hrs. were streaked onto LB. Candidates showing improved growth at 42°C relative

to parental LOS$^{(-)}$ strain were saved and sequenced (Dataset S2). Three non-identical isolates

($lpxA^*$ $pldA^*$, $lpxA^*$ $pldA^*$ $pbp1A^{ins}$, and $lpxA^*$ $pldA^*$ $pbp1A^*$) were selected for further study. **C.**

Kinetics of LOS$^{(-)}$ mutant growth in broth. Denoted LAC-4 strains were incubated in LB broth,

and mass increase was measured at one-hour intervals. Mean doubling time $\pm$ SEM of replicates

shown. **D-E.** Competition of LOS$^{(-)}$ mutants vs. WT during pulmonary infection. Denoted strains

were mixed approximately 1:1 and inoculated via oropharyngeal aspiration into mice. 24hpi,

mice were euthanized, lungs were removed, homogenized in cold PBS, and incubated on LB

solid medium in presence or absence of colistin. Geometric Mean CI $\pm$ SD of 8-9 replicates

determined as described (Materials and Methods). **C-E.** Statistical analysis of doubling time and

CI's were performed using one-way Anova followed by Dunnett's multiple comparison.

**$P < 0.01$, ***$P < 0.001$; ns, not significant.

Figure 4: Overexpression of *pmrB*-regulated genes leads to colistin resistance and

heteroresistance.

A. Susceptibility of *pmrB* and LOS$^-$ mutants to colistin. Colistin MIC determination via E-test

strip. MIC determined as described (Materials and Methods). **B.** Expression of *pmrAB*-regulated

genes in colistin-selected *pmrB* mutants. Transcript levels of *pmrC, eptA and naxD* was

quantified via RT-qPCR and displayed as expression relative to WT. Mean $\pm$ SEM from three

biological replicates shown. **C.** Lipid A from denoted strains was isolated, radiolabeled with ^{32}P

and separated by thin layer chromatography (TLC). Single and double modified forms refers to

the addition of phosphoethanolamine and/or galactosamine. **D.** Population analysis profiling of

pmrB and LOS$^-$ mutants**.** Cultures of denoted strains were serially diluted and plated in LB

supplemented with noted concentrations of colistin. Mean colony forming efficiency (CFE) was

calculated as described (Materials and Methods). Grey vertical line: clinical breakpoint for

resistance ($\geq$4µg/mL). **B.** Statistical analysis of gene expression changes were performed using

unpaired, two-tailed t-tests. ****P* $<$ 0.001.

Figure 5: Enrichment of *pmrB*(A236E) isolates with higher colistin resistance selects for

IS*Aba* activation of *eptA*.

A-B. Enrichment and stability of high colistin resistance strains derived from the *pmrB*(A236E)

background. Three lines (A, B, C) of the *pmrB*(A236E) mutant were selected for high resistance

by 24 hr. growth in either 2µg/mL (Lo) or 4µg/mL (Hi) colistin (Day 0/+COL). Each line was

subsequently passaged daily in antibiotic-free media for 7 days. Every 24 hrs., CFU were

quantified and colony forming efficiency (CFE) on LB containing 16µg/mL colistin plates was

determined, as described (Materials and Methods). Colonies growing at 16µg/mL colistin were

isolated at day 0, 7, 14 for further analysis. **C-D.** Location and orientation of IS*Aba* insertions in

colistin-enriched *pmrB*(A236E) isolates. Figures not drawn to scale. **E-F.** Expression of *eptA* in

the colistin-enriched *pmrB*(A236E) isolates. Representative isolates (in bold) were selected for

further analysis. Transcript levels of *eptA* in mutants relative to WT was quantified via RT-

qPCR. Mean $\pm$ SEM from three biological replicates shown. **C-F.** Isolate nomenclature refers to

"Line – Day collected – Level of colistin exposure at day 0". **E-F.** Statistical analysis of gene

expression changes were performed using one-way Anova followed by Dunnett's multiple

comparison. $*P < 0.05$, $**P < 0.01$, $****P < 0.0001$.

Figure 6: Colistin susceptible *pmrB*(A236E) A-0-Hi pseudorevertants emerge after

continued passaging in the absence of drug.

A. Stability of colistin resistance in the absence of antibiotic selection. Three lines of the colistin-

resistant *pmrB*(A236E) A-0-Hi isolate were passaged in antibiotic-free media for 14 days. Every

24 hrs, colony forming units on LB plates containing 0 or 16μg/mL colistin was measured. CFE

was calculated as described (Materials and Methods). **B-D.** 18 *pmrB*(A236E) A-0-Hi

pseudorevertants (PR) were collected for further analysis by picking colonies growing on LB

plates after 14 days of antibiotic-free passaging. **B.** Population analysis profiling of

pmrB(A236E) A-0-Hi pseudorevertants. Cultures of denoted strains were serially diluted and

plated on LB agar supplemented with noted concentrations of colistin. CFE was calculated as

described (Materials and Methods). **C.** Schematic indicating identity and location of mutations in

representative *pmrB*(A236E) A-0-Hi pseudorevertants detected by linear amplicon sequencing of

pmrCAB region. **D.** Growth rates of *pmrB*(A236E) A-0-Hi pseudorevertants in broth. Mean

doubling time + SEM of three biological replicates shown. Statistical analysis of doubling time

was performed using One-way Anova followed by Dunnett's multiple comparison. $*P < 0.05$,

$****P < 0.0001$; ns, not significant. Kinetics shown in *SI Appendix,* Fig. S8A.

Figure 7. Model for the two-step acquisition of colistin resistance.

A. Activation of the PmrB histidine kinase (via signal or mutation) leads to phosphorylation of

PmrA. PmrA transcription factor binds to PmrA binding site in the *pmrCAB*, *eptA* and *naxD*

promoter regions. PmrC and EptA add phosphoethanolamine (PEtN) to the 1 and 4' phosphate

groups on lipid A. NaxD deacetylates *N*-acetylgalactosamine (GalNAc) to galactosamine (GalN),

which is then added to the 1- phosphate on lipid A by an unknown enzyme. These modifications

reduce the negative charge of the lipid A portion of lipooligosaccharide which reduce its affinity

to colistin. Mutations that strongly activate this system [ex. *pmrB*(T235I)] lead to resistance

while mutations that weakly activate this system [ex. *pmrB*(A236E)] lead to heteroresistance. **B.**

Heteroresistant *pmrB* mutant becomes fully colistin-resistant through insertion sequence

transposition. The *eptA* gene is poorly activated by the *pmrB*(A236E) mutation, as the upstream

region is TA rich and is silenced by H-NS (Sensitive). IS*Aba* transposition to the region

upstream of *eptA* increases the expression of this gene leading to resistance (Path 1). IS*Aba*

transposition into *hns* derepresses TA rich regions of the genome, including *eptA*, and leads to

resistance (Path 2). **C.** Continued passaging in the absence of drug selects for colistin sensitive

pseudorevertants that acquire loss of function mutations in *pmrAB*.

References:

1.   Centers for Disease Control and Prevention (U.S.), "Antibiotic resistance threats in the
United States, 2019" (Centers for Disease Control and Prevention (U.S.), 2019).

2.   A. K. Cain, M. Hamidian, Portrait of a killer: Uncovering resistance mechanisms and global
spread of Acinetobacter baumannii. *PLOS Pathogens* **19**, e1011520 (2023).

3.   D. Wong, *et al.*, Clinical and Pathophysiological Overview of Acinetobacter Infections: a
Century of Challenges. *Clinical microbiology reviews* **30**, 409–447 (2017).

4.   C. Ayoub Moubareck, D. Hammoudi Halat, Insights into Acinetobacter baumannii: A
Review of Microbiological, Virulence, and Resistance Traits in a Threatening Nosocomial
Pathogen. *Antibiotics* **9**, 119 (2020).

5.   WHO bacterial priority pathogens list, 2024: Bacterial pathogens of public health
importance to guide research, development and strategies to prevent and control

antimicrobial resistance. Available at:
https://www.who.int/publications/i/item/9789240093461.

6.   B. Isler, Y. Doi, R. A. Bonomo, D. L. Paterson, New Treatment Options against
Carbapenem-Resistant Acinetobacter baumannii Infections. *Antimicrobial Agents and*
Chemotherapy **63**, 10.1128/aac.01110-18 (2018).

7.   D. Landman, C. Georgescu, D. A. Martin, J. Quale, Polymyxins Revisited. *Clinical*
Microbiology Reviews **21**, 449–465 (2008).

8.   A. L. H. Kwa, C. Loh, J. G. H. Low, A. Kurup, V. H. Tam, Nebulized Colistin in the
Treatment of Pneumonia Due to Multidrug-Resistant Acinetobacter baumannii and
Pseudomonas aeruginosa. *Clinical Infectious Diseases* **41**, 754–757 (2005).

9.   Z. Zhang, *et al.*, The efficacy and safety of inhaled antibiotics for pneumonia: A systematic
review and meta-analysis. *Pulmonary Pharmacology & Therapeutics* **86**, 102315 (2024).

10. J. H. Moffatt, M. Harper, J. D. Boyce, "Mechanisms of Polymyxin Resistance" in
Polymyxin Antibiotics: From Laboratory Bench to Bedside, J. Li, R. L. Nation, K. S. Kaye,
Eds. (Springer International Publishing, 2019), pp. 55–71.

11. M. D. Adams, *et al.*, Resistance to Colistin in Acinetobacter baumannii Associated with
Mutations in the PmrAB Two-Component System. *Antimicrobial Agents and*
Chemotherapy **53**, 3628–3634 (2009).

12. N. Yamada, *et al.*, PmrAB, the two-component system of Acinetobacter baumannii,
controls the phosphoethanolamine modification of lipooligosaccharide in response to metal
ions. *Journal of Bacteriology* **206**, e00435-23 (2024).

13. L. A. Arroyo, *et al.*, The pmrCAB operon mediates polymyxin resistance in Acinetobacter
baumannii ATCC 17978 and clinical isolates through phosphoethanolamine modification of
lipid A. *Antimicrobial agents and chemotherapy* **55**, 3743–51 (2011).

14. C.-Y. Chin, K. A. Gregg, B. A. Napier, R. K. Ernst, D. S. Weiss, A PmrB-Regulated
Deacetylase Required for Lipid A Modification and Polymyxin Resistance in Acinetobacter
baumannii. *Antimicrobial agents and chemotherapy* **59**, 7911–4 (2015).

15. E. J. Yoon, *et al.*, Trajectory of genetic alterations associated with colistin resistance in
Acinetobacter baumannii during an in-hospital outbreak of infection. *Journal of*
Antimicrobial Chemotherapy **77**, 69–73 (2021).

16. D. D. Lucas, *et al.*, Emergence of High-Level Colistin Resistance in an Acinetobacter
baumannii Clinical Isolate Mediated by Inactivation of the Global Regulator H-NS. (2018).
https://doi.org/10.1128/AAC.02442-17.

17. Y. Charretier, *et al.*, Colistin Heteroresistance and Involvement of the PmrAB Regulatory
System in Acinetobacter baumannii. *Antimicrobial Agents and Chemotherapy* **62**,
10.1128/aac.00788-18 (2018).

18.   D. I. Andersson, H. Nicoloff, K. Hjort, Mechanisms and clinical relevance of bacterial
853   heteroresistance. *Nature Reviews Microbiology 2019 17:8* **17**, 479–496 (2019).

854   19.   V. I. Band, D. S. Weiss, Heteroresistance: A cause of unexplained antibiotic treatment
855   failure? *PLOS Pathogens* **15**, e1007726 (2019).

856   20.   V. I. Band, D. S. Weiss, Heteroresistance to beta-lactam antibiotics may often be a stage in
857   the progression to antibiotic resistance. *PLoS Biol* **19**, e3001346 (2021).

858   21.   A. Pal, D. I. Andersson, Bacteria can compensate the fitness costs of amplified resistance
859   genes via a bypass mechanism. *Nat Commun* **15**, 2333 (2024).

860   22.   H. Nicoloff, K. Hjort, B. R. Levin, D. I. Andersson, The high prevalence of antibiotic
861   heteroresistance in pathogenic bacteria is mainly caused by gene amplification. *Nature
862   Microbiology 2019 4:3* **4**, 504–514 (2019).

863   23.   K. N. Kang, *et al.*, Colistin heteroresistance in Enterobacter cloacae is regulated by PhoPQ-
864   dependent 4-amino-4-deoxy-l-arabinose addition to lipid A. *Molecular microbiology* **111**,
865   1604 (2019).

866   24.   K. Hjort, H. Nicoloff, D. I. Andersson, Unstable tandem gene amplification generates
867   heteroresistance (variation in resistance within a population) to colistin in Salmonella
868   enterica. *Molecular Microbiology* **102**, 274–289 (2016).

869   25.   L. Bardet, *et al.*, Deciphering Heteroresistance to Colistin in a Klebsiella pneumoniae
870   Isolate from Marseille, France. *Antimicrobial Agents and Chemotherapy* **61**,
871   10.1128/aac.00356-17 (2017).

872   26.   S. P. Doijad, *et al.*, Resolving colistin resistance and heteroresistance in Enterobacter
873   species. *Nat Commun* **14**, 140 (2023).

874   27.   M. J. Powers, M. S. Trent, Phospholipid retention in the absence of asymmetry strengthens
875   the outer membrane permeability barrier to last-resort antibiotics. *Proceedings of the
876   National Academy of Sciences of the United States of America* **115**, E8518–E8527 (2018).

877   28.   M. J. Powers, M. S. Trent, Expanding the paradigm for the outer membrane: Acinetobacter
878   baumannii in the absence of endotoxin. *Molecular Microbiology* **107**, 47–56 (2018).

879   29.   G. J. D. Silva, S. Domingues, Interplay between Colistin Resistance, Virulence and Fitness
880   in Acinetobacter baumannii. *Antibiotics* **6** (2017).

881   30.   J. Y. Lee, E. S. Chung, K. S. Ko, Transition of colistin dependence into colistin resistance
882   in Acinetobacter baumannii. *Scientific Reports 2017 7:1* **7**, 1–11 (2017).

883   31.   J. S. Hawley, C. K. Murray, J. H. Jorgensen, Development of colistin-dependent
884   Acinetobacter baumannii-Acinetobacter calcoaceticus complex [1]. *Antimicrobial Agents
885   and Chemotherapy* **51**, 4529–4530 (2007).

32. J. M. Boll, *et al.*, A penicillin-binding protein inhibits selection of colistin-resistant,
lipooligosaccharide-deficient Acinetobacter baumannii. *Proceedings of the National*
Academy of Sciences of the United States of America **113**, E6228–E6237 (2016).

33. W. Huo, *et al.*, Immunosuppression broadens evolutionary pathways to drug resistance and
treatment failure during Acinetobacter baumannii pneumonia in mice. *Nat Microbiol* **7**,
796–809 (2022).

34. K. W. Bruhn, *et al.*, Host fate is rapidly determined by innate effector-microbial
interactions during Acinetobacter baumannii bacteremia. *The Journal of infectious diseases*
211, 1296–305 (2015).

35. A. Maure, E. Robino, C. Van der Henst, The intracellular life of *Acinetobacter baumannii*.
Trends in Microbiology **31**, 1238–1250 (2023).

36. H.-Y. Ou, *et al.*, Complete genome sequence of hypervirulent and outbreak-associated
Acinetobacter baumannii strain LAC-4: epidemiology, resistance genetic determinants and
potential virulence factors. *Scientific Reports* **5** (2015).

37. S. C. Valentine, *et al.*, Phenotypic and Molecular Characterization of Acinetobacter
baumannii Clinical Isolates from Nosocomial Outbreaks in Los Angeles County, California.
J Clin Microbiol **46**, 2499–2507 (2008).

38. G. Harris, *et al.*, A Mouse model of acinetobacter baumannii-associated pneumonia using a
clinically isolated hypervirulent strain. *Antimicrobial Agents and Chemotherapy* **57**, 3601–
3613 (2013).

39. R. J. North, Cyclophosphamide-facilitated adoptive immunotherapy of an established tumor
depends on elimination of tumor-induced suppressor T cells. *Journal of Experimental*
Medicine **155**, 1063–1074 (1982).

40. A. F. Zuluaga, *et al.*, Neutropenia induced in outbred mice by a simplified low-dose
cyclophosphamide regimen: characterization and applicability to diverse experimental
models of infectious diseases. *BMC Infect Dis* **6**, 55 (2006).

41. MR01Ed2E | Polymyxin Breakpoints for Enterobacterales, Pseudomonas aeruginosa, and
Acinetobacter spp., 2nd Edition. *Clinical & Laboratory Standards Institute*. Available at:
https://clsi.org/standards/products/microbiology/companion/mr01/.

42. L. A. Arroyo, *et al.*, The pmrCAB operon mediates polymyxin resistance in Acinetobacter
baumannii ATCC 17978 and clinical isolates through phosphoethanolamine modification of
lipid A. *Antimicrobial Agents and Chemotherapy* **55**, 3743–3751 (2011).

43. R. López-Rojas, M. García-Quintanilla, G. Labrador-Herrera, J. Pachón, M. J. McConnell,
Impaired growth under iron-limiting conditions associated with the acquisition of colistin
resistance in *Acinetobacter baumannii*. *International Journal of Antimicrobial Agents* **47**,
473–477 (2016).

44. T. Vogwill, R. C. MacLean, The genetic basis of the fitness costs of antimicrobial
resistance: a meta-analysis approach. *Evolutionary applications* **8**, 284–95 (2015).

45. A. H. Melnyk, A. Wong, R. Kassen, The fitness costs of antibiotic resistance mutations.
Evolutionary Applications **8**, 273 (2015).

46. M. J. Culyba, D. V. Tyne, Bacterial evolution during human infection: Adapt and live or
adapt and die. *PLOS Pathogens* **17**, e1009872 (2021).

47. B. W. Simpson, *et al.*, Acinetobacter baumannii Can Survive with an Outer Membrane
Lacking Lipooligosaccharide Due to Structural Support from Elongasome Peptidoglycan
Synthesis. *mBio* (2021). https://doi.org/10.1128/MBIO.03099-21.

48. J. W. Skudlarek, *et al.*, Cerastecin Inhibition of the Lipooligosaccharide Transporter MsbA
to Combat Acinetobacter baumannii: From Screening Impurity to In Vivo Efficacy. *J. Med.*
Chem. **67**, 15620–15675 (2024).

49. K. Novović, B. Jovčić, Colistin Resistance in Acinetobacter baumannii: Molecular
Mechanisms and Epidemiology. *Antibiotics* **12**, 516 (2023).

50. R. López-Rojas, *et al.*, Impaired virulence and in vivo fitness of colistin-resistant
Acinetobacter baumannii. *The Journal of infectious diseases* **203**, 545–8 (2011).

51. M. E. Wand, L. J. Bock, L. C. Bonney, J. M. Sutton, Retention of virulence following
adaptation to colistin in Acinetobacter baumannii reflects the mechanism of resistance.
Journal of Antimicrobial Chemotherapy **70**, 2209–2216 (2015).

52. A. Beceiro, *et al.*, Biological cost of different mechanisms of colistin resistance and their
impact on virulence in Acinetobacter baumannii. *Antimicrobial agents and chemotherapy*
58, 518–26 (2014).

53. S. Gerson, *et al.*, Diversity of amino acid substitutions in PmrCAB associated with colistin
resistance in clinical isolates of *Acinetobacter baumannii*. *International Journal of*
Antimicrobial Agents **55**, 105862 (2020).

54. E. X. Sherman, J. E. Wozniak, D. S. Weiss, Methods to Evaluate Colistin Heteroresistance
in Acinetobacter baumannii. *Methods in molecular biology (Clifton, N.J.)* **1946**, 39 (2019).

55. N. D. Maio, *et al.*, Comparison of long-read sequencing technologies in the hybrid
assembly of complex bacterial genomes. *Microbial Genomics* **5**, e000294 (2019).

56. H. R. Noel, J. R. Petrey, L. D. Palmer, Mobile genetic elements in Acinetobacter antibiotic-
resistance acquisition and dissemination. *Annals of the New York Academy of Sciences*
1518, 166–182 (2022).

57. C. Zampaloni, *et al.*, A novel antibiotic class targeting the lipopolysaccharide transporter.
Nature **625**, 566–571 (2024).

58.   K. Novović, B. Jovčić, Colistin Resistance in Acinetobacter baumannii: Molecular
957   Mechanisms and Epidemiology. *Antibiotics (Basel)* **12**, 516 (2023).

958   59.   L. Lin, *et al.*, Inhibition of LpxC protects mice from resistant Acinetobacter baumannii by
959   modulating inflammation and enhancing phagocytosis. *mBio* **3** (2012).

960   60.   G. E, H. W, H.-B. J, I. RR, Acinetobacter baumannii: Envelope Determinants That Control
961   Drug Resistance, Virulence, and Surface Variability. *Annual review of microbiology* **73**,
962   481–506 (2019).

963   61.   S. Gerson, *et al.*, Investigation of Novel pmrB and eptA Mutations in Isogenic
964   Acinetobacter baumannii Isolates Associated with Colistin Resistance and Increased
965   Virulence In Vivo. *Antimicrobial Agents and Chemotherapy* **63**, 10.1128/aac.01586-18
966   (2019).

967   62.   C. Cooper, *et al.*, H-NS is a bacterial transposon capture protein. *Nat Commun* **15**, 7137
968   (2024).

969   63.   C. J. Dorman, H-NS, the genome sentinel. *Nat Rev Microbiol* **5**, 157–161 (2007).

970   64.   A. O. Olaitan, S. Morand, J.-M. Rolain, Mechanisms of polymyxin resistance: acquired and
971   intrinsic resistance in bacteria. *Frontiers in microbiology* **5**, 643 (2014).

972   65.   E. S. Snitkin, *et al.*, Genomic insights into the fate of colistin resistance and Acinetobacter
973   baumannii during patient treatment. *Genome Res.* **23**, 1155–1162 (2013).

974   66.   S. Y. Ko, *et al.*, PmrAB controls virulence-associated traits and outer membrane vesicle
975   biogenesis in *Acinetobacter baumannii*. *Microbial Pathogenesis* **185**, 106434 (2023).

976   67.   S. Subashchandrabose, *et al.*, Acinetobacter baumannii Genes Required for Bacterial
977   Survival during Bloodstream Infection. *mSphere* **1**, e00013-15 (2015).

978   68.   N. Wang, E. A. Ozer, M. J. Mandel, A. R. Hauser, Genome-wide identification of
979   Acinetobacter baumannii genes necessary for persistence in the lung. *mBio* **5**, e01163-14
980   (2014).

981   69.   P. K. Jangir, L. Ogunlana, R. C. MacLean, Evolutionary constraints on the acquisition of
982   antimicrobial peptide resistance in bacterial pathogens. *Trends in Microbiology* **29**, 1058–
983   1061 (2021).

984   70.   V. C. Thai, K. A. Stubbs, M. Sarkar-Tyson, C. M. Kahler, Phosphoethanolamine
985   Transferases as Drug Discovery Targets for Therapeutic Treatment of Multi-Drug Resistant
986   Pathogenic Gram-Negative Bacteria. *Antibiotics (Basel)* **12**, 1382 (2023).

987   71.   J. C. Henderson, *et al.*, The Power of Asymmetry: Architecture and Assembly of the Gram-
988   Negative Outer Membrane Lipid Bilayer. *Annual Review of Microbiology* **70**, 255–278
989   (2016).

72. C. D. M. Wijers, *et al.*, Identification of Two Variants of Acinetobacter baumannii Strain
ATCC 17978 with Distinct Genotypes and Phenotypes. *Infection and Immunity* **89** (2021).

73. E. Geisinger, R. R. Isberg, Antibiotic Modulation of Capsular Exopolysaccharide and
Virulence in Acinetobacter baumannii. *PLOS Pathogens* **11**, e1004691 (2015).

74. J. H. Moffatt, *et al.*, Colistin Resistance in Acinetobacter baumannii Is Mediated by
Complete Loss of Lipopolysaccharide Production. *Antimicrobial Agents and Chemotherapy*
54, 4971–4977 (2010).

75. T. B. Nielsen, J. Yan, B. Luna, B. Spellberg, Murine Oropharyngeal Aspiration Model of
Ventilator-associated and Hospital-acquired Bacterial Pneumonia. *J Vis Exp* 57672 (2018).
https://doi.org/10.3791/57672.

76. Y.-W. Lin, *et al.*, Elucidating the Pharmacokinetics/Pharmacodynamics of Aerosolized
Colistin against Multidrug-Resistant Acinetobacter baumannii and Klebsiella pneumoniae
in a Mouse Lung Infection Model. *Antimicrobial agents and chemotherapy* **62** (2018).

77. J. E. Choby, *et al.*, Copy number flexibility facilitates heteroresistance to increasing
antibiotic pressure and threatens the beta-lactam pipeline. *Nat Commun* **16**, 5721 (2025).

78. D. E. Deatherage, J. E. Barrick, "Identification of Mutations in Laboratory-Evolved
Microbes from Next-Generation Sequencing Data Using breseq" in *Engineering and*
Analyzing Multicellular Systems: Methods and Protocols, L. Sun, W. Shou, Eds. (Springer,
2014), pp. 165–188.

79. S. Koren, *et al.*, Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

80. M. J. Powers, C. M. Herrera, A. T. Tucker, B. W. Davies, M. S. Trent, "Isolation of Lipid
Cell Envelope Components from Acinetobacter baumannii" in *Acinetobacter Baumannii:*
Methods and Protocols, I. Biswas, P. N. Rather, Eds. (Springer, 2019), pp. 233–252.

1014

A

Immune-depleted: lines A, B, C

-72 hr   IP: Cyclophosphamide

-24 hr   IP: Cyclophosphamide

0 hr   OP: AB inoculum

4hpi   OP: Colistin

24hpi   Homogenize

Overnight   Count and freeze

16 passages

Immunocompetent: lines A, B, C

OP: AB inoculum

OP: Colistin

Homogenize

Count and freeze

16 passages

B   Immune-depleted

Lung bacterial load (CFU)

Expected load without colistin

Passage

C   Immunocompetent

Lung bacterial load (CFU)

Expected load without colistin

Passage

Line A
Line B
Line C

D   Immune-depleted

Colony forming efficiency at 4µg/mL colistin

pmrB(T235I)

Passage

E   Immunocompetent

Colony forming efficiency at 4µg/mL colistin

pmrB(A236E)

Limit of detection

Passage

Line A
Line B
Line C