Disrupting Akt-Wnt/β-catenin signaling suppresses glioblastoma stem cell growth and tumor progression in immunocompetent mice

Md. Marzan Sarkar[1,2,5], Nathan Gonsalves[1,2,5], Amin Davarzani[1,3], Emma Mitchell[1,4], Amar M. Singh[1,2,5], Lohitash Karumbaiah[1,2,5*], and Steven L. Stice[1,2,5*]

[1]Regenerative Bioscience Center, Edgar L. Rhodes Center for Animal and Dairy Science, University of Georgia, Athens, GA 30602

[2]College of Agricultural and Environmental Sciences, University of Georgia, Athens, GA 30602

[3]Department of Electrical and Computer Engineering, College of Engineering, University of Georgia, Athens, GA 30602

[4]Department of Biomedical Physiology, College of Veterinary Medicine, University of Georgia, Athens, GA 30602

[5]Department of Animal and Dairy Science, University of Georgia, Athens, GA 30602

*Corresponding Authors: Steven L. Stice and Lohitash Karumbaiah

E-mail address: sstice@uga.edu (S.L. Stice). lohitash@uga.edu (L. Karumbaiah)

Abstract

Glioblastoma (GBM) is an aggressive primary malignant brain tumor in adults with limited therapeutic options and a median survival of 12–18 months post-diagnosis. The PI3K/Akt and Wnt/β-catenin signaling pathways promote GBM cell growth and survival, invasiveness and therapeutic resistance. We hypothesize that inhibiting Akt and β-catenin, which are key central regulatory components of the PI3K/Akt and Wnt/β-catenin pathway, will suppress GBM growth and progression. Mouse glioma cells (CT-2A) and human patient–derived glioma stem cells (N08-30) were treated with MK-2206 or iCRT3 to evaluate effects on cell viability, apoptosis, β-catenin transcriptional activity, and Wnt gene and protein expression *in vitro. In vivo*, luciferase-expressing CT-2A cells were intracranially implanted in C57BL/6J mice to assess the impact of MK-2206 treatment on tumor progression using bioluminescence imaging. Both MK-2206 and iCRT3 reduced cell viability in a dose-dependent manner, with MK-2206 demonstrating greater potency in reducing viability, inducing apoptosis, and inhibiting Wnt signaling. MK-2206 significantly suppressed tumor growth *in vivo* and reduced expression of phosphorylated Akt and GSK-3β in tumor tissues, confirming disruption of the Akt and Wnt/β-catenin signaling axis. In summary, MK-2206 outperformed iCRT3 efficacy *in vitro,* and suppressed GBM progression *in vivo.* These findings suggest that Akt inhibition via MK-2206 may offer a promising therapeutic strategy for treating GBM characterized by dysregulation of PI3K/Akt or Wnt/β-catenin pathways.

Key words: MK-2206, CT-2A, glioblastoma, Glioma Stem Cells, Akt, β-catenin.

Introduction

Glioblastoma (GBM) is the most commonly occurring malignant brain tumor in adults [1]. Patients diagnosed with GBM have a low median rate of survival of 12–18 months, and standard-of-care (SOC) treatment consists of maximally safe surgical brain tumor resection and concurrent chemoradiotherapy [1-5]. A major drawback of the SOC chemotherapeutic agent temozolomide (TMZ) which is a DNA alkylating agent [6], is the development of chemoresistance [7], observed in unmethylated tumors. This is due to the enhanced activity of the DNA repair enzyme O^6-methylguanine methyltransferase (MGMT), which account for approximately 50% of GBM patients [8-11]. The administration of TMZ chemotherapy in conjunction with low-intensity tumor treating fields (TTFields), or TMZ, in combination with anti-angiogenic agents such as bevacizumab, have been recently explored. However, these therapies have not yielded any significant survival benefits to GBM patients [5, 12-14].

PI3K/Akt signaling pathway is frequently found to be hyperactive in GBM [15]. Studies have demonstrated hyperactivation of Akt in 50% of GBM cases, while mutations or deletions in phosphatase and tensin homolog (PTEN), a negative regulator of PI3K/Akt signaling, is found in 36–44% of GBM cases [3, 16-21]. Hyperactive PI3K/Akt pathway promotes cell survival and uncontrolled cell proliferation, apoptosis inhibition, drug resistance, immune evasion of glioma cells, and ultimately contributes to an immunosuppressive environment [22-24]. Furthermore, high levels of Wnt/β-catenin activity have been observed in radio- and chemo-resistant GBM cells, and studies have suggested that TMZ treatment activates the Wnt/β-catenin signaling pathway, resulting in TMZ- and radio- resistance in GBM [4]. This is consistent with observations that Wnt/β-catenin signaling is strongly connected with poor clinical outcomes, with low patient-survival rates [25-27].

The dynamic interplay between the PI3K/Akt and Wnt/β-catenin pathways contributes to GBM aggressiveness, posing a significant challenge for effective therapeutic targeting. Akt hyperactivation occurs in most human cancers, including GBM, can be triggered by Wnt activation of epidermal growth factor receptor (EGFR) signaling, resulting in PI3K activation, which promotes β-catenin–dependent tumor immune evasion [27, 28]. Hyperactive Akt phosphorylates and inhibits GSK-3β, which results in enhanced β-catenin accumulation and reciprocally, β-catenin/T□cell factor-4 (TCF-4) interaction, directly regulates Akt1 in glioma [29, 30]. Furthermore, loss of APC activity from β-catenin destruction complex activates mTORC1 by β-catenin accumulation [31, 32] and mTORC1 promotes Akt activity [33, 34]. β-catenin, the major transcriptional driver of Wnt/β-catenin pathway, is also responsible for growth and maintenance of GBM cells through activation of Wnt target genes [35, 36]. These data demonstrate that Akt and β-catenin play an important role in the crosstalk of PI3K/Akt and Wnt/β-catenin signaling pathways, facilitating convergent pathway signaling that promotes GBM cell proliferation, survival, and invasion. As GBM is characterized by abnormal activation of the PI3K/Akt and Wnt/β-catenin signaling pathways, blocking a common effector for both pathways, such as Akt or β-catenin or a dual-targeting strategy may prove to be more effective, than conventional single pathway targeting therapies. Therefore, we hypothesize that inhibiting Akt and β-catenin, that are key central regulatory components of the PI3K/Akt and Wnt/β-catenin pathway, will suppress GBM growth and progression.

We assessed the inhibitory effects of the small molecule inhibitors MK-2206 2HCl (Akt and β-catenin inhibitor), and iCRT3 (β-catenin inhibitor), on the growth, proliferation, apoptosis, and β-catenin transcriptional activity of CT-2A mouse glioma cells and N08-30 human glioma stem cells, *in vitro*. MK-2206 is a potent and selective allosteric Akt inhibitor demonstrating *in vitro*

and *in vivo* antitumor activity as a single agent [37] across cancer types, including solid tumors. iCRT3 disrupts the interaction between β-catenin and TCF of the Wnt signaling pathway. We also evaluated the efficacy of MK-2206 in reducing the progression of aggressive, syngeneic CT-2A GBM tumors in immunocompetent C57BL/6J mice. Immunocompetent animal models provide a physiologically relevant tumor microenvironment with an intact host immune system [38]. CT-2A cells lack functional PTEN, which is associated with the activation of PI3/Akt and subsequent activation of the Wnt/β-catenin pathway [29, 39]. CT-2A tumors parallel human high-grade gliomas (hHGG) both have glioma stem cell markers, high mitotic index and cell density, nuclear polymorphism, hemorrhagic pseudo palisading necrosis, and microvascular proliferation [40, 41]. Tumor-induced immunosuppression is also exacerbated by PTEN deficiency [3]. An immunocompetent CT-2A model is ideally suited for determining therapeutic effectiveness of interventions that slow the progression of GBM caused by PTEN deficiency. Overall, our findings demonstrate that, MK-2206 outperformed iCRT3 *in vitro,* and suppressed CT-2A tumor growth, *in vivo*. Akt inhibitors such as MK-2206, holds promise as a therapeutic candidate for the specific treatment of PTEN-deficient GBM, where Akt and Wnt/β-catenin signaling is upregulated.

Materials and Methods

Cell Lines and Cell Culture

CT-2A mouse glioma cells (Millipore-Sigma, Cat. No.: SCC194) which harbor a PTEN mutation, were cultured in Dulbecco's Modified Eagle's Medium (DMEM), high glucose (Cytiva, Cat. No.: SH30081.02), supplemented with 10% fetal bovine serum (Neuromics, Cat. No.: FBS002.) and 1% penicillin/streptomycin (Gibco, Cat. No.: 15070-063).

Human glioblastoma cells (N08-30) were isolated from primary human GBM tissue according to procedures approved by Emory University Institutional Review Board protocol 45796 (D.J.B.). N08 cells were maintained in Neurobasal-A Medium (Gibco, Cat. No.:10888-022) containing 1% penicillin-streptomycin and 0.5% L-glutamine (Gibco, Cat. No.: 35050061, supplemented with 1% N-2 supplement (Gibco, Cat. No.: 17502048), 2% B-27 supplement (Gibco, Cat. No.: 17504044), rhFGF (10 ng/ml, R&D system Cat. No.: 233-FB-025/CR), hEGF (20 ng/ml Gibco, Cat. No.: AF100151MG), and 0.4% heparin (STEMCELL, Cat. No.: 07980). Cells were incubated at 37°C with 5% CO_2 and 95% relative humidity. Cells were fed with fresh medium, every 2 days, unless passaged, or harvested for use in *in vitro* and *in vivo* assays.

MTS Cell Viability Assay

CT-2A cells were seeded in 96-well plates at a density of 10K cells/well and left to attach for 24 h in a 37°C cell culture incubator. Subsequently, cells were treated with or without drug/siRNA media and incubated for an additional 48h. Then 20 μL MTS reagent (Abcam, Cat. No.: ab197010) was directly added to each well, and cells were incubated for 2-3 h. Absorbance at 490 nm was measured by using a microplate reader (SpectraMax iD5, Molecular Device). Absorbance values are directly proportional to the number of viable cells.

Detection of Cell Apoptosis

CT-2A cells were seeded on a 6-well plate at 600K per well to attach for ~24h and then treated with or without drug media and incubated for another 24 h. Apoptotic cells were determined using the Annexin V-FITC/PI Apoptosis Kit (Abnova, Cat.: KA3805). All procedures were performed according to the user manual with slight modification. Briefly, cells were washed with 4 mL of cold PBS and resuspended in 100 μL of 1X binding buffer. Then the binding buffer was removed from the cell pellet, and the cell pellet was resuspended again in cold 1X binding buffer. 5 μL of Annexin V-FITC was added to 100 μL cell suspension (120K cells) in each appropriate tube and incubated for 11 minutes at room temperature in the dark. Then 5 μL of PI solution was added and incubated for 6 min at room temperature in dark. Finally, the binding buffer with Annexin V-FITC and PI was removed, and cells were resuspended in 100 μL 1X binding buffer and analyzed by flow cytometry (Quanteon Flow Cytometer, Agilent) within 1 hour.

Caspase 3/7 Activity

Caspase-3/7 activity was measured by Caspase-Glo® 3/7 Assay System (Promega, Cat.: G8091) according to the manufacturer instructions with slight modifications. Briefly, cells were seeded in white wall clear bottom 96-well plates at a density of 20K cells/well, allowed to attach for ~24h and then treated with or without drug containing media (100 μL) and incubated for another 6 h. Cell culture plate was removed from incubator and allowed to equilibrate to room temperature for ~10 min. Then 100μl of Caspase-Glo® 3/7 Reagent was added to each well and mixed well by pipetting and incubated for 1 hour at RT. Finally, luminescence was measured following treatment using a microplate reader (SpectraMax iD5, Molecular Device). Here, luminescence is directly proportional to the caspase-3/7 activity.

Engineering CT-2A Mouse Glioma Cells

Pre-made lentivirus (GenTarget Inc, Cat. No.: LVP010-P-PBS) constitutively expressing Luciferase (Luc) reporter under the 8xTcf Wnt responsive promoter (mCMV) was used to infect CT-2A cells. This lentivirus also contains the puromycin selection marker, under the RSV promoter, and the cells were selected with 1 µg/ml puromycin. This cell line was used for all subsequent *in vitro* experiments. CT-2A-Luc-Puro cells were generated by transducing cells with EF1-GFP-T2A-fLuc-Puro lentivirus (SignaGen Laboratories, Cat. No.: SL100329) at an MOI of 50. Cells were selected and maintained with 1 µg/ml puromycin for >2 weeks to generate stable cell lines. This cell line was used for *in vivo* experiments.

Plasmid and siRNA Transfection

CT-2A cells were transfected using Lipofectamine RNAiMAX (ThermoFisher, Cat. No.: 13778030) by Akt siRNA (ON-TARGET plus siRNAs for Akt1, 2, 3 were purchased from Dharmacon). N08-30 cells were co-transfected using Lipofectamine 3000 (Invitrogen, Cat. No.:100022050) by 200 ng firefly luciferase expressing TOP and FOP-flash reporter plasmids (Millipore, MA, USA). Transfections were performed according to the manufacturer's protocol. Briefly, cells were seeded in 24 well plate at 80K cells/well and allowed to attach. At 60-70% confluent, fresh media was added, and then they were transfected with siRNA/plasmid using transfection reagent. The concentrations of siRNAs or plasmid were chosen based on dose-response studies for knockdown efficiency.

TOPFlash/FOPFlash Reporter Assay

To evaluate the β-catenin transcriptional activity in human GBM cells, we used a pair of luciferase reporter constructs, TOPFlash and FOPFlash (a gift from Dr. Ken-ichi Takemaru, Stony Brook University). TOPFlash contains three copies of TCF/LEF binding site, and

FOPFlash contains a mutated TCF/LEF binding site. Briefly, N08-30 cells were seeded in a precoated 24-well plate with Geltrex™ at 80K cells/well and allowed to attach. At 60-70% confluent, fresh drug media was added, and cells were transfected with TOP and FOPFlash plasmid (200 ng) using Lipofectamine 3000 reagent (Invitrogen), according to the manufacturer's protocol. After 48 h of incubation, cells were collected and lysed with 1X passive lysis buffer (Promega, Cat. No.: E1941) using standard protocol conditions. From each treatment group, 10 µL cell lysate was directly added to each well of a 96 well white opaque plate. 50 µL Luciferase Assay Reagent II (LARII, Promega, Cat. No.: E1500) was added, and luminescence signals were detected immediately using a microplate reader (SpectraMax iD5, Molecular Device). Luciferase activity was normalized to total protein amount.

Luciferase Reporter Assay

To evaluate the β-catenin transcriptional activity in mouse glioma cells, we created a stable CT-2A cell line expressing luciferase signal as described in tumor cell engineering section. Briefly, CT-2A cells were seeded in a 24-well plate at 80K cells/well and allowed to attach. At 60-70% confluent, fresh drug media was added, or cells were transfected with Akt siRNA (100 ng) using Lipofectamine RNAiMAX reagent (Invitrogen, Cat. No.: 13778030) according to the manufacturer's protocol. After 48 h of incubation, cells were collected and lysed with 1X passive lysis buffer. From each treatment group, 10 µL cell lysate was directly added to each well of a 96 well white opaque plate. 50 µL LARII was added, and luminescence signals were measured immediately using a microplate reader (SpectraMax iD5, Molecular Device). Luciferase activity was normalized to total protein amount.

Western Blotting

Cells were washed with cold 1X PBS and lysed with RIPA buffer (Thermo Scientific, Waltham, Cat. No.: J63306-AK) supplemented with Halt Protease & Phosphatase Inhibitor cocktail (Thermo Scientific, Cat. No.:1861281) on ice for 30-minutes. Cell lysates were centrifuged at 12,000 RPM for 20 minutes at 4□C and supernatant was used to determine protein concentration by Pierce BCA protein assay kit (Thermo Fisher Scientific, Cat. No.: 23227). Proteins (10–30 μg) were separated through electrophoresis, using Mini-PROTEAN TGX Stain-Free Precast Gels (BIO-RAD, Cat. No.: 4568094) and then transferred to nitrocellulose membrane (Bio-Rad, Cat. No.: 1704271) using a turbo transfer system (BIO-RAD). The membranes were incubated with the respective antibodies from Cell Signaling Technology (pAKT-S473, Cat No.: 4060T; Akt pan, Cat No.: 4691T; pGSK3-S9, Cat No.: 5558T; GSK3β, Cat No.: 12456T; and β-actin, Cat No.: 4970S) in 5% BSA in TBST at 4°C overnight, followed by incubation with secondary antibodies at room temperature for 1 h. Finally, protein bands were developed using Clarity western ECL substrate (BIO-RAD, Cat No.: 170-5060) using ChemiDoc MP imaging system (BIO-RAD). Protein amount was normalized to their respective β-actin loading control and band intensities were quantified by ImageJ software.

Quantitative Real-Time PCR

Total RNA was extracted from drug treated cells using Total RNA Kit (Cat. No.: R6834-01; Omega BIO-TEK, Norcross, GA, USA) according to the manufacturer's instructions, and concentrations were measured using a Nano Drop 800 (Thermo Scientific). cDNA was synthesized using iScript^{TM} reverse transcription supermix (BIO-RAD; Cat.: 1708840). Primers were purchased from ThermoFisher and the assay ID of the primer sequences are: Myc; Mm00487804_m1, Ccnd1; Mm00432359_m1, Axin2; Mm00443610_m1 and GAPDH; Mm99999915_g1. qRT-PCR was performed in Quant Studio 5 Real Time PCR System (Applied

Biosystem) using TaqMan™ Universal Master Mix II, no UNG (Fisher Sci. Cat. No.: 44-400-40) according to the manufacturer's protocol. The $2^{-\Delta\Delta CT}$ method was used to calculate the relative gene expression levels normalized to GAPDH and compared to control samples.

Animal Studies

All animal studies were approved by the University of Georgia Institutional Animal Care and Use Committee (IACUC), and all protocols were followed according to the National Institutes of Health's Guide for the Care and Use of Laboratory Animals. Briefly, male C57BL/6J mice (6-8 weeks old, 18–30 g) were purchased from the Jackson Laboratory. The mice were housed in a temperature-controlled and light-controlled environment in the animal facility. Luciferase-expressing CT-2A cells (~250K cells resuspended in 7 μL DMEM medium) were injected in the right frontal cortex at coordinates, +1.0 mm (AP), +1.5 mm (ML), and +3.0 mm (DV), relative to bregma, to induce mouse brain tumors. Bioluminescence imaging (BLI) was performed to confirm tumor uptake on day 7, and mice were randomly divided into two groups (12 mice/group; MK-2206 and vehicle control). Intraperitoneal drug treatments were started on day 8, according to the following regimen: MK-2206, 100 mg/kg body weight/day, and vehicle control (20% β-captisol) at the same frequency until day 28. Tumor growth and luciferase activity were monitored every 7 days using BLI imaging in NEWTON 7 (VILBER smart imaging). Animal were anaesthetized using 1.5% isoflurane and, D-luciferin potassium salt (GOLDBIO, Cat. No.: LUCK-100) was injected (IP, 150mg/kg) 12-13 minutes prior to *in vivo* imaging for BLI imaging. Body weight and overall behavior were monitored daily for the first 7 days and then every other day until terminal euthanasia.

Brain Tissue Processing and Histology

At the 29-day time point, the mice were deeply anesthetized using 3-4% isoflurane gas and transcardially perfused with Phosphate Buffered Saline (PBS, pH 7.2) for 15-20 min to perfuse blood from vasculature in brain tissue. Post-perfusion, brains was carefully extracted, flash frozen on dry-ice, and stored at −80°C, until cryosectioning. The initial 0.5 μm from olfactory bulb (16 μm thick, 31 sections) was discarded, and total of 120 coronal sections (16 μm thick) were serially collected on 30 slides (4 sections per slide) along the rostro-caudal axis. The tissue sections were processed for hematoxylin and eosin (H&E) staining, and the stained slides were imaged at 20X magnification using a digital slide scanner (Aperio AT2) by the University of Georgia Histology Laboratory.

Immunohistochemistry

1 slide per animal (4 sections per slide) was used for immunostaining. Briefly, sections were fixed using 4% paraformaldehyde in 0.4 M sucrose for 20 minutes after rinsing with PBS and blocked using 5% BSA in PBST (0.5% Triton-X 100 in PBS) for 1 hour at room temperature (RT). Then sections were incubated with primary antibody (pAKT-S473, Cat No.: 4060T; pGSK3-S9, Cat No.: 5558; Cell Signaling Technology) overnight, at 4 °C. Next day, the sections were rinsed in PBS, blocked again for 1 hour at RT, and incubated with Alexa Fluor 488 Goat-anti-rabbit (Cat No.: A11008, Invitrogen) secondary antibody for 3 hours at RT. Finally, sections were stained with nuclear stain Hoechst 33342 (Invitrogen, Dilution 1:200 from 1 mg/ml stock), airdried and cover slipped with Fluoromount-G Mounting Medium (Southern Biotech, AL) and stored at −20°C until imaged. The immunohistochemically stained brain sections were imaged using the Zeiss LSM880 confocal microscope. Images were acquired at 20X magnification. The acquired images were analyzed using CellProfiler [42] and plotted as area normalized to

Hoechst[+] cells. A total of 5 mice from each treatment group were randomly selected for immunostaining studies.

Statistical analysis

GraphPad Prism 9.5.1 was used to perform all statistical analyses and plot the data. The data was tested for normality and equal variance assumptions using the Shapiro-Wilk test and F-test, respectively. Group comparisons were performed using One-way ANOVA with Tukey's multiple comparison test. Unpaired two-tailed Student's t-tests were used to compare data between two groups. p-values <0.05 were considered statistically significant.

Results

1. Akt and β-catenin inhibitors promoted apoptosis, and suppressed the growth and proliferation of CT-2A cells, *in vitro*

To investigate the effect of Akt (MK-2206) and β-catenin (iCRT3) inhibitors on PTEN mutated mouse glioma (CT-2A) cells (Figure **1A**), we performed the MTS cell viability assay. We found that MK-2206 and iCRT3 suppressed cell growth and proliferation in a dose-dependent manner following 48h of treatment when compared to vehicle control group (Figure **1B & 1C**). The half-maximal inhibitory concentrations (IC_{50}) values, for cell viability inhibition, were found to be 8.8 µM for MK-2206, and 45 µM for iCRT3.

We measured Caspase-3/7 activity to determine whether MK-2206 and iCRT3 cell growth inhibition led to increased apoptosis of CT-2A cells. A small, but non-significant increase in caspase- 3/7 activity was observed in iCRT3-treated cells following 6h treatment, while MK-2206 treatment significantly increased caspase activity, compared to vehicle controls (Figure **1D**).

To confirm increased apoptosis in inhibitor treated cells, the percentage of apoptotic cells were measured using Annexin V+ staining. Following a 24h treatment, the percent apoptotic cells significantly increased in both, 10 µM MK-2206 and 50 µM iCRT3 treated cells, compared to vehicle controls (Figure **1E**).

Taken together, Akt/β-catenin pathway contributes to uncontrolled cell growth, proliferation, and apoptosis inhibition in CT-2A cells, which could be inhibited by treating these cells with the Akt or β-catenin inhibitors.

2. Akt and β-catenin inhibitors suppressed transcriptional activity of β-catenin in CT-2A cells

We compared the effects of MK-2206 and iCRT3 on suppressing the Wnt/β-catenin signaling pathway. To determine whether MK-2206 and iCRT3 impacts β-catenin/TCF transcriptional - activity, we used luciferase expressing stable CT-2A cell lines having 8X TCF binding cites for β-catenin. When compared to vehicle, both MK-2206 and iCRT3 treated CT-2A cells (48h) exhibited significantly decreased luciferase reporter activity, indicating that Akt and β-catenin inhibitors downregulated β-catenin/TCF transcriptional activity in CT-2A cells (Figure **2A**).

Western blot analysis was performed to investigate the effects of MK-2206 and β-catenin treatment on phosphorylated Akt (p-AKT) and phosphorylated GSK-3β (p-GSK-3β) levels. CT-2A cells were treated with MK-2206 and iCRT3 for 1h, lysed and total protein was extracted for western blot analysis. MK-2206 treatment significantly down-regulated both p-AKT and p-GSK-3β proteins in CT-2A cells, compared to untreated vehicle controls (Figure **2B**). iCRT3 showed no significant effects on p-AKT or p-GSK-3β levels, which is in line with its mechanism of action of preventing TCF- β-catenin binding.

Given the positive correlation between hyperactive Akt and activation of Wnt/β-catenin signaling pathway, we hypothesized that the expression of known Wnt/β-catenin target genes *cyclin D1*, *c-myc* and *Axin2* would be transcriptionally regulated by the Akt/β-catenin pathway in CT-2A cells. Results indicate that Akt inhibitor MK-2206 treatment significantly downregulated the expression of *cyclin D1*, *c-myc* and *Axin2*. Though β-catenin iAxin2iCRT3 treatment significantly reduced the expression of *Axin2*, it concomitantly enhanced the expression of *c-Myc* and induced no significant changes in *cyclin D1,* when compared to vehicle control, suggesting that MK-2206 can block Wnt/β-catenin signaling more effectively than iCRT3 (Figure **2C**).

These results suggest that Akt activation resulting from PTEN mutation correlates with the levels of GSK-3β phosphorylation and Wnt/β-catenin target gene expression in GBM cells. Taken together, the above results suggest that Akt/β-catenin pathway inhibition by MK-2206 can repress the proliferation of CT-2A cells by inhibiting PI3K/Akt and Wnt/β-catenin signaling pathways and the expression of associated genes.

3. Akt and β-catenin inhibitors suppressed human glioma stem cell growth, β-catenin transcriptional activity and promoted apoptosis, *in vitro*

The effects of Akt and β-catenin inhibitors were then evaluated on N08-30, patient-derived glioma stem cell line, carrying both EGFR amplification and PTEN mutations. To investigate the inhibitory effects of MK-2206 and iCRT3 on N08-30, the MTS cell viability assay was performed. MK-2206 and iCRT3 suppressed glioma stem cell growth in a dose-dependent manner following 48h of treatment compared to the vehicle control group (Figure **3A & 3B**). IC$_{50}$ values of MK-2206 for inhibiting cell viability was found to be 9.8 μM, and >100 μM for iCRT3. These results indicated that iCRT3 is less cytotoxic to patient-derived glioma stem cells, compared to MK-2206.

The ability of MK-2206 and iCRT3 to induce glioma stem cell apoptosis was evaluated by quantifying the percentage of Annexin V+ cells and caspase 3/7 activity. As shown in Figure **3C** and **3D**, following treatment, caspase 3/7 activity and percent apoptotic cells were significantly increased in MK-2206 treated cells, compared to vehicle controls and iCRT3. iCRT3 treatment neither enhanced caspase 3/7 activity, nor Annexin V+ cells, compared to vehicle controls.

To determine whether MK-2206 and iCRT3 impacts β-catenin/TCF-interaction in N08-30 human glioma stem cells, we transfected the cells with TOPflash and FOPflash reporter plasmids. When compared to the vehicle control, MK-2206 and iCRT3 treated cells significantly

decreased TOPflash reporter activity, indicating that both inhibitors downregulated β-catenin/TCF-induced transcription (Figure **3E**). There was no change in the FOPflash reporter activity, which served as mutant negative control.

Taken together, Akt inhibition by MK-2206 outperformed β-catenin inhibition by iCRT3 by suppressing patient-derived glioma stem cell growth, and inducing apoptosis *in vitro*, while both MK-2206 and iCRT3 significantly downregulated β-catenin/TCF-induced transcriptional activity in these cells.

4. Akt siRNA suppressed CT-2A cell growth, transcriptional activity of β-catenin, and reduced Wnt target gene expression

In our previous experiments, while both Akt inhibitor, MK-2206, and β-catenin inhibitor, iCRT3, showed promise, MK-2206 consistently outperformed iCRT3, prompting us to focus on Akt mediated Wnt/β-catenin pathway inhibition in future studies. To confirm whether the inhibitory effect of MK-2206 was specifically mediated through Akt inhibition in our previous findings, we performed siRNA mediated knockdown of Akt in CT-2A cells.

Before assessing the functional impact from the loss of *Akt* isoforms on CT-2A cell growth, we first quantified their transcript levels (Figure **4A**). We observed that *Akt1* had the highest expression level, followed by *Akt2*, while *Akt3* exhibited the lowest levels. The Akt isoforms (*Akt1*, *Akt2*, *Akt3)* were then treated with siRNAs individually or in combination. Subsequent viability assessments demonstrated that knockdown of multiple isoforms, including *Akt3*, resulted in a significant reduction in cell growth (Figure **4B**).

To determine whether knocking down of individual Akt isoforms affects β-catenin/TCF-interaction and Wnt/β-catenin target gene expression, we performed luciferase reporter assay and

qRT-PCR in CT-2A cells. Silencing any individual *Akt* isoform led to a significant decrease in β-catenin/TCF-mediated transcription (Figure **4C**), with *Akt3* showing the most potent effect.

Individual Akt knockdown had no effect on target gene expression (data not shown). However, when all three *Akt* isoforms were knocked down, we observed a significant reduction in *cyclin D1*, and *c-myc* (Figure **4D & 4E**). These findings confirmed that the previous effects of Akt inhibition by MK-2206 were due to the inhibition of the AKT ruling out nonspecific mechanisms.

5. MK-2206 suppressed glioma tumor progression in immunocompetent mice

To evaluate the inhibitory effect of MK-2206 on tumor progression *in vivo*, we performed orthotopic inoculations of luciferase expressing mouse CT-2A cells into immunocompetent C57BL/6J mice. CT-2A-Luciferase (CT-2A-Luc) cells were intracranially inoculated in the right frontal cortex of C57BL/6J mice. Tumor uptake was confirmed on Day 7 following an intraperitoneal (IP) D-luciferin injection (150mg/kg body weight) and subsequent bioluminescence (BLI) imaging. Mice were then randomly divided into two groups: one receiving daily IP injections of the MK-2206 (100mg/kg body weight) and vehicle control group receiving 20% β-captisol (Figure **5A**).

Mice body weights and any abnormalities (e.g., coordination, tremor, head tilting, eye position, ruffled fur) were monitored throughout the study period. Mice body weight was found to trend downwards in the MK-2206-treated mice; however, the observed weight loss was not statistically significant. (Figure **5B).**

Tumor progression was monitored over time using BLI imaging (Figure **5C**). Our results demonstrated that the tumor size (BLI signal) of the MK-2206-treated group was significantly

reduced compared to vehicle group (Figure **5D**). At study endpoint of 29-days, mice were euthanized, and tumor tissues were collected for histological analysis. Representative hematoxylin and eosin (H&E) staining confirmed the tumor burden reduction in MK-2206 treated group compared to vehicle control group (Figure **5E)** which was consistent with BLI results.

6. **Akt inhibition suppressed downstream protein phosphorylation in tumor tissue**

To determine whether intraperitoneal MK-2206 administration inhibited the expression of the components of Akt and Wnt/β-catenin signaling pathway, tumor tissue sections were analyzed by immunohistochemistry for p-AKT and p-GSK3β. The levels of p-AKT-Ser473 and p-GSK-3β-Ser9 were significantly downregulated in MK-2206 treated mice, compared to vehicle controls, indicating that MK-2206 treatment suppressed Akt activity in GBM tumors, *in vivo* (Figure **6A** & **6B** respectively).

Altogether, our *in vivo* data demonstrated that Akt inhibition by MK-2206 significantly suppressed glioma tumor progression, likely by inhibiting both PI3K/Akt and Wnt/β-catenin signaling. This supports the notion that Akt inhibitors may have therapeutic utility for GBM with elevated PI3K/Akt and Wnt/β-catenin signaling.

Discussion

PI3K/Akt and Wnt/β-catenin signaling are two key drivers of GBM malignancy and therapeutic resistance [20, 27, 29, 43]. Genetic alterations in the PI3K/Akt/mTOR signaling cascade, a key driver of cell survival and apoptosis inhibition, were detected in 88% of all GBM cases [44]. While expression of multiple proteins in the pathway are dysregulated, hyperactivation of Akt alone occurs in approximately 50% of GBM cases [16, 17, 22]. Additionally, β-catenin, the

major transcriptional driver of Wnt/β-catenin pathway is considered a prognostic marker, due to enhanced mRNA and protein expression in GBM [45, 46]. The crosstalk between these two pathways contributes to GBM's invasive phenotype. Akt-mediated Wnt/β-catenin activation drives the transcriptional activation of Wnt target genes such as *c-Myc, Cyclin D1,* and *Axin2,* which are responsible for GBM growth, proliferation, and immune evasion [27, 36, 43, 47]. However, the role of Akt mediated Wnt/β-catenin role in glioma stem cell growth and proliferation has not been sufficiently elucidated. To address this knowledge gap, we evaluated the therapeutic potential of inhibiting Akt and β-catenin signaling in mouse and human patient-derived glioma stem cells, *in vitro,* and validated findings using a syngeneic, immunocompetent mouse GBM model *in vivo.*

We compared the inhibitory effects of MK-2206, a potent and selective allosteric Akt inhibitor [37]; and iCRT3, a β-catenin-TCF interaction inhibitor [48], in both mouse glioma cells (CT-2A) and patient-derived glioma stem cells (N08-30). The enhanced apoptotic response and reduced cell viability we observed in MK-2206 treated group, explains the central role of Akt in regulating pro-survival pathways in GBM. Mechanistically, Akt inhibition promotes apoptosis by reducing the phosphorylation of pro-apoptotic proteins such as BAD and BIM [49, 50] and by activating apoptotic gene expression via FOXO3a dephosphorylation [51]. Additionally, Akt inhibition suppresses NF-κB activity and reduce the expression of anti-apoptotic targets like *Bcl-xL* [52, 53]. These mechanisms highlight MK-2206's pro-apoptotic efficacy over iCRT3 in GBM models.

Overexpression of *c-Myc* and *cyclin D1*, the downstream targets of Akt and Wnt/β-catenin signaling contributes to uncontrolled cell proliferation and inhibition of apoptosis in gliomas [54-57]. p-Akt promotes cell survival and proliferation by modulating downstream targets including

mTOR, FOXO3a and MDM2, while simultaneously inhibiting GSK-3β phosphorylation [58]. GSK-3β inhibition stabilizes β-catenin and increases the expression of oncogenic Wnt/β-catenin target genes such as *c-Myc* and *cyclin D1* [59, 60]. In our findings, MK-2206 downregulated p-Akt and p-GSK-3β in mouse glioma cells, but iCRT3 lacked this effect, indicating limited target modulation. Similar to our results, previous studies using MK-2206 as single agent or in combination have shown that Akt inhibition leads to decreased phosphorylation of Akt contributing to reduced GBM cell proliferation and enhanced apoptosis [61, 62]. The unexpected upregulation of *c-Myc,* by β-catenin inhibitor iCRT3 might be due to the compensatory effects of other signaling cascades [63]. These findings emphasize Akt's role as a central regulator linking PI3K/Akt and Wnt/β-catenin signaling in GBM. Further evidence for therapeutic Akt inhibition is supported by the recent (2023) FDA-approval of the pan-Akt inhibitor Capivasertib (AZD5363) [64, 65]. Both MK-2206 and Capivasertib act as pan-Akt inhibitor [62, 66, 67] and suppress cell growth and proliferation in a range of cancers through targeting downstream signaling pathways including mTORC1, GSK-3β, and FOXO3a [62, 68, 69]. In preclinical and Phase I clinical studies, Capivasertib has demonstrated radio sensitizing effects and reduced cell viability targeting Akt and GSK-3β phosphorylation in oral and breast cancer model [70, 71] indicating a mechanistic profile overlapping with MK-2206 [67]. Results from these studies have demonstrated promising anti-tumor activity across multiple cancers by suppressing Akt signaling [72, 73]. Our findings with MK-2206 similarly support the therapeutic value of targeting Akt, in PTEN-deficit Akt-driven glioblastoma.

From our *in vitro* studies, the Akt inhibitor MK-2206 consistently outperformed iCRT3, and therefore we limited our investigations to studying Akt mediated Wnt/β-catenin pathway inhibition in suppressing GBM cell growth, *in vivo*. Previous studies investigating the therapeutic

effects of the Akt inhibitor MK-2206 in GBM models were often limited by the use of animals with low tumor burden or immunodeficient backgrounds, limiting the clinical relevance of tumor-host interactions [27, 62, 74, 75]. Consequently, the single-agent therapeutic potential of MK-2206 against highly invasive GBM tumors in immunocompetent animals has not yet been fully investigated.

In our study, we employed a syngeneic, immunocompetent C57BL/6J mouse model with established intracranial CT-2A tumors and monitored treatment effects using BLI imaging. MK-2206 treatment significantly suppressed tumor progression compared to vehicle group, which was consistent with previously reported findings in immunodeficient mouse models [62]. H&E staining of MK-2206-treated tumor tissues further supported tumor suppression revealing reduced hyper-nucleated and necrotic area, and a reduced midline shift, which are distinguishing features of aggressive and invasive GBM phenotypes [76, 77]. Additionally, decreased phosphorylation of Akt and GSK-3β in tumor tissue, confirming GSK-3β–mediated Wnt/β-catenin signaling inhibition by Akt inhibitor MK-2206. Overall, our *in vitro* and *in vivo* findings support the therapeutic potential of targeting hyperactive Akt with MK-2206 in PTEN-deficient GBM driven by aberrant Akt-Wnt/β-catenin axis.

AUTHOR CONTRIBUTIONS

MMS, AS, LK and SLS contributed to the study conception and design; MMS performed all *in vitro* experiments; AS created luciferase expressing CT-2A cell line; MMS, NG and AD performed *in vivo* tumor inoculation study; MMS conducted all *in vivo* imaging and analysis; MMS and NG performed immunohistochemical staining and imaging, and AD performed IHC analysis; EM assisted with cell culture and caspase 3/7 assay; MMS wrote the original draft and, AS, NG, LK and SLS contributed to writing and revision of the manuscript.

ACKNOWLEDGEMENTS & FUNDING

We thank UGA's Biomedical Core Facility (BMC) for providing us with access to confocal microscopes and *in vivo* Bioluminescence Imaging facilities. We also thank Shanmathi Ramasubramanian for assisting during pilot animal study. We also thank Dr. Ken-ichi Takemaru, Stony Brook University, for providing luciferase reporter plasmids. This research was supported by the Georgia Research Alliance, the National Science Foundation Engineering Research Center for Cell Manufacturing Technologies (NSF No. EEC-1648035, awarded to Steven Stice).

DISCLOSURES

The authors declare that there are no conflicts of interest.

REFERENCES

1. Wirsching, H.G., E. Galanis, and M. Weller, *Glioblastoma.* Handb Clin Neurol, 2016. **134**: p. 381-97.
2. Hashemi, M., et al., *Progress in targeting PTEN/PI3K/Akt axis in glioblastoma therapy: Revisiting molecular interactions.* Biomed Pharmacother, 2023. **158**: p. 114204.
3. Pearson, J.R.D. and T. Regad, *Targeting cellular pathways in glioblastoma multiforme.* Signal Transduct Target Ther, 2017. **2**: p. 17040.
4. Ma, Z., et al., *WNT signaling modulates chemoresistance to temozolomide in p53-mutant glioblastoma multiforme.* Apoptosis, 2022. **27**(1-2): p. 80-89.
5. Tomar, V.S., V. Patil, and K. Somasundaram, *Temozolomide induces activation of Wnt/beta-catenin signaling in glioma cells via PI3K/Akt pathway: implications in glioma therapy.* Cell Biol Toxicol, 2020. **36**(3): p. 273-278.
6. Lee, S.Y., *Temozolomide resistance in glioblastoma multiforme.* Genes Dis, 2016. **3**(3): p. 198-210.
7. Alonso, M.M., et al., *Adenovirus-based strategies overcome temozolomide resistance by silencing the O6-methylguanine-DNA methyltransferase promoter.* Cancer Res, 2007. **67**(24): p. 11499-504.
8. Aldape, K., et al., *Glioblastoma: pathology, molecular mechanisms and markers.* Acta Neuropathol, 2015. **129**(6): p. 829-48.
9. Yang, S.H., et al., *Metformin treatment reduces temozolomide resistance of glioblastoma cells.* Oncotarget, 2016. **7**(48): p. 78787-78803.
10. Erasimus, H., et al., *DNA repair mechanisms and their clinical impact in glioblastoma.* Mutat Res Rev Mutat Res, 2016. **769**: p. 19-35.
11. Wick, W., et al., *MGMT testing--the challenges for biomarker-based glioma treatment.* Nat Rev Neurol, 2014. **10**(7): p. 372-85.
12. Gregory, J.V., et al., *Systemic brain tumor delivery of synthetic protein nanoparticles for glioblastoma therapy.* Nat Commun, 2020. **11**(1): p. 5687.
13. Zou, Y., et al., *Single siRNA Nanocapsules for Effective siRNA Brain Delivery and Glioblastoma Treatment.* Adv Mater, 2020. **32**(24): p. e2000416.
14. Vazquez Cervantes, G.I., et al., *New Immunotherapeutic Approaches for Glioblastoma.* J Immunol Res, 2021. **2021**: p. 3412906.
15. Colardo, M., M. Segatto, and S. Di Bartolomeo, *Targeting RTK-PI3K-mTOR Axis in Gliomas: An Update.* Int J Mol Sci, 2021. **22**(9).
16. Crespo* S, K.M., Arcaro A., *The role of the PI3K/AKT/mTOR pathway in brain tumor metastasis.* J Cancer Metastasis Treat, 2016. **2**: p. 80-9.
17. Langhans, J., et al., *The effects of PI3K-mediated signalling on glioblastoma cell behaviour.* Oncogenesis, 2017. **6**(11): p. 398.
18. Brennan, C.W., et al., *The somatic genomic landscape of glioblastoma.* Cell, 2013. **155**(2): p. 462-77.
19. Cancer Genome Atlas Research, N., *Comprehensive genomic characterization defines human glioblastoma genes and core pathways.* Nature, 2008. **455**(7216): p. 1061-8.
20. Carracedo, A. and P.P. Pandolfi, *The PTEN-PI3K pathway: of feedbacks and cross-talks.* Oncogene, 2008. **27**(41): p. 5527-41.
21. Liu, Y., et al., *PTEN deficiency confers colorectal cancer cell resistance to dual inhibitors of FLT3 and aurora kinase A.* Cancer Lett, 2018. **436**: p. 28-37.

22. Yap, T.A., et al., *First-in-man clinical trial of the oral pan-AKT inhibitor MK-2206 in patients with advanced solid tumors.* J Clin Oncol, 2011. **29**(35): p. 4688-95.

23. Sonoda, Y., et al., *Akt pathway activation converts anaplastic astrocytoma to glioblastoma multiforme in a human astrocyte model of glioma.* Cancer Res, 2001. **61**(18): p. 6674-8.

24. Parsa, A.T., et al., *Loss of tumor suppressor PTEN function increases B7-H1 expression and immunoresistance in glioma.* Nat Med, 2007. **13**(1): p. 84-8.

25. Anastas, J.N. and R.T. Moon, *WNT signalling pathways as therapeutic targets in cancer.* Nat Rev Cancer, 2013. **13**(1): p. 11-26.

26. Nusse, R. and H. Clevers, *Wnt/beta-Catenin Signaling, Disease, and Emerging Therapeutic Modalities.* Cell, 2017. **169**(6): p. 985-999.

27. Du, L., et al., *beta-Catenin induces transcriptional expression of PD-L1 to promote glioblastoma immune evasion.* J Exp Med, 2020. **217**(11).

28. Hu, T. and C. Li, *Convergence between Wnt-beta-catenin and EGFR signaling in cancer.* Mol Cancer, 2010. **9**: p. 236.

29. Ding, Q., et al., *Erk associates with and primes GSK-3beta for its inactivation resulting in upregulation of beta-catenin.* Mol Cell, 2005. **19**(2): p. 159-70.

30. Chen, L., et al., *beta-catenin/Tcf-4 complex transcriptionally regulates AKT1 in glioma.* Int J Oncol, 2011. **39**(4): p. 883-90.

31. Liu, J., et al., *Wnt/beta-catenin signalling: function, biological mechanisms, and therapeutic opportunities.* Signal Transduct Target Ther, 2022. **7**(1): p. 3.

32. Valvezan, A.J., et al., *Adenomatous polyposis coli (APC) regulates multiple signaling pathways by enhancing glycogen synthase kinase-3 (GSK-3) activity.* J Biol Chem, 2012. **287**(6): p. 3823-32.

33. Saxton, R.A. and D.M. Sabatini, *mTOR Signaling in Growth, Metabolism, and Disease.* Cell, 2017. **168**(6): p. 960-976.

34. Sun, K., et al., *The PI3K/AKT/mTOR signaling pathway in osteoarthritis: a narrative review.* Osteoarthritis Cartilage, 2020. **28**(4): p. 400-409.

35. Lee, Y., et al., *WNT signaling in glioblastoma and therapeutic opportunities.* Lab Invest, 2016. **96**(2): p. 137-50.

36. Vallee, A., et al., *Thermodynamics in Gliomas: Interactions between the Canonical WNT/Beta-Catenin Pathway and PPAR Gamma.* Front Physiol, 2017. **8**: p. 352.

37. Coleman, N., et al., *Clinical Development of AKT Inhibitors and Associated Predictive Biomarkers to Guide Patient Treatment in Cancer Medicine.* Pharmgenomics Pers Med, 2021. **14**: p. 1517-1535.

38. Sharma, P., et al., *Tumor microenvironment in glioblastoma: Current and emerging concepts.* Neurooncol Adv, 2023. **5**(1): p. vdad009.

39. Marsh, J., P. Mukherjee, and T.N. Seyfried, *Akt-dependent proapoptotic effects of dietary restriction on late-stage management of a phosphatase and tensin homologue/tuberous sclerosis complex 2-deficient mouse astrocytoma.* Clin Cancer Res, 2008. **14**(23): p. 7751-62.

40. Martinez-Murillo, R. and A. Martinez, *Standardization of an orthotopic mouse brain tumor model following transplantation of CT-2A astrocytoma cells.* Histol Histopathol, 2007. **22**(12): p. 1309-26.

41. Oh, T., et al., *Immunocompetent murine models for the study of glioblastoma immunotherapy.* J Transl Med, 2014. **12**: p. 107.

42. Stirling, D.R., et al., *CellProfiler 4: improvements in speed, utility and usability.* BMC Bioinformatics, 2021. **22**(1): p. 433.

43. Nager, M., et al., *beta-Catenin Signalling in Glioblastoma Multiforme and Glioma-Initiating Cells.* Chemother Res Pract, 2012. **2012**: p. 192362.

44.   Li, X., et al., *PI3K/Akt/mTOR signaling pathway and targeted therapy for glioblastoma.* Oncotarget, 2016. **7**(22): p. 33440-50.

45.   Sareddy, G.R., et al., *Activation of Wnt/beta-catenin/Tcf signaling pathway in human astrocytomas.* Neurochem Int, 2009. **55**(5): p. 307-17.

46.   Liu, C., et al., *Wnt/beta-Catenin pathway in human glioma: expression pattern and clinical/prognostic correlations.* Clin Exp Med, 2011. **11**(2): p. 105-12.

47.   Kahn, M., *Can we safely target the WNT pathway?* Nat Rev Drug Discov, 2014. **13**(7): p. 513-32.

48.   Yan, M., G. Li, and J. An, *Discovery of small molecule inhibitors of the Wnt/beta-catenin signaling pathway by targeting beta-catenin/Tcf4 interactions.* Exp Biol Med (Maywood), 2017. **242**(11): p. 1185-1197.

49.   Datta, S.R., et al., *Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery.* Cell, 1997. **91**(2): p. 231-41.

50.   Yuan, Z., et al., *BIM-mediated AKT phosphorylation is a key modulator of arsenic trioxide-induced apoptosis in cisplatin-sensitive and -resistant ovarian cancer cells.* PLoS One, 2011. **6**(5): p. e20586.

51.   Santo, E.E., et al., *FOXO3a is a major target of inactivation by PI3K/AKT signaling in aggressive neuroblastoma.* Cancer Res, 2013. **73**(7): p. 2189-98.

52.   Cahill, K.E., R.A. Morshed, and B. Yamini, *Nuclear factor-kappaB in glioblastoma: insights into regulators and targeted therapy.* Neuro Oncol, 2016. **18**(3): p. 329-39.

53.   Wang, Z., G. Luo, and Z. Qiu, *Akt inhibitor MK-2206 reduces pancreatic cancer cell viability and increases the efficacy of gemcitabine.* Oncol Lett, 2020. **19**(3): p. 1999-2004.

54.   Zhang, X., et al., *The effect of cyclin D expression on cell proliferation in human gliomas.* J Clin Neurosci, 2005. **12**(2): p. 166-8.

55.   Kotliarova, S., et al., *Glycogen synthase kinase-3 inhibition induces glioma cell death through c-MYC, nuclear factor-kappaB, and glucose regulation.* Cancer Res, 2008. **68**(16): p. 6643-51.

56.   Pu, P., et al., *Downregulation of Wnt2 and beta-catenin by siRNA suppresses malignant glioma cell growth.* Cancer Gene Ther, 2009. **16**(4): p. 351-61.

57.   Cully, M., et al., *Beyond PTEN mutations: the PI3K pathway as an integrator of multiple inputs during tumorigenesis.* Nat Rev Cancer, 2006. **6**(3): p. 184-92.

58.   Liu, R., et al., *PI3K/AKT pathway as a key link modulates the multidrug resistance of cancers.* Cell Death Dis, 2020. **11**(9): p. 797.

59.   Farago, M., et al., *Kinase-inactive glycogen synthase kinase 3beta promotes Wnt signaling and mammary tumorigenesis.* Cancer Res, 2005. **65**(13): p. 5792-801.

60.   Lecarpentier, Y., et al., *Multiple Targets of the Canonical WNT/beta-Catenin Signaling in Cancers.* Front Oncol, 2019. **9**: p. 1248.

61.   Narayan, R.S., et al., *The allosteric AKT inhibitor MK2206 shows a synergistic interaction with chemotherapy and radiotherapy in glioblastoma spheroid cultures.* BMC Cancer, 2017. **17**(1): p. 204.

62.   Cheng, Y., et al., *MK-2206, a novel allosteric inhibitor of Akt, synergizes with gefitinib against malignant glioma via modulating both autophagy and apoptosis.* Mol Cancer Ther, 2012. **11**(1): p. 154-64.

63.   Han, L., et al., *Inactivation of PI3K/AKT signaling inhibits glioma cell growth through modulation of beta-catenin-mediated transcription.* Brain Res, 2010. **1366**: p. 9-17.

64.   in *LiverTox: Clinical and Research Information on Drug-Induced Liver Injury*. 2012: Bethesda (MD).

65.   Nierengarten, M.B., *FDA approves capivasertib with fulvestrant for breast cancer.* Cancer, 2024. **130**(6): p. 835-836.

66. Andrikopoulou, A., et al., *"The emerging role of capivasertib in breast cancer".* Breast, 2022. **63**: p. 157-167.

67. Hirai, H., et al., *MK-2206, an allosteric Akt inhibitor, enhances antitumor efficacy by standard chemotherapeutic agents or molecular targeted drugs in vitro and in vivo.* Mol Cancer Ther, 2010. **9**(7): p. 1956-67.

68. Maynard, J., et al., *2-Deoxy-2-[18F]fluoro-D-glucose positron emission tomography demonstrates target inhibition with the potential to predict anti-tumour activity following treatment with the AKT inhibitor AZD5363.* Mol Imaging Biol, 2013. **15**(4): p. 476-85.

69. Davies, B.R., et al., *Preclinical pharmacology of AZD5363, an inhibitor of AKT: pharmacodynamics, antitumor activity, and correlation of monotherapy activity with genetic background.* Mol Cancer Ther, 2012. **11**(4): p. 873-87.

70. Lang, L., et al., *Circumventing AKT-Associated Radioresistance in Oral Cancer by Novel Nanoparticle-Encapsulated Capivasertib.* Cells, 2020. **9**(3).

71. Robertson, J.F.R., et al., *Proliferation and AKT Activity Biomarker Analyses after Capivasertib (AZD5363) Treatment of Patients with ER(+) Invasive Breast Cancer (STAKT).* Clin Cancer Res, 2020. **26**(7): p. 1574-1585.

72. Sobsey, C.A., et al., *mTORC1-Driven Protein Translation Correlates with Clinical Benefit of Capivasertib within a Genetically Preselected Cohort of PIK3CA-Altered Tumors.* Cancer Res Commun, 2024. **4**(8): p. 2058-2074.

73. Mejia-Rodriguez, R., et al., *Combined treatments with AZD5363, AZD8542, curcumin or resveratrol induce death of human glioblastoma cells by suppressing the PI3K/AKT and SHH signaling pathways.* Biochem Biophys Rep, 2023. **33**: p. 101430.

74. Cui, X., et al., *Blockage of EGFR/AKT and mevalonate pathways synergize the antitumor effect of temozolomide by reprogramming energy metabolism in glioblastoma.* Cancer Commun (Lond), 2023. **43**(12): p. 1326-1353.

75. Haddad, A.F., et al., *Mouse models of glioblastoma for the evaluation of novel therapeutic strategies.* Neurooncol Adv, 2021. **3**(1): p. vdab100.

76. Brat, D.J., et al., *Pseudopalisades in glioblastoma are hypoxic, express extracellular matrix proteases, and are formed by an actively migrating cell population.* Cancer Res, 2004. **64**(3): p. 920-7.

77. Qin, X., et al., *Peri-tumoral brain edema associated with glioblastoma correlates with tumor recurrence.* J Cancer, 2021. **12**(7): p. 2073-2082.

FIGURE LEGENDS

Figure 1: Akt and β-catenin inhibitors reduce CT-2A cell growth and proliferation.

(**A**) Schematic of PI3/AKT and Wnt/β-catenin pathway regulation of GBM growth. (**B, C**) MK-2206 and iCRT3 dose-dependent changes in CT-2A cell growth and viability, respectively; (n=5) after 48h treatment. (**D**) Caspase 3/7 activity following 6 h treatment of 10 μM MK-2206 or 50 μM iCRT3 (n= 5). (**E**) Flow cytometric analysis of CT-2A cells co-stained with propidium iodide (PI) and Annexin V following 24 h of vehicle, MK-2206, or iCRT3 treatment (n=3). Data represent mean ± SD from three or more replicates. Statistical significance is determined by one-way ANOVA with Tukey's HSD test; where * represents $p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 2: Akt and β-catenin inhibitors suppress transcriptional activity of β-catenin in CT-2A cells.

(**A**) Transcriptional activity of β-catenin by luciferase reporter assay using Akt inhibitor MK (5μM) and β-catenin inhibitor iCRT3 (25μM) following 48h treatment (n=9) of CT-2A cells. (**B**) Immunoblotting of cell lysates from cells treated with MK-2206(10 μM) and iCRT3 treatment (50 μM) for 1h (n=3) compared to vehicle control. Densitometric analysis of Western blot bands using ImageJ. (**C**) qRT-PCR analysis of *Myc, Ccnd1,* and *Axin2* mRNA levels following 36-hour treatment with MK (10 μM) and iCRT3 (50μM) (n=3). Data represent mean ± SD from three or more replicates. Statistical significance is determined by one-way ANOVA with Tukey's HSD test; where * represents $p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 3: Akt and β-catenin inhibitors reduce growth & proliferation of N08-30 human GBM cells.

(A, B) MK-2206 and iCRT3 dose-dependent changes in N08-30 cell growth and viability, respectively; (n=5) after 48h treatment. **(C)** Caspase 3/7 activity following 6 h treatment of 10 µM MK-2206 or 50 µM iCRT3 (n= 5). **(D)** Flow cytometric analysis of N08-30 cells co-stained with propidium iodide (PI) and Annexin V following 24 h of vehicle, MK-2206, or iCRT3 treatment (n=3). **(E)** Transcriptional activity of β-catenin analysis by both Akt inhibitor MK (5µM) and β-catenin inhibitor iCRT3 (25µM) following 48h treatment (n=3) of N08-30 cells. Data represent mean ± SD from three or more replicates. Statistical significance is determined by one-way ANOVA with Tukey's HSD test; where * represents $p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 4: Akt siRNAs suppress mouse CT-2A mouse glioma cell growth and Wnt/β-catenin signaling.

(**A**) Level of Akt mRNA isoform expression in CT-2A cells. (**B**) Cell growth and viability assay following 48h treatment by Akt siRNA (n= 3) in CT-2A cells. (**C**) Transcriptional activity of β-catenin using Akt siRNA following 48h treatment (n=3) in CT-2A cells. (**D**) qRT-PCR analysis of Myc, Ccnd1, and Axin2 mRNA levels following 48 h treatment with AKT siRNA (n=3). The results shown above represent three or more replicates. Data represent mean ± SD from three or more replicates. Statistical significance is determined by one-way ANOVA with Tukey's HSD test; where * represents $p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 5: Akt inhibition suppressed tumor progression significantly.

(A) Schematic of *in vivo* study. **(B).** Effect of intraperitoneal administration of MK-2206 on mice body weight (vehicle, n=9 and MK-2206, n=11). **(C&D)** Tumor progression analysis in vehicle (n=9) vs MK-2206 (n=11) treated group. **(E)** Representative H&E staining with tumor bulk from

vehicle and treatment group. Red arrow indicates the midline, and discontinuous black boarder indicates hyper nucleated area. Data represent mean ± SD or SEM from three or more replicates. Statistical significance is determined by Student's t-test or by two-way repeated measures ANOVA; where * represents p <0.05, **p <0.01, ***p <0.001.

Figure 6: MK-2206 treatment suppressed the p-AKT and p-GSK-3β expression in tumor tissues.

p-AKT-ser473 (**6A**) and p-GSK3-β-ser9 (**6B**) expression in MK-2206 treated mice compared to vehicle group (n= 5 mice per group). Data represent mean ± SD from three or more replicates. Statistical significance is determined by Student's t-test; where * represents p <0.05, **p <0.01, ***p <0.001.

FIGURES

Figure 1

Figure 2:

Figure 3:

Figure 4:

Figure 5:

Figure 6: