Emergence and stabilisation of a neo-Y chromosome in nematode species with rare males

Brice Letcher[1], Lewis Stevens[2], Nathanaelle Saclier[3], Andrew Hsiao[1], Manuela Kieninger[2], Caroline Launay[1], Eva Wenger[1], Mark Blaxter[2], Marie Delattre[1]

　　1: LBMC, ENS de Lyon and CNRS, Lyon, France
　　2: Tree of Life, Wellcome Sanger Institute, Cambridge, United Kingdom
　　3: ISEM, Université Montpellier, France

Abstract

How do sex chromosomes and modes of reproduction evolve? In nematodes, many modes of reproduction exist ('regular' sexuality, hermaphroditism, parthenogenesis), but most species have a XX/X0 system: females have two X chromosomes and males have one. In rare cases, Y chromosomes have emerged in males, but without associated changes in reproductive mode. Here we demonstrate that, in *Mesorhabditis* nematodes, a new Y chromosome co-emerged with an unconventional mode of reproduction called autopseudogamy. In this system, females produce clonal females plus rare (~10%) males, needed for fertilisation only. These males do not contribute to the female genome and may thus be 'genetically useless' - except for their Y. Analysing the Y chromosomes of two autopseudogamous species, we found two conserved features: i) they accumulated male-beneficial genes, and ii) they display a strong fertilisation drive, in that mainly Y-bearing sperm fertilise female oocytes. Both features are, evolutionarily speaking, favourable in the context of autopseudogamy, making the Y chromosome a possible stabilisor of this mode of reproduction.

Teaser

A Y chromosome stabilises a reproductive system in which males are rare and do not contribute to the genetic makeup of females.

Introduction

Many species of multicellular eukaryotes rely on genetic sex determination, using a set of genes that trigger a sex determination cascade to produce males and females (and/or hermaphrodites). These 'master switch' genes can be located on a sex-limited chromosome, such as the Y of males in mammals (XX/XY system) or the W of females in birds (ZZ/ZW), but they can also be located on the X or Z chromosomes or on autosomes (1,2). Notably, some species have a XX/X0 system in which males have just a single X chromosome and sex is determined by cellular processes measuring the X-to-autosome ratio (3,4). This system is found in the model nematode *Caenorhabditis elegans* and thought to be ancestral to all nematodes (5).

In XX/X0 species, new Y chromosomes can arise *de facto* through simple X-autosome fusions (Fig. 1A). Initially identical to the fused autosome, the unfused autosome can gradually lose recombination and diverge in males, becoming a 'true' Y chromosome (Fig. 1A). So far, this type of neo-Y chromosome emergence has been found in two clades, filarial parasites (6,7) and within the *Pristionchus* genus (family Rhabditidae) (8). However, it does

not appear associated with changes in sex-determination (remaining X-to-autosome ratio based) or in reproductive mode, so that it is unclear if and how these neo-Ys will persist. Indeed, evolutionary reconstructions of ancestral chromosomal states in nematodes have inferred frequent X-autosome fusions (5), but most extant species are XX/X0, not XX/XY, suggesting neo-Ys may arise but also be lost relatively frequently.

Here, we study sex chromosome evolution in the genus *Mesorhabditis* (also in family Rhabditidae) in which a peculiar mode of reproduction called autopseudogamy has evolved, from gonochoristic (sexual) ancestors. Intriguingly, we had previously found that the autopseudogamous species *M. belari* is XX/XY (9). In autopseudogamous species of *Mesorhabditis*, males are rare in the population (2-14% of all individuals, depending on the species (10)) and produced by the 'regular' (sexual) mixing of paternal and maternal genomes. By contrast, females are produced clonally, using the maternal genome only. In both cases, oocytes need to be fertilised by sperm; however, in ~90% of fertilised oocytes, the sperm DNA is not used, and a modified meiotic programme gives rise to clonal females (11). We illustrate autopseudogamy in Fig. 1B.

Fig. 1. Introductory concepts: X-autosome fusions and autopseudogamy. **A) Neo-Y emergence by X-autosome fusion**. A female diploid for the X chromosome (blue) and one pair of autosomes (orange) is shown on the left, and a male diploid for the autosome pair and haploid for the X is shown on the right. In this XX/X0 system, if an autosome fuses to the X, individuals diploid for the fusion become females, while individuals haploid for the fusion become males. In the latter case, the unfused autosome may persist in males, and keep recombining with its fused counterpart. The unfused autosome thus forms a neo-Y chromosome *de facto*. If recombination gradually ceases for this pair, the unfused autosome diverged specifically in males and becomes a 'true' Y chromosome (bottom-right cell). **B) Autopseudogamy**. In this mode of reproduction, found in several species of *Mesorhabditis*, females produce two types of embryos (each female produces both types): in gynogenetic embryos (left-hand cell), oocytes do not use the sperm DNA (green pronucleus), and modify their meiotic divisions to remain diploid (2n inside the yellow female pronucleus). This gives rise to clonal females. By contrast, in amphimictic embryos, oocytes do use the sperm DNA (right-hand cell). In this case, only males are produced (see main text for why).

In this system - also called gynogenesis - the DNA of males is thus never transmitted to females. Conversely, the male genome rapidly converges to the female genome, and rare males may only be produced by females to trigger the production of new clonal females (9). We hypothesised that in such a system, a Y chromosome may be favoured, as it is the only DNA not diluted in the female genome. To test this, we reconstructed the sex chromosomes of two gonochoristic ('regular' sexual) and two autopseudogamous (hereafter, for simplicity, *pseudosexual*) species spanning the phylogenetic diversity of *Mesorhabditis* (including *M. belari*, which we reassembled for this study) and analysed their evolutionary history. We focussed in particular on the putative functions, if any, encoded by the Y chromosome.

Results

Sexual species are XX/X0 and pseudosexual species are XX/XY in *Mesorhabditis*

We have previously established that pseudosexuality arose a single time in *Mesorhabditis* (10) and that one pseudosexual species, *M. belari*, has male-specific genome regions clearly consistent with a Y chromosome (9). To establish the origin of the Y chromosome, and whether it correlates with the emergence of pseudosexuality, we selected four species spanning phylogenetic diversity of *Mesorhabditis*: two sexual species, *M. spiculigera*, and *M. longespiculosa*, and two pseudosexual species, *M. belari* and *M. monhystera* (Fig. 2A).

Using long-read DNA-Seq data plus Hi-C for scaffolding, we assembled each species' genome to high contiguity (N50 values of 514kbp-1.8Mbp, for genome sizes 140-240Mbp) and completeness (BUSCO completeness values of 81-84%). We note that our assemblies are not chromosome-level, however, in large part due to unbridged long repeat arrays (see Methods for details).

To test for the presence of sex chromosomes, we generated short-read DNA-Seq data from separated pools of males and females that we mapped to predicted genes in the genomes (focussing on genes as they are easier to map to, due to lower repeat content; see Methods for details). After normalising for sequencing depth, we compared male and female coverages: genes with twice the coverage in females should be on the X, and coverage only in males should be on the Y. Genes with equal coverage in both should be on autosomes (or on a portion of the Y that has not yet diverged from the X - we return to this later). This approach clearly demonstrated that the two sexual *Mesorhabditis* species are XX/X0 - as are most other species in the *Rhabditidae* family (5,12) - while the two pseudosexual species are XX/XY (Fig. 2B). Thus, at the phylogenetic level, a Y chromosome co-emerged with pseudosexuality in *Mesorhabditis* (Fig. 2A, yellow star).

Fig. 2. Co-emergence of a neo-Y chromosome and pseudosexual reproduction in *Mesorhabditis*. **A)** Phylogeny of *Mesorhabditis* species, built from conserved protein-coding genes found across all species (see Methods). Numbers at the nodes indicate the fraction of trees supporting each node, out of 1000 bootstrapped trees. Coloured boxes show mode of reproduction (as identified in the lab (10)), with autopseudogamous species shaded in orange and sexual species in blue. These clearly support a single emergence of autopseudogamy (that we refer to as pseudosexuality), marked with an orange star on the phylogeny. The four species in bold are those for which we identified the sex chromosomes using assembled genomes and male/female DNA-Seq: the data are consistent with a single emergence of a neo-Y chromosome on the branch indicated by the orange star. **B)** Identification of sex chromosomes using male/female DNA-Seq. Sequencing reads

were aligned to predicted genes in the genomes of the four focal species (*M. belari*, *M. monhystera*, *M. longespiculosa*, *M. spiculigera*): genes with the same count in males and females are autosomal (red curve), twice the coverage in females are on the X (green curve), and coverage in males but no coverage in females are on the Y (blue curve). Genes consistent with a Y chromosome are only seen in the two pseudosexual species (*M. belari* and *M. monhystera*).

Evolution of the X chromosome in *Mesorhabditis*: turnover and fusions

To understand how the sex chromosomes of *Mesorhabditis* evolved, we assigned entire assembled scaffolds to X, Y or autosomal status, by comparing male/female coverage ratios along entire scaffolds. As genes are easier to map to, we also used male/female coverage ratios along genes to check for consistency. As an example, a X-assigned scaffold for *M. monhystera* is shown in Fig. 3: almost all genes are clearly X-assigned (Fig. 3A), and the entire scaffold has coverage consistent with an X chromosome (Fig. 3B). Almost all large assembled scaffolds (>100kbp) could be confidently assigned in this way, and when not, visual inspection was used to resolve the scaffold's status (see Methods). For the two pseudosexuals, we obtained two X-assigned scaffolds, while for the sexuals, we obtained 6 (*M. spiculigera*) and 20 (*M. longespiculosa*). The latter are more fragmented overall, likely due to higher heterozygosity in sexual species, and due to lack of Hi-C data for assembly specifically for *M. longespiculosa* (see Methods).

To infer the evolutionary origin of the sex chromosomes in our genome assemblies, we used Nigon elements. Nigon elements are the inferred ancestral chromosomes of nematodes in order Rhabditina (to which *Mesorhabditis* belongs), defined based on the observation that groups of genetic markers have mostly remained on the same chromosomes through evolutionary time. Previous work defined seven Nigon elements (A, B, C, D, E, N, X), with X being the ancestrally X-linked element (5,13), and assigned highly-conserved genes to each element (5). We thus 'painted' each of our confidently-assigned scaffolds by the Nigon-assigned genes found on it (if any). As an example, our X-assigned scaffold of *M. monhystera* is mostly composed of Nigon B genes (Fig. 3C).

A) Gene coverage-based assignments

B) Scaffold-wide coverages

C) Nigon hits

Fig. 3. **Example X-assigned scaffold in *M. monhystera* and its Nigon painting. A)** Ratio of female to male DNA-Seq counts for genes on this specific scaffold: most genes have a ratio consistent with an X chromosome (blue dots). **B)** Normalised female and male DNA-Seq coverage along the entire scaffold (not just genes). Coverage was normalised so that diploid coverage should have a value of 1 (y-axis). Females have diploid coverage, and males haploid coverage, on this scaffold, again consistent with an X chromosome. **C)** Nigon painting of this X-assigned scaffold. Almost all Nigon elements along this scaffold are Nigon B assigned (orange dots). Note the y-axis values are meaningless here and used only for spreading out the points.

We initially expected to see many Nigon X hits across our X-assigned scaffolds, as its component genes were previously found on the X chromosomes of 15 nematode genomes (5); however, no *Mesorhabditis* genome was used in this definition set. Instead, we found that Nigon X elements were majoritarily on autosomes in all four *Mesorhabditis* species (68% for *M. spiculigera* and >90% for the other three *Mesorhabditis* species; Fig 4A and Fig. S1). Conversely, Nigon X elements were only partly present on the X chromosome of *M. spiculigera* (32%), and were near-absent from the X chromosome of the three other *Mesorhabditis* species (<3%).

Instead, we found the X chromosomes of all four species is majoritarily (54-70%) composed of Nigon C elements (blue, Fig. 4A). Assuming the ancestral X chromosome was composed of Nigon X (based on (5)), this clearly suggests the ancestral X chromosome of *Mesorhabditis* fully transitioned away into Nigon C, likely following a fusion event - except in the lineage leading to *M. spiculigera* (Fig. 4B).

Fig. 4. X chromosome turnover and fusions with autosomes in *Mesorhabditis*. A) For each scaffold assignment class (autosome, X and unclear, x-axis; see Methods for assignment criteria), the relative fraction of each Nigon element is shown as coloured bars (y-axis). Each panel shows one of the four focal species; numbers above coloured bars show the total number of Nigon elements per assignment and species. All X-assigned scaffolds have a significant fraction of Nigon C elements (blue), and the two pseudosexual species uniquely have Nigon C elements on their X chromosomes. **B)** The same tree as in Fig. 2 is shown, with a yellow star marking the origin of pseudosexuality. To the right of the tree leaves, putative X chromosomes are shown for our four focal species: each horizontal coloured bar represents one X-assigned scaffold, coloured by the relative distribution of Nigon elements on this scaffold, and black horizontal bars represent unbridged repeat arrays between scaffolds, preventing full assembly (see main text for details). To the left, at the tree nodes, the inferred ancestral state of the X chromosome at different points in time is shown. We infer an ancestral Nigon C fusion to the X at the base of all *Mesorhabditis*, as all four of our species have

Nigon C painted X-assigned scaffolds. Fusion of Nigon E (green) and Nigon N (light purple) likely occurred in the lineage leading to *M. spiculigera*, and fusion of a Nigon B autosome (orange) occurred in the ancestor of all pseudosexuals. The Nigon B fusion co-occurs with the emergence of pseudosexuality.

For the two pseudosexuals, two X-assigned scaffolds were assembled in total, each composed of Nigon B (orange) and Nigon C (blue). We thus hypothesised that, in the ancestor of pseudosexuals, part of an autosome (Nigon B) fused to the X chromosome (Nigon C; Fig. 4B). As we did not assemble a single X-assigned scaffold, however, we looked for evidence of physical linkage. We found such evidence using our Hi-C data, but it was relatively weak, especially for *M. belari* (see Methods and Tables S1/S2). We thus experimentally tested and confirmed linkage of the two X-assigned scaffolds in *M. belari*, by genotyping animals at two pairs of heterozygous loci (see Methods for details and Fig. S2).

Only part of Nigon B fused to the X chromosome in pseudosexuals, as the majority of Nigon B elements remain on autosomes in both *M. belari* and *M. monhystera* (Fig. S1). A single fusion event likely occurred in the common ancestor of both species, as the Nigon B elements on the fused autosome were conserved and co-linear between *M. belari* and *M. monhystera* (Fig. S3). We also found two additional fusions involving the X chromosome, both in the sexual species *M. spiculigera*: Nigon E (green) and Nigon N (light purple) (Fig. 4B and Fig. S4). For *M. longespiculosa*, all X-assigned scaffolds were Nigon C (blue).

Overall, the X chromosome has thus undergone fusions with various autosomes in different *Mesorhabditis* species, including a full replacement of the ancestral X chromosome by autosomal elements before the emergence of autopseudogamy.

Emergence of the neo-Y chromosome in pseudosexuals

In an XX/X0 sex-determining system, X-autosome fusions can give rise to neo-Y chromosomes in males (Fig. 1A). While initially identical to the fused copy, if the unfused copy stops recombining in males it diverges from its autosomal counterpart, eventually showing haploid coverage in males.

In *M. belari*, the X-fused, Nigon B scaffold (orange in Fig. 4B) showed a pattern of male/female coverages exactly consistent with this scenario (Fig. 5A, upper panel). The right-hand part of this scaffold had the expected pattern for an X chromosome, with diploid coverage in females and haploid coverage in males, but the left-hand part was diploid in both males and females. This is compatible with the scenario shown in Fig. 1A, in which a neo-Y arises by fusion followed by divergence: here, one part of the Nigon B fusion still aligns with the X of females, making it diploid in males, while the remaining part would have diverged, making it haploid in males. To further test this scenario, we mapped Y-assigned scaffolds to the rest of the genome, and found that the best hits occurred at the junction of the diploid-haploid transition zone on this X scaffold (lower panel of Fig. 5A).

These data are thus consistent with the neo-Y emerging following the Nigon B autosome fusion with the *Mesorhabditis* ancestral X chromosome (Fig. 5B). In *M. belari*, most of the neo-Y scaffolds did not map to the rest of the genome, showing it has substantially diverged in sequence from its fused counterpart in males - with the exception of the diploid region in males. Interestingly, in *M. monhystera*, no such diploid region was found, with the

corresponding 'B' fusion being entirely haploid in males (Fig. 2A). This suggests the neo-Y of *M. monhystera* has fully diverged from its unfused counterpart, and possibly no longer recombines with the X.

Fig. 5. Neo-Y emergence following X-to-autosome fusion in the pseudosexuals. A) One of the two X-assigned scaffolds of *M. belari* (the one that fused to the ancestral X - see Fig. 4B) shows a mixture of autosome-like coverage (left-hand portion of the upper panel: females and males both have a coverage of 1, indicating diploid coverage) and X-like coverage (rest of the coverage plot, with diploid coverage in females and haploid coverage in males). To test whether the neo-Y chromosome of *M. belari* is related to this diploid region in males, we aligned Y-assigned scaffolds to the rest of the genome: the best hits genome-wide were found on this scaffold, and are displayed in the lower panel. The y-axis shows the percent identity of each alignment hit, the width of the bar represents how long the hit is, and the hit is coloured by which Y-assigned scaffold it was found in. The hist cluster in the transition zone between diploid and haploid coverage in males, consistent with the neo-Y originating from this entire scaffold. **B)** Evolutionary scenario for neo-Y origin in pseudosexuals. After the fusion of an ancestral autosome (orange) to the ancestral X (blue) in the lineage leading to pseudosexuals,

the unfused autosome (orange) only ever occurs in males (assuming males remain defined by being haploid for the ancestral X, in blue). This neo-Y then diverged away from its autosomal ancestor: both are depicted as black rectangles, but they are of different sizes and do not share any large blocks of sequence homology (see Methods). Only in *M. belari* is a diploid region still found in males (shown as an orange portion on the neo-Y), suggesting recombination may still be occurring in this region.

We further tested this scenario using DNA stainings of meiotic cells in male gonads of the two pseudosexual species. In *M. monhystera*, we found two lagging chromosomes during metaphase and anaphase of meiosis I, suggesting they are not paired (Fig. S5). By contrast, in *M. belari* for which we expect proper pairing of sex chromosomes through the small diploid region (Fig. 5A), we did not observe any lagging chromosomes (Fig. S5). Lagging chromosomes were also absent in gonads of *M. monhystera* females, where the two X chromosomes are expected to pair (Fig, S5). This strongly suggests the unpaired chromosomes in *M. monhystera* males are the X and the Y.

Characterisation of the neo-Y chromosomes

Overall neo-Y degeneration
We could confidently assign 20 scaffolds spanning 7.8Mbp to the Y chromosome of *M. belari*, and 152 scaffolds spanning 17Mbp to the Y chromosome of *M. monhystera*. Despite the large size of the neo-Ys, we found very few predicted genes with RNA-Seq expression: 155 for *M. belari* and 319 for *M. monhystera*. In both species, gene density (number of genes per kbp) was ~8-10 fold lower on the Y than on X and autosomes. In addition, genes on the Y had more copies (=more paralogs), and were part of fewer Y-specific families, compared to genes on the X or autosomes (Tables S3 and S4). The Y chromosomes were also highly enriched for repeated elements compared to the X and autosomes (Table S5), and had a higher fraction of genes with no RNA-Seq support (see Methods). Altogether this is consistent with pseudogenisations, gene duplications and accumulation of repeated sequences on the neo-Y of pseudosexuals, as often seen during the evolution of Y chromosomes (14).

No evidence for changes in genetic sex determination in the pseudosexuals
In the XX/X0 nematode *C. elegans*, specific genes serve to 'count' the ratio between X chromosomes and autosomes, leading to the specification of males or hermaphrodites. In *C. elegans*, these are *sex-1* and *fox-1* located on the X chromosome, and *sea-1* and *sea-2* located on autosomes (15). Together these control the dose of the 'master' sex regulator gene *xol-1* (15). To check for any changes in sex determination pathways, we looked for these genes in our four focal *Mesorhabditis*. We could find orthologs for *sex-1*, *fox-1* and *sea-2* in all four species, and no orthologs in any of the four species for *sea-1* and *xol-1* (see Methods). Surprisingly, all three orthologs - including the X-linked *sex-1* and *fox-1* in *C. elegans* - were located on autosomes in our four species. As we found no differences in gene presence/absence or localisation between the sexual and pseudosexual species, we hypothesise that the pseudosexual species still rely on X:autosome dosage for sex determination and not their Y chromosome.

Conserved male-beneficial gene families on the neo-Y
We next characterised the genes on the neo-Y chromosomes, in particular looking for genes that could explain its maintenance through an extended evolutionary period. Based on a recent time-calibrated phylogeny of the clade Rhabditina (16), to which *Mesorhabditis*

belongs, we estimate the neo-Y emerged ~45 million years ago (see Methods). We thus hypothesised that, in the absence of factors ensuring its maintenance, the neo-Y could easily have been lost.

We first looked for any orthologous families with members on the Y chromosomes of both *M. belari* and *M. monhystera*, as such genes could have maintained the neo-Y prior to the split of the two species. Using whole sequence-based homology (with OrthoFinder (17)), we found just one such family that we called 'hen-1-like', as members are orthologous to the *C. elegans* gene *hen-1*. In *C. elegans hen-1* codes for an LDL receptor domain-containing secretory protein involved in sensory perception (18), and the members in *M. belari* and *M. monhystera* were similarly annotated as 'lipoprotein receptors' (using eggNOG (19)). In *M. belari*, the family had two members on autosomes and 15 on the Y, and in *M. monhystera*, one member on an autosome and 4 on the Y, consistent with a Y-specific expansion. In both species, the Y-specific copies bear specific amino acid substitutions compared to the autosomal copies that are highly conserved across the Y copies. The substitutions were specific to each species, however, suggesting the duplications on the Y may have occurred independently in the two species (i.e., after the split of *M. belari* and *M. monhystera*; Fig. S6).

Because the hen-1-like genes could be cell surface-exposed and/or secreted, we wondered if they were involved in oocyte-sperm interactions. However, comparing RNA-Seq data of gonads with whole adults in *M. belari*, we found that most of the expression was not in gonads (estimated 1-5% of the total expression in gonads), consistent with a function in somatic tissues, maybe for sensory perception as found in *C. elegans*. We thus hypothesise the hen-1-like genes on the Y chromosome encode male-beneficial functions, but are not involved in a fertilisation drive, as we expect such genes to be transcribed in the germline. We note that the autosomal copies already have a male expression bias, of 16% in *M. belari* to 600% in *M. monhystera*.

As just one family of orthologs was common to the neo-Y chromosomes, we expanded our search to shared functional domains. We found only a few shared functional terms, of which 'lipoprotein receptor' - corresponding to the hen-1-like family - and 'promoting spermatogenesis'. For the latter, we found two main gene families, one in each pseudosexual species, encoding zinc-binding transcription factors with homology to the *C. elegans* gene *tra-1*, a master regulator of germline and somatic sex differentiation (20,21). The homology to *tra-1* was particularly clear for the *M. monhystera* family (copies aligned well to *C. elegans tra-1*), and more distant for the *M. belari* family (only domain-level homology was found) (Fig. S7). In both families, 8-9 copies were found on the neo-Y chromosome, and 1-4 copies were found on the X chromosome, and the Y copies had species-specific substitutions, again consistent with male-specific evolution on the neo-Y (Fig. S7). Here also, the non-Y copies (on the X chromosome) had a strong male expression bias, of 278% in *M. monhystera* and with almost no female expression in *M. belari*.

While *tra-1* was initially identified as a strongly feminising gene (notably, XX mutants for *tra-1* become fertile males (22)), other work has in fact shown that it also plays a role in spermatogenesis (20,21) as well as male behaviour, including exploration, through expression in specific neurons (23). We thus similarly hypothesise that the neo-Y tra-1-like copies encode male-beneficial functions.

Convergent fertilisation drives on the neo-Ys
In *M. belari*, Y-bearing sperm have a strong fertilisation drive: we previously found (using DNA-FISH with a Y-specific probe) that ~87% of fertilised oocytes are fertilised by Y-bearing sperm (9). We tested whether *M. monhystera* also has a fertilisation drive, using mounted one-cell embryos for DIC live imaging. Of 11 amphimictic embryos (where the sperm DNA is used; Fig. 1B) that developed into adults, 11 were male. At a p-value threshold of 0.05, this is incompatible with a fertilisation drive below 74% (chi-square goodness-of-fit test; see Methods).

To identify candidate drivers, we looked for protein-coding genes with annotations consistent with a role in the fertilisation drive - e.g., cell-surface exposed, or with a role in meiosis for segregating proteins specifically to Y-bearing sperm. As only the tra-1-like and hen-1-like families were shared across the Y chromosomes of the two pseudosexuals, these genes were all species-specific. We identified just four families with interesting annotations, but all had closely-related paralogs on the X or autosomes, making them unlikely candidates. We next identified genes occurring exclusively on the Y (i.e. with no paralogs on the X or autosomes): 91 genes in 10 families in *M. belari* and 63 genes in 17 families in *M. monhystera*. Of these, some genes code for putative DNA-binding (helix-loop helix, zinc finger, TATA-box-binding domains), RNA-binding, transmembrane, and structural proteins (vimentin, condensin, kinetochore protein). The list of these loci is provided with this paper (see Methods).

The Y chromosome may not in itself encode a protein directly responsible for the fertilisation drive, however. Notably, we found - by detection of an antibody against the active form of RNA polymerase II - that in male gonads of *M. belari*, spermatocytes are transcriptionally silent from the onset of meiosis I, i.e., prior to segregation of the neo-Y (Fig. S8). This suggests that a factor responsible for the drive may not be expressed specifically in Y-bearing sperm after they have been individualised (though we note immunostaining may not mean that Y-bearing sperm are fully transcriptionally silent). Instead, factors responsible for the Y fertilisation drive could be expressed earlier, and elsewhere (e.g. on the X or autosomes), and co-segregate with the Y during meiosis, perhaps through recognition of a non-coding element (e.g., a repeat array) on the Y.

To test this idea, we looked for repeated arrays of small (between ~3bp and 120bp units) sequences called satellites, as these have been implicated in meiotic drives in other systems (24,25). We found several satellites highly specific to the Y chromosomes of both *M. belari* and *M. monhystera*, as well as a few satellites specific to their X chromosomes, but none of these satellites were in common between the two species (see Methods). We return to this in the Discussion.

Search for mitochondrial genome rearrangements

Finally, because the neo-Y chromosome is the only DNA that is transmitted exclusively through males, we also analysed the other DNA with uniparental transmission: the mitochondrial genome. We specifically wanted to test for genomic changes associated with the pseudosexuals - similarly to the neo-Y emergence. For this we assembled the mitochondrial genomes of all *Mesorhabditis* species shown in Fig. 2A using long-read data (mostly Nanopore; see Methods). The assembled mitochondrial genomes all contain 12

protein coding genes (*atp-8* being absent, as in most other nematodes (26)), and are relatively small, ranging 13-15kbp. We found two large-scale mitochondrial genome translocations along the phylogeny. The first is common to all pseudosexual species except *M. monhystera* and *M. sp. JU4397*, with the gene order of the latter being conserved with *C. elegans* (26). We also found a translocation in one of the sexual species, *M. irregularis*. We concluded that mitogenome translocations are frequent in this genera but are not linked to pseudosexuality (Fig. S9).

Discussion

Using genomics, we reconstituted the evolution of the sex chromosomes in the *Mesorhabditis* genus. Unexpectedly, we found X chromosomes that have entirely lost the features common to the X chromosome of other rhabditid nematodes (Nigon X); this could call for a redefinition of these common features in Rhabditina. Perhaps as a consequence of this evolutionary transition, we found that among well-known genes that count the X:autosome ratio in *C. elegans*, that are present on the X and on autosomes in *C. elegans*, all identified orthologs were found on autosomes in *Mesorhabditis*. Because we found no presence/absence differences between sexual (XX/X0) and pseudosexual (XX/XY) species, we hypothesise the Y chromosome of pseudosexual species does not play a role in sex determination. However, how the X:autosome ratio is counted in *Mesorhabditis* remains to be explored.

Our data suggests a single X-autosome fusion gave rise to a neo-Y chromosome in pseudosexual species of *Mesorhabditis*. Since then, the neo-Y maintained a region of homology with the fused X in *M. belari*, but not in *M. monhystera*. Chromosome fusion has been shown to drastically reduce recombination at the fusion point in *C. elegans* (27): consistent with this, the diploid in region in *M. belari* is distal to the fusion, so that recombination may gradually have been lost from the fusion point. In *M. belari*, this 'pseudoautosomal' region (PAR) could mediate recombination and proper chromosome segregation between the X and Y during male meiosis. In this case, to what extent proper X/Y segregation is achieved in *M. monhystera* in the absence of a PAR remains to be investigated.

While the X chromosome has been especially prone to past fusions with autosomes in Rhabditina (5), most rhabditid nematodes are XX/X0, not XX/XY (12), suggesting any neo-Ys have mostly been lost. Here by contrast we found a neo-Y that has been maintained since its emergence, ~45 millions of years ago. To account for this, we exclude a role for the neo-Y in sex determination (as discussed above), or through haploinsufficiency of the fused autosome in males (where males haploid for genes on the fused/unfused autosome pair are not viable), because there is no longer any diploid region in *M. monhystera*. Instead, maintenance could be explained by the presence of male- or sperm-beneficial genes on the Y chromosome.

In support of this, we found two families of genes on the Y of both pseudosexual species, both of which could encode male-beneficial functions as they are predicted to either promote spermatogenesis or be involved in learning and sensory perception. Evolutionarily, both families underwent duplications on the Y and accumulated Y-specific substitutions. We found these substitutions are species-specific, so that they may have emerged only after the

speciation of *M. belari* and *M. monhystera*. Alternatively, species-specific substitutions could have been fixed following gene conversions between the Y paralogs, a frequent phenomenon on Y chromosomes (28).

We also found that both Y chromosomes are likely involved in a strong oocyte fertilisation drive. We did not find any conserved protein-coding genes or satellite repeats that could underlie these drives. This suggests that either an ancestral neo-Y drive diverged in each species (e.g. through an arms race with the X chromosome), or that these drives emerged convergently. Mechanistically, we do not know whether a factor facilitates fertilisation by Y-bearing sperm or inhibits fertilisation by X-bearing sperm. A factor that poisons oocytes that use the DNA of X-bearing sperm could also be involved, explaining the absence of sexual females (9). Our observed drives could occur through expression of Y or X-bearing sperm following male meiosis (though they appear transcriptionally silent at this stage), or preferential segregation of a cytoplasmic factor to Y or X-bearing sperm during male meiosis. The latter has been observed before: in the nematode genus *Auanema*, proteins involved in sperm motility co-segregate with the X chromosome during male meiosis (29), and in *D. melanogaster*, the well-known driver Stellate poisons Y-bearing sperm, also during male meiosis (30).

In light of our results, we hypothesise that autopseudogamy as a mode of reproduction may favour the emergence and stabilisation of a Y chromosome. Indeed, the main feature of autopseudogamy is the production of a majority of clonal females, through the 'rejection' by the female oocyte of sperm DNA following fertilisation (also called gynogenesis). There are two main consequences to this. The first is that males no longer contribute to the female genome, so that any male-beneficial genes can rapidly disappear - except if they are encoded on a Y chromosome. The second is that a fertilisation drive of Y-bearing sperm ensures that when oocytes use the sperm DNA (=amphimictic embryos), males are mostly produced. If rare X-bearing amphimictic embryos are additionally poisoned, only males will ever be produced by sexual reproduction. In *M. belari*, we indeed never observed mature sexual females by genotyping (9). In turn, preventing the production of sexual females may stabilise gynogenesis. In the future, we can test these hypotheses by checking whether dead embryos are XX amphimictic by genotyping, and testing for a correlation between the strength of fertilisation drive and the rate of amphimictic embryos in *Mesorhabditis*.

More broadly, the relative timing of autopseudogamy and neo-Y chromosome emergence remains unknown. Future denser sampling and comparative genomics in *Mesorhabditis* could resolve this, e.g. by finding sexual species with a proto-Y, or pseudosexual species without a Y. In addition, despite our reasoning and results, we cannot definitively exclude that the Y has no selected function, and was simply never lost over time by random genetic drift. To truly test this, forward (unbiased) or reverse (starting with our candidates) genetic screens for Y genes affecting male fitness, and/or whose sperm exhibits more or less fertilisation bias, will be needed.

Materials and Methods

Nematode culture and sequencing

All strains were maintained at 20°C on enriched plates (Nematode Growth Media supplemented with 7 g/L agarose, 5 g/L bacto-peptone and seeded with *E. coli* OP50). We sequenced the following *Mesorhabditis* strains: JU2817 (*M. belari*), JU2855 (*M. monhystera*), DF5017 (*M. longespiculosa*), AF72 (*M. spiculigera*), JU2848 (*M. littoralis*), JU2864 (*M. simplex*), JU3143 (*M. okuensis*), ANA126 (*M. sp.* ana126), JU3174 (*M. franseni*), JU3347 (*M. vernalis*), JU2902 (*M. bifurcata*), JU3003 (*M. paucipapillata*), JU4397 (*M. sp.* JU4397), SB321 (*M. irregularis*).

To produce worm pellets, worms were synchronised by bleaching as previously described in (9) and then grown as above. When plates were nearly starved, worms were harvested and washed in M9 + 0,01% Tween20. Worm pellets were stored at -20°C.

DNA Sequencing: *M. belari* and *M. spiculigera*

We performed high molecular weight DNA extraction of a 40 mg frozen pellet of *M. belari* using the Monarch® HMW DNA Extraction Kit for Tissue (NEB) with the following modifications. 50 µl of the lysis buffer mix was used for pestling the frozen worm pellet. We added the rest of the lysis buffer and transferred the mix in a 2 ml DNA LoBind® Tube (Eppendorf). The mix was digested at 56 ℃ in the Thermomix C (Eppendorf) at 750 rpm for 2 hrs. After RNA digestion we spun the sample for 5 min at 2500 rcf and transferred the supernatant to a new 1.5 ml DNA LoBind® Tube (Eppendorf) leaving about 30 µl behind. We added 30 µl of lysis buffer to the mix and put the sample for 5 min on ice. We added 300 µl cold Protein Separation solution and inverted the sample at about 20 rpm for 1 to 2 min. We put the sample for 10 min on ice. We spun the sample at 16.000 rcf for 25 min at room temperature. We transferred the supernatant to a new 2 ml DNA LoBind® Tube (Eppendorf) and added 2 glass beads. We added 550 µl Isopropanol and inverted the sample several times before we put it in a wheel for 5 min at 7 rpm. From here we followed the recommended protocol steps. In total, we extracted 2.640 ng high molecular weight DNA. 2,500 ng of this DNA was sheared in two rounds to an average size of 21.1 kb with a Megaruptor 3 (setting 28 followed by setting 29 in the second shearing round) (Diagenode). The sheared DNA was SPRI cleaned with 0.81x of ProNex beads (Promega).

For the DNA extraction of *M. spiculigera* we used CTAB digestion followed by Phenol-Chloroform extraction. We used a worm pellet of 50 mg for extraction. We pestled the frozen worm pellet in 300 µl CTAB lysis buffer (2% w/v) and added 10 µl Proteinase K (800 U/ml). After 1 hour digestion in a Thermomixer C (Eppendorf) at 56 ℃, 550 rpm, we added 10 µl RNaseA and left it for another hour in the Thermomixer. After a total of 2 hours we added 1x Phenol/Chloroform/Isoamylalcohol maix (25:24:1) and mixed well with a bore tip. We centrifuged at 14,000 rcf for 10 min at 4 ℃. We took the supernatant and performed two Chloroform washes with 1x Chloroform/Isoamylalcohol (49:1). We centrifuged 14,000 rcf for 10 min at 4 ℃. We then transferred the supernatant to a new 1.5 ml tube and added 0.7x Isopropanol. We inverted the mix and incubated it for 10 min at room temperature. We centrifuged at 14,000 rcf for 10 min at 4 ℃. We washed the pellet twice with 70% Ethanol and centrifuged for 5 min 14,000 at 4 ℃. The pellet was eluted in a volume of 200 µl elution buffer. We then mixed the extracted DNA several times with a bore pipette tip and let the DNA elute overnight at room temperature before proceeding to QC measurements. We extracted 4,500 ng DNA. We sheared the DNA to an average size of 21.4 kb with a Megaruptor 3 (setting 29) (Diagenode). The sheared DNA was SPRI cleaned with 0.81x of ProNex beads (Promega).

The PacBio libraries were prepared by the Sequencing Operations core at the Wellcome Sanger Institute.

Hi-C library preparation and sequencing were performed by the Scientific Operations: Sequencing Operations core at the Wellcome Sanger Institute. A 25 mg pellet of adult-enriched nematodes of *M. belari* strain JU2817 and *M. spiculigera* strain AF72 was processed using the Arima Hi-C version 2 kit following the manufacturer's instructions. An Illumina library was prepared using the NEBNext Ultra II DNA Library Prep Kit and sequenced on one-eighth of a NovaSeq S4 lane using paired-end 150 bp sequencing.

DNA Sequencing: all other *Mesorhabditis* species

For DNA extraction, 100-200 µl of nematode pellets were snap frozen in liquid nitrogen and thawed at 37°C three times for breakage of the cuticle. These were vortexed in 600 µl of Cell Lysis Solution (Qiagen ref. 158906) and 6 µl of proteinase K (17 mg/ml; Sigma-Aldrich), then incubated three hours at 65°C with gentle shaking. After addition of 40 µl of RNAse A (5 mg/ml; Roche), samples were incubated 1 hr at 37°C, then 5 min on ice with 200 µl of Protein Precipitation Solution (Qiagen ref. 158910). After 10 min centrifugation at 13000 rpm, 4°C, the supernatant was precipitated with 600 µl of Isopropanol, incubated 10 minutes at room temperature and centrifuged 30 min at 13000 rpm, 4°C. The pellet was washed twice with 70% ethanol, centrifuged, dried 1h under a hood, then dissolved in 30 µl of water.

For sequencing, *M. littoralis*, *M. simplex*, *M. okuensis* and *M. sp.* ANA126 were processed by IGFL (Lyon, France), with library preparation using Native Barcoding Kit 24 V14 and sequencing on an ONT MinION MK1C machine (R10 flow cell).
M. longespiculosa, *M. franseni*, *M. vernalis*, *M. bifurcata*, *M. paucipapillata*, *M. sp.* JU4397 and *M. irregularis* were also processed by IGFL, with the same library preparation and sequencing on an ONT PromethION machine (R10 flow cell).
The *M. monhystera* sample was processed by BGI (China) for high-fidelity (HiFi) library preparation and sequencing on a PacBio Revio machine. In addition a worm pellet of *M. monhystera* was processed by the MGX platform (Montpellier France) for Hi-C library preparation (Arima-HiC kit) and paired-end sequencing (2x150bp) on an Illumina NovaSeq 6000 machine.

For *M. belari*, *M. monhystera*, *M. longespiculosa* and *M. spiculigera*, we also separately sequenced pools of male and female worms. *M. belari* and *M. spiculigera* samples were processed by the MGX platform with library preparation using Illumina's DNA prep kit and paired-end sequencing (2x250bp) on an Illumina NovaSeq 6000 machine. *M. monhystera* and *M. longespiculosa* were processed by the GenomEast platform (France): *M. longespiculosa* was sequenced on an Illumina HiSeq 4000 (2x100bp) and *M. monhystera* on an Illumina NextSeq 2000 (2x100bp).

RNA Sequencing

For *M. belari* and *M. longespiculosa*, we prepared RNA from entire populations. 100 µl of nematode pellets were vortexed in 500 µl Trizol. The suspension was frozen in liquid nitrogen and transferred to 37°C water bath (4-5 times), then vortexed at room temperature for 30 s, let rest 30 s (5 times). 100 µl of chloroform (24:1 CHCl :isoamyl alcohol) was added, tubes were vigorously shaken for 15 s, incubated at room temperature for 2-3 min, then centrifuged 15 min at 13000 rpm, 4°C. The upper phase was transferred to a new tube with

250 µl of isopropanol. After mixing and incubation at room temperature for 10 min, the tube was centrifuged for 15 min at 13000 rpm, 4°C. The pellet was washed with 75% ethanol, then centrifuged for 5 min at 13000 rpm, 4°C. After removing the supernatant and the pellet was air-dried for 15-20 min, then dissolved with 50-100 µl of RNAse-free water.

For *M. belari*, *M. monhystera, M. longespiculosa* and *M. spiculigera*, we sequenced RNA from single adult males and adult females using the Smart-Seq2 protocol adapted for single worms and described in (31), with the following modifications: 1µl of proteinase K at 17mg/ml, SuperaseIn instead of RNAsin ribonuclease inhibitor from Promega, and an extra step of snap freezing in liquid nitrogen before worm lysis.

All samples were processed by the GenomEast platform, with library preparation using Nextera XT and paired-end sequencing (2x100bp) on an Illumina HiSeq 4000. For *M. monystera* and *M. spiculigera*, the reads from males and females were pooled for gene prediction.

Genome assembly

For three species (*M. belari*, *M. monhystera*, *M. spiculigera*) we had PacBio HiFi and Hi-C data for genome assembly. We first assembled the HiFi reads into contigs representing a single, unphased haplotype with hifiasm v0.19.5 (32) (-s 0.55 --primary). Haplotypic duplications (heterozygous regions collapsed into a single contig) were removed from the primary assembly with purge_dups v1.2.6 (https://github.com/dfguan/purge_dups). We then mapped the Hi-C reads to the purged assembly with bwa v0.7.17 (33) (bwa mem -SP5M) and scaffolded the contigs with yahs v1.2a.2 (34). We then removed any contaminant scaffolds (e.g. bacteria coming from intestines of sequenced worms) with BlobToolKit v4.3.9 (35), based on abnormal coverage of the HiFi data or GC content and aberrant genes characteristic of non-nematode clades.

For the remaining species in the *Mesorhabditis* group (Fig. 2A), we had Nanopore data only. These were assembled with Flye v2.9.2 (36) (-g 150m), with two modes depending on how our Nanopore data was generated: --nano-raw for the MinION data and --asm-coverage 50 --nano-hq for the PromethION P2 data. For *M. longespiculosa*, we further purged and decontaminated the assembly as above.

For all assemblies, quality control was performed using BUSCO v5.8.0 (37) (with the genes in the 'nematoda_odb10' lineage) and Merqury v1.3 (38). For visual inspection of the scaffolding results, Hi-C contact maps were produced and visualised using Juicer (39). For metrics on the finished assemblies, see Table 1 below.

Mesorhabditis phylogeny

After running BUSCO on each of our assembled genomes - plus *C. elegans*, used as an outgroup -, we found 422 genes present in single-copy across all assemblies (out of 3131 BUSCO searched genes). For each gene, we produced a multiple-sequence alignment of all protein-coding sequences with mafft v7.525 (40) and trimmed the alignment with clipkit v2.3.0 (41) to remove aligned positions with too many gaps (-m smart-gap). We concatenated all alignments into a supermatrix with seqkit v2.4.0 (42), from which we built a phylogeny using IQ-TREE v2.3.6 (43), with auto-detection of the best-fitting substitution

model (here, VT+F+R7) and 1000 bootstrap replicates (-B 1000). Trees were visualised and rendered with FigTree (https://github.com/rambaut/figtree/).

Gene prediction

We predicted protein-coding genes in the four focal species of this study (*M. belari*, *M. monhystera*, *M. longespiculosa* and *M. spiculigera*) using the tool BRAKER3 v3.0.7.5 (44). As input for prediction, we gave BRAKER3 a set of known proteins: the 3131 proteins in BUSCO 'nematoda_odb10', the proteins in BUSCO 'metazoa_odb10', and the existing proteomes of *C. elegans* and *M. belari* obtained from WormBase ParaSite release 18 (WBPS18) (45). We also gave BRAKER3 RNA-Seq sequencing data we obtained for each species: population-wide for *M. belari* and *M. longespiculosa*, and of adult males and females for *M. monhystera* and *M. spiculigera*. The RNA-Seq reads were mapped to our assembled genomes using STAR v2.7.10b (46) prior to use with BRAKER3. This resulted in 16717 predicted genes for *M. spiculigera*, 19410 for *M. longespiculosa*, 26653 for *M. belari* and 22542 for *M. monhystera*.

Assigning genes and scaffolds to X, Y, or autosomal status

To assign individual genes a status we used our male and female DNA-Seq data to count the number of sequencing reads coming from each gene with Kallisto v0.50.1 (47). For comparability between males and females, we used the size factor normalisation in DESeq2 (48). We assigned a status to each gene by taking the closest expected regression line of male (m) vs female (f) counts for X, Y or autosomal genes. In other words, we assigned each gene to the status minimising the following squared errors e:

- Autosomal gene: expectation: $f = m$, $e = (f - m)^2$
- X gene: expectation $f = 2m$, $e = (f - 2m)^2$
- Y gene: expectation $f = 0$, squared error $e = f^2$

Genes with low read counts are hard to distinguish this way (Fig. 2B). We thus reassigned genes to a fourth status, 'unclear', if they had fewer than 30 read count support in either males (for X, Y, or autosomal genes) or females (for X or autosomal genes). To ensure confident assignment of Y genes, we also set any Y-assigned gene with >5 read count support in females to unclear.

To assign entire assembled scaffolds to X, Y or autosomal status, we used two approaches. In the first, we reused our assignments of individual genes found along each scaffold: a scaffold was assigned to a given status if >50% of its component genes were assigned to that status (including 'unclear'), and to 'unclear' if no gene assignment group matched this criterion, or the scaffold had no genes.

In the second approach, we used coverages of male and female DNA-Seq reads computed along the entire scaffolds. Reads were mapped to assembled genomes with minimap2 v2.26 (49) (-x sr for short-read preset), duplicates marked with GATK's picard v4.3.0 (50). To ensure coverage was computed only for unique and well-mapped regions of the genome, reads with MAPQ<20 (corresponding to a >=1% probability of the read being mapped to the wrong location, as estimated by minimap2) or marked as 'SECONDARY', 'SUPPLEMENTARY' or 'DUP' were filtered out with samtools v1.17 (51). We computed coverage along 1kbp non-overlapping windows of scaffolds with bedtools v2.31.0 (52), and normalised coverages so that a value of 1 corresponded to the expected coverage on a

diploid chromosome. For this, we divided the value in each window by the median coverage along all 1kbp windows in scaffolds of size >1Mbp, assuming that most of the genome is composed of autosomes. We used large scaffolds only for the median as we observed these mostly had uniform coverage along their entire length. For assignment, we define the median normalised female coverage as n_f and that for males n_m, and the normalised female/male ratio $r = n_f / n_m$. We then used the following ad-hoc rules:

- Autosomal scaffold: $0.6 < n_f < 1.4$ and $0.6 < n_m < 1.4$ and $0.7 < r < 1.3$
- X scaffold: $0.6 < n_f < 1.4$ and $0.6 < n_m < 1.4$ and $1.5 < r < 2.5$
- Y scaffold: $0.05 < n_m$ and $n_f < 0.001$
- Unclear scaffold: neither of the above

Note that for Y assignment, a value close to 0.5 is expected in males, as they should be haploid in males. However, due to their high repeat content, most coverage windows had very low coverage values, as reads did not map confidently to a single location. When this constraint was lifted (i.e., no longer filtering reads by MAPQ), a value of ~0.5 was restored in males: an example is shown in Fig. S10. We filtered including MAPQ, and with a low minimum coverage in males (0.05), to ensure that almost no coverage in uniquely mappable regions was found in females (0.001 threshold above).

We then proceeded to an inspection of these assignments to produce a final classification. If both gene and scaffold coverage-based classification were in agreement, we used this classification (1397 out of 1895 scaffolds). In all remaining cases, either the gene-based or the coverage-based classification were 'unclear'. In most cases we set the scaffold to 'unclear' classification (379 of the 498 disagreeing scaffolds set to unclear), especially when few genes were on that scaffold. We note that many of these 379 scaffolds are in fact eliminated portions of the genome, due to Programmed DNA Elimination in *Mesorhabditis* nematodes (53): their normalised coverages in males and females were typically <0.6 because eliminated portions of chromosomes are only sequenced in germline cells, which form a minority of sequenced individuals. For the remaining scaffolds, we nonetheless assigned eliminated scaffolds if they contained many genes, to ensure accurate Nigon painting-based analyses, and to ensure we captured all genes on the Y chromosomes. All our classifications, and the data supporting them, are publicly available: see data availability section.

Summary of genome assemblies and scaffold assignments

In Table 1 below, we summarise overall assembly and scaffold assignment statistics.

	M. belari	*M. monhystera*	*M. longespiculosa*	*M. spiculigera*
Assembly span (#scaffolds)	241 Mbp (454)	211 Mbp (441)	138 Mbp (725)	147 Mbp (275)
Assembly N50	6.81 Mbp	10.2 Mbp	441 Kbp	10.88 Mbp
%BUSCO genes found (%single-copy, %duplicated)	83.6 (77.8, 5.8)	84.1 (79.9, 4.2)	83.1 (71.8, 11.3)	81.1 (62.2, 16.9)
% autosome-assigned	63.5	70.8	80.9 (111.5 Mbp,	70.7 (103.9

(span, #scaffolds)	(153.2Mbp, 22)	(149.4Mbp, 16)	394)	Mbp, 21)
%X-assigned (span, #scaffolds)	6.3 (15.1Mbp, 2)	8.1 (17.1Mbp, 2)	8.32 (11.5Mbp, 20)	15.4 (22.6Mbp, 6)
%Y-assigned (span, #scaffolds)	3.2 (7.8Mbp, 20)	8.1 (17.0Mbp, 152)	NA	NA
%unclear-assigned (span, #scaffolds)	27.0 (65.3Mbp, 410)	13.0 (27.4Mbp, 271)	10.8 (14.9 Mbp, 311)	13.9 (20.5 Mbp, 248)

Table 1. Summary of assembly statistics and how much of each assembly was assigned to X/Y/autosomal/unclear status, for the four focal species of this study.

We note that the main reason many small scaffolds were of unclear assignment in all four species is because of the process of Programmed DNA Elimination (PDE) that occurs in the genus *Mesorhabditis* (53). During PDE, somatic cells undergo the systematic destruction of parts of their genome during early embryonic development. As mixed-stage, whole individuals were sequenced here, both somatic and germline cells are captured and the genome assembly contains a mixture of eliminated (=present only in germline cells) and retained (=present in both somatic and germline cells) DNA. Most of the unclear-assigned scaffolds are eliminated: they have lower coverage and are highly enriched for long arrays of repetitive DNA, making contiguous assembly challenging. We defer a more detailed explanation of how we classified scaffolds as retained or eliminated to a future study on PDE in *Mesorhabditis*. However, these scaffolds have almost no BUSCO hits, and we manually checked that they did not correspond to X or Y scaffolds so as not to miss any possibly important genes, especially on the Y.

We also predicted the fraction of X, Y and autosomal-assigned scaffolds covered by repeated elements, using EarlGrey v3.1 (54): these data are shown in Table S5.

Nigon painting

To paint our scaffolds with Nigon elements, we used the approach defined in (5), in which a subset of 2298 genes out of the 3131 BUSCO genes that are part of the 'nematoda_odb10' lineage were each assigned a Nigon letter (A, B, C, D, E, N, X). This letter was determined on the basis of gene co-occurrence patterns in chromosome-level assemblies of 12 species in the order *Rhabditina* (5). The correspondence table between BUSCO gene ID and Nigon letter is publicly available here:
https://github.com/tolkit/nemADSQ/blob/main/example/gene2Nigon_busco20200927.tsv.gz.

For visualisation, we used the same colour code for Nigon letters as in (5).

Testing for X chromosome linkage
Testing for X linkage using Hi-C
To test whether the two X-assigned scaffolds of *M. belari* and *M. monhystera* were physically linked, we mapped the Hi-C sequencing data to our assembled genomes using bwa v0.7.17 (bwa mem -SP5M) and extracted contacts using two approaches. The first and most permissive is using pairix v0.3.9 (bam2pairs) (55), as all aligned reads are used to compute contacts, including reads aligning to repetitive parts of the genome. The second is using pairtools v1.1.2 (56), that parses contacts in a more sophisticated manner and allows

filtering aligned reads by MAPQ, a score proportional to the probability that a read is misaligned (essentially due to sequence repetition in the genome). We computed contacts using pairix, and pairtools for reads with minimum MAPQ of 1 and 20.

To test for linkage, we then extracted all contacts overlapping the first or last 50kbp of each assembled scaffold and counted the number of contacts supporting each possible pair of scaffolds. For a given scaffold pair, four possible extremity contacts are possible, based on whether the contacts occur at the start or end (i.e., 5' or 3' end) of each scaffold. For each species, we extracted the top 10 pairs by inter-scaffold count, normalised by how much contact occurs at the extremities themselves (=within-contact count), and looked for whether the two X-assigned scaffolds consistently came on top. This was clearly the case for *M. monhystera*, even after filtering for reads with MAPQ>=20 (Tables S1 and S2).

Testing for X linkage by PCR-based genotyping
To further test for physical linkage in *M. belari*, we used a PCR-based approach. We first identified heterozygous polymorphisms on each of the two X-assigned scaffolds (scaffold 7 and scaffold 16) whose size difference can be easily distinguished on agarose gels. We targeted variants at the 3' end of these scaffolds as based on the Hi-C data, this is the likely position of these scaffolds' linkage. Using IGV (57) we found a heterozygous 182-bp insertion on scaffold 7 (at position 8,594,943, in the intron of gene g23849), and a heterozygous 124-bp insertion on scaffold 16 (at position 5,122,107, in the intron of gene g8032). We designed PCR primers amplifying both the short and long alleles, with expected amplicon sizes of 395bp and 577bp for scaffold 7 (forward primer 5'-TCACCCGTTCTTCCCATTACA-3', reverse primer 5'-TGGCTCGGTAGGGGAAATTT-3') and 437bp and 541bp for scaffold 16 (forward primer 5'-TGTTTCCCATGTAAGTTGAGCA-3', reverse primer 5'-TCTTCTTCAGGACCGATCTCC-3'). We also designed primers to amplify a homozygous and unique region of the neo-Y (on scaffold 31, of size 436bp; forward primer 5'-TGGCTTTTAGATCAATGGAACCT-3', reverse primer 5'-GCAACGGTGATTTTCAAGCA-3'), to confirm genotyped individuals were male. We confirmed using PCR on wild-type individual *M. belari* males and females (strain JU2817) that the X-linked markers were indeed heterozygous in females, and haploid in males with the short or long allele, and that only males were positive for the Y marker (data not shown, see results below).

We then isolated a single adult gravid female on an agar plate (containing nematode growth medium and seeded with *E. coli* strain OP50), and lysed 27 individual male descendents of this initial female (mostly second-generation males, as a given female only produces ~5 males across her lifespan), plus one second-generation female as a control (lysis: 10µl worm lysis buffer + 1% proteinase K for each worm). We genotyped all these individuals for the two X and one neo-Y markers by PCR.

The results are shown in Fig. S2. The female amplified both X markers at the expected size and not the Y marker, while males amplified the Y marker, and one of the X markers, at the expected sizes. A total of 6 males were discarded for analysis as at least one of the X loci did not amplify strongly, leaving 21 males. Of these, and denoting 'S_7L_{16}' to mean 'short allele for scaffold 7 and long allele for scaffold 16', we obtained the following marker combination counts: 10 S_7L_{16}, 11 L_7S_{16}, 0 S_7S_{16}, 0 L_7L_{16}.

Under the null hypothesis that these two markers are on different chromosomes, all four combinations should occur at the ~25% frequency due to Mendelian segregation at meiosis. By contrast if the markers are on the same chromosome, two of these combinations should be highly over-represented among males due to genetic linkage, and the other two combinations arise only after maternal X chromosome recombination exactly between the two markers. Under the null, the probability of observing data at least as extreme as ours (p-value) is 10e-04 (goodness-of-fit test on a one-dimensional contingency table, df = 3). We thus concluded that these two markers are on the same chromosome, and that a single X chromosome occurs in germline cells of *M. belari*. We note that for 4 of the 6 discarded males, the faint bands also supported the S_7L_{16} or L_7S_{16} combinations as found above; one male had two bands for the scaffold 7 locus; and the remaining male did not amplify any locus (Fig. S2).

Assessing homology between neo-Ys and the rest of the genome

All Y-assigned scaffolds were extracted and aligned to the rest of the genome using minimap2 v2.26 (49) (`-x map-ont`). The resulting alignment files (PAF format) were processed in R and alignment blocks filtered for a MAPQ>20, number of aligned bases >5000 nucleotides and a percent identity > 50%. For the *M. belari* Y chromosome, the largest summed alignment blocks totalled 422 kbp against an X-assigned scaffold (scaffold_16; shown in Fig. 5B). The next longest alignments spanned 235kbp on an autosome (scaffold_20), but they were almost all composed of a single repeated element on scaffold_20 shared across many Y-assigned scaffolds. For *M. monhystera*, the largest summed alignment blocks totalled only 29kbp on an autosome (scaffold_11).

We also aligned the Y-assigned scaffolds between the two pseudosexual species, also using minimap2. Applying the same filters, no alignment blocks were found.

Search for sex determining genes

We identified the well-known X:autosome ratio counting genes from *C. elegans* based on (15): the two X-linked counters *sex-1* and *fox-1*, the two autosomal counters *sea-1* and *sea-2* and the integrator *xol-1*. These were extracted from the *C. elegans* proteome downloaded from Wormbase Parasite Release 18, and the longest isoform of each protein sequence searched against each of our four *Mesorhabditis* genome assemblies using miniprot v0.13-r248 (58). No hits were found for *sea-1* or *xol-1*. For the remaining genes, a single hit was found across all four species: all hits were on autosomes, and consistent in terms of Nigon elements. This table is provided on Zenodo (see data availability section).

Age estimation of the ancestral neo-Y

Our results support neo-Y emergence following a single X-autosome fusion event in the ancestor of the two pseudosexual species *M. belari* and *M. monhystera* (Fig. 4B). As a conservative estimate, we thus used the speciation point between *M. belari* and *M. monhystera* in our *Mesorhabditis* phylogeny (Fig. 2A) as the latest possible time of origin of the neo-Y.

We then used a recently time-calibrated phylogeny of the nematode clade Rhabditina (also called clade V), to which *Mesorhabditis* belongs (16). Their calibration relied on mutation rates and generation times as estimated in the genus *Caenorhabditis*. In their phylogeny (Fig. S3 of (16)), the two species *M. belari* and *M. spiculigera* diverged ~86 million years ago. From this, we compared the point of speciation of *M. belari* and *M. monhystera* with that of *M. belari* and *M. spiculigera*, that forms the leftmost point in our *Mesorhabditis* phylogeny (Fig. 2A). We obtained two different age estimates, depending on whether we used *M. belari* or *M. spiculigera* as the reference for the present - e.g., due to different substitution rates in the two species - : 39.7 million years using *M. belari* and 51.45 million years using *M. spiculigera*. We thus estimated the age of the neo-Y at ~45 million years.

Testing for a fertilisation drive in *M. monhystera*

To test for a fertilisation drive, we isolated young gravid females, dissected their embryos, and mounted them on agarose pads for live DIC imaging. We looked for one-cell embryos containing a decondensed male pronucleus, indicative of an amphimictic embryo, before placing these on individual enriched plates, and identifying the sex of the resulting adult individual phenotypically. Of 16 amphimictic embryos, 5 did not hatch, and 11 hatched to become adult males. We then performed a chi-square goodness-of-fit test in R, testing the null hypothesis that viable amphimictic embryos were fertilised by X- and Y-bearing sperm at 50% frequency, in which case males and females are also expected at 50% frequency. At a p-value threshold of 0.05, observing 11/11 males is incompatible with a Y-bearing fertilisation rate below 74%.

Analysis of genes on the neo-Y chromosomes

To exclude pseudogenes, we only analysed genes with some evidence of expression in either adult males or adult females. We used three adult male and two adult female samples for *M. belari* and *M. monhystera*, and one male and one female sample for *M. longespiculosa* and *M. spiculigera* (fewer biological replicates as they do not have any Y genes). Reads were trimmed using fastp v0.23.2 (59) (--cut_mean_quality 20 --cut_by_quality3 --qualified_quality_phred 15 --length_required 50) and transcript counts obtained using Kallisto v0.50.1 (47), mapped against all predicted coding sequences (see 'Gene prediction' section above). Counts were normalised using the size-factor normalisation function of DESeq2 (48) in R.

We then filtered out genes with less than a mean normalised RNA-Seq count of 10 in either males or females. For *M. belari*, this removed 6.2%, 16.8% and 36.7% of all X, autosomal and Y genes respectively, and for *M. monhystera* this removed 5.8%, 11.2% and 28.2% of all X, autosomal and Y genes respectively. We note that Y-assigned genes also more frequently had no RNA-Seq support at all: ~7% of all genes for both species, compared to 0.4-1.5% for X and autosomal genes. In *M. belari*, 9 Y-assigned genes had significant RNA-Seq support in females and were discarded from analysis.

Search for male-beneficial or candidate fertilisation driver genes

To analyse patterns of orthology and paralogy of genes on the neo-Y chromosomes, we first predicted families of homologs using OrthoFinder v2.5.5 (17). We used the predicted protein sequences of our four focal *Mesorhabditis* species, plus the proteome of *C. elegans* obtained from WormBase ParaSite WBPS18 (45) to identify any known *C. elegans* homologs for genes of interest. OrthoFinder groups genes into families called 'Orthogroups'

that include both orthologs (homologs across species), and paralogs (homologs in the same species, i.e. arising from a gene duplication). We then looked for Orthogroups with members on the neo-Y of both pseudosexuals (*M. belari* and *M. monhystera*) and Orthogroups with members exclusively on the neo-Y of either pseudosexual (i.e., no paralogs on autosomes or the X chromosome).

To functionally annotate genes, we ran eggNOG v2.1.12 (19) on the proteomes of our four focal species, and also used a table of known functions of *C. elegans* genes, parsed from the WormBase ParaSite proteome file, for when a *C. elegans* ortholog was found with OrthoFinder.

To analyse the evolution of specific gene families, we used the same methods as for the *Mesorhabditis* phylogeny (see above): alignments were produced using mafft, trimmed using clipkit, and trees produced using IQ-TREE. Alignments were visualised using seaview version 5 (60).

We used orthology to *C. elegans* and domain-level annotations from eggNOG to look for species-specific genes that could be candidates for the fertilisation drive of the neo-Y chromosomes. In *M. belari*, we found three different families, encoding smc-5 genes (involved in meiosis), condensins (involved in chromosome compaction), and bub-1 genes (a kinetochore protein, involved in chromosome segregation), all of which could be linked to meiosis, and thus to biased segregation of proteins to Y-bearing sperm. In *M. monhystera*, we found one family encoding PRG-1 genes, part of the Argonaute family, which has been tied to a meiotic driver in *D. melanogaster* (61). However, all of these had closely-related paralogs on the X or autosomes, bearing only a few Y-specific substitutions, making them unlikely candidates.

As most genes occurring exclusively on the neo-Y of either pseudosexual had no annotations from eggNOG, we further ran hh-suite v3.3.0 (62) (hhblits -d uniclust30 -p 90 -noaddfilter -e 0.0000001) on each of these genes to obtain further domain-level annotations. We then filtered out any hits with a homology probability <90%, E-value >1, or with no associated functional terms.

All alignment files and annotation tables of our candidates of interest are provided on Zenodo (see data availability section).

Analysis of satellite repeats

We used two approaches to identify satellite sequences. In the first, we used Tandem Repeats Finder (trf) v4.10.0-rc.2 (63) (2 7 7 80 10 50 2000 -h -ngs) to identify satellites in the Y- and X-assigned assembled scaffolds of *M. belari* and *M. monhystera*. We filtered the raw outputs of trf to include only satellites with a minimum size of 5bp, tandem copy number of 10, and minimum identity to the consensus repeat of 80%. Our candidate satellites were then searched in the entire assemblies using seqkit v2.4.0 (seqkit locate), and satellites with a minimum total copy number of 40, and with at least 70% total occurrences on Y- or X-assigned scaffolds (compared to the rest of the genome) were extracted. This yielded 32 Y-enriched and 5 X-enriched satellites in *M. belari* and 112 Y-enriched satellites in *M. monhystera*. None of the satellites were shared between species.

Because satellite repeats are hard to assemble, we also used k-seek (64) to discover satellites from sequencing reads directly. Detection was performed on our male and female Illumina DNA-Seq reads (used for assigning scaffolds, see above) for *M. belari* and *M. monhystera*. All satellite counts were normalised by the median coverage of the reads along our genome assemblies, so that counts correspond to the number of satellites per 1x sequenced position. We then looked for satellites with a count of at least 40 in males (corresponding roughly to the number of copies in the genome) and with >4 times more counts in males compared to females. This yielded 9 Y-enriched and 8 Y-enriched satellites in *M. belari* and *M. monhystera*, respectively (after excluding 5 Y-enriched satellites in *M. belari* already found using the previous approach). Again, none of these were shared between species. To be less stringent, we also found no satellite with >2 times the counts in males, and at least 10 copies, across both species.

All the species-specific satellites we identified across both approaches are provided on Zenodo (see data availability section).

Analysis of mitochondrial genomes

Mitochondrial genome assembly
To reconstruct the mitochondrial genomes, we re-used the DNA-Seq data for the same *Mesorhabditis* species that we used to build our phylogeny (Fig. 2A). Reads were cleaned using fastp (59). The genomes of all species were assembled using Flye (36). Mitochondrial genomes were then assembled and annotated using MitoHiFi (65) with the default annotation tool (66) and using the *C. elegans* mitochondrial genome as reference (NC_001328.1). To ensure absence of contamination, mitochondrial genome coverage was measured. For this, the contig identified as the mitochondrial sequence in the Flye assemblies was removed and replaced by the assembled mitochondrial genome from MitoHiFi, as the former tends to contain multiple copies of the mitochondrial genome (due to its circularity). Reads were mapped against the assembly using minimap2 (49) and the coverage estimated using samtools (51).

Mitochondrial phylogeny
To reconstruct the mitochondrial phylogeny, we used the annotated genes from MitoHiFi. For each species, 12 mitochondrial protein coding genes were aligned with PRANK (67) using a codon model and sites ambiguously aligned were removed with Gblocks (68). The 13 genes were concatenated and and a phylogenetic tree was built using PhyML v3.0 (69) under a GTR+G+I model with 100 bootstrap replicates and was rooted using the clade [*M. sp* ana0126*, M. irregularis and M. spiculigera*].

Reproducibility
Analyses were performed using Nextflow workflows (70), with individual analysis steps run using Charliecloud containers (71) or Conda environments built tools available on Bioconda (72). All used tools are thus publicly available via Bioconda. The workflows and their configuration and analysis code supporting this work are publicly available at https://doi.org/10.5281/zenodo.15407996.

Data Availability

All assembled genomes and raw sequencing data are available at the ENA under project accession xx.

Data tables underlying the analyses and figures in the paper are publicly available at https://doi.org/10.5281/zenodo.15407996.

Author Contributions

Conceptualisation: BL, MD
Data Curation: BL
Funding acquisition: MD
Investigation: BL, LS, NS, AH, EW, MD
Methodology: BL, LS, NS, MD
Software: BL
Visualisation: BL, NS
Resources: MK, CL, MB
Writing - original draft: BL, MD
Writing - review & editing: BL, LS, NS, MB, MD

Competing interests

Authors declare that they have no competing interest.

Acknowledgements

For computing, we gratefully acknowledge the resources of the HPC at ENS de Lyon (managed by the Pôle Scientifique de Modélisation Numérique (73)) on which all workflows were deployed and run.

This work was supported by a grant from the Agence Nationale de la Recherche: ANR-19-CE02-0012.

We also thank Quentin Helleu and Cécile Courret for productive discussions.

References

1. Gilbert SF. Chromosomal Sex Determination in Mammals. In: Developmental Biology 6th edition [Internet]. Sinauer Associates; 2000 [cited 2025 Apr 9]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK9967/

2. Haag E. The evolution of nematode sex determination: C. elegans as a reference point for comparative biology. WormBook [Internet]. 2005 [cited 2025 Apr 9]; Available from: http://www.wormbook.org/chapters/www_evolutionsexdetermin/evolutionsexdetermin.html

3. Madl JE, Herman RK. Polyploids and sex determination in Caenorhabditis elegans. Genetics. 1979 Oct;93(2):393–402.

4. Stothard P, Pilgrim D. Sex-determination gene and pathway evolution in nematodes. BioEssays. 2003 Mar;25(3):221–31.

5. Gonzalez de la Rosa PM, Thomson M, Trivedi U, Tracey A, Tandonnet S, Blaxter M. A telomere-to-telomere assembly of *Oscheius tipulae* and the evolution of rhabditid

nematode chromosomes. Rockman M, editor. G3 GenesGenomesGenetics. 2021 Mar 10;11(1):jkaa020.

6. Stevens L, Kieninger M, Chan B, Wood JMD, Rosa PG de la, Allen J, et al. The genome of Litomosoides sigmodontis illuminates the origins of Y chromosomes in filarial nematodes. PLOS Genet. 2024 Jan 16;20(1):e1011116.

7. Post R. The chromosomes of the Filariae. Filaria J. 2005 Nov 2;4(1):10.

8. Yoshida K, Witte H, Hatashima R, Sun S, Kikuchi T, Röseler W, et al. Rapid chromosome evolution and acquisition of thermosensitive stochastic sex determination in nematode androdioecious hermaphrodites. Nat Commun. 2024 Nov 7;15(1):9649.

9. Grosmaire M, Launay C, Siegwald M, Brugière T, Estrada-Virrueta L, Berger D, et al. Males as somatic investment in a parthenogenetic nematode. Science. 2019 Mar 15;363(6432):1210–3.

10. Launay C, Félix MA, Dieng J, Delattre M. Diversification and hybrid incompatibility in auto-pseudogamous species of Mesorhabditis nematodes. BMC Evol Biol. 2020 Aug 18;20(1):105.

11. Blanc C, Saclier N, Le Faou E, Marie-Orleach L, Wenger E, Diblasi C, et al. Cosegregation of recombinant chromatids maintains genome-wide heterozygosity in an asexual nematode. Sci Adv. 2023 Aug 25;9(34):eadi2804.

12. Carlton PM, Davis RE, Ahmed S. Nematode chromosomes. Genetics. 2022 May 1;221(1):iyac014.

13. Tandonnet S, Koutsovoulos GD, Adams S, Cloarec D, Parihar M, Blaxter ML, et al. Chromosome-Wide Evolution and Sex Determination in the Three-Sexed Nematode Auanema rhodensis. G3 GenesGenomesGenetics. 2019 Apr 1;9(4):1211–30.

14. Bachtrog D. The Y Chromosome as a Battleground for Intragenomic Conflict. Trends Genet. 2020 Jul 1;36(7):510–22.

15. Farboud B, Nix P, Jow MM, Gladden JM, Meyer BJ. Molecular antagonism between X-chromosome and autosome signals determines nematode sex. Genes Dev. 2013 May 15;27(10):1159–78.

16. Picao-Osorio J, Bouleau C, Gonzalez De La Rosa PM, Stevens L, Fekonja N, Blaxter M, et al. Evolution of developmental bias explains divergent patterns of phenotypic evolution [Internet]. 2025 [cited 2025 Mar 25]. Available from: http://biorxiv.org/lookup/doi/10.1101/2025.02.04.636455

17. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome Biol. 2019 Nov 14;20(1):238.

18. Ishihara T, Iino Y, Mohri A, Mori I, Gengyo-Ando K, Mitani S, et al. HEN-1, a Secretory Protein with an LDL Receptor Motif, Regulates Sensory Integration and Learning in Caenorhabditis elegans. Cell. 2002 May 31;109(5):639–49.

19. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, et al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res. 2019 Jan 8;47(D1):D309–14.

20.   Schedl T, Graham PL, Barton MK, Kimble J. Analysis of the role of tra-1 in germline sex determination in the nematode Caenorhabditis elegans. Genetics. 1989 Dec;123(4):755–69.

21.   Guo Y, Chen X, Ellis RE. Evolutionary Change within a Bipotential Switch Shaped the Sperm/Oocyte Decision in Hermaphroditic Nematodes. PLOS Genet. 2013 Oct 3;9(10):e1003850.

22.   Hodgkin J. A genetic analysis of the sex-determining gene, tra-1, in the nematode Caenorhabditis elegans. Genes Dev. 1987 Sep 1;1(7):731–45.

23.   Lawson HN, Wexler LR, Wnuk HK, Portman DS. Dynamic, Non-binary Specification of Sexual State in the C. elegans Nervous System. Curr Biol. 2020 Sep 21;30(18):3617-3623.e3.

24.   Herbette M, Wei X, Chang CH, Larracuente AM, Loppin B, Dubruille R. Distinct spermiogenic phenotypes underlie sperm elimination in the Segregation Distorter meiotic drive system. Bosco G, editor. PLOS Genet. 2021 Jul 6;17(7):e1009662.

25.   Courret C, Ogereau D, Gilbert C, Larracuente AM, Montchamp-Moreau C. The Evolutionary History of Drosophila simulans Y Chromosomes Reveals Molecular Signatures of Resistance to Sex Ratio Meiotic Drive. Mol Biol Evol. 2023 Jul 1;40(7):msad152.

26.   Kern EMA, Kim T, Park JK. The Mitochondrial Genome in Nematode Phylogenetics. Front Ecol Evol [Internet]. 2020 Aug 4 [cited 2025 Apr 17];8. Available from: https://www.frontiersin.orghttps://www.frontiersin.org/journals/ecology-and-evolution/articles/10.3389/fevo.2020.00250/full

27.   Henzel JV, Nabeshima K, Schvarzstein M, Turner BE, Villeneuve AM, Hillers KJ. An Asymmetric Chromosome Pair Undergoes Synaptic Adjustment and Crossover Redistribution During Caenorhabditis elegans Meiosis: Implications for Sex Chromosome Evolution. Genetics. 2011 Mar 1;187(3):685–99.

28.   Rozen S, Skaletsky H, Marszalek JD, Minx PJ, Cordum HS, Waterston RH, et al. Abundant gene conversion between arms of palindromes in human and ape Y chromosomes. Nature. 2003 Jun;423(6942):873–6.

29.   Shakes DC, Neva BJ, Huynh H, Chaudhuri J, Pires-daSilva A. Asymmetric spermatocyte division as a mechanism for controlling sex ratios. Nat Commun. 2011 Jan 18;2(1):157.

30.   Meng X, Yamashita YM. Intrinsically weak sex chromosome drive through sequential asymmetric meiosis. Sci Adv. 2025 May 7;11(19):eadv7089.

31.   Serra L, Chang D, Macchietto M, Williams K, Murad R, Lu D, et al. Adapting the Smart-seq2 Protocol for Robust Single Worm RNA-seq. BIO-Protoc [Internet]. 2018 [cited 2025 May 13];8(4). Available from: https://bio-protocol.org/e2729

32.   Cheng H, Concepcion GT, Feng X, Zhang H, Li H. Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. Nat Methods. 2021 Feb;18(2):170–5.

33.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. ArXiv13033997 Q-Bio [Internet]. 2013 Mar 16 [cited 2020 Mar 28]; Available from:

http://arxiv.org/abs/1303.3997

34. Zhou C, McCarthy SA, Durbin R. YaHS: yet another Hi-C scaffolding tool [Internet]. Bioinformatics; 2022 Jun [cited 2022 Nov 11]. Available from: http://biorxiv.org/lookup/doi/10.1101/2022.06.09.495093

35. Challis R, Richards E, Rajan J, Cochrane G, Blaxter M. BlobToolKit – Interactive Quality Assessment of Genome Assemblies. G3 GenesGenomesGenetics. 2020 Feb 18;10(4):1361–74.

36. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat graphs. Nat Biotechnol. 2019 Apr 1;1.

37. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 2021 Oct 1;38(10):4647–54.

38. Rhie A, Walenz BP, Koren S, Phillippy AM. Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. Genome Biol. 2020 Sep 14;21(1):245.

39. Durand NC, Shamim MS, Machol I, Rao SSP, Huntley MH, Lander ES, et al. Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell Syst. 2016 Jul 27;3(1):95–8.

40. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002 Jul 15;30(14):3059–66.

41. Steenwyk JL, Buida TJ, Li Y, Shen XX, Rokas A. ClipKIT: A multiple sequence alignment trimming software for accurate phylogenomic inference. Hejnol A, editor. PLOS Biol. 2020 Dec 2;18(12):e3001007.

42. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation. PLOS ONE. 2016 Oct 5;11(10):e0163962.

43. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol. 2015 Jan 1;32(1):268–74.

44. Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, et al. BRAKER3: Fully Automated Genome Annotation Using RNA-Seq and Protein Evidence with GeneMark-ETP, AUGUSTUS and TSEBRA [Internet]. bioRxiv; 2023 [cited 2023 Aug 23]. p. 2023.06.10.544449. Available from: https://www.biorxiv.org/content/10.1101/2023.06.10.544449v1

45. Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. WormBase ParaSite − a comprehensive resource for helminth genomics. Mol Biochem Parasitol. 2017;215:2–10.

46. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013 Jan 1;29(1):15–21.

47. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq

quantification. Nat Biotechnol. 2016 May;34(5):525–7.

48.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014 Dec 5;15(12):550.

49.   Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018 Sep 15;34(18):3094–100.

50.   Picard toolkit [Internet]. Broad Institute, GitHub repository. Broad Institute; 2019. Available from: https://broadinstitute.github.io/picard/

51.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of SAMtools and BCFtools. GigaScience. 2021 Feb 1;10(2):giab008.

52.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics. 2010 Mar 15;26(6):841–2.

53.   Rey C, Launay C, Wenger E, Delattre M. Programmed DNA elimination in Mesorhabditis nematodes. Curr Biol. 2023 Sep;33(17):3711-3721.e5.

54.   Baril T, Galbraith J, Hayward A. Earl Grey: A Fully Automated User-Friendly Transposable Element Annotation and Analysis Pipeline. Mol Biol Evol. 2024 Apr 1;41(4):msae068.

55.   Lee S, Bakker CR, Vitzthum C, Alver BH, Park PJ. Pairs and Pairix: a file format and a tool for efficient storage and retrieval for Hi-C read pairs. Bioinformatics. 2022 Mar 4;38(6):1729–31.

56.   Open2C, Abdennur N, Fudenberg G, Flyamer IM, Galitsyna AA, Goloborodko A, et al. Pairtools: From sequencing data to chromosome contacts. PLOS Comput Biol. 2024 May 29;20(5):e1012164.

57.   Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative genomics viewer. Nat Biotechnol. 2011 Jan;29(1):24–6.

58.   Li H. Protein-to-genome alignment with miniprot. Bioinformatics. 2023 Jan 1;39(1):btad014.

59.   Chen S. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication using fastp. iMeta. 2023;2(2):e107.

60.   Gouy M, Tannier E, Comte N, Parsons DP. Seaview Version 5: A Multiplatform Software for Multiple Sequence Alignment, Molecular Phylogenetic Analyses, and Tree Reconciliation. In: Katoh K, editor. Multiple Sequence Alignment: Methods and Protocols [Internet]. New York, NY: Springer US; 2021 [cited 2025 Jul 2]. p. 241–60. Available from: https://doi.org/10.1007/978-1-0716-1036-7_15

61.   Gell SL, Reenan RA. Mutations to the piRNA Pathway Component Aubergine Enhance Meiotic Drive of Segregation Distorter in Drosophila melanogaster. Genetics. 2013 Mar 1;193(3):771–84.

62.   Steinegger M, Meier M, Mirdita M, Vöhringer H, Haunsberger SJ, Söding J. HH-suite3 for fast remote homology detection and deep protein annotation. BMC Bioinformatics. 2019 Sep 14;20(1):473.

63.   Benson G. Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res. 1999 Jan 1;27(2):573–80.

64.   Wei KHC, Grenier JK, Barbash DA, Clark AG. Correlated variation and population differentiation in satellite DNA abundance among lines of Drosophila melanogaster. Proc Natl Acad Sci. 2014 Dec 30;111(52):18793–8.

65.   Uliano-Silva M, Ferreira JGRN, Krasheninnikova K, Blaxter M, Mieszkowska N, Hall N, et al. MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio high fidelity reads. BMC Bioinformatics. 2023 Jul 18;24(1):288.

66.   Allio R, Schomaker-Bastos A, Romiguier J, Prosdocimi F, Nabholz B, Delsuc F. MitoFinder: Efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics. Mol Ecol Resour. 2020;20(4):892–905.

67.   Löytynoja A, Goldman N. Phylogeny-Aware Gap Placement Prevents Errors in Sequence Alignment and Evolutionary Analysis. Science. 2008 Jun 20;320(5883):1632–5.

68.   Castresana J. Selection of Conserved Blocks from Multiple Alignments for Their Use in Phylogenetic Analysis. Mol Biol Evol. 2000 Apr 1;17(4):540–52.

69.   Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. Syst Biol. 2010 May 1;59(3):307–21.

70.   Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C. Nextflow enables reproducible computational workflows. Nat Biotechnol. 2017 Apr;35(4):316–9.

71.   Priedhorsky R, Randles T. Charliecloud: unprivileged containers for user-defined software stacks in HPC. In: Proceedings of the International Conference for High Performance Computing, Networking, Storage and Analysis [Internet]. New York, NY, USA: Association for Computing Machinery; 2017 [cited 2025 Mar 25]. p. 1–10. (SC '17). Available from: https://dl.acm.org/doi/10.1145/3126908.3126925

72.   Grüning B, Dale R, Sjödin A, Chapman BA, Rowe J, Tomkins-Tinch CH, et al. Bioconda: sustainable and comprehensive software distribution for the life sciences. Nat Methods. 2018 Jul;15(7):475–6.

73.   Quemener E, Corvellec M. SIDUS—the solution for extreme deduplication of an operating system. Linux J. 2013 Nov 1;2013(235):3:3.