Plasmids link antibiotic resistance genes and phage defense systems in *E. coli*

Weronika Ślesak[1,*], Piotr Jedryszek[1], Daniel Cazares[1,**], William Matlock[1,**], R. Craig MacLean[1,2,*]

[1]Department of Biology, University of Oxford, 11a Mansfield Road, Oxford, UK

[2]All Souls College, High Street, Oxford, UK

*corresponding authors: weronika.slesak@sjc.ox.ac.uk; craig.maclean@biology.ox.ac.uk

**equal contribution

Abstract

Phage therapy has been proposed as an alternative to antibiotics to treat resistant infections. However, we have a limited understanding of how antibiotic resistance genes (ARGs) associate with bacterial phage defense systems (PDSs). Here, we explore the relationship between ARGs and PDSs in a sample of 2,559 plasmids originating from 1,044 *E. coli* isolates, representing a snapshot of clinical and non-clinical diversity in Oxfordshire, UK (2008-2020). In total, we identify 3,193 ARGs and 14,013 PDSs (180 unique types). We demonstrate that *E. coli* plasmids are enriched for ARGs and PDSs (both $p<0.001$), with a bias towards toxin-antitoxin, abortive infection, and restriction-modification systems (all $q<0.025$). We proceed to show that ARGs and PDSs are physically linked by plasmids ($p<0.001$). Together, our results suggest that phage therapy may inadvertently select for antibiotic resistant bacteria, and that antibiotic use may similarly drive resistance to phage.

Keywords: *Escherichia coli*; bacteriophage; anti-bacterial agent

Introduction

Bacterial defense systems can protect the cell against invaders, such as plasmids and phage[1]. Plasmids are extrachromosomal replicons often able to transfer between bacterial hosts via conjugation[2]. Phage, and in particular, lytic phage, are bacterial viruses that, like plasmids, rely on the host cell for replication, but unlike plasmids, ultimately kill the host during the infection cycle[3]. Plasmids are often associated with antimicrobial resistance genes (ARGs), which can confer resistance to the host cell, sometimes at a low fitness cost[4,5]. Moreover, plasmids can also encode phage defense systems (PDSs), such as toxin-antitoxin, restriction modification, and CRISPR-Cas systems, blurring the boundary between selfish genetic elements and mutualistic partners[6–9].

One unresolved issue is the extent to which ARGs and PDSs co-occur on plasmids. This is important because the physical linkage of different resistance determinants can drive co-selection: selective pressure for one trait (phage resistance) may inadvertently maintain another (antibiotic resistance), and vice versa. Co-selection is already well documented for ARGs and other resistance genes, such as those against heavy metals or biocides, where a single stressor can select for multi-resistant bacteria[10,11]. Some examples have been reported of ARGs and PDSs co-localising on the same mobile genetic element, yet whether this occurs more generally remains to be seen[12–14]. Understanding these patterns is particularly relevant as phage therapy gains traction as a strategy to combat antibiotic resistant infections[15,16].

In this study, we investigated the relationship between ARGs and PDSs in a large sample of 2,559 plasmids from 1,044 clinical and non-clinical *E. coli* isolates[17]. We first annotate the genomes for ARGs and PDSs, then explore (i) their overall enrichment on plasmids versus chromosomes, (ii) the enrichment of specific PDS types on plasmids versus chromosomes, and (iii) their linkage on plasmids.

Methods

Dataset curation

Isolates and associated genome assemblies were from an existing, pre-collated dataset[17]. The original dataset contained Enterobacterales species collected from within 60km in Oxfordshire from 2008-2020, covering multiple clinical and non-clinical niches: human bloodstream infection (BSI), livestock faeces (cows, pigs, poultry, and sheep), and wastewater (influent, effluent, and rivers upstream of effluent)[18,19]. All BSI isolates and a majority of non-BSI isolates were not selectively cultured for antimicrobial resistance. Species were determined with matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF). Authors used a hybrid approach (short- and long-reads) to recover reference-grade assemblies. We selected all the *E. coli* genomes where the chromosome and all plasmids were fully circularised (n= 1,044). Isolate and assembly metadata was also downloaded.

Genome annotation

We annotated for PDSs using the tool DefenseFinder (v. 2.0.0) with default paramaters, which queries against a database of known PDSs[20–22]. DefenseFinder also denotes when a system is "complete" ("systems" output), meaning all component genes have been annotated. We only considered complete systems in this study. For ARGs, we used pre-existing annotations from AMRFinderPlus (v. 3.10.18) with default parameters[23,24]. Plasmid mobility predictions (conjugative, mobilisable, or non-mobilisable) also used pre-existing annotations from MOB-typer from MOB-suite (v. 3.03) with default parameters. We annotated for integrons using IntegronFinder 2.0 with default parameters[25]. For gene annotations we used Prodigal (v. 2.6.3)[26] with default parameters.

Testing for the enrichment of ARGs and PDS genes on plasmids

We first calculated the total number of plasmid genes as a fraction of the total number of genes in the genome. Similarly, we calculated the fraction of just ARGs and PDSs genes on plasmids. These proportions were compared using a Pearson's chi-squared test with Yates' continuity correction. We conducted this across the whole dataset (see Results), but also separately for BSI-associated and non-BSI-associated plasmids (Figure S5).

Testing for the over-representation of PDS families on plasmids

First, for each PDS type, we calculated a binary presence/absence score for each replicon. We next wanted to generate a null distribution that accounted for some genomes containing more plasmids than others. To do this, we performed $b=10,000$ label permutations of replicon type (plasmid or chromosome) within each genome. After each permutation, we recorded the number of plasmid-labelled contigs carrying each PDS. This generated a one-sided p-value$=(n+1)/(b+1)$, where n was the number of permutations with a plasmid count at least as extreme as observed ($\geq$ for the plasmid-enrichment test; $\leq$ for the chromosome-enrichment test) and $b=10,000$. Separate p-values were obtained for plasmid (H$_1$: observed $>$ null) and chromosome (H$_1$: observed $<$ null) bias. To mitigate against multiple tests increasing the rate of Type I errors, we applied both the Benjamini–Hochberg (BH) and the Benjamini–Yekutieli (BY) false-discovery-rate procedures. Families with BH-adjusted q < 0.025 for the relevant one-sided test (plasmid or chromosome) were called significant; BY-adjusted q-values are also reported when q < 0.025 (providing a stricter, dependence-robust correction).

Testing the association between ARGs and PDSs on plasmids

If two plasmids are vertical descendants, or share large portions of their sequence from a recombination event, they might contain many of the same genes. This can be problematic

when testing for the association of ARGs and PDSs, as it introduces pseudo-replicates to the

model. We minimised the risk of pseudo-replication by genetically dereplicating plasmids in

the dataset. We first took the *k*-mer plasmid clusters generated by the original authors[17]. We

then randomly selected a representative from each cluster. In total, we recovered *n*=712

genetically representative plasmids.

We next predicted the count of ARGs per plasmid with a generalised additive mixed-model

(GAMM) using the mgcv library[27]. Model choice was guided by Akaike Information Criterion

(AIC), which balances goodness-of-fit and parsimony. We found that a Tweedie family (log-

link) GAMM with a log-link family outperformed analogous GAMM/GLMMs with Poisson

and negative binomial families.

The model used the fixed effects (i) plasmid PDS count, (ii) plasmid GC content (mean

centred), and (iii) sampling niche (BSI-associated, livestock-associated, or wastewater-

associated). We used plasmid replicon type as a random intercept to control for plasmid

lineage. We also included an offset of $\log_{10}$-scaled plasmid length (bp). Originally, we also

included plasmid integron count and plasmid mobility (non-mobilisable, mobilisable,

conjugative) as fixed-effects, but these were dropped since they were insignificant (likelihood-

ratio test *p*-value>0.05). Plasmid PDS count and GC content initially used thin-plate regression

splines ($k = 4$), but by assessing the effective degrees of freedom and plots of the smooth terms,

we concluded they were consistent with linearity, so the smooth terms were dropped.

Lastly, we checked for multicollinearity and correlations among predictors. For numeric-

numeric fixed-effect pairs, we computed Spearman's rank correlation, and for numeric-

categorical fixed-effect pairs, we calculated the Kruskal-Wallis effect size. No issues were

identified.

Of the *n*=712 genetically representative plasmids, 23% (167/712) had inconclusive plasmid

replicon typing by PlasmidFinder (recorded as "NA"). Nonetheless, re-running the GAMM

with these plasmids excluded did not meaningfully change the coefficient estimates (95% CIs

overlapped).

Code and data availability

The data and metadata used in this study will be stably archived on Zenodo. The code to

reproduce the analysis will be made available on GitHub, and also archived on Zenodo.

Supplementary File 1 contains the permutation test output. Supplementary File 2 contains the

GAMM diagnostics.

Results

Plasmids are enriched for antimicrobial resistance genes and phage-defense system genes.

We first curated a sample of *n*=1,044 *E. coli* isolates, a common human commensal and

pathogen, with *n*=2,559 plasmids (see Methods). Isolates were sampled within a geographically

(<60km in Oxfordshire, UK) and temporally (2008-2020) restricted frame, and represented

both clinical (human bloodstream infection [BSI]) and non-clinical (livestock-associated and

wastewater-associated) niches, totaling 52% (547/1,044) BSI, 42% (433/1,044) livestock-

associated, and 6% (64/1,044) wastewater-associated. Livestock-associated isolates were

further divided into 32.8% (142/433) cattle-associated, 28.9% (125/433) pig-associated, 11.8%

(51/433) poultry-associated, and 26.5% (115/433) sheep-associated. In total, 89% (938/1,044)

of isolates carried a plasmid. Isolates carried a median of 2 plasmids (range: 0-16), with the

most common replicon types being small colicinogenic-type plasmids (see Methods).

We next annotated our genomes for ARGs and PDSs (see Methods). In total, we identified

n=14,013 PDS hits with 180 distinct system types. Every chromosome carried at least one PDS,

contrasted with only 41% (1,051/2,559) of plasmids. Comparatively, ARGs were less common,

with 3,193 identified in total. Overall, 27% (286/1,044) of chromosomes and 20% (521/2,559)

of plasmids carried at least one ARG, respectively. Of plasmids that carried at least one ARG

or PDS, the three top replicon types were IncF-types: IncFIB, Col156-IncFIB-IncFII, and

IncFIB-IncFIC, which together accounted for 27% (325/1,225) of that plasmid subset.

Distributions of ARGs and PDSs were variable across *E. coli* niches (Figures S1–S4).

Livestock-associated plasmids (cattle, pig, poultry, and sheep) carried defense systems in 44%

(475/1,080) of cases but carried ARGs in only 13% (143/1,080). Within this group, pig-

associated plasmids were distinct: they encoded ARGs roughly four-fold more frequently: 25%

(101/403) of pig plasmids versus 6% (42/677) of plasmids from the other three livestock

groups, while their PDS prevalence remained comparable (39% [158/403] versus 47%

[317/677]). Wastewater-associated plasmids showed intermediate prevalence: 35% (53/150)

encoded PDSs and 19% (29/150) carried ARGs. BSI-associated plasmids exhibited the greatest

combined load outside pigs, with 39% (523/1,329) containing PDSs and 26% (349/1,329)

carrying ARGs.

Next, we tested whether ARGs and PDSs were enriched on plasmids versus chromosomes

(Figure 1; see Methods). Of the n=4,999,278 annotated coding sequences in our dataset, only

3.42% (170,778/4,999,278) were found on plasmids. By contrast, 8.66% of all PDS genes

(2,474/28,581) and 75.76% of all ARGs (2,419/3,193) were plasmid-encoded. Pearson's χ^2

tests (Yates-corrected) confirmed that both categories were highly over-represented on

plasmids relative to the genomic baseline (ARGs: $\chi^2 = 49{,}943.4$, df = 1, $p < 0.001$; PDSs: $\chi^2 =$

$2{,}343.8$, df = 1, $p < 0.001$). The magnitude of enrichment was therefore ~2.5-fold for PDS

genes and ~22-fold for ARGs. Surprisingly, this trend carried for both clinical and non-clinical

isolates (Figure S5).

Figure 1. Plasmids are hotspots for ARGs and PDSs in *E. coli*. Bars show the percentage of genes in that category residing on plasmids. Both comparisons were significant at the p<0.001 level (Pearson χ^2 tests with Yates correction).

Plasmids and chromosomes are associated with distinct repertoires of phage-defense

systems.

We next explored whether the types of PDSs varied between plasmids and chromosomes. We

conducted a permutation test for each PDS (see Methods) and found 8% (15/180) of the PDS

families preferentially resided on plasmids (Figure 2). These 15 families made up 42.46%

(1,343/3,163) of defense-system occurrences on plasmids. In contrast, 20% (36/180) PDSs

were associated with chromosomes (Figure S6). Tests for the remaining 72% (129/180) PDSs

were inconclusive.

We found that TIR-III and Pif systems and two small uncharacterised families (HEC-09, DS-

28) were exclusively associated with plasmids in this dataset. Conversely, toxin-antitoxin

module MazEF and the cyclic-oligonucleotide producer CBASS were enriched more modestly

(~0.25, Figure 2A). When enrichment was quantified as absolute surplus (Figure 2B), the toxin-

antitoxin complex Mok-Hok-Sok was an outlier, occurring on plasmids 661 times versus the

approximately 100 occurrences expected by chance. Progressively smaller and significant

surpluses were observed for the remaining systems.

Figure 2. Defense systems associated with plasmids. (A) The x-axis shows the proportion of contigs that carried the PDS type. Each point represents a PDS type whose frequency on plasmids was greater than expected when contig labels (plasmid/chromosome) were randomly shuffled within isolates under two one-tailed permutation tests (see Methods). Open grey circles indicate p<0.05 significance under Benjamini-Hochberg (BH) correction only. Filled blue circles indicate significance under both BH and Benjamini-Yekutieli (BY) correction, which does not assume independence. The dashed line marks the genome-wide plasmid share of defence systems ($\bar{\pi} \approx 0.16$). **(B)** Bars count the excess number of plasmid-labelled contigs carried by each type ("observed – expected", where the expectation equals the type's total contig count $\times \bar{\pi}$). Bars are coloured the same as points in **(A)**. Only PDS families that were significantly over-represented on plasmids in the within-isolate permutation test are shown.

Phage-defense system count predicts ARG gene count on plasmids independent of plasmid

size.

Lastly, we explored the distribution of ARGs and PDSs within the genomes. We tested whether

ARGs and PDSs co-occurred on the same plasmids by constructing a generalised additive

mixed-model (GAMM) with a Tweedie family and log-link (see Methods). Briefly, we used

plasmid ARG count as a response with fixed effects (i) plasmid PDSs count, (ii) plasmid GC

content (mean centred), and (iii) sampling niche (BSI-associated, livestock-associated, or

wastewater-associated). We used plasmid replicon type as a random intercept to control for

plasmid lineage. To avoid pseudo-replication of plasmid sequences, we used single

representatives from clusters generated by genetic similarity. We also included an offset of

$\log_{10}$-scaled plasmid length (bp). Our total sample size was $n=712$ genetically representative

plasmids. The model output is detailed in Table 1 and visualised in Figure 3a. Further, Figure

3b visualises the ARG-PDS co-occurrence heatmap for all plasmids ($n=2,559$) in the dataset.

We found that the count of PDSs was a strong positive predictor of ARG count: each additional

phage-defense system increased the chance of an ARG being present on a plasmid by 48%

(95% CI=[18%, 86%]; $p<0.001$). Similarly, each 1% increase in plasmid GC content above

average increased ARG count by 37% (95% CI=[28%, 48%]; $p<0.001$). We found no signal

from isolate sampling niche. The random-effect smooth term for plasmid replicon type was

consistent with substantial non-linear variation (edf = 25.7, $F = 1.21$, $p<0.001$). Overall, the

model demonstrated good fit with adjusted $R^2=0.40$.

All isolates (1044/1044) carried a chromosomal PDS, meaning that all isolates with plasmid-

borne ARG (429/1,044) had the potential for dual resistance. Additionally, of the 41%

(429/1,044) of isolates carrying an ARG-positive plasmid, 88% (378/429) also carried at least

one plasmid-borne PDS. In 75% (324/429) of these cases, they were both on the same plasmid

(Figure 4). Further, Figure S7 shows the relative frequencies of ARGs, PDSs, and their co-

localisation on plasmids, and Figure S8 shows how these frequencies vary with plasmid size in

clinical versus non-clinical *E. coli*, consistent with clinical isolates, unlike non-clinical isolates,

often carrying small plasmids (<13kbp) that link ARGs and PDSs.

Figure 3. PDSs and ARGs co-occur on plasmids. **(A)** Forest plot of GAMM predictor coefficients as fold-changes with 95% confidence intervals ($\log_{10}$-scaled axis). The vertical dashed line at x=1 marks the reference. **(B)** Heatmap of frequency of co-localisation of ARGs and PDSs on same plasmid (n= 347). Cell colour indicates count.

Covariate		Coefficient (95% CI)	Odds ratio (95% CI)	*p*-value
(Intercept)		-6.58 (-7.25, -5.91)	0.00 (0.00, 0.00)	<0.001
PDS count		0.39 (0.16, 0.62)	1.48 (1.18. 1.86)	<0.001
Plasmid GC content (mean-centred)		0.32 (0.24, 0.39)	1.37 (1.28, 1.48)	<0.001
Niche	BSI-associated	0.72 (0.19, 1.25)	2.05 (1.21, 3.48)	0.008
	Wastewater-associated	0.48 (-0.43, 1.39)	1.61 (0.65, 4.00)	0.3

Table 1. Parameter estimates for the GAMM. For niche, livestock-associated was the reference level. Odds ratios were calculated by taking the exponent.

Number of isolates bearing at least one plasmid containing:

Figure 4. Overlap of ARGs and PDSs on plasmids within *E. coli* isolates. Euler diagram illustrating the intersection between sets of *E. coli* isolates that carried (i) at least one plasmid with a PDS, (ii) at least one plasmid with an ARG, and (iii) at least one plasmid with both an ARG and a PDS. The size of each ellipse is proportional to the set size. The total sample size was *n*=884 isolates.

Discussion

We have analysed the distribution of ARGs and PDSs in a diverse sample of *E. coli* genomes.

We demonstrate that plasmids are enriched for both types of features, disproportionately carry

specific PDS families compared to the chromosome, and physically link ARGs and PDSs in

the genome. Overall, our findings suggest a fundamental role for plasmids in the dissemination

of PDSs on top of known associations with ARGs.

Our study has limitations. Whilst *E. coli* represents a predominant clinical pathogen, plasmids

are widespread across bacterial taxa and frequently cross species boundaries[28]. Future work

should expand these modelling approaches to incorporate a broader range of bacterial

hosts. We also did not model the co-occurrence of specific ARGs and PDSs. Capturing such

associations requires modelling frameworks that account for both vertical inheritance and

horizontal transfer events (such as those mediated by transposons) as well as methods capable

of resolving multivariate networks of association. Without such controls, pairwise tests risk

conflating indirect correlations with direct genetic linkage, as well as inflating Type I errors[29].

The co-localisation of ARGs and PDSs on plasmids can facilitate joint horizontal transfer and

coordinated regulation. We hypothesise that plasmids are an ideal platform for co-localising

ARGs and PDSs due to their variable copy number and potential for rapid evolution. In clinical

isolates, multicopy plasmids carrying $bla_{\text{TEM-1}}$ can amplify phenotypic resistance and

accelerate adaptive evolution of the gene[30,31]. In our dataset, $bla_{\text{TEM-1}}$ was often co-localised

with toxin-antitoxin systems such as RnlAB, PARIS, and AbiQ[32–34]on small (>13kb)

colicinogenic-type plasmids in the BSI-associated isolates. Analogously, plasmids carrying

PDSs may experience rapid and fluctuating selection in response to episodic lytic phage

attacks, favouring those that can quickly up or downregulate their PDSs[35,36].

Phage pressure alone can maintain mobile genetic elements carrying ARGs even in the absence of antibiotics. For example, in *Vibrio cholerae*, episodic phage challenge increased conjugation frequencies of integrative conjugative elements carrying both ARGs and PDSs[14]. In another study, *E. coli* IncF plasmids carrying antibiotic resistance reduced susceptibility to coliphage infection and persisted for around 10 days without antibiotic selection[37]. This suggests that simply curbing antibiotic use may be insufficient to reverse the selection for antibiotic and/or phage resistance. Future work should monitor plasmid gene frequencies, copy-number dynamics, and host-fitness effects under isolated antibiotic, isolated phage, and combined selection regimes, ideally incorporating varied dosing schedules and temporal patterns.

Funding

This work was supported by the UKRI Frontiers grant (EP/Y031067/1) to RCM. WS is supported by the Wellcome Trust grant (218514/Z/19/Z), Merck Sharp and Dohme Corp., and Janssen Pharmaceutica NV. PJ is supported by funding from the Biotechnology and Biological Sciences Research Council UKRI-BBSRC grant (BB/T008784/1).

Acknowledgements

The authors thank Rachel Wheatley and Liam P. Shaw for the valuable help and discussions.

Conflicting interests

The authors declare no conflicting interests.

Bibliography

1. Koonin, E. V., Makarova, K. S. & Wolf, Y. I. Evolutionary Genomics of Defense Systems in Archaea and Bacteria. *Annu Rev Microbiol* **71**, (2017).

2. Coluzzi, C., Garcillán-Barcia, M. P., De La Cruz, F. & Rocha, E. P. C. Evolution of Plasmid Mobility: Origin and Fate of Conjugative and Nonconjugative Plasmids. *Mol Biol Evol* **39**, (2022).

3. Egido, J. E., Costa, A. R., Aparicio-Maldonado, C., Haas, P.-J. & Brouns, S. J. J. Mechanisms and clinical importance of bacteriophage resistance. *FEMS Microbiol Rev* **46**, (2022).

4. Partridge, S. R., Kwong, S. M., Firth, N. & Jensen, S. O. Mobile genetic elements associated with antimicrobial resistance. *Clinical Microbiology Reviews* vol. 31 Preprint at https://doi.org/10.1128/CMR.00088-17 (2018).

5. Vogwill, T. & Maclean, R. C. The genetic basis of the fitness costs of antimicrobial resistance: A meta-analysis approach. *Evol Appl* **8**, (2015).

6. Siedentop, B., Rüegg, D., Bonhoeffer, S. & Chabas, H. My host's enemy is my enemy: plasmids carrying CRISPR-Cas as a defence against phages. *Proceedings of the Royal Society B: Biological Sciences* **291**, (2024).

7. Picton, D. M. *et al.* The phage defence island of a multidrug resistant plasmid uses both BREX and type IV restriction for complementary protection from viruses. *Nucleic Acids Res* **49**, (2021).

8. Vassallo, C. N., Doering, C. R., Littlehale, M. L., Teodoro, G. I. C. & Laub, M. T. A functional selection reveals previously undetected anti-phage defence systems in the E. coli pangenome. *Nat Microbiol* **7**, (2022).

9. Leroux, M. & Laub, M. T. Toxin-Antitoxin Systems as Phage Defense Elements. *Annual Review of Microbiology* vol. 76 Preprint at https://doi.org/10.1146/annurev-micro-020722-013730 (2022).

10. Wales, A. D. & Davies, R. H. Co-selection of resistance to antibiotics, biocides and heavy metals, and its relevance to foodborne pathogens. *Antibiotics* vol. 4 Preprint at https://doi.org/10.3390/antibiotics4040567 (2015).

11. Pal, C., Bengtsson-Palme, J., Kristiansson, E. & Larsson, D. G. J. Co-occurrence of resistance genes to antibiotics, biocides and metals reveals novel insights into their co-selection potential. *BMC Genomics* **16**, (2015).

12. Mahata, T. *et al.* Gamma-Mobile-Trio systems define a new class of mobile elements rich in bacterial defensive and offensive tools. *bioRxiv* (2024).

13. Botelho, J. Defense systems are pervasive across chromosomally integrated mobile genetic elements and are inversely correlated to virulence and antimicrobial resistance. *Nucleic Acids Res* **51**, (2023).

14. LeGault, K. N. *et al.* Temporal shifts in antibiotic resistance elements govern phage-pathogen conflicts. *Science (1979)* **373**, (2021).

309   15.   BiomX, Inc. BiomX Announces Positive Topline Results from Phase 2 Trial Evaluating
310   BX211 for the Treatment of Diabetic Foot Osteomyelitis (DFO).
311   https://ir.biomx.com/news-events/press-releases/detail/130/biomx-announces-positive-
312   topline-results-from-phase-2-trial.

313   16.   Chan, B. K. *et al.* Personalized inhaled bacteriophage therapy for treatment of
314   multidrug-resistant Pseudomonas aeruginosa in cystic fibrosis. *Nat Med* **31**, 1494–1501
315   (2025).

316   17.   Matlock, W. *et al.* Enterobacterales plasmid sharing amongst human bloodstream
317   infections, livestock, wastewater, and waterway niches in Oxfordshire, UK. *Elife* **12**,
318   (2023).

319   18.   Shaw, L. P. *et al.* Niche and local geography shape the pangenome of wastewater-and
320   livestock-associated Enterobacteriaceae. *Sci Adv* **7**, (2021).

321   19.   Lipworth, S. *et al.* Ten-year longitudinal molecular epidemiology study of Escherichia
322   coli and Klebsiella species bloodstream infections in Oxfordshire, UK. *Genome Med* **13**,
323   (2021).

324   20.   Tesson, F. *et al.* Systematic and quantitative view of the antiviral arsenal of prokaryotes.
325   *Nat Commun* **13**, (2022).

326   21.   Néron, B. *et al.* MacSyFinder v2: Improved modelling and search engine to identify
327   molecular systems in genomes. *Peer Community Journal* **3**, (2023).

328   22.   Tesson, F. *et al.* A Comprehensive Resource for Exploring Antiphage Defense:
329   DefenseFinder Webservice,Wiki and Databases. *Peer Community Journal* **4**, (2024).

330   23.   Feldgarden, M. *et al.* AMRFinderPlus and the Reference Gene Catalog facilitate
331   examination of the genomic links among antimicrobial resistance, stress response, and
332   virulence. *Sci Rep* **11**, (2021).

333   24.   Robertson, J. & Nash, J. H. E. MOB-suite: software tools for clustering, reconstruction
334   and typing of plasmids from draft assemblies. *Microb Genom* **4**, (2018).

335   25.   Néron, B. *et al.* IntegronFinder 2.0: Identification and Analysis of Integrons across
336   Bacteria, with a Focus on Antibiotic Resistance in Klebsiella. *Microorganisms* **10**,
337   (2022).

338   26.   Hyatt, D. *et al.* Prodigal: Prokaryotic gene recognition and translation initiation site
339   identification. *BMC Bioinformatics* **11**, (2010).

340   27.   Pedersen, E. J., Miller, D. L., Simpson, G. L. & Ross, N. Hierarchical generalized
341   additive models in ecology: An introduction with mgcv. *PeerJ* **2019**, (2019).

342   28.   Redondo-Salvo, S. *et al.* Pathways for horizontal gene transfer in bacteria revealed by a
343   global map of their plasmids. *Nat Commun* **11**, (2020).

344   29.   Kim, P. J. & Price, N. D. Genetic co-occurrence network across sequenced microbes.
345   *PLoS Comput Biol* **7**, (2011).

346   30.   San Millan, A., Escudero, J. A., Gifford, D. R., Mazel, D. & MacLean, R. C. Multicopy
347   plasmids potentiate the evolution of antibiotic resistance in bacteria. *Nat Ecol Evol* **1**,
348   0010 (2016).

31.   Matlock, W. *et al.* E. coli phylogeny drives co-amoxiclav resistance through variable
expression of blaTEM-1. Preprint at https://doi.org/10.1101/2024.08.12.607562 (2024).

32.   Deep, A., Liang, Q., Enustun, E., Pogliano, J. & Corbett, K. D. Architecture and
infection-sensing mechanism of the bacterial PARIS defense system. (2024)
doi:10.1101/2024.01.02.573835.

33.   Koga, M., Otsuka, Y., Lemire, S. & Yonesaki, T. Escherichia coli rnlA and rnlB
compose a novel toxin-antitoxin system. *Genetics* **187**, (2011).

34.   Pecota, D. C. & Wood, T. K. Exclusion of T4 phage by the hok/sok killer locus from
plasmid R1. *J Bacteriol* **178**, (1996).

35.   Hall, A. R., Scanlan, P. D., Morgan, A. D. & Buckling, A. Host-parasite coevolutionary
arms races give way to fluctuating selection. *Ecol Lett* **14**, 635–642 (2011).

36.   Betts, A., Gifford, D. R., MacLean, R. C. & King, K. C. Parasite diversity drives rapid
host dynamics and evolution of resistance in a bacteria-phage system. *Evolution (N Y)*
70, 969–978 (2016).

37.   Montelongo Hernandez, C., Putonti, C. & Wolfe, A. J. Urinary Plasmids Reduce
Permissivity to Coliphage Infection. *Microbiol Spectr* **11**, (2023).

365