## *Title:* **Physical exercise and motor learning: A scoping review**

*Running title:* **Physical exercise and motor learning**

*Authors:* Layale Youssef[1,2,3#], Amanda O'Farrell[1,2,3], Denis Arvisais[4], Jason L. Neva[1,2,3]*,

Benjamin Pageaux[1,2,3]*

[1] École de kinésiologie et des sciences de l'activité physique (EKSAP), Faculté de médecine, Université de Montréal, Montréal, Québec, Canada.

[2] Centre de Recherche de l'Institut Universitaire de Gériatrie de Montréal (CRIUGM), Montréal, Québec, Canada

[3] Centre Interdisciplinaire de Recherche sur le Cerveau et l'Apprentissage (CIRCA), Montréal, Québec, Canada

[4] Bibliothèques des Sciences de la Santé, Université de Montréal, Québec, Canada.

* Both authors contributed equally

# *Corresponding Author:*
Layale Youssef
4545 Queen Mary Road, Montreal, QC, Canada
layale.youssef@umontreal.ca

## **Abstract**

Physical exercise, as an adjunct to motor task practice, can enhance motor learning by inducing neurophysiological changes known to facilitate this process. This preregistered scoping review mapped current knowledge on physical exercise's effect on motor learning. The search strategy used indexed terms and keywords across several databases: MEDLINE, EMBASE, SPORTDiscus, Web of Science, PsycINFO, CINAHL Complete, ERIC and Dissertations & Theses Global (ProQuest). A total of 66 sources were included, comprising 62 experimental studies and 4 review articles. Most studies involved healthy populations (83%), while fewer focused on clinical populations (17%). Aerobic exercise was the predominant type used, with lower-limb cycling the most common approach (73%). Among various intensity levels, high-intensity exercise was most frequently investigated; however, in some studies, reported exercise intensity did not accurately reflect the method prescribed (6%). In contrast, resistance exercise was rarely examined (0.6%), highlighting a notable gap in existing literature. While some studies assessed both motor skill acquisition and a retention test (55%), others only employed a retention test (45%). Motor learning was measured at different time points including short-term or long-term delayed retention tests, or both. This underscores the need for consistent inclusion of long-term delayed retention tests to better capture lasting effects of physical exercise on motor learning. This review highlights gaps in the literature, including underrepresentation of clinical populations, inaccurate reporting of exercise intensity, scarce research on resistance exercise, and limited assessment of long-term retention tests, all of which may affect interpretation of the effect of physical exercise on motor learning.

**Keywords: aerobic exercise, resistance exercise, motor skill acquisition, retention test, performance**

## Key points

- This scoping review provides perspectives for future research on the effects of physical exercise on motor learning.
- Despite the crucial role of motor learning for clinical populations, a smaller proportion of these population groups was evaluated compared to healthy populations.
- Lower-limb cycling was the most commonly used physical exercise.
- High-intensity exercise was the most frequently studied intensity level.
- Resistance exercise was underrepresented in the field.
- Accurate prescription of exercise intensity is essential to ensure reliable findings.
- Future research should more consistently use longer-term retention tests to better assess the lasting effects of physical exercise on motor learning.

## 1. Introduction

Although the terms "physical activity" and "physical exercise" are sometimes used in similar contexts, they refer to different types of physical movements [1]. Physical activity is defined as any body movement produced by skeletal muscles resulting in energy expenditure [2]. In contrast, physical exercise consists of performing planned and structured physical activity to enhance physical fitness or maintain overall health, and can be either chronic or acute [3]. There are different types of physical exercise (e.g., aerobic or resistance), such as running, cycling, or swimming for aerobic exercise, and weightlifting or bodyweight exercises for resistance exercise, which could involve different modes of muscular contractions (i.e., concentric, eccentric, or isometric). Aerobic exercise primarily targets the cardiovascular system and metabolic health, leading to improvement in endurance performance. Resistance exercise primarily targets the neuromuscular system and is essential for developing muscular strength and power. Both aerobic exercise and resistance exercise are important for maintaining functional independence. Moreover, physical exercise intensity may vary (e.g., low, moderate, and high), and the methodology used to prescribe exercise intensity is different among studies. Importantly, physical exercise is not only beneficial for maintaining physical function and overall health, as it has also been shown to exert beneficial effects on brain health [4, 5].

Recently, numerous studies in the field of exercise science have highlighted how external interventions, such as physical exercise, can enhance nervous system function [6]. More specifically, physical exercise has been shown to induce significant improvements in cognitive function [5], cognitive performance [7] and memory [8]. Furthermore, physical exercise has been associated with changes in neuroplasticity [9-13]. Interestingly, neuroplasticity changes, including changes in motor cortex excitability, have been observed during the process of motor learning [14-16]. These observations suggest that physical exercise may act as a priming mechanism for motor learning by facilitating these neuroplasticity changes. Consequently, incorporating physical exercise as an adjunct intervention could be an effective strategy to enhance motor learning [17, 18].

A motor skill refers to the ability to perform a goal-directed movement that enables an individual to achieve a specific outcome with accuracy and efficiency, using minimal time and effort [19]. Although often used interchangeably, the terms motor performance and motor learning represent distinct concepts. Motor performance refers to the observable execution of a voluntary action, whereas motor learning involves practice-driven processes that lead to relatively lasting improvements in movement ability [20]. Motor learning consists of two fundamental aspects: skill acquisition and skill maintenance. Skill acquisition involves developing movement control through practice, and skill maintenance ensures long-

term skill retention and adaptability of the learned skill under different contexts [21]. To study these

processes, several motor learning categories have been established, including sequence learning, motor

adaptation, motor acuity, and associative learning (for more details, see Figure S1 in supplementary

material 1).

Despite the well-established benefits of physical exercise on physical function and cognition [5, 7,

22], its impact on motor learning remains unclear, with divergent findings contributing to this uncertainty.

For example, some studies found that an acute bout of aerobic exercise only improves the acquisition of

motor skills [23, 24] whereas other studies showed improvements in both motor acquisition and motor skill

retention [25-27]. Another group of studies found an enhancement only in motor skill retention [28-31]. In

contrast, some studies have found no effect on both skill acquisition and retention [32-37]. Interestingly,

the literature includes a wide range of exercise modalities (e.g., continuous and interval training), types

(e.g., aerobic and resistance), and intensities (e.g., low, moderate, and high), as well as varied delays in

timing of retention tests (i.e., ranging from hours to days), all of which may contribute to the divergent

effects of physical exercise on motor learning. Also, within aerobic exercise, multiple forms have been

used, including lower-limb cycling, running, walking, and upper-limb cycling. This diversity in exercise

types and protocols adds to the variability observed across studies and underscores the need for careful

comparison when interpreting results.

Considering these discrepancies, mapping the existing literature is crucial to better understand the

effect of physical exercise on motor learning. Understanding how different types and intensities of physical

exercise could impact motor learning is crucial for developing targeted exercise protocols that address the

specific needs of various populations, whether clinical populations in rehabilitation settings or healthy

individuals in sport. To optimize the effects of physical exercise on motor learning, it is important to

recognize that these populations have distinct needs, and therefore, tailored exercise protocols may be

necessary to enhance motor learning.

This scoping review aims to explore and map the existing literature on the impact of physical

exercise on motor learning in humans. It examines the various populations studied, the types of physical

exercise implemented, and the timing of exercise interventions, specifically whether they were performed

before or after motor task practice. Additionally, this scoping review will also identify the different motor

learning categories used across studies.

## 2. Methods

This scoping review was pre-registered on the Open Science Framework (OSF) Website (https://osf.io/4mv53) and was conducted according to the JBI scoping review guidelines and recommendations [38].

### 2.1 Search strategy

The database search was conducted by a librarian (DA) in collaboration with LY and consultation with BP and JN. The search strategy included a combination of keywords and terms indexed across various databases, including MEDLINE (Ovid), EMBASE (Ovid), PsycINFO (Ovid), SportDiscus with Full Text (Ebsco), CINAHL Complete (Ebsco), ERIC (ProQuest), Web of Science (Clarivate), and Dissertations & Theses Global (ProQuest). The complete search equations are provided in supplementary material 2. The search process was conducted in February 2024 and updated in March 2025. The second search led to the inclusion of two additional experimental studies. The search was limited to articles in English and French, with no restrictions on the publication year. All identified sources were uploaded in Endnote (Clarivate Analytics, PA, USA) and duplicates were removed. The screening of all sources was then managed using Covidence software [39] and followed three steps: (1) remaining duplicates removal, (2) title and abstract screening, and (3) full-text evaluation. Two researchers (LY and AOF) independently reviewed articles at each stage, and any disagreements were resolved through discussion with a third researcher (BP). A detailed flowchart outlining the selection process of sources can be found in Figure 1.

### 2.2 Inclusion Criteria

All fully published, peer-reviewed articles evaluating the effect of physical exercise on motor learning were included. Both experimental studies and reviews that incorporated experimental studies were considered. The primary focus was the impact of physical exercise on motor learning, assessed in either laboratory or field settings. Studies were included if they involved clearly defined motor tasks with assessments of skill acquisition, with or without retention tests. However, following the classical definition of motor learning [20], skill maintenance over time was not considered in studies where a retention test was not included. Furthermore, all types of physical exercise interventions (aerobic or resistance) at any intensity level (low, moderate, high) were included. Finally, the review included studies involving all types of human populations (healthy and clinical) of all ages.

164      **Fig . 1 Flowchart illustrating the selection process of sources.**

**2.3 Data extraction**

The extracted data included details about the evaluated populations, the physical exercise interventions, the motor learning parameters and additional key findings relevant to the objectives of this scoping review. Data extraction was carried out by two independent reviewers (LY and AOF), with any disagreements resolved through discussion. If consensus could not be reached, JN and BP were consulted to provide a final resolution. A data extraction table was used to record the information from the included studies. The initial table was tested on two articles by LY and AOF, then slightly refined before reaching its final version presented in supplementary material 3. For the 62 experimental studies reviewed, the following data were collected: population characteristics (age and type), details of the physical exercise performed (type, duration, intensity) based on American College of Sports Medicine (ACSM) guidelines [40], timing of exercise relative to motor task practice (before or after acquisition), motor learning category (sequence, motor acuity, motor adaptation, associative learning) as well as the result as reported in each study (i.e, effect of physical exercise on skill acquisition and/or skill retention).

Although we followed the pre-registration for the review, two minor deviations were made during the data extraction process to improve methodological clarity. First, we introduced an additional distinction between studies regarding the methodology for exercise prescription. We decided to dissociate prescriptions based on measured versus estimated parameters (e.g., estimated vs measured maximal heart rate). This refinement was not initially planned but was added during data extraction to better capture variability in exercise prescription methods across studies. Second, regarding the motor learning categories [21], we originally intended to use predefined categories such as sequence learning and adaptation learning. However, after examining the motor tasks used in the included studies, we found that these predefined categories did not fully capture the distinct characteristics of the motor learning constructs investigated. Therefore, we refined our categorization to better reflect the outcomes being evaluated. The categorization used is detailed in Figure S1 of supplementary material 1.

In this scoping review, the number of experimental conditions exceeds the number of studies, as several studies included multiple experimental conditions. For example, some studies involved different exercise intensities or different populations. As a result, the mapping of the literature was also conducted based on the number of experimental conditions relevant to each research question. In the manuscript, $n$ refers to a subtotal of the number of participants, $k$ refers to a subtotal of the number of studies, $g$ refers to a subtotal of the number of groups, and $e$ refers to a subtotal of the number of experimental conditions.

When these letters are capitalized, it refers to the grand total (e.g., N = the total number of participants
including all studies and all experimental conditions).

*Motor learning categorization and results*

To classify the motor learning categories used across studies, we identified four main categories
based on task characteristics and learning objectives [21]. *Sequence learning*, often referred to as
sequence-specific learning, was defined for motor tasks that involve comparing performance on a
repeated sequence or sequences with that of random sequences (e.g., serial reaction time task). This
comparison is necessary to identify learning that is specific to the practiced sequence rather than general
improvements in other aspects of the motor skill. However, if only a repeated sequence or sequences
were practiced without including random sequences for comparison, the motor task was categorized as
motor acuity. *Motor acuity* refers to improvements in the accuracy and/or precision of performing a
particular movement or sequence of movements through practice (e.g., Rho task). In this case, the focus
is on refining the ability to execute the motor task more precisely and/or accurately. *Motor adaptation*
included tasks in which participants adjusted their movements in response to external perturbation (e.g.,
visuomotor rotation task). Finally, *associative learning* referred to tasks requiring participants to form
arbitrary associations between stimuli and responses (e.g., conditional learning task). The primary
objective of each category is shown in Figure S1 of supplementary material 1. This classification provides
a clear view of the distribution of motor learning categories across the included studies and the
identification of gaps in literature.

To provide a comprehensive understanding of the effect of physical exercise on motor learning,
we examined skill acquisition and skill retention separately based on when physical exercise was
performed relative to the motor task. The results for both skill acquisition and skill retention were
reported in studies that performed physical exercise prior to the motor task, while only results from
retention tests were reported in studies that conducted physical exercise after the motor task.
Furthermore, the timing of retention tests varied across studies. To address this potential challenge in the
synthesis of findings, we categorized the reported results from retention tests into three time windows:
before 24 hours, at 24 hours, and after 24 hours. Consequently, findings from some studies appear in
multiple retention time windows. Results were categorized based on exercise intensity and separated
according to population type. However, due to the complexity of intervention types and timing, all forms
of physical exercise were grouped together in the outcome of this scoping review, regardless of when the

exercise was performed. To accurately assess the effects of physical exercise on motor learning, a

comparison with a control condition (i.e., without physical exercise) is essential and was included in the

majority of the studies. Based on the results reported by the authors in the experimental studies, motor

learning effects were categorized as enhanced, no difference, or attenuated performance. Although

studies without a control condition were included in the scoping review and their data were extracted,

their findings were not considered in the outcome of this scoping review.

## 235   3. Results

### 236   3.1 Included studies

Following the systematic search in various databases, 66 sources were included (see Figure 1).

Following the JBI guidelines for categorization, 62 primary sources were experimental studies and 4

secondary sources were review articles [1 narrative review [17], 2 systematic reviews [41, 42], and 1

systematic review with meta-analysis [18]]. A timeline of the included sources is presented in Figure 2a.

The research interest on the effect of physical exercise on motor learning first appeared as early as 1970,

but after a long gap, studies in this area began to emerge more consistently starting in 2009, reaching a

peak in 2023 with 13 studies published that year. A world map illustrating the geographical distribution

of the included studies is presented in Figure 2b. This scoping review revealed a worldwide interest in

the effects of physical exercise on motor learning, with most studies conducted in Canada (k = 16), the

United States (k = 11), Germany (k = 10) and Denmark (k = 6).

We extracted data from 62 experimental studies to address our specific research questions. The

following sections address our review questions, detailing the populations studied, the types of physical

exercise implemented, the motor learning parameters investigated, and the timing of physical exercise in

relation to motor task practice. An overview of the main characteristics of all experimental studies is

presented in Table 1. Detailed characteristics of all experimental studies are provided in Table S1 of

supplementary material 1.

**Table 1.** Characteristics of the included studies related to the review questions

| Study | (1) What types of population were investigated? | (2) What types of physical exercise were performed to impact motor learning? (Type, Intensity) | | (3) What types of motor learning categories were used? | (4) When was the physical exercise performed relative to motor task practice? |
|---|---|---|---|---|---|
| **Andrews et al. (2023) [43]** | Huntington's diseases | Lower-limb cycling | Moderate | Motor acuity | Before |
| **Andrushko et al. (2023) [44]** | Stroke | Lower-limb cycling | High | Sequence | Before |
| **Angulo-Barroso et al. (2019) [45]** | Healthy children | Running | High | Motor adaptation | Before |
| **Baird et al. (2018) [32]** | Healthy adults | Lower-limb cycling | Low and high | Sequence | Before |
| **Bartz and Smith (1970) [46]** | Healthy adults | Walking | Moderate | Motor adaptation | Before |
| **Bonuzzi et al. (2023) [47]** | Stroke and healthy adults | Lower-limb cycling | Moderate | Sequence | Before and after |
| **Bosch et al. (2020) [48]** | Healthy adults | Lower-limb cycling | Moderate and high | Sequence | After |
| **Chan et al. (2023) [49]** | Parkinson's disease | Lower-limb cycling | High | Sequence | After |
| **Charalambous et al. (2018) [50]** | Stroke | Treadmill walking and total body exercise | High | Motor adaptation | Before and after |
| **Charalambous et al. (2019) [33]** | Healthy adults | Whole-body cycling | High | Motor adaptation | Before |
| **Chen et al. (2023) [51]** | Healthy adults | Treadmill running | Moderate | Motor acuity | Before |
| **Coco et al. (2016) [52]** | Healthy adults | Lower-limb cycling | High | Motor acuity | After |
| **Cristini et al. (2023) [53]** | Healthy adults | Lower-limb cycling | High | Sequence | After |
| **Dal Maso et al. (2018) [54]** | Healthy adults | Lower-limb cycling | High | Motor acuity | After |
| **Duchesne et al. (2016) [55]** | Parkinson's disease and healthy older adults | Lower-limb cycling | High | Sequence | After |
| **Ferrer-Uris et al. (2017) [28]** | Healthy adults | Shuttle run | High | Motor adaptation | Before and after |
| **Ferrer-Uris et al. (2018) [56]** | Healthy children | Shuttle run | High | Motor adaptation | Before and after |
| **Greeley et al. (2021) [57]** | Healthy older adults | Lower-limb cycling | High | Sequence | Before |
| **Helm et al. (2017) [58]** | Healthy adults | Upper-limb cycling | High | Motor adaptation | Before |
| **Holman and Staines (2021) [59]** | Healthy adults | Lower-limb cycling | High | Motor adaptation | After |
| **Hubner et al. (2018) [60]** | Healthy older adults | Lower-limb cycling | Moderate | Motor acuity | Before |
| **Hung et al. (2021) [61]** | Healthy adults | Lower-limb cycling | High | Motor acuity | After |
| **James and Wang (2023) [34]** | Healthy adults | Lower-limb cycling | Moderate | Motor adaptation | Before |
| **Jespersen et al. (2023) [29]** | Healthy adults | Lower-limb cycling | Moderate and high | Sequence | Before and after |
| **Khan et al. (2022) [62]** | Healthy adults | Lower-limb cycling | High | Motor acuity | After |
| **Kuo et al. (2023a) [63]** | Healthy adults | Lower-limb cycling | Low, moderate and high | Sequence | After |
| **Kuo et al. (2023b) [64]** | Healthy adults | Lower-limb cycling | Moderate | Sequence | After |
| **Lehmann et al. (2020) [27]** | Healthy adults | Lower-limb cycling | High | Motor adaptation | Before |
| **Loras et al. (2020) [65]** | Healthy adults | Lower-limb cycling | Low and moderate | Motor acuity | Before |
| **Lundbye-Jensen et al. (2017) [66]** | Healthy children | Floorball and running | High | Motor acuity | After |
| **Mang et al. (2014) [25]** | Healthy adults | Lower-limb cycling | High | Sequence | Before |
| **Mang et al. (2016) [30]** | Healthy adults | Lower-limb cycling | High | Sequence | Before |
| **Munz et al. (2021) [67]** | Healthy children | Lower-limb cycling | High | Motor acuity | After |
| **Nepveu et al. (2017) [68]** | Stroke | Lower-limb cycling | High | Motor acuity | After |

258    **Table 1.** *Continued*

| Study | (1) What types of population were investigated? | (2) What types of physical exercise were performed to impact motor learning? (Type, Intensity) | | (3) What types of motor learning categories were used? | (4) When was the physical exercise performed relative to motor task practice? |
|---|---|---|---|---|---|
| **Neva et al. (2019) [26]** | Healthy adults | Lower-limb cycling | Moderate | Motor adaptation | Before |
| **Ostadan et al. (2016) [69]** | Healthy adults | Lower-limb cycling | High | Sequence | After |
| **Opie and Semmler (2019) [70]** | Healthy adults | Lower-limb cycling | Moderate and high | Motor acuity | Before |
| **Perini et al. (2016) [71]** | Healthy adults | Lower-limb cycling | Moderate | Motor acuity | Before |
| **Pixa et al. (2021) [35]** | Healthy adults | Lower-limb cycling | Low and high | Motor acuity | Before |
| **Quaney et al. (2009) [72]** | Stroke | Lower-limb cycling | Moderate | Sequence | Before |
| **Quinlan et al. (2021) [36]** | Healthy adults | Lower-limb cycling | High | Motor acuity | Before |
| **Rhee et al. (2016) [73]** | Healthy adults | Lower-limb cycling | High | Motor acuity | After |
| **Roig et al. (2012) [74]** | Healthy adults | Lower-limb cycling | High | Motor acuity | Before and after |
| **Roig-Hierro and Batalla (2023) [75]** | Healthy adults | Shuttle run | High | Motor acuity | After |
| **Singh et al. (2016) [76]** | Healthy adults | Lower-limb cycling | Moderate | Motor adaptation | Before |
| **Skriver et al. (2014) [77]** | Healthy adults | Lower-limb cycling | High | Motor acuity | Before |
| **Snow et al. (2016) [23]** | Healthy adults | Lower-limb cycling | Moderate | Motor acuity | Before |
| **Statton et al. (2015) [24]** | Healthy adults | Running | High | Motor acuity | Before |
| **Stavrinos and Coxon (2017) [31]** | Healthy adults | Lower-limb cycling | High | Motor acuity | Before |
| **Steib et al. (2018) [78]** | Parkinson's disease | Lower-limb cycling | Moderate | Motor adaptation | Before |
| **Stranda et al. (2019) [79]** | Healthy adults | Lower-limb cycling | Moderate | Motor acuity | Before |
| **Swarbrick et al. (2020) [80]** | Healthy adults | Lower-limb cycling | Low and high | Motor acuity | Before |
| **Taylor et al. (2024) [81]** | Healthy older adults | Lower-limb cycling | High | Motor acuity | Before |
| **Thomas et al. (2016) [82]** | Healthy adults | Lower-limb cycling | Low and high | Motor acuity | After |
| **Thomas et al. (2017) [83]** | Healthy adults | Strength training, Circuit training and Hockey training | High | Motor acuity | After |
| **Thompson et al. (2023) [84]** | Stroke | Whole-body cycling | High | Motor adaptation | After |
| **Tomporowski and Pendleton (2018) [85]** | Healthy adults | Dancing | Moderate | Motor acuity | Before and after |
| **Wanga et al. (2023) [86]** | Healthy adults | Lower-limb cycling | Low and high | Sequence | After |
| **Wanner et al. (2020) [37]** | Healthy adults | Lower-limb cycling | Low, moderate, and high | Motor adaptation | Before |
| **Wanner et al. (2021) [87]** | Parkinson's disease | Lower-limb cycling | Moderate | Motor adaptation | Before |
| **Williams et al. (1985) [88]** | Healthy adults | Upper-limb cycling | Low and high | Motor acuity | Before |
| **Yildirim et al. (2024) [89]** | Healthy adults | Running | Low and moderate | Motor acuity | Before and after |

262
263
**Fig. 2 Temporal and geographical distribution of included sources.**
(Panel a) Temporal distribution of all included sources. Orange bars represent experimental studies, while blue gradient bars represent other types of sources (narrative review; systematic review; systematic review with meta-analysis). (Panel b) Geographical distribution of all included sources. The format used is the following: total number [other types of sources*]. The total number of sources is shown first; when sources other than experimental studies are included, their number is indicated in brackets. Darker colors indicate a higher number of sources reported from each country. n refers to the number of participants reported from experimental studies conducted in that country.

**3.2 What types of populations were investigated?**

The majority of included studies investigated healthy populations (g = 53), while only a small portion of studies (g = 11) focused on clinical populations (see Figure 3). Among the studies including healthy populations, 45 studies involved young healthy adults (studies are listed in Table 1). Research involving other age groups was limited, with only 4 studies investigating healthy children [45, 56, 66, 67] and 4 studies focusing on healthy older adults [57, 60, 81, 90]. Within the studies investigating clinical populations, three neurological conditions were identified with 6 studies investigating individuals with stroke [44, 47, 50, 68, 72, 84], followed by 4 studies investigating individuals with Parkinson's disease [49, 78, 87, 90]. Additionally, a single study focused on individuals with Huntington's disease [43].

**Fig. 3 Types of populations evaluated across all included experimental studies.**
Distribution of the 1,479 participants reported across the 62 experimental studies. Participants were categorized as either healthy or clinical populations, with each category further divided into subcategories. Green gradients represent healthy populations, and orange gradients represent clinical populations. k refers to the number of experimental studies, G refers to the overall number of reported groups, g refers to the number of reported subgroups.

**3.3 What types and intensities of physical exercise were used?**

Among the 62 experimental studies, a total of 81 experimental conditions were conducted (E =
81). Interestingly, the vast majority involved aerobic exercise (e = 79), while only a small number focused
on resistance exercise (e = 2) (see Figure 4a). Of the aerobic exercise conditions, lower-limb cycling was
the main type of exercise used (e = 59). Other aerobic exercise modalities included running [24, 28, 45,
51, 56, 66, 75, 89], walking [46, 50], upper-limb cycling [58, 88], combined upper and lower-limb
cycling (whole-body cycling) [33, 84], and whole-body aerobic exercise [50]. Additionally, a small
number of studies employed less common physical exercise, including floorball [66], dancing [85], and
hockey [83].

Of the 62 experimental studies included, 34 prescribed exercise intensity based on measured
physiological parameters reflecting individual capacity while 28 relied on estimated parameters (see
Figure 4b). No study used the perception of effort or other perceptual responses to prescribe the exercise
intensity. According to ACSM guidelines, the 81 experimental conditions were further classified as high
intensity (e = 48), moderate intensity (e = 23), and low intensity (e = 10) (see Figure 4c). When combining
exercise type and intensity, lower-limb cycling at high intensity was the most commonly used
combination, followed by lower-limb cycling at moderate intensity (see Figure 4d).

Among the 81 experimental conditions, five showed discrepancies between the reported and
actual exercise intensities, as defined by the ACSM guidelines. Specifically, two conditions were
reported as moderate intensity, but the prescribed workloads—70% age-predicted heart rate reserve [59]
and 65-85% age-predicted maximal heart rate with a reported average of 84% maximal heart rate [24]—
fall within the high-intensity range. In another study, two conditions were labeled as moderate and high
intensity, but the prescriptions of 50% age-predicted maximal heart rate and 75% age-predicted maximal
heart rate respectively, actually correspond to low and moderate intensity based on ACSM guidelines
[65]. Also, one condition was described as low intensity, although the exercise was performed at 50%
age-predicted heart rate reserve, which is interpreted as moderate intensity (Opie & Semmler, 2019).

346
347

348
349

**Fig. 4 Types and intensities of physical exercise reported across all included experimental studies.**
(Panel a) Distribution of the 81 experimental conditions according to exercise type, categorized as aerobic or resistance. Aerobic exercise is further subcategorized by specific types. (Panel b) Prescription of exercise intensities across the experimental studies, pink refers to measured capacity and turquoise refers to estimated capacity. Pink and turquoise gradients refer to measured and estimated exersie parameters respectively. (Panel c) Exercise intensities used across all experiments where red refers to high intensity, yellow refers to moderate intensity and green refers to low intensity. (Panel d) Heatmap showing the combination of exercise type and intensity where darker purple indicates a higher number of experimental conditions using the specific combination. Others refers to all exercise types different from lower-limb cycling, running and walking. E refers to the total number of experimental conditions reported across the 62 included experimental studies, e refers to the number of experimental conditions per exercise type category and n refers to the number of participants.

**3.4 Which categories of motor learning were investigated, and when was physical exercise**
**performed relative to motor task practice?**

Three motor learning categories were used across the 62 included experimental studies (see
Figure 5a): 30 involved motor acuity, 16 focused on sequence learning, and 16 addressed motor
adaptation; no study investigated associative learning (studies are listed in Table 1).

Among the 73 experimental conditions related to the timing of the exercise performed (E = 73),
42 involved physical exercise performed before motor task practice and 31 after (see Figure 5b). In
healthy adults (e = 51), 30 conditions implemented exercise before motor task practice and 21 after.
Among older healthy adults (e = 4), three conditions performed exercise before and one after motor task
practice. In healthy children (e = 6), two conditions included exercise before and four after motor task
practice. In individuals with stroke (e = 7), four conditions involved exercise before and three after motor
task practice. In individuals with Parkinson's disease (e = 4), two conditions implemented exercise before
and two after motor task practice. Finally, the single condition involving individuals with Huntington's
disease (e = 1) performed exercise before motor task practice (see Figure 5c).

**Fig. 5 Motor learning categories and timing of exercise relative to motor task practice.**
(Panel a) Distribution of motor learning categories evaluated across the 62 experimental studies. (Panel b) Timing of exercise in relation to motor task practice across the 73 experimental conditions, (Panel c) Timing of exercise in relation to motor task practice across the 73 experimental conditions shown by population type. Light blue indicates exercise performed before the motor task practice, and light red indicates exercise performed after the motor task practice. Before = exercise performed prior to motor task practice; After = exercise performed after the motor task practice; n refers to the number of participants.

**3.5 Effects of physical exercise on motor learning**

The effects of physical exercise on motor learning, as reported by the authors in each included

experimental study, were categorized based on acquisition and retention tests (enhanced, no difference,

or attenuated) relative to a control condition. Results for the retention tests were separated into three time

windows: less than 24 hours, at 24 hours, and more than 24 hours. These results were categorized by

exercise intensity (high, moderate and low) and by population type (see Figure 6). Detailed results for

each study are also provided in Table S1 of supplementary material 1.

*3.5.1 Skill acquisition*

A total of 49 experimental conditions (E = 49) assessed the effects of physical exercise on motor

skill acquisition (see Figure 6a). In healthy children (e = 3; all high intensity), no significant effects were

observed. In healthy adults (e = 35), the majority of experimental conditions showed no effect (e = 28;

across low, moderate, and high intensities), while six conditions reported enhancement (e = 6; three with

high intensity, three with moderate intensity), and one moderate-intensity condition led to attenuated

performance. In healthy older adults (e = 3), two conditions showed no effect (high and moderate

intensity), and one high-intensity condition led to attenuated performance. Among individuals with

stroke (e = 4), one moderate-intensity condition enhanced acquisition, while the others (two high

intensity and one moderate intensity) showed no effect. In individuals with Parkinson's disease, three

conditions (e = 3; 2 moderate and 1 high intensity) reported no difference. Finally, the single

experimental condition involving individuals with Huntington's disease (e = 1) reported improvement

following a moderate-intensity exercise.

*3.5.2 Retention tests less than 24h*

A total of 28 experimental conditions (E = 28) assessed the effects of physical exercise on

performance within 24 hours following the exercise intervention (see Figure 6b). In healthy children (e

= 3), two experimental conditions showed enhanced performance (both at high intensity), and one

condition at high intensity showed no difference. In healthy adults (e = 23), five conditions demonstrated

enhanced performance (three at high intensity, two at moderate intensity), 17 showed no difference

(across all three intensities), and one high-intensity condition showed attenuated performance at

retention. In healthy older adults (e = 2), one condition showed enhanced performance (high intensity)

and one showed no difference (moderate intensity). Notably, no experimental conditions involving

clinical populations were reported in this retention time category.

*3.5.3 Retention tests at 24h*

A total of 49 experimental conditions (E = 49) assessed the effects of physical exercise on

performance at 24 hours following the exercise intervention (see Figure 6c). In healthy children (e = 4),

three conditions showed enhanced performance (all at high intensity), and one condition showed

attenuated performance (high intensity). In healthy adults (e = 36), 12 conditions demonstrated enhanced

performance (nine at high intensity, two at moderate intensity, and one at low intensity), while 24

conditions showed no difference (across all intensities). In healthy older adults (e = 2), both conditions

showed no difference (one at high intensity and one at moderate intensity). Among individuals with

stroke (e = 4), one condition showed enhanced performance (high intensity), and three conditions showed

no difference (all at high intensity). In individuals with Parkinson's disease (e = 3), one condition showed

enhanced performance (moderate intensity), and two conditions showed no difference (one at high

intensity and one at moderate intensity).

*3.5.4 Retention tests more than 24h*

A total of 31 experimental conditions (E = 31) assessed the effects of physical exercise on

performance beyond 24 hours following the exercise intervention (see Figure 6d). In healthy children (e

= 3), all conditions showed enhanced performance (all at high intensity). In healthy adults (e = 21),

fourteen conditions showed enhanced performance (ten at high intensity, two at moderate intensity, and

two at low intensity), and seven conditions showed no difference (three at high intensity, three at

moderate intensity, and one at low intensity). In healthy older adults one condition showed no difference

(high intensity). Among individuals with stroke (e = 2), one condition showed no difference (moderate

intensity), and one condition showed attenuated performance (moderate intensity). In individuals with

Parkinson's disease (e = 3), all conditions showed enhanced performance (two at high intensity and one

at moderate intensity).

**Fig. 6 Effects of physical exercise interventions on motor learning.**
Colored dots represent individual experimental conditions, with their position indicating whether motor learning was enhanced, not different, or attenuated compared to a control condition. Only experiments including a control condition were considered. All population types investigated are represented. Reported results related to acquisition are shown only for experiments in which exercise was performed before motor task practice. Reported retention test results are shown regardless of whether exercise was performed before or after motor task practice. (Panel a) Reported results from acquisition. (Panel b) Reported results on motor learning from retention tests conducted within 24 hours following exercise, (Panel c) Reported results on motor learning from retention tests conducted at 24 hours following exercise (Panel d) Reported results on motor learning from retention tests conducted after 24 hours following exercise. Green dots indicate low-intensity exercise, orange dots indicate moderate-intensity exercise, and red dots indicate high-intensity exercise.

**3.6 Review articles previously published**

Among the included sources, 4 secondary sources were review articles. Bonuzzi and colleagues conducted a systematic review on aerobic exercise and motor learning in healthy individuals, highlighting that exercise performed before or after motor task practice can improve motor learning [41]. Hubner and colleagues conducted a systematic review examining the effects of physical activity level and acute bouts of physical exercise on motor performance and motor learning of upper limb tasks in older people. This systematic review highlighted the benefits of a high physical activity level in the early phases of motor learning [42]. Taubert and colleagues performed a narrative review and discussed behavioral and neurobiological evidence in favor of aerobic exercise as a neuromodulator strategy to promote motor learning and neuroplasticity [17]. Wanner and colleagues performed a systematic review and meta-analysis and evaluated how acute aerobic exercise affects the encoding and consolidation of motor memory, highlighting the importance of timing and intensity [91].

**4. Discussion**

To the best of our knowledge, this is the first scoping review to explore and map the existing
literature on the use of physical exercise to prime motor learning. We included studies involving any
population type, any form of physical exercise, and whether the physical exercise was performed before
or after any motor learning task. Although our scoping review focuses largely on experimental studies
(62 included studies), it should be noted that four previous reviews have explored specific aspects of the
topic (1 narrative review, 2 systematic reviews and 1 systematic review with meta-analysis). These
focused reviews each addressed important aspects of the topic, and our scoping review synthesis builds
on and integrates these contributions by examining a broader range of populations, exercise types, motor
learning categories and exercise timing in a single comprehensive framework.

The main outcomes of this scoping review were the following: (1) the evaluated population
consisted primarily of healthy individuals, with limited focus on clinical populations and children; (2)
the predominant type of physical exercise used was lower-limb cycling, potentially due to its ease of
implementation; (3) resistance exercise was rarely used, highlighting a significant gap in the literature
regarding its potential role in the promotion of motor learning; (4) in several studies, the prescribed
exercise intensity did not align accurately with ACSM guidelines, which may limit the ability to draw
accurate conclusions about the effects of specific exercise intensities on motor learning; (5) many studies
conducted a single retention test on the same day as the practice session; however, according to the
classical definition of motor learning, demonstrating maintenance in motor skill improvement over time
via delayed retention tests is essential [92], and hence these findings may not truly reflect motor learning.

**4.1 Limited representation of clinical populations**

Most of the included studies involved healthy individuals, with children and clinical populations
being significantly less evaluated. One reason could be due to practical limitations, as recruiting healthy
adults participants is usually less complicated and requires less time and resources. Yet, in order to
determine whether the results obtained in healthy individuals are transferable to children and individuals
living with specific health-related conditions, it is essential to replicate the interventions in children and
clinical populations. This is particularly of interest for children as this population is crucially developing
motor skills needed for their adult life. It is also crucial for clinical populations given that many
interventions improving motor learning are designed for rehabilitation purposes [93], and the clinical
relevance of these interventions should be tested within the clinical populations of interest. Furthermore,

given the limited number of studies involving clinical populations, the division across three distinct

conditions (stroke, Parkinson's disease, and Huntington's disease) further limits the applicability of the

findings to specific clinical settings, as each subgroup is represented by only a small number of studies.

Importantly, the underlying mechanisms of motor learning may differ between healthy

individuals and clinical populations due to different factors including impaired neuroplasticity, altered

sensory feedback, or compensatory motor strategies [94, 95]. For instance, in individuals with stroke,

cortical reorganization and the recruitment of alternative neural pathways may affect how motor skills

are re-acquired [96]. Similarly, in Parkinson's disease, basal ganglia and sensorimotor-related

dysfunction may alter feedback-based learning mechanisms [97]. These differences imply that exercise

interventions that are effective in healthy individuals may not produce the same effects in the context of

clinical conditions, making it essential to investigate tailored strategies specific to each clinical

population. Therefore, prioritizing research in populations with motor dysfunction and rehabilitation

needs including individuals with stroke, or other populations living with neurodegenerative disorders,

would be particularly valuable. These population groups could benefit the most from interventions that

can enhance motor learning, recovery and maintenance of functional independence. By targeting clinical

populations, we can develop more effective, evidence-based interventions that address their specific

motor learning profiles.

## 4.2 Types of aerobic exercise: a focus on lower-limb cycling and the need for diversity

Among the included studies, the predominant form of aerobic exercise was lower-limb cycling.

This type of aerobic exercise is likely favored due to its ease of implementation, low risk of injury, and

practicality in experimental settings. Particularly, lower-limb cycling effectively induces an aerobic

response by engaging the leg muscles, the largest muscle group in the body, resulting in a substantial

cardiovascular activation [98, 99]. Also, lower-limb cycling allows the upper limbs to remain at rest,

which helps prevent the development of neuromuscular fatigue that could negatively impact motor

learning when upper-limb motor tasks are used [100].

Other types of aerobic exercise such as running, walking, dancing and upper-limb cycling were

considerably less represented. Including these diverse forms of aerobic exercise in future research is

important for several reasons. Different types may activate distinct neuromuscular and cardiovascular

responses, influencing task-specific adaptations, and offer practical alternatives for various populations.

For example, running engages more complex coordination, while upper-limb cycling may be particularly

suitable for individuals with lower-limb limitations. Dancing, which is often practiced in groups and
especially popular among older adults also deserves attention, as it not only combines demands on the
aerobic system and coordination [101, 102], but has also been associated with enhanced cognitive
function [103, 104]. Dancing also may boost motivation and adherence to regular engagement in physical
activity through social interaction [105]. In addition, music, which is an integral part of most dance
interventions, has itself been linked to improved cognitive function [106] and positive affect [107],
further supporting the potential of dance-based activities in promoting motor learning. Such variety of
physical exercise could help reveal how the specific demands of different types of exercise might affect
motor learning.

Importantly, all of these reported aerobic types of physical exercise primarily relied on concentric
muscle contractions, where the muscles shorten while producing a force. None of the included studies
investigated aerobic exercise involving primarily eccentric contractions, like eccentric cycling, in which
muscles lengthen while producing a force.

**4.3 Eccentric and resistance exercise: underexplored avenues in motor learning research**

Interestingly, none of the included studies investigated the effects of physical exercise primarily
involving eccentric muscle contractions on motor learning. Eccentric contractions, which involve muscle
lengthening while resisting a workload, are known to induce distinct neural adaptations compared to
concentric contractions [108, 109]. Despite this, research focusing specifically on eccentric-based
physical exercise remains limited, representing a notable gap in the literature. Specifically, eccentric
contractions have been associated with a prolonged decrease in intracortical inhibition [109], a
neurophysiological marker of altered motor cortex excitability. Also, it has been shown that eccentric
cycling elicits greater activation in sensorimotor-related cortical areas of the frontal and parietal lobes
during movement planning and execution [108], as well as increased activation in cognitive-related brain
regions [110]. Given the involvement of these brain regions in motor learning, eccentric cycling could
induce potential benefits to motor learning.

In fact, recent preliminary evidence from our laboratory [111] supports this idea. This study,
which was not included in the present scoping review due to its preprint status at the time of submission,
suggests that eccentric cycling enhances motor learning more than concentric cycling in healthy young
adults. These findings suggest that compared to concentric cycling, eccentric cycling may offer specific
advantages that promote motor learning. Beyond its potential effects on motor learning, eccentric cycling

also provides physiological advantages. For a similar workload, eccentric cycling results in a lower
cardiovascular, respiratory, and metabolic cost compared to concentric cycling [112-115]. Eccentric
cycling is therefore a particularly appealing exercise modality for clinical populations with reduced
cardiorespiratory capacity [e.g., [116]], as well as for older adults [(e.g., [117]]. Since lower-limb cycling
is already the most employed exercise type in the reviewed literature, integrating eccentric cycling could
allow researchers to build on already established protocols while expanding the range of muscular
contractions studied. Finally, other eccentric-focused exercise types like downhill walking or running
also deserve attention. These exercise types involve distinct neuromuscular demands [118] and a lower
energy cost [119], which could be beneficial in rehabilitation. Given the potential benefits of eccentric
exercise on motor learning, its integration into future studies evaluating the effects of physical exercise
on motor learning appears both feasible and promising [120].

Beyond eccentric-focused types of exercise, our findings also raise important considerations
about resistance exercise, which often incorporates eccentric contractions and shares similar
physiological mechanisms [121]. These findings also highlight a significant gap in the motor learning
literature, which is the underrepresentation of resistance exercise as priming exercise. Of the 81
experimental conditions included, only two involved resistance exercise, despite the fact that this exercise
type is widely used in rehabilitation and training. Resistance exercise may influence motor learning
through several neural mechanisms. It has been shown that resistance exercise increases muscle
activation and neural drive to muscles, leading to improved force production capacity [122]. Resistance
exercise can also modulate corticospinal excitability, with evidence indicating that resistance training
induces neuroplastic changes within the motor cortex and corticospinal tract [123, 124]. Furthermore,
resistance exercise may enhance proprioceptive input, increasing sensory feedback, which is an essential
component of motor control and learning [125]. Supporting these observations, cerebral blood flow
during force exertion has been shown to correlate with activity in brain areas such as the primary motor
cortex, and the supplementary motor area [126]. These regions are also integral to motor learning
processes, suggesting a neuroanatomical overlap between areas recruited during resistance exercise and
those underlying motor learning. However, the direct role of the effects of resistance exercise on motor
learning remains largely unexplored. Investigating how resistance exercise influences motor learning
represents a promising avenue for future research. Taken together, these observations highlight the need
for future studies to incorporate both eccentric and resistance exercise types to better understand their
potential roles in enhancing motor learning across various populations.

**4.4 Prescription of exercise intensities**

Although all studies reported the exercise intensity used, we standardized intensity categorization

based on the exercise prescription details provided using the ACSM guidelines [40]. In 5 experimental

conditions, discrepancies were observed between the prescribed intensity and the actual workload

performed by participants. Some exercise appeared either lighter or more intense than prescribed. For

instance, Holman et al. (2021) and Statton et al. (2015) reported using moderate-intensity exercise, yet

the prescribed workloads correspond to high-intensity levels according to ACSM guidelines [24, 59].

Similarly, Opie and Semmler (2019) described their exercise protocol as low-intensity, although it aligns

with moderate-intensity based on ACSM classifications [70]. In contrast, Loras et al. (2020) referred to

their two exercise conditions as moderate and high intensity exercise, but these would correspond to low

and moderate intensity, respectively, under ACSM classifications [65]. First, this inconsistency suggests

that following established exercise guidelines is essential for drawing accurate conclusions about the

effects of specific exercise intensities on motor learning. Second, it allows replicability across studies

which is an essential factor for building a consistent and reliable body of evidence. Additionally, while

many studies relied on predicted values (e.g., age-predicted maximal heart rate, age-predicted heart rate

reserve, estimated VO2max), and although we recognize the practicality of using predicted values in

certain research settings, prescribing intensities based on measured capacities are preferable to ensure

that the targeted intensity is truly achieved for each participant. Interestingly, none of the studies included

in this review employed self-paced exercise protocols. This is notable, as self-paced exercise, for example

prescribed with the intensity of perceptual responses such as effort [127, 128], when matched for

workload with externally prescribed intensities [129], has been proposed to enhance affective responses,

which can be beneficial for promoting adherence to rehabilitation programs [130].

**4.5 Uneven representation of motor learning categories**

While the studies included in this review examined various motor learning categories such as

sequence learning, motor acuity, and motor adaptation, none investigated the effects of physical exercise

on associative learning, which is a fundamental form of motor learning involving stimulus-response

associations. Given its relevance in real-world contexts and rehabilitation settings [131], investigating

how physical exercise influences associative learning could offer valuable insights. For instance, tasks

such as learning to respond appropriately to traffic signals while driving rely heavily on stimulus-

response associations.

Additionally, the field would benefit from more balanced research across motor learning
categories. The current emphasis on motor acuity (n = 30) contrasts with the limited number of studies
evaluating sequence learning and motor adaptation (both n = 16). Sequence learning, for example, is
highly relevant for activities like playing a musical instrument or typing on a keyboard, while motor
adaptation is crucial in situations such as adjusting movements to a new sports technique or learning to
walk on uneven surfaces. More research in these underrepresented categories could allow for more robust
meta-analyses and potentially clarify whether the effects of physical exercise vary depending on the
motor learning type.

**4.6 Timing of retention tests**

When measuring skill retention, different time points were reported across the studies falling in
our defined time windows: less than 24 hours, at 24 hours, and more than 24 hours following motor task
practice. Some studies included two time points [29, 33, 35, 49, 54, 60, 62, 83, 86, 87, 89, 132] or all
three time points [28, 45, 56, 59, 66, 74, 77, 80, 85], which provides a clearer understanding of the effects
on motor learning. Other studies only assessed short-term retention at less than 24 hours [31, 34, 46, 48,
63, 64, 69, 81, 133]. However, measuring skill retention only in the short term (e.g., same day as initial
practice) may not be sufficient to assess motor learning, as this may primarily reflect temporary rather
sustained performance improvements [92]. According to Schmidt et al. (2018), observing performance
after a sufficient delay provides stronger evidence of stable and relatively permanent changes in the
learner's capability, which is the hallmark of motor learning [92].

Ideally, studies could include both short-term and longer-term retention tests, with the 24-hour
time point used as a standard. This would help determine whether performance improvements due to
practice are maintained over time, facilitate comparisons across studies, and provide insight into the
evolution of motor skill consolidation, which is the process by which newly acquired motor skills become
stable and resistant to interference. Using consistent methods to measure skill retention would also help
improve the quality of future studies in the field of physical exercise and motor learning.

**4.7 Limitations**

This scoping review has several limitations, particularly regarding the reported results of motor learning, due to the complexity of categorization. First, for retention tests, all studies were grouped together regardless of whether physical exercise was performed before or after skill acquisition, as the heterogeneity in study designs made it difficult to establish clear subcategories for the reported results on motor learning. Second, no differentiation was made based on the type of exercise for the reported results on motor learning, which limits the ability to attribute specific effects to a particular type of exercise. Third, because we examine the effects of physical exercise in general and as reported by the authors, exercise frequency was not considered, and no distinction was made between acute and chronic interventions. Future studies with more data may allow for a more refined mapping of the literature addressing these aspects.

**4.8 Perspectives for future studies**

This scoping review highlighted several key areas for future research on the effects of physical exercise on motor learning. First, accurately prescribing exercise intensity is essential. Many studies showed inconsistencies between the reported and the actual prescribed intensity, which can affect the reliability of findings. Using measured values rather than predicted ones, and adhering to established guidelines like the ACSM, would improve consistency and accuracy across studies. As self-paced exercise are proposed to enhance affective responses [127, 128] and have positive effect on exercise adherence, future studies should consider testing the possibility to prime motor learning with exercise prescribed based on perceptual responses, with particular interest on the perception of effort [127, 128]. Furthermore, although most studies have concentrated on concentric lower-limb cycling, it is essential to explore the impact of other types of exercise, such as eccentric cycling, on motor learning. Additionally, resistance exercise was rarely used. Integrating resistance exercise modalities will enable us to better understand its potential role in enhancing motor learning across various populations. Moreover, measuring skill retention in motor learning studies in exercise science is another area to improve. Studies testing retention only within a few hours may not fully capture the achievement and maintenance of motor learning. Therefore, including longer retention time points, particularly at 24 hours and more, might help in providing a more comprehensive understanding of the effects of physical exercise on motor learning. Finally, more studies are needed where children, older adults and clinical populations are involved. Increasing research on clinical populations is especially important as motor learning can play a critical role in rehabilitation and recovery.

**5. Conclusion**

In conclusion, while the field of physical exercise and motor learning is growing, there are several
key areas that require improvement. Future research should prioritize accurate and consistent reporting
of exercise intensity and consider longer retention intervals to better capture lasting motor learning
effects and, consequently, better understand the impact of physical exercise on motor learning.
Additionally, greater inclusion of children, older adults and clinical populations is essential, as these
population groups may benefit most from motor learning interventions but remain underrepresented in
the current literature. Expanding the range of exercise modalities, more particularly types of exercise
involving eccentric contractions and resistance-based exercise, may also reveal unique advantages
relevant to motor learning in both healthy and clinical populations. Addressing these gaps will help in
building a more robust and applicable body of evidence to inform both clinical and non-clinical practice.

## Author contributions

**Layale Youssef:** Conceptualization, Methodology, Formal analysis, Investigation, Writing – Original Draft, Visualization. **Amanda O'Farrell:** Formal analysis, Investigation. **Denis Arvisais:** Methodology, Investigation. **Jason L. Neva:** Conceptualization, Writing – Review & Editing, Visualization, Supervision. **Benjamin Pageaux:** Conceptualization, Writing – Review & Editing, Visualization, Supervision.

## Funding

LY is supported by Centre de Recherche de l'Institut Universitaire de Gériatrie de Montréal (CRIUGM), the Faculté de médecine at Université de Montréal and the Fonds de Recherche du Québec - Nature et Technologies. AOF is supported by internal scholarships from Université de Montréal. BP is supported by the Chercheur Boursier Junior 1 award from the Fonds de Recherche du Québec - Santé. JLN is supported by the Chercheur Boursier Junior 1 award from the Fonds de Recherche du Québec - Santé (FRQS #313769).

766                                                    References
767
1. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness:
definitions and distinctions for health-related research. Public Health Rep. 1985;100(2):126-31.

2. Ndahimana D, Kim E-KJCnr. Measurement methods for physical activity and energy expenditure: a
review. 2017;6(2):68-80.

3. Kenney WL, Wilmore JH, Costill DL. Physiology of sport and exercise: Human kinetics; 2022.

4. Cotman CW, Berchtold NC. Exercise: a behavioral intervention to enhance brain health and plasticity.
Trends in Neurosciences. 2002;25(6):295-301.

5. Phillips C, Baktir MA, Srivatsan M, Salehi A. Neuroprotective effects of physical activity on the brain:
a closer look at trophic factor signaling. Front Cell Neurosci. 2014;8:170.

6. Hillman CH, Erickson KI, Kramer AF. Be smart, exercise your heart: exercise effects on brain and
cognition. Nat Rev Neurosci. 2008;9(1):58-65.

7. Yanagisawa H, Dan I, Tsuzuki D, Kato M, Okamoto M, Kyutoku Y, et al. Acute moderate exercise
elicits increased dorsolateral prefrontal activation and improves cognitive performance with Stroop test.
Neuroimage. 2010;50(4):1702-10.

8. Erickson KI, Voss MW, Prakash RS, Basak C, Szabo A, Chaddock L, et al. Exercise training increases
size of hippocampus and improves memory. Proc Natl Acad Sci U S A. 2011;108(7):3017-22.

9. Andrews SC, Curtin D, Hawi Z, Wongtrakun J, Stout JC, Coxon JP. Intensity Matters: High-intensity
Interval Exercise Enhances Motor Cortex Plasticity More Than Moderate Exercise. Cereb Cortex.
2020;30(1):101-12.

10. Neva JL, Brown KE, Mang CS, Francisco BA, Boyd LA. An acute bout of exercise modulates both
intracortical and interhemispheric excitability. Eur J Neurosci. 2017;45(10):1343-55.

11. Youssef L, Harroum N, Francisco BA, Johnson L, Arvisais D, Pageaux B, et al. Neurophysiological
effects of acute aerobic exercise in young adults: a systematic review and meta-analysis. Neurosci
Biobehav Rev. 2024;164:105811.

12. Singh AM, Duncan RE, Neva JL, Staines WR. Aerobic exercise modulates intracortical inhibition
and facilitation in a nonexercised upper limb muscle. BMC Sports Science, Medicine and Rehabilitation.
2014;6(1):23.

13. Neva JL. Exercise-induced neuroplasticity. In: Grafman JH, editor. Encyclopedia of the Human Brain
(Second Edition). Oxford: Elsevier; 2025. p. 706-29.

14. Cirillo J, Todd G, Semmler JG. Corticomotor excitability and plasticity following complex
visuomotor training in young and old adults. Eur J Neurosci. 2011;34(11):1847-56.

15. Coxon JP, Peat NM, Byblow WD. Primary motor cortex disinhibition during motor skill learning. J
Neurophysiol. 2014;112(1):156-64.

16. Holland L, Murphy B, Passmore S, Yielder P. Time course of corticospinal excitability changes
following a novel motor training task. Neurosci Lett. 2015;591:81-5.

17. Taubert M, Villringer A, Lehmann N. Endurance exercise as an "endogenous" neuro-enhancement
strategy to facilitate motor learning. Frontiers in Human Neuroscience Vol 9 2015, ArtID 692. 2015;9.

18. Wanner P, Cheng F-H, Steib S. Effects of acute cardiovascular exercise on motor memory encoding
and consolidation: A systematic review with meta-analysis. Neuroscience and Biobehavioral Reviews.
2020;116:365-81.

19. Knapp B. Skill in Sport: The Attainment of Proficiency: Routledge & K. Paul; 1963.

20. Schmidt RA, Lee TD. Motor control and learning: A behavioral emphasis, 5th ed. Champaign, IL,
US: Human Kinetics; 2011. ix, 581-ix, p.

21. Krakauer JW, Hadjiosif AM, Xu J, Wong AL, Haith AM. Motor Learning. Compr Physiol.
2019;9(2):613-63.

22. McMorris T, Tomporowski P, Audiffren M. Exercise and cognitive function: John Wiley & Sons;
2009.

23. Snow NJ, Mang CS, Roig M, McDonnell MN, Campbell KL, Boyd LA. The effect of an acute bout
of moderate-intensity aerobic exercise on motor learning of a continuous tracking task. PLoS ONE Vol
11(2), 2016, ArtID e0150039. 2016;11(2).

24. Statton MA, Encarnacion M, Celnik P, Bastian AJ. A single bout of moderate aerobic exercise
improves motor skill acquisition. PLoS ONE Vol 10(10), 2015, ArtID e0141393. 2015;10(10).

25. Mang CS, Snow NJ, Campbell KL, Ross CJ, Boyd LA. A single bout of high-intensity aerobic
exercise facilitates response to paired associative stimulation and promotes sequence-specific implicit
motor learning. J Appl Physiol (1985). 2014;117(11):1325-36.

26. Neva JL, Ma JA, Orsholits D, Boisgontier MP, Boyd LA. The effects of acute exercise on visuomotor
adaptation, learning, and inter-limb transfer. Experimental Brain Research. 2019;237(4):1109-27.

27. Lehmann N, Villringer A, Taubert M. Colocalized white matter plasticity and increased cerebral
blood flow mediate the beneficial effect of cardiovascular exercise on long-term motor learning. The
Journal of Neuroscience. 2020;40(12):2416-29.

28. Ferrer-Uris B, Busquets A, Lopez-Alonso V, Fernandez-Del-Olmo M, Angulo-Barroso R. Enhancing
consolidation of a rotational visuomotor adaptation task through acute exercise. PLoS ONE.
2017;12(4):e0175296.

29. Jespersen L, Maes KM, Ardenkjær-Skinnerup N, Roig M, Bjorndal JR, Beck MM, et al. Acute
exercise performed before and after motor practice enhances the positive effects on motor memory
consolidation. Neurobiology of Learning and Memory. 2023;205:10.

30. Mang CS, Snow NJ, Wadden KP, Campbell KL, Boyd LA. High-Intensity Aerobic Exercise
Enhances Motor Memory Retrieval. Medicine & Science in Sports & Exercise. 2016;48(12):2477-86.

31. Stavrinos EL, Coxon JP. High-intensity interval exercise promotes motor cortex disinhibition and
early motor skill consolidation. Journal of Cognitive Neuroscience. 2017;29(4):593-604.

32. Baird JF, Gaughan ME, Saffer HM, Sarzynski MA, Herter TM, Fritz SL, et al. The effect of energy-
matched exercise intensity on brain-derived neurotrophic factor and motor learning. Neurobiology of
Learning and Memory. 2018;156:33-44.

33. Charalambous CC, French MA, Morton SM, Reisman DS. A single high-intensity exercise bout
during early consolidation does not influence retention or relearning of sensorimotor locomotor long-
term memories. Experimental Brain Research. 2019;237(11):2799-810.

34. James R, Wang JS. The effects of a single bout of moderate-intensity aerobic exercise on visuomotor
adaptation and its savings. Neurobiology of Learning and Memory. 2023;204:7.

35. Pixa NH, Hubner L, Kutz DF, Voelcker-Rehage C. A Single Bout of High-Intensity Cardiovascular
Exercise Does Not Enhance Motor Performance and Learning of a Visuomotor Force Modulation Task,
but Triggers Ipsilateral Task-Related EEG Activity. Int J Environ Res Public Health. 2021;18(23):27.

36. Quinlan C, Rattray B, Pryor D, Northey JM, Coxon J, Cherbuin N, et al. A short-term intervention
of high-intensity exercise and anodal-tDCS on motor learning in middle-aged adults: An RCT. Frontiers
in Human Neuroscience Vol 15 2021, ArtID 661079. 2021;15.

37. Wanner P, Muller T, Cristini J, Pfeifer K, Steib S. Exercise Intensity Does not Modulate the Effect
of Acute Exercise on Learning a Complex Whole-Body Task. Neuroscience. 2020a;426:115-28.

38. Peters MDJ, Marnie C, Tricco AC, Pollock D, Munn Z, Alexander L, et al. Updated methodological
guidance for the conduct of scoping reviews. JBI Evid Synth. 2020;18(10):2119-26.

39. Innovation VH. Covidence systematic review software. 2023.

40. ACSM. Amercian College of Sports Medecine's Guidelines for Exercise Testing and Prescription.
10th ed2018.

41. Bonuzzi GMG, Torriani-Pasin C. Cardiovascular exercise and motor learning in non-disabled
individuals: A systematic review with a behavioral emphasis. Motriz-Revista De Educacao Fisica.
2022;Revista de Educacao Fisica. 28(no pagination)(e1022005221).

42. Hubner L, Voelcker-Rehage C. Does physical activity benefit motor performance and learning of
upper extremity tasks in older adults? - A systematic review. Eur Rev Aging Phys Act. 2017;14:19.

43. Andrews SC, Kämpf L, Curtin D, Hinder M, Wenderoth N, Stout JC, et al. A single bout of moderate-
intensity aerobic exercise improves motor learning in premanifest and early Huntington's disease. Front
Psychol. 2023;14:9.

44. Andrushko JW, Rinat S, Greeley B, Larssen BC, Jones CB, Rubino C, et al. Improved processing
speed and decreased functional connectivity in individuals with chronic stroke after paired exercise and
motor training. Sci. 2023;13(1):14.

45. Angulo-Barroso R, Ferrer-Uris B, Busquets A. Enhancing children's motor memory retention through
acute intense exercise: Effects of different exercise durations. Frontiers in Psychology Vol 10 2019,
ArtID 2000. 2019;10.

46. Bartz DW, Smith LE. Effect of moderate exercise on the performance and learning of a gross motor
skill. Perceptual and Motor Skills. 1970;31(1):187-90.

47. Bonuzzi GMG, Bastos FH, Schweighofer N, Wade E, Winstein CJ, Torriani-Pasin C. Moderate-
intensity cardiovascular exercise performed before motor practice attenuates offline implicit motor
learning in stroke survivors but not age-matched neurotypical adults. Experimental Brain Research.
2023;241(8):2019-32.

48. Bosch B, Bringard A, Logrieco MG, Lauer E, Imobersteg N, Thomas A, et al. Effect of acute physical
exercise on motor sequence memory. Sci. 2020;10(1):15322.

49. Chan ST, Tai CH, Wang LY, Luh JJ, Lee YY. Influences of Aerobic Exercise on Motor Sequence
Learning and Corticomotor Excitability in People With Parkinson's Disease. Neurorehabilitation and
Neural Repair. 2023;37(1):37-45.

50. Charalambous CC, Alcantara CC, French MA, Li X, Matt KS, Kim HE, et al. A single exercise bout
and locomotor learning after stroke: physiological, behavioural, and computational outcomes. J Physiol
(Lond). 2018;596(10):1999-2016.

51. Chen C, Ryuh YJ, Tomporowski PP, Lamberth J, Pan Z, Conners DA. Acute Effect of Skill Random
Practice Versus Cardiovascular Exercise on Motor Learning of a Golf Putting Task. International Journal
of Sport Psychology. 2023;54(2):140-51.

52. Coco M, Cavallari P, Perciavalle V. Effects of an exhaustive exercise on motor skill learning and on
the excitability of primary motor cortex and supplementary motor area. Medicine (United States).
2016;95 (11) (no pagination)(e2978).

53. Cristini J, Kraft VS, De las Heras B, Rodrigues L, Parwanta Z, Hermsdorfer J, et al. Differential
effects of acute cardiovascular exercise on explicit and implicit motor memory: The moderating effects
of fitness level. Neurobiology of Learning and Memory. 2023;205:1-12.

54. Dal Maso F, Desormeau B, Boudrias M-H, Roig M. Acute cardiovascular exercise promotes
functional changes in cortico-motor networks during the early stages of motor memory consolidation.
NeuroImage. 2018;174:380-92.

55. Duchesne C, Gheysen F, Bore A, Albouy G, Nadeau A, Robillard ME, et al. Influence of aerobic
exercise training on the neural correlates of motor learning in Parkinson's disease individuals.
NeuroImage: Clinical. 2016;12:559-69.

56. Ferrer-Uris B, Busquets A, Angulo-Barroso R. Adaptation and retention of a perceptual-motor task
in children: Effects of a single bout of intense endurance exercise. Journal of Sport & Exercise
Psychology. 2018;40(1):1-9.

57. Greeley B, Chau B, Jones CB, Neva JL, Kraeutner SN, Campbell KL, et al. Multiple bouts of high-intensity interval exercise reverse age-related functional connectivity disruptions without affecting motor learning in older adults. Sci. 2021;11(1):17108.

58. Helm EE, Matt KS, Kirschner KF, Pohlig RT, Kohl D, Reisman DS. The influence of high intensity exercise and the Val66Met polymorphism on circulating BDNF and locomotor learning. Neurobiology of Learning and Memory. 2017;144:77-85.

59. Holman SR, Staines WR. The effect of acute aerobic exercise on the consolidation of motor memories. Experimental Brain Research. 2021;239(8):2461-75.

60. Hubner L, Godde B, Voelcker-Rehage C. Acute Exercise as an Intervention to Trigger Motor Performance and EEG Beta Activity in Older Adults. Neural Plast. 2018;2018:4756785.

61. Hung A, Roig M, Gillen JB, Sabiston CM, Swardfager W, Chen JL. Aerobic exercise and aerobic fitness level do not modify motor learning. Sci. 2021;11(1):12.

62. Khan A, Singh J, Neary JP, Mang CS. Exercise Effects on Motor Skill Consolidation and Intermuscular Coherence Depend on Practice Schedule. Brain Sciences. 2022;12(4):17.

63. Kuo HI, Hsieh MH, Lin YT, Kuo MF, Nitsche MA. A single bout of aerobic exercise modulates motor learning performance and cortical excitability in humans. Int J Clin Health Psychol. 2023;23(1):100333.

64. Kuo HI, Hsieh MH, Lin YT, Nitsche MA. Acute Aerobic Exercise at Different Intensities Modulates Motor Learning Performance and Cortical Excitability in Sedentary Individuals. eNeuro. 2023;10(11).

65. Loras H, Haga M, Sigmundsson H. Effect of a Single Bout of Acute Aerobic Exercise at Moderate-to-Vigorous Intensities on Motor Learning, Retention and Transfer. Sports. 2020;8(2):13.

66. Lundbye-Jensen J, Skriver K, Nielsen JB, Roig M. Acute exercise improves motor memory consolidation in preadolescent children. Frontiers in Human Neuroscience Vol 11 2017, ArtID 182. 2017;11.

67. Munz M, Baving L, Prehn-Kristensen A. Sleep following intense physical exercise stabilizes motor learning in typically developing boys. Mental Health and Physical Activity Vol 20 2021, ArtID 100365. 2021;20.

68. Nepveu JF, Thiel A, Tang A, Fung J, Lundbye-Jensen J, Boyd LA, et al. A Single Bout of High-Intensity Interval Training Improves Motor Skill Retention in Individuals With Stroke. Neurorehabil Neural Repair. 2017;31(8):726-35.

69. Ostadan F, Centeno C, Daloze JF, Frenn M, Lundbye-Jensen J, Roig M. Changes in corticospinal excitability during consolidation predict acute exercise-induced off-line gains in procedural memory. Neurobiol Learn Mem. 2016;136:196-203.

70. Opie GM, Semmler JG. Acute Exercise at Different Intensities Influences Corticomotor Excitability and Performance of a Ballistic Thumb Training Task. Neuroscience. 2019;412:29-39.

71. Perini R, Bortoletto M, Capogrosso M, Fertonani A, Miniussi C. Acute effects of aerobic exercise promote learning. Sci Rep. 2016;6:25440.

72. Quaney BM, Boyd LA, McDowd JM, Zahner LH, Jianghua H, Mayo MS, et al. Aerobic exercise improves cognition and motor function poststroke. Neurorehabilitation and Neural Repair. 2009;23(9):879-85.

73. Rhee J, Chen J, Riechman SM, Handa A, Bhatia S, Wright DL. An acute bout of aerobic exercise can protect immediate offline motor sequence gains. Psychological Research. 2016;80(4):518-31.

74. Roig M, Skriver K, Lundbye-Jensen J, Kiens B, Nielsen JB. A single bout of exercise improves motor memory. PLoS ONE Vol 7(9), 2012, ArtID e44594. 2012;7(9).

75. Roig-Hierro E, Batalla A. Acute exercise on complex motor memory retention: the role of task cognitive demands. International Journal of Sport and Exercise Psychology. 2023:15.

76. Singh, Neva JL, Staines WR. Aerobic exercise enhances neural correlates of motor skill learning. Behav Brain Res. 2016;301:19-26.

77. Skriver K, Roig M, Lundbye-Jensen J, Pingel J, Helge JW, Kiens B, et al. Acute exercise improves motor memory: Exploring potential biomarkers. Neurobiology of Learning and Memory. 2014;116:46-58.

78. Steib S, Wanner P, Adler W, Winkler J, Klucken J, Pfeifer K. A single bout of aerobic exercise improves motor skill consolidation in Parkinson's disease. Frontiers in Aging Neuroscience Vol 10 2018, ArtID 328. 2018;10.

79. Stranda H, Haga M, Sigmundsson H, Loras H. The Effect of Aerobic Exercise on Speed and Accuracy Task Components in Motor Learning. Sports. 2019;7(3):10.

80. Swarbrick D, Kiss A, Trehub S, Tremblay L, Alter D, Chen JL. HIIT the Road Jack: An Exploratory Study on the Effects of an Acute Bout of Cardiovascular High-Intensity Interval Training on Piano Learning. Front Psychol. 2020;11:17.

81. Taylor EM, Cadwallader CJ, Curtin D, Chong TTJ, Hendrikse JJ, Coxon JP. High-intensity acute exercise impacts motor learning in healthy older adults. npj Science of Learning. 2024;9(1):9.

82. Thomas R, Beck MM, Lind RR, Korsgaard Johnsen L, Geertsen SS, Christiansen L, et al. Acute Exercise and Motor Memory Consolidation: The Role of Exercise Timing. Neural Plasticity. 2016;2016:6205452.

83. Thomas R, Flindtgaard M, Skriver K, Geertsen SS, Christiansen L, Korsgaard Johnsen L, et al. Acute exercise and motor memory consolidation: Does exercise type play a role? Scandinavian Journal of Medicine & Science in Sports. 2017;27(11):1523-32.

84. Thompson ED, Bhat S, French MA, Morton S, Pohlig RT, Reisman DS. Effects of an Acute High Intensity Exercise Bout on Retention of Explicit, Strategic Locomotor Learning in Individuals With Chronic Stroke. Neurorehabilitation and Neural Repair. 2023;37(9):628-39.

85. Tomporowski PD, Pendleton DM. Effects of the timing of acute exercise and movement complexity on young adults' psychomotor learning. Journal of Sport & Exercise Psychology. 2018;40(5):240-8.

86. Wanga F, Jiang Y, Hou L. Effects of different exercise intensities on motor skill learning capability and process. Sci Sports. 2023;38(2):9.

87. Wanner P, Winterholler M, Gasner H, Winkler J, Klucken J, Pfeifer K, et al. Acute exercise following skill practice promotes motor memory consolidation in Parkinson's disease. Neurobiology of Learning and Memory Vol 178 2021, ArtID 107366. 2021;178.

88. Williams L, Pottinger P, Shapcott D. Effects of exercise on choice reaction latency and movement speed. Perceptual and Motor Skills. 1985;60(1):67-71.

89. Yildirim MS, Guclu-Gunduz A, Ozkul C, Korkmaz S. Investigating the acute effect of low and moderate intensity aerobic exercise on whole-body task learning and cognition in young adults. Eur J Neurosci. 2024;60(6):5203-16.

90. Duchesne C, Lungu O, Nadeau A, Robillard ME, Boré A, Bobeuf F, et al. Enhancing both motor and cognitive functioning in Parkinson's disease: Aerobic exercise as a rehabilitative intervention. Brain Cogn. 2015;99:68-77.

91. Wanner P, Cheng F, Steib S. Effects of acute cardiovascular exercise on motor memory encoding and consolidation: A systematic review with meta-analysis. Neuroscience and Biobehavioral Reviews. 2020b;116:365-81.

92. Schmidt RA, Lee TD, Winstein C, Wulf G, Zelaznik HN. Motor control and learning: A behavioral emphasis: Human kinetics; 2018.

93. Krakauer JW. Motor learning: its relevance to stroke recovery and neurorehabilitation. Curr Opin Neurol. 2006;19(1):84-90.

94. Boyd LA, Winstein CJ. Explicit Information Interferes with Implicit Motor Learning of Both Continuous and Discrete Movement Tasks After Stroke. 2006;30(2):46-57.

95. Buma FE, Lindeman E, Ramsey NF, Kwakkel GJN, repair n. Functional neuroimaging studies of early upper limb recovery after stroke: a systematic review of the literature. 2010;24(7):589-608.

96. Nudo RJJCoin. Mechanisms for recovery of motor function following cortical damage. 2006;16(6):638-44.

97. Boyd LA, Winstein CJJL, memory. Providing explicit information disrupts implicit motor learning after basal ganglia stroke. 2004;11(4):388-96.

98. Poole DC, Richardson RSJSM. Determinants of oxygen uptake: implications for exercise testing. 1997;24:308-20.

99. Scott K, Howley ET, Quindry J. Exercise physiology: theory and application to fitness and performance: McGraw-Hill; 2021.

100. Ranieri F, Di Lazzaro V. The role of motor neuron drive in muscle fatigue. Neuromuscul Disord. 2012;22 Suppl 3:S157-61.

101. Keogh JW, Kilding A, Pidgeon P, Ashley L, Gillis DJJoa, activity p. Physical benefits of dancing for healthy older adults: a review. 2009;17(4):479-500.

102. Limanskaya OV, Yefimova OV, Kriventsova IV, Wnorowski K, Bensbaa AJPeos. The coordination abilities development in female students based on dance exercises. 2021;25(4):249-56.

103. Karpati FJ, Giacosa C, Foster NEV, Penhune VB, Hyde KL. Dance and the brain: a review. 2015;1337(1):140-6.

104. Hewston P, Kennedy CC, Borhan S, Merom D, Santaguida P, Ioannidis G, et al. Effects of dance on cognitive function in older adults: a systematic review and meta-analysis. 2021;50(4):1084-92.

105. Brustio PR, Liubicich ME, Chiabrero M, Rabaglietti EJGN. Dancing in the golden age: a study on physical function, quality of life, and social engagement. 2018;39(6):635-9.

106. Schellenberg EGJCDiPS. Music and cognitive abilities. 2005;14(6):317-20.

107. Campion M, Levita LJTJoPP. Enhancing positive affect and divergent thinking abilities: Play some music and dance. 2014;9(2):137-45.

108. Fang Y, Siemionow V, Sahgal V, Xiong F, Yue GH. Greater movement-related cortical potential during human eccentric versus concentric muscle contractions. J Neurophysiol. 2001;86(4):1764-72.

109. Latella C, Goodwill AM, Muthalib M, Hendy AM, Major B, Nosaka K, et al. Effects of eccentric versus concentric contractions of the biceps brachii on intracortical inhibition and facilitation. Scand J Med Sci Sports. 2019;29(3):369-79.

110. Borot L, Pageaux B, Laroche D, Vergotte G, Lepers R, Perrey S. Eccentric cycling involves greater mental demand and cortical activation of the frontoparietal network. 2024;34(1):e14517.

111. Youssef L, Pageaux B, Neva JL. Eccentric cycling improves motor learning more than concentric cycling. 2025:2025.05.08.652761.

112. Clos P, Laroche D, Stapley PJ, Lepers R. Neuromuscular and Perceptual Responses to Sub-Maximal Eccentric Cycling. Front Physiol. 2019;10:354.

113. Isner-Horobeti ME, Dufour SP, Vautravers P, Geny B, Coudeyre E, Richard R. Eccentric exercise training: modalities, applications and perspectives. Sports Med. 2013;43(6):483-512.

114. Peñailillo L, Blazevich A, Numazawa H, Nosaka K. Metabolic and muscle damage profiles of concentric versus repeated eccentric cycling. Med Sci Sports Exerc. 2013;45(9):1773-81.

115. Peñailillo L, Blazevich AJ, Nosaka K. Muscle fascicle behavior during eccentric cycling and its relation to muscle soreness. Med Sci Sports Exerc. 2015;47(4):708-17.

116. Pageaux B, Besson D, Casillas JM, Lepers R, Gremeaux V, Ornetti P, et al. Progressively increasing the intensity of eccentric cycling over four training sessions: A feasibility study in coronary heart disease patients. Ann Phys Rehabil Med. 2020;63(3):241-4.

117. Gault ML, Willems ME. Aging, functional capacity and eccentric exercise training. Aging Dis. 2013;4(6):351-63.

118. Garnier YM, Lepers R, Dubau Q, Pageaux B, Paizis C. Neuromuscular and perceptual responses to moderate-intensity incline, level and decline treadmill exercise. Eur J Appl Physiol. 2018;118(10):2039-53.

119. Vernillo G, Giandolini M, Edwards WB, Morin J-B, Samozino P, Horvais N, et al. Biomechanics and Physiology of Uphill and Downhill Running. Sports Medicine. 2017;47(4):615-29.

120. LaStayo P, Marcus R, Dibble L, Frajacomo F, Lindstedt SJJoAP. Eccentric exercise in rehabilitation: safety, feasibility, and application. 2014;116(11):1426-34.

121. Douglas J, Pearson S, Ross A, McGuigan M. Eccentric Exercise: Physiological Characteristics and Acute Responses. Sports Medicine. 2017;47(4):663-75.

122. Aagaard P, Simonsen EB, Andersen JL, Magnusson P, Dyhre-Poulsen P. Increased rate of force development and neural drive of human skeletal muscle following resistance training. 2002;93(4):1318-26.

123. Kidgell DJ, Pearce AJ. Corticospinal properties following short-term strength training of an intrinsic hand muscle. Human Movement Science. 2010;29(5):631-41.

124. Carroll TJ, Riek S, Carson RG. The sites of neural adaptation induced by resistance training in humans. 2002;544(2):641-52.

125. Proske U, Gandevia SC. The Proprioceptive Senses: Their Roles in Signaling Body Shape, Body Position and Movement, and Muscle Force. 2012;92(4):1651-97.

126. Dettmers C, Fink GR, Lemon RN, Stephan KM, Passingham RE, Silbersweig D, et al. Relation between cerebral activity and force in the motor areas of the human brain. 1995;74(2):802-15.

127. Pageaux B. Perception of effort in Exercise Science: Definition, measurement and perspectives. Eur J Sport Sci. 2016;16(8):885-94.

128. Beaume J-B, Di Domenico H, Bowen M, Hintzy F, Millet GY, Pageaux B, et al. Neuromuscular Fatigue Induced by Cycling at a Fixed Level of Perceived Effort: Effects of Different Purported Hypoxic Methods. 2025;35(2):e70021.

129. Ekkekakis P, Hall EE, Petruzzello SJ. The relationship between exercise intensity and affective responses demystified: to crack the 40-year-old nut, replace the 40-year-old nutcracker! Ann Behav Med. 2008;35(2):136-49.

130. Ekkekakis P, Parfitt G, Petruzzello SJ. The pleasure and displeasure people feel when they exercise
at different intensities: decennial update and progress towards a tripartite rationale for exercise intensity
prescription. Sports Med. 2011;41(8):641-71.

131. Le Pelley ME, Mitchell CJ, Beesley T, George DN, Wills AJJPb. Attention and associative learning
in humans: An integrative review. 2016;142(10):1111.

132. Thomas R, Johnsen LK, Geertsen SS, Christiansen L, Ritz C, Roig M, et al. Acute exercise and
motor memory consolidation: The role of exercise intensity. PLoS ONE Vol 11(7), 2016, ArtID
e0159589. 2016;11(7).

133. Coco M, Perciavalle V, Cavallari P, Perciavalle V. Effects of an Exhaustive Exercise on Motor Skill
Learning and on the Excitability of Primary Motor Cortex and Supplementary Motor Area. Medicine
(Baltimore). 2016;95(11):e2978.

1085

1086

# Figure captions

**Fig. 1 Flowchart illustrating the selection process of sources.**

**Fig. 2 Temporal and geographical distribution of included sources.**
(Panel a) Temporal distribution of all included sources. Orange bars represent experimental studies, while blue gradient bars represent other types of sources (narrative review; systematic review; systematic review with meta-analysis). (Panel b) Geographical distribution of all included sources. The format used is the following: total number [other types of sources*]. The total number of sources is shown first; when sources other than experimental studies are included, their number is indicated in brackets. Darker colors indicate a higher number of sources reported from each country. n refers to the number of participants reported from experimental studies conducted in that country.

**Fig. 3 Types of populations evaluated across all included experimental studies.**
Distribution of the 1,479 participants reported across the 62 experimental studies. Participants were categorized as either healthy or clinical populations, with each category further divided into subcategories. Green gradients represent healthy populations, and orange gradients represent clinical populations. k refers to the number of experimental studies, G refers to the overall number of reported groups, g refers to the number of reported subgroups.

**Fig. 4 Types and intensities of physical exercise reported across all included experimental studies.**
(Panel a) Distribution of the 81 experimental conditions according to exercise type, categorized as aerobic or resistance. Aerobic exercise is further subcategorized by specific types. (Panel b) Prescription of exercise intensities across the experimental studies, pink refers to measured capacity and turquoise refers to estimated capacity. Pink and turquoise gradients refer to measured and estimated exersie parameters respectively. (Panel c) Exercise intensities used across all experiments where red refers to high intensity, yellow refers to moderate intensity and green refers to low intensity. (Panel d) Heatmap showing the combination of exercise type and intensity where darker purple indicates a higher number of experimental conditions using the specific combination. Others refers to all exercise types different from lower-limb cycling, running and walking. E refers to the total number of experimental conditions reported across the 62 included experimental studies, e refers to the number of experimental conditions per exercise type category and n refers to the number of participants.

**Fig. 5 Motor learning categories and timing of exercise relative to motor task practice.**

(Panel a) Distribution of motor learning categories evaluated across the 62 experimental studies. (Panel

b) Timing of exercise in relation to motor task practice across the 73 experimental conditions, (Panel c)

Timing of exercise in relation to motor task practice across the 73 experimental conditions shown by

population type. Light blue indicates exercise performed before the motor task practice, and light red

indicates exercise performed after the motor task practice. Before = exercise performed prior to motor

task practice; After = exercise performed after the motor task practice; n refers to the number of

participants.

1126

**Fig. 6 Effects of physical exercise interventions on motor learning.**

Colored dots represent individual experimental conditions, with their position indicating whether motor

learning was enhanced, not different, or attenuated compared to a control condition. Only experiments

including a control condition were considered. All population types investigated are represented.

Reported results related to acquisition are shown only for experiments in which exercise was performed

before motor task practice. Reported retention test results are shown regardless of whether exercise was

performed before or after motor task practice. (Panel a) Reported results from acquisition. (Panel b)

Reported results on motor learning from retention tests conducted within 24 hours following exercise,

(Panel c) Reported results on motor learning from retention tests conducted at 24 hours following

exercise (Panel d) Reported results on motor learning from retention tests conducted after 24 hours

following exercise. Green dots indicate low-intensity exercise, orange dots indicate moderate-intensity

exercise, and red dots indicate high-intensity exercise.

1139

1140

1141 # **Supplementary Material 1**

1142

1143 **Figure S1.** Classification of the task categories investigated to infer motor learning. This figure illustrates
1144 four commonly used motor learning task categories, classified according to their primary objective. The
1145 examples shown represent typical experimental tasks used within each category to assess motor learning.

1146

*Studies cited in Figure S1:*

Alderman, R. B. (1965, 1965/05/01). Influence of Local Fatigue on Speed and Accuracy in Motor Learning. Research Quarterly. American Association for
Health, Physical Education and Recreation, 36(2), 131-140. https://doi.org/10.1080/10671188.1965.10614670

Balsters, J. H., & Ramnani, N. (2011, Feb 9). Cerebellar plasticity and the automation of first-order rules. Journal of Neuroscience, 31(6), 2305-2312.
https://doi.org/10.1523/jneurosci.4358-10.2011

Neva, J. L., Ma, J. A., Orsholits, D., Boisgontier, M. P., & Boyd, L. A. (2019, Apr). The effects of acute exercise on visuomotor adaptation, learning, and
inter-limb transfer [Empirical Study; Quantitative Study]. Experimental Brain Research, 237(4), 1109-1127.
https://doi.org/http://dx.doi.org/10.1007/s00221-019-05491-5

Robertson, E. M. (2007). The Serial Reaction Time Task: Implicit Motor Skill Learning? , 27(38), 10073-10075. https://doi.org/10.1523/JNEUROSCI.2747-
07.2007 %J The Journal of Neuroscience

1157

1158

1159 **Table S1:** Characteristics of the studies included in the scoping review.

| Study | Total N | Type of population (n of exercise groups; age ± SD years) | Physical exercise | Exercise parameters | Motor Learning Category | Exercise Timing to Acquisition | Retention | Exercise effect on motor learning compared to control — Acquisition | Retention |
|---|---|---|---|---|---|---|---|---|---|
| Andrews et al., 2023 | 20 | Huntington's disease (n = 10, 48.4 ± 13) | Lower-limb cycling | Moderate 20 min - CONT 50-55% age-predicted HRR | Motor acuity | Before | 7d | ↗ | ↔ |
| Andrushko et al., 2023 | 25 | Stroke patients (n = 14, 66 ± 11) | Lower-limb cycling (5 sessions) | High 23 min - HIIT 3min at 75% Wpeak 3 min at 10W | Sequence | Before | 24h | ↔ | ↔ |
| Angulo-Barroso et al., 2019 | 71 | Healthy children (n = 46, 9.2 ± 0.9) | Running | High 16 min (LONG) ; 5 min (SHORT) - HIIT LONG: 3 min at 85% estim. VO₂max ; 2 min at 60% estim. VO₂max SHORT: 2 min at 85% estim. VO₂max ; 1 min at 60% estim. VO₂max | Motor adaptation | Before | 1h, 24h, 7d | ↔ | ↗ for LONG and SHORT |
| Baird et al., 2018 | 48 | Healthy adults (n = 32, 23 ± 3.1) | Lower-limb cycling | Low CONT: 40% measured max resistance until 200 kcals were expended (average: 28.67 min) High CONT: 80% measured max resistance until 200kcals were expended (average 16.75 min) | Sequence | Before | 24h | ↔ | ↔ |
| Bartz et al., 1970 | 20 | Healthy adults (n = 10, 18-20 yrs) | Walking | Moderate 4 min - CONT HR target: 120-170 bpm | Motor adaptation | Before | <10 min | ↔ | ↔ |
| Bonuzzi et al., 2023 | 90 | Stroke patients (n = 15; 51.6 ± 9.2) Stroke patients (n = 15; 49.6 ± 15.9) Healthy adults (n = 15; 51.6 ± 9.2) Healthy adults (n = 15; 49.6 ± 15.9) | Lower-limb cycling (3 sessions) | Moderate 20 min - CONT 50-70% age-predicted HRR | Sequence | Before / After / Before / After | 10d | ↔ / ↔ / ↘ / ↔ | ↘ / ↔ / ↔ / ↔ |
| Bosch et al., 2020 | 15 | Healthy adults (n = 15; 23.7 ± 4.0) | Lower-limb cycling | Moderate 30 min - CONT 70% measured MHR High 15 min - CONT 80% measured MHR | Sequence | After | 2.5h | NA | ↗ for High ↔ for Moderate |
| Chan et al., 2023 | 30 | Parkinson's disease (n = 15; 62.4 ± 8.5) | Lower-limb cycling | High 20 min - CONT 60-75% measured HRR | Sequence | After | 1d, 7d | ↔ | ↗ at Day 7 |
| Charalambous et al., 2018 (active control) | 37 | Stroke patients (n=12; 55 ± 16) Stroke patients (n=12; 62 ± 10) | Treadmill walking Total body exercise | High 5 min - CONT 70-85% age-predicted MHR | Motor adaptation | Before / After | 24h | ↔ / ↔ | ↔ / ↔ |
| Charalambous et al., 2019 | 26 | Healthy adults (n = 13; 21.8 ± 0.5) | Whole-body cycling | High 5 min - CONT 77-94% age-predicted MHR | Motor adaptation | Before | 1d, 7d | ↔ | ↔ |
| Chen et al., 2023 | 24 | Healthy adults (n = 12; 20.9 ± 0.7) | Treadmill Running | Moderate 30 min - CONT 65-85% age-predicted MHR (average was 75%) | Motor acuity | Before | 24h | ↔ | ↔ |
| Coco et al., 2016 (no control) | 28 | Healthy adults (n = 28; 37.4 ± 5.8) | Lower-limb cycling | High CONT: 3 minutes of unloaded cycling with 30W increase every 3min until exhaustion | Motor acuity | After | < 1h | NA | ↘ |
| Cristini et al., 2023 | 56 | Healthy adults (n = 20; 23.9 ± 3.4) | Lower-limb cycling | High 16 min - HIIT 3 min at 85–90 % of measured PPO 2min at 25 % of PPO. | Sequence | After | 24h | NA | ↔ |
| Dal Maso et al., 2018 | 25 | Healthy adults (n = 12; 24.3 ± 3.6) | Lower-limb cycling | High 15 min - HIIT 3min at 90% measured HRR 2min at 50W | Motor acuity | After | 8h, 24h | NA | ↔ for 8h ↗ at 24h |
| Duchesne et al., 2015 | 39 | Parkinson's disease (n = 19; 59 ± 7.1) Healthy older adults (n = 20; 64 ± 8.2) | Lower-limb cycling (3x/week for 12 weeks) | High CONT Start: 20 min at 60% measured PPO End: 40 min at 80% measured PPO | Sequence | After / After | 12 weeks | NA | ↗ / ↗ |

1160
1161
1162
1163
CONT = continuous; HRR = Heart rate reserve; W = Watts; HIIT = high-intensity interval training; kcals = kilocalories; bpm = beats per minute; HR = Heart Rate; MHR = maximal heart rate; PPO = peak power output; estim. = estimated; Before = exercise was performed before the motor task practice; After = exercise was performed after the motor task practice; min = minutes; h = hours; d = days; NA = not available; ↗ = significant enhancement following exercise; ↘ = significant decrease following exercise ; ↔ = no change following exercise.

1164 **Table S1.** *continued*

| Study | Total N | Type of population (n of exercise groups; age ± SD years) | Physical exercise | Exercise parameters | Motor Learning Category | Exercise Timing to Acquisition | Retention | Exercise effect on motor learning compared to control | |
|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | | Acquisition | Retention |
| Ferrer-Uris et al., 2017 | 29 | Healthy adults (n = 10, 20.9 ± 1.8) | Shuttle run | High 13 min - HIIT 3min at 85% estimated VO2max 2min at 60% estimated VO2max | Motor adaptation | Before | 1h, 24h, 7d | ↔ | ↗ at 1h |
| | | Healthy adults (n = 10, 20.5 ± 1.8) | | | | After | | NA | ↗ at 1h |
| Ferrer-Uris et al., 2018 | 33 | Healthy children (n = 10, 9.2 ± 1.1) | Shuttle run | High 13 min - HIIT 3min at 85% estimated VO2max 2min at 60% estimated VO2max | Motor adaptation | Before | 1h, 24h, 7d | ↔ | ↗ |
| | | Healthy children (n = 12, 9.1 ± 0.8) | | | | After | | NA | ↔ |
| Greeley et al., 2021 | 44 | Healthy older adults (n = 19; 68.3 ± 9.2) | Lower-limb cycling (5 sessions) | High 23 min - HIIT 3min at 75% measured Wpeak 3min at 10W | Sequence | Before | 24h, 35 days | ↔ | ↔ |
| Helm et al., 2017 | 54 | Healthy adults (n = 27; 24.5 ± 2.8) | Upper-limb cycling | High 5 min - HIIT 1 min at 80% age-predicted MHR 1 min at half of cycling power | Motor adaptation | Before | 24h | ↔ | ↔ |
| Holman et al., 2021 | 33 | Healthy adults (n = 17; 21.6 ± 2.3) | Lower-limb cycling | High 5 min - CONT 70% age-predicted HRR | Motor adaptation | After | 30min, 24h, 7d | NA | ↗ at 7d |
| Hubner et al., 2018 | 38 | Healthy older adults (n = 17; 68.2 ± 3.2) | Lower-limb cycling | Moderate 25 min - CONT 60% measured Wpeak | Motor acuity | Before | 30 min, 24h | ↔ | ↔ |
| Hung et al., 2021 | 44 | Healthy adults (n = 22; 23 ± 2) | Lower-limb cycling | High 25 min - HIIT 1 min at 90% measured Wpeak ; 1 min 25% measured Wpeak | Motor acuity | After | 24h | NA | ↔ |
| James and Wang., 2023 | 16 | Healthy adults (n = 16; 24.6 ± 4.7 ) | Lower-limb cycling | Moderate 25 min - CONT 65-70% age-predicted MHR | Motor adaptation | Before | < 30min | ↔ | ↔ |
| Jespersen et al., 2023 | 64 | Healthy adults (n = 16; 24.1 ± 3.9) | Lower-limb cycling | Moderate 20 min - HIIT 3min at 45% measured Wpeak ; 2 min at 50W | Sequence | Before | <5 min, 7d | ↔ | ↗ at 7d |
| | | Healthy adults (n = 16; 23.8 ± 4.2) | | High 20 min - HIIT 3min at 90% measured Wpeak; 2 min at 50W | | After | | NA | ↗ at 7d |
| Khan et al., 2022 | 48 | Healthy adults (n = 23; 22.9 ± 4.9) | Lower-limb cycling | High 20 min - HIIT 3min at 90% measured Wpeak 2min at 50W | Motor acuity | After | 24h, 7d | NA | ↗ at 7d |
| Kuo et al., 2023a | 26 | Healthy adults (n = 26; 27.2 ± 1.0) | Lower-limb cycling | Low 20 min - CONT 40-60% age-predicted MHR Moderate 20 min - CONT 61-74% age-predicted MHR High 20 min - CONT 75-95% age-predicted MHR | Sequence | After | 0-5 min | NA | ↗ for Moderate |
| Kuo et al., 2023b | 20 | Healthy adults (n = 20; 26.4 ± 1.0) | Lower-limb cycling | Moderate 20 min - CONT 61-74% age-predicted MHR | Sequence | After | 0-5 min | NA | ↗ |
| Lehmann et al., 2020 | 31 | Healthy adults (n = 15; 23 ± 7) | Lower-limb cycling (7 sessions) | High 20 min - HIIT 2x3min at 170 (PWC) 4 min at 120 (PWC) | Motor adaptation | Before | 8 weeks | ↗ | ↗ |
| Loras et al., 2020 (no control) | 40 | Healthy adults (n = 40; 23.8 ± 2.0) | Lower-limb cycling | Low 25 min - CONT 50% age-predicted MHR Moderate 25 min - CONT 75% age-predicted MHR | Motor acuity | Before | 24h | ↔ | ↔ |
| Lundbye-Jensen et al., 2017 | 77 | Healthy children (n = 26; 10.4 ± 0.7) | Floorball | High 15 min - HIIT 3 min at high intensity 2 min at low intensity (measured via HR monitoring) | Motor acuity | After | 1h, 24h, 7d | ↔ | ↗ at 7d |
| | | Healthy children (n = 25; 10.8 ± 0.8) | Running | | | After | | | ↘ at 1h ↗ at 7d |
| Mang et al., 2014 | 16 | Healthy adults (n = 16; 23.9 ± 3.7) | Lower-limb cycling | High 20 min - HIIT 3min at 90% measured PPO 2min at 50W | Sequence | Before | 24h | ↗ | ↗ |
| Mang et al., 2016 | 16 | Healthy adults (n = 16; 25.7 ± 4.6) | Lower-limb cycling | High 20min - HIIT 3min at 90% measured PPO 2min at 50W | Sequence | Before | 24h | ↔ | ↗ |

1165
1166
1167
CONT = continuous; HRR = Heart rate reserve; W = Watts; HIIT = high-intensity interval training; HR = Heart Rate; MHR = maximal heart rate; PPO = peak power output; PWC = physical working capacity; Before = exercise was performed before the motor task practice; after = exercise was performed after the motor task practice; min = minutes; h = hours; d = days; NA = not available; ↗ = significant enhancement following exercise; ↘ = significant decrease following exercise ; ↔ = no change following exercise.

1168

1169

1170 **Table S1.** *continued*

| Study | Total N | Type of population (n of exercise groups; age ± SD years) | Physical exercise | Exercise parameters | Motor Learning Category | Exercise Timing to Acquisition | Retention | Exercise effect on motor learning compared to control | |
|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | | Acquisition | Retention |
| Munz et al., 2021 | 12 | Healthy children (n = 12; 10.2 ± 1.4) | Lower-limb cycling | High 40 min – CONT (4 blocks of 10 min) 85-90% of age-predicted MHR | Motor acuity | After | 24h | NA | ↗ |
| Nepveu et al., 2017 | 22 | Stroke patients (n = 11; 64.7 ± 11.6) | Lower-limb cycling | High 15 min - HIIT 3 min at 100% measured Wpeak 2min at 25% measured Wpeak | Motor acuity | After | 24h | NA | ↗ |
| Neva et al., 2019 | 17 | Healthy adults (n = 17; 24 ± 3) | Lower-limb cycling | Moderate 25 min - CONT 65-70% age-predicted MHR | Motor adaptation | Before | 24h | ↗ | ↗ |
| Ostadan et al., 2016 | 38 | Healthy adults (n = 25 ; 22.4 ± 3.9) | Lower-limb cycling | High 15 min - HIIT 3 min at 85-90% measured VO2peak 2 min at 25% measured Wpeak | Sequence | After | 8h | NA | ↔ |
| Opie and Semmler., 2019 | 13 | Healthy adults (n = 13 ; 24.0 ± 3.3) | Lower-limb cycling | Moderate 30 min - CONT 50% age-predicted HRR High 30 min - HIIT 77% age-predicted HRR | Motor acuity | Before | 24h | ↗ for Mod | ↗ for Mod |
| Perini et al., 2016 | 38 | Healthy adults (n = 18; 22.9 ± 2.2) | Lower-limb cycling | Moderate 30 min - CONT 70% measured MHR | Motor acuity | Before | NA | ↗ | NA |
| Pixa et al., 2021 | 27 | Healthy adults (n = 30; 23.7 ± 4.4) | Lower-limb cycling | Low 25 min - CONT 20% measured PPO High 23 min - HIIT 3 min at 90% measured Wpeak 2 min at 60% measured Wpeak | Motor acuity | Before | 15 min, 30min, 24h | ↔ | ↔ |
| Quaney et al., 2009 | 21 | Stroke patients (n = 19; 64.1 ± 12.3) | Lower-limb cycling (24 sessions) | Moderate 45 min - CONT 70% measured MHR | Sequence | Before | 8 weeks | ↗ | ↔ |
| Quinlan et al., 2021 | 29 | Healthy adults (n = 15; 45 ± 5) | Lower-limb cycling | High 20 min - HIIT 4 min at 90% measured Wpeak 2min of active recovery at 50-80W | Motor acuity | Before | 5-7d | ↔ | ↔ |
| Rhee et al., 2016 | 60 | Healthy adults (n = 22; 19.8 ± 1.3) | Lower-limb cycling | High 21 min - CONT 18min at 80% measured MHR 3min at 100% measured MHR | Motor acuity | After | 24h | NA | ↗ |
| Roig et al., 2012 | 48 | Healthy adults (n = 16; 24.1 [21-33]) | Lower-limb cycling | High 15 min – HIIT 3min at high intensity 2min at 50W Based on measured lactate level Workload (200-315) | Motor acuity | Before | 1h, 24h, 7d | ↔ | ↗ for 24h and 7d (higher when exercise is performed after) |
| | | Healthy adults (n = 16; 24.4 [20-30]) | | | | After | | NA | |
| Roig-Hierro et al., 2023 | 32 | Healthy adults (n = 30; 22.9 ± 0.3) | Shuttle run | High 13 min - HIIT 3 min at 80% age-predicted MHR 2min at 60% age-predicted MHR | Motor acuity | After | 7d | NA | ↗ |
| Singh et al., 2016 | 23 | Healthy adults (n = 13; avg = 27 yrs) | Lower-limb cycling | Moderate 20 min - CONT 65-70% age-predicted MHR | Motor adaptation | Before | NA | ↔ | NA |
| Skriver et al., 2014 | 32 | Healthy adults (n = 16; 24.1 ± 3.4) | Lower-limb cycling | High 20 min - HIIT 3min at high intensity 2min at 50W Based on measured lactate level Workload (200-315) | Motor acuity | Before | 1h, 24h, 7d | ↔ | ↗ for 24h and 7d |
| Snow et al., 2016 | 16 | Healthy adults (n = 16; 25.7 ± 3.1) | Lower-limb cycling | Moderate 30 min - CONT 60% measured VO2 peak | Motor acuity | Before | 24h | ↗ | ↔ |
| Statton et al., 2015 | 24 | Healthy adults (n = 8; 22.4 ± 1.1) | Running | High 30 min - CONT 65-85% age-predicted MHR (average was 84% MHR) | Motor acuity | Before | NA | ↗ | NA |
| | 20 | Healthy adults (n = 8; 24.9 ± 1.1) | Running (3 sessions) | | | | 24h | ↗ | ↔ |
| Stavrinos & Coxon 2017 | 24 | Healthy adults (n = 12; 24.8 ± 5.8) | Lower-limb cycling | High 20 min - HIIT 2min at 90% age-predicted HRR 3min at 50% age-predicted HRR | Motor acuity | Before | 5h | ↔ | ↗ |
| Steib et al., 2018 | 17 | Parkinson's disease (n = 17; 64.4 ± 6.2) | Lower-limb cycling | Moderate 30 min - CONT 60-70% measured MHR | Motor adaptation | Before | 24h | ↔ | ↗ |
| Stranda et al., 2019 | 26 | Healthy adults (n = 13; 25.3 ± 2.5) | Lower-limb cycling (12 sessions) | Moderate 20 min - CONT 65% age-predicted MHR | Motor acuity | Before | 7d | ↔ | ↔ |

1171
1172
1173 CONT = continuous; HRR = Heart rate reserve; W = Watts; HIIT = high-intensity interval training; HR = Heart Rate; MHR = maximal heart rate; PPO = peak power output; Before = exercise was performed before the motor task practice; After = exercise was performed after the motor task practice; min = minutes; h = hours; d = days; NA = not available; ↗ = significant enhancement following exercise; ↘ = significant decrease following exercise ; ↔ = no change following exercise.

1174 **Table S1.** *continued*

| Study | Total N | Type of population (n of exercise groups; age ± SD years) | Physical exercise | Exercise parameters | Motor Learning Category | Exercise Timing to Acquisition | Retention | Exercise effect on motor learning compared to control — Acquisition | Retention |
|---|---|---|---|---|---|---|---|---|---|
| Swarbrick et al., 2020 (no control) | 25 | Healthy adults (n = 12; 21.5 ± 2.1) | Lower-limb cycling | Low / 19 min - LIIT / 2 min at 8% Wpeak / 3 min at 12% Wpeak | Motor acuity | Before | 1h, 24h. 7d | ↔ | ↔ |
| | | Healthy adults (n = 13; 22.2 ± 3.1) | | High / 19 min - HIIT / 2 min at 60% measured Wpeak / 3 min at 90% measured Wpeak | | | | | |
| Taylor et al., 2024 (Active rest) | 23 | Healthy older adults (n = 11; 65.3 ± 6.3) | Lower-limb cycling | High / 20 min - HIIT / 3 min at 50% measured HRR / 2 min at 90% measured HRR | Motor acuity | Before | 6h | ↘ | ↗ |
| Thomas et al., 2016 | 36 | Healthy adults (n = 12; 23.5 ± 2.3) | Lower-limb cycling | Low / 17 min - LIIT / 2 min at 25% measured Wpeak / 3 min at 45% measured Wpeak | Motor acuity | After | 24h, 7d | NA | ↗ for High at 24h / ↗ for Low and High at 7d |
| | | Healthy adults (n = 12; 24.3 ± 2.3) | | High / 17 min - HIIT / 2 min at 60% measured Wpeak / 3 min at 90% measured Wpeak | | | | | |
| Thomas et al., 2017 | 40 | Healthy adults (n = 10; 24.7 ± 3.5) | Strength training | High / 45-50 min / Resistance training program / (4 RM and 8 RM) | Motor acuity | After | 1h, 24h | NA | ↘ at 1h / ↗ at 1d |
| | | Healthy adults (n = 10; 25.9 ± 4.5) | Circuit training | High / 45-50 min / (30% of 1 RM) | | After | | | |
| | | Healthy adults (n = 10; 25.9 ± 3.7) | Hockey training | High / 48 min / 5 x 7 min; 2 min rest | | After | | | |
| Thompson et al., 2023 | 45 | Stroke patients (n = 15; 68.1 ± 7.1) | Whole-body cycling | High / 5 min- CONT / 70-85% age-predicted MHR | Motor adaptation | After | 24h | NA | ↔ |
| | | Stroke patients (n = 15; 61.8 ± 7.5) | | High / 15 min - HIIT / 3 min at 70%-85% age-predicted MHR / 2 min at < 70% age-predicted MHR | | | | | |
| Tomporowski and Pendleton, 2018 | 32 | Healthy adults (n = 11; 22.4 ± 3.0) | Dancing | Moderate / 10 min - CONT / Simple Dance Dance Revolution / Intensity determined based on HR | Motor acuity | Before | 13min, 24h, 7d | ↔ | ↔ |
| | | Healthy adults (n = 10; 22.9 ± 2.9) | | Moderate / 10 min - CONT / Complex Dance Dance Revolution / Intensity determined based on HR | | After | | NA | ↗ at 7d |
| Wanga et al., 2023 | 33 | Healthy adults (n = 11; 20.5 ± 2.3) | Lower-limb cycling | Low / 23 min - HIIT / 3min at 45% measured Wpeak / 2min at 25% measured Wpeak | Sequence | After | 24h, 48h | NA | ↗ (higher for Low) |
| | | Healthy adults (n = 11; 20.3 ± 2.3) | | High / 23 min - HIIT / 2min at 60% measured Wpeak / 3min at 90% measured Wpeak | | | | | |
| Wanner et al., 2020 | 50 | Healthy adults (n = 17; 25.3 ± 2.7) | Lower-limb cycling | Low / 17 min - CONT / 25W | Motor adaptation | Before | 24h | ↔ | ↔ |
| | | Healthy adults (n = 18; 25.1 ± 2.3) | | Moderate / 17 min - HIIT / 2min at 25% measured Wpeak / 3min at 45% measured Wpeak | | | | | |
| | | Healthy adults (n = 15; 25.7 ± 3.6) | | High / 17 min - HIIT / 2min at 60% measured Wpeak / 3min at 90% measured Wpeak | | | | | |
| Wanner et al., 2021 | 17 | Parkinson's disease (n = 8; 59.3 ± 5.4) | Lower-limb cycling | Moderate / 25 min - CONT / 60% maximal Wpeak | Motor adaptation | Before | 1d, 7d | ↔ | ↗ at 7d |
| Williams et al., 1985 | 90 | Healthy adults (n = 30; 20.9 ± 1.9) | Arm cranking | Low / 5 min - CONT / WO maintained at 10W (30 rpm) / (110-125 bpm) | Motor acuity | Before | 24h | ↔ | ↔ |
| | | Healthy adults (n = 30; 20.9 ± 1.9) | | High / 5 min - CONT / WO maintained at 75W (90 rpm) / (165-180 bpm) | | | | | |
| Yildirim et al., 2024 | 70 | Healthy adults (n = 15; 20.5 ± 0.7) | Running | Low / 30 min- CONT / 57%-63% estimated MHR | Motor acuity | Before | 24h, 7d | ↔ | ↔ |
| | | Healthy adults (n = 15; 20.1 ± 0.5) | | | | After | | ↔ | ↔ |
| | | Healthy adults (n = 15; 20.8 ± 1.5) | | Moderate / 30 min- CONT / 64%-76% estimated MHR | | Before | | ↔ | ↔ |
| | | Healthy adults (n = 15; 20.9 ± 1.0) | | | | After | | ↔ | ↔ |

1175 CONT = continuous; HRR = Heart rate reserve; W = Watts; HIIT = high-intensity interval training; LIIT = light-intensity interval training; HR = Heart Rate; MHR = maximal
1176 heart rate; PPO = peak power output; rpm = revolutions per minute; bpm = beats per minute; RM = repetition maximum; Before = exercise was performed before the motor task
1177 practice; After = exercise was performed after the motor task practice; min = minutes; h = hours; d = days; NA = not available; ↗ = significant enhancement following exercise; ↘
1178 = significant decrease following exercise ; ↔ = no change following exercise.

# Supplementary Material 2

1179

## <u>Search equations</u>

1180

1181

**Web of Science Core Collection : A&HCI , ESCI , CPCI-SSH , CPCI-S , SCI-EXPANDED , SSCI**

1182

1183

**(exercise OR training) NEAR/5 ( "motor learning" OR "motor skill" OR "motor skills" OR "motor adaptation" OR "motor acuity" OR "motor coordination" OR "de novo learning" OR "motor control" OR "motor sequence" OR "motor behavior" OR "motor behaviour" OR "sensorimotor learning" OR "motor competence" OR "motor competences") (Topic)**

1184
1185
1186
1187

1188

-

1189

1190

**ERIC (ProQuest)**

1191

1192

((MAINSUBJECT.EXACT("Perceptual Motor Learning") OR MAINSUBJECT.EXACT("Movement Education") OR MAINSUBJECT.EXACT("Perceptual Motor Coordination") OR MAINSUBJECT.EXACT("Multisensory Learning") OR MAINSUBJECT.EXACT("Psychomotor Skills")) AND MAINSUBJECT.EXACT("Exercise")) AND PEER(yes)

1193
1194
1195

**OR**

1196

noft((exercise OR training) NEAR/5 ("motor learning" OR "motor skill" OR "motor skills" OR "motor adaptation" OR "motor acuity" OR "motor coordination" OR "de novo learning" OR "motor control" OR "motor sequence" OR "motor behavior" OR "motor behaviour" OR "sensorimotor learning" OR "motor competence" OR "motor competences")) AND PEER(yes)

1197
1198
1199
1200

1201

Limité par : Revu par les pairs

1202

1203

-

1204

**SPORTDiscus with Full Text**

1205

| # | Question |
|---|---|
| S7 | S5 OR S6<br>Opérateurs de restriction - Relu par un comité de lecture; Type de publication: Academic Journal<br>Opérateurs d'expansion - Appliquer des sujets équivalents<br>Modes de recherche - Booléen/Phrase |
| S6 | TI ( exercise OR training) N5 ( "motor learning" OR "motor skill" OR "motor skills" OR "motor adaptation" OR "motor acuity" OR "motor coordination" OR "de novo learning" OR "motor control" OR "motor sequence" OR "motor behavior" OR "motor behaviour" OR "sensorimotor learning" OR "motor competence" OR "motor competences") ) OR AB ( exercise OR training) N5 ( "motor learning" OR "motor skill" OR "motor skills" OR "motor adaptation" OR "motor acuity" OR "motor coordination" OR "de novo learning" OR "motor control" OR "motor sequence" OR "motor behavior" OR "motor behaviour" OR "sensorimotor learning" OR "motor competence" OR "motor competences") ) |
| S5 | S3 AND S4 |
| S4 | DE "MOTOR learning" OR DE "PERCEPTUAL motor learning" |
| S3 | S1 OR S2 |

S2         DE "EXERCISE" +

S1         DE "PHYSICAL training & conditioning" +

1206

1207 -

1208 Database(s): **APA PsycInfo** 1806 to March 2025 Week 2

1209 Search Strategy:

| # | Searches |
|---|---|
| 1 | athletic training/ |
| 2 | exp exercise/ |
| 3 | 1 or 2 |
| 4 | Motor Coordination/ or Motor Skills/ or Motor Control/ or Perceptual Motor Coordination/ or Perceptual Motor Learning/ or Gross Motor Skill Learning/ or Fine Motor Skill Learning/ |
| 5 | 3 and 4 |
| 6 | ((exercise or "physical training" or "endurance training" or "resistance training") adj6 ("motor learning" or "motor skill" or "motor skills" or "motor adaptation" or "motor acuity" or "motor coordination" or "de novo learning" or "motor control" or "motor sequence" or "motor behavior" or "motor behaviour" or "sensorimotor learning" or "motor competence" or "motor competences")).ab,id,ti. |
| 7 | 5 or 6 |
| 8 | limit 7 to human |

1210

1211 -

1212 Database(s): **Ovid MEDLINE(R) ALL** 1946 to March 18, 2025

1213 Search Strategy:

| # | Searches |
|---|---|
| 1 | ((exercise or "physical training" or "endurance training" or "resistance training") adj6 ("motor learning" or "motor skill" or "motor skills" or "motor adaptation" or "motor acuity" or "motor coordination" or "de novo learning" or "motor control" or "motor sequence" or "motor behavior" or "motor behaviour" or "sensorimotor learning" or "motor competence" or "motor competences")).ab,kf,ti. |
| 2 | exp animals/ not humans/ |
| 3 | 1 not 2 |

1214 -

1215

1216

1217 Database(s): **Embase** 1974 to 2025 March 18

1218 Search Strategy:

| # | Searches |
|---|---|
| 1 | ((exercise or "physical training" or "endurance training" or "resistance training") adj6 ("motor learning" or "motor skill" or "motor skills" or "motor adaptation" or "motor acuity" or "motor coordination" or "de novo learning" or "motor control" or "motor sequence" or "motor behavior" or "motor behaviour" or "sensorimotor learning" or "motor competence" or "motor competences")).ab,kf,ti. |
| 2 | exp exercise/ |
| 3 | resistance training/ or endurance training/ |
| 4 | 2 or 3 |
| 5 | motor learning/ |
| 6 | 4 and 5 |
| 7 | 1 or 6 |
| 8 | (exp animal/ or nonhuman/) not exp human/ |
| 9 | 7 not 8 |

1219

1220

# Supplementary Material 3

Data extraction sheet is attached to this manuscript on BioRxiv.