A Drosophila Tonic Motor Neuron Reinnervates Ectopic Muscles Fully

Deprived of Native Tonic and Phasic Inputs

Lizzy Olsen[1,2,3], Parinita Mitchelle Mandhyan[1,2], Sara Arredondo[2], Komal Kaur[1,2], Ankura Sitaula[1,2], Aref

Zarin[2,3*]

[1]Biology Graduate Program, Texas A&M University, College Station, United States

[2]Department of Biology, Texas A&M University, College Station, United States

[3]Texas A&M Institute for Neuroscience, Texas A&M University, College Station, United States

*For correspondence: azarin@bio.tamu.edu

Abstract

Motor neurons (MNs) form precise neuromuscular junctions (NMJs) during

development, but the extent to which individual MNs can reinnervate fully denervated

muscles in vivo remains poorly understood. In the Drosophila larva, each muscle is co-

innervated by a tonic and a phasic glutamatergic MN. Here, we show that the tonic MN1

undergoes robust heterosynaptic sprouting and forms ectopic NMJs when neighboring

muscles are deprived of both their native tonic and phasic inputs. This structural plasticity

is not induced by silencing, but instead requires the physical ablation of adjacent MNs.

Live imaging of the same MN1 axons in individual animals reveals that sprouting

initiates early and expands progressively across larval stages. In contrast, phasic MNs

show minimal remodeling, indicating that tonic MNs possess a greater intrinsic capacity

for neuroplasticity. Notably, as MN1 establishes new synapses on targets it does not

normally innervate, it redistributes pre-synaptic territory across both native and ectopic

muscles. These findings identify a subtype-specific, injury-induced rewiring program in

an intact motor circuit.

Introduction

Spinal cord injury, stroke, and neurodegenerative disease disrupt neural circuits, impairing motor

and cognitive function. Functional recovery may depend on heterosynaptic structural plasticity

(HTSP), in which intact neurons sprout new branches to reinnervate denervated targets (1-3).

Therapeutic strategies aimed at enhancing HTSP hold promise, but their success hinges on a

deeper understanding of the intrinsic and extrinsic factors that promote or constrain neuronal

sprouting (4). Neurons differ in properties such as firing dynamics (e.g., tonic versus phasic),

neurotransmitter identity, and synaptic architecture (e.g., large versus small boutons), but it

remains unclear whether all neuron types are equally capable of structural remodeling after

injury. This gap limits our ability to predict which neurons can compensate for lost inputs—and

under what conditions. A central question, therefore, is how physiologically distinct neurons

respond to the dysfunction or loss of their synaptic partners. Here, we examine the capacity for

post-injury axonal sprouting and neuromuscular junction (NMJ) remodeling in physiologically

distinct tonic and phasic motor neurons in Drosophila larvae.

The *Drosophila* larval neuromuscular system provides a genetically accessible and anatomically

tractable model for investigating neuron subtype-specific injury responses. Larval muscles are

innervated by two classes of glutamatergic excitatory motor neurons (MNs): tonic-firing type Ib

MNs, which form large boutons and typically innervate a single muscle, and phasic-firing type Is

MNs, which form smaller boutons and innervate multiple muscles (5-12). These MNs converge

on shared target muscles, creating a synaptic motif in which one tonic and one phasic MN co-

innervate the same postsynaptic muscle fiber. This configuration provides a unique opportunity to

directly compare how distinct MN subtypes respond to the same insult. Notably, convergence of

tonic and phasic inputs onto a common postsynaptic cell is an evolutionarily conserved circuit

motif observed across both vertebrate and invertebrate systems (13-16). Due to its simplicity and

powerful genetic toolkit, the *Drosophila* NMJ has long served as a leading model for dissecting

conserved mechanisms of synaptic development and plasticity at single-cell resolution (5, 6, 8,

10, 17-29).

Previous studies have shown that ablating—but not silencing—the phasic MN RP2 induces

structural expansion of the NMJ formed by its co-innervating tonic partner MN1 on Muscle 1,

while loss of MN1 has no effect on RP2 (7-10). These findings suggest that tonic MNs are more

structurally adaptive than phasic MNs. However, since only one MN type was disrupted in those

studies, the muscles retained partial excitatory input. Thus, whether complete denervation

prompts more robust or qualitatively different plastic responses—and which neurons are capable

of compensating—remains unknown.

To address this, we developed a genetic paradigm to simultaneously ablate both tonic and phasic MNs that co-innervate dorsal longitudinal muscles, creating a fully denervated group of target muscles. We found that the surviving tonic MN1, which normally innervates Muscle 1, exhibited robust axonal sprouting and formed ectopic NMJs on denervated muscles it does not typically target. In contrast, the phasic MN RP2 did not sprout following ablation of its tonic partners, and MN1 failed to sprout when neighboring MNs were chronically or acutely silenced rather than ablated. These results demonstrate that injury-induced HTSP is neuron-subtype specific and requires complete physical loss of input—not merely loss of activity—to initiate structural remodeling.

We further show that the ectopic NMJs formed by MN1 are functional: they restore calcium transients in previously denervated muscles and partially rescue crawling performance. Using live motor axon imaging, behavioral assays, and temporally controlled genetic ablation, we mapped the dynamics of MN1 sprouting across larval stages. Sprouting was initiated shortly after larval hatching and continued throughout larval development, revealing a broader plasticity window than previously appreciated.

Together, our findings identify a MN subtype-specific and injury-dependent form of HTSP that enables functional circuit compensation. This work uncovers key cellular and temporal principles governing adaptive circuit remodeling and provides a foundation for understanding how defined neurons sense local denervation and initiate compensatory growth programs.

Results

Tonic MN1 Sprouts and Forms Ectopic NMJs Following Complete Denervation of Neighboring Muscles

The *Drosophila* larval body comprises three thoracic segments (T1–T3) and nine abdominal segments (A1–A9). The central nervous system (CNS) consists of two brain lobes and a segmented ventral nerve cord (VNC), which functions analogously to the vertebrate spinal cord. Each VNC segment controls the muscles of its corresponding

body segment (**Fig. 1A**). In each abdominal segment, approximately 30 bilaterally

symmetrical muscle pairs are innervated by 28 pairs of tonic and two pairs of phasic

glutamatergic excitatory motor neurons (MNs) projecting from the segmental VNC (5,

30-36) (**Fig. 1A**).

In this study, we focused on the dorsal-longitudinal (DL) muscles, specifically Muscles 1,

2, 9, and 10 (**Fig. 1A**). These muscles receive inputs from a common phasic-firing MN-

Is, also known as RP2 or the common dorsal excitor (CDE), and four distinct tonic-firing

(Ib) MNs—MN1, MN2, MN9, and MN10—each innervating an individual muscle (**Fig.**

1A) (5, 30-36).

Previous studies have shown that apoptotic ablation of the phasic MN-Is (hereafter

referred to as **Is**) leads to structural expansion of the neuromuscular junction (NMJ)

formed by the co-innervating tonic MN1, increasing tonic input to Muscle 1. In contrast,

ablation of the tonic MN1 does not trigger structural changes in the NMJ formed by MN-

Is (7-10). These findings highlight intrinsic differences in how tonic and phasic MNs

respond to the loss of their co-innervating partners. However, these studies removed only

one MN class—either tonic or phasic—leaving the target muscles partially innervated.

Thus, it remained unclear whether complete removal of both excitatory inputs could

trigger compensatory sprouting by other MNs.

To address this, we genetically ablated either or both tonic and phasic MNs innervating

Muscles 2, 9, and 10 and examined structural remodeling by the bystander MN1 (**Fig.**

1B–D). For ablation, we used previously validated MN-LexA drivers (37, 38) to express

the pro-apoptotic genes *reaper (rpr)* and *head involution defective (hid)*, inducing cell

death in targeted MNs (39). The tonic MN1 was visualized using 94G06 (40) (MN1 Ib)-

Gal4 driving UAS-myr-GFP.

We found that MN1 did not form ectopic NMJs when only the phasic MN-Is was ablated

(Is >RPR) and only formed ectopic NMJs on Muscle 2 when tonic MNs 2, 9, and 10 were

ablated (CQ >RPR) (**Fig. 1 D,E**). In contrast, concurrent ablation of both tonic and phasic

MNs (CQ ∪ Is > RPR) triggered robust sprouting of MN1, which extended axons and

formed de novo NMJs on fully denervated Muscles 2, 9, and 10 (**Fig. 1 D,E**).

Quantitative morphometric analysis (41, 42) (**Fig. 1 C**) revealed that MN1 in CQ ∪ Is

>RPR larvae showed a significant increase in overall NMJ area across all four DL

muscles (1, 2, 9, and 10) compared to controls, while average bouton size and total

bouton number remained unchanged (**Fig. 1 D,E**). Importantly, the total MN1 NMJ area

was redistributed across the DL muscles, with reduced allocation to its native target

Muscle 1 compared to controls—suggesting an upper limit on the total synaptic area

MN1 can establish (**Fig. 1 D,E**). Finally, we quantified total axonal and NMJ branch

points. CQ > RPR and CQ ∪ Is > RPR larvae showed a significant increase in axon

branch points (GFP-only), consistent with a sprouting phenotype, whereas NMJ branch

points (Dlg+GFP) were not significantly altered (**Fig. 1 D,E**).

Together, these results demonstrate that robust heterosynaptic sprouting of MN1 is

selectively triggered by complete loss of both tonic and phasic excitatory inputs to

neighboring muscles.

Phasic MN-Is Maintains Its Normal Innervation After Ablation of Co-innervating

Tonic MNs

Our earlier experiments demonstrated that the tonic MN1 exhibits robust heterosynaptic

sprouting when neighboring muscles are fully deprived of both tonic and phasic inputs.

We next asked whether the phasic MN-Is possesses a comparable capacity for structural

remodeling and NMJ expansion following the loss of its tonic co-innervators.

Previous studies, in which only MN1 was ablated, reported no structural changes in

NMJs formed by the phasic MN-Is on Muscle 1 (7-10). To extend these findings, we

ablated additional tonic MNs (MN1, MN2, MN9, and MN10), thereby fully depriving the

dorsal-longitudinal (DL) muscles 1, 2, 9, and 10 of all tonic innervations (**Fig. 2A**). In

both control and ablated groups, the phasic MN-Is was visualized using myr-GFP driven

by an enhancer element cloned from the 27E09-Gal4 driver, which selectively targets

phasic MNs.

Following tonic MN ablation (CQ ∪ MN1 >RPR), the intact phasic MN-Is retained its

normal innervation pattern on Muscles 1, 2, 9, and 10 (**Fig. 2A**). Apart from a modest

reduction in the number of 2nd- and 3rd-order axonal branches in CQ ∪ MN1 >RPR

larvae, there were no significant differences in bouton number, area, or size between

control and ablated groups (**Fig. 2B, C**). Despite complete loss of tonic excitatory input,

the phasic MN-Is did not exhibit ectopic sprouting or NMJ expansion on these muscles.

This lack of response contrasts sharply with the extensive sprouting observed in MN1

under similar conditions, underscoring a fundamental difference in the plasticity potential

of tonic versus phasic MNs. Consistent with earlier studies (7-10), these results suggest

that phasic MN-Is lacks the intrinsic capacity for compensatory structural remodeling

following the loss of its co-innervating tonic partners.

MN1 Sprouting is Induced by Ablation but Not by Chronic or Acute Silencing of

Neighboring MNs

To confirm that the robust sprouting observed in MN1 results specifically from the loss

of neighboring motor neurons—and not from artifacts of the LexA/AOP genetic

system—we repeated the ablation experiments using the Gal80[ts]/GAL4/UAS system.

Gal80[ts] inhibits GAL4 activity at 24°C but not at 29°C (43, 44).

Using previously characterized CQ and Is GAL4 drivers (7, 36-38, 40, 45), we targeted

the tonic MN2, MN9, and MN10 along with the phasic MN-Is for UAS-reaper (rpr)-

mediated ablation. MN1 was labeled with MN1-lexA >AOP-myr-GFP. Animals raised

continuously at 29°C showed robust MN1 sprouting and ectopic NMJs on denervated

Muscles 2, 9, and 10, whereas controls maintained at 24°C (with functional Gal80 protein

blocking GAL4 activity) showed no sprouting (**Fig. 3A-B, Supp Fig S1**). These findings

confirm that MN1 sprouting is specifically triggered by the physical loss of neighboring

MNs rather than by potential genetic artifacts.

We next tested whether silencing the activity of neighboring MNs—without their

physical removal—would be sufficient to induce MN1 sprouting. For acute silencing, we

used the light-activated chloride channel GtACR1, which causes chloride influx and

neuronal hyperpolarization upon light exposure (46, 47). Previous works have

demonstrated the efficacy of GtACR1 in silencing MNs and other neurons in Drosophila

larvae (37, 45, 48, 49). In CQ ∪ Is >GtACR1 larvae, MN1 maintained its normal

innervation of Muscle 1 and did not form ectopic NMJs on Muscles 2, 9, or 10. However,

we observed a modest increase in MN1 NMJ area and bouton number compared to

controls, without changes in bouton size (**Fig. 3A-B, Supp Fig S1**).

We then tested chronic silencing using botulinum neurotoxin-C (BoNT-C), which blocks

both evoked and spontaneous neurotransmitter release (27, 28). In CQ ∪ Is >BoNT-C

larvae, MN1 did not form ectopic NMJs, but the number of boutons on Muscle 1 was

significantly increased relative to controls, despite no changes in NMJ area or bouton

size. (**Fig. 3A-B Supp Fig S1**).

We then expressed tetanus toxin light chain (TeTxLC), which blocks evoked but not

spontaneous neurotransmitter release (50). In CQ ∪ Is >TeTxLC larvae, MN1 again did

not form ectopic NMJs, but showed substantial NMJ expansion on its native target,

Muscle 1 (**Fig 3 A-C Supp Fig S1**). This included significantly increased NMJ area,

branching, and bouton number, along with a marked reduction in average bouton size

(**Fig 3 B-C Supp Fig S1**). These features are consistent with satellite bouton formation

previously observed following TeTxLC-mediated silencing of MN-Is (7, 8). Notably, Dlg

staining revealed that the NMJs of TeTxLC-silenced tonic MNs maintained normal

morphology and did not exhibit the extensive NMJ overgrowth observed in the bystander

MN1 (**Fig. 3A)**, suggesting that the sprouting phenotype is specific to the unaffected

neuron responding to loss of presynaptic activity in neighboring inputs.

Taken together, these results demonstrate that heterosynaptic sprouting of MN1 requires
the physical ablation of neighboring MNs. In contrast, silencing of neighboring MNs—
whether acute or chronic—does not trigger ectopic sprouting, though it can modulate
bouton number and NMJ structure at the native target.
208

Ectopic MN1 Sprouting Restores Muscle Activation and Partially Rescues Crawling
Behavior

Thus far, we have shown that concurrent ablation—but not silencing—of neighboring
tonic and phasic MNs triggers robust sprouting of the bystander MN1, resulting in
ectopic NMJs on fully denervated Muscles 2, 9, and 10. To determine whether these
ectopic NMJs are functional and capable of activating the muscles they innervate, we
performed in vivo muscle calcium imaging in intact, forward-crawling larvae.

As a baseline, we first examined muscle calcium activity in larvae where all tonic and
phasic MNs innervating dorsal muscles 1, 2, 9, and 10 were either genetically ablated
(CQ-lexA, 27E09-lexA; 94G06-lexA >RPR) or acutely silenced (CQ-lexA, 27E09-lexA;
94G06-lexA >GtACR1) (**Fig 4 A-C**). In both groups, the dorsal-longitudinal (DL)
muscles failed to show calcium transients or contraction during forward crawling,
confirming complete loss of excitatory input. In contrast, ventral and lateral muscles with
intact MN innervation exhibited robust calcium activity and contractions (**Fig 4 A,B;**
Video S1, Video S2, Video S3). These observations validate that (i) both GtACR1-
mediated silencing and RPR-mediated ablation effectively eliminate MN output and (ii)
the MN driver lines selectively target the intended MNs without affecting off-target MNs.

We then repeated these experiments but spared MN1. In the acute silencing group (CQ-
lexA, 27E09-lexA >GtACR1), only Muscle 1 displayed calcium transients and
contraction during crawling, consistent with MN1 maintaining its normal innervation
(**Fig 4 A,B; Video S4**). By contrast, in the ablation group (CQ-lexA, 27E09-lexA
>RPR), all four dorsal muscles (1, 2, 9, and 10) exhibited calcium transients and
contraction, demonstrating that ectopic NMJs formed by MN1 are capable of restoring
contractile activity in denervated muscles (**Fig 4 A,B; Video S5**).

To directly test whether the sprouted MN1 drives these responses, we acutely silenced

MN1 (94G06-Gal4 >GtACR1) in larvae where neighboring MNs had been ablated (CQ-

lexA, 27E09-lexA >RPR). Silencing MN1 abolished calcium activity in all four dorsal

muscles during crawling, confirming that the sprouted MN1 terminals are functionally

active and required for muscle contraction (**Fig 4 A,B; Video S6**).

Next, we investigated whether MN1 sprouting contributes to behavioral recovery. We

quantified forward crawling by measuring peristalsis efficiency (distance traveled per

peristalsis) and peristalsis duration (time to complete a forward wave). Control larvae

displayed a peristalsis efficiency of ~0.8 mm per wave. Larvae lacking dorsal MN inputs

(CQ-lexA, 27E09-lexA; 94G06-lexA >RPR) showed markedly reduced efficiency, but

those with intact MN1 (CQ-lexA, 27E09-lexA >RPR) performed significantly better,

indicating partial restoration of crawling (**Fig 4 D,E; Video S7**).

Notably, CQ-lexA, 27E09-lexA >RPR larvae also outperformed CQ-lexA, 27E09-lexA

>GtACR1 larvae (**Fig 4 D,E; Video S7**), suggesting that ablation-induced sprouting of

MN1 (occurring in the ablated group but not in the silenced group) underlies this

improvement. However, because 27E09-lexA also targets ventral-projecting phasic MNs,

it is possible that structural plasticity of ventral tonic MNs contributed to recovery. To

isolate MN1's contribution, we silenced the sprouted MN1 in CQ-lexA, 27E09-lexA

>RPR larvae using 94G06-Gal4 >GtACR1. This manipulation significantly reduced

peristalsis efficiency compared to unsilenced CQ-lexA, 27E09-lexA >RPR larvae (**Fig 4**

D,E; Video S7), directly demonstrating that MN1 sprouting is adaptive and improves

locomotion.

Across all genotypes, we observed an inverse relationship between peristalsis efficiency

and peristalsis duration: larvae traveling shorter distances per peristaltic wave also

exhibited slower propagation of contractions across body segments, resulting in longer

peristalsis durations (**Fig 4 E; Video S7**).

Together, these results establish that MN1 ectopic sprouting restores muscle contractility
and partially rescues motor behavior, highlighting its role as a compensatory response
following injury.

Ectopic MN1 Sprouting Occurs Primarily During Larval Stages

In *Drosophila*, embryogenesis lasts ~22–24 hours at 25°C, after which the embryo
hatches into a first instar larva (L1) capable of foraging, feeding, and performing
locomotor behaviors such as forward crawling. By hatching, core motor circuits are
already established, with motor neurons forming functional synapses onto their target
muscles. The L1 stage lasts ~25 hours, followed by molting into second (L2) and third
instar (L3) stages, each lasting ~24–25 hours before pupation. Our initial analyses of
MN1 sprouting and ectopic NMJ formation were performed in L3 larvae. To better define
the timeline of these structural changes, we examined MN1 remodeling across the L1–L3
transition in animals with ablated tonic and phasic MNs.

Using live imaging of MN1 labeled with tdTomato, we tracked axonal branching in
individual larvae at L1, L2, and L3 stages. Between imaging sessions, larvae were
recovered and returned to food to continue development. This allowed us to quantify and
distinguish 1st-, 2nd-, and 3rd-order axon branches extending from MN1's main shaft
(**Fig. 5A schematic**).

At L1, MN1 axons in ablated larvae already exhibited a twofold increase in 1st-order
branches compared to controls, while 2nd-order branching was comparable between
groups. By L3, ablated larvae showed significantly higher numbers of 1st-order branches
relative to controls. Within the ablated group, both 1st- and 2nd-order branch numbers
increased significantly from L1 to L3, whereas in controls only 2nd-order branching
increased across development. These findings suggest that while MN1 sprouting in
ablated animals is initiated early, the majority of axon branching and ectopic NMJ
formation occurs progressively during the L1–L3 transition (**Fig. 5A–B; Supp. Fig.**).

To test whether MN1 sprouting can be induced exclusively during larval stages, we performed temporally restricted ablation. Since MN-Gal4 drivers (e.g., CQ-Gal4, 27E09-Gal4) activate UAS-rpr/hid from embryogenesis onward, we used Gal80^{ts} to suppress Gal4 activity at 24°C and permit it at 29°C (43, 44). For larval-only ablation, embryos were maintained at 24°C until hatching and then shifted to 29°C from L1 through L3 prior to NMJ analysis. As reference points, we used data from embryonic ablation (continuous 29°C) and control animals (continuous 24°C) described earlier in **Fig. 3**. Larvae subjected to larval-stage ablation showed MN1 sprouting and ectopic NMJ formation, though to a lesser extent than embryonically ablated animals (**Fig. 5C–D**).

Taken together, these results indicate that MN1 sprouting is initiated early but progresses substantially throughout larval development, with the L1–L3 period representing a key window for ectopic NMJ formation following injury.

Discussion

Subtype-Specific Plasticity Revealed Through Complete Denervation

Our study reveals a robust, subtype-specific capacity for structural plasticity within an established motor circuit following injury. We demonstrate that tonic MN1, but not phasic MN-Is, undergoes extensive axonal sprouting and ectopic NMJ formation when neighboring tonic and phasic MNs are ablated. These ectopic NMJs restore muscle contractility and contribute to improved locomotor performance, providing functional evidence that the observed plasticity is adaptive. In contrast, when neighboring MNs are functionally silenced without ablation, MN1 does not sprout, indicating that complete physical loss—not merely loss of synaptic activity—is required to trigger heterosynaptic structural plasticity (HTSP).

Our approach builds on earlier observations by Chang and Keshishian (1996), who reported that embryonic denervation could lead to some ectopic innervation from nearby neurons (51). However, they utilized laser-mediated MN ablation, which resulted in variable levels of denervation and often lacked control over which neurons contributed to reinnervation. In contrast, we used targeted genetic ablation to achieve reproducible, complete denervation of specific muscles, revealing that sprouting arises specifically from the identified tonic MN1. Furthermore,

we show that the newly formed NMJs are functional and that sprouting mainly occurs after larval

hatching and continues across larval stages, defining a broader temporal window for plasticity.

In more recent studies investigating NMJ plasticity, either tonic or phasic MNs were disrupted,

but never both. As a result, target muscles continued to receive partial excitatory input from

surviving MNs, allowing for homeostatic compensatory changes within an already innervated

context. Thus, the effects of completely removing all excitatory input from a given muscle and

testing whether other MNs can sprout to reinnervate these fully denervated targets remained

unknown. To address this gap, we developed genetic tools enabling the simultaneous ablation of

both tonic and phasic MNs innervating specific muscles, creating a fully denervated environment.

This strategy revealed a previously unrecognized capacity for bystander MN1 to form de novo

ectopic NMJs and reestablish motor output. This subtype specificity aligns with prior findings

that tonic Ib MNs—but not phasic Is MNs—exhibit structural and functional compensation

following input loss (7, 8, 10). Our findings extend this framework by demonstrating that MN1

can not only expand existing synaptic terminals but also innervate novel postsynaptic targets,

forming functional ectopic NMJs on muscles it does not typically contact. This level of

remodeling goes beyond classic homeostatic plasticity and reflects a more profound form of

structural reorganization.

Importantly, MN1 sprouting occurred only following the physical ablation of neighboring MNs

and not when those MNs were chronically or acutely silenced. Our findings further support that

tonic MNs possess a greater intrinsic capacity for structural remodeling in response to injury

compared to phasic MNs. This finding agrees with Han et al. (2022), who showed that botulinum

toxin silencing—even when eliminating both evoked and spontaneous transmission—does not

induce structural compensation, whereas physical loss does (27). These results emphasize that

circuit rewiring is not merely a response to inactivity but likely requires injury-associated cues

such as axonal degeneration or glial signaling.

Our injury model appears to diverge from CNS critical period constraints. Ackerman et al. (2021)

silenced MN1 and MN-Is and examined dendritic remodeling within the same MNs, identifying

an early embryonic window during which activity loss induces dendritic growth (52). In contrast,

we demonstrate that MN1 can undergo peripheral axonal remodeling beginning 0–4 h after larval

hatching and continuing throughout larval stages, highlighting that different compartments of the

same neuron (dendrites vs. axons) follow distinct temporal rules for plasticity.

Intrinsic Differences in Plasticity Between Tonic MN1 and Phasic MNs

MN1 displays a markedly greater intrinsic capacity for structural plasticity compared to its phasic

counterparts, such as the phasic MN-Is. This enhanced plasticity is likely underpinned by a

multilayered set of molecular, structural, genetic, and functional differences at the synaptic,

transcriptional, and cellular level (28, 38, 53, 54).

Structurally, tonic MN1 forms larger synaptic boutons with expansive, donut-shaped

active zones and a 4.4-fold larger neuromuscular junction area at muscle 1 containing

3.8-fold more active zones (AZ), creating an ideal substrate for growth and remodeling,

unlike the compact active zones of phasic neurons optimized for rapid transmission rather

than plasticity (13, 27, 28, 38, 54). Functionally, tonic MN1 displays lower basal synaptic

output and more "silent" active zones possibly conducive to sprouting, whereas phasic

MN-Is show high initial release probability (Pr) that depresses with repetitive stimulation

(54). Importantly, high-Pr phasic MN-Is synapses show denser voltage-gated calcium channel

(VGCC) clustering and tighter **vesicle** coupling—architectural features that could enhance

release efficiency but may potentially constrain structural remodeling and growth (54).

At the molecular level, MN1 expresses significantly elevated levels of growth-promoting genes

including Wnt4 (40-fold higher), Pyramus, Toll-6, and SiaT—establishing a growth-permissive

transcriptional program and environment (38). Conversely, phasic neurons upregulate calcium

buffering proteins like Cbp53E (30-fold higher) and vesicle trafficking chaperones that constrain

remodeling (38). MN1's superior calcium handling—maintaining higher resting calcium levels

while enabling rapid extrusion after activity—prevents growth inhibition and supports sprouting,

unlike phasic neurons that accumulate excessive calcium during stimulation with slower

recovery. Furthermore, MN1's distinct post-translational modifications, cytoskeletal regulation,

and broader distribution of scaffolding proteins like Bruchpilot could likely create a cellular

environment permissive to structural reorganization and functional recovery after injury,

fundamentally distinguishing it from phasic neurons that are intrinsically optimized for high-

fidelity transmission rather than adaptive structural change.

Evolutionary Parallels in Injury-Induced Plasticity

Our findings reflect a conserved strategy for injury-induced plasticity across species. In the

mammalian spinal cord, both interneurons and descending corticospinal tract (CST) neurons can

sprout following injury, particularly when targets are fully or partially denervated (55). Similar to

our findings in Drosophila, this rewiring often originates from spared neurons, such as CST axons

that form new connections post-lesion (56, 57). Notably, Nakamura et al. showed that collateral

sprouting from spared CST axons—rather than long-distance regeneration of damaged neurons—

was sufficient to restore motor control (57), echoing our demonstration that MN1 can reinnervate

denervated targets to restore crawling.

Additional studies highlight that promoting plasticity, not just regeneration, may be key for

functional recovery (58, 59). These studies emphasize the role of the injury environment,

including target denervation and glial modulation, in supporting sprouting (60, 61). Vertebrate

work shows that denervated muscle and Schwann cells transiently emit growth-permissive cues

that support reinnervation (62). Brown and Holland (1979) demonstrated that partial denervation

of mouse soleus muscle triggers sprouting from intact motor axons, and that this response is

suppressed by direct electrical stimulation of denervated muscle—indicating that specific signals

from denervated targets, not simply inactivity, drive structural plasticity (63). Keynes et al.

(1983) further confirmed that structurally intact, but denervated, endplates are required for nodal

sprouting (64).

These parallels suggest that neuron-intrinsic identity, injury-induced cues, and the structural and

molecular context of the target together shape the capacity for adaptive circuit remodeling. Our

Drosophila model provides a genetically tractable system to dissect these mechanisms with

single-cell resolution and causal precision.

Conclusion and Perspective

Our study reveals that injury-induced heterosynaptic sprouting in the Drosophila motor circuit is

triggered by complete loss of presynaptic input, is neuron-subtype specific, and contributes to

functional recovery. These findings lay a foundation for uncovering the molecular logic of

plasticity and circuit repair.

Drosophila offers a powerful model for dissecting the cellular and genetic mechanisms

underlying this process. Future studies can use this system to identify the intrinsic growth

properties of specific neurons and the extrinsic cues that promote or restrict sprouting. These

insights may guide therapeutic strategies to enhance plasticity in motor disorders such as ALS

and spinal cord injury, and help test whether similar principles apply in vertebrate systems.

Materials and Methods

Fly Husbandry

427   All Crosses were reared at 25°C at 50% humidity with a 12 hr light/dark cycle. For crosses with

428   GTACR, larvae were fed with all-trans retinol (ATR) and kept in constant darkness until they

429   were ready to be imaged or dissected. Crosses involving RPR were at 29°C to ensure ablation

430   efficiency. For Larval ablation using Gal80ts, Embryos were kept at 25°C and as larvae hatched,

431   they transferred to 29°C to induce ablation. A list of stocks used can be found in Table S1.

Fillet Dissection and Immunohistochemistry

434   L3 larvae were used for fillet dissection, cutting on the ventral side. Dissections were fixed in 4%

435   PFA, blocked with equal parts 5% Goat serum and 5% Donkey serum in PBST, and stained with

436   primary and secondary antibodies. See SI Materials and methods and Table S2 for more details.

NMJ Imaging and Quantification

Airyscan Z-stacks were acquired using a Zeiss LSM900 Confocal with a 40x Objective and

processed with 3D Airyscan processing with the automatic filter. Postsynaptic Dlg of the neuron

of interest were isolated using a mask of the marked neuron and the area, bouton number and

bouton size were determined using a Fiji plugin developed by the Schneck lab. Branching points

were counted and separated based on colocalization with Dlg and the order of branching. See SI

Materials and Methods for more details.

Behavioral Assays

447   Larvae were placed in groves in a 2% agarose gel and recorded

448   (30 frames per second) using a smartphone mounted on a Zeiss Stemi 305 Microscope as they

449   crawled 1 cm. The number of peristalsis and frames for each peristalsis were counted to

450   determine the efficiency and duration, respectively. For analysis of overall crawling behavior,

451   Larvae were placed on a 2% agarose gel and recorded using a smartphone above the arena for 1

minute. The WrmTrck Plugin was utilized to determine the distance traveled and for each stage

the ablatio group was normalized by the average of the control. See SI Materials and Methods for

more details.

Intact Larvae Imaging and Muscle Calcium Imaging Quantification

Larvae were immobilized between a 2% agarose gel pad and a coverslip and imaged with 40x,

20x, or 10x, depending on the size of the larvae. Z-stacks of the marked axons at L0, L1, L2, and

L3 were used to count the number of branching points. Time series of larvae undergoing

peristalsis were analyzed using a custom MATLAB script to determine the intensity of the

GCamp signal during peristalsis. See SI Materials and Methods for more details.

Statistical Analysis

All statistical analysis was performed using R in RStudio. Averages were plotted with SEM error

bars and where applicable individual samples were added as points. Number of replicates and

animals is indicated in figure legends. Kruskal Wallis and Dunn post hoc were used for all NMJ

quantification and behavioral data, except the data in Figure 2 which used a t-test. To compare the

L1 and L3 behavioral deficits, a t-test was used. Significance was established with $* = p<0.05$, $**$

$= p<0.01$, $*** = p<0.001$, $****=p<0.0001$.

Acknowledgments

We are grateful to the Bloomington Drosophila Stock Center, Chris Doe, Troy Littleton, Dion Dickman, Keiko Hirono, Sarah Ackerman for generously providing transgenic fly lines. We also thank Lauren Carlisle and Elaina Hildner for their invaluable technical support. This research was funded by a grant from Texas A&M University.

References

1. N. Aderinto, M. O. AbdulBasit, G. Olatunji, T. Adejumo, Exploring the transformative influence of neuroplasticity on stroke rehabilitation: a narrative review of current evidence. *Ann Med Surg (Lond)* **85**, 4425-4432 (2023).
2. M. Bonanno, R. De Luca, A. M. De Nunzio, A. Quartarone, R. S. Calabro, Innovative Technologies in the Neurorehabilitation of Traumatic Brain Injury: A Systematic Review. *Brain Sci* **12** (2022).

3.   C. T. Werner, C. J. Williams, M. R. Fermelia, D. T. Lin, Y. Li, Circuit Mechanisms of
Neurodegenerative Diseases: A New Frontier With Miniature Fluorescence Microscopy.
Front Neurosci **13**, 1174 (2019).
4.   D. Tome, R. D. Almeida, Remaking a connection: molecular players involved in post-
injury synapse formation. *Neural Regen Res* **20**, 1719-1720 (2025).
5.   K. P. Menon, R. A. Carrillo, K. Zinn, Development and plasticity of the Drosophila larval
neuromuscular junction. *Wiley interdisciplinary reviews. Developmental biology* **2**, 647-
670 (2013).
6.   Z. L. Newman *et al.*, Input-Specific Plasticity and Homeostasis at the Drosophila Larval
Neuromuscular Junction. *Neuron* **93**, 1388-1404.e1310 (2017).
7.   N. A. Aponte-Santiago, K. G. Ormerod, Y. Akbergenova, J. T. Littleton, Synaptic Plasticity
Induced by Differential Manipulation of Tonic and Phasic Motoneurons in Drosophila.
The Journal of neuroscience : the official journal of the Society for Neuroscience **40**,
6270-6288 (2020).
8.   Y. Wang, M. Lobb-Rabe, J. Ashley, V. Anand, R. A. Carrillo, Structural and Functional
Synaptic Plasticity Induced by Convergent Synapse Loss in the Drosophila
Neuromuscular Circuit. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience **41**, 1401-1417 (2021).
9.   C. C. Yifu Han, Pragya Goel, Kaikai He, Cristian Pinales, Chris Buser, Dion Dickman,
Botulinum neurotoxin accurately separates tonic vs phasic transmission and reveals
heterosynaptic plasticity rules in Drosophila. *bioRxiv*
https://doi.org/10.1101/2022.02.15.480429 (2022).
10.   Y. Wang *et al.*, Glial Draper signaling triggers cross-neuron plasticity in bystander
neurons after neuronal cell death in Drosophila. *Nature communications* **14**, 4452
(2023).
11.   Z. Lu *et al.*, High-Probability Neurotransmitter Release Sites Represent an Energy-
Efficient Design. *Current biology : CB* **26**, 2562-2571 (2016).
12.   G. A. Lnenicka, H. Keshishian, Identified motor terminals in Drosophila larvae show
distinct differences in morphology and physiology. *Journal of neurobiology* **43**, 186-197
(2000).
13.   N. A. Aponte-Santiago, J. T. Littleton, Synaptic Properties and Plasticity Mechanisms of
Invertebrate Tonic and Phasic Neurons. *Front Physiol* **11**, 611982 (2020).
14.   R. P. Dum, T. T. Kennedy, Physiological and histochemical characteristics of motor units
in cat tibialis anterior and extensor digitorum longus muscles. *Journal of*
neurophysiology **43**, 1615-1630 (1980).
15.   S. W. Kuffler, E. M. Vaughan Williams, Properties of the 'slow' skeletal muscles fibres of
the frog. *The Journal of physiology* **121**, 318-340 (1953).
16.   H. Atwood, Parallel 'phasic' and 'tonic' motor systems of the crayfish abdomen. *The*
Journal of experimental biology **211**, 2193-2195 (2008).
17.   C. A. Frank *et al.*, New approaches for studying synaptic development, function, and
plasticity using Drosophila as a model system. *The Journal of neuroscience : the official*
journal of the Society for Neuroscience **33**, 17560-17568 (2013).
18.   G. W. Davis, M. Müller, Homeostatic control of presynaptic neurotransmitter release.
Annu Rev Physiol **77**, 251-270 (2015).
19.   K. P. Harris, J. T. Littleton, Transmission, Development, and Plasticity of Synapses.
Genetics **201**, 345-375 (2015).
20.   O. Genc, G. W. Davis, Target-wide Induction and Synapse Type-Specific Robustness of
Presynaptic Homeostasis. *Current biology : CB* **29**, 3863-3873 e3862 (2019).

21. P. Goel *et al.*, Homeostatic scaling of active zone scaffolds maintains global synaptic
strength. *Journal of Cell Biology* **218**, 1706-1724 (2019).
22. T. D. James, D. J. Zwiefelhofer, C. A. Frank, Maintenance of homeostatic plasticity at the
Drosophila neuromuscular synapse requires continuous IP(3)-directed signaling. *eLife* **8**
(2019).
23. C. A. Frank, T. D. James, M. Müller, Homeostatic control of Drosophila neuromuscular
junction function. *Synapse* **74**, e22133 (2020).
24. X. Li *et al.*, Autocrine inhibition by a glutamate-gated chloride channel mediates
presynaptic homeostatic depression. *Science Advances* **7**, eabj1215 (2021).
25. C. J. Yeates, C. A. Frank, Homeostatic Depression Shows Heightened Sensitivity to
Synaptic Calcium. *Front Cell Neurosci* **15**, 618393 (2021).
26. N. S. Armstrong, C. A. Frank, The calcineurin regulator Sarah enables distinct forms of
homeostatic plasticity at the Drosophila neuromuscular junction. *Front Synaptic*
Neurosci **14**, 1033743 (2022).
27. Y. Han *et al.*, Botulinum neurotoxin accurately separates tonic vs. phasic transmission
and reveals heterosynaptic plasticity rules in Drosophila. *eLife* **11**, e77924 (2022).
28. K. He *et al.*, Physiologic and Nanoscale Distinctions Define Glutamatergic Synapses in
Tonic vs Phasic Neurons. *The Journal of neuroscience : the official journal of the Society*
for Neuroscience **43**, 4598-4611 (2023).
29. T. Wang, C. A. Frank, Using Electrophysiology to Study Homeostatic Plasticity at the
Drosophila Neuromuscular Junction. *Cold Spring Harb Protoc* 10.1101/pdb.top108393
(2024).
30. M. Landgraf, T. Bossing, G. M. Technau, M. Bate, The origin, location, and projections of
the embryonic abdominal motorneurons of Drosophila. *The Journal of neuroscience : the*
official journal of the Society for Neuroscience **17**, 9642-9655 (1997).
31. B. Hoang, A. Chiba, Single-cell analysis of Drosophila larval neuromuscular synapses.
Developmental biology **229**, 55-70 (2001).
32. M. Landgraf, V. Jeffrey, M. Fujioka, J. B. Jaynes, M. Bate, Embryonic origins of a motor
system: motor dendrites form a myotopic map in Drosophila. *PLoS biology* **1**, E41 (2003).
33. M. Landgraf, S. Thor, Development of Drosophila motoneurons: specification and
morphology. *Seminars in cell & developmental biology* **17**, 3-11 (2006).
34. M. F. Zwart *et al.*, Selective Inhibition Mediates the Sequential Recruitment of Motor
Pools. *Neuron* **91**, 615-628 (2016).
35. A. A. Zarin, J. P. Labrador, Motor axon guidance in Drosophila. *Seminars in cell &*
developmental biology 10.1016/j.semcdb.2017.11.013 (2017).
36. A. A. Zarin, B. Mark, A. Cardona, A. Litwin-Kumar, C. Q. Doe, A multilayer circuit
architecture for the generation of distinct locomotor behaviors in Drosophila. *eLife* **8**
(2019).
37. P. C. Cooney *et al.*, Neuromuscular basis of Drosophila larval rolling escape behavior.
Proceedings of the National Academy of Sciences of the United States of America **120**,
e2303641120 (2023).
38. S. K. Jetti *et al.*, Molecular logic of synaptic diversity between Drosophila tonic and
phasic motoneurons. *Neuron* **111**, 3554-3569 e3557 (2023).
39. L. Goyal, K. McCall, J. Agapite, E. Hartwieg, H. Steller, Induction of apoptosis by
Drosophila reaper, hid and grim through inhibition of IAP function. *The EMBO journal*
19, 589-597 (2000).
40. J. J. Perez-Moreno, C. J. O'Kane, GAL4 Drivers Specific for Type Ib and Type Is Motor
Neurons in Drosophila. *G3 (Bethesda, Md.)* **9**, 453-462 (2019).

41.   B. Nijhof *et al.*, A New Fiji-Based Algorithm That Systematically Quantifies Nine Synaptic
585   Parameters Provides Insights into Drosophila NMJ Morphometry. *PLoS computational*
586   *biology* **12**, e1004823 (2016).

587   42.   A. Castells-Nobau *et al.*, Two Algorithms for High-throughput and Multi-parametric
588   Quantification of Drosophila Neuromuscular Junction Morphology. *J Vis Exp*
589   10.3791/55395 (2017).

590   43.   S. E. McGuire, P. T. Le, A. J. Osborn, K. Matsumoto, R. L. Davis, Spatiotemporal rescue of
591   memory dysfunction in Drosophila. *Science (New York, N.Y.)* **302**, 1765-1768 (2003).

592   44.   M. L. Suster, L. Seugnet, M. Bate, M. B. Sokolowski, Refining GAL4-driven transgene
593   expression in Drosophila with a GAL80 enhancer-trap. *Genesis* **39**, 240-245 (2004).

594   45.   A. Carreira-Rosario *et al.*, MDN brain descending neurons coordinately activate
595   backward and inhibit forward locomotion. *eLife* **7** (2018).

596   46.   A. S. Mauss, C. Busch, A. Borst, Optogenetic Neuronal Silencing in Drosophila during
597   Visual Processing. *Scientific reports* **7**, 13823 (2017).

598   47.   F. Mohammad *et al.*, Optogenetic inhibition of behavior with anion channelrhodopsins.
599   *Nature methods* **14**, 271-274 (2017).

600   48.   A. Sitaula, Y. Huang, A. Zarin, Application of a Dual Optogenetic Silencing-Activation
601   Protocol to Map Motor Neurons Driving Rolling Escape Behavior in Drosophila Larvae.
602   *Bio Protoc* **14**, e5131 (2024).

603   49.   A. A. Z. Yuhan Huang, A premotor microcircuit to generate behavior-specific muscle
604   activation patterns in Drosophila larvae. *bioRxiv*
605   https://doi.org/10.1101/2022.08.18.504452 (2022).

606   50.   S. T. Sweeney, K. Broadie, J. Keane, H. Niemann, C. J. O'Kane, Targeted expression of
607   tetanus toxin light chain in Drosophila specifically eliminates synaptic transmission and
608   causes behavioral defects. *Neuron* **14**, 341-351 (1995).

609   51.   T. N. Chang, H. Keshishian, Laser ablation of Drosophila embryonic motoneurons causes
610   ectopic innervation of target muscle fibers. *The Journal of neuroscience : the official*
611   *journal of the Society for Neuroscience* **16**, 5715-5726 (1996).

612   52.   S. D. Ackerman, N. A. Perez-Catalan, M. R. Freeman, C. Q. Doe, Astrocytes close a motor
613   circuit critical period. *Nature* **592**, 414-420 (2021).

614   53.   T. H. Nguyen *et al.*, scRNA-seq data from the larval Drosophila ventral cord provides a
615   resource for studying motor systems function and development. *Dev Cell* **59**, 1210-1230
616   e1219 (2024).

617   54.   A. T. Medeiros, S. J. Gratz, A. Delgado, J. T. Ritt, K. M. O'Connor-Giles, Ca(2+) channel
618   and active zone protein abundance intersects with input-specific synapse organization
619   to shape functional synaptic diversity. *eLife* **12** (2024).

620   55.   T. Hagg, Collateral sprouting as a target for improved function after spinal cord injury. *J*
621   *Neurotrauma* **23**, 281-294 (2006).

622   56.   N. Weidner, A. Ner, N. Salimi, M. H. Tuszynski, Spontaneous corticospinal axonal
623   plasticity and functional recovery after adult central nervous system injury. *Proceedings*
624   *of the National Academy of Sciences of the United States of America* **98**, 3513-3518
625   (2001).

626   57.   Y. Nakamura *et al.*, Modulation of Both Intrinsic and Extrinsic Factors Additively
627   Promotes Rewiring of Corticospinal Circuits after Spinal Cord Injury. *The Journal of*
628   *neuroscience : the official journal of the Society for Neuroscience* **41**, 10247-10260
629   (2021).

630   58.   S. F. Kazim *et al.*, Corticospinal Motor Circuit Plasticity After Spinal Cord Injury:
631   Harnessing Neuroplasticity to Improve Functional Outcomes. *Mol Neurobiol* **58**, 5494-
632   5516 (2021).
633   59.   M. Chen, B. Zheng, Axon plasticity in the mammalian central nervous system after
634   injury. *Trends Neurosci* **37**, 583-593 (2014).
635   60.   J. P. Kapfhammer, Axon sprouting in the spinal cord: growth promoting and growth
636   inhibitory mechanisms. *Anat Embryol (Berl)* **196**, 417-426 (1997).
637   61.   S. Bolivar, X. Navarro, E. Udina, Schwann Cell Role in Selectivity of Nerve Regeneration.
638   *Cells* **9** (2020).
639   62.   I. C. Maier, M. E. Schwab, Sprouting, regeneration and circuit formation in the injured
640   spinal cord: factors and activity. *Philosophical transactions of the Royal Society of*
641   *London. Series B, Biological sciences* **361**, 1611-1634 (2006).
642   63.   M. C. Brown, R. L. Holland, A central role for denervated tissues in causing nerve
643   sprouting. *Nature* **282**, 724-726 (1979).
644   64.   R. J. Keynes, W. G. Hopkins, M. C. Brown, Sprouting of mammalian motor neurones at
645   nodes of Ranvier: the role of the denervated motor endplate. *Brain research* **264**, 209-
646   213 (1983).

Figure 1. The tonic MN1 axon sprouts and forms ectopic NMJs on fully denervated

dorsal-longitudinal (DL) muscles.

(A) Schematic diagrams of the Drosophila larval nervous system and body wall muscles.

This study focuses on four DL muscles (1, 2, 9, and 10), each receiving phasic input from

a common phasic MN-Is and tonic input from distinct tonic Ib MNs (MN1, MN2, MN9,

and MN10).

(B) Target muscles of the Gal4 and LexA MN drivers used in this study. The union

symbol (∪) indicates fly lines carrying two drivers simultaneously. For example, in *Is ∪*

CQ larvae, both tonic and phasic MNs innervating muscles 2, 3, 4, 9, and 10 are targeted.

(C) Workflow illustrating the NMJ analysis pipeline [adapted from (41, 42)].

(D) Representative NMJ immunostaining visualizing the bystander MN1 axon in

different genotypes: control, phasic MN-Is ablated (*Is>RPR*), tonic MN2, MN9, MN10

ablated (*CQ>RPR*), and both tonic and phasic MNs ablated (*Is ∪ CQ>RPR*). Schematics

(left) depict ablated MNs and their effects on MN1. In controls, MN1 exclusively

innervates muscle 1 (top row), and this pattern is unchanged after phasic MN-Is ablation

(second row). Ablation of tonic MNs (CQ>RPR) induces sprouting of MN1 onto muscle

2 (M2). Co-ablation of tonic and phasic MNs (*Is ∪ CQ>RPR*) induces robust MN1

sprouting, forming ectopic NMJs on all nearby denervated muscles (M2, M9, M10) in

addition to its native target (M1). Postsynaptic sites are marked by Dlg (magenta) and

muscles by phalloidin (blue). MN1 is labeled with UAS-myr-GFP driven by 94G06-

Gal4; targeted MN ablation is achieved using the LexA/OP-RPR system.

(E) Quantification of MN1 axon and NMJ parameters across groups shown in (D).

(i) Total MN1 area on DL muscles 1, 2, 9, and 10. *Is ∪ CQ>RPR* larvae exhibit

significantly increased total MN1 surface area relative to control and other ablation

groups.

(ii) Total number of MN1 boutons on DL muscles. No significant differences across

groups.

(iii) Average bouton size (area/bouton number). No significant differences across groups.

(iv) Proportional distribution of MN1 NMJ area on individual DL muscles. In CQ>RPR

larvae, MN1 ectopically innervates muscle 2; in *Is ∪ CQ>RPR* larvae, MN1 innervates

muscles 2, 9, and 10 in addition to its native target Muscle 1. The area on Muscle 1 is

significantly decreased compared to control group.

(v) Average number of motor axon branching points (green) and NMJ branching points.

CQ>RPR and *Is ∪ CQ>RPR* groups exhibit significantly increased axon branching

compared to control group.

For control, *Is>RPR*, and *Is ∪ CQ>RPR*, n = 8 larvae (16 hemisegments). For *CQ>RPR*,

n = 7 larvae (13 hemisegments). Data points represent single hemisegments. Error bars:

±SEM. Statistical analysis: Kruskal-Wallis with Dunn's post-hoc test (*p<0.05;

**p<0.01; ***p<0.001; ****p<0.0001*).

A

Muscles (Phalloidin) MN-Is>GFP Dlg

Control (no driver)

MN1 — Muscle 1
MN9 — Muscle 9
MN2 — Muscle 2
MN10 — Muscle 10
MN Is

25 μm

MN1^{U}CQ >RPR

MN1 — Muscle 1
MN9 — Muscle 9
MN2 — Muscle 2
MN10 — Muscle 10
MN Is

25 μm

B

MN-Is area on DL muscles

Total Area (μm²)

Gal4: none   MN1^{U}CQ

UAS-RPR

MN-Is Bouton Size

Area/Bouton (μm2)

Gal4: none   MN1^{U}CQ

UAS-RPR

MN-Is boutons on DL muscles

Bouton Number

Gal4: none   MN1^{U}CQ

UAS-RPR

MN-Is Percent of Area

% Total Area

Gal4: none   MN1^{U}CQ

UAS-RPR

1
9
2
10

C

MN-Is branching on four DL muscles 1,2,9,10

Position
Axon 1st   Axon Total
Axon 2nd   NMJ
Axon 3rd   Axon+NMJ

Branch Point Number

*
**
***

Gal4: none   MN1^{U}CQ

UAS-RPR

Axon branch classification

2nd
1st
3rd

Figure 2. Phasic MN-Is does not exhibit structural expansion following ablation of tonic MNs innervating dorsal-longitudinal (DL) muscles.

(A) Representative NMJ immunostaining showing the bystander phasic MN-Is axon in control larvae and in larvae where all tonic Ib MNs innervating DL muscles (MN1, MN2, MN9, MN10) were ablated (*MN1 ∪ CQ>RPR*). Postsynaptic sites are labeled with Dlg (magenta), muscles with phalloidin (blue), and MN-Is axons with myr-GFP driven by the 27E09 enhancer. Ablation of tonic MNs was achieved using the 94G06-Gal4 ∪ CQ-Gal4 / UAS-RPR system.

(B) Quantification of Is NMJ parameters shows no significant differences in total NMJ area, bouton size, bouton number, or NMJ area distribution between control and *MN1 ∪ CQ>RPR* larvae. Each data point represents a single hemisegment.

(C) Ablation of tonic MNs results in a significant reduction in 2nd- and 3rd-order axonal branches of the bystander MN-Is.

For both control and *MN1 ∪ CQ>RPR* groups, n = 6 larvae (12 hemisegments). Data points represent individual hemisegments. Error bars indicate mean ± SEM. Statistical analysis: Kruskal-Wallis with Dunn's post-hoc test ($p<0.05$; $*p<0.01$; $**p<0.001$; $***p<0.0001$).

Figure 3. The bystander MN1 exhibits distinct structural responses to different perturbations of neighboring MNs.

(A–C) Representative images (A) and quantitative analyses (B–C) of MN1 axons and NMJs across genotypes. MN1 is labeled with MN1-lexA >AOP-myr-GFP.

In control animals (CQ ∪ Is-Gal4/UAS-RPR, Gal80^{ts} raised at 24°C), MN1 shows normal axon and NMJ morphology. In the ablation group (CQ ∪ Is-Gal4/UAS-RPR, Gal80^{ts} raised at 29°C), MN1 exhibits robust sprouting, forming ectopic NMJs on denervated muscles 2, 9, and 10.

In the acute silencing group (CQ ∪ Is-Gal4/UAS-GtACR1), MN1 NMJ area and bouton number are elevated relative to controls, but no ectopic sprouting onto muscles 2, 9, and 10 is observed.

In the chronic silencing group targeting both evoked and spontaneous release (CQ ∪ Is-Gal4/UAS-BoNT-C), MN1 shows increased bouton numbers without ectopic sprouting.

In the chronic silencing group targeting only evoked release (CQ ∪ Is-Gal4/UAS-TeTxLC), MN1 NMJ branching, area, and bouton number are markedly increased compared to controls and other perturbations. However, bouton size is significantly reduced, and no ectopic sprouting onto muscles 2, 9, and 10 occurs. Notably, the overall NMJ morphology of TeTxLC-silenced tonic and phasic MNs (white arrows) appears normal based on Dlg staining.

Data points represent single hemisegments. Ablation group: n=7 larvae (14 hemisegments). All other groups: n=6 larvae (12 hemisegments). Error bars: ±SEM. Kruskal-Wallis with Dunn's post-hoc test (*p<0.05; **p<0.01; ***p<0.001; ****p<0.0001).

A

MN1^{u}CQ^{u}Is (Control) | MN1^{u}CQ^{u}Is > RPR | MN1^{u}CQ^{u}Is > GtACR1 | CQ^{u}Is > GtACR1 | CQ^{u}Is > RPR | CQ^{u}Is > RPR; MN1 > GtACR1

Time

B

Muscle GCaMP6f (DF/F)

Muscle: 1, 9, 2, 10

MN1^{u}CQ^{u}Is (Control) | MN1^{u}CQ^{u}Is > RPR | MN1^{u}CQ^{u}Is > GtACR1 | CQ^{u}Is > GtACR1 | CQ^{u}Is > RPR | CQ^{u}Is > RPR; MN1 > GtACR1

C

Phalloidin   HRP   Dlg

MN1^{u}CQ^{u}Is > RPR

D

Peristalsis = 0 | Peristalsis = 5 | Peristalsis = 10

MN1^{u}CQ^{u}Is (Control)
MN1^{u}CQ^{u}Is > RPR
MN1^{u}CQ^{u}Is > GtACR1
CQ^{u}Is > GtACR1
CQ^{u}Is > RPR
CQ^{u}Is > RPR; MN1 > GtACR1

E

Peristalsis Efficiency

MN1^{u}CQ^{u}Is (Control) | MN1^{u}CQ^{u}Is > RPR | MN1^{u}CQ^{u}Is > GtACR1 | CQ^{u}Is > GtACR1 | CQ^{u}Is > RPR | CQ^{u}Is > RPR; MN1 > GtACR1

Peristalsis Duration

MN1^{u}CQ^{u}Is (Control) | MN1^{u}CQ^{u}Is > RPR | MN1^{u}CQ^{u}Is > GtACR1 | CQ^{u}Is > GtACR1 | CQ^{u}Is > RPR | CQ^{u}Is > RPR; MN1 > GtACR1

Figure 4. Ectopic NMJs formed by sprouting MN1 are functional and improve larval locomotion.

(A) Muscle calcium imaging in intact, forward-crawling larvae of different genotypes. Each column shows still images from three adjacent abdominal segments at successive time points during a forward peristaltic wave. GCaMP6f fluorescence (expressed under the 44H10 muscle-specific enhancer) increases during muscle contraction. Cyan rectangles mark dorsal-longitudinal (DL) muscles in the top panels. In control larvae (MN1 ∪ CQ ∪ Is-lexA/ Aop-GTACR ATR (-)), DL muscles contract sequentially, with GCaMP6f signals propagating from posterior (right) to anterior (left) during crawling. Ablation of all tonic and phasic MNs (MN1 ∪ CQ ∪ Is-lexA/Aop-RPR) abolishes DL muscle activity, leaving ventral and lateral muscles unaffected. Similarly, acute silencing of these MNs (MN1 ∪ CQ ∪ Is-lexA/Aop-GtACR1 ATR (+)) eliminates DL muscle activity. In CQ ∪ Is-lexA/Aop-GtACR1 ATR (+) larvae, only Muscle 1 remains active, consistent with preserved MN1 input. In contrast, CQ ∪ Is-lexA/Aop-RPR larvae exhibit robust calcium activity and contractions in all four DL muscles, indicating functional reinnervation by sprouted MN1. Acute silencing of the sprouted MN1 in CQ ∪ Is-lexA/Aop-RPR; MN1-Gal4>UAS-GtACR1 ATR (+) larvae abolishes DL muscle activity. Frames are taken from Supplementary Video X.

(B) Quantification of GCaMP6f signal in DL muscles (1, 2, 9, and 10) confirms observations in (A). n = 5 larvae per genotype (10 hemisegments total). Statistical analysis: Kruskal–Wallis test with Dunn's post-hoc correction (*$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$).

(C) Immunostaining of DL muscles in MN1 ∪ CQ ∪ Is-lexA/Aop-RPR larvae verifies complete ablation of targeted MNs. Anti-HRP (green) labels the nearby sensory neurons, and no motor axons or Dlg puncta are detected in denervated DL muscles.

(D) Representative crawling trajectories of larvae from the genotypes in (A). Each row shows larval positions at the start line (Peristalsis = 0), after 5 peristalses, and after 10 peristalses. Among all groups, control larvae (MN1 ∪ CQ ∪ Is-lexA/+) ranked first in distance traveled, followed by CQ ∪ Is-lexA/Aop-RPR larvae, which ranked second. In

contrast, MN1 ∪ CQ ∪ Is-lexA/Aop-GtACR1 larvae showed the least forward

progression.

(E) Quantification of peristalsis efficiency (distance traveled per peristaltic wave) and

peristalsis duration (time required to complete a wave). Control larvae (MN1 ∪ CQ ∪ Is-

lexA/+) exhibited the highest peristalsis efficiency and shortest duration, followed by CQ

∪ Is-lexA/Aop-RPR larvae. Silencing the sprouted MN1 in CQ ∪ Is-lexA/Aop-RPR;

MN1-Gal4>UAS-GtACR1 larvae reduced locomotor performance to levels comparable

with other two silenced groups. n = 40 larvae per group, with each dot representing an

individual animal. Statistical analysis: One-way ANOVA with Tukey post-hoc test

(*$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$).

Figure 5. MN1 axon sprouting dynamics and timing of injury-induced plasticity

(A) Live imaging of MN1 axons across larval stages. Each row shows stills from the same hemisegment at L1, L2, and L3.

(i) In a control larva (CQ ∪ Is-Gal4/+; MN1-lexA >Aop-tdTomato), MN1 innervates its native target muscle 1 (cyan box) by L1. The axon terminal arborizes further between L1–L3, with the main shaft bifurcating beneath muscle 1 to form a T-shaped NMJ.

(ii–vi) In ablated larvae (CQ ∪ Is-Gal4/UAS-RPR; MN1-lexA >Aop-tdTomato), diverse sprouting patterns are observed:

(ii) New axon branches (white arrow) emerge at L2 and grow further by L3.

(iii–v) Small axon buds appear at L1 (white arrows) and expand progressively in L2 and L3. Some branches (white asterisks) are already present at L1 and undergo marked growth during later stages.

(vi) In rare cases, MN1 fails to reach its native target muscle 1 and instead branches prematurely to innervate other DL muscles.

(B) Quantification of 1st- and 2nd-order axon branches in L1 and L3 stages for controls and ablated animals. At L1, ablated larvae show ~2× more 1st-order branches than controls, with comparable 2nd-order branching. By L3, ablated larvae exhibit significantly more 1st-order branches than controls. Within ablated larvae, both branch types increase markedly from L1 to L3; in controls, only 2nd-order branching increases over time. Data points represent single hemisegments (L1 control: n=5 larvae, 57 hemisegments; L1 ablation: n=5 larvae, 56 hemisegments; L3 control: n=5 larvae, 57 hemisegments; L3 ablation: n=5 larvae, 54 hemisegments). Error bars: ±SEM. Kruskal-Wallis with Dunn's post-hoc test (*$p<0.05$; **$p<0.01$; ***$p<0.001$; ****$p<0.0001$).

(C) Timing of ablation influences sprouting. Post-embryonic (larval) ablation of MNs (CQ ∪ Is-Gal4/UAS-RPR,Gal80^{ts} shifted from 24°C to 29°C at L1) still triggers MN1 sprouting but less robustly than embryonic ablation (CQ ∪ Is-Gal4/UAS-RPR,Gal80^{ts} raised at 29°C throughout). No sprouting occurs in control larvae maintained at 24°C.

(D) Quantification of MN1 bouton size, bouton number, axon area, and branch

complexity across groups in (C). Data points represent single hemisegments. Control

group: n = 6 larvae (12 hemisegments). For embryonic ablation: n = 7 larvae (14

hemisegments). Embryonic and control datasets are the same as in **Fig. 3**. For larval

ablation, n = 7 larvae (13 hemisegments). Error bars: ±SEM. Kruskal-Wallis with Dunn's

post-hoc test (*$p<0.05$; **$p<0.01$; ***$p<0.001$; ****$p<0.0001$).

A Control   **24°C** Is^UCQ Gal4>RPR Gal80ts

MN1 Dlg

Morphometric Area Outlining

Bouton Number Quantification

10µm

B Ablation   **29°C** Is^UCQ Gal4>RPR Gal80ts

MN1 Dlg

10µm

D Chronic silencing   Is^UCQ Gal4>BontC

MN1 Dlg

10µm

C Acute silencing   Is^UCQ Gal4>GtACR1

MN1 Dlg

10µm

E Chronic silencing   Is^UCQ Gal4>TeTxLC

MN1 Dlg TNT

MN1 Dlg

10µm

10µm

Fig. S1. Workflow illustrating the NMJ analysis pipeline [adapted from (41, 42)] in different larval groups. Morphological responses of MN1 to distinct manipulations of neighboring motor neurons.

Representative images illustrate the structural remodeling of MN1 following ablation or silencing of adjacent tonic and phasic motor neurons (MNs), as visualized with membrane-targeted GFP and Dlg immunostaining.

(A) In temperature-controlled **unmanipulated controls** (CQ ∪ Is-Gal4 > UAS-RPR, Gal80ts; raised entirely at 24°C), MN1 displays its typical morphology, forming a compact NMJ exclusively on its native target, Muscle 1. No ectopic projections or abnormal branching are detected.

(B) In the **ablation group** (CQ ∪ Is-Gal4 > UAS-RPR, Gal80ts; raised at 29°C), selective apoptosis of neighboring MNs induces a dramatic response in MN1. The axon extends beyond its native territory and establishes new, ectopic NMJs on denervated Muscles 2, 9, and 10, demonstrating robust sprouting and target reinnervation.

(C) In the **acute silencing condition** (CQ ∪ Is-Gal4 > UAS-GtACR1; light-exposed), MN1 remains restricted to Muscle 1. Although NMJ area and bouton number are elevated relative to controls, MN1 does not form branches onto adjacent muscles.

(D) In the **chronic silencing group targeting both evoked and spontaneous release** (CQ ∪ Is-Gal4 > UAS-BoNT-C), MN1 similarly exhibits increased bouton numbers but no ectopic sprouting. The NMJ remains confined to its native muscle target, and axon branching remains limited.

(E) In the **chronic silencing group that blocks only evoked transmission** (CQ∪Is-Gal4 > UAS-TeTxLC), MN1 shows the most extensive remodeling without sprouting. NMJ area, bouton number, and terminal branching complexity are all significantly increased compared to controls and other silencing groups. However, bouton size is notably reduced. Importantly, the tonic and phasic MNs targeted by TeTxLC appear morphologically normal (white arrows), suggesting that structural changes in MN1 are not due to degeneration of neighboring neurons but rather reflect a plastic response to altered circuit activity.

Scale bars and genotype-specific labeling are provided in each panel. This figure complements the quantitative analysis shown in **main Figure 3**.