Title

Pangenome of U.S. ex-PVP and Wild Sorghum Reveals Structural Variants and Selective Sweeps Shaping Adaptation and Trait Improvement

Running Title

Pangenome of U.S. ex-PVP and Wild Sorghum

Authors

Justine K. Kitony [1], Emily R. Murray [1], Kelly Colt [1], Ryan C. Lynch [1], Nicholas Allsing [1], Nolan T. Hartwick [1], Tiffany Duong [1], Jocelyn Saxton [2], Nadia Shakoor [2], Todd P. Michael [1,3]

Affiliation
1. Plant Molecular and Cellular Biology Laboratory, Salk Institute for Biological Studies, La Jolla, CA, USA
2. Donald Danforth Plant Science Center, Olivette, MO, USA
3. Science and Conservation, San Diego Botanical Garden, Encinitas, CA, USA

Corresponding Author

Todd P. Michael, toddpmichael@gmail.com, tmichael@salk.edu

Abstract

Sorghum is a cereal crop grown for food, feed, and biofuel, yet the genomic diversity of elite commercial lines remains underexplored. Here, we assembled a sorghum pangenome from long-read sequences of 46 U.S. ex-Plant Variety Protection Act (ex-PVP) cultivars and a haplotype-resolved assembly of one wild accession. The pangenome revealed largely conserved genomic architecture across elite lines but identified presence–absence variations (PAVs) in 28% of gene families, including genes involved in stress responses and starch metabolism, such as *STARCH BRANCHING ENZYME I (SBE1)*. Selective sweep analyses detected strong signals at key flowering-time loci (*Ma1/SbPRR37, Ma2, Ma6*) and circadian/light regulators (*PHOT1, DET1, XAP5*), with *PHYA* showing copy number variations (CNVs). Analysis of 24 wild accessions, beyond the core pangenome panel, detected CNVs in *MULTIDRUG AND TOXIC COMPOUND EXTRUSION (MATE)* transporters, potentially reflecting selection or breeding for enhanced herbicide detoxification. These results highlight breeding-shaped regions and position the pangenome as a resource for sorghum improvement.

Main
Sorghum (*Sorghum bicolor*) is a C_4 monocot renowned for its exceptional resilience to drought,
heat, and low-fertility soils [1]. As the fifth most important cereal crop globally, it is cultivated for
diverse purposes—including food, animal feed, alcoholic beverages, and biofuel—driven by its
nutritional value (high protein, essential micronutrients, gluten-free) and adaptability to abiotic
stresses. These attributes make sorghum a vital crop for enhancing food security and promoting
sustainable agriculture in the face of climate change [1–5].

Sorghum, originating in Africa, was introduced to the U.S. in the late 19th century through
germplasm collections of sweet, grain, and sudangrass types [6–9]. Early public breeding
programs played a key role in improvement, with notable successes such as the development of
the drought-tolerant 'Atlas' variety in 1936 [8]. The discovery of cytoplasmic male sterility (CMS)
and corresponding fertility restorer (Rf) genes in the mid-20th century enabled commercial hybrid
production, boosting yields by up to 40% by the 1950s [7,10].

From the 1960s to 1980s, U.S. sorghum breeding shifted toward the private sector. Companies
like Pioneer Hi-Bred began developing proprietary hybrids protected for 20 years under the
1970 Plant Variety Protection Act (PVPA). These hybrids featured dwarf stature (suitable for
mechanical harvesting), early maturity, stay-green traits (e.g., SC35), and resistance to
anthracnose, greenbug, downy mildew, and smut [7,11]. As a result, the United States emerged as
the world's leading exporter of sorghum, accounting for approximately 15% of global production
[12–14].

Most genomic studies in sorghum have relied on the 'BTx623' reference genome [1,15], which
represents a single grain-type line and fails to capture the full extent of genetic variation within
cultivated germplasm. This reference bias limits the discovery of presence–absence variations
(PAVs), SVs, and CNVs that underlie important agronomic traits [16–18]. Pangenome approaches
provide a broader framework for characterizing SVs across diverse accessions [19,20]. Short-
read-based pangenomes in sorghum have identified 35,719 genes, with ~53% showing PAVs
and 79 loci absent from the reference genome [21,22]. Long-read-based pangenomes have
uncovered even greater diversity—44,079 gene families across 16 accessions, including 222.6
Mb of novel sequence [20]. Similarly, high-quality assemblies from ten bioenergy sorghum
genotypes uncovered thousands of SVs, including those associated with traits like iron transport
[23]. Accessibility of long-read sequencing (e.g., PacBio HiFi, ONT) is now enabling detection of
novel gene content and SVs associated with traits [24,25].
80
From early domestication to modern crop improvement, selection—both natural and artificial—
has left footprints across plant genomes. These genomic signatures reveal how crops have
been shaped for key agronomic traits across time and environments. In wheat, genomic scans
have uncovered loci controlling flowering time [26]; in rice, genes conferring pathogen resistance
[27]; and in maize and cotton, variants linked to yield and stress tolerance [28,29]. Sorghum is no
exception. Selective sweeps have been reported at loci such as *SHATTERING1 (SH1)*, which
affects seed dispersal, and the *DRY NAC* transcription factor, involved in sugar accumulation
[30–32]. Beyond selection on classical domestication and yield-related genes, natural and artificial
selection has also acted on sorghum's internal time-of-day (TOD) regulatory network (circadian
rhythms), which governs key developmental processes such as vegetative growth and
reproductive timing [33–35]. These rhythms also influence the optimal timing of agricultural inputs

and support sorghum's adaptation across broad latitudinal ranges. Selection at the *Ma1* locus,
encoding *PSEUDO-RESPONSE REGULATOR 37 (SbPRR37)*, exemplifies adaptation to
temperate climates and remains a target in modern breeding for bioenergy [36]. Yet many regions
under selection remain unresolved due to the underrepresentation of elite commercial
germplasm genomes and the complexity of sorghum's breeding history [37–39].
97
98 Here we present a long-read-based sorghum pangenome constructed from 46 U.S. ex-PVP
99 lines and wild accessions, revealing approximately 28% PAVs in gene families, localized SVs,
100 selection signatures in TOD loci, and CNVs associated with key agronomic traits. This resource
101 enhances our understanding of sorghum adaptation and provides a foundation for genomic
102 characterization and crop improvement through modern breeding approaches [38,40,41].
103
Results
High-Quality Genome Assemblies of Ex-PVP and Wild Sorghum Accessions
We assembled high-quality genomes for 46 elite ex-PVP lines and both haplotypes of a wild
sorghum accession, 'PI156549.' Genome sizes ranged from 610 to 779 megabases (Mb), with
confirmed gene models ranging from 30,000 to 33,000 per genome (Fig. 1a). The new
assemblies achieved levels of contiguity comparable to the 'BTx623' reference, with N50 values
reaching up to 19 Mb (Fig. 1b, Supplementary Fig. 1a-c, Supplementary Table 1), reflecting
reference-quality assemblies across the dataset.
Assembly quality was assessed using the LTR Assembly Index (LAI), with scores ranging from
20 to 30, consistent with reference-grade assemblies (Fig. 1c, Supplementary Table 1). BUSCO
analysis confirmed high completeness of both genome assemblies (92.60–99.10% complete
BUSCOs) and predicted proteins (91.70–98.50%), supporting an accurate gene space view
across diverse genotypes (Supplementary Table 2).
Total interspersed repeat content showed a strong correlation with genome size (R^2 = 0.834;
Fig. 1d), underscoring the dominant role of repeats in genome expansion. Centromeric tandem
repeat expansions also correlated with genome size (R^2 = 0.747; Supplementary Fig. 1d).
Transposable elements (TEs), which make up ~65% of each genome assembly (Fig. 1e),
dominate sorghum genome composition and have broadly shaped its structure, regulation, and
adaptability [42]. Among these, retroelements (LTRs) were the most abundant class, followed by
DNA transposons and rolling-circle elements. Notably, a substantial proportion (~15–20%) of
the TEs content remained unclassified across genomes, even when using high-resolution
annotations from Helixer [43]. This unclassified fraction highlights the presence of highly divergent
or lineage-specific elements that escape conventional TEs classification approaches [44,45].
Analysis based on K-mer profiles revealed genetic clustering among the ex-PVP lines, with
some accessions showing closer relatedness but overall forming a cohesive group. The wild
accession formed a distinct outgroup (Fig. 1f), consistent with its contribution to novel K-mers in
the PanKmer collector's curve, which shows an open pangenome (Supplementary Fig. 1e,
Supplementary Table 3).
132

Fig. 1: Genome Assembly Metrics, Transposable Elements (TEs), and Genome
Architecture of the U.S. Ex-PVP and Wild Sorghum Pangenome.
a, Genome length (Mb; right y-axis) and gene count (left y-axis) across assembled accessions.
b, Assembly contiguity measured by N50 values (Mb). **c**, Example LTR Assembly Index (LAI)
distribution across chromosomes for the wild accession 'PI156549' Haplotype 1, with a global
mean of 23.88 and a standard deviation (SD) of 5.37. **d**, Correlation between total interspersed
repeat content and genome size. **e**, TEs classes expressed as a percentage of genome size.
Colors represent TEs categories as shown in the legend. **f**, Clustered heatmap of adjacency
values based on pairwise K-mer comparisons between genomes. The ex-PVP lines exhibit
largely conserved nucleotide identity, forming a homogeneous cluster distinct from the wild
accession haplotypes (red box). Pangenome accessions include elite ex-PVP lines and wild
sorghum samples. Full sample and sequencing details are in Supplementary Table 1; source
data are available in the associated data file.

Sorghum Pangenome Captures Gene Diversity Missed by the BTx623 Reference
Using the constructed pangenome, we characterized gene content variation that would be
overlooked when relying solely on the 'BTx623' reference. In total, 36,004 gene families
(orthogroups) were identified (Supplementary Table 4), comprising 71.5% core genes, 2.2%
soft-core genes, 16.7% dispensable genes, and 9.6% private genes (Fig. 2a). Gene
accumulation curves showed continued pangenome growth and core genome decline,
approaching a plateau that suggests most gene diversity was captured (Fig. 2b). A power law fit
of novel gene family discovery (a = 4.13, R² = 0.999) confirmed a highly closed sorghum
pangenome (Fig.2c) [46]. Comparison to the 'BTx623' reference revealed 6,389 additional gene
families, many absent from the reference but present in wild and elite ex-PVP lines (Fig. 2d).
Gene PAVs analysis across the 50 genomes revealed shared gene family patterns among the
ex-PVPs (Fig. 2e), with genespace visualization further highlighting overall gene conservation
across chromosomal segments (Fig. 2f).

The 6,389 pangenome exclusive orthogroups (Fig. 2d) include key KEGG functions absent from
the reference 'BTx623' (Fig. 2g), such as chloroplast electron transport components—ATP
SYNTHASE SUBUNIT BETA (ATPB), NADH DEHYDROGENASE SUBUNIT H (NDHH), and
NADH DEHYDROGENASE SUBUNIT F (NDHF); stress response regulators—GLUTATHIONE
S-TRANSFERASE TAU 1 (GSTU1) and ABSCISIC ALDEHYDE OXIDASE 3 (AAO3); and
specialized metabolism genes—FLAVONOID 3'-HYDROXYLASE (F3'H), SALICYLIC ACID
CARBOXYL METHYLTRANSFERASE (SAMT) and TREHALOSE-6-PHOSPHATE SYNTHASE
1 (TPS1). Notably detected is STARCH BRANCHING ENZYME I (SBE1), which controls starch
branching and may harbor allelic variants preferentially retained in ex-PVP lines for improved
grain quality or digestibility traits [47]. In total, 982 KEGG orthology (KOs) associations were
identified across these orthogroups (Supplementary Table 5). These KOs suggest hidden
variation in energy metabolism, detoxification, and hormone-mediated stress adaptation,
echoing findings in other crops where pangenome analyses revealed adaptive traits beyond
reference genomes [48]. These findings also demonstrate the value of the pangenome in
uncovering agronomically relevant diversity that is missed by single-reference approaches [49].

177
178
Fig. 2: Pangenome-Wide Gene Presence–Absence Variation Reveals Genes Missing from
the BTx623 Reference.
a, Classification of 36,004 gene families into core (green), soft-core (blue), dispensable
(orange), and private (red). **b,** Collectors curves showing pan- and core-genome orthogroups
dynamics across genomes sampled. **c,** Power law fit of novel gene family discovery indicating a
closed sorghum pangenome as evidenced by the plateauing collectors curve **d,** Stacked bar
plot comparing gene categories in the 'BTx623' reference (left) and the full pangenome (right),
the red bar reveals 6,389 additional gene families present in the pangenome. **e,** Gene family
presence–absence variation across accessions highlights shared patterns among ex-PVP lines.
Yellow indicates presence; black indicates absence. **f,** Genespace riparian plot illustrating

macro-synteny across sorghum genomes, revealing broad genic similarity and a mostly
conserved structure with a few notable rearrangements. **g,** a bar plot of enriched KEGG
orthology (for genes in pangenome-exclusive orthogroups). The orthologies include ATP
synthase subunit beta (atpB), NADH dehydrogenase subunits (ndhH, ndhI, ndhF, ndhB, ndhG),
E3 ubiquitin-protein ligase SIAH1 (SIAH1), ribosomal protein L36a (RPL36a), DNA-directed
RNA polymerase subunit alpha (rpoA), Flavonoid 3'-hydroxylase (F3'H), Salicylic acid carboxyl
methyltransferase (SAMT), Cinnamoyl-CoA reductase 1 (CCR1), Pyrimidine 5'-nucleotidase
(PYR5), Protein tyrosine phosphatase 1 (PTP1), Abscisic aldehyde oxidase 3 (AAO3),
ribosomal protein S3a (RPS3a), Ornithine decarboxylase 1 (ODC1), Trehalose-6-phosphate
synthase 1 (TPS1), Starch branching enzyme I (SBEI), and Glutathione S-transferase tau 1
(GSTU1), suggesting functional convergence in stress response, metabolism, and development.
The SBEI gene, which plays a role in starch metabolism and grain quality, is highlighted in red.
h, A bar plot showing how much each genome contributes to the pangenome-exclusive
orthogroups. Source data are provided in the Source Data file.
203

Structural Variants Underpin Genomic Diversity and Adaptive Traits in Elite ex-PVP
Sorghum Lines
A total of 1,118,641 SVs were identified across the pangenome, including insertions, deletions,
duplications, inversions, and translocations (Fig. 3a), with the highest individual SVs (42,096)
found in wild haplotype 1 (Supplementary Table 6). Insertions and deletions had median sizes
of 319 bp and 346 bp, respectively, reflecting a predominance of short to medium-sized INDELs
with potential regulatory or coding impacts (Fig. 3b). SVs also overlapped promoter and exon
regions, with mean lengths of 1,744 bp and 2,000 bp, respectively (Fig. 3c), suggesting
functional consequences for gene expression and/or protein activity. Macro-synteny analysis
revealed broadly conserved chromosomal architecture with lineage-specific large-scale
rearrangements, including a ~5 Mb inversion on chromosome 5 annotated to a wall-associated
kinase (WAK) gene associated with cold tolerance (Fig. 3d). Pairwise comparisons across the
pangenome highlighted extensive SV-based divergence, particularly between wild and elite ex-
PVP accessions (Fig. 3e, Supplementary Table 7). SVs were unevenly distributed across the
genome, with higher densities in pericentromeric and telomeric regions, consistent with
suppressed recombination and TE accumulation (Fig. 3f) [50].

Among key SVs, a 105 bp deletion was detected in *Sobic.001G445700.v5.1*, a gene
homologous to *Arabidopsis thaliana β-1,3-GLUCANASE A6* gene. The deletion removes a
conserved glycosidase domain (Fig. 3g), likely impairing its role in cell wall remodeling and
defense. The variant is predominantly present in elite ex-PVP lines and absent in wild
accessions, which retain the full gene (Fig. 3h), suggesting this deletion may reflect relaxed
defense investment under high-input agricultural systems (e.g., pesticide applications). This
pattern could result from reduced pathogen pressure in managed environments or represent a
trade-off favoring growth and yield over defense. Whole-genome alignments further confirmed
genotype-specific loss of functional sequence, as evidenced by the absence of mapped reads in
lines carrying the deletion (Fig. 3i). These findings highlight SVs as a distinct and functionally
relevant layer of genomic variation contributing to cultivated sorghum diversity and productivity.

Fig. 3: Structural Variants in the Sorghum Pangenome Impact Gene Function and Stress-Related Pathways.

a, Total number of structural variants (SVs) identified across the 50-genome pangenome; wild haplotype 1 exhibited the highest SV count (42,096), highlighting elevated genomic diversity. **b**, Length distribution of insertions and deletions (INDELs), highlighting the prevalence of short to medium-sized variants. **c**, Distribution of SVs lengths within promoter and exon regions, with potential impacts on gene regulation and coding sequences. **d**, Macro-synteny karyotype plot illustrating conserved chromosomal structure and large-scale rearrangements across sorghum genomes, including a ~5 Mb inversion on chromosome 5 linked to a wall-associated kinase (WAK) candidate gene associated with cold tolerance. **e**, Distribution of pairwise genomic differences across the pangenome, illustrating inter-accession genomic diversity. **f**, Genome-wide SVs density along sorghum chromosomes, with enriched regions corresponding to pericentromeric and telomeric zones. **g**, Pangenome graph and 3D structural visualization of the *Sobic.001G445700.v5.1* protein, a homolog of the *Arabidopsis thaliana β-1,3-GLUCANASE A6* gene involved in plant defense and cell wall remodeling. A 105 bp deletion in this gene disrupts the conserved glycosidase domain (black), likely impairing enzymatic function and suggesting a potential breeding signature for reduced defense investment under high-input agricultural systems. **h**, Presence-absence pattern of the *Sobic.001G445700.v5.1* SVs across the pangenome, distinguishing elite lines from wild accessions. **i**, In silico validation of the structural variant, based on whole-genome read alignments viewed in IGV, confirmed genotype-specific

258   loss of functional sequence—observed in accessions with the 105 bp deletion
259   (ExPVP_PI543243, ExPVP_PI601557) compared to those without (ExPVP_PI543246,
260   Wild_PI565549). Source data are provided in the Source Data file.
261
Selection Targets Circadian and Photoperiod Genes in U.S. Sorghum
We aligned the whole genome sequences of 46 ex-PVP and 25 wild sorghum accessions
(Supplementary Table 1) to the 'BTx623' reference genome. SNPs were called across all
accessions, and after quality filtering, 34,035 high-confidence biallelic SNPs were retained for
population structure analysis. Using this dataset, ADMIXTURE analysis (K = 1–10) identified K
= 2 as the optimal number of ancestral populations based on the lowest cross-validation error
(Supplementary Fig. 2a-b). PCA on the same SNP dataset further detected three genetic
clusters, separating ex-PVP lines from two distinct wild subgroups (wild1 and wild2;
Supplementary Fig. 2c-f). Based on this population structure, we performed genome-wide
selection scans using FST, XP-CLR, and π-ratio to identify genomic regions under differential
selection between wild and cultivated accessions.

Selective sweep analysis revealed that XP-CLR and π-ratio identified the most candidates,
while overlapping regions from all three methods defined smaller, high-confidence gene sets
under strong selection (Supplementary Tables 8-9). Multiple sweep regions differentiated wild
and elite ex-PVP lines (Fig. 4a, Supplementary Fig. 3b-d), with strong signals co-localizing with
known domestication genes involved in auxin transport, seed dormancy, and aromatic amino
acid biosynthesis (Fig. 4b). These pathways are critical for traits such as plant architecture,
seed dispersal, and germination.

Although a *β-1,3-GLUCANASE* showed structural disruption in ex-PVP lines (Fig. 3g-i),
genome-wide sweeps reveal positive selection on other immune-related pathways, such as
PATTERN RECOGNITION RECEPTOR (PRRs) signaling and innate immune activation,
suggesting a shift in defense strategy rather than an overall reduction. Enriched GO terms also
indicate that modern breeding emphasized immune signaling, xenobiotic transport,
phospholipase activity, and RNA export (Fig. 4b)—reflecting a broader focus on stress
resilience and metabolic efficiency under high-input agricultural systems.

Fig. 4: Selection Signatures During Sorghum Improvement Reveal Functional Divergence and Adaptive Loci.

a, Genome-wide selective sweep signals (FST, π, XP-CLR) across chromosomes comparing 25 wild vs. 46 ex-PVP accessions. The bottom track shows large structural variants identified across the chromosomes. The red arrow marks the position of a *WALL-ASSOCIATED RECEPTOR KINASE* gene *(Sobic.005G179200)* on chromosome 5 (~71 Mb), a member of a gene family known to regulate cell wall integrity and stress signaling under cold conditions. The blue arrow indicates the *PSEUDO-RESPONSE REGULATOR 37* gene (*SbPRR37*, *Sobic.006G057866*) on chromosome 6 (~41 Mb), a key regulator of photoperiod sensitivity and flowering time in sorghum. **b**, Gene Ontology (GO) enrichment of genes within top 5% sweep regions, highlighting domestication and improvement functions (e.g., auxin transport, immune signaling, dormancy) from various population comparisons and methods shown on the x-axis. **c**, Copy number variation of the *MULTIDRUG AND TOXIC COMPOUND EXTRUSION (MATE)* gene between representative wild and ex-PVP lines, showing approximately fourfold higher copy number in the ex-PVP lines. **d**, methylation percentage across *MATE* gene loci in ex-PVP, indicating low levels across the gene body region. **e**, methylation percentage across *MATE* loci in wild accession showing dynamic levels across the gene body region. Both accessions exhibit higher methylation levels in the promoter region. Source data are provided in the Source Data file.

Enrichment analysis of genes under selection highlighted key regulators of circadian rhythm and flowering time, including *PRR7* and *PRR95*, and *PRR95*, core components of the circadian clock, as well as *CO* and *PHOT1*, which mediate photoperiodic flowering (Supplementary Table 10). Genes involved in seasonal development—such as *FRIGIDA (FRI), VRN2B,* and *VRN3B*—were also enriched, suggesting coordinated selection on flowering pathways. Additional targets included *REGULATORY-ASSOCIATED PROTEIN OF TOR 1A (RAPTORA), TARGET OF*

RAPAMYCIN (TOR), GRAIN NUMBER, PLANT HEIGHT, AND HEADING DATE 7 / MATURITY
6 (GHD7/MA6), and *XAP5 CIRCADIAN TIMEKEEPER (XCT)*, implicating broader roles for
metabolic and growth regulation in adaptation [51]. These loci lie within sweep regions analogous
to those reported in other crops—for example, selection on *DRIVERS OF THE CIRCADIAN*
CLOCK (DOC) in barley has been shown to modulate rhythmicity and reproductive traits under
domestication [52]. The co-selection of these interconnected regulators underscores a
multifaceted strategy favoring clock-modulated traits, offering breeders a powerful entry point to
fine-tune flowering time, growth rate, and environmental responsiveness in modern sorghum
improvement programs.
324
Further analysis of genomic regions associated with flowering time, photoperiod sensitivity, and
circadian regulation revealed a strong signal at the *SbPRR37 locus (Sobic.006G057866)*, the
classical *MATURITY 1 (Ma1)* gene in sorghum. Its closest *Arabidopsis thaliana* ortholog is
PRR7 (Fig. 5a, Supplementary Table 11). We detected CNVs at *SbPRR37*, with two copies
present in most cultivated genotypes but only one in wild haplotype 2, suggesting reduced
redundancy or differential selection pressure in wild populations. Additional sweep intervals
contained other circadian and flowering regulators, including *PRR7* (Sobic.001G411400),
PRR95 (*Sobic.002G275100*), *Ma2* (*Sobic.002G302700*), and *GDH7/Ma6* (*Sobic.006G004400*),
supporting the hypothesis that multiple maturity genes (*Ma1–Ma6*) have been targets of
selection during domestication and improvement [36].
335
Several other sweep regions harbored genes involved in florigen signaling and circadian
entrainment, such as *VERNALIZATION 3B (VRN3B, Sobic.003G173032)*, *CONSTANS (CO,*
Sobic.010G115800), and *PHOTOTROPIN 1 (PHOT1, Sobic.008G001000)*. Selection signals
also overlapped with key metabolic and photomorphogenic regulators, including *TARGET OF*
RAPAMYCIN (TOR) (Sobic.009G109200), *REGULATORY-ASSOCIATED PROTEIN OF TOR*
*1A (*RAPTORa) (Sobic.005G008800), *XAP5 CIRCADIAN TIMEKEEPER (XCT)*
(Sobic.002G277600), and *DE-ETIOLATED 1 (DET1)* (Sobic.007G058100). Structural variation,
including PAVs and CNVs, was also evident at the *PHYTOCHROME A (PHYA)* and
*FLOWERING LOCUS T (*FT)*, alongside enrichment of core circadian components. Notably,
regulators such as *CIRCADIAN CLOCK ASSOCIATED 1 (CCA1)* appeared to be dicot-
specific—present in *Arabidopsis thaliana* and *Brassica napus* but absent from the sorghum
genome (Fig. 5).
348
349
350

Figure 5. Time-of-Day (TOD) Gene Variation and Model of the Sorghum Circadian Clock Network.

a, Variation in circadian clock and flowering-time genes reveals distinct selection pressures in cultivated sorghum. Copy number expansion of the circadian photoreceptor PHYTOCHROME A (PHYA) and the clock regulator PSEUDO-RESPONSE REGULATOR 37 (SbPRR37, Sobic.006G057866)—orthologous to PRR7 (AT5G02810) and PRR3 (AT5G60100) in Arabidopsis thaliana—reflects selection on the circadian system, while duplication of the florigen gene FLOWERING LOCUS T (FT) indicates selection on flowering-time pathways. PHYA and FT are highlighted in red boxes. These patterns suggest targeted adaptation to seasonal and latitudinal environments via independent tuning of clock and flowering components. **b**, Schematic representation of interlocked transcriptional feedback loops that drive circadian rhythms, allowing coordination with environmental signals such as light and temperature. The central area illustrates core clock regulatory interactions, adapted from Bendix et al. (2015) [36]. Yellow and black regions represent day and night, respectively. Arrows with pointed ends indicate activation; diamond-tipped arrows indicate repression. The orange-shaded middle layer denotes signaling and physiological pathways directly regulated by the circadian clock, while the outer blue-shaded ring highlights traits influenced by clock-controlled outputs. See supplementary notes for details. Source data are provided in the Source Data file.

Copy Number Expansion and Differential Methylation of the MATE gene in ExPVP lines
CNVs analysis based on read depth revealed four copies of the *MATE*
(*Sobic.003G403000.v5.1*) gene in ex-PVP lines compared to a single copy in the wild accession
(Fig. 4c). This copy number expansion coincides with a marked reduction in DNA methylation
across the gene body in the ex-PVP lines, whereas the wild accession exhibited higher and
more variable methylation across the same region (Fig. 4d). The lower methylation in ex-PVP
accessions, particularly within the gene body (Supplementary Fig. 4), suggests epigenetic
reprogramming likely associated with transcriptional activation of duplicated *MATE* copies. In
contrast, the higher and fluctuating methylation levels in the wild accession may reflect
transcriptional repression or tighter regulation (Fig. 4e). These patterns are consistent with
selective amplification and epigenetic remodeling of *MATE* in elite germplasm to support traits
such as xenobiotic transport or abiotic stress response [53].
393
394 **Discussion**
395 This study explores SVs, gene PAVs, and epigenomic remodeling in sorghum, uncovering
396 previously unrecognized diversity linked to key agronomic traits and environmental adaptation, a
397 variation often missed when relying on a single reference genome [1,20,46,54].
398
A substantial fraction of TEs in the pangenome remained unclassified, likely reflecting lineage-
specific or divergent TEs families underrepresented in existing repeat libraries [44,45]. These
findings are consistent with recent large-scale pangenome efforts in sorghum, where more than
200 Mb of novel sequence were uncovered across cultivated and wild accessions [20,21,55]. Our
analysis identified 28% of genes displaying PAVs, highlighting genomic diversity among elite
lines. This proportion is comparable to elite maize accessions, where ~32% of gene transcripts
in elite inbred lines display PAVs and are linked to stress- and environment-responsive
functions as well as heterosis [56].
407
408 The PAVs suggest that multiple gene combinations may have been retained or selected to
409 optimize adaptation and productivity across diverse growing environments. We discovered
410 *SBE1* gene among pangenome-exclusive genes. In cereals like maize and rice, *SBE1* is
411 associated with grain digestibility and quality, suggesting that its presence in elite U.S. sorghum
412 likely reflects selection for processing traits and end-use preferences [57].
413
Despite the pangenome showing strong genome-wide collinearity and preserved telomere
architecture, consistent with counter-selection against large-scale rearrangements [58], we
identified a notable ~5 Mb inversion on chromosome 5 in ex-PVP lines. This region includes a
WAK gene, part of a family known to regulate cell wall integrity and stress signaling in cold,
drought, and salinity conditions [59,60]. In rice and Arabidopsis, WAK genes play central roles in
early abiotic stress responses and cell wall remodeling, suggesting that this inversion may
represent targeted selection for cold tolerance in temperate breeding programs.
421
422
423

Our genome-wide selective sweep analysis revealed strong signals of selection on circadian
and photoperiodic regulators, extending beyond canonical domestication genes [30]. We detected
a sweep at *SbPRR37 (Sobic.006G057866)*, the gene underlying the *Ma1* locus, a major
repressor of flowering under long-day (LD) conditions. Recessive *ma1* alleles historically
enabled early flowering and adaptation to temperate latitudes [61,62], while dominant *Ma1* alleles
have been favored in modern biomass sorghum lines for delayed flowering under LD conditions
by increasing expression of *SbCO* which repress *SbFT* thus extended vegetative growth [63,64].
Our observation of CNVs at *SbPRR37*—with two copies in most cultivated lines and one in wild
haplotype 2—suggests differential selective pressure on photoperiod response across gene
pools.

Additional sweeps at *PRR7, PRR95, Ma2, Ma6, VRN3b,* and *CO* underscore the polygenic
nature of flowering-time control, integrating both classical *Ma* loci and newly implicated
regulators. Notably, genes such as *PHYA* exhibited copy number expansion in ex-PVP lines,
which may enhance photoperiod sensitivity or stabilize developmental transitions under variable
seasonal cues. The detection of enriched selection signals at *PHOT1, DET1,* and *XAP5*
suggests that upstream regulatory pathways have been knowingly or unknowingly targeted
during sorghum improvement. This highlights the importance of fine-tuned seasonal and
environmental sensing in shaping phenological plasticity. Similar patterns have been observed
in barley, where a natural variation at *Ppd-H1*, a *PRR7* ortholog, alters the circadian expression
of flowering time genes without disrupting core clock components, enabling latitude-specific
flowering responses [65]. In maize, selection on *ZmCCT9*, a diurnally regulated repressor of the
florigen *ZCN8*, delays flowering under LD conditions and enhances adaptation to higher
latitudes [66].

Beyond structural and genetic variation, we observed epigenomic divergence at functionally
relevant loci, consistent with evidence that epigenetic modifications can be subject to selection
and contribute to rapid adaptation, as demonstrated in *Arabidopsis thaliana* [67]. Notably, MATE
transporter genes exhibited both a fourfold copy number expansion and reduced gene-body
methylation in ex-PVP lines. These transporters mediate the efflux of organic acids, xenobiotics,
and flavonoids, with established roles in herbicide detoxification and aluminum tolerance in
cereals [68–71]. The observed gene-body hypomethylation may enhance expression under high-
input agricultural conditions, suggesting that coordinated structural and epigenetic remodeling
has been a feature of modern sorghum breeding.

Together, these results support a model in which sorghum's adaptation to temperate climates
and agricultural intensification has been shaped by diverse molecular mechanisms—including
targeted selection on flowering-time genes, clock genes, and epigenetic reprogramming of
stress and detoxification genes. Our study complements and extends recent sorghum
pangenome analyses [20,21,55,72], which established foundational catalogs of structural and gene-
content diversity across races and wild sorghum relatives. While those efforts focused on broad
germplasm panels, our study focuses on ex-PVP cultivars—historically important but under-
characterized—and adds critical insights into how modern breeding has reshaped the genome.
467

Although this study offers major insights, limitations remain. The ex-PVP panel studied here
represents lines from 1976–1992 and may not reflect recent commercial diversity. The
functional impacts of structural variants and epialleles also require validation. Future work
should integrate long-read resequencing, qPCR, and CRISPR to resolve complex loci.
Additionally, field-based time-series transcriptomics are needed to link genomic variation with
gene expression under natural conditions, improving our understanding of trait-environment
interactions and guiding precision breeding.
475
476 In conclusion, this study demonstrates how modern breeding has shaped the sorghum genome
477 through targeted structural, regulatory, and epigenetic changes, offering new opportunities for
478 trait dissection and breeding innovation. The resulting pangenome resource enhances our
479 ability to explore adaptation, gene-trait relationships, and the development of resilient, climate-
480 smart sorghum varieties to meet global food, feed, and bioenergy demands.
481
Methods
Plant Materials: Seeds for 46 elite ex-Plant Variety Protection (ex-PVP) sorghum lines were
obtained from the USDA Germplasm Resources Information Network (GRIN). These
accessions originated from the breeding programs of Pioneer Hi-Bred International, Inc. (n =
39), Novartis Seeds, Inc. (n = 1), Cargill Wheat Research Farm (n = 1), Holden's Foundation
Seeds, Inc. (n = 1), Walter Moss Seed Company, LLC (n = 1), Ring Around Products, Inc. (n =
1), and Northrup, King & Company (n = 2). Wild sorghum accessions were also sourced from
GRIN and selected to represent diverse geographic and taxonomic origins for comparative
genomic analyses. RNA-seq was performed on two sorghum accessions, 'PI329478' and
'PI510757', using eight tissues per genotype: seedling, 3-leaf, 5-leaf, tiller, boot, panicle with
anthers, root, and dough-stage panicle. Total RNA was extracted, quality-checked, and used to
construct poly(A)-selected libraries, which were sequenced on an Illumina platform to produce
paired-end reads (Supplementary Table 12).
High-Molecular-Weight (HMW) DNA Extraction and Sequencing: HMW genomic DNA was
extracted from 3-week-old seedlings using Carlson lysis buffer and Qiagen Genomic-tips,
following the Oxford Nanopore Technologies (ONT) 'Plant leaf gDNA' protocol for Arabidopsis.
Samples were ground in liquid nitrogen, followed by two chloroform:isoamyl alcohol (24:1)
washes. Total Pure NGS beads (Omega Bio-tek) were used for cleanup. DNA was size-
selected for fragments >10–25 kb using the ONT Short Fragment Eliminator Kit (EXP-SFE001)
and quality assessed using the Agilent Genomic DNA ScreenTape system (5067-5365) or
Femto Pulse Genomic DNA 165 kb Kit (FP-1002-0275). HiFi circular consensus sequencing
(CCS) libraries were prepared using the PacBio protocol (PN 101-853-100 v5). Genomic DNA
was sheared using Covaris g-TUBEs to a modal fragment size of ~18 kb, and fragments <10 kb
were excluded using BluePippin S1 High Pass 6–10 kb cassettes.

Genome Assembly and Annotation: Ex-PVP genomes were assembled from Oxford
Nanopore Technologies (ONT) reads using Flye (v2.9.2), followed by polishing with Racon
(v1.5.0) and Pilon (v1.24) using Illumina short reads. Wild genomes were assembled using
Hifiasm (v0.16.1-r159) with PacBio HiFi reads. Assemblies were scaffolded using RagTag
(v2.1.0) with the BTx623 reference genome [73]. For the wild accession 'PI156549', Hi-C reads
were additionally aligned to contigs using the Juicer pipeline (v1.6.2), followed by scaffolding
with 3D-DNA (v180419). Manual curation was performed using Juicebox Assembly Tools
(v1.11.08). Genome assembly quality was evaluated using BUSCO (v5.4.3) with the
embryophyta_odb10 dataset and the LTR Assembly Index (LAI), which provides a metric for
assembly continuity and repeat completeness based on intact long terminal repeat
retrotransposons. Gene models were predicted using Helixer, a deep learning-based annotation
tool trained on plant genomes [43].

Transposable Elements (TEs) Detection: TEs were annotated using EDTA (v2.0.0) with the
flags --sensitive 1 and --anno 1, which integrates both structure- and homology-based
approaches. The pipeline incorporates multiple tools, including LTRharvest, LTR_FINDER,
LTR_retriever, Generic Repeat Finder, TIR-learner, HelitronScanner, TEsorter, RepeatModeler,
and RepeatMasker. Interspersed and tandem repeats were further annotated using
RepeatMasker (v4.1.1) with the custom EDTA-derived TE library, and tandem repeats were
identified with TRF (v4.09).

Large Structural Variant Detection: Structural variants (SVs) were identified using a
combination of reference-based and assembly-based approaches based on 46 exPVP and wild
accession PI156549. SYRI (https://github.com/schneebergerlab/syri) was first used to
characterize large-scale structural rearrangements, including translocations and inversions. For
assembly-based SV calling, Svim-asm (v1.0.3) was applied to genome assemblies to detect
insertions, deletions, and complex events. Additionally, CuteSV (v2.1.2) was used to extract
duplications (DUP), inversions (INV), and breakends (BND). SV results were refined and
benchmarked using Truvari (v5.3.0), and nonredundant SV sets were merged using SURVIVOR
(v1.0.7). Whole-genome alignments were performed using Minimap2 (v2.29-r1283) to support
SV analysis. Pangenome graph construction and structural diversity visualization were
performed using PGGB (v0.6.0).

Synteny and Collinearity Analysis: Conducted at both genome and gene levels using D-
Genies (v1.4) with whole-genome alignments from Minimap2 (v2.22), and MCScan in the JCVI
package (v1.2.7) based on CDS alignments of longest isoforms generated using Last (v1418).

Identification of Orthologs: The interspecific all orthogroup gene pairs were determined using
OrthoFinder (v2.5.4, ortho_opt: -M msa -A mafft -T fasttree)

Single-Nucleotide Variant Calling, Admixture Analysis, and Population Structure: Genome
assemblies from 46 ex-PVP lines and 25 wild sorghum accessions (Supplementary Table 1)
were aligned to the BTx623 v5 reference genome [74], and variants were called using FreeBayes
(v1.3.6) with a ploidy-aware setting (-g 1000). Variants were filtered to retain sites with quality
score (QUAL) > 30, yielding 494,424 SNPs and 33,132 INDELs (Source:
Fig4_Sbic2groups_wild_cultivated_variants.v1.freebayes.filt.vcf.gz). Filtered SNPs were further
refined to retain only biallelic sites with MAF between 0.01 and 0.99 and no more than 10%
missing data, resulting in 34,035 high-quality SNPs. These were used for population structure
analysis. ADMIXTURE (v1.3.0) was run with five-fold cross-validation for K = 1 to 10, identifying
K = 2 as the optimal number of ancestral populations based on the lowest CV error
(Supplementary Fig. 2a-b). Population structure was also assessed using principal component
analysis (PCA) in PLINK (v1.90b7.7). PCA separated accessions into three distinct clusters,
with PC1 and PC2 explaining 30.34% and 11.42% of the variation, respectively (Supplementary
Fig. 2c-f).
Selection Signature Identification: We performed four pairwise comparisons using filtered
SNPs derived from whole-genome sequencing data: #1. Ex-PVP lines vs. all wild accessions.
#2. Ex-PVP lines vs. Wild1. #3. Ex-PVP lines vs. Wild2. #4. Wild1 vs. Wild2. For each
comparison, we employed three complementary methods to detect selective sweeps: FST
(fixation index):
Population differentiation was estimated using the Weir and Cockerham FST method
implemented in VCFtools. Analyses were performed using sliding windows of 100 kb with 10 kb
steps (--fst-window-size 100000 --fst-window-step 10000). Windows with FST > 0.3 were
considered indicative of strong differentiation. π-ratio (nucleotide diversity ratio): Nucleotide
diversity (π) was computed separately for each group using VCFtools (--window-pi 100000 --
window-pi-step 10000), and the ratio π exPVP / π wild was calculated. Windows with a π-ratio <
0.5 were interpreted as signals of reduced genetic diversity in the ex-PVP group, consistent with
selection. XP-CLR (Cross-Population Composite Likelihood Ratio): XP-CLR scores were
computed using XP-CLR (v1.1.2, https://github.com/hardingnj/xpclr), using a window size of 100
kb and step size of 10 kb. Windows in the top 1% (99th percentile) of XP-CLR scores were
considered candidate regions under strong selection. We further defined high-confidence sweep
regions using FST & XP-CLR intersection and FST & π-ratio & XP-CLR intersection. Only
windows meeting all three thresholds were retained in the final sweep set, enhancing
confidence that these represent genuine targets of selection during domestication or
improvement.
Circadian Clock, Photoperiod, and Flowering Time Genes: Sorghum core circadian clock,
light signaling, and flowering time orthologs genes were identified using known *Arabidopsis*
thaliana genes as the seed [75]

Gene Ontology (GO) Enrichment Analysis: Genes located within or overlapping candidate sweep regions (±100 kb) were extracted by intersecting sweep windows with mRNA coordinates from the Sorghum bicolor BTx623 v5.1 reference annotation (Phytozome). This was done using the PyRanges Python library. Genes from each metric-specific sweep set (e.g., FST-only, XP-CLR-only, triple-intersection) were analyzed separately. GO enrichment analysis was conducted using the GOATOOLS Python library (v1.2.3) with the find_enrichment.py script. GO term associations were generated from functional annotations predicted by eggNOG-mapper, and formatted into id2gos style association files. The GO DAG structure was derived from go-basic.obo. Each gene list was tested against a common background of all BTx623 genes with GO annotations. Statistically significant GO terms (corrected p-value < 0.05, using Benjamini-Hochberg FDR) were retained and reported.

Data Availability

All relevant datasets are provided in the main text and supplementary materials. Raw sequencing reads are publicly available under NCBI BioProject PRJNA1283509.
 Analysis scripts and pipelines are available on GitHub at https://github.com/jkitony/Pangenome-of-U.S.-ex-PVP-and-Wild-Sorghum.
 Genome assemblies and annotations can be accessed via the Salk Institute's resource portal (https://resources.michael.salk.edu/), Sorghumbase (https://www.sorghumbase.org/), and Figshare (https://doi.org/10.6084/m9.figshare.29261795.v4).
Source data for all figures are provided via Figshare.

Acknowledgements

We would like to thank the Donald Danforth Plant Science Center, the United States Department of Agriculture (USDA), and the Germplasm Resources Information Network (GRIN) for providing resources used in this study. This work was supported through the Salk Harnessing Plants Initiative (HPI) with funding from the TED Audacious, Bezos Earth Fund, and Hess Corporation.

References

1. Paterson, A. H. *et al.* The Sorghum bicolor genome and the diversification of grasses. *Nature* **457**, 551–556 (2009).

2. Khoddami, A. *et al.* Sorghum in foods: Functionality and potential in innovative products. *Crit. Rev. Food Sci. Nutr.* (2023) doi:10.1080/10408398.2021.1960793.

3. George, T. T. *et al.* Trends and progress in sorghum research over two decades, and implications for global food security. *S. Afr. J. Bot.* **151**, 960–969 (2022).

4. House, L. R. A Guide to Sorghum Breeding Second Edition. 220 (1985).

5. Prasad, P. V. V., Staggenborg, S. A. & Ristic, Z. Impacts of drought and/or heat stress on physiological, developmental, growth, and yield processes of crop plants. in *Response of Crops to Limited Water* 301–355 (American Society of Agronomy and Soil Science Society of America, Madison, WI, USA, 2015).

6. Conner, A. B. *Hybrid Vigor in Sorghum*. (Texas Agricultural and Mechanical College System, 1927).

7. Reddy, P. S. & Reddy, B. V. S. History of sorghum improvement. in *Breeding Sorghum for Diverse End Uses* 61–75 (Elsevier, 2019).

8. Bird, J. A. The Northwestern Miller. *The Miller Publishing Company* **191**, (1937).

9. Wayne Smith, C. & Frederiksen, R. A. *Sorghum: Origin, History, Technology, and Production*. (John Wiley & Sons, 2000).

10. Klein, R. R. *et al.* The effect of tropical sorghum con version and inbred development on genome diversity as revealed by high-resolution genotyping. *Crop Sci* **48**, S12–S26 (2008).

11. Frederiksen, R. A. Proceedings of the Consultative Group Discussion on Research Needs and Strategies for Control of Sorghum Root and Stalk Rot Diseases. *International Crops Research Institute for the Semi-Arid Tropics* (1983).

12. Shahbandeh, M. Sorghum production by leading country worldwide 2023/24. *Statista* (2024).

13. USDA. National Agricultural Statistics Service. *USDA* https://www.nass.usda.gov (2021).

14. Shakoor, N., Lee, S. & Mockler, T. C. High throughput phenotyping to accelerate crop

breeding and monitoring of diseases in the field. *Curr. Opin. Plant Biol.* **38**, 184–192 (2017).

15. McCormick, R. F. *et al.* The Sorghum bicolor reference genome: improved assembly, gene

annotations, a transcriptome atlas, and signatures of genome organization. *Plant J.* **93**,

338–354 (2018).

16. Zhang, Z. *et al.* Major impacts of widespread structural variation on sorghum. *Genome Res.*

34, 286–299 (2024).

17. Lye, Z. N. & Purugganan, M. D. Copy number variation in domestication. *Trends Plant Sci.*

24, 352–365 (2019).

18. Zhang, D. *et al.* Chromosomal inversion at the DG1 promoter drives double-grain spikelets

and enhances grain yield in sorghum. *Nat. Plants* **11**, 453–467 (2025).

19. Lynch, R. C. *et al.* Domesticated cannabinoid synthases amid a wild mosaic cannabis

pangenome. *Nature* (2025) doi:10.1038/s41586-025-09065-0.

20. Tao, Y. *et al.* Extensive variation within the pan-genome of cultivated and wild sorghum.

Nat. Plants **7**, 766–773 (2021).

21. Ruperao, P. *et al.* Sorghum pan-genome explores the functional utility for genomic-assisted

breeding to accelerate the genetic gain. *Front. Plant Sci.* **12**, 666342 (2021).

22. Boatwright, J. L. *et al.* Sorghum Association Panel whole-genome sequencing establishes

cornerstone resource for dissecting genomic diversity. *Plant J.* **111**, 888–904 (2022).

23. Voelker, W. G. *et al.* Ten new high-quality genome assemblies for diverse bioenergy

sorghum genotypes. *bioRxiv* (2022) doi:10.1101/2022.09.26.509520.

24. Liu, F. *et al.* Unravelling sorghum functional genomics and molecular breeding: past

achievements and future prospects. *J. Genet. Genomics* (2024)

doi:10.1016/j.jgg.2024.07.016.

25. Deng, Y. *et al.* A complete assembly of the sorghum BTx623 reference genome. *Plant*

Commun. **5**, 100977 (2024).

26. Semagn, K. *et al.* Genetic diversity and selective sweeps in historical and modern

Canadian spring wheat cultivars using the 90K SNP array. *Sci. Rep.* **11**, 1–16 (2021).

27. Chen, S. *et al.* Genome-wide study of an elite rice pedigree reveals a complex history of

genetic architecture for breeding improvement. *Sci. Rep.* **7**, 1–12 (2017).

28. Yamasaki, M. *et al.* A Large-Scale Screen for Artificial Selection in Maize Identifies

Candidate Agronomic Loci for Domestication and Crop Improvement. *Plant Cell* **17**, 2859–

2872 (2005).

29. Han, Z. *et al.* Genomic signatures and candidate genes of lint yield and fibre quality

improvement in Upland cotton in Xinjiang. *Plant Biotechnol. J.* **18**, 2002–2014 (2020).

30. Lin, Z. *et al.* Parallel domestication of the Shattering1 genes in cereals. *Nat. Genet.* **44**,

720–724 (2012).

31. Li, X. & Yu, J. Unraveling the sorghum domestication. *Mol. Plant* **15**, 791–792 (2022).

32. Zhang, L.-M. *et al.* Sweet Sorghum Originated through Selection of Dry, a Plant-Specific

NAC Transcription Factor Gene. *Plant Cell* **30**, 2286–2307 (2018).

33. Creux, N. & Harmer, S. Circadian rhythms in plants. *Cold Spring Harb. Perspect. Biol.* **11**,

a034611 (2019).

34. Michael, T. P. Time of day analysis over a field grown developmental time course in rice.

Plants **12**, 166 (2022).

35. McClung, C. R. Circadian clock components offer targets for crop domestication and

improvement. *Genes (Basel)* **12**, 374 (2021).

36. Bendix, C., Marshall, C. M. & Harmon, F. G. Circadian clock genes universally control key

agricultural traits. *Mol. Plant* **8**, 1135–1152 (2015).

37. Yan, H. *et al.* The genetic diversity of 69 widely used Chinese sorghum hybrids released

between the 1970s and 2010s. *Agronomy (Basel)* **14**, 2180 (2024).

38. Maazou, A.-R. S. *et al.* Optimizing use of U.S. Ex-PVP inbred lines for enhancing

agronomic performance of tropical Striga resistant maize inbred lines. *BMC Plant Biol.* **22**, 286 (2022).

39. Hamblin, M. T. *et al.* Challenges of detecting directional selection after a bottleneck: lessons from Sorghum bicolor. *Genetics* **173**, 953–964 (2006).

40. Jordan, D. R., Mace, E. S., Cruickshank, A. W., Hunt, C. H. & Henzell, R. G. Exploring and exploiting genetic variation from unadapted sorghum germplasm in a breeding program. *Crop Sci.* **51**, 1444–1457 (2011).

41. Coffman, S. M., Hufford, M. B., Andorf, C. M. & Lübberstedt, T. Haplotype structure in commercial maize breeding programs in relation to key founder lines. *Theor. Appl. Genet.* **133**, 547–561 (2020).

42. Deneweth, J., Van de Peer, Y. & Vermeirssen, V. Nearby transposable elements impact plant stress gene regulatory networks: a meta-analysis in A. thaliana and S. lycopersicum. *BMC Genomics* **23**, 18 (2022).

43. Stiehler, F. *et al.* Helixer: cross-species gene annotation of large eukaryotic genomes using deep learning. *Bioinformatics* **36**, 5291–5298 (2021).

44. Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. *Genome Biol.* **20**, 275 (2019).

45. Wicker, T. *et al.* A unified classification system for eukaryotic transposable elements. *Nat. Rev. Genet.* **8**, 973–982 (2007).

46. Zhao, K. *et al.* Pangenome analysis reveals structural variation associated with seed size and weight traits in peanut. *Nat. Genet.* 1–12 (2025).

47. Biswas, S. *et al.* Increasing the level of resistant starch in 'Presidio' rice through multiplex CRISPR-Cas9 gene editing of starch branching enzyme genes. *Plant Genome* **16**, e20225 (2023).

48. Zhou, Z. *et al.* Resequencing 302 wild and cultivated accessions identifies genes related to domestication and improvement in soybean. *Nat. Biotechnol.* **33**, 408–414 (2015).

49. Della Coletta, R., Qiu, Y., Ou, S., Hufford, M. B. & Hirsch, C. N. How the pan-genome is

changing crop genomics and improvement. *Genome Biol.* **22**, 3 (2021).

50. Liao, Z. *et al.* Structural variations in papaya genomes. *BMC Genomics* **22**, 335 (2021).

51. Zhang, H., Kumimoto, R. W., Anver, S. & Harmer, S. L. XAP5 CIRCADIAN TIMEKEEPER

regulates RNA splicing and the circadian clock by genetically separable pathways. *Plant*

Physiol. **192**, 2492–2506 (2023).

52. Prusty, M. R. *et al.* Genetic loci mediating circadian clock output plasticity and crop

productivity under barley domestication. *New Phytol.* **230**, 1787–1801 (2021).

53. Ku, Y.-S., Cheng, S.-S., Cheung, M.-Y. & Lam, H.-M. The roles of multidrug and toxic

compound extrusion (MATE) transporters in regulating agronomic traits. *Agronomy (Basel)*

12, 878 (2022).

54. Mace, E. S. *et al.* Whole-genome sequencing reveals untapped genetic potential in Africa's

indigenous cereal crop sorghum. *Nat. Commun.* **4**, 2320 (2013).

55. Marla, S. R. *et al.* Mining sorghum pangenome enabled identification of new dw3 alleles for

breeding stable-dwarfing hybrids. *G3 (Bethesda)* **15**, (2025).

56. Hirsch, C. N. *et al.* Draft assembly of elite inbred line PH207 provides insights into genomic

and transcriptome diversity in maize. *Plant Cell* **28**, 2700–2714 (2016).

57. Xia, H., Yandeau-Nelson, M., Thompson, D. B. & Guiltinan, M. J. Deficiency of maize

starch-branching enzyme I results in altered starch fine structure, decreased digestibility

and reduced coleoptile growth during germination. *BMC Plant Biol.* **11**, 95 (2011).

58. Lian, Q. *et al.* A pan-genome of 69 Arabidopsis thaliana accessions reveals a conserved

genome structure throughout the global species range. *Nat. Genet.* **56**, 982–991 (2024).

59. Kohorn, B. D. & Kohorn, S. L. The cell wall-associated kinases, WAKs, as pectin receptors.

Front. Plant Sci. **3**, 88 (2012).

60. Wu, Q. *et al.* CsWAK12, a novel cell wall-associated receptor kinase gene from Camellia

sinensis, promotes growth but reduces cold tolerance in Arabidopsis. *Front. Plant Sci.* **15**,

1420431 (2024).

61. Quinby, J. R. Fourth maturity gene locus in sorghum 1. *Crop Sci.* **6**, 516–518 (1966).

62. Quinby, J. R. The genetics of sorghum improvement. *J. Hered.* **66**, 56–62 (1975).

63. Murphy, R. L. *et al.* Coincident light and clock regulation of pseudoresponse regulator

protein 37 (PRR37) controls photoperiodic flowering in sorghum. *Proc. Natl. Acad. Sci. U.*

S. A. **108**, 16469–16474 (2011).

64. Mullet, J. *et al.* Energy sorghum--a genetic model for the design of C4 grass bioenergy

crops. *J. Exp. Bot.* **65**, 3479–3489 (2014).

65. Campoli, C., Shtaya, M., Davis, S. J. & von Korff, M. Expression conservation within the

circadian clock of a monocot: natural variation at barley Ppd-H1 affects circadian

expression of flowering time genes, but not clock orthologs. *BMC Plant Biol.* **12**, 97 (2012).

66. Huang, C. *et al.* ZmCCT9 enhances maize adaptation to higher latitudes. *Proc. Natl. Acad.*

Sci. U. S. A. **115**, E334–E341 (2018).

67. Schmid, M. W. *et al.* Contribution of epigenetic variation to adaptation in Arabidopsis. *Nat.*

Commun. **9**, 4446 (2018).

68. Maron, L. G. *et al.* Aluminum tolerance in maize is associated with higher*MATE1*gene copy

number. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 5241–5246 (2013).

69. Magalhaes, J. V. *et al.* A gene in the multidrug and toxic compound extrusion (MATE)

family confers aluminum tolerance in sorghum. *Nat. Genet.* **39**, 1156–1161 (2007).

70. Du, Z. *et al.* Genome-wide characterization of MATE gene family and expression profiles in

response to abiotic stresses in rice (Oryza sativa). *BMC Ecol. Evol.* **21**, 141 (2021).

71. Zhou, G., Delhaize, E., Zhou, M. & Ryan, P. R. The barley MATE gene, HvAACT1,

increases citrate efflux and Al(3+) tolerance when expressed in wheat and barley. *Ann. Bot.*

112, 603–612 (2013).

72. Chen, C. *et al.* A comprehensive omics resource and genetic tools for functional genomics

research and genetic improvement of sorghum. *Mol. Plant* **18**, 703–719 (2025).

73. Alonge, M. *et al.* RaGOO: fast and accurate reference-guided scaffolding of draft genomes.

Genome Biol. **20**, 224 (2019).

74. Gladman, N. *et al.* SorghumBase: a web-based portal for sorghum genetic information and

community advancement. *Planta* **255**, 35 (2022).

75. Michael, T. P. Core circadian clock and light signaling genes brought into genetic linkage

across the green lineage. *Plant Physiol.* **190**, 1037–1056 (2022).

803