Quantifying the effect of groundwater fauna and temperature on the ecosystem service of microbial carbon degradation

Susanne I. Schmidt[1], Nora Rütz[2], Jürgen Marxsen[2,3]

[1]Department of Lake Research, Helmholtz Centre for Environmental Research (UFZ), 39114 Magdeburg, Germany
5   [2]Institute of Animal Ecology, Justus Liebig University, 35392 Giessen, Germany
[3]Limnological River Station of the Max Planck Institute for Limnology, 36110 Schlitz, Germany

Correspondence to: Susanne I. Schmidt (Susanne.I.Schmidt@ufz.de; Susanne_I_Schmidt@yahoo.de)

10   **Abstract.** The groundwater ecosystem fulfils functions that humankind relies upon, e.g. for sustainable drinking water production. Quantification of these services is lacking so far, and thus, it is not possible to predict scenarios of e.g. future climates. Based on values from a comprehensive groundwater ecosystem study comprising four zones of varying land use and groundwater / surface water exchange, we parameterized a quantitative dynamic food web model (recharged organic carbon, microorganisms using the biodegradable fraction of this carbon, and fauna grazing on the microorganisms). With the model

15   satisfactorily reflecting the field values, we calculated five further scenarios, three of which without fauna (mortality e.g. due to contamination, sudden peaks of temperature etc.). Two of the "fauna" and two of the "no fauna" scenarios were run with temperature elevated by 1.5°C and 3°C, respectively. The ecosystem service of carbon degradation was expressed as the difference in carbon concentration between the beginning of the simulation and the end of the simulation. In most scenarios, remaining carbon increased over time. The remaining carbon in some scenarios was up to 660 times as high in the no fauna

20   scenarios compared to the reference case. Fauna was thus shown to fulfil a service by promoting microbial carbon degradation that may be substantial. Sustainable drinking water production is more reliable and less costly, the more active the groundwater fauna in the production area is. This model set up can serve to test other cases of varying physical and chemical variations and disturbances.

1 Introduction

Ecosystems fulfil functions that humans rely on. Groundwater is no exception (Avramov et al., 2010). The food web in groundwater (Malard et al., 2023) is often truncated (Gibert and Deharveng, 2002), but may encompass several trophic levels (Ercoli et al., 2019; Malard et al., 2023), depending on the respective environmental conditions. One of the functions that these food webs perform is processing organic carbon. This function of self-cleaning in groundwater, i.e. degrading carbon

35 compounds to ultimately the gases CO_2 or CH_4, is probably most relevant to humans. These functions are one prerequisite for sustainable drinking water production, which is one of the 17 goals (goal 6) of the 2030 Agenda for Sustainable Development (United Nations General Assembly, 2017).

Some groundwater food webs are qualitatively well described, in terms of e.g. "who eats whom" (Bradford, 2009; Bradford et

40 al., 2010; Ercoli et al., 2019; Francois et al., 2016; Leijs et al., 2015). Quantitative evaluations are a lot scarcer (but see e.g. Foulquier et al., 2011). To this date, no quantitative evaluation of ecosystem services in groundwater and their susceptibility to temperature existed. However, we urgently need to quantify the impacts of temperature on groundwater ecosystem services since we may have to future-proof drinking water production if groundwater ecosystems change their functionality due to increasing temperatures and changing climates and land uses.

Generally, at the bottom of a food web are producers. In groundwater, where light is lacking, however, primary production is only possible in the form of chemolithoautotrophy. While this is more prevalent than previously thought (Kellermann, 2008), heterotrophy, i.e. using organic resources that enter the groundwater ecosystem with groundwater recharge, will dominate in most places (Herrmann et al., 2015). Where heterotrophic degradation of imported organic material yields sufficient energy, a

50 further level of the food web may be supported, i.e. grazers consuming producers (we use the term producers here to encompass autotrophs and heterotrophs feeding on imported organics). Where even the producer biomass becomes substantial, a further level may be added, i.e. predators preying on grazers (Ercoli et al., 2019; Saccò et al., 2019). In Middle European aquifers, there are few taxa with just one feeding mode – most fauna will be grazing for at least part of their life cycle (Hose et al., 2022). It can be assumed that the biomass and energy in a respective level reduces to 10 % of the underlying level's biomass

55 and energy (Lindeman, 1942; Odum, 1957). The loss is due to respiration which occurs due to mobility, reproduction, hunting etc. (Kozłowski, 1992), and in a full budget, would result in outgassing of CO_2 (or CH_4 in an anoxic system; Herrmann and Taubert, 2022; Nydahl et al., 2020) from the respective compartment, as a result of respiration (Moinet et al., 2023). Under which circumstances which complexity of the food web will result, is still largely unknown and thus, impossible to parameterize.

It is well-known that grazing either reduces (Kota et al., 1999) or advances (Mattison et al., 2005) bacterial biodegradation of contaminants. Thus, adding levels to the food web might increase or it might compromise remediation and attenuation.

While structuring food webs with just producers, grazers, and predators is grossly simplifying, this pattern has successfully been used e.g. in conceptualizing biomanipulation where e.g. predatory, piscivorous fish are added to a water body which harbours producing algae, grazing zooplankton and zooplankton-consuming planktivorous fish (Carpenter et al., 1985; Vrba et al., 2024). In such systems, commonly, the planktivorous fish reduce the zooplankton so that the algae are not grazed upon, may grow to harmful algal blooms and thus, deteriorate the water quality (Dadi et al., 2023). By adding piscivorous fish, the planktivorous fish are reduced, thus, more zooplankton remains which grazes more efficiently the algal blooms (Triest et al., 2016). Adding piscivorous fish is thus one measure for improving water quality. Is such management possible in groundwater as well? Or, vice versa, would the loss of grazing fauna result in lower water quality?

In drinking water production, it is advantageous to have raw water with low organic substances so that little respiration (which would reduce the dissolved oxygen) takes place in the pipes on the way to the customers. Where dissolved oxygen is low or lacking, respiration of organic substances may lead to CH_4 which may be harmful to organisms and might thus limit degradation further. Anoxic conditions, resulting from respiration of excessive carbon sources, may lead to reduction of substances, and thus, dissolution of compounds which are bound in macro molecules at oxic conditions. E.g. aluminium, iron, etc. may become soluble at anoxic conditions (Dadi et al., 2023), and may be toxic to organisms. Thus, high raw water quality, with low levels of carbon, begins in the aquifers with a community of active organisms, degrading the recharged carbon and ensuring high oxic state throughout the drinking water production.

In addition to reduced conditions leading to undesired dissolved compounds, recently, brownification has become increasingly a problem (de Wit et al., 2016). One source of high molar mass organic compounds colouring the water yellow or brown is forest die back (Kopáček et al., 2023). The combination of dry summers stressing trees, and bark beetles making use of higher temperatures to produce more offspring than at colder climates (Seidl and Rammer, 2017) have led to expansive forest die back, e.g. in the Hercynian mountains (Musolff et al., 2024) in Germany and the Bavarian and Šumava mountains at the German-Czech boundary (Schmidt et al., 2024). This sudden increase of biomass is broken down to organic dissolved compounds many of which are undesired in drinking water production.

But higher temperatures have more, and more direct, effects than just favouring bark beetle generation times and dry summers. Dissolved oxygen solubility in water decreases with increasing temperatures (Retter et al., 2020), and brownification is believed to be exacerbated by increased temperatures (Blanchet et al., 2022). For these, and further, reasons, the temperature regime in a groundwater body is important when evaluating the current and potential future state of an aquifer. Globally, air temperature has increased by 1.5 °C compared to pre-industrial times in 2025 (https://climate.copernicus.eu/copernicus-2024-

first-year-exceed-15degc-above-pre-industrial-level; accessed 2 May 2025). The temperature increase in groundwater is delayed (Benz et al., 2017) and depends strongly on soil and geological properties and hydrological processes (Rau et al., 2015), but has, e.g. in Austria, reached approximately 0.4 ± 0.5 K per 10 years, meaning that an increase of 1.5 °C might be observable within 40 years. Benz et al. (2024) predicted that between 2000 and 2099, groundwater will warm on average by 2.1°C, under a medium emission pathway. It is thus important to estimate which effect these temperature increases will have on the groundwater ecosystem and the functions it fulfils.

On top of increased carbon input and higher likelihood for reduced conditions, increasing temperatures also impact groundwater organisms directly. E.g., increasing temperatures were related to increased microbial biomass in a temperature plume in Southern Germany, with the microbial community changing qualitatively (Brielmann et al., 2009). A change in microbial ecosystem function might follow but has so far rarely been characterized.

The microbial community is, however, the main food source for fauna. It is well known that groundwater fauna are less sensitive to some contaminants such as heavy metals than their surface relatives (Mösslacher, 2000), but are very sensitive towards temperature. Faunal diversities decreased with increasing temperatures within a temperature plume (Brielmann et al., 2009). This has implications on ecosystem services. Avramov et al. (2013) demonstrated the response of crustacean stress hormones to increasing temperatures. Groundwater niphargids and asellids differed in the temperature range they tolerated, with niphargids covering a wider range (Brielmann et al., 2011). Spengler and Hahn (2018) identified a threshold value for middle European groundwater systems of 13 °C above which fauna community changed drastically in composition and presumably function. Avramov (2013) and Brielmann et al. (2011) performed tests analogous to toxicity tests and extrapolated lethal temperatures for various durations of exposure. Like for the favoured temperature ranges, niphargids survived longer at higher temperatures than asellids. The lethal temperatures from these experiments can be used to parameterize fauna temperature-related mortality in models. Effects of temperature on the sublethal level, e.g. dissolved oxygen consumption, have also been quantified (Di Lorenzo & Galassi, 2017) and should be included in future models – here, we concentrate on the lethal temperature effect on fauna only.

In extreme cases, in summary, a temperature increase of a few °C may be lethal to groundwater fauna. Groundwater fauna is thus threatened from extinction, like, globally, a quarter of freshwater fauna (Sayer et al., 2025). In contrast to climate opportunists following along the preferred temperature range (Dormann et al., 2018), groundwater fauna's expansion or retreat is limited by the aquifers' physical boundaries. For the same reason, any faunal community not coping with increased temperatures will probably not be replaced by a higher-temperature-adapted community. With fauna's ecosystem function lacking, any effect of microorganisms being more active at higher temperatures, and thus degrading more carbon, may be (more than) compromised by the lack of grazing.

Since it is hard to derive the relevant data sets on the required temporal and spatial resolution, while controlling for different environmental settings globally, one way to approach answers to these questions are models. Here, we present a model based on data from one of the most comprehensive data sets on a groundwater ecosystem (Husmann and Marxsen, 1988; Marxsen et al., 2021). Marxsen et al. (2021) grouped the wells in the River Fulda floodplain into four groups: one group combining wells near the river Fulda and in intense exchange with the river ("R"), one group covering wells that were characterized by a plume of a carbon source from the near-by village ("P"), one group characterized by agriculture with elevated nitrogen and phosphorus concentrations ("A"), and a group representing wells of a mixing zone being situated in-between the other three zones and combining characteristics of all zones ("M"). While many groundwater taxa are ubiquitous, neighbouring aquifer zones may differ markedly in fauna taxa composition (Di Lorenzo, Amalfitano, et al., 2025). The River Fulda data set thus offers to study reactions to scenarios of increased temperatures and / or setting fauna to 0 as if after a calamity, in different environmental exchange conditions, under the same climate. The data set allowed to check whether the developed model represented a range of measured field conditions. Once satisfied with the model, it is possible to change parameters and / or exclude fauna biomass to run scenarios representing warmer climates, and comparing ecosystem functions (here: degradation of organic carbon) when fauna is present versus absent. Fauna being absent might also be a consequence of climate change, since, as described, fauna does not survive well elevated temperatures.

The hypotheses were: 1) Without fauna, a groundwater ecosystem shows lower carbon degradation, thus, higher remaining organic carbon. 2) An increase in temperature by 1.5°C or even 3°C leads to an increase in microbial degradation, and thus, lower remaining organic carbon. 3) Fauna is compromised more by increasing temperatures, than microorganisms profit from a temperature increase; thus, a scenario of modelling fauna at a temperatures increased by 1.5 °C and more so at 3°C will decrease the degradation of organic carbon, leading to higher organic carbon remaining than in the reference scenario with fauna at current temperature.

2 Study area and methods

The data that the modelling was based on, were taken from a field study in the Johannisaue, River Fulda floodplain, described in Husmann and Marxsen (1988) and Marxsen et al. (2021). This study is to this date one of the most comprehensive (chemical and physical conditions, prokaryotes and fauna; microbial growth rate measurements) and longest-running studies (measurements between 1978 and 1984), including dividing prokaryote cells into size fractions allowing detailed biomass distributions. Marxsen et al. (2021) had performed a k-means cluster analysis on the physical and chemical properties, assuming four groups. They characterized the four groups based on the chemical and physical properties of the groundwater sampled in the wells within the groups, and the wells' position in the floodplain (compare Fig. SI 1). These groups are used here for different exchange situations at the same climate.

Groundwater temperature was only measured at the time of sampling in the wells in the river Fulda floodplain, i.e. at the maximum monthly. Therefore, we took the daily temperatures from the DWD (German weather service) station Fulda Horas (1526, 255.00 m a.s.l.; geographical latitude 50.5335; longitude 9.6750; on the east shore of the river at the north of the investigated area). From the relationship of the minima and maxima of the *in situ* groundwater temperatures, compared to the minima and maxima of the Fulda Horas air temperature, we derived daily groundwater temperatures separately for the four groups of wells (Fig. SI 2). These were the basis for the scenario runs where temperature was one variable influencing maximum microbial growth rate and microbial half saturation concentration (refer to SI 2 for further explanations). We used the precipitation data from the same meteorological station 1526: Fulda Horas for estimating recharge (SI 3). We then modelled carbon import into groundwater by multiplying the respective daily recharge volume from precipitation with an average carbon concentration. All carbon sources, and all organism compartments, were transformed to mol COD, i.e. chemical oxygen demand, to allow directly comparing compartments. See SI 4 for these calculations.

For modelling microbial and faunal growth, we chose hybrid Monod/ Verhulst models (Schlogelhofer et al., 2021), combining resource-dependent growth with the limiting factor of a carrying capacity. For microbial growth, the equation reads

$$\frac{d MO_{het}}{dt} = \mu_{max,MO\_het} \, c_{MO_{het}} \left(1 - \frac{c_{MO_{het}}}{K_{MO_{het}}}\right)\left(\frac{c_{BOC}}{K_{S,BOC}+c_{BOC}}\right) \qquad \text{eq. 1}$$

where c_{BOC} is the concentration of biodegradable organic carbon (BOC; see below), c_{MO_het} is the concentration of heterotrophic microorganisms, μ_{max,MO_het} the maximum growth rate of heterotrophic microorganisms, K_{MO_het} the carrying capacity, and $K_{S,BOC}$ the half saturation concentration (Schlogelhofer et al., 2021). We derived growth rate and carrying capacity from the maximum biomass (dry mass) measured in the wells (SI 5, 6, 7). Modelled biomasses unlimited by a carrying capacity tend to increase exponentially because natural limiting processes such as heterogeneity of the environment are not incorporated in the model. The equation for faunal grazing on microorganisms was built accordingly. For fauna, the model was used with c_{MO_het} instead of c_{BOC}, c_{fauna} instead of c_{MO_het}, $\mu_{max,fauna}$ instead of μ_{max,MO_het}, K_{fauna} instead of K_{MO_het} for the carrying capacity, and K_{S,MO_het} instead of K_{BOC} for the half saturation concentration.

With the paradigm "Everything is everywhere but the environment selects" (Baas Becking, 1934) still holding (de Wit and Bouvier, 2006), microorganisms can be assumed to change in composition when environmental conditions change. The functionality largely remains the same, due to functional redundancy (Taubert et al., 2019). For microorganisms, there is no mortality in the stricter sense. Instead, maximum growth rate and carrying capacity are parameters usually describing growth quite well. Since most bioreactions are most efficient at temperatures between 25 and 35 degrees (Kong et al., 2020), temperature increases in groundwater can be expected to lead to increases in microbial functions up to ca. 35 °C, which will probably never be reached in groundwater unless under geothermal or anthropogenic influences. Therefore, we chose the

simplistic approach of modelling temperature increasing microbial growth rate, based on Schmidt et al. (2018). Due to the lack of knowledge on parameters for faunal growth rates, we chose a linearly increasing temperature-induced mortality instead, based on data in Avramov (2013; see below and SI 7).

Models were run separately for the four well groups derived in Marxsen et al. (2021). Some characteristics, such as air temperature, precipitation, TOC (total organic carbon) content in precipitation, fauna growth parameters μ_{max}, half saturation concentration K_s, and carrying capacity for microorganisms and fauna, fauna mortality rate, fauna excretion rate, were the same for all four groups, because group-wise measurements were lacking. The intercepts of detritus, DOM, heterotrophic microorganisms' biomass (dry mass), microbial growth rate parameters μ_{max}, half saturation concentration K_s, and fauna dry mass were the group-wise values from the start of the observation period, respectively.

The values for particulate organics measured in the River Fulda floodplain (Marxsen et al., 2021) were assumed here to be representative for detritus, and it was assumed that continuously, a fraction of 0.0001 per day would become biologically available, and this form was referred to as BOC here.

Since the remaining BOC concentrations after the modelled period of 3.5 years might just be a hot or cold spots or moments (McClain et al., 2003), we fitted trends over the modelled period to approximate overall development of carbon in the respective compartment.

All simulations, statistical evaluations, and plots were done in R 4.5.0 (R Core Team, 2025). Colours were chosen to be distinguishable by colour blind people using the package "ggokabeito" (Barrett, 2021).

3 Results

Precipitation and air temperature showed the expected seasonality, with the years 1978 having been drier, and the year 1981 wetter than the other years (Fig. 1 (a)). Figure 1 (b)-(e) show the measurement points and the simulations results (lines) over the investigated period. Figure SI 3 in SI 7 includes the trends added to Fig. 1 (b)-(e) as an example. The other figures with measured values and trends are not shown because the figures would have been largely redundant with the comparison of trends across scenarios (see below). The trend parameters of all scenarios are given in Table S1 in SI 9.

Figure 1: Measured and modelled carbon compartments in the river Fulda floodplain for which Marxsen et al. (2021) had divided groundwater wells into four groups: river-near = "R", mixing zone = "M", plume = "P", agricultural area = "A". (a) Air temperature (black) and precipitation in Fulda Horas meteorological station 1526 (blue) for the four years covering the study period. (b), (c), (d), (e): measured (dots) and modelled (lines) carbon compounds in the four well groups (refer to text and SI Fig. 1). This scenario is referred to as the "Reference" scenario. Please note that the two highest measured values in fauna dry mass ($1.3*10^{-4}$ mol COD L^{-1} in P03, group "R"; $1.8*10^{-5}$ mol COD L^{-1} in P26, group "A"; both on 2 January 1979) were omitted from the plot to enable visualization of the modelled time course.

235   The simulations (lines) in Fig. 1 (b), (c), (d), and (e) reflect the measured values (points) well in general. We therefore used this model and changed only the parameters for fauna and temperature for the scenarios of "no fauna" and/or "+ 1.5 °C", and/or "+ 3 °C", respectively. The respective overviews of the same format as Fig. 1 are included in the Supporting Information as Fig. SI 4, SI 5, SI 6, SI 7, and Fig. SI 8.

240   The modelled detritus did not change much during the modelled time span (Fig. 1 (b)). Event-based recharge in response to precipitation seemed to contribute little (Fig. 1 (b)). Modelled BOC first decreased then increased in the mixing "M" and the plume "P" zones, and in the mixing "M" zone and the agricultural "A" zone reached higher values at the end of the modelled time span than the measured values (Fig. 1 (c)). The increasing trend was consistent with an apparent increasing trend in the *in situ* values over the study period (Fig. 1 (c)). The microbial dry mass decreased temporarily in all scenarios (delayed in the

245   plume "P" zone), but to a different degree and for different time spans (Fig. 1 (d)). After about a third of the modelled time span, microbial dry mass started rising steadily and seemed to saturate around the maximum value measured which was set as the carrying capacity in all but the "plume" zone where the rise was slowest. Modelled fauna dry mass peaked within the first months in all zones, and decreased sharply from the average start values (note that two maxima values are not shown; see caption of Fig. 1) in the river-near "R" zone and the agricultural "A" zone (Fig. 1 (e)). For most of the modelled period fauna

250   dry mass stayed at low values.

An overview over the start and end values of the modelled run, and of the linear fit, along with the fit parameters, is given in SI 9, Table S 1. The comparison of all trends is given in Fig. 2 for BOC, for microbial dry mass in Fig. 3, and for fauna dry mass in Fig. 4. In all zones, the "no fauna" scenarios ("No fauna", "No fauna +1.5°C", "No fauna + 3°C") were

255   indistinguishable from each other and BOC trend rose to the highest values per zone (Fig. 2). The BOC trends in the three scenarios with fauna were lower than those in the "no fauna" scenarios, but rather indistinguishable from each other, particularly in the river-near "R" and the plume "P" zones. Where distinguishable at all, it was the "+3°C" scenario which showed the trend with a slope tending towards that of the "no fauna" scenarios. In the river-near "R" zone and the plume "P" zone, the trend for the three fauna scenarios decreased.

Figure 2: Modelled BOC in the river Fulda floodplain for the six scenarios which are given in the legend: "Reference": input variables according to Marxsen et al. (2021); "No fauna": setting all fauna variables to 0; "+1.5°C": raising all groundwater temperatures by 1.5°C; "No fauna +1.5°C": setting all fauna variables to 0 and raising all groundwater temperatures by 1.5°C; "+3°C: raising all groundwater temperatures by 3°C; "No fauna +3°C": setting all fauna variables to 0 and raising all groundwater temperatures by 3°C. The four sub plots depict the four aquifer groups: river-near = "R", mixing zone = "M", plume = "P", agricultural area = "A"; see text and SI 1.

Modelled microbial dry mass trends, like the BOC trends, were indistinguishable and showed the lowest slopes for the "no fauna" scenarios (Fig. 3). In the river-near "R" and in the agricultural "A" zones, the three fauna scenario trends were almost indistinguishable from each other. In the mixing "M" zone, the "+3°C" scenario trend was in-between that of "no fauna" and the two other "fauna" scenarios. Only in the plume "P" zone was the reference scenario that with the lowest slope of the three "fauna" scenarios (Fig. 3). In summary, microbial growth was steepest under the presence of fauna.

Figure 3: Modelled microbial dry mass in the river Fulda floodplain for the six scenarios: "Reference": input variables according to Marxsen et al. (2021); "No fauna": setting all fauna variables to 0; "+1.5°C": raising all groundwater temperatures by 1.5°C; "No fauna +1.5°C": setting all fauna variables to 0 and raising all groundwater temperatures by 1.5°C; "+3°C: raising all groundwater

temperatures by 3°C; "No fauna +3°C": setting all fauna variables to 0 and raising all groundwater temperatures by 3°C. Note the different scales on the y axis. The four sub plots depict the four aquifer groups: river-near = "R", mixing zone = "M", plume = "P", agricultural area = "A". Note the different scales of the y axes.

Faunal dry mass was of course 0 in all "no fauna" scenarios and is indicated in Fig. 4 only for completeness. Modelled fauna dry mass trends decreased in all "fauna" scenarios. The reference scenario trend was always the one with the highest intercept and steepest decrease, and, vice versa, the "+3°C" scenario was the one that came closest to the "no fauna" scenario zero trends.

Figure 4: Modelled fauna dry mass in the river Fulda floodplain for the six scenarios: "Reference": input variables according to Marxsen et al. (2021); "No fauna": setting all fauna variables to 0; "+1.5°C": raising all groundwater temperatures by 1.5°C; "No fauna +1.5°C": setting all fauna variables to 0 and raising all groundwater temperatures by 1.5°C; "+3°C": raising all groundwater temperatures by 3°C; "No fauna +3°C": setting all fauna variables to 0 and raising all groundwater temperatures by 3°C. Note the different scales on the y axis. The four sub plots depict the four aquifer groups: river-near = "R", mixing zone = "M", plume = "P", agricultural area = "A". Note the different scales of the y axes. The zero lines for the scenarios without fauna are only added for completion.

The ecosystem service of carbon degradation was expressed as the concentration of mol COD of biodegradable organic carbon remaining in the system compared to the start value of the simulation, with column "Modelled conc. at end minus modelled conc. time 0" being the difference between the columns "Modelled conc. time end" and "Modelled conc. time 0" (SI 9, Table S1). We divided this difference by the value in column "Modelled conc. time 0" to yield the value in the column "% change at end compared to modelled conc. time 0". BOC decreased by almost 100% to very low values in all river-near "R" and mixing zone "M" scenarios (SI 9, Table S1). This shows that it depends on the start condition and measured microbial rates per zone what the outcome of the simulation will be – there is no general behaviour. We then related the "Modelled conc. time end" of any scenario to the reference case. A value below 1 indicates a decrease in BOC compared to the reference case and was only observed in the river-near "R" zone at "+1.5°C" (SI 9, Table S1). In many cases, BOC rose slightly, indicated by

310 values just above 1, compared to the reference, e.g. in the other three zones in the "+1.5°C". The increase compared to the reference case was highest in the "no fauna" scenarios with factors of 1.37 to 661, i.e. the ecosystem service was so diminished that the remaining BOC increased to more than the 600 times the reference case outcome (SI 9, Table S1).

4 Discussion

This is to our knowledge the first attempt at developing a process-based model of the groundwater food web, and it is the first
315 attempt at comparing scenarios of excluding fauna from the system. There have been laboratory studies exposing fauna to increasing temperature (e.g. Brielmann et al., 2011; Di Lorenzo and Reboleira, 2022; Spengler, 2017), but to our knowledge this has never been translated to a trophic dynamic model, even if just restricting to heterotrophic microorganisms, as done here. One reason for such attempts not having been done before, lies in the difficulty of parameterizing such models. Our knowledge of the groundwater ecosystems' processes and dependencies is still patchy. Thus, the choice and development of
320 appropriate parameters is crucial. If the reference case is not appropriately parameterized, using it as a base for further scenarios is not feasible.

4.1 Modelling the reference case with fauna at the temperature of the observation period

Largely, the model set up, including the parameters, proved to be appropriate for representing the measured data. Modelled
325 detritus in the reference case with fauna at the temperatures of the observation period stayed in the same order as most of the measured values and was a constant source of carbon in abundance. We assumed here that detritus would largely stay in the catchment and thus not threaten drinking water production. We further assumed that BOC is the more influential factor for food webs and drinking water production, as e.g. in Vogt et al. (2023), and thus, we focussed on BOC and its pathway in the following.

Modelled BOC concentrations, designed to represent the bioavailable carbon fraction, mostly increased during the investigated period of 3.5 years in the mixing "M" and the agricultural "A" zone and rose slightly beyond the measured values in the "M" zone. The model thus did not indicate a quasi steady state. This may be an artefact of the combination of model parameters and needs to be studied further. However, this increase was in the order of 0.00015 mol COD L^{-1} over the investigated period
335 of 3.75 years, while the highest values of detritus measured in the field were around 0.05 mol COD L^{-1}. Thus, this BOC increase was negligible for the catchment-scale budget. It just means that, seen over the whole 3.5 year modelling period, less of the biologically available carbon pool was taken up by microorganisms than was refilled from precipitation, physical breakdown, and import from the soil. Assuming that the ecosystem was in a quasi steady state, the trends would have been expected to be even closer to 0 in the reference model case, but for a simplistic food web model, we believe that the current
340 approach is sufficiently close to the measured values. It further ensured that carbon was not limiting at the bottom of the food

web over the whole period. This enabled comparison of the degradation of this resource being in over abundance. It reflects the conditions in a high exchange zone such as a river floodplain. The less exchange with the surface, i.e. the deeper and the more secluded aquifers are, the more dependent on carbon import they will be (Schmidt et al., 2017; Schmidt and Hahn, 2012). Such scenarios will be part of the next development steps.

In the models, temporarily, modelled BOC was reduced to near-zero during the growth phase of the microorganisms (see below) in the river-near "R" and the plume "P" zones. This indicates feedback among the carbon fractions' concentrations – their scales seemed to be appropriate to reflect dynamic ecosystem behaviour. It also means that a longer-term simulation might have yielded different end concentrations and trends. This may be the subject of further model applications.

Modelled microbial dry mass showed temporary reductions to near-zero concentrations in the first modelled year. The early quite sudden dips might reflect moments of feeding by increasing fauna biomass (see below). The highest values for microbial dry mass had been measured in the agricultural "A" zone and concurrently, carrying capacity was set to be at this high level. In parallel, modelled BOC rose steadily and did not seem to be reduced by the high microbial dry mass. It is likely that the groundwater microbial community under agricultural land use exhibits growth and degradation characteristics much different from other zones (e.g. Dibbern et al., 2014), because subsurface carbon characteristics differ under the different land uses (Lehmann et al., 2021). It might be necessary to adapt the parameters and maybe even the model set-up for agricultural zones in future work. Another source of variation may be the distribution of organic carbon within the aquifer – in a porous environment, it might be physically separate from the microbial cells (Schmidt et al. 2017). To take such micro scale resource heterogeneity into account, requires a model which parameterizes at least two spatial dimensions (Schmidt et al. 2018), instead of the spatial 0D approach taken here.

Modelled fauna dry mass showed a rather sudden peak in the mixing "M" and plume "P" zone which might mean that some generation time term is missing in the model. Sudden decreases might point to mortality of organisms which then become detritus, but the sudden increase is physiologically not possible and unrealistic in resource-poor groundwater. All modelled fauna dry mass approached 0 over time, which might be an artefact due to the parameters not allowing more growth and grazing. This needs to be studied in further modelling attempts. With the end values being lower than the start values, the trends were all negative. The fauna trends decreasing to below 0 are an artefact and most likely due to high dry masses at the start, dropping down suddenly. In the cases where the dry mass stayed at elevated levels over at least a certain period at the start, and were not just outliers, i.e. in the reference case, the drop down was steeper than in scenarios where fauna decreased at the beginning. The reason for the fauna dry masses not increasing later when the microbial dry mass was at carrying capacity, i.e. when food was presumably not limiting, requires further investigation. Experiments on feeding and growth rates, building up on the published experiments on reproductive rates in Rütz et al. (2022) from central Germany, are still scarce and hardly available for middle-European temperate weather groundwater environments. We used data from three studies which had been

375 conducted at 14 and 17°C (Di Lorenzo et al., 2025), at 15°C (Navel et al., 2011), and at 10°C (Mermillod-Blondin et al., 2025). Thus, only the study by Mermillod-Blondin et al. (2025) was under conditions somewhat comparable with the River Fulda floodplain. The study by Navel et al. (2011) was even conducted with a 12 h light 12 h dark cycle because of the studied interaction with a surface amphipod. Both elevated temperature (see above) and light (Simčič and and Brancelj, 2007) may be stressful for stygobites though, and we did not deem the parameters appropriate for our model. This reduced the available

380 parameters. Considering these caveats, for the sake of a first simplistic food web model, we considered the model set-up sufficient for the purpose.

4.2 Modelling the lack of fauna

In agreement with the first hypothesis, all "no fauna" scenarios resulted in the highest remaining BOC, both in models, and in

385 the fitted trend lines, and resulted in mostly lower microbial dry mass, regardless of the zone. Less microbial growth because of the lack in rejuvenating grazing (Mattison et al., 2005) will have resulted in less BOC being degraded and thus, the BOC trends remaining at higher levels (see also above). It is interesting that this pattern emerges clearly from the rather simple food web modelling with the hybrid formula of combining the Monod model with the Verhulst model (Schlogelhofer et al., 2021).

390 The opposite effect of microbial dry mass being reduced due to grazing had also been observed (Kota et al., 1999). We lack understanding of the circumstances under which one pattern dominates. Also, the cited publications refer to grazing by protozoa, while we lacked quantitative field data for protozoa and assumed that the grazing can be subsumed under a multicellular "fauna" dry mass term, based on the fauna appraisal in the River Fulda floodplain. To date, there is no field or laboratory study quantifying multicellular fauna, protozoa, and microorganisms in the same samples.

Depending on the zone, remaining BOC concentrations decreased to the point of vanishing in one scenario (river-near "R" case at "+1.5°C"), but increased in all other cases. This increase was up to more than 600-fold over the 3.5 year observation period. Without groundwater fauna, drinking water production will have to compensate by intensified treatment of organics which raises costs and insecurity.

4.3 Modelling an increase of groundwater temperature

In contrast to the second hypothesis, an increase in temperature by 1.5°C or 3°C in the "with fauna" scenarios only led to lower remaining BOC in the 1.5°C scenario in the river-near "R" zone. Also, microbial growth trends did not show a clear pattern, with in some zones increasing beyond the reference case, in other decreasing. Unlike the BOC and microbial dry mass trends,

405 fauna dry mass trends in the three "fauna" scenarios showed a consistent pattern in all zones, with the reference case trends starting at the highest values and decreasing most sharply, and the "+3°C" scenarios showing the lowest absolute slopes. High

BOC values apparently did not translate to higher faunal dry mass although the modelled fauna dry mass never reached carrying capacity. This needs to be investigated further.

410 **4.4 Modelling the lack of fauna at an increase of groundwater temperature**

In partial agreement with the third hypothesis, the combination of increased temperatures and lack of fauna did lead to reduced degradation and higher remaining BOC compared to the reference scenario, but the difference to the BOC increase in the "no fauna at recent temperature" was negligible and thus, the reduced degradation was almost only due to lack of faunal grazing.

415 While the trends in the microbial dry mass were lowest in all "no fauna" scenarios, concurrent with the lack in rejuvenating grazing (Mattison et al., 2005; and see above), there were more differences among zones than with the BOC trends. Namely, the microbial dry mass trend slope in the mixing "M" zone was lowest for the "+3°C" scenario, indicating that in this zone at such elevated temperatures, the decrease in faunal grazing due to elevated temperature might not have been outweighed by increased microbial uptake with increasing temperature. The plume "P" zone and the agricultural "A" zone trends in microbial 420 dry mass were similar to each other, despite different intercepts. This is all the more surprising as the "A" zone harboured the highest BOC. It thus depended on the start values and the group-wise microbial growth rates, what the influence of temperature and faunal grazing were. The differences among groups were, however, largely smaller than the differences between "fauna" and "no fauna" scenarios.

425 The fact that the net microbial biomass in the elevated temperature scenarios in the "fauna" scenarios did not change clearly compared to the reference scenario, indicates that any temperature-induced increase in microbial biomass due to increased degradation and growth apparently was literally eaten up by fauna, despite the fauna being hampered itself by increased temperature. The effect of losing fauna was stronger than that of increased temperatures. The risk of losing fauna altogether increases with increasing temperatures, and thus, the risk of running into "no fauna" scenarios with clearly decreased 430 degradation increases.

4.5 Reduced ecosystem services

Here, we focused on degradation of BOC as a main ecosystem service relevant to drinking water production for humankind. In most scenarios, including all "no fauna" scenarios, BOC concentration increased, i.e. the remaining carbon increased, which 435 is an indicator of a reduced ecosystem service. Thus, the ecosystem service was reduced mainly under the lack of fauna. Losing groundwater fauna would mean losing a substantial part of the microbially-driven ecosystem service of carbon degradation.

This ecosystem service of degrading organic carbon comes at costs. E.g. the oxidation of carbon to CO_2 requires oxygen. There are a few laboratory studies quantifying the uptake of oxygen by groundwater fauna, e.g. Di Lorenzo et al. (2024), Hervant (1997), Hervant et al. (1998, 1999), Malard and Hervant (1999), and Mermillod-Blondin et al. (2008). Future studies should make dissolved oxygen a reaction partner in the organisms' uptake models, in order to evaluate where the provision of ecosystem services may be hampered by the aquifer's oxidation state.

4.6 Full budget

Full accounts of the productivity of groundwater systems with or without fauna, and at different temperatures, require incorporating CO_2 into the budget. While dissolved bicarbonate had been measured in the River Fulda aquifer study, some of the CO_2 (and CH_4 if produced; Herrmann and Taubert, 2022) will gas out. Such carbon loss from the system needs to be measured with the appropriate methods. Isotopes of the carbon atom may assist in such calculations (Schwab et al., 2019). A study on all carbon fractions in a groundwater ecosystem is still lacking, – the AQUADIVA project (Küsel et al., 2016) offers probably the most complete data set on all compartments. But important grazer groups such as the rotifera (Ejsmont-Karabin and Karpowicz, 2025) and protozoa (Sinclair et al., 1993) require different sampling methods than the larger fauna and are thus often not included in groundwater grazer accounts. For a full budget of the standing stocks and processes in the groundwater ecosystem, these compartments need to be included in future field studies and the models based on their results.

4.7 Extrapolating to the global scale

The total groundwater volume in the upper two km of the continental crust was modelled, based on tritium, to be approximately 22.6 Mio. km^3 (Gleeson et al., 2016). Of these, Gleeson et al. (2016) estimated that between 0.1 and 5 Mio. km^3 are less than 50 years old. Abbott et al. (2019) drew the conclusion that between 300 and 1200 km^3, with an average of 630 000 km^3, are renewable groundwater ("young" and mostly fresh). This "younger" groundwater is often the resource used for drinking water production.

We found increases in the BOC concentrations from the reference scenario to increased temperature and no fauna scenarios, with the increase being over 660 fold in the case of the "+3°C no fauna" scenario in the river-near "R" group, which was a difference of $1.6*10^{-5}$ mol L^{-1} over the observation period of 3.5 years, i.e. $4.6*10^{-6}$ mol L^{-1} $year^{-1}$. Assuming that this difference would be representative for global conditions, multiplying with the average of 630 000 km^3 for renewable groundwater (Abbott et al. 2019) and with a molar mass of 227 g mol^{-1} for humic acid (see SI 4), this translates to globally, 653,760,000 tons $year^{-1}$ not degraded by the groundwater ecosystem, when fauna is lacking. This is the additional load of COD that drinking water production needs to reduce globally in the worst case under a +3°C scenario when and if groundwater fauna has become extinct.

Extrapolating like this assumes that the situation in the Fulda floodplain is representative for the whole world, which is unlikely to hold true, for various reasons. One step in making such scenario accessible for the global situation is to use spatially resolved temperature scenarios such as by Benz et al. (2017, 2024) to extrapolate groundwater ecosystem services to the national, continental, and global scale. Another important step is to make not only the microbial degradation rate dependent on

475 temperature, but also the fauna grazing rate, not only the fauna mortality.

4.8 Necessary steps for further development

In this modelling study we considered different kinds of energies as equal, due to a lack in better knowledge and data. This approach may lead to errors (Odum and Odum, 2000). Odum and Odum (2000) suggested to express everything in an

480 ecosystem in the solar energy used to make each item. However, in an ecosystem where the sun does not shine, and where instead not only the import of plant material, but also chemolithoautotrophy without solar energy provide resources at the basis of the food web (Herrmann et al., 2015; Kellermann et al., 2012), compromises must be found.

Many temperature-related processes are not considered here, namely environmental changes, such as lower solubility of

485 dissolved oxygen, higher dissolution of ions (Retter et al., 2020), all of which may be detrimental to organisms. Thus, the net optimum for groundwater microbial activity is probably lower than the physiological activity optimum itself. However, we lack parameters for groundwater ecosystems to include such dependencies into models.

This study was based on a plethora of assumptions most of which might trigger discussions. Also, here, only average

490 compartment carbon concentrations were modelled. It is relevant to focus on the whole range of likely conditions, i.e. include minima and maxima. However, because of error propagation, this is not a straightforward process and requires diligent development. The present study aims to serve as a starting point for future developments.

Climate change does not happen everywhere in the same way – it not only translates in increased temperatures, but also in

495 changed recharge patterns which were not considered here. One next step is to use spatially resolved temperature scenarios and chemical reaction solvers such as PHREEQ-CE to extrapolate groundwater ecosystem services under climate scenarios to the national, continental, and global scale.

5 Conclusion

500   This first study on quantifying the change in ecosystem services at scenarios of losing groundwater fauna, and of increased temperature, showed that the partial loss of ecosystem services (BOC degradation) without fauna increased remaining BOC more than 660-fold, based on the 3.5 year observation period. While there were differences among aquifer zones that were characterized by differing patterns in nutrient and carbon background, the general patterns were the same within this catchment. This means that biomanipulation by protecting and promoting groundwater fauna for ameliorating water quality, might be a

505   promising concept. The fauna's role might still be underestimated. This has implications for drinking water production which will have to compensate the lack of fauna's ecosystem services by increased treatment which raises costs and insecurity.

Code and data availability

    Parts of the raw data from the River Fulda floodplain study had been published in Marxsen et al. (2021). The raw data and

510   code used in this publication are available on https://github.com/suisch/gtm_River_Fulda_floodplain.

Supplement

    The supplement related to this article is available in the file SI_Groundwater_ecosystem_model.pdf.

Author contributions

    SIS: conceptualization; SIS: methodology; SIS, JM, and NR: data curation; SIS, JM, and NR: writing – original draft

515   preparation; SIS, JM, and NR: validation; SIS, JM, and NR: writing – review and editing.

Competing interests

    The contact author has declared that none of the authors has any competing interests.

Acknowledgements

    SIS thanks Carola Winkelmann, Thomas Petzoldt, Martin Thullner, Jan Ulrich Kreft, Muhammed Shikhani, Martin Schultze,

520   and Susanne Worischka for discussions on modelling.

Financial support

This study was supported by a grant (Hu 41/17) from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation; www.dfg.de) to Dr. Siegfried Husmann (deceased).

References

525 Abbott, B. W., Bishop, K., Zarnetske, J. P., Minaudo, C., Chapin, F. S., Krause, S., Hannah, D. M., Conner, L., Ellison, D., Godsey, S. E., Plont, S., Marçais, J., Kolbe, T., Huebner, A., Frei, R. J., Hampton, T., Gu, S., Buhman, M., Sayedi, S. S., Ursache, O., Chapin, M., Henderson, K. D., and Pinay, G.: Human domination of the global water cycle absent from depictions and perceptions, Nat. Geosci., 12, 533–540, https://doi.org/10.1038/s41561-019-0374-y, 2019.

Avramov, M.: Sensitivity and Stress of Groundwater Invertebrates to Toxic Pollution and Changes in Temperature, PhD, 2013.

530 Avramov, M., Schmidt, S. I., Griebler, C., Hahn, H. J., and Berkhoff, S.: Dienstleistungen der Grundwasserökosysteme, KW Korresp. Wasserwirtsch., 3, 74–81, https://doi.org/10.3243/kwe2010.02.001, 2010.

Avramov, M., Rock, T. M., Pfister, G., Schramm, K.-W., Schmidt, S. I., and Griebler, C.: Catecholamine levels in groundwater and stream amphipods and their response to temperature stress., Gen. Comp. Endocrinol., 194, 110–117, https://doi.org/10.1016/j.ygcen.2013.09.004, 2013.

535 Baas Becking, L. G. M.: Geobiologie of Inleiding Tot de Milieukunde, Van Stockum & Zoon N.V., The Hague, 1934.

Blanchet, C. C., Arzel, C., Davranche, A., Kahilainen, K. K., Secondi, J., Taipale, S., Lindberg, H., Loehr, J., Manninen-Johansen, S., Sundell, J., Maanan, M., and Nummi, P.: Ecology and extent of freshwater browning - What we know and what should be studied next in the context of global change, Science of The Total Environment, 812, 152420, https://doi.org/10.1016/j.scitotenv.2021.152420, 2022.

540 Barrett, M.: Package: ggokabeito: "Okabe-Ito" Scales for "ggplot2" and "ggraph," 2021.

Benz, S. A., Bayer, P., and Blum, P.: Global patterns of shallow groundwater temperatures, Env. Res Lett, 12, 034005, 2017.

Benz, S. A., Irvine, D. J., Rau, G. C., Bayer, P., Menberg, K., Blum, P., Jamieson, R. C., Griebler, C., and Kurylyk, B. L.: Global groundwater warming due to climate change, Nat. Geosci., 1–7, https://doi.org/10.1038/s41561-024-01453-x, 2024.

Bradford, T.: Modes of Speciation in Subterranean Diving Beetles from a Single Calcrete Aquifer in Central Western Australia, 545 PhD thesis, University of Adelaide, 2009.

Bradford, T., Adams, M., Humphreys, W. F., Austin, A. D., and Cooper, S. J. B.: DNA barcoding of stygofauna uncovers cryptic amphipod diversity in a calcrete aquifer in Western Australia's arid zone, Mol. Ecol. Resour., 10, 41–50, https://doi.org/10.1111/j.1755-0998.2009.02706.x, 2010.

Brielmann, H., Griebler, C., Schmidt, S. I., Michel, R., and Lueders, T.: Effects of thermal energy discharge on shallow 550 groundwater ecosystems, FEMS Microbiol. Ecol., 68, 273–286, https://doi.org/10.1111/j.1574-6941.2009.00674.x, 2009.

Brielmann, H., Lueders, T., Schreglmann, K., Ferraro, F., Avramov, M., Hammerl, V., Blum, P., Bayer, P., and Griebler, C.: Oberflächennahe Geothermie und ihre potenziellen Auswirkungen auf Grundwasserökosysteme, Grundwasser, 16, 77–91, https://doi.org/10.1007/s00767-011-0166-9, 2011.

Carpenter, S. R., Kitchell, J. F., and Hodgson, J. R.: Cascading trophic interactions and lake productivity: Fish predation and 555 herbivory can regulate lake ecosystems, BioScience, 35, 634–639, 1985.

Dadi, T., Schultze, M., Kong, X., Seewald, M., Rinke, K., and Friese, K.: Sudden eutrophication of an aluminum sulphate treated lake due to abrupt increase of internal phosphorus loading after three decades of mesotrophy, Water Res., 235, 119824, https://doi.org/10.1016/j.watres.2023.119824, 2023.

Di Lorenzo, T. and Galassi, D.: Effect of temperature rising on the stygobitic crustacean species *Diacyclops belgicus*: Does global warming affect groundwater populations?, Water, 9, 951, https://doi.org/10.3390/w9120951, 2017.

Di Lorenzo, T. and Reboleira, A. S. P. S.: Thermal acclimation and metabolic scaling of a groundwater asellid in the climate change scenario, Sci. Rep., 12, 17938, https://doi.org/10.1038/s41598-022-20891-4, 2022.

Di Lorenzo, T., Mori, N., and Simčič, T.: Metabolic rates of groundwater species as a function of body mass and temperature, Subterr. Biol., 49, 53–74, https://doi.org/10.3897/subtbiol.49.125131, 2024.

Di Lorenzo, T., Amalfitano, S., Galassi, D. M. P., Melita, M., Zoppini, A., Parrone, D., Ghergo, S., Rossi, D., Tabilio Di Camillo, A., and Preziosi, E.: Geochemical and microbial factors driving crustacean assemblages in adjacent aquifer units within the same aquifer, Biogeosciences, 22, 1237–1256, https://doi.org/10.5194/bg-22-1237-2025, 2025a.

Di Lorenzo, T., Tabilio Di Camillo, A., Iepure, S., Galassi, D. M. P., Mori, N., and Simčič, T.: Oxygen consumption and carbon budget in groundwater-obligate and surface-dwelling *Diacyclops* species (Crustacea Copepoda Cyclopoida) under temperature variability, Environments, 12, 32, https://doi.org/10.3390/environments12010032, 2025b.

Dibbern, D., Schmalwasser, A., Lueders, T., and Totsche, K. U.: Selective transport of plant root-associated bacterial populations in agricultural soils upon snowmelt, Soil Biol. Biochem., 69, 187–196, https://doi.org/10.1016/j.soilbio.2013.10.040, 2014.

Dormann, C. F., Bobrowski, M., Dehling, D. M., Harris, D. J., Hartig, F., Lischke, H., Moretti, M. D., Pagel, J., Pinkert, S., Schleuning, M., Schmidt, S. I., Sheppard, C. S., Steinbauer, M. J., Zeuss, D., and Kraan, C.: Biotic interactions in species distribution modelling: 10 questions to guide interpretation and avoid false conclusions, Glob. Ecol Biogeogr, 27, 1004–1016, https://doi.org/10.1111/geb.12759, 2018.

Ejsmont-Karabin, J. and Karpowicz, M.: Rotifera Occurrence and Diversity in the Groundwater Wells of a Major European Wetland Complex, Int. Rev. Hydrobiol., 110, 22–28, https://doi.org/10.1002/iroh.70014, 2025.

Ercoli, F., Lefebvre, F., Delangle, M., Godé, N., Caillon, M., Raimond, R., and Souty-Grosset, C.: Differing trophic niches of three French stygobionts and their implications for conservation of endemic stygofauna, Aquat. Conserv. Mar. Freshw. Ecosyst., 29, 2193–2203, https://doi.org/10.1002/aqc.3227, 2019.

Foulquier, A., Malard, F., Mermillod-Blondin, F., Montuelle, B., Dolédec, S., Volat, B., and Gibert, J.: Surface water linkages regulate trophic interactions in a groundwater food web, Ecosystems, 14, 1339–1353, https://doi.org/10.1007/s10021-011-9484-0, 2011.

Francois, C. M., Mermillod-Blondin, F., Malard, F., Fourel, F., Lécuyer, C., Douady, C. J., and Simon, L.: Trophic ecology of groundwater species reveals specialization in a low-productivity environment, Funct. Ecol., 30, 262–273, https://doi.org/10.1111/1365-2435.12484, 2016.

Gibert, J. and Deharveng, L.: Subterranean ecosystems: A truncated functional biodiversity, BioScience, 52, 473–481, https://doi.org/doi: 10.1641/0006-3568, 2002.

Gleeson, T., Befus, K. M., Jasechko, S., Luijendijk, E., and Cardenas, M. B.: The global volume and distribution of modern groundwater, Nat. Geosci., 9, 161–164, https://doi.org/10.1038/ngeo2590, 2016.

Herrmann, M. and Taubert, M.: Biogeochemical Cycling of Carbon and Nitrogen in Groundwater—Key Processes and Microbial Drivers☆, in: Encyclopedia of Inland Waters (Second Edition), edited by: Mehner, T. and Tockner, K., Elsevier, Oxford, 412–427, https://doi.org/10.1016/B978-0-12-819166-8.00087-6, 2022.

Herrmann, M., Rusznyák, A., Akob, D. M., Schulze, I., Opitz, S., Totsche, K. U., and Küsel, K.: Large fractions of CO_2-fixing microorganisms in pristine limestone aquifers appear to be involved in the oxidation of reduced sulfur and nitrogen compounds, Appl. Environ. Microbiol., 81, 2384–2394, https://doi.org/10.1128/AEM.03269-14, 2015.

Hervant, F.: Reponses et adaptations a l'hypoxie severe chez des Crustaces souterrains et superficiels, Geobios, 21, 75–79, 1997.

Hervant, F., Mathieu, J., and Messana, G.: Oxygen consumption and ventilation in declining oxygen tension and posthypoxic recovery in epigean and hypogean crustaceans, J. Crustac. Biol., 18, 717–727, 1998.

Hervant, F., Garin, D., Mathieu, J., and Freminet, A.: Lactate metabolism and glucose turnover in the subterranean crustacean niphargus virei during post-hypoxic recovery, J. Exp. Biol., 202, 579–92, 1999.

605 Hose, G. C., Chariton, A., Daam, M. A., Di Lorenzo, T., Galassi, D. M. P., Halse, S. A., Reboleira, A. S. P. S., Robertson, A. L., Schmidt, S. I., and Korbel, K. L.: Invertebrate traits, diversity and the vulnerability of groundwater ecosystems, Funct. Ecol., 39, 2200–2214, https://doi.org/10.1111/1365-2435.14125, 2022.

Husmann, S. and Marxsen, J.: Besiedlung und Milieufaktoren im Grundwasser der Weserterrasse und der Fuldaaue, DVWK, Bedeutung biologischer Vorgänge für die Beschaffenheit des Grundwassers, Heft 80. Schriftenreihe des Deutschen Verbandes 610 für Wasserwirtschaft und Kulturbau e.V., Parey. Hamburg, 1988.

Kellermann, C., Selesi, D., Lee, N., Hügler, M., Esperschütz, J., Hartmann, A., and Griebler, C.: Microbial CO_2 fixation potential in a tar-oil-contaminated porous aquifer., FEMS Microbiol. Ecol., 81, 172–87, https://doi.org/10.1111/j.1574-6941.2012.01359.x, 2012.

Kellermann, C. S.: Autotrophy in groundwater ecosystems, PhD thesis, Ludwig-Maximilians-Universität München, 100 pp. 615 pp., 2008.

Khurana, S., Heße, F., Hildebrandt, A., and Thullner, M.: Predicting the impact of spatial heterogeneity on microbially mediated nutrient cycling in the subsurface, Biogeosciences, 19, 665–688, https://doi.org/10.5194/bg-19-665-2022, 2022.

Kong, F., Ren, H.-Y., Pavlostathis, S. G., Nan, J., Ren, N.-Q., and Wang, A.: Overview of value-added products bioelectrosynthesized from waste materials in microbial electrosynthesis systems, Renew. Sustain. Energy Rev., 125, 109816, 620 https://doi.org/10.1016/j.rser.2020.109816, 2020.

Kopáček, J., Grill, S., Hejzlar, J., Porcal, P., and Turek, J.: Tree dieback and subsequent changes in water quality accelerated the climate-related warming of a central European forest lake, J. Water Clim. Change, jwc2023581, https://doi.org/10.2166/wcc.2023.581, 2023.

Kota, S., Borden, R. C., and Barlaz, M. A.: Influence of protozoan grazing on contaminant biodegradation, FEMS Microbiol. 625 Ecol., 29, 179–189, https://doi.org/10.1016/S0168-6496(99)00010-0, 1999.

Kozłowski, J.: Optimal allocation of resources to growth and reproduction: Implications for age and size at maturity, Trends Ecol. Evol., 7, 15–19, https://doi.org/10.1016/0169-5347(92)90192-E, 1992.

Küsel, K., Totsche, K. U., Trumbore, S. E., Lehmann, R., Steinhäuser, C., and Herrmann, M.: How deep can surface signals be traced in the critical zone? merging biodiversity with biogeochemistry research in a Central German Muschelkalk landscape, 630 Front. Earth Sci., 4, 1–18, https://doi.org/10.3389/feart.2016.00032, 2016.

Lehmann, K., Lehmann, R., and Totsche, K. U.: Event-driven dynamics of the total mobile inventory in undisturbed soil account for significant fluxes of particulate organic carbon, Sci. Total Environ., 756, 143774, https://doi.org/10.1016/j.scitotenv.2020.143774, 2021.

Leijs, R., Bradford, T., Mitchell, J. G., Humphreys, W. F., Cooper, S. J. B., Goonan, P., and King, R. A.: The Evolution of 635 Epigean and Stygobitic Species of *Koonunga* Sayce, 1907 (Syncarida: Anaspidacea) in Southern Australia, with the Description of Three New Species, Plos One, 10, e0134673, https://doi.org/10.1371/journal.pone.0134673, 2015.

Lindeman, R. L.: The trophic-dynamic aspect of ecology, Ecology, 23, 399–417, 1942.

Malard, F. and Hervant, F.: Oxygen supply and the adaptations of animals in groundwater, Freshw. Biol., 41, 1–30, https://doi.org/10.1046/j.1365-2427.1999.00379.x, 1999.

640 Malard, F., Griebler, C., and Retaux, S.: Groundwater Ecology and Evolution, 2nd Edition., Academic Press, 2023.

Mermillod-Blondin, F., Poggiale, J.-C., Tolla, C., Auger, P., Thuiller, W., and des Châtelliers, M. C.: Using a mathematical model to simulate the influence of tubificid worms (Oligochaeta) on oxygen concentrations in hyporheic sediments, Fundam. Appl. Limnol., 172, 135–145, https://doi.org/10.1127/1863-9135/2008/0172-0135, 2008.

645 Marxsen, J., Rütz, N. K., and Schmidt, S. I.: Organic carbon and nutrients drive prokaryote and metazoan communities in a floodplain aquifer, Basic Appl. Ecol., 51, 43–58, https://doi.org/10.1016/j.baae.2020.12.006, 2021.

Mattison, R. G., Taki, H., and Harayama, S.: The soil flagellate *Heteromita globosa* accelerates bacterial degradation of alkylbenzenes through grazing and acetate excretion in batch culture, Microb. Ecol., 49, 142–150, https://doi.org/10.1007/s00248-003-0226-5, 2005.

McClain, M. E., Boyer, E. W., Dent, C. L., Gergel, S. E., Grimm, N. B., Groffman, P. M., Hart, S. C., Harvey, J. W., Johnston, 650 C. A., Mayorga, E., McDowell, W. H., and Pinay, G.: Biogeochemical Hot Spots and Hot Moments at the Interface of Terrestrial and Aquatic Ecosystems, Ecosystems, 6, 301–312, https://doi.org/10.1007/s10021-003-0161-9, 2003.

Mermillod-Blondin, F., Douady, C. J., François, C. M., Hervant, F., and Simon, L.: Environmental organic carbon availability determines metabolic and consumption rates in freshwater isopods, Oikos, e11149, https://doi.org/10.1002/oik.11149, 2025.

Moinet, G. Y. K., Hijbeek, R., van Vuuren, D. P., and Giller, K. E.: Carbon for soils, not soils for carbon, Glob. Change Biol., 655 29, 2384–2398, https://doi.org/10.1111/gcb.16570, 2023.

Mösslacher, F.: Sensitivity of groundwater and surface water crustaceans to chemical pollutants and hypoxia: implications for pollution management, Arch. Für Hydrobiol., 149, 51–66, https://doi.org/10.1127/archiv-hydrobiol/149/2000/51, 2000.

Musolff, A., Tarasova, L., Rinke, K., and Ledesma, J. L. J.: Forest Dieback Alters Nutrient Pathways in a Temperate Headwater Catchment, Hydrol. Process., 38, e15308, https://doi.org/10.1002/hyp.15308, 2024.

660 Navel, S., Simon, M., Lécuyer, C., Fourel, F., and Mermillod-Blondin, F.: The shredding activity of gammarids facilitates the processing of organic matter by the subterranean amphipod *Niphargus rhenorhodanensis*, Freshw. Biol., 56, 481–490, https://doi.org/10.1111/j.1365-2427.2010.02513.x, 2011.

Nydahl, A. C., Wallin, M. B., Laudon, H., and Weyhenmeyer, G. A.: Groundwater carbon within a boreal catchment: spatiotemporal variability of a hidden aquatic carbon pool, J. Geophys. Res. Biogeosciences, 125, e2019JG005244, 665 https://doi.org/10.1029/2019JG005244, 2020.

Odum, H. T.: Trophic Structure and Productivity of Silver Springs, Florida, Ecol. Monogr., https://doi.org/10.2307/1948571, 1957.

Odum, H. T. and Odum, E. P.: The energetic basis for valuation of ecosystem services, Ecosystems, 3, 21–23, https://doi.org/10.1007/s100210000005, 2000.

670 R Core Team: R: A Language and Environment for Statistical Computing. https://www.R-project.org/. Accessed 1 March 2025, 2025.

Rau, G. C., Cuthbert, M. O., Andersen, M. S., Baker, A., Rutlidge, H., Markowska, M., Roshan, H., Marjo, C. E., Graham, P. W., and Acworth, R. I.: Controls on cave drip water temperature and implications for speleothem-based paleoclimate reconstructions, Quat. Sci. Rev., 1–18, https://doi.org/10.1016/j.quascirev.2015.03.026, 2015.

675 Retter, A., Karwautz, C., and Griebler, C.: Groundwater microbial communities in times of climate change, Curr. Issues Mol. Biol., 41, 509–538, https://doi.org/10.21775/cimb.041.509, 2020.

Rütz, N. K., Marxsen, J., and Wolters, V.: Long-term cultivation of the groundwater amphipod *Niphargus aquilex* (Crustacea), Hydrobiologia, 850, 269–281, https://doi.org/10.1007/s10750-022-05058-6, 2022.

Saccò, M., Blyth, A. J., Humphreys, W. F., Kuhl, A., Mazumder, D., Smith, C., and Grice, K.: Elucidating stygofaunal trophic 680 web interactions via isotopic ecology, PLOS ONE, 14, e0223982, https://doi.org/10.1371/journal.pone.0223982, 2019.

Sayer, C. A., Fernando, E., Jimenez, R. R., Macfarlane, N. B. W., Rapacciuolo, G., Böhm, M., Brooks, T. M., Contreras-MacBeath, T., Cox, N. A., Harrison, I., Hoffmann, M., Jenkins, R., Smith, K. G., Vié, J.-C., Abbott, J. C., Allen, D. J., Allen,

G. R., Barrios, V., Boudot, J.-P., Carrizo, S. F., Charvet, P., Clausnitzer, V., Congiu, L., Crandall, K. A., Cumberlidge, N., Cuttelod, A., Dalton, J., Daniels, A. G., De Grave, S., De Knijf, G., Dijkstra, K.-D. B., Dow, R. A., Freyhof, J., García, N.,
685 Gessner, J., Getahun, A., Gibson, C., Gollock, M. J., Grant, M. I., Groom, A. E. R., Hammer, M. P., Hammerson, G. A., Hilton-Taylor, C., Hodgkinson, L., Holland, R. A., Jabado, R. W., Juffe Bignoli, D., Kalkman, V. J., Karimov, B. K., Kipping, J., Kottelat, M., Lalèyè, P. A., Larson, H. K., Lintermans, M., Lozano, F., Ludwig, A., Lyons, T. J., Máiz-Tomé, L., Molur, S., Ng, H. H., Numa, C., Palmer-Newton, A. F., Pike, C., Pippard, H. E., Polaz, C. N. M., Pollock, C. M., Raghavan, R., Rand, P. S., Ravelomanana, T., Reis, R. E., Rigby, C. L., Scott, J. A., Skelton, P. H., Sloat, M. R., Snoeks, J., Stiassny, M. L. J., Tan,
690 H. H., Taniguchi, Y., Thorstad, E. B., Tognelli, M. F., Torres, A. G., Torres, Y., Tweddle, D., Watanabe, K., Westrip, J. R. S., Wright, E. G. E., Zhang, E., and Darwall, W. R. T.: One-quarter of freshwater fauna threatened with extinction, Nature, 1–8, https://doi.org/10.1038/s41586-024-08375-z, 2025.

Schlogelhofer, H. L., Peaudecerf, F. J., Bunbury, F., Whitehouse, M. J., Foster, R. A., Smith, A. G., and Croze, O. A.: Combining SIMS and mechanistic modelling to reveal nutrient kinetics in an algal-bacterial mutualism, PLOS ONE, 16,
695 e0251643, https://doi.org/10.1371/journal.pone.0251643, 2021.

Schmidt, S. I. and Hahn, H. J.: What is groundwater and what does this mean to fauna? – An opinion, Limnologica, 42, 1–6, https://doi.org/10.1016/j.limno.2011.08.002, 2012.

Schmidt, S. I., Cuthbert, M. O., and Schwientek, M.: Towards an integrated understanding of how micro scale processes shape groundwater ecosystem functions, Sci. Total Environ., 592, 215–227, https://doi.org/10.1016/j.scitotenv.2017.03.047, 2017.

700 Schmidt, S. I., Kreft, J.-U., Mackay, R., Picioreanu, C., and Thullner, M.: Elucidating the impact of micro-scale heterogeneous bacterial distribution on biodegradation, Adv. Water Resour., 116, 67–76, https://doi.org/10.1016/j.advwatres.2018.01.013, 2018.

Schmidt, S. I., Fluksová, H., Grill, S., and Kopáček, J.: The significance of tree height as a predictor of tree mortality during bark beetle outbreaks in a small catchment, Forests, 15, 803, https://doi.org/10.3390/f15050803, 2024.

705 Schwab, V. F., Nowak, M. E., Elder, C. D., Trumbore, S. E., Xu, X., Gleixner, G., Lehmann, R., Pohnert, G., Muhr, J., Küsel, K., and Totsche, K. U.: 14C-free carbon is a major contributor to cellular biomass in geochemically distinct groundwater of shallow sedimentary bedrock aquifers, Water Resour. Res., 55, 2104–2121, https://doi.org/10.1029/2017WR022067, 2019.

Seidl, R. and Rammer, W.: Climate change amplifies the interactions between wind and bark beetle disturbances in forest landscapes, Landsc. Ecol., 32, 1485–1498, https://doi.org/10.1007/s10980-016-0396-4, 2017.

710 Simčič, T. and and Brancelj, A.: The effect of light on oxygen consumption in two amphipod crustaceans–the hypogean *Niphargus stygius* and the epigean *Gammarus fossarum*, Mar. Freshw. Behav. Physiol., 40, 141–150, https://doi.org/10.1080/10236240701452465, 2007.

Sinclair, J. L., Kampbell, D. H., Cook, M. L., and Wilson, J. T.: Protozoa in subsurface sediments from sites contaminated with aviation gasoline or jet fuel, Appl Env. Microbiol, 59, 467–472, 1993.

715 Spengler, C.: Die Auswirkungen von anthropogenen Temperaturerhöhungen auf die Crustaceagemeinschaften im Grundwasser -Versuch einer Prognose zur Klimaerwärmung und lokalen Wärmeeinträgen, University of Koblenz-Landau, 2017.

Spengler, C. and Hahn, J.: Thermostress: Ökologisch begründete, thermische Schwellenwerte und Bewertungsansätze für das Grundwasser, KW Korresp. Wasserwirtsch., 11, 521–525, https://doi.org/10.3243/kwe2018.09.001, 2018.

720 Taubert, M., Stähly, J., Kolb, S., and Küsel, K.: Divergent microbial communities in groundwater and overlying soils exhibit functional redundancy for plant-polysaccharide degradation, PLOS ONE, 14, e0212937, https://doi.org/10.1371/journal.pone.0212937, 2019.

Triest, L., Stiers, I., and Van Onsem, S.: Biomanipulation as a nature-based solution to reduce cyanobacterial blooms, Aquat. Ecol., 50, 461–483, https://doi.org/10.1007/s10452-015-9548-x, 2016.

725 United Nations General Assembly (Ed.): Transforming our world: the 2030 Agenda for Sustainable Development. 17 Sustainable Development Goals, in: A New Era in Global Health, Springer Publishing Company, New York, NY, https://doi.org/10.1891/9780826190123.ap02, 2017.

Vogt, R. D., Porcal, P., Hejzlar, J., Paule-Mercado, M. C., Haaland, S., Gundersen, C. B., Orderud, G. I., and Eikebrokk, B.: Distinguishing between sources of natural dissolved organic matter (DOM) based on its characteristics, Water, 15, 3006, 730 https://doi.org/10.3390/w15163006, 2023.

Vrba, J., Šorf, M., Nedoma, J., Benedová, Z., Kröpfelová, L., Šulcová, J., Tesařová, B., Musil, M., Pechar, L., Potužák, J., Regenda, J., Šimek, K., and Řeháková, K.: Top-down and bottom-up control of plankton structure and dynamics in hypertrophic fishponds, Hydrobiologia, 851, 1095–1111, https://doi.org/10.1007/s10750-023-05312-5, 2024.

de Wit, H. A., Valinia, S., Weyhenmeyer, G. A., Futter, M. N., Kortelainen, P., Austnes, K., Hessen, D. O., Räike, A., Laudon, 735 H., and Vuorenmaa, J.: Current browning of surface waters will be further promoted by wetter climate, Environ. Sci. Technol. Lett., 3, 430–435, https://doi.org/10.1021/acs.estlett.6b00396, 2016.

de Wit, R. and Bouvier, T.: 'Everything is everywhere, but, the environment selects'; what did Baas Becking and Beijerinck really say?, Environ. Microbiol., 8, 755–758, https://doi.org/10.1111/j.1462-2920.2006.01017.x, 2006.