Functional response predicts invasiveness but not trophic impact

Marine Courtois[1*], Chloé Souques [1*], Yann Voituron [1], Loïc Teulier [1], Vincent Médoc [2‡],

François-Xavier Dechaume-Moncharmont [1‡]

[1]Universite Claude Bernard Lyon 1, CNRS, ENTPE, LEHNA, UMR 5023, Villeurbanne, F-69100

[2] Equipe de Neuro-Ethologie Sensorielle (ENES), Centre de Recherche en Neurosciences de Lyon (CRNL), CNRS, INSERM, Universite de Lyon/Saint-Etienne, Saint-Etienne, France

* Equal contribution

‡ Equal contribution

Marine Courtois (marine.courtois@inrae.fr), ORCID : 0009-0002-3623-1460

Chloé Souques (chloe.souques@ univ-lyon1.fr), ORCID 0009-0004-4482-5829

Yann Voituron (yann.voituron@univ-lyon1.fr), ORCID 0000-0003-0572-7199

Loïc Teulier (loic.teulier@univ-lyon1.fr), ORCID 0000-0001-7779-7634

Vincent Médoc (vincent.medoc@univ-st-etienne.fr), ORCID 0000-0002-4888-1914

François-Xavier Dechaume-Moncharmont (fx.dechaume@univ-lyon1.fr), ORCID 0000-0001-7607-8224

Short title: Invasive species sampling range matters

Keywords: Alien species, Attack rate, Biological invasions, Ecological filtering, Feeding rates, Freshwater fish, Handling time, Impact prediction, Temperature.

Article type: Letter

Words in abstract: 150

Words in main text: 5157

Number of references: 109

Number of figures: 2 + Supplementary figures: 3

Number of Tables: 2 + Supplementary tables: 4

Corresponding author: Marine Courtois (marine.courtois@inrae.fr)

Statement of authorship: MC and CS equally contributed to this work. FXDM, LT and VM

conceived the original idea and developed the protocol. MC and CS conducted bibliographical

research and data collection. MC and FXDM performed the statistical analyses. MC and CS led

the writing of the manuscript largely supported by FXDM, LT, VM and YV. All authors

analyzed the data, contributed critically to the MS and gave final approval for publication.

Data accessibility statement: For the purpose of review, the raw dataset files and the complete

R script are available in a public repository on Gitlab at the following address:

https://gitlab.com/marinecourtois/fr_invasive_status. Submission to a permanent repository,

such as Zenodo, will occur upon the eventual final acceptance of the MS.

ABSTRACT. Biological invasions are recognized as one of the primary drivers of biodiversity

erosion mainly because invasive species wield greater trophic impact than their non-invasive

counterparts. The experimental paradigm for assessing the trophic impact of invasive species

is the functional response test. This test is frequently conducted on invasive species sampled

within their native ranges, implicitly assuming that such measures can be extrapolated to the

invasive range. Using a meta-analysis of 269 functional responses in 45 fish species, we

confirm that invasive species exhibited higher functional responses than non-invasive species.

However, this pattern holds true only when considering invasive species sampled in their native

range. Invasive species studied in their invasive range displayed functional responses

comparable to non-invasive species. These results raise questions on the use of functional

responses to predict invasiveness and trophic impact and on the mechanisms leading to trait

convergence between invasive and non-invasive species in introduced ecosystems.

Graphical abstract:

Comparative functional response between invasive and non-invasive species

INTRODUCTION

The interplay between species traits and the biological invasion processes continues to raise fundamental questions about ecological adaptation and species success and has far-reaching implications for biodiversity conservation and ecosystem resilience. Certain traits, such as selfing capacity in plants (Razanajatovo et al. 2016), physiology (Christensen et al. 2021) or personality in animals (Behrens et al. 2020; Chapple et al. 2012; Jolles et al. 2020) can be positively selected by the invasion processes, rendering the species that possess these traits potentially successful invaders. Post-establishment, species traits may be altered in response to new selective pressures within the recipient ecosystem, thereby inducing alterations in the environmental niche (niche shift, Broennimann et al. 2007; Guisan et al. 2014). For instance, the absence of natural enemies, including competitors, predators and parasites, has been identified as an important driver of invasion success (the Enemy Release Hypothesis, Colautti et al. 2004; Heger & Jeschke 2014; Keane & Crowley 2002; Meijer et al. 2016). Random variations in species traits may also be attributed to the founder effect, whereby introduced populations experience a genetic bottleneck whose magnitude is determined by population size, introduction frequency, and distance from source population (Dlugosch & Parker 2008; Dlugosch et al., 2015; Simberloff 2009). As a consequence of these non-mutually exclusive adaptive and neutral processes, species traits can diverge not only between native and invasive geographic ranges but also between introduced populations within the invasive range.

Among the traits likely to promote invasion success, those associated with resource acquisition have been extensively studied. Consumers exhibiting high feeding rates and/or broad diets are more likely to satisfy their nutritional needs in novel environments, but also a higher probability of disrupting the balance of local communities. Predation by invasive species is recognized as a major threat to biodiversity and has been estimated to cause a 21% decline in species richness in aquatic habitats (Mollot et al. 2017; Reid et al. 2019). In the invasion

ecology terminology, the Resource Consumption Hypothesis posits that invasive species are

more efficient consumers than non-invasive species (Faria et al. 2025; Ricciardi et al. 2013).

However, whether this represents a general pattern and functions as a cause or consequence of

the invasion process remains to be elucidated.

A validated methodology for quantifying trophic interaction strength consists in modeling

the functional response, which is defined as the relationship between *per* capita consumption

rate and resource density (Holling 1959). The shape of this response provides insights into

potential consequences for resource populations, with certain shapes indicating more

destabilizing effect than others (type-II versus type-III functional responses, Daugaard et al.

2019; Murdoch & Oaten 1975; Uszko et al. 2015). Additionally, it provides an overview on

consumers' behavior and how they attempt to maximize energy intake while minimizing costs

under conditions of resource limitation (Galarowicz & Wahl 2005; Oyugi et al. 2012; Murray

et al. 2013). The most prevalent functional response shape is characterized by a saturation curve

(Holling 1959) wherein the maximal consumption rate is constrained by handling time, which

is defined as the time required for a consumer to subdue, ingest, and digest a resource item

(DeLong 2021). The consumption rate at low resource densities provides an indicative of

consumer efficiency and is referred to as the space clearance rate or attack rate (DeLong 2021;

Jeschke et al 2002; Hassell & May 1973; Holling 1959).

Under the Resource Consumption Hypothesis invasive species are predicted to have

higher functional responses than non-invasive species (Bollache et al. 2008; Dick et al. 2013,

2014). If invasive species are predisposed toward strong trophic impacts, populations of the

invasive species in the donor area should exhibit higher functional responses than trophic

counterparts exhibiting no invasive tendency (Dick et al. 2014). Under these circumstances,

functional responses could be used as a screening mechanism for species invasiveness.

Conversely, if trophic impact is amplified post invasion in response to new environmental

conditions, such as prey naïveté (Cox & Lima, 2006; Sih et al. 2010), introduced populations

(IS-IR) should exhibit higher functional responses than the non-invasive (NIS) counterparts

with which they will compete (Xu et al. 2016). From this alternative perspective, comparative

functional response is a valuable tool for quantifying the trophic impacts on local resources.

Among the growing number of functional response comparisons made on plants,

invertebrates, and vertebrates (Alexander et al. 2014; Barrios-O'Neill et al. 2014; Ens et al.

2021; Funk & Vitousek 2007; Howard et al. 2018; 2022; Kattler et al. 2023; Landi et al. 2022),

several studies have reported that invasive species exhibit higher space clearance rates and

lower handling times than non-invasive species (*i.e.,* trophic analogs from the introduced range

without invasive status). A recent large-scale comparative study across a broad range of taxa

suggests that this may be a general pattern (Faria et al. 2025), although such studies remain

relatively scarce. This scarcity is primarily due to the absence of unifying experimental

framework, resulting in a divergence among functional response experiments, particularly in

aspects related to the characteristics of the interacting species (size, mass, density),

experimental protocol (duration of the test, area, volume and shape of the experimental arena)

and the environmental conditions (most notably temperature, which is of primary importance

for ectotherm species). This heterogeneity posed considerable challenges to comparative

functional responses until the development of a database that systematically collected

standardised functional response data from studies conducted worldwide. The FoRAGE

database (Uiterwaal et al. 2018) facilitates the investigation of the role of species and

environmental characteristics as potential drivers of functional response parameters. Using over

300 fish functional responses from this database, Buba et al. (2022) found that the alien status

was not a strong predictor of functional response parameters. However, the majority of the

functional responses for invasive species were obtained from populations of their native range.

These findings support the hypothesis that increased trophic impact may emerge post-invasion

as a consequence of new selective pressures, yet it leaves unresolved the broader question of

whether invasive and non-invasive species inherently differ in their functional response.

The objectives of this study are twofold. Firstly, we addressed the prevailing hypothesis

that invasive species exhibit higher functional responses than non-invasive species. Secondly,

the study aimed to establish whether functional responses are consistent when studies are based

on invasive species sampled in either their native or their introduced range. Assuming that

ecological novelty (*e.g.*, prey naïveté) exerts a more significant influence than preadaptation in

increasing the trophic impact of invasive species, we predicted that, invasive species from their

introduced range would demonstrate a higher functional response (*i.e.*, higher space clearance

rate, lower handling time, or both) than non-invasive species and invasive species from their

native range. Should such a discrepancy be observed, it would imply that results obtained from

tests carried out on individuals sampled in their native range should be interpreted with caution

when employed to predict the trophic impact of a species in its invasive range.

We conducted this comparative analysis in 45 freshwater fish species including 23 non-

invasive species and 22 invasive species sampled either in their native range or their introduced

range. Freshwater fish species are particularly sensitive to biological invasions and alterations

to their trophic networks (Albert et al. 2021; Reid et al. 2019; Sayer et al. 2025). They are

among the species for which the most data on functional responses are available (Uiterwaal et

al. 2018). The analysis of phylogenetic inertia was facilitated by the focus on a single,

homogeneous taxonomic group.

MATERIAL AND METHOD

Functional response data. The functional responses parameters used in this analysis were

obtained from the FoRAGE database (version 5 on 20 December 2024, Uiterwaal et al. 2018,

Uiterwaal et al. 2022), which is an extensive compilation of functional responses derived from

extant literature. These functional response parameters have been standardized to ensure the

comparability of the space and time units (see details below). The median values were reported

alongside the 95% quantiles of the bootstrapped parameter distributions. The database, when

available, also reported information regarding predators (species, body mass, life stage, juvenile

or adult), prey (species, body mass, life stage), and the original experimental conditions

(temperature, arena dimension). A total of 353 observations of functional responses in

freshwater fish species were identified in FoRAGE database (See PRISMA flowchart, Fig. S1).

Their functional responses were fitted in the database with a type II model using standardized

methods (Uiterwaal et al. 2018). The type II functional response is described by Holling's disk

equation (Eq. 1, Holling 1959):

164
$$F = \frac{aD}{1 + aDh}$$

This functional response equation relates consumption rate F (number of preys consumed per

unit of time) to prey density D (number of preys per unit of space, where space is generally a

volume of water in the case of fish species). The functional response is determined by two two

key parameters, a and h. The clearance rate or attack rate a (DeLong 2021; Holling 1959)

defines the volume of space cleared of prey per predator. It serves as a measure of predatory

efficiency at low prey densities (Jeschke et al. 2002). The handling time h corresponds to the

time required for a predator to capture, handle, and digest a prey item (DeLong 2021).

In the majority of functional response studies, the units of the parameters a and h remain

undefined. Authors typically compared responses of several populations within the same

species and the same study using identical experimental setup. As the primary focus of these

studies was on the significance of the difference between homogeneous experimental groups,

direct comparisons were made between the numerical values of the estimates, albeit without a

clear definition of the units. Nevertheless, it is imperative to unambiguously define these units

in order to allow meaningful comparisons across studies. In order to express the functional

response in homogeneous units, the focus was placed on studies based on 3D arena, i.e. where

space units were volumes (See PRISMA flowchart, Fig. S1).

In accordance with the standardized value reported in FoRAGE database, the space

clearance rate a was expressed in cubic meters per day per predator (units: $m^3.day^{-1}$). The

handling time h was expressed in time unit times number of predator per number of prey (units:

day). Our primary analyses were conducted using these *per capita* units (hereafter referred to

as 'standard units'), which focused on the number of prey individuals removed from the

environment. However, the number of prey items consumed varied by several orders of

magnitude, depending on whether the prey species in question are, for instance, Daphnia or

vertebrate species. From the perspectives of functional ecology or predator physiology, the

consequences of consuming one prey item are not equivalent when the predator and prey

species strongly differ in size. To account for substantial variability in body size, analyses were

repeated after converting a and h into prey and predator biomass (hereafter referred to as

'biomass units'). In these secondary analyses, biomass clearance rate a was expressed in cubic

meters per kilogram of predator per day (units: $m^3.kg^{-1}.day^{-1}$), and biomass handling time h was

expressed in time unit times kilogram of predator per kilogram of prey per day (units: day).

The analyses presented herein were based on a total of 269 estimates of a and h, reported

in a total of 45 freshwater fish species from 40 studies (See PRISMA flowchart, Fig. S1).

Interspecific differences were analyzed through a comparison of their mean values of the

functional response parameters a, or h, but also the Functional Response Ratio FRR = a/h,

which has recently emerged as a reliable metric for balancing the two functional response

parameters (Faria et al. 2025; Cuthbert et al. 2019b). Due to missing information about prey

body mass in the original articles, the analyses based on biomass units were conducted on 254

estimates for the handling time h and FRR a/h.

Invasive status and geographic range of sampling. The functional response database was

completed with the invasive status of each freshwater fish species specified as 'invasive' or

'non-invasive'. For this purpose, we relied on the Global Register of Introduced and Invasive

Species (GRIIS, Pagad et al. 2018), conceived within the framework of the Global Invasive

Alien Species Information (GIASI) partnership of the Convention on Biological Diversity

(CBD). It provides the first country-by-country checklists of invasive species. This

classification has been established on the basis of the number of countries in which the species

have been introduced and their ecological impact in the countries into which they were

introduced. The classification of a species as 'invasive' was determined by its introduction into

at least one country outside the limits of its native geographic range, and by the potential threat

to native biodiversity that its spread poses, which is referred to as "evidence of impact" in

GRIIS database. Following this classification, our study was based on a total of 22 invasive

species from 31 studies (168 observations functional response estimates for a and h), and 23

non-invasive species from 19 studies (101 observation of a and h). Furthermore, we

distinguished between the geographical regions ('native range' vs 'invasive range') from which

fish were sampled. All non-invasive species (NIS) were sampled in their native range.

However, for invasive species, it was necessary to differentiate between studies conducted on

invasive species sampled in their native range (IS-NR, donor population, 13 species from 20

studies, 107 estimates) and those conducted on invasive species sampled in their invasive range

(IS-IR, introduced population, 11 species from 13 studies, 61 estimates). The classification of

each functional response data in either NIS, IS-NR or IS-IR was determined by comparing the

sampling geographic range reported in each original study with the corresponding invasive

status and native range referenced in the GRIIS database. All observations identified in the

FoRAGE database that did not report information regarding the sampling area of the fish

species have been excluded from the analyses (see PRISMA flowchart, Fig. S1).

Effect of the temperature. Previous study (Buba et al. 2022) reported an effect of the

temperature at which the functional response assays were carried out. The temperature was thus

taken into account in the analyses. All observations identified in the FoRAGE database that did

not report information regarding the experimental temperature have been excluded from the

analyses (see PRISMA flowchart, Fig. S1).

Phylogenetic tree. Species that share a common ancestor often possess inherited traits due to

descent rather than independent adaptation. Some traits exhibit a strong phylogenetic signal,

meaning they are conserved along lineages. Consequently, closely related species cannot be

considered as independent data points. In addition, certain taxa may be overrepresented in the

literature, potentially biasing analyses. Therefore, comparative analyses must incorporate

phylogeny to enhance their robustness and reduce potential biases (Cinar 2022; Lajeunesse et

al. 2013). Our analyses took into account the phylogenetic relationships between the fish

species. The phylogenetic correlation matrix was integrated at species-level in all our models.

This matrix was derived from the phylogenetic tree (Fig. 1) constructed for the 45 freshwater

fish species present in our dataset, with the R package 'FishPhyloMaker' (Nakamura et al.

2021). Within this tree, invasive species were distributed across all families or orders, with no

single taxon accounting for a disproportionately large number of species (Fig. 1). Lynch's

formula was applied to estimate the phylogenetic signal (or phylogenetic heritability) for each

functional response parameter expressed in both standard and biomass units (Garamszegi

2014).

Figure 1: *Phylogenetic tree of the 45 freshwater fish species used in the analyses. Each colored circle in front of the species names illustrates the status of the species: non-invasive species (NIS, blue circle, all were sampled in their native range), invasive species sampled either in their native range (IS-NR, donor populations, orange circle) or in their invasive area (IS-IR, introduced populations, red circle). Some invasive species have been sampled both in their native range and in their invasive range depending on the specific study undertaken (IS-NR and IS-IR, bicolor circle). Taxonomic rank is reported at the order (in grey in dotted boxes) or family level (in bold black, on the right side).*

Statistical analysis. All analyses were carried out separately for each functional response

parameters (a, h, FRR, both in the standard units and in biomass units). Due to the wide

variation between species, the distributions of these parameters were highly skewed. We used

log transformation to normalize data and make it symmetric, thus meeting the assumptions of

normality and homoscedasticity. There was no missing data in our datasets, thereby enabling

model selection based on information theory (Nakagawa & Freckleton 2011) to test whether

invasive status (non-invasive vs. invasive species) or sampling range (NIS, IS-NR, IS-IR)

explained the differences in functional response parameters.

The phylogenetic correlation matrix derived from our phylogenetic tree was used as random

variable in all models using a phylogenetic mixed model approach (Hadfield & Nakagawa

2010) with the R package MCMCglmm for Bayesian mixed models (Hadfield 2010), assuming

a Brownian model of trait evolution (Hadfield & Nakagawa 2010). The Markov-chain Monte

Carlo (MCMC) was used with 3×10^5 iterations, 300 burn-in period and a thinning interval of

100 and an inverse-Wishar prior. Prior type had no impact on the conclusions that were drawn

from the analyses.

For each functional response parameter (a, h, or FRR) and for each unit (standard or

biomass unit), several competitive models were considered. The simplest model, 'Model 0',

solely took into account phylogenetic correlation matrix as random variable. No discrimination

was made between invasive status (non-invasive vs. invasive species) or sampling status (NIS,

IS-NR, IS-IR). 'Model 1' considered the invasive status, invasive (IS) versus non-invasive

(NIS) species, as fixed effect, irrespective of the geographic area in which the species had been

sampled. 'Model 2' considered the sampling status as fixed effect. This composite variable was

a three-level factor: non-invasive species (NIS, which were always sampled in their native

range), invasive species sampled in their native range (IS-NR, donor population), and invasive

species sampled in their invasive range (IS-IR, introduced population). Because Model 1 and

Model 2 were not nested, it was not possible to directly compare them using likelihood ratio

test. They were compared using information criterion (Burnham & Anderson, 2004). Model

selection was performed using the deviance information criterion (DIC, Spiegelhalter et al.

2002). The models with the lowest DIC value were considered the best models. Models with

ΔDIC > 2 were considered to have significantly worse fit, and models with ΔDIC < 2 were

considered to have equivalent support to the best model (Barnett et al. 2010). All these analyses

were repeated separately for each functional response parameter (a, h, FRR) both in standard

units and in biomass units.

The temperature was taken into account as fixed variable. 'Model 3' considered the

temperature, the sampling status, and their interaction as fixed effects. The effects of the

explanatory variables and their interaction term were analyzed by comparing the DIC of the

nested models, using the same model selection procedure as described above.

Data processing and analysis were carried out with the R programming language

(version 4.4.3, R Development Core Team 2025).

RESULTS

Effect of the invasive status. The functional response parameters expressed in standard units

varied significantly according to the invasive status (non-invasive vs. invasive species). Space

clearance rate a (posterior mean = -1.07, 95% credible interval 95%CrI = [-1.63; -0.55],

pMCMC = 6.7×10^{-04}, Fig. 2A), as well as FRR = a/h (posterior mean = -1.8, 95%CrI = [-2.83;

-0.94], pMCMC = 3.0×10^{-04}) were significantly higher for invasive species than for non-

invasive species, and handling time h (posterior mean = 0.82, 95%CrI = [0.18; 1.54], pMCMC

= 0.022) were significantly lower for invasive species than non-invasive. Similar effects,

though less pronounced, were found for functional response parameters expressed in biomass

units: space clearance rate a (posterior mean = -0.49, 95%CrI = [-0.94; -0.028], pMCMC =

0.043), FRR a/h (posterior mean = -1.12, 95%CrI = [-2.01; -0.26], pMCMC = 0.019), and

handling time h (posterior mean = 0.60, lower and upper 95%CrI= [-0.010; 1.23], pMCMC =

0.056).

Effect of the sampling status. The ΔDIC > 6 in the model selection procedure (Table 1)

indicated strong evidence that the 'Model 2' based on sampling status (NIS, IS-NR, IS-IR) was

a substantially better fit of the data compared to 'Model 1' based solely on the invasive status

(non-invasive vs. invasive species) for all functional response parameters expressed in standard

units. When invasive species were separated into two distinct groups, donor populations (IS-

NR) and introduced populations (IS-IR), donor populations showed higher space clearance rate

a (posterior mean = 0.88, 95%CrI = [0.49; 1.27], pMCMC < 3×10^{-04} ; Fig. 2B), lower handling

time h (posterior mean = -1.18, 95%CrI = [-1.89; -0.44], pMCMC = 1.3×10^{-03} ; Fig. S2A) and

higher FRR a/h (posterior mean = 2.11, 95%CrI = [1.22; 2.97], pMCMC < 3×10^{-04}, Fig. S2C)

than introduced populations. Furthermore, introduced populations and native species did not

differ in space clearance rate a (posterior mean = -0.56, 95%CrI = [-1.15; -0.02], pMCMC =

0.062, Fig. 2B), handling time h (posterior mean = 0.038, 95%CrI = [-0.83; 0.82], pMCMC =

0.92, Fig. S2A) and FRR a/h (posterior mean = -0.51, lower and upper 95%CrI = [-1.55; 0.50],

pMCMC = 0.33, Fig. S2C). Fully consistent results were observed for the three parameters a,

h, and FRR expressed in biomass units (Table S1, Fig. S3A,C,E).

Table 1. *Model selection table for the influence of sampling status on each functional response*
parameter (space clearance rate a, handling time h, and functional response ratio FRR = a/h) expressed
in standard units. The competitive models were ranked according to their DIC. For a detailed
description of the three models, refer to the main text.

Model	Clearance rate a		Handling time h		FRR = a/h	
	DIC	ΔDIC	DIC	ΔDIC	DIC	ΔDIC
Model 2 (sampling status)	579.48	0	1052.58	0	1105.65	0
Model 1 (invasive status)	597.94	18.46	1058.67	6.09	1121.22	15.57
Model 0 (intercept only)	598.97	19.49	1059.72	7.14	1124.03	18.38

Effect of the temperature. For the study of space clearance rate *a*, there was substantial

support for an effect of the interaction between the sampling status (NIS, IS-NR, IS-IR) and

temperature (ΔDIC > 12 in Table 2). In non-invasive species (NIS), *a* decreased as a function

of the temperature (posterior mean slope = -0.095, 95%CrI = [-0.136; -0.0526], pMCMC <

3×10^{-04}, Fig. 2C), whereas introduced populations appeared to react positively, but non

significantly, to temperature (Table S3). The same pattern was observed for the functional

response ratio *a/h* (ΔDIC > 2 in Table 2, Fig. S2D, Table S3). By contrast, the handling time *h*

(Table 2, Fig. S2B, Table S3) was unaffected by the temperature (ΔDIC > 3 in Table 2). The

analyses based on the dataset in biomass units led to consistent results (Table S2, Fig. S3B,D,F,

Table S4).

Table 2. *Model selection table for the effect of sampling status and experimental temperature on each*

functional response parameter (space clearance rate a, handling time h, and functional response ratio

FRR = a/h) expressed in standard units. The full model ('Model 3', see main text) and all the nested

models were ranked according to their DIC. The fixed variables considered were the experimental

temperature (T°), the sampling status (NIS, IS-NR, IS-IR), and their interaction term T°:status between

Temperature and sampling status. Degree of freedom (df) for each model is also reported.

Parameter	Nested model			df	DIC	ΔDIC
a	T°	Status	T°: Status	8	565.9	0
	T°	Status		6	578.8	12.89
		Status		5	579.3	13.39
				3	599.0	33.05
	T°			4	601.0	35.01
h		Status		5	1052.6	0
	T°	Status		6	1054.2	1.62
	T°	Status	T°: Status	8	1055.7	3.11
				3	1059.8	7.18
	T°			4	1061.7	9.04
FRR = *a/h*	T°	Status	T°:Status	8	1103.4	0
		Status		5	1105.6	2.18
	T°	Status		6	1107.8	4.38
				3	1124.0	20.64
	T°			4	1126.6	23.17

348
Figure 2: *The functional response parameters are expressed in standard units (log-transformed).* **A)**
Boxplot of space clearance rate as a function of the invasive status, non-invasive vs. invasive species.
B) *Boxplot of space clearance rate as a function of the sampling status, which was either non-invasive*
(NIS, in blue, all were sampled in their native range), invasive species sampled in their native range
(IS-NR, donor populations, in orange) or invasive species sampled in their invasive range (IS-IR,
introduced populations, in red). **C)** *Interaction between sampling status and temperature.* **D)** *Barplot*
of the phylogenetic signal for the various functional response parameters (space clearance rate a,
handling time h and functional response ratio FRR = a/h) expressed in standard or biomass units. These
signals were calculated using Lynch's phylogenetic heritability formula (Garamszegi, 2014). The
sample size (N) is indicated below each group when relevant. The sample size differed for the analyses
in biomass unit due to unreported values. Each data point corresponds to the report in a specific article
of one estimated value of clearance rate for one species. Multiple data points may correspond to the
same species studied across different articles. The mixed-effect model incorporates multiple imputations
*and accounts for phylogenetic inertia. "***" indicates pMCMC < 0.001, and NS pMCMC > 0.05.*

363
364

Phylogenetic signals. The phylogenetic signals were greater than 0.50 for all functional response parameters, with the strongest effect observed for the space clearance rate ($a = 0.93$, 95%CI = [0.89; 0.96] in standard units, $a = 0.82$, 95%CI = [0.72; 0.91] in biomass units, Fig. 2D). A phylogenetic signal of 0 indicates phylogenetic independence, whereas a signal of 1 indicates very strong variation in the parameter with phylogeny (Freckleton et al. 2002). These results underscore the importance of including phylogeny in comparative analysis of functional response.

DISCUSSION

The results presented here compiled over 269 functional responses from 45 freshwater fish species to test whether invasive species are stronger consumers than non-invasive species while accounting for the area where invasive species were tested, phylogeny and temperature. Overall, invasive species exhibited higher functional responses compared to non-invasive species, but surprisingly this difference was driven by the invasive species tested in their native range. These results question the paradigm that drives the comparison of functional response between invasive and non-invasive species and suggest new hypotheses regarding how the invasion process can make species traits evolve.

When invasive species were not differentiated based on whether they were sampled in their native (donor population, IS-NR) or invasive range (introduced population, IS-IR), it was found that they had overall higher functional responses than non-invasive species, with significantly higher space clearance rates and a tendency for lower handling times. The results remained valid when the functional responses were expressed in 'standard units' (per capita) or in 'biomass units', which accounted for the wide variability in body mass of prey and predator species across studies. These results are consistent with the prediction and underlying hypothesis of the comparative functional response approach, which states that invasive species

are better consumers than non-invasive species. This explains their success in becoming invasive and their impact on the invaded ecosystems. (see the Resource Consumption Hypothesis, Faria et al. 2025 ; Dick et al. 2014 ; Ricciardi et al. 2013). Until the present work, this hypothesis had been supported by empirical functional response studies in various taxa (Alexander et al. 2014; Barrios-O'Neill et al. 2014; Crookes et al. 2019; Cuthbert et al. 2019a; Howard et al. 2018 ; Laverty et al. 2015; Paterson et al. 2015; Mofu et al. 2019; Madzivanzira et al. 2021) but validation from more than one large-scale comparative approach was needed (Faria et al. 2025). However, when the geographic range in which the functional responses of invasive species were sampled was taken into account, it appeared that the general pattern of higher consumption rate in invasive species compared to non-invasive species was largely driven by studies based on donor populations (IS-NR).

Invasive species from their native range indeed exhibited significantly higher space clearance rates a and lower handling times h than non-invasive species, which is in line with the pre-invasion scenario in which some species would be predisposed to be invasive (Cadotte et al. 2018; Hulcr et al. 2017; Schlaepfer et al. 2010; Suarez & Tsutsui 2008). In the previous and unique comparative study also using the FoRAGE database, 309 functional responses involving freshwater and marine predatory fishes were compiled to test the role of size, activity level, temperature and invasive status as potential predictors of functional response parameters (Buba et al. 2022). Invasive species were not classified based on their study area, but it happened that the majority were investigated in their native range, allowing direct comparison with our study. Although their results were qualitatively consistent with ours, they only found a minimal connection between space clearance rates and invasive status. The reason could be that the authors did not account for phylogenetic inertia.

The high functional response of donor populations (IS-NR) suggests that comparative functional responses could be used to screen species for invasiveness with the aim of identifying

likely future invaders and preventing invasion (Hulcr et al. 2017; Van Kleunen et al. 2010).

Theory can help explain why species predisposed to invasiveness would not cause imbalance

to their community and remain unnoticed. When interspecific competition is greater than

intraspecific competition, superior competitors are predicted to exclude inferior competitors

(competitive exclusion, Bøhn et al. 2008; Gause 1934; Hardin 1960; Paine 1966; Tilman 1994).

According to optimal foraging theory, the most successful and therefore most frequently

encountered species should be the preferred resources of higher trophic-level natural enemies

such as predators and parasites (Ishii & Shimada 2012; Lacher et al. 1982; Pyke 1984). As a

result, natural enemies promote coexistence at the lower trophic level by decreasing the strength

of competitive exclusion (Chase et al. 2002; Chesson & Kuang 2008; Stump & Chesson 2017).

Competitors would exhibit their high potential once released from their natural enemies, as

occurs in the invasive range or in the experimental setting of functional response derivation.

Contrary to our prediction that prey novelty in the invasive range would benefit

introduced populations (IS-IR) through increased predation rates, their functional response ratio

was, on average, equivalent to that of non-invasive species (NIS) and lower than that of donor

population (IS-NR). This decrease was mainly driven by smaller space clearance rates and

could result from environmental constraints operating in the recipient ecosystem and leading to

trait convergence between introduced populations (IS-IR) and resident non-invasive species

(Craven et al. 2018). Comparative functional response studies involving donor (IS-NR) and

introduced (IS-IR) populations of the same invasive species, while also accounting for the time

elapsed since introduction, could help identify which mechanisms could explain trait

convergence in the recipient ecosystems. Indeed, environmental filtering (Gaston et al. 2009;

Keddy 1992), intra- and inter- generational phenotypic plasticity (within or between

individuals), or genetic changes (Marzluff 2012) could all be relevant factors. Whether invasive

species actually perform better in their introduced range remains controversial depending on

the taxa studied (Parker et al. 2013).

A limitation of our study is that we estimated global differences between IS-NR and IS-

IR, without direct comparison of donor (IS-NR) and introduced (IS-IR) populations within the

same species. Data are currently lacking to perform such Comparative Functional Response

approaches (CFR, Cuthbert et al. 2019b; Faria et al. 2023, 2025). Only three invasive species

have been studied in both their native and invasive ranges.

This result also challenges the general assumption of a positive relationship between

resource intake and performance (Brown et al. 2004; DeLong et al. 2021 ; Kooijman 2010), in

that metabolically efficient species might not be those with high feeding rates, and therefore

high functional responses, but those able to maintain viable metabolic rates while reducing the

amount of resources required. Such "thrifty" species would have more chances to persist in

ecosystems where ecological niches are occupied and become integrated into the local trophic

network. Despite the bunch of literature on the crucial role of physiology in shaping the

evolutionary trajectories and biogeographic distributions of non-invasive species (Somero

2002, 2005), there are to date few studies aimed at assessing the physiology of invasive species.

Compared to non-invasive species, they would be more eurytherms (Kelley 2014, but see Boher

et al. 2018), have higher mass-specific metabolic rates (Lagos et al. 2017), and be more efficient

in both energy conversion and food intake (Kooijman 2010; Taylor & Dunn 2018) with

ecomorphological advantages for feeding efficiency (Luger et al. 2020). Whether these findings

are congruent with our results needs further research on the link between metabolism and

functional response parameters.

Another important finding of our study concerns the negative consequences that climate

change could have on recipient communities. The relationship between space clearance rate

and temperature was indeed significantly negative in non-invasive species but tended to be

positive in invasive species tested in their introduced range. This means that a warmer

temperature could exacerbate the difference in functional response and likely the asymmetry in

competition between non-invasive (NIS) and invasive (IS-IR) species in recipient ecosystems.

Our result brings support to the literature suggesting that invasive species would often be less

vulnerable to warming because of predisposing physiological patterns, including energetic

performance and thermal tolerance (Bates et al. 2013; Kelley 2014; Komoroske et al. 2021;

Penk et al. 2019), their establishment being favored in certain ecosystemic contexts (Keller &

Shea 2021; Sorte et al. 2010; Stachowicz et al. 2002).

To conclude, our study calls into question the shortcut that invasive species have higher

functional responses than non-invasive species, without considering whether they were studied

in the native or introduced range. Ecological filters leading to trait convergence (at least from

the functional response perspective) between invasive populations and resident non-invasive

species seem to be at work in recipient communities and a promising avenue of research would

be to compare species traits, with a focus on metabolism, between donor and introduced

populations of the same invasive species but also between a recently introduced invasive

population with one that has been established for longer. From a broader perspective, this comes

down to asking how the invasion process shapes ecological niches through time (Ackerly 2003;

Broennimann et al. 2007; Mukherjee et al. 2012; Stiels et al. 2015; Wiens & Graham 2005).

Furthermore, the absence of differential functional response between invasive species in the

introduced range and non-invasive species raises concerns about the exclusive use of

comparative functional responses to predict the trophic impact of an invasive species.

Functional response should be used in combination with other metrics, such as species

abundance, to better predict impact (Dickey et al. 2020; Laverty et al. 2017). Our study also

reveals that global warming might exacerbate asymmetric competition between invasive and

non-invasive species as space clearance rate was found to decrease with temperature in non-

invasive species. We focused on freshwater fishes towards which functional response literature

is largely uneven and biased, probably because freshwater research offers ideal model species

and experimental conditions for functional response derivation. Extending our approach to

other taxa is needed to know to what extent our results reflect general patterns.

Acknowledgements. This work was supported by the Heatwaves project and the

BiodiverSâone project of the Agence de l'Eau Rhône-Méditerranée-Corse (AE-RMC).

498
499   **Competing interests.** No competing interests declared.
500

REFERENCES

Ackerly, D.D. (2003). Community Assembly, Niche Conservatism, and Adaptive Evolution in Changing Environments. International Journal of Plant Sciences, 164, S165–S184.

Albert J.S., Destouni G., Duke-Sylvester S.M., Magurran A.E., Oberdorff T., Reis R.E., et al. (2021) Scientists' warning to humanity on the freshwater biodiversity crisis. Ambio 50:85–94

Alexander, M.E., Dick, J.T.A., Weyl, O.L.F., Robinson, T.B. & Richardson, D.M. (2014). Existing and emerging high impact invasive species are characterized by higher functional responses than natives. Biol. Lett., 10, 20130946.

Barnett, A.G., Koper, N., Dobson, A.J., Schmiegelow, F. & Manseau, M. (2010). Using information criteria to select the correct variance-covariance structure for longitudinal data in ecology: Selecting the correct variance-covariance. Methods in Ecology and Evolution, 1, 15–24.

Barrios-O'Neill, D., Dick, J.T.A., Emmerson, M.C., Ricciardi, A., MacIsaac, H.J., Alexander, M.E., et al. (2014). Fortune favours the bold: a higher predator reduces the impact of a native but not an invasive intermediate predator. Journal of Animal Ecology, 83, 693–701.

Bates, A.E., McKelvie, C.M., Sorte, C.J.B., Morley, S.A., Jones, N.A.R., Mondon, J.A., et al. (2013). Geographical range, heat tolerance and invasion success in aquatic species. Proc. R. Soc. B., 280, 20131958.

Behrens J.W., von Friesen L.W.., Brodin T, Ericsson P., Hirsch P..E, Persson A., et al. (2020) Personality- and size-related metabolic performance in invasive round goby (*Neogobius melanostomus*). Physiology and Behavior 215:112777

Boher, F., Jaksic, F.M., Martel, S.I., Orellana, M.J. & Bozinovic, F. (2018). Does thermal physiology explain the ecological and evolutionary success of invasive species? Lessons from ladybird beetles. Evolutionary Ecology Research, 19.

Bøhn, T., Amundsen, P. A., & Sparrow, A. (2008). Competitive exclusion after invasion?. Biological Invasions, 10(3), 359-368.

Bollache, L., Dick, J.T.A., Farnsworth, K.D. & Montgomery, W.I. (2008). Comparison of the functional responses of invasive and native amphipods. Biol. Lett., 4, 166–169.

Broennimann, O., Treier, U.A., Müller-Schärer, H., Thuiller, W., Peterson, A.T. & Guisan, A. (2007). Evidence of climatic niche shift during biological invasion. Ecology Letters, 10, 701–709.

Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004). TOWARD A METABOLIC THEORY OF ECOLOGY. *Ecology*, 85, 1771–1789.

Buba, Y., DeLong, J.P. & Belmaker, J. (2022). Synthesizing drivers of fish functional responses across species. Fish and Fisheries, 23, 376–391.

Burnham, K.P. & Anderson, D.R. (Eds.). (2004). Model Selection and Multimodel Inference. Springer New York, New York, NY.

Cadotte, M.W., Campbell, S.E., Li, S., Sodhi, D.S. & Mandrak, N.E. (2018). Preadaptation and Naturalization of Nonnative Species: Darwin's Two Fundamental Insights into Species Invasion. *Annu. Rev. Plant Biol.*, 69, 661–684.

Chapple D.G., Simmonds S.M., Wong B.B.M. (2012) Can behavioral and personality traits influence the success of unintentional species introductions? Trends in Ecology and Evolution 27:57–64

Chase, J.M., Abrams, P.A., Grover, J.P., Diehl, S., Chesson, P., Holt, R.D., *et al.* (2002). The interaction between predation and competition: a review and synthesis. *Ecology Letters*, 5, 302–315.

Chesson, P. & Kuang, J.J. (2008). The interaction between predation and competition. *Nature*, 456, 235–238.

Christensen E.A.F., Norin T., Tabak I., van Deurs M., Behrens J.W. (2021) Effects of temperature on physiological performance and behavioral thermoregulation in an invasive fish, the round goby. Journal of Experimental Biology 224:jeb237669

563 Cinar, O., Nakagawa, S. & Viechtbauer, W. (2022). Phylogenetic multilevel meta-analysis: A simulation study on the
564 importance of modelling the phylogeny. Methods Ecol Evol, 13, 383–395.
565
566 Colautti, R. I., Ricciardi, A., Grigorovich, I. A., & MacIsaac, H. J. (2004). Is invasion success explained by the enemy
567 release hypothesis?. Ecology Letters, 7(8), 721-733.
568
569 Cox, J. G., & Lima, S. L. (2006). Naiveté and an aquatic–terrestrial dichotomy in the effects of introduced predators.
570 Trends in Ecology & Evolution, 21(12), 674-680.
571
572 Craven, D., Eisenhauer, N., Pearse, W.D., Hautier, Y., Isbell, F., Roscher, C., et al. (2018). Multiple facets of biodiversity
573 drive the diversity–stability relationship. Nat Ecol Evol, 2, 1579–1587.
574
575 Crookes, S., DeRoy, E.M., Dick, J.T.A. & MacIsaac, H.J. (2019). Comparative functional responses of introduced and
576 native ladybird beetles track ecological impact through predation and competition. Biol Invasions, 21, 519–529.
577
578 Cuthbert, R.N., Callaghan, A. & Dick, J.T.A. (2019a). A novel metric reveals biotic resistance potential and informs
579 predictions of invasion success. *Sci Rep*, 9, 15314.
580
581 Cuthbert, R. N., Dickey, J. W. E., Coughlan, N. E., Joyce, P. W. S. & Dick, J. T. A. (2019b). The functional response
582 ratio (FRR): advancing comparative metrics for predicting the ecological impacts of invasive alien species. Biological
583 Invasions 21, 2543–2547
584
585 Daugaard, U., Petchey, O. L., & Pennekamp, F. (2019). Warming can destabilize predator–prey interactions by shifting
586 the functional response from Type III to Type II. Journal of Animal Ecology, 88(10), 1575-1586.
587
588 DeLong, J.P. (2021). Predator ecology: Evolutionary ecology of the functional response. Oxford University Press.
589
590 DeLong, J.P., Uiterwaal, S.F. & Dell, A.I. (2021). Trait-Based Variation in the Foraging Performance of Individuals.
591 *Front. Ecol. Evol.*, 9, 649542.
592
593 Dick, J.T.A., Alexander, M.E., Jeschke, J.M., Ricciardi, A., MacIsaac, H.J., Robinson, T.B., et al. (2014). Advancing
594 impact prediction and hypothesis testing in invasion ecology using a comparative functional response approach. Biol
595 Invasions, 16, 735–753.
596
597 Dick, J.T.A., Gallagher, K., Avlijas, S., Clarke, H.C., Lewis, S.E., Leung, S., et al. (2013). Ecological impacts of an
598 invasive predator explained and predicted by comparative functional responses. Biol Invasions, 15, 837–846.
599
600 Dickey, J.W.E., Cuthbert, R.N., South, J., Britton, J.R., Caffrey, J., Chang, X., et al. (2020). On the RIP: using Relative
601 Impact Potential to assess the ecological impacts of invasive alien species. NeoBiota 55, 27–60.
602
603 Dlugosch K.M., Parker I.M. (2008). Founding events in species invasions: genetic variation, adaptive evolution, and the
604 role of multiple introductions. Molecular Ecology 17:431–49
605
606 Dlugosch K.M., Anderson S.R., Braasch J., Cang F.A., Gillette H. (2015). The devil is in the details: genetic variation
607 in introduced populations and its contributions to invasion. Mol. Ecol. 24:2095–111
608
609 Ens, N.J., Lim, E.G., Howard, B.R. & Eastham, T.M. (2021). A comparison of the predatory impacts of an invasive and
610 native crab species using a functional response approach. Biological Invasions, 23, 2329–2336.
611
612 Faria, L., Cuthbert, R. N., Dickey, J. W. E., Jeschke, J. M., Ricciardi, A., Dick, J. T. A. & Vitule, J. R. S. (2023). The
613 rise of the functional response in invasion science: a systematic review. NeoBiota 85, 43–79
614
615 Faria, L., Cuthbert, R.N., Dickey, J.W.E., Jeschke, J.M., Ricciardi, A., Dick, J.T.A., et al. (2025). Non-native species
616 have higher consumption rates than their native counterparts. Biological Reviews, 100, 1163-1180
617
618 Freckleton, R.P., Harvey, P.H. & Pagel, M. (2002). Phylogenetic analysis and comparative data: A test and review of
619 evidence. *The American Naturalist*, 160, 712–726.
620
621 Funk, J.L. & Vitousek, P.M. (2007). Resource-use efficiency and plant invasion in low-resource systems. Nature, 446,
622 1079–1081.
623
624 Galarowicz, T.L. & Wahl, D.H. (2005). Foraging by a young-of-the-year piscivore: the role of predator size, prey type,
625 and density. Can. J. Fish. Aquat. Sci., 62, 2330–2342.
626

627 Garamszegi, L.Z. (2014). Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology.
628 Concepts and Practice.
629
630 Gaston, K.J. (2009). Geographic range limits of species. Proc. R. Soc. B., 276, 1391–1393.
631
632 Gause, G. F. (1934). The struggle for existence.
633
634 Guisan, A., Petitpierre, B., Broennimann, O., Daehler, C., & Kueffer, C. (2014). Unifying niche shift studies: insights
635 from biological invasions. Trends in Ecology & Evolution, 29(5), 260-269.
636
637 Hadfield, J.D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R
638 Package. J. Stat. Soft., 33.
639
640 Hadfield, J.D. & Nakagawa, S. (2010). General quantitative genetic methods for comparative biology: phylogenies,
641 taxonomies and multi-trait models for continuous and categorical characters. Journal of Evolutionary Biology, 23, 494–
642 508.
643
644 Hardin, G. (1960). The Competitive Exclusion Principle: An idea that took a century to be born has implications in
645 ecology, economics, and genetics. *Science*, 131, 1292–1297.
646
647 Hassell, M.P. & May, R.M. (1973). Stability in Insect Host-Parasite Models. The Journal of Animal Ecology, 42, 693.
648
649 Heger, T. & Jeschke, J.M. (2014). The enemy release hypothesis as a hierarchy of hypotheses. Oikos, 123, 741–750.
650
651 Holling, C.S. (1959). Some characteristics of simple types of predation and parasitism. Can Entomol, 91, 385–398.
652
653 Howard, B., Wong, D., Aguiar, V., Desforges, J., Oishi, E., Stewart, J., et al. (2022). Effects of perceived competition
654 and water temperature on the functional responses of invasive and native crabs. Mar. Ecol. Prog. Ser., 684, 69–78.
655
656 Howard, B.R., Barrios-O'Neill, D., Alexander, M.E., Dick, J.T.A., Therriault, T.W., Robinson, T.B., et al. (2018).
657 Functional responses of a cosmopolitan invader demonstrate intraspecific variability in consumer-resource dynamics.
658 PeerJ, 6, e5634.
659
660 Hulcr, J., Black, A., Prior, K., Chen, C. Y., & Li, H. F. (2017). Studies of ambrosia beetles (Coleoptera: Curculionidae)
661 in their native ranges help predict invasion impact. Florida Entomologist, 100(2), 257-261.
662
663 Ishii, Y., & Shimada, M. (2012). Learning predator promotes coexistence of prey species in host–parasitoid systems.
664 *Proceedings of the National Academy of Sciences*, *109*(13), 5116-5120.
665
666 Jeschke, J.M., Kopp, M. & Tollrian, R. (2002). Predator functional responses: discriminating between handling and
667 digesting prey. Ecological Monographs, 72, 95–112.
668
669 Jolles J.W., King A.J., Killen S.S. (2020). The role of individual heterogeneity in collective animal behaviour. Trends
670 in Ecology and Evolution 35:278–291
671
672 Kattler, K.R., Oishi, E.M., Lim, E.G., Watkins, H.V. & Côté, I.M. (2023). Functional responses of male and female
673 European green crabs suggest potential sex-specific impacts of invasion. PeerJ, 11, e15424.
674
675 Keddy, P.A. (1992). Assembly and response rules: two goals for predictive community ecology. J Vegetation Science,
676 3, 157–164.
677
678 Keller, J.A. & Shea, K. (2021). Warming and shifting phenology accelerate an invasive plant life cycle. Ecology, 102,
679 e03219.
680
681 Kelley, A.L. (2014). The role thermal physiology plays in species invasion. Conservation Physiology, 2, cou045–
682 cou045.
683
684 Komoroske, L.M., Jeffries, K.M., Whitehead, A., Roach, J.L., Britton, M., Connon, R.E., et al. (2021). Transcriptional
685 flexibility during thermal challenge corresponds with expanded thermal tolerance in an invasive compared to native fish.
686 Evolutionary Applications, 14, 931–949.
687
688 Kooijman, S.A.L.M. (2010). *Dynamic Energy and Mass Budgets in Biological Systems*. 2nd edn. Cambridge University
689 Press.
690

Lacher, T.E., Willig, M.R. & Mares, M.A. (1982). Food Preference as a Function of Resource Abundance with Multiple Prey Types: An Experimental Analysis of Optimal Foraging Theory. *The American Naturalist*, 120, 297–316.

Lagos, M.E., White, C.R. & Marshall, D.J. (2017). Do invasive species live faster? Mass-specific metabolic rate depends on growth form and invasion status. Functional Ecology, 31, 2080–2086.

Lajeunesse, M.J., Rosenberg, M.S. & Jennions, M.D. (2013). 17. Phylogenetic Nonindependence and Meta-analysis. In: Handbook of Meta-analysis in Ecology and Evolution. Princeton University Press, pp. 284–299.

Landi, P., McCoy, M.W. & Vonesh, J.R. (2022). Predicting invasive consumer impact via the comparative functional response approach: linking application to ecological theory. Biol Invasions, 24, 3565–3579.

Laverty, C., Dick, J.T.A., Alexander, M.E. & Lucy, F.E. (2015). Differential ecological impacts of invader and native predatory freshwater amphipods under environmental change are revealed by comparative functional responses. Biol Invasions, 17, 1761–1770.

Laverty, C., Green, K.D., Dick, J.T.A., Barrios-O'Neill, D., Mensink, P.J., Médoc, V., et al. (2017). Assessing the ecological impacts of invasive species based on their functional responses and abundances. Biol Invasions, 19, 1653–1665.

Luger, A.M., South, J., Alexander, M.E., Ellender, B.R., Weyl, O.L.F. & Nagelkerke, L.A.J. (2020). Ecomorphology of largemouth bass relative to a native trophic analogue explains its high invasive impact. Biol Invasions, 22, 2223–2233. Madzivanzira, T.C., South, J. & Weyl, O.L.F. (2021). Invasive crayfish outperform Potamonautid crabs at higher temperatures. *Freshwater Biology*, 66, 978–991.

Marzluff, J.M. (2012). 18. Urban Evolutionary Ecology. In: Urban Bird Ecology and Conservation. University of California Press, pp. 287–308.

Meijer, K., Schilthuizen, M., Beukeboom, L., & Smit, C. (2016). A review and meta-analysis of the enemy release hypothesis in plant–herbivorous insect systems. PeerJ, 4, e2778.

Mofu, L., South, J., Wasserman, R.J., Dalu, T., Woodford, D.J., Dick, J.T.A., et al. (2019). Inter-specific differences in invader and native fish functional responses illustrate neutral effects on prey but superior invader competitive ability. Freshwater Biology, 64, 1655–1663.

Mollot, G., Pantel, J.H. & Romanuk, T.N. (2017). The Effects of Invasive Species on the Decline in Species Richness. In: Advances in Ecological Research. Elsevier, pp. 61–83.

Mukherjee, A., Williams, D.A., Wheeler, G.S., Cuda, J.P., Pal, S. & Overholt, W.A. (2012). Brazilian peppertree (Schinus terebinthifolius) in Florida and South America: evidence of a possible niche shift driven by hybridization. Biol Invasions, 14, 1415–1430.

Murray, G.P.D., Stillman, R.A., Gozlan, R.E. & Britton, J.R. (2013). Experimental Predictions of The Functional Response of A Freshwater Fish. Ethology, 119, 751–761.

Nakagawa, S. & Freckleton, R.P. (2011). Model averaging, missing data and multiple imputation: a case study for behavioural ecology. Behav Ecol Sociobiol, 65, 103–116.

Nakamura, G., Richter, A. & Soares, B.E. (2021). FishPhyloMaker: An R package to generate phylogenies for ray-finned fishes. Ecological Informatics, 66, 101481.

Oyugi, D.O., Cucherousset, J., Baker, D.J. & Britton, J.R. (2012). Effects of temperature on the foraging and growth rate of juvenile common carp, Cyprinus carpio. Journal of Thermal Biology, 37, 89–94.

Pagad, S., Genovesi, P., Carnevali, L., Schigel, D. & McGeoch, M.A. (2018). Introducing the Global Register of Introduced and Invasive Species. Sci Data, 5, 170202.

Paine, R.T. (1966). Food Web Complexity and Species Diversity. *The American Naturalist*, 100, 65–75.

Parker, J.D., Torchin, M.E., Hufbauer, R.A., Lemoine, N.P., Alba, C., Blumenthal, D.M., et al. (2013). Do invasive species perform better in their new ranges? Ecology, 94, 985–994.

Paterson, R.A., Dick, J.T.A., Pritchard, D.W., Ennis, M., Hatcher, M.J. & Dunn, A.M. (2015). Predicting invasive species impacts: a community module functional response approach reveals context dependencies. Journal of Animal Ecology, 84, 453–463.

Penk, M.R. & Williams, M.A. (2019). Thermal effluents from power plants boost performance of the invasive clam Corbicula fluminea in Ireland's largest river. Science of The Total Environment, 693, 133546.

Pyke, G.H. (1984). Optimal Foraging Theory: A Critical Review. *Annu. Rev. Ecol. Syst.*, 15, 523–575.

R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. (2025).

Razanajatovo, M., Maurel, N., Dawson, W., Essl, F., Kreft, H., Pergl, J., et al. (2016). Plants capable of selfing are more likely to become naturalized. Nature Communications, 7(1), 13313.

Reid, A.J., Carlson, A.K., Creed, I.F., Eliason, E.J., Gell, P.A., Johnson, P.T.J., et al. (2019). Emerging threats and persistent conservation challenges for freshwater biodiversity. Biological Reviews, 94, 849–873.

Ricciardi, A., Hoopes, M. F., Marchetti, M. P., & Lockwood, J. L. (2013). Progress toward understanding the ecological impacts of nonnative species. Ecological Monographs, 83(3), 263-282.

Sayer, C.A., Fernando, E., Jimenez, R.R., Macfarlane, N.B.W., Rapacciuolo, G., Böhm, M., *et al.* (2025). One-quarter of freshwater fauna threatened with extinction. *Nature*, 638, 138–145.

Schlaepfer, D.R., Glättli, M., Fischer, M. & Van Kleunen, M. (2010). A multi-species experiment in their native range indicates pre-adaptation of invasive alien plant species. *New Phytologist*, 185, 1087–1099.

Sih, A., Bolnick, D. I., Luttbeg, B., Orrock, J. L., Peacor, S. D., Pintor et al. (2010). Predator–prey naïveté, antipredator behavior, and the ecology of predator invasions. *Oikos*, 119(4), 610-621.

Simberloff, D. (2009). The role of propagule pressure in biological invasions. *Annual review of ecology, evolution, and systematics*, 40(1), 81-102.

Somero, G.N. (2002). Thermal Physiology and Vertical Zonation of Intertidal Animals: Optima, Limits, and Costs of Living. Integrative and Comparative Biology, 42, 780–789.

Somero, G.N. (2005). Linking biogeography to physiology: Evolutionary and acclimatory adjustments of thermal limits. Front Zool, 2, 1.

Sorte, C.J.B., Williams, S.L. & Zerebecki, R.A. (2010). Ocean warming increases threat of invasive species in a marine fouling community. Ecology, 91, 2198–2204.

Spiegelhalter, D.J., Best, N.G., Carlin, B.P. & Van Der Linde, A. (2002). Bayesian measures of model complexity and fit. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 64, 583–639.

Stachowicz, J.J., Terwin, J.R., Whitlatch, R.B. & Osman, R.W. (2002). Linking climate change and biological invasions: Ocean warming facilitates nonindigenous species invasions. Proc. Natl. Acad. Sci. U.S.A., 99, 15497–15500.

Stephens, D.W. & Krebs, J.R. (1986). *Foraging theory*. Monographs in behavior and ecology. Princeton University Press, Princeton, New Jersey.

Stiels, D., Gaißer, B., Schidelko, K., Engler, J.O. & Rödder, D. (2015). Niche shift in four non-native estrildid finches and implications for species distribution models. Ibis, 157, 75–90.

Stump, S.M. & Chesson, P. (2017). How optimally foraging predators promote prey coexistence in a variable environment. *Theoretical Population Biology*, 114, 40–58.

Suarez, A.V. & Tsutsui, N.D. (2008). The evolutionary consequences of biological invasions. *Molecular Ecology*, 17, 351–360.

Tilman, D. (1994). Competition and Biodiversity in Spatially Structured Habitats. *Ecology*, 75, 2–16.

Taylor, N.G. & Dunn, A.M. (2018). Predatory impacts of alien decapod Crustacea are predicted by functional responses and explained by differences in metabolic rate. Biol Invasions, 20, 2821–2837.

817
818 Uiterwaal, S.F., Lagerstrom, I.T., Lyon, S.R. & DeLong, J.P. (2018). Data paper: FoRAGE (Functional Responses from
819 Around the Globe in all Ecosystems) database: a compilation of functional responses for consumers and parasitoids.
820
821 Uiterwaal, S.F., Lagerstrom, I.T., Lyon, S.R. & DeLong, J.P. (2022). FoRAGE database: A compilation of functional
822 responses for consumers and parasitoids. Ecology, 103, e3706.
823
824 Uszko, W., Diehl, S., Pitsch, N., Lengfellner, K., & Müller, T. (2015). When is a type III functional response stabilizing?
825 Theory and practice of predicting plankton dynamics under enrichment. Ecology, 96(12), 3243–3256.
826
827 Van Kleunen, M., Dawson, W., Schlaepfer, D., Jeschke, J.M. & Fischer, M. (2010). Are invaders different? A conceptual
828 framework of comparative approaches for assessing determinants of invasiveness. *Ecology Letters*, 13, 947–958.
829
830 Wiens, J.J. & Graham, C.H. (2005). Niche Conservatism: Integrating Evolution, Ecology, and Conservation Biology.
831 Annu. Rev. Ecol. Evol. Syst., 36, 519–539.
832
833 Xu, M., Mu, X., Dick, J.T.A., Fang, M., Gu, D., Luo, D., *et al.* (2016). Comparative Functional Responses Predict the
834 Invasiveness and Ecological Impacts of Alien Herbivorous Snails. *PLoS ONE*, 11, e0147017.
835
836

837 **SUPPLEMENTARY MATERIAL**

839
840 ***Figure S1***. *PRISMA flowchart showing the number of observations (O, i.e. number of estimated*
841 *functional response curves reporting the space clearance rate* a *and the handling time* h*), the number*
842 *of articles (A), and the number of freshwater fish species as predator (S).*

844
Figure S2. *The functional response parameters are expressed in standard units (log-transformed).* **A)**
Boxplot of handling time (h) as a function of the sampling status, which was either non-invasive (NIS,
in blue, all were sampled in their native range), invasive species sampled in their native range (IS-NR,
donor populations, in orange) or invasive species sampled in their invasive range (IS-IR, introduced
populations, in red). **B)** *Effect of the interaction between sampling status and temperature on the*
handling time. **C)** *Boxplot of functional response ratio (FRR = a/h) as a function of the sampling status.*
D) *Effect of the interaction between sampling status and temperature on the functional response ratio*
(FRR = a/h). The sample size (N) is indicated below each group when relevant. Each data point
corresponds to the report in a specific article of one estimated value of clearance rate for one species.
Multiple data points may correspond to the same species studied across different articles. The mixed-
*effect model incorporates multiple imputations and accounts for phylogenetic inertia. "***" indicates*
*pMCMC < 0.001, "**" pMCMC < 0.01 and NS pMCMC > 0.05.*

Figure S3. *The functional response parameters are expressed in biomass units (log-transformed).* **A)** *Boxplot of space clearance rate as a function of the sampling status, which was either non-invasive (NIS, in blue, all were sampled in their native range), invasive species sampled in their native range (IS-NR, donor populations, in orange) or invasive species sampled in their invasive range (IS-IR, introduced populations, in red).* **B)** *Effect of the interaction between sampling status and temperature on the space clearance rate (a).* **C)** *Boxplot of handling time (h) as a function of the sampling status.* **D)** *Effect of the interaction between sampling status and temperature on the handling time (h).* **E)** *Boxplot of functional response ratio (FRR = a/h) as a function of the sampling status.* **F)** *Effect of the interaction between sampling status and temperature on the functional response ratio (FRR = a/h). The sample size (N) is indicated below each group when relevant. Each data point corresponds to the report in a specific article of one estimated value of clearance rate for one species. Multiple data points may correspond to the same species studied across different articles. The mixed-effect model incorporates multiple imputations and accounts for phylogenetic inertia. "***" indicates pMCMC < 0.001, "**" pMCMC < 0.01, "*" pMCMC < 0.05, and NS pMCMC > 0.05.*

872 ***Table S1.*** *Model selection table for the influence of sampling status on each functional response*
873 *parameter (space clearance rate a, handling time h, and functional response ratio FRR = a/h) expressed*
874 *in biomass units. The competitive models were ranked according to their DIC. For a detailed description*
875 *of the three models, refer to the main text.*

Model	Clearance rate a		Handling time h		FRR = a/h	
	DIC	ΔDIC	DIC	ΔDIC	DIC	ΔDIC
Model 2: sampling status	627.61	0	1035.15	0	1103.01	0
Model 1: invasive status	656.45	28.84	1035.88	0.73	1113.16	10.15
Model 0: intercept	654.51	26.90	1036.99	1.84	1114.46	11.45

Table S2. *Model selection table for the effect of sampling status and experimental temperature on each functional response parameter (space clearance rate a, handling time h, and functional response ratio FRR = a/h) expressed in biomass units. The full model ('Model 3', see main text) and all the nested models were ranked according to their DIC. The fixed variables considered were the experimental temperature (T°), the sampling status (NIS, IS-NR, IS-IR), and their interaction term T°:status. Degree of freedom (df) for each model is also reported.*

Parameter	Nested model			df	DIC	ΔDIC
a	T°	Status	T°:Status	8	609.7	0
		Status		5	627.5	17.85
	T°	Status		6	629.3	19.64
				3	654.6	44.94
	T°			4	656.1	46.47
h		Status		5	1035.2	0
				3	1037.0	1.83
	T°	Status		6	1038.0	2.81
	T°			4	1040.2	5.06
	T°	Status	T°:Status	8	1040.9	5.72
FRR *a/h*	T°	Status	T°:Status	8	1100.8	0
		Status		5	1103.0	2.21
	T°	Status		6	1104.9	4.60
				3	1114.4	13.58
		T°		4	1116.7	15.87

Table S3. *Effect of the experimental temperature for each functional response parameter (space clearance rate a, handling time h, and functional response ratio FRR = a/h) expressed in standard units, and for each sampling status (NIS, IS-NR, IS-IR). The posterior mean of the slope with 95% credible interval, and pMCMC are reported. "***" indicates pMCMC < 0.001, "**" pMCMC < 0.01, "*" pMCMC < 0.05.*

Parameter	Sampling status	Posterior mean	95% CrI	pMCMC
a	NIS	-0.095	[-0.14, -0.052]	$< 3 \times 10^{-04}$ ***
	IS-NR	0.0078	[-0.046, 0.056]	0.767
	IS-IR	0.056	[-0.0073, 0.11]	0.072
h	NIS	0.054	[-0.019, 0.13]	0.15
	IS-NR	-0.044	[-0.19, 0.087]	0.52
	IS-IR	0.020	[-0.11, 0.16]	0.76
FRR *a/h*	NIS	-0.13	[-0.11, -0.011]	0.014 *
	IS-NR	0.069	[-0.062, 0.20]	0.32
	IS-IR	0.025	[-0.12, 0.18]	0.75

Table S4. *Effect of the experimental temperature for each functional response parameter (space clearance rate a, handling time h, and functional response ratio FRR = a/h) expressed in biomass units, and for each sampling status (NIS, IS-NR, IS-IR). The posterior mean of the slope with 95% credible interval, and pMCMC are reported. "*" pMCMC < 0.05.*

Parameter	Sampling status	Posterior mean	95% CrI	pMCMC
a	NIS	-0.060	[-0.11, -0.011]	0.017 *
	IS-NR	0.025	[-0.029, 0.075]	0.34
	IS-IR	0.056	[-0.0034, 0.12]	0.072
h	NIS	0.018	[-0.061, 0.090]	0.61
	IS-NR	-0.010	[-0.17, 0.14]	0.96
	IS-IR	0.024	[-0.087, 0.14]	0.68
FRR *a/h*	NIS	-0.074	[-0.19, 0.035]	0.20
	IS-NR	0.067	[-0.12, 0.24]	0.51
	IS-IR	0.030	[-0.11, 0.19]	0.69