Membrane rupture and independent extension of sister membranes drive

cytokinesis in *C. elegans* embryos

Jingjing Liang[1]*, Tingrui Huang[1,2,3], Xun Huang[1,2], and Mei Ding[1,2]*

[1] Institute of Genetics and Developmental Biology, Chinese Academy of Sciences,

Beijing 100101, China

[2] Taikang Center for Life and Medical Sciences, School of Basic Medical Sciences,

Wuhan University, Wuhan 430071, China

[3] University of Chinese Academy of Sciences, Beijing 100049, China

* Correspondence: jjliang@genetics.ac.cn; mding@genetics.ac.cn

Abstract

Cell division, the physical separation of a cell into two daughter cells, is classically thought to require continuous plasma membrane invagination driven by equatorial contractile forces. Here, we challenge this paradigm by uncovering an unexpected mechanism in *C. elegans* embryos: the plasma membrane ruptures as early as metaphase. Instead of invaginating as a continuous sheet, the separated membranes extend independently to partition the cytoplasm. The membrane rupture and independent extension of sister membranes create membrane discontinuities—widely observed in electron microscopy (EM) sections. Without plasma membrane coverage, the exposed cytoplasmic regions are shielded either by plasma membrane from neighboring, likely kinship, cells or extracellular matrix components, preventing massive leakage. Our findings overturn the long-standing assumption that cytokinesis requires persistent membrane integrity, and offer fresh insights into cell division, particularly in fast dividing embryonic cells.

Introduction

The plasma membrane integrity is the corner stone of cell biology. Based upon, , current models further propose that cell division should also occur with uninterrupted membrane integrity and the invagination of this intact membrane, driven by a contractile ring of actin filaments, myosin II, and regulatory proteins (Pollard and O'Shaughnessy, 2019), with vesicle fusion supplying new membrane to accommodate shape change (Gerien and Wu, 2018), progressively deepens until pinching the cell into two daughter cells (Figure 1A). Critically, the foundational premise of uninterrupted membrane integrity during cytokinesis has never been rigorously tested until our unexpected discovery in *C. elegans* embryos.

After fertilization, the single cell *C. elegans* embryo begins a series of highly stereotyped cell divisions. During this phase, cells undergo rapid division (15-30 minutes per cycle), with each cycle alternating between S (DNA synthesis) and M (mitosis and cytokinesis) phases while bypassing G1and G2 gap phases until the 28-cell stage (Edgar and McGhee, 1988; Deppe et al., 1978; Sulston et al., 1983; Bao et al., 2008; Pintard and Bowerman, 2019). Around this stage, the worm embryos are laid by the mother. Subsequent development continues outside the mother, encapsulated within a protective, multi-layer extracellular matrix known as the eggshell (Stein, 2018). This impermeable eggshell is established earlier, by the end of anaphase II of meiosis, and undergoes its final modifications during the embryo's first few mitotic divisions (Bembenek et al., 2007; Zhang et al., 2005).

During routine ultrastructural analysis (Zhu et al., 2022), we found that rather than maintaining membrane continuity, the dividing *C. elegans* embryonic cells routinely exhibit plasma membrane discontinuities. Further analysis revealed that the

plasma membrane of the embryonic cells ruptures as early as metaphase and sister

membrane primarily extend independent from each other to achieve cytoplasmic

partition (Figure 1B), a mechanism never before documented. This discovery

challenges the canonical view of cytokinesis at its core, revealing an unanticipated

plasticity in cell division mechanisms and prompting a re-evaluation of how

membrane dynamics are regulated across species.

Results

Widespread membrane discontinuities in dividing *C. elegans* embryonic cells

Our study was initially prompted by the curious observation that the plasma

membrane of embryonic cells in *C. elegans* does not appear to be continuously intact.

We refer to these membrane discontinuities as membrane openings (MOs). They are

particularly evident at the embryo's periphery. As shown in Figure 1C, the plasma

membrane is absent in the region marked by the white dashed line. Membrane

interruptions are also be observed in the embryo's central regions. For instance,

Figure 1D shows that Cell 3 contains a MO facing both Cell 1 and Cell 2.

Additionally, we noted instances where a single plasma membrane (SPM) appears to

traverse the cytoplasm, seemingly partitioning it into separate compartments across

adjacent cells (Figure 1D) (yellow arrows). According to the conventional view, the

plasma membrane should remain intact throughout the cell cycle—whether during

division or interphase—and each cell should be enclosed by its own membrane.

Consequently, between two adjacent cells, we would expect to observe two distinct

plasma membranes. The presence of a single continuous membrane, however,

suggests the existence of regions lacking a complete boundary. For instance, Cell 4 in

Figure 1D clearly contains such membrane discontinuities. Collectively, these

findings challenge the established notion that the plasma membrane remains intact

over the entire lifetime of a cell.

To rule out potential artifacts inherent to electron microscopy (EM) techniques,

we performed a series of control experiments. First, we prepared four independent

sample batches. Membrane discontinuities were consistently observed in all batches,

indicating that the phenomenon is not specific to a single preparation. Second, we

examined a publicly available historical collection of *C. elegans* embryo EM images

(Wormatlas) and identified MOs in several instances, particularly at the embryonic

periphery. This confirmation in an independent dataset suggests that MOs are genuine

cellular structures rather than procedural artifacts. Third, we analyzed embryos across

various developmental stages. MO-containing cells were observed in both intrauterine

and already laid embryos at early stages (Figure S1). However, as embryos reached

the comma stage (Figure S2)—a period characterized by the onset of elongation and a

decline in active cell proliferation—the incidence of MOs dropped dramatically

(observed in 0/13, 0/17, 0/30, and 0/30 cells examined, respectively). These results

demonstrate that plasma membrane discontinuities are associated with specific stages

of *C. elegans* embryogenesis and are not a product of random membrane tearing.

The early stage of *C. elegans* embryonic cells is under rapid and extensive cell

division. Within approximately 100 minutes, a two-cell embryo develops into an

embryo containing nearly 30 cells (Bao et al., 2008). Given this dynamic

proliferation, we suspected that the formation of membrane discontinuities might be

associated with cell division events. Therefore, to systematically examine the

formation of membrane discontinuities, we collected EM images of 60 early embryos

from 9 gravid adults. By examining the entire circumference of 425 cells in single

high-pressure freezing transmission electron microscopy (HPF-TEM) sections (Figure

S2) (Modern Electron Microscopy Methods for C. elegans, 2012), we found that MOs

in 159 cells distributed across 57 embryos, representing 95% of the total sampled

embryos (Figure 1E). Among these 159 affected cells, 89% (141/159) exhibited a

single membrane discontinuity, while 13 cells had two openings and 5 cells had three.

We analyzed the 159 cells exhibiting MOs based on nuclear and cytoplasmic

morphology (Figure 1E). Of these, 64 cells appeared in EM sections containing only

cytoplasmic regions—9 of which completely lacked discernible plasma membranes.

The remaining 95 cells contained one or more nuclei or condensed chromosomes.

Among them, 10 cells were clearly in metaphase stage (defined by a single

chromosomal cluster). As shown in Figure 2A, a representative metaphase cell (MC)

displayed a MO (dashed white line) spanning roughly one-third of the cell's

circumference. High-magnification imaging revealed that in contrast to the wrinkled

plasma membranes (Figure 2A, enlarged boxed area displayed in 1), the large MO

exhibited a smoother surface morphology, and the internal cytoplasm remained

enclosed within the cell (Figure 2A, enlarged boxed areas displayed in 2 and 3). MOs

were also observed in anaphase cells (AC) (n=5; characterized by <10 chromosomal

clumps; Figure 2B, enlarged in 2) and telophase cells (n=5; defined by two nuclei

sharing cytoplasm). Intriguingly, one telophase-like cell contained three nuclei within

a common cytoplasm (Figure S2, marked by &).

Because EM sectioning captures cells at random stages of the cell cycle, we

sought to systematically assess the relationship between mitosis and MOs. To do so,

we analyzed 20 cells with unambiguous division features, selected based on nuclear

morphology—such as aligned chromosomes (metaphase) or discrete clusters

(anaphase)—and cytoplasmic sharing, indicated by two or more nuclei within a

common cytoplasm (telophase). Strikingly, 19 out of 20 cells (95%) revealed MOs

upon tracing the plasma membrane in single EM sections (Figure S3). This near-

ubiquitous association strongly suggests that membrane discontinuities are an integral

feature of cell division in early *C. elegans* embryos.

Membrane rupture in cells at metaphase

The observed membrane discontinuities suggest that the plasma membrane of early *C.*

elegans embryos may rupture during cell division. However, because TEM provides

only static, two-dimensional images, we could not determine the three-dimensional

nature of these ruptures. To better understand this phenomenon, we acquired

continuous serial sections and performed 3D reconstruction.

Since membrane discontinuities appear at metaphase—well before nuclear or

cytoplasmic division—we performed continuous EM sectioning to trace the entire

plasma membrane of a metaphase cell (Figures 2C). Through 3D reconstruction, we

identified a continuous plasma membrane rupture extending over 8,940 nm, which is

more than two-thirds of the cell's circumference (Figure 2D). Notably, one end of the

membrane rupture appeared long and narrow, while the other was wide and exhibited

cytoplasmic leakage (Figure 2D and Video S1). Further analysis revealed that the

leaked cytoplasm remained confined to a relatively localized region (Video S1).

During metaphase, chromosomes align at the equatorial plane, a key region for

subsequent segregation. To explore the spatial relationship between membrane

rupture and chromosomal organization, we further reconstructed the entire condensed

chromosome region (Figures 2C). We found that this large-scale membrane rupture

largely encircled the chromosomal disc (Figure 2D and Video S1), spatially aligning

with the future segregation zone.

The detachment of sister membranes

According to the conventional model, cytokinesis in animal cells begins with the

formation of a membrane invagination. As this invagination deepens, it gives rise to

two back-to-back membrane layers, ultimately leading to the physical separation of

the two daughter cells. In the context of membrane rupture, however, plasma

membrane disruption results in the separation of two putative sister membranes. We

then asked how the subsequent cytokinesis process proceeds with detached sister

membranes. To address this question, we examined 51 dividing cells. In each of these

cells, two well-separated nuclei were observed, indicating that they were at telophase.

We found that while a minority of cells (6/51) exhibited classical membrane

invagination (Figure 3A), including one embryo transitioning from the one-cell to the

two-cell stage, the vast majority (45/51) displayed a striking detached membrane

configuration (Figure 3B). In these cases, sister membrane tips were completely

separated from each other (Figures 3B and 3C, marked with *), forming distinct, non-

contiguous structures that defy conventional models of cytokinesis. This predominant

pattern suggests that membrane detachment—rather than continuous invagination—

may represent the primary mode of cytoplasmic separation in *C. elegans* embryos. Of

note, the membrane rupture simultaneously generates individual sister membranes

with severed edges. Indeed, as shown in Figure 3B, the tips of both free-ended edges

(*) of Cell 1 are visible in the EM section.

In the detached membrane configuration, cytoplasmic components—such as

glycogen granules and ribosomal particles—were clearly visible surrounding the

severed edges of the two sister membranes (Figure 3C). This observation is

particularly significant because the classical invagination model cannot generate

single plasma membrane barriers between adjacent cytoplasmic domains. Instead,

independent extension of severed sister membranes could explain the formation of

cytoplasm-immersed membranes. To further test this hypothesis, we traced 12

cytoplasm-embedded membranes, including those separating Cell 1 from Cells 2 and

3, and those separating Cell 4 from 3 (displayed on Figure 3D, enlarged in 1 and 2). In

all cases, the cytoplasmic regions flanking these membranes belonged to cytosolically

interconnected sister cells. Together, these results demonstrate that *C. elegans*

embryonic cytokinesis relies on detached sister membrane extension rather than

invagination mediated by intact membranes.

Tracing from the severed edges back to their origins, we found that the sister

membranes were frequently anchored side-by-side near the cytosol-immersed region

(Figures 4A-4D). Within this region, gap junction plaques were often observed (57 in

94 cases) at membrane contact sites (Figure 4B), though their potential role in sister

membrane cohesion is unclear. These contact sites alternated with regions where the

two sister membranes separated, revealing lightly stained intercellular spaces in EM

images. Beyond the anchored region, the separation between sister membranes

became evident and each membrane exhibited the characteristic features of a typical

plasma membrane—with one side facing the cytosol and the other exposed to the

extracellular environment (Figure 4D).

Cellular vesicles were observed near free membrane edges (Figures 4A-4C).

Some cytosol-immersed membrane tips appeared connected to vesicles (Figure 4A),

with instances of direct contact (Figure 4B) and possible fusion events (Figures 4B

and 4C). Vesicles also clustered along the anchoring area, as well as the separated

sister plasma membranes (Figure 4D). EM analysis revealed no morphological

differences between vesicles associated with the cytosol-immersed region and those

near conventional regions of the sister membranes (Figures 4B and 4D).

Analysis of a tri-nucleated cell during cytokinesis

To gain more comprehensive insight into membrane organization during cytokinesis,

we further applied the AutoCUTS-SEM (automatic collector of ultrathin sections

scanning electron microscopy) workflow (Li et al., 2017) to acquire serial EM

sections of a telophase cell. Through continuous tracing, we discovered that this cell

actually contained three independent nuclei. We color-coded each membrane region

according to its associated nucleus, with relative sizes estimated from EM sections to

ensure proportional representation (Figure 5A). The 3D model demonstrated that the

division plane between the yellow and blue cells was roughly perpendicular to that

between the purple cell and the other two (Figure 5A). The presence of three nuclei

within a shared cytoplasm suggests that one of the daughter cells entered the next

round of cell division without completing the previous one. This phenomenon—

concurrent divisions—appears to be common in early *C. elegans* embryos, as our EM

analysis identified more than 10 cells with three or four nuclei. For example, Figure

3D shows an EM section of a cell containing four nuclei.

Using consecutive EM sections, we traced membrane integrity in this tri-

nucleated cell. We found that this tri-nucleated cell was enveloped by three distinct

plasma membranes, each containing a single rupture site. These three ruptures

converged to form a "Y"-shaped exposed cytoplasmic region spanning >351 sections

(21,060 nm; Figures 5A and 5B and Video S2). Throughout these 351 consecutive

sections, we detected no continuity between adjacent membranes, indicating that membrane independence had been established by this stage. Consecutive tracing further showed that the MOs—frequently observed in single EM sections (Figures 1 and 2)—could arise from either the separation between sister membranes (Figure 5C) or from separated edges of the plasma membrane within the same cell (Figure 5D). Additionally, multiple openings within a single cell might represent different cross-sectional planes of a common membrane rupture (Figure 5E).

In all three cells (where cytoplasmic separation was incomplete), the free membrane edges were in close proximity (marked by *; Figures 5F and 5G), suggesting they were in late cytokinesis and that membrane extension mediated by detached membranes could persist into this stage. Taken together, these findings demonstrate that this tri-nucleated cell was undergoing concurrent cytokinesis, with three detached membranes independently extending toward a common cytoplasmic region to achieve cytoplasm individualization (Figure 5H). A similar mechanism may operate in four-nucleated cells (Figure 3D, for instance), in which two detached daughter membranes—without completing the last round of cell division—undergo further rupture, forming four granddaughter membranes. These membranes then extend independently to complete cell division (Figure 5H).

Capturing the membrane rupture and independent extension of detached sister membranes in living embryos

While high resolution HPF-TEM analysis revealed widespread membrane discontinuities in early *C. elegans* embryos, this static imaging approach cannot capture dynamic membrane behavior. To investigate whether optical microscopy can visualize membrane ruptures and the movement of severed sister membranes, we

employed a fluorescence-labeled membrane marker, GFP::PH(PLC1δ1) (see Methods

and Materials for details). Our live imaging detected both putative membrane ruptures

(Figure 6A) and free-ended sister membrane structures (Figures 6B and 6C),

providing additional evidence that membrane rupture and independent extension of

detached sister membranes underlie cytokinesis in *C. elegans* embryos (see

Discussion).

Sealing the exposed cytosol by neighboring membranes and extracellular matrix

layers

The integrity of the plasma membrane is essential for cell survival, as its disruption

can lead to cytoplasmic leakage and, if severe, result in cell death. However,

widespread cytoplasmic leakage has not been observed in wild-type *C.*

elegans embryos. Based on our EM observations, we propose two mechanisms

through which dividing embryonic cells may seal or cover exposed cytosolic regions.

First, the sharing of plasma membrane with neighboring cells may protect the

exposed cytosol. As repeatedly shown in EM images in this study (Figures 1, 3 and 4),

a single elongated free-ended membrane from one cell consistently helps define

boundaries with adjacent cells. These neighboring cells may be direct sister cells

derived from the same mother cell, or cousin cells originating from mother cells that

were once sisters. In cases where three nuclei share a common cytosol, the adjacent

cell could even be an aunt or nephew cell.

In addition, eggshell components may seal cytoplasmic regions lacking

membrane coverage. The *C. elegans* eggshell is composed of six distinct extracellular

matrix layers (Olson et al., 2012; Benenati et al., 2009; Rappleye et al., 1999;

Schierenberg and Junkersdorf, 1992; Stein, 2018). Among these, the fifth

permeability barrier layer (PBL), a lipid-rich structure, serves as the osmotic and

permeability barrier for the developing embryo (Olson et al., 2012; Benenati et al.,

2009). Normally, the PBL encloses the peri-embryonic layer (PEL), which separates it

from the embryonic plasma membrane (Olson et al., 2012; Benenati et al., 2009)

(Figures 7A and 7B). In contrast, in regions where the plasma membrane was absent,

we observed the PBL adhering directly to the exposed cytoplasm (red triangles and

dashed red lines, Figures 7B and 7C), bypassing the PEL. This phenomenon was also

evident in other EM images (Figures 1C and Figure 2, box 1). PBL-mediated sealing

is not always complete. When the PBL was not accessible or only loosely covered

membrane-free regions, minor cytoplasmic leakage occurred (Figure 7D). In cases

where PBL sealing was ineffective or absent, cytoplasmic material appeared to be

retained within a confined region, likely by the PEL (Figure 7D). However, the

mechanisms by which extracellular matrix components protect the exposed

cytoplasmic region remain unknown.

Discussion

In this study, we discovered that the plasma membrane of *C. elegans* embryonic cells

undergo rupture at metaphase. Following this rupture, the detached sister membranes

extend rather independently from each other to complete cytokinesis. To date, the

ultrastructural dynamics of plasma membrane integrity during cell division remained

unexplored in other species. We anticipate that our findings will catalyze comparative

studies across different species or under various pathological conditions, to determine

whether this mechanism represents a fundamental paradigm or a context-specific

strategy. Our work established the first mechanistic framework to guide such future

investigations.

The advantage of membrane rupture and independent sister membrane

extension

Cell division involves both nuclear and cytoplasmic division. However, due to the

difficulty of visualizing cell boundaries in living embryos, previous studies in *C.*

elegans have largely emphasized nuclear events. Here, we observed that early *C.*

elegans embryonic cells can contain three or four nuclei within a shared cytoplasm,

indicating that cytoplasmic division may be uncoupled from nuclear division. This

phenomenon is not unique to *C. elegans*—it has also been reported in other species

such as *Drosophila*, where the early embryo undergoes 13 rapid nuclear divisions

without cytokinesis before cellularization occurs. It has been proposed that by

minimizing or delaying cytokinesis, embryos can accelerate nuclear multiplication,

thereby supporting the rapid pace of early development. In this context, the embryonic

cells of *C. elegans* appear to adopt a notably rapid—though seemingly risky—strategy

for cell proliferation, yet one with clear developmental benefits.

First, membrane rupture may help alleviate the rise in intracellular pressure

during division (Stewart et al., 2011; Hiramoto, 1963). We observed that the size of

membrane discontinuities appeared to peak during metaphase, primarily around the

equatorial plane-consistent with localized pressure surges at this stage. Notably,

ruptures were also detected in tri-nucleated cell clusters, where all three cells

exhibited membrane discontinuities during late cytokinesis. This suggests that even

though the relative size of the ruptures decreases as mitosis proceeds, membrane

discontinuities may persist throughout the cell cycle. Maintaining an opening in the

plasma membrane during division likely reduces internal pressure, thereby conserving

energy and resources that would otherwise be expended on resisting membrane

tension. These conserved resources could then be redirected toward essential

processes such as chromosome segregation and membrane synthesis—a particular

advantage for rapidly dividing embryonic cells.

Of note, 3D reconstruction revealed that the metaphase rupture does not fully

bisect the membrane (Figure 2). However, given the limitations of our EM

sampling—neither random nor continuous imaging provides complete coverage—we

cannot exclude the possibility that some ruptures completely sever the membrane.

Regardless of the extent of separation, the rupture produces two distinct membrane

domains. Since this occurs as early as metaphase, before nuclear or cytoplasmic

division, the physical independence of the plasma membranes may serve as an early

marker of nascent daughter cells, preceding other cellular landmarks.

Following membrane rupture, the prospective sister membranes become

separated, raising the question of how cytokinesis proceeds with severed membranes.

Analysis of single EM sections revealed classical membrane invagination in only a

few cases (6 out of 51; Figure 3A). Notably, one such instance occurred during the

transition from a one-cell to a two-cell embryo. This suggests that while a subset of *C.*

elegans embryonic cells employs the membrane invagination and extension model,

the majority do not. Interestingly, membrane discontinuities were observed in one-cell

embryos (Figure S1), indicating that membrane rupture can co-occur with

invagination. Alternatively, in dividing cells, the back-to-back invagination may be

transient and may subsequently transition to free-ended extension. At present, distinguishing between these hypotheses remains challenging. A definitive resolution will likely require advanced imaging techniques capable of high-resolution, real-time visualization at the EM level.

Unlike a continuous bilayer, a free-ended membrane offers greater flexibility during cytokinesis. Without being physically constrained by its sister, each membrane can extend in different directions as needed. This independence may be particularly advantageous in accommodating asynchrony between nuclear and cytoplasmic division, as seen in early *C. elegans* embryos where multinucleated cells form through incomplete cytokinesis. In such cases, independent membrane growth from two directions facilitates cytoplasmic separation more effectively than a single continuous membrane would allow. Such autonomous movement may also explain the divergent orientations of sister membranes frequently observed in EM sections.

The expansion of the plasma membrane is generally driven by vesicle fusion. In this study, vesicles seem observed randomly distributed beneath the extending sister membranes. EM analysis further showed no morphological differences between vesicles near cytosol-immersed regions and those at conventional membrane sites. Since vesicles can fuse with the membrane at various locations, we speculate that membrane growth may be enhanced under these conditions—potentially accelerating cell cleavage during early embryogenesis. However, this remains speculative and requires further experimental validation.

Finally, when the separated membrane edges meet within daughter cells, full cytoplasmic separation is achieved. In the classical invagination-extension model, cytokinesis requires both membrane scission and fusion. In the detached-membrane

mode described here, however, cytokinesis proceeds without a continuous membrane structure. This raises important questions: Is membrane scission dispensable in this context? Could bypassing scission accelerate cell separation? If so, might such a mechanism offer a selective advantage during rapid embryonic development? These questions remain open and merit further investigation.

Limitations and perspectives

In this study, the application of HPF-TEM has enabled unprecedented ultrastructural visualization of dividing cell surfaces. Our findings reveal a cytoplasmic division mechanism that challenges current models and raises important new questions regarding the plasticity of cytokinesis.

HPF-TEM offers a significant advantage for membrane studies by capturing cellular structures in a near-native state. Through ultra-rapid freezing under high pressure, it suppresses ice crystal formation and preserves water in a vitreous state, thereby minimizing chemical artifacts and maintaining the integrity of lipids and proteins. This method provides a faithful snapshot of membrane architecture and dynamics(Moor et al., 1980). In our experiments, the consistent observation of limited number of membrane discontinuities-generally one per dividing cell-that are closely associated with cell division strongly suggests these are genuine biological structures rather than preparation-induced artifacts. While one might argue that such features could reflect EM-related artifacts specific to the plasma membrane during division, the correlative evidence presented here supports their biological relevance.

Using fluorescence-labeled membrane markers, we identified membrane ruptures and free-ended sister membrane structures in live embryos (Figure 6). Similar "abnormal" or "fuzzy" membrane signals have in fact been reported

previously(Harrell and Goldstein, 2011). However, interpretating these structures has

remained challenging for both technical and conceptual reasons. Technically, the

rapid kinetics of membrane extension-about 10-30 seconds during metaphase and less

than 3 minutes during cytokinesis-combined with cell motility, introduce inherent

spatiotemporal ambiguities. Conceptually, the longstanding dominance of the

membrane invagination model has likely shaped—and potentially constrained—

interpretations of membrane dynamics. This paradigm may have led researchers to

prioritize observations consistent with invagination, while dismissing ambiguous

structures such as free-ended extensions as artifacts, rather than considering them as

potential elements of alternative mechanisms. Furthermore, theoretical and

computational models of cytokinesis have largely been built on invagination-centric

assumptions, further narrowing the conceptual framework.

A key limitation is our current inability to capture 3D dynamics of these

membrane structures in real time, a challenge that awaits imaging technologies with

sufficient spatiotemporal resolution at near -EM levels.

Therefore, while our findings propose a transformative model for cytokinesis,

they open a new field of inquiry. Fully validating this mechanism and exploring its

conservation and implications will require an integrated approach combining

advanced live-cell imaging, biophysical modeling, and genetic tools.

Methods and Materials

Examination of *C. elegans* embryonic cells

Wild-type N2 worms were cultured on OP50-seeded NGM plates at 22°C (Brenner, 1974). To obtain images of early embryos, gravid hermaphrodites were collected and sectioned. At low magnification (×4000), individual embryonic cells were identified by the presence of a nucleus. In sections where the nucleus was not visible, cytoplasmic regions—whether covered or not covered by plasma membrane—were counted as individual cells if they were separated from surrounding cells. A total of 425 cells were examined (summarized in Figure 1C), with one EM section analyzed per cell. For embryos with fewer than 10 cells (one or two rows of cells), every cell in the given section was examined. For embryos with three or more rows of cells, only cells at the periphery were analyzed. At higher magnification (×30,000), the entire surface of each cell was examined, revealing 159 cells with one or more membrane openings. Higher magnification improved the resolution of cytoplasmic continuity, reducing potential misidentification of multinucleated cells or interconnected cytoplasmic regions that might have been miscounted at lower magnification. A total of 60 early embryos were serially sectioned as completely as possible, with each embryo yielding at least 20 sections. While Figure 1C presents summary data, images from additional sections were also used for further analyses. Since these additional sections may capture different cells, the total number of embryonic cells examined likely exceeds 425.

Additionally, we examined eight embryos laid outside the worm. Among these, four pre-comma stage embryos each contained at least one MO-positive cell. In another pre-comma stage embryo, no MOs were detected after examining 17 cells.

Analysis of one comma-stage embryo (19 cells examined) and two 1.5-fold-stage

embryos (>30 cells analyzed each) revealed no MOs.

Identification of MOs

The entire cell surface of individual embryonic cells was examined using electron

microscopy at 30,000× magnification. The isolated cytoplasmic regions without

plasma membrane coverage were defined as MOs. Sites where the plasma membrane

terminated and reappeared were marked as putative MOs. In periphery region of

worm embryos, in the non-plasma-membrane-covering regions, the PEL was

generally absent, while the PBL typically extended over the exposed cytoplasmic

surface. Plasma membranes often appeared wrinkled, whereas cytoplasmic regions

covered by the PBL exhibited a smoother appearance. Smooth, non-plasma-

membrane-covered regions overlaid by the PBL are readily identifiable as MOs.

When exposed cytoplasm was not tightly covered by the PBL, cytoplasmic leakage—

still confined to a specific area by the PEL—could also indicate a MO. Given the

variable morphology and frequent cytoplasmic leakage, membrane discontinuity size

could not serve as a reliable criterion for defining MOs. However, to ensure

robustness in our analysis, we considered only linear membrane discontinuities

exceeding 200 nm as genuine openings. Each MO was examined independently at

least three times. When necessary, adjacent sections were also reviewed to conform an

MO's presence, and identification was finalized only after all three assessments were

positive.

High-pressure freezing and transmission electron microscopy

Basic protocols for high-pressure freezing of a range of organisms, including *C.*

elegans, have been described (Weimer, 2006; McDonald, 2007; Manning and

Richmond, 2015). Briefly, adult worms were loaded into carriers and cryofixed using

a Leica Microsystems EM ICE high-pressure freezer, followed by automatic cooling

in liquid nitrogen. The carriers used for freezing (Leica Microsystems, Germany;

catalog nos. 16770153 and 16770152) were coated with a non-stick layer of 1-

hexadecene (Sigma-Aldrich). After high-pressure freezing (HPF), the samples were

transferred under liquid nitrogen to a Leica AFS-2 freeze-substitution unit and

incubated at −90°C for 72 h in a freeze-substitution solution consisting of acetone

with 2% (wt/vol) osmium tetroxide. The temperature was then raised at a rate of

8°C/h for 4 h, held at −60°C for 12 h, increased again at 5°C/h for 6 h, held at −30°C

for 10 h, and finally raised to 10°C at a rate of 4°C/h for 10 h. Samples were washed

four times in acetone, stained with 1% uranyl acetate for 1 h, and rinsed three times

with pure acetone. They were then infiltrated stepwise in increasing concentrations of

SPI-Chem™ resin (2:1 resin:acetone for 3 h, 1:1 for 5 h), followed by two incubations

in 100% fresh resin (4 h each). The samples were placed in fresh resin in embedding

molds and polymerized in a 60°C oven for 3 days. Ultrathin sections (60 nm) were cut

using a diamond knife (Diatome) on an ultramicrotome (Leica Ultracut UC7) and

collected on slot copper grids (EMS). The sections were visualized using a Hitachi

HT7700 transmission electron microscope operating at 80 kV. Images were recorded

with a Gatan 832 (4k × 2.7k) CCD camera. Data from four independent preparations

were analyzed in this study.

3D reconstruction

The metaphase 3D model was prepared using the following steps. First, resin blocks

containing gravid adults were prepared via high-pressure freezing and freeze

substitution. Continuous ultrathin sections (60 nm thick) were manually collected

using a Leica UC-7 ultramicrotome and mounted on slot grids for TEM imaging. A

total of 112 serial sections were collected, though 3 sections were lost due to

obstruction, resulting in 822 TEM images. Among these, 105 sections contained

metaphase-stage cell information. To examine the plasma membrane integrity, high-

magnification imaging was performed. Since a single high-magnification image could

not cover the entire cell, multiple overlapping images were captured per section and

stitched together using ImageJ to generate a complete composite image of the cellular

morphology at each section. In total, 292 images were stitched. The 105 stitched

images were then aligned using ImageJ, and 99 of these aligned images were used for

3D reconstruction using Imaris software. About a quarter of the cell lacked membrane

discontinuities. For this portion, we estimated its size and shape based on EM sections

to ensure proportional representation. Of note, single MOs smaller than 200 nm that

appeared continuously across more than 3 consecutive sections were considered the

endpoint of the membrane rupture. Based on this, the actual length of the membrane

rupture could be underestimated at one end, while the other end was clearly defined.

The ultrastructural 3D study of the tri-nucleated cell was performed using an

AutoCUTS device (Li et al., 2017) and a Helios Nanolab 600i dual-beam scanning

electron microscope (FEI). Adult worms were treated with high-pressure freezing,

freeze-substitution, and embedding as described above. Subsequently, 351 serial

sections (60 nm thick) were collected using the AutoCUTS device, and scanning

electron microscopy imaging was performed. All datasets were analyzed using

IMARIS 8 software. After Z-stacks were aligned, the membrane rupture was manually

traced and reconstructed.

Live imaging of plasma membrane dynamics using GFP::PH

The plasma membrane was labeled using GFP::PH(PLC1δ1). The strain carrying this

marker was obtained from the Caenorhabditis Genetics Center and was originally

generated in Dr. Karen Faye Oegema's laboratory. Gravid adult hermaphrodites

expressing this marker were collected and washed with M9 buffer. Embryos were then

released from the gravid adult worms using a 1 mL syringe needle and transferred

onto a 2% agarose pad. Fluorescence imaging was performed on a Zeiss LSM-980

microscope with a 63×/1.42 oil objective. Only one embryo at a time was imaged,

starting at the 2- to 4-cell stage. Images were acquired at 4 or 8-second intervals, with

total imaging duration typically kept under 30 minutes to ensure embryo viability.

Acknowledgements

We thank the Center for Biological Imaging at the Institute of Biophysics, Chinese Academy of Sciences (CAS) for technical support. We are grateful to Drs. John White, Andrew Chisholm, and Zhuo Du for their encouragement and insightful suggestions. This work was supported by grants from the National Key R&D Program of China (2024YFA1803401, 2021YFA0805802, and 2024YFA1306100) and the Natural Science Foundation of China (32170790 and 32321004).

Author contributions

Study concept and design: MD, JL. EM analyses: JL, TH, MD. Drafting and major editing of original manuscript: MD, JL, TH, XH. All authors reviewed, revised, and approved the final version of the paper.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to Mei Ding

Figure legends

Figure 1. Membrane discontinuities in *C. elegans* embryonic cells.

A, Schematic drawing of a metaphase cell and the classical membrane invagination model. B, Schematic drawing of a metaphase cell with a ruptured plasma membrane and a cytokinesis model showing that the detached sister membranes mediate membrane extension in *C. elegans* embryonic cells. C, A membrane opening (MO, indicated by white dashed line) in an embryonic cell at the embryo's periphery. D, Membrane openings in the embryo's central regions. Yellow arrows point to single plasma membranes (SPM) traversing the cytoplasm. Plasma membranes are in blue. E, Summary of the single EM section survey of plasma membrane continuity. Orange lines outline the plasma membranes. Solid gray circles/ellipses enclosed by dashed lines represent evenly stained nucleoid regions. Condensed chromosome structures are colored gray-blue. Light brown areas represent cytoplasmic regions.

Figure 2. Metaphase membrane rupture.

A, Membrane discontinuities or openings (MO) (white dashed line) in a metaphase cell (MC). Enlarged views of the boxed regions (1, 2 and 3) are shown. Yellow arrows indicate the plasma membrane (PM). B, MOs (marked by 1 and 2) are detected in anaphase (AC) and other early embryonic cells; enlarged views of the MOs are shown (white dashed lines). All yellow arrows point from the interior of the cell toward the plasma membrane. C and D, Three-dimensional (3D) reconstruction of a metaphase cell. Representative EM images are shown in (C). Purple lines outline the plasma membranes. White dashed lines outline the exposed cytoplasmic region. Blue lines outline the condensed chromosome region. D, The 3D reconstruction of the metaphase cell. The plasma membrane (PM) and condensed chromosome region are

displayed.

Figure 3. The detached severed sister membranes.

A, Representative image of a classical back-to-back membrane arrangement. B,

Representative image of detached severed sister membranes. Diagrams of the

corresponding cytokinesis models are displayed on the right. C, Tips of two detached

sister membranes. D, Cytosol-immersed single membranes separating the cytoplasmic

regions of two adjacent cells. Enlarged views of regions 1 and 2 are shown below.

Plasma membranes (PM) are indicated by yellow arrows (A, B and D) or blue lines.

Asterisks indicate membrane tips.

Figure 4. Cellular vesicles on detached sister membranes.

A, A vesicle localized at the tip of a cytosol-immersed membrane. B, Vesicles

clustering near the cytosol-immersed region of sister membranes. Contact or fusion

between vesicles and membranes is indicated. The rectangular inset shows an

enlarged view of a gap junction (GJ) plaque. C, Putative fusion events between

vesicles and free-ended membranes. D, Vesicles found near the separated sister

membranes. Plasma membranes are in blue.

Figure 5. Membrane rupture on a tri-nucleated cell.

A, The 3D model of a tri-nucleated cell showing the membrane rupture (black). B,

Reconstruction of the membrane ruptures and adjacent plasma membranes. Individual

membranes, each associated with a distinct nucleus, are color-coded (blue, yellow,

and purple). C and D, Membrane openings (MO, dashed white lines) arise either from

separation between sister membranes (C) or between the severed membrane edges of

the same cell (D). E, In a single EM section, the multiple openings originate from the

same rupture. F and G, The opposing severed edges are close to each other in the

yellow, purple and blue cells. Plasma membranes (PM) are indicated in blue.

Asterisks indicate membrane tips. H, Schematic diagram of the formation of cells

with three or four nuclei.

Figure 6. Detection of membrane disruptions by optical microscopy.

A, The putative membrane rupture and extension (*). B, Two extending free-ended

sister membranes (*). Insets are enlarged views. C, Possible free-ended edges of

detached membranes, probably from multiple cells (*). Below are proposed models

illustrating the membrane disruptions that could explain the anomalous GFP signals.

Images were taken at 4-second intervals for (A and B) and 8-second intervals for (C).

Green signal is GFP::PH(PLC1δ1).

Figure 7. The exposed cytoplasmic regions are covered by extracellular matrix

layers.

A-C, The exposed cytoplasmic region is covered by the permeability barrier layer

(PBL, red triangles and red dashed lines). PM, plasma membrane (yellow arrows and

blue lines). The boxed region in (A) is enlarged in (B), and the boxed region in (B) is

enlarged in (C). D, The exposed cytoplasmic region can also be confined by the peri-

embryonic layer (PEL).

Supplementary Figure Legends

Figure S1. Membrane discontinuities in externally laid *C. elegans* embryos.

A, An externally laid embryos with a membrane discontinuity (white box). B, Higher-

magnification view of the membrane opening (MO) (white dashed line). Yellow

arrow, plasma membrane; asterisk (*), condensed chromosome.

Figure S2. The wide distribution of cells with membrane discontinuities in early

***C. elegans* embryos.**

A, Schematic diagram of *C. elegans* embryogenesis. B-N, Representative low-

magnification EM images of early embryos within adult hermaphrodites. Red lines

outline cells contain membrane openings. In (F), a tri-nucleated cell is marked with &.

(L) and (M) show embryos during the transition from the one-cell to the two-cell

stage.

Figure S3. Membrane discontinuities in cells during mitosis.

A-C, Membrane discontinuities or openings (MO, white or black dashed lines) in cells

at metaphase (A), anaphase (B) or telophase (C). Yellow arrows points to plasma

membranes (PMs).

Video S1. 3D reconstruction of the metaphase cell in Figure 2.

The plasma membrane is shown in red, the condensed chromosome is in blue, and

exposed cytoplasm is in green.

Video S2. 3D reconstruction of the tri-nucleated cell in Figure 5.

3D reconstruction of membrane ruptures and adjacent plasma membranes. Individual

membranes, each associated with a distinct nucleus, are color-coded (purple, yellow,

or blue).

Bao, Z., Z. Zhao, T.J. Boyle, J.I. Murray, and R.H. Waterston. 2008. Control of cell cycle timing during C. elegans embryogenesis. *Developmental Biology*. 318:65–72. doi:10.1016/j.ydbio.2008.02.054.

Bembenek, J.N., C.T. Richie, J.M. Squirrell, J.M. Campbell, K.W. Eliceiri, D. Poteryaev, A. Spang, A. Golden, and J.G. White. 2007. Cortical granule exocytosis in *C. elegans* is regulated by cell cycle components including separase. *Development*. 134:3837–3848. doi:10.1242/dev.011361.

Benenati, G., S. Penkov, T. Müller-Reichert, E.V. Entchev, and T.V. Kurzchalia. 2009. Two cytochrome P450s in Caenorhabditis elegans are essential for the organization of eggshell, correct execution of meiosis and the polarization of embryo. *Mechanisms of Development*. 126:382–393. doi:10.1016/j.mod.2009.02.001.

Brenner, S. 1974. THE GENETICS OF *CAENORHABDITIS ELEGANS*. *Genetics*. 77:71–94. doi:10.1093/genetics/77.1.71.

Deppe, U., E. Schierenberg, T. Cole, C. Krieg, D. Schmitt, B. Yoder, and G. Von Ehrenstein. 1978. Cell lineages of the embryo of the nematode Caenorhabditis elegans. *Proc. Natl. Acad. Sci. U.S.A.* 75:376–380. doi:10.1073/pnas.75.1.376.

Edgar, L.G., and J.D. McGhee. 1988. DNA synthesis and the control of embryonic gene expression in C. elegans. *Cell*. 53:589–599. doi:10.1016/0092-8674(88)90575-2.

Gerien, K.S., and J.-Q. Wu. 2018. Molecular mechanisms of contractile-ring constriction and membrane trafficking in cytokinesis. *Biophys Rev*. 10:1649–1666. doi:10.1007/s12551-018-0479-3.

Harrell, J.R., and B. Goldstein. 2011. Internalization of multiple cells during C. elegans gastrulation depends on common cytoskeletal mechanisms but different cell polarity and cell fate regulators. *Developmental Biology*. 350:1–12. doi:10.1016/j.ydbio.2010.09.012.

Hiramoto, Y. 1963. Mechanical properties of sea urchin eggs. I. Surface force and elastic modulus of the cell membrane. *Experimental Cell Research*. 32:59–75. doi:10.1016/0014-4827(63)90069-7.

Li, X., G. Ji, X. Chen, W. Ding, L. Sun, W. Xu, H. Han, and F. Sun. 2017. Large scale three-dimensional reconstruction of an entire Caenorhabditis elegans larva using AutoCUTS-SEM. *Journal of Structural Biology*. 200:87–96. doi:10.1016/j.jsb.2017.09.010.

Manning, L., and J. Richmond. 2015. High-Pressure Freeze and Freeze Substitution

Electron Microscopy in C. elegans. *In* C. elegans. D. Biron and G. Haspel, editors. Humana Press, Totowa, NJ. 121–140.

McDonald, K. 2007. Cryopreparation Methods for Electron Microscopy of Selected Model Systems. *In* Methods in Cell Biology. Elsevier. 23–56.

Modern Electron Microscopy Methods for C. elegans. 2012. *In* Methods in Cell Biology. Elsevier. 93–149.

Moor, H., G. Bellin, C. Sandri, and K. Akert. 1980. The influence of high pressure freezing on mammalian nerve tissue. *Cell Tissue Res.* 209. doi:10.1007/BF00237626.

Olson, S.K., G. Greenan, A. Desai, T. Müller-Reichert, and K. Oegema. 2012. Hierarchical assembly of the eggshell and permeability barrier in *C. elegans*. *Journal of Cell Biology*. 198:731–748. doi:10.1083/jcb.201206008.

Pintard, L., and B. Bowerman. 2019. Mitotic Cell Division in *Caenorhabditis elegans*. *Genetics*. 211:35–73. doi:10.1534/genetics.118.301367.

Pollard, T.D., and B. O'Shaughnessy. 2019. Molecular Mechanism of Cytokinesis. *Annu. Rev. Biochem.* 88:661–689. doi:10.1146/annurev-biochem-062917-012530.

Rappleye, C.A., A.R. Paredez, C.W. Smith, K.L. McDonald, and R.V. Aroian. 1999. The coronin-like protein POD-1 is required for anterior-posterior axis formation and cellular architecture in the nematode Caenorhabditis elegans. *Genes & Development*. 13:2838–2851. doi:10.1101/gad.13.21.2838.

Schierenberg, E., and B. Junkersdorf. 1992. The role of eggshell and underlying vitelline membrane for normal pattern formation in the early C. elegans embryo. *Roux's Arch Dev Biol*. 202:10–16. doi:10.1007/BF00364592.

Stein, K.K. 2018a. The C. elegans eggshell. *WormBook*. 1–36. doi:10.1895/wormbook.1.179.1.

Stewart, M.P., J. Helenius, Y. Toyoda, S.P. Ramanathan, D.J. Muller, and A.A. Hyman. 2011. Hydrostatic pressure and the actomyosin cortex drive mitotic cell rounding. *Nature*. 469:226–230. doi:10.1038/nature09642.

Sulston, J.E., E. Schierenberg, J.G. White, and J.N. Thomson. 1983. The embryonic cell lineage of the nematode Caenorhabditis elegans. *Developmental Biology*. 100:64–119. doi:10.1016/0012-1606(83)90201-4.

Weimer, R.M. 2006. Preservation of *C. elegans* Tissue Via High-Pressure Freezing and Freeze-Substitution for Ultrastructural Analysis and Immunocytochemistry. *In* C. elegans. Humana Press, New Jersey. 203–222.

Zhang, Y., J.M. Foster, L.S. Nelson, D. Ma, and C.K.S. Carlow. 2005. The chitin
synthase genes chs-1 and chs-2 are essential for C. elegans development and
responsible for chitin deposition in the eggshell and pharynx, respectively.
Developmental Biology. 285:330–339. doi:10.1016/j.ydbio.2005.06.037.

Zhu, J., S.M. Lam, L. Yang, J. Liang, M. Ding, G. Shui, and X. Huang. 2022. Reduced
phosphatidylcholine synthesis suppresses the embryonic lethality of seipin
deficiency. *Life Metabolism*. 1:175–189. doi:10.1093/lifemeta/loac021.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Sup Figure 1

Sup Figure 2

A

in utero development (150 min)
First cleavage (40 min)

eggs laid at gastrula
(28-cell)

Comma

1.5-fold

Sup Figure 3