Uranium mining fuels evolution in deep groundwater microbiomes

Wei Xiu[1], Till L.V. Bornemann[2], Tianjing Zhang[1], André R. Soares[2], Luping Xie[1], Guoxi

Lian[3], Haicheng Weng[4], Bing Yang[4], Ruixuan Gai[1], Zhipeng Gao[1], Di Zhang[1],

Jonathan R Lloyd[5], Huaming Guo[1]*, and Alexander J. Probst[2]*

1 State Key Laboratory of Geomicrobiology and Environmental Changes (GMEC),

China University of Geosciences, Beijing 100083, People's Republic of China

2 Environmental Metagenomics, Research Center One Health Ruhr, University

Alliance Ruhr, Faculty of Chemistry, University of Duisburg-Essen, Germany

3 Nuclear and Radiation Safety Center, Ministry of Ecology and Environment, Beijing,

100082, China

4 Beijing Research Institute of Chemical Engineering and Metallurgy CNNC, No.145

Jiukeshu, Tongzhou District, Beijing 101149, China

5 Williamson Research Laboratories for Molecular Environmental Science,

Department of Earth and Environmental Sciences, The University of Manchester,

Manchester M13 9PL, United Kingdom

* to whom correspondence should be addressed: Prof. Alexander J. Probst

(alexander.probst@uni-due.de); Prof. Huaming Guo (hmguo@cugb.edu.cn)

Abstract

Microbial communities in the deep biosphere underpin global biogeochemical cycles, yet their evolutionary responses to extensive anthropogenic disturbances remain unclear. Uranium in-situ leaching (ISL), a dominant mining strategy, creates persistent radiochemical and redox gradients in aquifers, providing a unique system for studying subsurface microbial evolution. Here, by exploiting metagenomics and metatranscriptomics in examining microbial responses through various mining stages at neutral U ISL sites, we show that geochemical changes accompany significant alterations in microbial community composition, oxidative stress, biogeochemical cycling, and overexpression of DNA repair genes. Comparative metagenomics and metatranscriptomics revealed elevated non-synonymous to synonymous polymorphism ratios and altered nucleotide diversity correlated with radiochemical gradients, underscoring the influence of environmental stress in promoting microbial evolution. Our results demonstrate that neutral ISL drives adaptive genetic and functional diversification of subsurface microbiomes and extend core microbial evolutionary frameworks to anthropogenically modified subsurface environments, redefining the deep biosphere as a dynamic, responsive system.

Introduction

Microbial consortia in the deep subsurface adapted to nutrient-scarce, anoxic environments[1–4] mediate critical biogeochemical processes and help maintain groundwater quality[5–8]. However, anthropogenic stresses arising from mining, groundwater extraction, and energy production are perturbing these fragile environments at increasing rates[9]. Recent studies have shown that environmental stresses, including heavy metal contamination and altered pH, affect groundwater microbial communities[10,11]. Mining operations are particularly disruptive, as they rapidly subject pristine subsurface ecosystems to heavy metals, radioactive contaminants, altered redox conditions, and electrochemical gradients[9]. Notably, a recent study in the abyssal seafloor showed that deep-sea mining can cause biological disruptions persisting for decades[12], underscoring the need to assess long-term impacts of subsurface resource extraction on microbiomes. Although distinct from the deep subsurface examined here, this finding underscores how resource extraction can impose long-lasting impacts on microbial communities in Earth's hidden biomes.

In-situ leaching (ISL), a widely used uranium (U) mining technique, is based on injecting oxidizing agents into aquifers to mobilize U by oxidizing insoluble U(IV) to soluble U(VI). In addition to fundamentally altering groundwater chemistry[13], this activity imposes profound chemical and radiological stress on microbial communities that have evolved over eons under stable, reducing conditions. Whereas the environmental consequences of ISL are well documented mostly via 16S rRNA gene analysis[11,14–16], the evolutionary responses of microbial communities to these stressors remain poorly understood. Indeed, while mining-driven alterations to groundwater chemistry are expected to disrupt the microbial community structure,

functional potential, and genetic diversity[17–20], the question remains as to whether

anthropogenic stress can drive microbial evolution in deep aquifers.

To this end, we implement innovative metagenomic and metatranscriptomic

strategies to examine microbial community dynamics and evolutionary adaptations at

a neutral-pH ISL area ($CO_2 + O_2$ ISL). It involved sampling aquifers spanning mining

spanning pre-mining, active mining, and post-mining phases. Leveraging this rare

sampling opportunity to study microbial adaptation as a response to mining and

geochemical changes, we show that U exposure and radiochemical stress drive the

restructuring of microbial communities, oxidative stress response and DNA repair, and

alterations in genetic diversity. Elevated pN/pS ratios (the ratio of the rates of non-

synonymous (pN) and synonymous (pS) polymorphism) and shifts in nucleotide

diversity implicate mining-induced environmental gradients in significantly altering

microbial evolutionary processes. Our findings yield new insights into how deep

subsurface microbiomes structurally and evolutionally adapt to anthropogenic stress,

bearing implications for global biogeochemical cycles, ecosystem resilience, and

bioremediation strategies. As mining operations continue to intersect with fragile

groundwater ecosystems, understanding the evolutionary implications to subsurface

microbial communities will be of critical importance to predicting the long-term impacts

of resource extraction from Earth's deep biosphere. The findings of this investigation

underscore the consequences of uranium ISL on the evolution of subsurface life,

offering a novel framework for appraising microbial resilience and adaptation in

mining-impacted deep subsurface ecosystems.

RESULTS

Geochemical and radiological alterations during and following Uranium ISL

Neutral pH uranium *in-situ* leaching (U ISL) commences while the aquifer is in

its natural state (pre-mining phase), proceeds through the injection of leaching

solutions (mining phase), and concludes with natural restoration efforts upon cessation

of mining activities (post-mining phase; *Fig. 1*). We monitored the all phases of neutral-

pH ISL, having collected 80 groundwater samples from four distinct mining sites (S1,

S2, S3a, and S3b; *Figs. 1b-c*). These sites underwent neutral ISL for varying durations,

creating a unique time-resolved dataset with considerable variation in their redox

conditions, from neutral-reducing pre-mining (BG, baseline groundwater) to weakly

acid-oxidizing during mining (MT, mining groundwater), and back to neutral-reducing

post-mining (PMI, PMIS, and PMS, representing initial, intermediate, and stable post-

mining phases; *Figs. 1b–c*).

In the neutral-reducing pre-mining phase, low concentrations of total dissolved

uranium (TU; 0.44 ± 0.31 mg/L*; Fig. 2b*) were detected, indicative of ongoing U

mineralization under reducing conditions. This environment was rich in electron

donors, including dissolved organic carbon (DOC, 5.26 ± 3.51 mg/L), Fe(II) (0.85 ±

1.11 mg/L), ammonium (0.73 ± 0.70 mg/L), and sulfide (0.92 ± 1.18 mg/L) (*Fig. 2* and

Table S1). These sources likely fuelled the reduction of Fe(III), and sulfate, and

enhanced the reductive immobilization of U in the aquifer.

During the oxidizing mining phase, the injection of $CO_2 + O_2$ mobilized U from

ore deposits. Total dissolved uranium (TU) concentrations increased rapidly after *in-*

situ leaching activities commenced (from MT0.25 to MT1), peaking after roughly four

years (MT4), followed by a gradual decline thereafter (from MT5 to MT20) (*Fig. 2b*).

The oxidation of U(IV) to U(VI) and the subsequent formation of soluble uranyl-

carbonate complexes were the primary drivers of TU increase[21,22], as evidenced by

strong positive correlations between TU and bicarbonate concentrations (Spearman

correlation, ρ = 0.43, n = 80, p < 0.0001) (*Fig. S1*). Significant increases in sulfate and

nitrate were also observed during this phase, both positively correlated with TU

(sulfate-TU, ρ = 0.65, n = 80, p < 0.0001; nitrate-TU, ρ = 0.69, n = 80, p < 0.0001) (*Fig.*

S1), likely resulting from the oxidation of pyrite and ammonium, respectively[21,22].

Upon ISL decommissioning, the groundwater redox potential (Eh) decreased

significantly, likely due to the degradation of dissolved organic matter (DOM;

particularly protein-like and humic substances[23]). This transition from oxic to anoxic

conditions was further evidenced by a decline in TU, sulfate, and sulfate/Cl ratios

declined, while Fe(II) concentrations increased. This   suggested the microbial

reduction of Fe(III) and sulfate, alongside the reductive natural attenuation of U(VI)[23].

TU concentrations did not return to pre-mining levels (*Fig. 2*), however, indicative of

incomplete restoration of the aquifers' original geochemical conditions.

As U and its decay products emit α and β radiation, changes in groundwater

radioactivity can serve as indicators of U mobilization and contamination. Monitoring

α and β activity therefore during mining operations, with elevated gross (α + β) values

indicating radiogenic element mobilization, while declining levels after mining suggest

the efficacy of natural attenuation processes in limiting further U migration. Here, gross

(α + β) radiation levels increased significantly after mining activities commenced, but

decreased following ISL decommissioning (Figs. *2c-e*). These changes in radioactivity

correlated strongly with TU concentrations (ρ = 0.24, n = 80, p < 0.0001; Fig. *S1*),

underscoring the coupling between uranium mobilization and radiogenic activity during

mining and attenuation phases.

Figure 1 | Sampling locations and study overview. *(a) Geographic location of the Songliao Basin and broader regional context, modified from a previously published source[24]. (b) Schematic layout of study sites along a uranium ISL chronosequence: Site 1, unmined aquifer; Site 2, recent ISL (0.25–1 yr); Site 3a, intermediate ISL (4–6 yr); and Site 3b, long-term ISL (15–20 yr), which after decommissioning transitioned through post-mining phases of iron-reducing (PMI), mixed iron- and sulfate-reducing (PMIS), and sulfate-reducing (PMS) conditions. Wells shown with two names indicate that the same well was sampled at different time points. The background map shows oxidized sandstone (yellow), reduced sandstone (grey), uranium-bearing sandstone (pink), uranium ore deposits (red), and ISL units (arrow pointers). (c) Numbers of groundwater samples collected for hydrochemical, metagenomic, and*

Figure 2 | Geochemical profiles of groundwater across neutral-pH ISL phases.

Boxplots show geochemical parameters through the stage of neutral-pH ISL, including

pre-mining (BG), mining (MT), and post-mining (PM). Sampling groups include: Site 1

(unmined aquifer, S1_BG); Site 2 (unmined aquifer, S2_BG; recent ISL at 0.25-1

years: S2_MT0.25 and S2_MT01); Site 3a (unmined aquifer, S3a_BG; intermediate

ISL, 4-6 years: S3a_MT04, S3a_MT05, and S3a_MT06); and Site 3b (unmined

aquifer, S3b_BG; long-term ISL, 15-20 years: S3b_MT15 and S3b_MT20; and

decommissioned ISL under iron-reducing, S3b_PMI, mixed iron- and sulfate-reducing,

S3b_PMIS, and sulfate-reducing, S3b_PMS conditions). Panels represent (a – p) Eh,

gross (α+β), pH, DOC, bicarbonate, sulfide, sulfate, sulfate/Cl, Fe(II), ammonium,

nitrite, nitrate, and total Fe. Colored backgrounds denote different ISL phases. Colors

indicate study sites: light blue (Site 1), light green (Site 2), medium green (Site 3a),

and beige (Site 3b).

Thousands of metagenome-assembled genomes reveal microbiome changes

and enhanced co-occurence resulting from *in-situ* mining activity

Sixty groundwater samples representative of the pre-mining, mining, and post-

mining conditions across more than 20 years of neutral-pH ISL were metagenomically

sequenced (*Fig. 1*). Nonpareil[25] analysis indicated that sequencing coverage,

exceeding 75%, was of sufficient depth to accurately reflect the aquifer microbiome

diversity (*Fig. S2*). To assess both bacterial and archaeal diversity, we focused on

analyzing the ribosomal protein S3 (*rpS3*) gene[26], a single-copy, universally conserved

marker that provides high phylogenetic resolution for both bacteria and archaea,

yielding 12,972 sequences that clustered (at 99% similarity) into 6,487 representative

sequences. Taxonomic assignments based on the Genome Taxonomy Database

(GTDB, r220) grouped 6,431 rpS3 sequences into 215 distinct phyla (21 archaeal, 194

bacterial; *Fig. S3*). Groundwater microbial community composition and alpha diversity,

based on *rpS3* gene profiles, were strongly influenced by geochemical gradients,

particularly uranium concentration and radiochemical activity, with higher diversity and

distinct community structures observed under elevated stress conditions (*Figs. S4 and*

S5).

Through bioinformatic analysis, we identified 4,036 medium-quality

metagenome-assembled genomes (MAGs; completeness > 50%, contamination <

10%)[27], from which 2,255 strain-cluster representative MAGs (strain-cluster rMAGs)

were obtained via dereplicating at strain-level (99% ANI). These included 1,165 high-

quality and 1,090 medium-quality strain-cluster rMAGs bearing a median

completeness of 96.5%, contamination of 1.7%, and N50 of 50,891 (Fig. *3* and *Table*

S2). These MAGs represented 64 of the 215 phyla detected and accounted for roughly

35% of the total phylum-level diversity detected via *rpS3*, thus providing a robust

representation of the predominant microbial communities at Qianjiadian sites (*Figs.*

S3, S6, and S7). The phyla comprising the remaining 6.9% of rpS3-based abundance

were likely absent from the strain-cluster rMAGs because their low abundance

hindered successful genome reconstruction (*Fig. 2a*).

Neutral-pH ISL significantly altered microbial diversity in the mining aquifer, as

revealed by *rpS3* gene profiles. Microbial diversity showed strong positive correlations

with uranium concentration and radioactivity, with prolonged neutral-pH

ISL(S3a_MT04 and S3a_MT05) associated with increased Shannon and Invsimpson

indices (ANOVA, $p < 0.05$; Tukey HSD post-hoc test; *Fig. S4*). Microbial community

composition shifted markedly across the ISL chronosequence, as shown by non-

metric multidimensional scaling (NMDS) and sample-to-sample correlation analyses

of strain-cluster rMAGs (*Figs. 3e, 4a–d*). Communities from pre-mining, mining, and

post-mining stages formed distinct clusters, with ANOSIM indicating significant

separation between stages (R = 0.72, $p < 0.001$; Fig. 4d), reflecting divergent

community trajectories during and after ISL. Network analysis of relative abundances

of strain-cluster rMAGs revealed positive correlations, accounting for 90.4% of all edges, defined as significant correlations between taxa, dominating microbial interactions (*Fig. 4e*). Four distinct modules were identified, with microbial compositions of Module 1 remaining stable throughout neutral-pH ISL, indicative of a core microbial community. By contrast, the microbial compositions of Modules 2 and 4 were more prevalent during the early stages of neutral-pH ISL (indicative of adaptation), while that of Module 3 associated primarily with post-mining (indicative of recovery and/or attenuation; *Figs. 4f-i*). These results highlight the significant shifts in microbial community structure, diversity, and ecological interactions imposed by neutral-pH ISL. The observed tendency towards positive interactions emphasizes the dynamic and evolving nature of subsurface microbiomes in response to acute environmental stress, although such co-occurrence may arise from a variety of ecological processes .

Figure 3 | Characteristics and distribution of strain-cluster representative metagenome-assembled genomes (strain-cluster rMAGs). *(a) Rank abundance of strain-cluster rMAGs across phyla, with the top 100 depicted in more detail. The displayed abundances represent the average strain-cluster rMAG abundances across all samples. (b) Strain-cluster rMAGs completeness and contamination by quality tier. (c) Distribution of log(N50) values of strain-cluster rMAGs. (d) Genome size comparison between archaeal and bacterial strain-cluster rMAGs. (e) Heatmap of*

strain-cluster rMAGs Z-score normalized relative abundance across sampling sites,

clustered by similarity. Sampling groups include: Site 1 (unmined aquifer, S1_BG);

Site 2 (unmined aquifer, S2_BG; recent neutral-pH ISL at 0.25-1 years: S2_MT0.25

and S2_MT01); Site 3a (unmined aquifer, S3a_BG; intermediate neutral-pH ISL, 4-6

years: S3a_MT04, S3a_MT05, and S3a_MT06); and Site 3b (unmined aquifer,

S3b_BG; long-term neutral-pH ISL, 15-20 years: S3b_MT15 and S3b_MT20; and

decommissioned neutral-pH ISL under iron-reducing, S3b_PMI, mixed iron- and

sulfate-reducing, S3b_PMIS, and sulfate-reducing, S3b_PMS conditions).

Figure 4 | Microbial community structure and modularity across *neutral-pH ISL*

phases. *(a) Heatmap of sample-to-sample Spearman's rank correlations of strain-*

cluster rMAG abundances across early (blue), mid (green), and late (red) neutral-pH

ISL stages. (b) Taxonomic composition of top 12 phyla across samples, with the

sample order consistent with that in the heatmap. (c) Non-metric multidimensional

scaling (NMDS) plot of community structure based on strain-cluster rMAGs (stress =

0.141). (d) Analysis of Similarity (ANOSIM) showing significant differences between

groups (R = 0.7217, p = 0.001). (e) Network of strain-cluster rMAGs co-occurrence,

with positive (grey) and negative (red) Spearman correlations grouped into four

modules. (f–i) Distribution of the abundance of module-specific taxa across neutral-pH

ISL phases.

Neutral-pH ISL impacts gene expression linked to DNA repair, oxidative stress, and biogeochemical cycling

To investigate microbial metabolic adaptations in response to long-term neutral-pH ISL, we analyzed gene expression from 45 metatranscriptomic samples representative of pre-mining, mining, and post-mining conditions over a 20-year period. By integrating gene expression profiles with our suite of 2,255 de-replicated and functionally annotated strain-cluster rMAGs, we identified widespread overexpression (coverage increase at least 2-fold) of genes involved in DNA repair, oxidative stress responses, and biogeochemical cycling (*Fig. S8-S14*). A corresponding Spearman correlation heatmap revealed coherent expression patterns and their associations with key environmental drivers (*Fig. 5a*). Principal coordinate analysis (PCoA) identified key environmental variables, such as Fe(II), gross alpha radioactivity, and bicarbonate, as major drivers of gene expression during different neutral-pH ISL phases (*Figs. 5b-d*). Specifically, Fe(II), gross radioactivity (alpha and beta) and redox potential (Eh) associated with the overexpression of DNA repair genes

(*Fig. 5b*), while Fe(II) and nitrate correlated with oxidative stress response gene

expression (*Fig. 5c*). Changes in bicarbonate, Fe(II), Eh and pH affected genes

involved in biogeochemical cycling (*Fig. 5d*). These findings indicate that both

geochemical and radiochemical gradients jointly shape microbial metabolic responses

during neutral-pH ISL.

Microbial populations at the 20-year ISL site exhibited extensive DNA damage

responses, as shown by the overexpression of genes involved in homologous

recombination, nucleotide excision repair, and mismatch repair pathways (*Figs. 5a*

and 5e), reflecting adaptation to the persistently oxidative, metal-rich, and radiological

conditions maintained during prolonged neutral-pH ISL. The abundance of a key

overexpressed gene (*recA*) involved in homologous DNA repair[28], was moderately

correlated with gross ($\alpha + \beta$) radioactivity ($\rho = 0.52$, n = 32, $p < 0.05$) (*Fig. 5e*). Similarly,

U concentrations showed moderate associations with increased expression of *soxY*,

a gene involved in the sulfur oxidation (Sox) system ($\rho = 0.44$, n = 26, $p < 0.05$), as

well as with genes linked to ammonium oxidation, including *amoA* ($\rho = 0.43$, n = 8, p

< 0.05) and *amoB* ($\rho = 0.43$, n = 10, $p < 0.05$) (*Fig. 5e*). Functional genes associated

with iron, nitrogen, and sulfur redox transformations, processes that regulate electron

flow and thereby influence U speciation and mobility, were widely distributed across

100 strain-cluster rMAGs. These included *Cyc2* cluster genes mediating Fe(II)

oxidation in Gallionella spp. (potentially affecting U(IV) oxidation), *amoAB* genes for

ammonia oxidation in *Nitrosarchaeum* spp., and *soxY* genes for sulfur oxidation in

Pseudomonadota spp. (*Fig. 5e*). This functional consistency, potentially underpinned

by redundancy and convergent adaptation, points to environmental filtering as a key

mechanism structuring microbial communities under sustained geochemical stress[29–

33].

Figure 5 | Correlation and environmental drivers of overexpressed genes in metatranscriptomic datasets. *(a) Spearman correlation heatmap illustrates associations between hydrogeochemicals and the expression levels of overexpressed genes (defined as transcript coverage > 2× the average 30S coverage), detected in at least six samples across the dataset. Rows and columns are clustered by gene expression patterns. Color scale denotes Spearman correlation coefficients (red = positive, blue = negative), which the p values were adjusted using the BH method. The right-side bar plot indicates the number of strain-cluster rMAGs expressing each gene. Colored annotations indicate functional categories: oxidative stress response, DNA repair, carbon cycling, nitrogen cycling, sulfur cycling, iron cycling, oxygen metabolism, and hydrogeochemicals. Principal coordinate analysis (PCoA) of MAGs based on expression of DNA repair genes (b), oxidative stress response genes (c), genes involved in carbon, nitrogen, sulfur, and iron (C-N-S-Fe) cycling (d). Arrows indicate environmental variables significantly correlated with expression patterns. Points are colored and shaped by sample groups. (e) Heatmap showing presence or absence of overexpressed genes across strain-cluster rMAGs. Gene abundance (FPKM) is shown by color intensity, with taxa classified by phylum. The MAGs were clustered based on the presence of the functional genes.*

Uranium ISL-induced stress drives genetic diversification in subsurface aquifers

Genetic diversity and evolutionary pressures were assessed for species-cluster rMAGs by calculating non-synonymous to synonymous polymorphism (pN/pS) ratios, which indicate the balance between neutral variation and adaptive selection, and by appraising nucleotide diversity via PCoA to capture overall genetic variation among

populations. Only genes with greater than 5× coverage in both metagenomic and

metatranscriptomic datasets were included, and analyses focused on dereplicated

species-cluster rMAGs (95% ANI) containing more than three clustered members

(1,733 species-cluster rMAGs). This ensured robust comparisons across neutral-pH

ISL phases. PCoA revealed that, based on metagenomic and metatranscriptomic

data, both nucleotide diversity (Fig. 6g, i) and pN/pS ratios (Fig. 6h, j) were influenced

by groundwater geochemistry. Nucleotide diversity was positively associated with

bicarbonate, gross (α + β) radioactivity, sulfate, DOC, and Eh, and negatively with

sulfide and pH (*bioenv* correlation analysis R^2, 0.27). Similar patterns were observed

for pN/pS ratios, although with slightly reduced explanatory strength (*bioenv*

correlation analysis R^2, 0.31). These results indicate that U concentrations, redox

conditions, and radiochemical stressors all contribute to alterations in microbiome

genetic diversity and evolutionary trajectory during and following neutral-pH ISL.

Across metagenomic datasets, no significant differences were observed in

pN/pS ratios or nucleotide diversity during neutral-pH ISL. By contrast,

metatranscriptomic data revealed elevated pN/pS ratios at both mining-impacted and

post-mining sites, with median values increasing from 0.585 in S3a_BG to 0.608 in

S3a_MT05, and from 0.380 in S3b_BG to 0.503 in S3b_MT15 and 0.491 in S3b_PMS,

respectively (*Figs. 6b, c*). Notably, non-synonymous SNPs were detected in 481 genes

that were transcriptionally overexpressed and functionally linked to oxidative stress

responses, DNA repair, and cycling of carbon, nitrogen, sulfur, and iron (*Fig. S15b*),

suggesting that positive selection preferentially targeted actively transcribed genes

under stress. Notably, pN/pS ratios exceeding 1 were predominantly found in the

metatranscriptomic datasets, with 7 species-cluster rMAGs in BG, 47 species-cluster

rMAGs in MT, and 5 species-cluster rMAGs in PM, while only 5 species-cluster rMAGs

(four in MT and one in PM) exhibited pN/pS ratios exceeding 1 in both metagenomic

and metatranscriptomic datasets (*Fig. S16*). These results suggest that elevated

genetic variation during ISL is linked to transcriptionally active taxa responding to

combined radiochemical and geochemical perturbations, including oxidative stress

and the influx of new substrates such as oxygen from leaching solutions, rather than

a simple genome-wide increase in mutation rates. This framework also explains why

post-mining communities, despite residual radioactivity, exhibit reduced diversity as

redox conditions stabilize and substrate inputs diminish. The fact that no significant

changes were detected in the early mining phase (*Fig. 6a*) suggests that the

extent/duration of mining plays a major role in the process of natural selection.

However, it should be noted that only one sample (BG6 in S2 site) from the

background aquifer was available during the early mining phase, which may limit the

conclusions drawn from these data and underscores the need for further sampling to

better understand evolutionary dynamics at this stage.

Metatranscriptomic data revealed distinct evolutionary responses to neutral-pH

ISL across mining stages, marked by dynamic shifts in nucleotide diversity and

species-level selection. During the mid-mining phase (MT in S3a site), nucleotide

diversity significantly increased relative to both background and other mining stages

(*Fig. 6d–f*), coinciding with peak uranium and radioactivity levels. This suggests that

sustained radiochemical and oxidative stress promoted elevated genetic variation,

potentially by enabling the persistence of diverse genotypes capable of tolerating

extreme conditions. By contrast, the early and late mining phases exhibited reduced

nucleotide diversity, likely reflecting genetic bottlenecks driven by acute oxidative

perturbations and the selective loss of more vulnerable populations. Notably, post-

mining samples also showed diminished nucleotide diversity, despite the fact that both

TU and radioactivity levels remained elevated. These findings suggest that both the

intensity and duration of oxidative and radiochemical stress play a major role in

shaping genetic diversity in mining-impacted aquifers.

Figure 6 | pN/pS ratio and nucleotide diversity. *(a-c) Boxplots of pN/pS ratio from*

metagenomic (metaG) and metatranscriptomic (metaT) datasets spanning pre-mining,

mining, and post-mining phases; (d-f) Boxplots of nucleotide diversity from metaG and

metaT data across selected neutral-pH ISL stages; (g-j) PCoA plots showing the

influence of geochemical parameters on pN/pS ratio and nucleotide diversity derived

from metaG (g, h) and metaT (i, j) datasets.Statistical significance was assessed using

the Kruskal–Wallis test followed by Dunn's post‑hoc test with Benjamini–Hochberg

(BH) correction for multiple comparisons (*p < 0.05, **p < 0.01, and ***p < 0.001).

Differences in pN/pS ratios and nucleotide diversity between metaG and metaT for

each group were assessed using two‑sample t‑tests (*p < 0.05, **p < 0.01, and ***p <

0.001).

To better understand the genomic basis of adaptive responses, we identified 31 species-cluster rMAGs that exhibited consistent increases in nucleotide diversity from background (BG) to mining-impacted (MT) conditions (*Figs. 7a-b*). These rMAGs spanned diverse taxonomic groups, such as *Bacteroidales*, *Gallionella*, *Hyphomonas*, and *Kuenenia*, suggesting that the response to mining-related stress was phylogenetically widespread (*Fig. 7c*). Gene-level analysis showed that elevated nucleotide diversity was spread across many genes within each species-cluster rMAG, rather than being limited to a few specific regions, suggesting widespread mutational pressure under long-term environmental stress (*Fig. 7c*). Functional categorization of highly variable genomic regions showed enrichment in genes associated with information processing, regulatory networks, and cellular signaling, functions that are likely essential for coping with radiochemical and oxidative stress. Notably, the majority of these diversity increases were detected in metatranscriptomic data, reinforcing the interpretation that selection is preferentially acting on actively expressed genes. These patterns suggest that under prolonged mining stress, a

subset of microbial populations, those that remain transcriptionally active, exhibit

elevated genetic diversity, potentially as a bet-hedging strategy to enhance adaptive

flexibility. However, the absence of a community-wide increase in nucleotide diversity

suggests that evolutionary responses to ISL are concentrated in metabolically active,

redox-cycling taxa (e.g., iron-oxidizing Gallionella and nitrogen-oxidizing

Nitrosarchaeum), while many other lineages remain genetically stable or enter

dormancy under extreme stress. This indicates that anthropogenic disturbance does

not uniformly accelerate microbial evolution, but instead drives targeted adaptation in

functional groups central to biogeochemical cycling. This decoupling between

metagenomic and metatranscriptomic signals highlights that evolutionary responses

are concentrated within metabolically active fractions of the community, rather than

reflecting broad community-wide genomic shifts. Notably, the functional categories

enriched among highly variable regions, such as information processing, regulatory

systems, and cellular signaling, further suggest that selection is acting on genes

directly relevant to stress resilience. Taken together, these findings reveal elevated

diversity among transcriptionally active strains in mining-impacted aquifers,

underscoring how long-term ISL introduces radiochemical stress and geochemical

perturbations, including altered redox conditions and new electron donors and

acceptors, that collectively shape the active microbial fraction. While the overall

genomic potential of the community may remain conserved, sustained environmental

disturbance appears to activate and favor adaptive subpopulations, with responses

driven not only by stress tolerance but also by shifts in substrate availability and redox

gradients.

Figure 7 | Genome-wide nucleotide diversity patterns across mining stages in representative rMAGs. (a) Number of rMAGs with increased nucleotide diversity from background (BG) to mining-impacted (MT) conditions in metagenomic (metaG) and metatranscriptomic (metaT) datasets. (b) Ternary plot showing the distribution of rMAG-level nucleotide diversity across BG, MT, and post-mining (PM) samples, with points colored by dataset and sized by diversity magnitude. (c) Gene-level nucleotide diversity profiles for four representative rMAGs (*Bacteroidales*, *Hyphomonas*, *Kuenenia*, *Gallionella*), showing elevated and widespread diversity in MT, particularly

in transcriptionally active genes. Gene function distributions are indicated by functional

category bar plots.

Discussion

The global resurgence of nuclear energy has renewed interest in U ISL, a

minimally invasive yet chemically intensive extraction method poised to dominate

future uranium supply chains[34,35]. Given its operational efficiency and limited surface

disturbance, U ISL is expected to remain a dominant extraction method where

geological conditions permit in the near future[13]. While the geochemical and

biodiversity impacts of U ISL are well documented[13,17–21,23,36–38], its evolutionary

consequences for subsurface microbiomes remain underexplored. Here, we present

metagenomic and metatranscriptomic evidence that neutral-pH U ISL exerts sustained

selective pressure on microbial populations, driving *in situ* evolution in deep aquifers.

Neutral-pH ISL introduces persistent redox and geochemical gradients to

subsurface ecosystems by injecting oxidants into anoxic strata, mobilizing uranium,

radionuclides, and heavy metals. By examining a neutral-pH ISL area with aquifers

having undergone different time periods of leaching (up to 20-years), we reveal that

these sustained geochemical disturbances go beyond restructuring of microbial

community composition and alter transcriptional activity, along with an increased

genetic diversity illuminating long-term ecological and evolutionary consequences for

deep biosphere microbiomes. These transformations are shaped by sustained

selection pressures imposed by neutral-pH ISL, including oxidative stress, redox-

active metal toxicity, and low-dose radiation, stressors known to destabilize cellular

systems and promote adaptive traits linked to stress tolerance, DNA repair, and redox

regulation[39–41]. Correspondingly, we detected increased taxonomic diversity and

ecological modularity in neutral-pH ISL-impacted aquifers, signatures consistent with

stress-induced diversification[11,42–46]. Temporal succession patterns revealed dynamic

microbial responses, with oxidative stress-tolerant taxa dominating early neutral-pH

ISL stages and anaerobic fermenters and metal reducers emerging post-mining,

reflecting adaptation to evolving geochemical conditions.

Crucially, these ecological transitions were accompanied by molecular

signatures of adaptation. Genes associated with oxidative damage, heavy metal

toxicity, and radiation exposure were broadly overexpressed throughout all neutral-pH

ISL stages (*Figs. S8-S14*), suggesting a constitutive stress response. Yet, some

strains showed a substantial change in nucleotide diversification along with neutral U

ISL enabling the conclusion that repair mechanisms may have been insufficient to

counteract accumulated genomic damage. Consequently, elevated nucleotide

diversity, particularly in mid-mining metatranscriptomes, supports the hypothesis that

mutational input exceeded repair capacity, consistent with models of stress-induced

mutagenesis[47–50]. This model posits that environmentally triggered DNA damage,

coupled with error-prone repair, contributes to increased mutation rates. While DNA

repair pathways buffer deleterious mutations, they may also limit adaptive potential by

constraining the fixation of beneficial variants[51,52]. In our study, elevated nucleotide

diversity together with transcriptional activation of DNA repair and oxidative stress–

response genes indicates that adaptive evolution is concentrated in metabolically

active redox-cycling taxa rather than across the whole community. This resolves the

modest community-wide transcriptional changes yet strong gene-level positive

selection signals (Fig. S15b), highlighting that long-term ISL creates evolutionary

hotspots where radiochemical stress and altered substrate availability jointly promote

adaptation in key biogeochemical taxa.

These patterns are further supported by pN/pS ratios exceeding one, a

canonical signal of positive selection, detected predominantly in metatranscriptomes

(47 species-cluster rMAGs in mid-mining and 5 in post-mining), with only limited

overlap to respective metagenomes. Notably, the species-cluster rMAGs exhibiting

elevated nucleotide diversity spanned diverse taxa and showed widespread gene-

level variation in information processing and regulatory functions, underscoring that

selection acts primarily on transcriptionally active populations. This layered response,

genome-wide mutation accumulation coupled with transcription-linked selection,

highlights the dual role of environmental stress in both generating genetic variability

and filtering for functional adaptation, consistent with eco-evolutionary rescue

models[53–55], wherein microbial populations rebound from stress-induced bottlenecks

through the acquisition and retention of adaptive mutations.

Our study provides rare empirical evidence that stress-induced mutagenesis

and transcription-linked selection operate in natural deep subsurface microbiomes

over decadal timescales. Although such mechanisms are well characterized in

laboratory systems[56–62] and diverse surface environments[47,63–66], their manifestation in

oligotrophic, energy-limited subsurface ecosystems[67–70] has remained largely

speculative due to assumptions of metabolic quiescence and evolutionary stasis[71–74].

Our findings challenge this paradigm, demonstrating that chronic anthropogenic stress

in engineered aquifers can generate sustained mutational input and facilitate

directional selection. The convergence of elevated nucleotide diversity, DNA repair

activation, and positive selection in expressed genes reveals a coordinated

evolutionary response to long-term environmental disturbance. These observations

extend core theoretical frameworks, including stress-induced mutagenesis and eco-

evolutionary rescue, from controlled systems to complex, anthropogenically modified

subsurface environments, redefining the deep biosphere as a dynamic, evolutionarily

responsive system.

MATERIAL AND METHODS

Study area, sampling strategy, and sample collection. The study area located in

the Qianjiadian U ore deposit within the QJD sag of the Kailu Depression in the

southwestern Songliao Basin, northeast China, represents a typical sandstone-hosted

uranium system[75]. The stratigraphy comprises the Qingshankou (K_2qn), Yaojia (K_2y),

and Nenjiang (K_2n) formations, overlain by Neoproterozoic and Quaternary alluvial

deposits. The uranium mineralization is primarily hosted in the K_2y Formation, which

consists of red, yellow, gray, and mineralized gray sandstones, with widespread

carbonate cementation (dolomite, siderite, ankerite, and calcite). Pitchblende and

coffinite are the dominant uranium minerals, while Fe-Ti oxides, natural organic matter

(NOM), clay minerals, and pyrite serve as additional uranium hosts[36,76–78].

Groundwater flow follows a southwest-to-northeast trajectory within a confined

Cretaceous sandstone aquifer (~298 m depth, 35 m thickness), with permeability

coefficients averaging 0.145 m/day[21]. The aquifer is hydraulically isolated by a low-

permeability mudstone-dominated aquitard, with limited connectivity to adjacent

hydrogeological units.

The Qianjiadian uranium deposit has been a key site for advancing in-situ recovery

(ISR) technology in China, serving as the first large-scale demonstration of CO_2 + O_2

leaching in a deep, low-permeability sandstone-hosted system. During the neutral-pH

ISL process, the mixed reagent of dissolved O_2 (oxidant) and CO_2 (complexing agent)

was injected into the ore-bearing aquifer, and subsequently, U-containing groundwater

was pumped out and U was retrieved using ion exchange resins[21,79]. The effluent from

the reverse osmosis system was reused as a leaching solution after being mixed with

CO_2 and O_2 to maintain carbonate buffering and oxidizing conditions[80]. The orebody,

located within confined sandstone aquifers at depths ranging from 220 to 460 m,

presents complex hydrogeological conditions that influence solute transport, aquifer

reactivity, and microbial community dynamics. The unique geological and

hydrogeochemical setting of the Qianjiadian deposit, combined with its prolonged

history of in situ leaching (ISR) operations, establishes it as a natural laboratory for

investigating subsurface fluid-rock-microbe interactions, uranium mobilization

processes, and the long-term geochemical and microbial responses to neutral-pH ISL

as well as post-mining restoration.

To investigate the hydrogeochemical and microbial responses to neutral-pH ISL

mining, groundwater samples were collected from the Qianjiadian (QJD) sandstone-

hosted uranium deposit during two field campaigns in October-November 2021 and

August–October 2022 (Figure 1). A total of 80 groundwater samples were obtained

from sites representing different stages of neutral-pH ISL, spanning ore-bearing

aquifers at depths of 220–460 m. Sampling locations encompassed neutral-pH ISL

sites spanning multiple operational durations: 0.25 years (MT0.25, 8 samples), 1 year

(MT1, 3 samples), 4 years (MT4, 6 samples), 5 years (MT5, 7 samples), 6 years (MT6,

6 samples), 15 years (MT15, 6 samples), and 20 years (MT20, 4 samples). Post-

mining phases were sampled under distinct redox conditions, including iron-reducing

(PMI, 4 samples), mixed iron- and sulfate-reducing (PMIS, 2 samples), and sulfate-

reducing (PMS, 3 samples) conditions. Background aquifer samples (BG, 19 total)

were collected across the chronosequence, including non-mining areas (5 samples),

early-stage sites (MT0.25 and MT1, 1 sample), mid-stage sites (MT4 and MT5, 5

samples), and late-stage/post-mining sites (MT15, MT20, and PM, 8 samples).

Background wells were present at all sites and sampled concurrently with mining and

post-mining wells, providing site-specific baseline controls. Detailed

hydrogeochemical data for MT20, MT4, MT5, PMI, PMIS, PMS, and BGPM are

reported in our previous studies[21–23,36].

Prior to sample collection using a deep groundwater sampler (Geosub 2, Geotech,

USA), monitoring and post-mining wells were pumped for at least 20 minutes until

water temperature, electrical conductivity (EC), and oxidation-reduction potential

(ORP) stabilized. Groundwater was then pumped after well flushing and approximately

45 ± 0.5 L of water was filtered through 0.2-µm sterile cellulose nitrate membrane

filters (Sartorius, Göttingen, Germany) to capture microbial organisms from each

groundwater well. All filtered membranes were frozen at -80 °C for high-throughput

sequencing. Groundwater samples for physicochemical analysis were collected in 5 L

sterile PET bottles and stored at -20 °C.

Chemical Analysis. Major anions (Cl^-, NO_3^-, and SO_4^{2-}) were measured using ion

chromatography (Dionex ICS-600, Thermo Fisher Scientific, Massachusetts, USA),

while major cations (Na^+, K^+, Mg^{2+}, and Ca^{2+}) were determined using inductively

coupled plasma optical emission spectrometry (ICP-OES, Optima 8300, PerkinElmer,

Massachusetts, USA). Trace elements, including U, Fe, and other transition metals,

were quantified using inductively coupled plasma mass spectrometry (ICP-MS, 7900

ICP-MS, Agilent Technologies, California, USA). Alkalinity was measured by Gran

titration using a Metrohm 888 Titrando (Metrohm, Herisau, Switzerland). The total

organic carbon (TOC) and dissolved organic carbon (DOC) concentrations were

determined using a TOC analyzer (TOC-L, Shimadzu, Kyoto, Japan). The gross alpha

and gross beta radioactivity of groundwater samples were measured using a low-

background α/β meter (LB6008, Beijing Yida Measurement Technology Co., Ltd.,

Beijing, China) following the thick source method for radioactivity analysis.

Measurements of gross alpha and gross beta radioactivity were conducted at the

China Nuclear Mining Science and Technology Corporation, Beijing, China, to assess

radiological characteristics associated with uranium mobilization.

DNA extraction and sequencing. Total metagenomic DNA was extracted from

biomass retained on 0.22-µm sterile cellulose nitrate membrane from 60 groundwater

samples, selected from a total of 80 samples retrieved from different neutral-pH ISL

stages in ore-bearing aquifers (see above and *Figure 1*). DNA extraction was

performed using the FastDNA® SPIN Kit for Soil (MP Biomedicals, Solon, OH, USA)

following the manufacturer's protocol. The concentration and purity of the DNA

extracts were measured using a TBS-380 fluorometer (Turner Biosystem, USA) and a

NanoDrop 2000 spectrophotometer (Thermo Scientific, USA), respectively. The

quality of the DNA extracts was checked using 1%-agarose gel electrophoresis.

Paired-end sequencing was performed on the Illumina HiSeq XTen platform (Illumina,

Inc., San Diego, CA, USA) at Majorbio Bio-Pharm Technology Co., Ltd., China. DNA

extracts were fragmented to an average size of approximately 400 bp using a Covaris

M220 ultrasonicator (Gene Company Limited, China). Paired-end libraries were

constructed using the NEXTFLEX Rapid DNA-Seq Kit (Bioo Scientific, Austin, TX,

USA), with adapters containing the full complement of sequencing primer hybridization

sites ligated to the blunt end of fragments.

Metagenome processing and MAG recovery. Raw reads were trimmed and quality-

filtered using fastp v0.19.7[81]. The microbial diversity and its coverage based on quality-

filtered raw metagenomic reads was estimated by Nonpareil v3.4.1[25] in "kmer" mode.

High-quality reads from individual samples were assembled into scaffolds using

metaSPAdes version 3.15.5 (--k-list 21, 33, 55, 77, 99, and 127)[82]. Open-Reading

Frames on scaffolds were predicted using prodigal (v2.6.3)[83] in "meta" mode. MAGs

were recovered from each assembly using the binning module (parameters: -metabat2

-maxbin2 -concoct) within metaWRAP v1.3 (metabat2 v2.12.1, MaxBin2 v.2.2.6, and

concoct v0.4.0)[84–87], which bins based on reads mapped only from the corresponding

sample. Quality of genome bins (completeness and contamination) was evaluated

using CheckM2 (v2.0.11)[88]. The best MAG representative for each sample was

determined using DASTool v1.1.7 with default parameters[89]. Finally, a total of 4036

MAGs were obtained with completeness >50% and contamination <10%. Bins were

further consolidated via dereplication at strain level using dRep (v.3.2.2) at 99% ANI[90]

resulting in 2,255 strain-cluster representative MAGs (strain-cluster rMAGs).

Taxonomic classification of strain-cluster rMAGs and phylogenetic tree

generation. Taxonomies were assigned to each strain-cluster rMAG using GTDB-Tk

v2.1.1 (https://github.com/Ecogenomics/GtdbTk, database release 220)[91]. Conserved

marker genes (ribosomal proteins and RNA polymerase subunits) were identified

using MarkerFinder (-m ribo_rnap, -c) (https://github.com/faylward/markerfinder) and

concatenated into a supermatrix, which were used for constructing a maximum-

likelihood phylogenetic tree via IQ-TREE 2 (v 2.2.0.3)[92] with 1,000 ultrafast

bootstrapping iterations. ModelFinder[93] determined Qpfam+I+I+R10 to be the best

model. The phylogenetic tree was visualized using the Interactive Tree of Life (iTOL)[94].

Coverage estimation and metabolic potential prediction. Clean reads from each

sample were mapped to strain-cluster rMAGs using strobealign (v0.13.0)[95] in CoverM

(v0.7.0; default parameters)[96]. The metabolic potential of strain-cluster rMAGs was

determined using METABOLIC v4.0[97]. We constructed two databases, one containing

96 genes associated with DNA repair mechanisms and one containing 32 genes

implicated in microbial responses to oxidative stress, with all genes being sourced

from NCBI (https://www.ncbi.nlm.nih.gov/). Genes of strain-cluster rMAGs were

annotated against these databases using DIAMOND BLAST[98] with an E-value

threshold of $1e^{-5}$ and the parameter -k 1 to retain only the top scoring hit. To ensure

the specificity of these annotations, a reverse BLAST was subsequently conducted by

aligning database hits back against the original amino acid sequences of the

corresponding strain-cluster rMAGs using the same E-value threshold. Only reciprocal

hits that aligned to the same ORFs as the original query were retained, ensuring one-

to-one correspondence. Cases with multiple high-scoring matches were excluded to

avoid ambiguous annotations. Final hits were filtered based on a similarity threshold

of >50%, ensuring high-confidence functional assignments. These curated datasets

facilitated a refined analysis of microbial resilience and adaptation strategies in diverse

environmental contexts.

Community profiling based on the *rpS3* marker gene. ORFs representing the *rpS3*

marker gene were identified using hmmsearch[99] against custom Hidden Markov

Models (HMMs) for Archaea and Bacteria (E-value 1e-28;

https://github.com/AJProbst/rpS3_trckr). Sequences were pooled and clustered at

99% similarity using vsearch[100], representing species-level clusters[101]. Representative

sequences were selected following a tiered approach: (1) centroids that could be

extended by 1000 bp in both directions, (2) non-centroids that were similarly

extendable, and (3) the longest available sequence when neither of the first two criteria

were met. To improve abundance estimates, *rpS3* gene sequences of representative

rpS3 genes were extended by up to 1,000 bp in both directions on their scaffold of

origin if possible. This was done as mapping on just the approximately 1,000 bp long

rpS3 gene results in poor mapping on the gene edges and this has progressively more

impact when the total length of the sequence is short. The resulting *rpS3* gene

sequences, ideally extended in both directions by 1,000 bp, are henceforth called

rpS3-extended. The workflow for clustering, extracting, and calculating coverage is

available   at   https://github.com/ProbstLab/publication-

associated_scripts/tree/main/FigueroaGonzalez_Bornemann_etal.   The rpS3 genes

in rpS3-extended sequences (in amino acid format) were taxonomically classified

using DIAMOND[98] (E-value ≤ 1e^{-5}) against the GTDB r220 protein database,

assigning the best hit to be the taxonomy. If no hit meeting the search criteria was

found, "Unclassified" was assigned. Abundance was calculated by mapping reads of

all metagenomic samples to rpS3-extended sequences and calculating the average

coverage per sample. Additionally, the breadth, i.e., the number of nucleotide positions

in rpS3-extended having a coverage of at least 1, was determined. The coverage of

an rpS3-extended sequence in a given sample was set to zero if the breadth of the

sequence in that sample was less than 95%, indicating spurious mapping instead of

actual presence of the rpS3-extended-represented species. Relative coverage values

were transformed into percentages (%).

RNA extraction, sequencing, and metatranscriptomics. Total RNA was isolated

and purified using Soil RNA Extraction Kit (Majorbio, Shanghai, China) following the

manufacturer's procedure. The integrity and quantity of the extracted RNA were

measured with a NanoDrop 2000 spectrophotometer (Thermo Scientific, MA, USA)

and an Agilent 5300 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA). The RNA

was subjected to standard Illumina library preparation with an Illumina® Stranded

mRNA Prep, Ligation (Illumina, San Diego, CA, USA), and rRNA was depleted using

a RiboCop rRNA Depletion Kit for Mixed Bacterial Samples (Lexogen, USA). Depleted

RNA was extracted from biomass retained on 0.22 µm sterile cellulose nitrate

membranes from 60 groundwater samples (same to metagenomic groundwater

samples) and finally library preparation of RNA extraction from 45 groundwater

samples were successfully constructed (Figure 1). Sequencing was performed on an

Illumina Novaseq6000 sequencer (Illumina, San Diego, CA, USA) with paired-end

sequencing method at Majorbio Bio-Pharm Technology Co., Ltd. (Shanghai, China).

Quality control of the raw data generated by sequencing was carried out with fastp

v0.19.7[81]. SortMeRNA (v4.3.6) was used to remove non-coding RNA sequences

(tRNA, tmRNA, 5S, 16S, 18S, 23S, and 28S)[102]. The proportion of reads removed per

sample is summarized in Supplementary Table S3 and Figure S17, providing an

overview of rRNA/tRNA content prior to filtering.

Gene overexpression in each metatranscriptome. InStrain (v1.8.1)[103] with the

parameter (--min_cov 0) was used on strain-cluster rMAGs recovered from the same

sample to get the coverage of each gene using the clean reads from each

metagenome and each metatranscriptome. The gene coverage in each strain-cluster

rMAG was received by using the cluster database generated when dereplicate

genomes at 99% ANI using dRep (v3.2.2)[90]. The mean coverage of 30S ribosomal

proteins were used to calculate the normalization factor of each gene from the strain-

cluster rMAGs in each metatranscriptome sample. The genes with normalization factor

greater than two were defined as overexpressed genes.

Species-level clustering and calculation of evolutionary metrics. Filtered reads

from each metagenome and metatranscriptome were mapped to the best MAGs

output from DAStool of the same sample using strobealign (v0.13.0)[95] in CoverM

(v0.7.0; default parameters)[96]. Genetic annotation of MAGs was performed with

Prodigal (v2.6.3; –p meta)[83] for the gene module of inStrain (v1.8.1)[103]. Population

statistics and nucleotide metrics, including nucleotide diversity (SNVs/kbp) and

nonsynonymous to synonymous mutation ratios (pN/pS) were calculated from these

mappings using the profile module of the inStrain program (v1.8.1; –database mode;

default parameters)[103] at genome and gene levels. The 4036 MAGs reconstructed

from all individual assemblies (n = 60) were combined and dereplicated for species-

level clustering using dRep (v3.2.2)[90] at 95% ANI. A total of 1733 MAGs with the

highest genome quality from each species cluster were designated as the

representative species (species-cluster rMAGs). The species-cluster rMAGs with over

three members in the cluster database generated when dereplicate genomes at 95%

ANI were selected for the final calculation of microdiversity metrics such as coverage,

breadth, SNPs density, nucleotide diversity, and pN/pS ratios.

For SNP calling, the number of quality-filtered reads mapping to the position had to be

at least 5× coverage and 5% SNP frequency, and the variant based on reads had to

be lower than the expected sequencing error rate ($1e^{-6}$). Genome-level evolutionary

metrics based on both metagenomic and metatranscriptomic data from mining sites

were compared with those from background aquifer samples at the same location

(*Table S4, S5*). To evaluate the impact of radioactivity and chemical stress on

microbial genomes, 41 species rMAGs present in at least two groups (BG, MT, and

PM) and in a minimum of three samples from each group were selected. Gene

annotations for these 41 species rMAGs were performed (*Table S6*) using

MetaCerberus v1.3.1[104]. Gene-level nucleotide diversity of microbial genes within

these 41 rMAGs was then analyzed using both metagenomic and metatranscriptomic

data across the BG, MT, and PM groups (*Table S7,S8*).

Statistical analyses and data visualization. Spearman correlation analyses were

performed in the R platform (http://cran.r-project.org), with p-values adjusted using the

Benjamini-Hochberg procedure at a minimum significance threshold of 0.05[105]. Co-

occurrence networks were constructed using the *igraph* package with default

parameters[106]. Only Spearman correlations with p-values < 0.001 and $|\rho| > 0.6$

between pairs of rMAGs were considered statistically significant, contributing to the

construction of a reliable network. The resulting correlation network was imported into

Gephi (v0.9.2) for visualization[107]. The Spearman correlations between over-

expressed genes from rMAGs and hydrogeochemical variables were visualized using

the *pheatmap* package in R (v4.4.2)[108]. To assess sample similarity and identify key

factors driving clustering patterns, Principal Coordinates Analysis (PCoA) was

conducted based on the Bray-Curtis dissimilarity of microbial community

compositions. Analysis of Similarity (ANOSIM) was employed to evaluate significant

differences among sample groups.

ACKNOWLEDGEMENTS

This study was supported by the National Natural Science Foundation of China grant

nos. 42130509 (to H.G.) and 42072273 (to W.X.), the Deep Earth Probe and Mineral

Resources Exploration–National Science and Technology Major Project grant no.

2024ZD1000601 (to W.X.), the Fundamental Research Funds for the Central

Universities grant no. 590223006 (to W.X.) and the 111 Project grant no. B20010 (to

H.G.); and by the European Research Council Synergy Grant "Archean Park" under

grant no. 101118631 (to A.J.P.) and the German Research Foundation grant no. PR

1603/4-1 (to A.J.P.). We thank Prof. Rizlan Bernier-Latmani (Environmental

Microbiology Laboratory, École Polytechnique Fédérale de Lausanne, Switzerland) for

valuable comments on the manuscript and for providing access to laboratory

resources.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

Ethics Statement

All sampling and sample collection were performed with permission from the local

environmental authorities and in compliance with all applicable regulations governing

environmental research in China. No human subjects or animal samples were involved

in this study.

CODE AVAILABILITY STATEMENT

All code used in this manuscript is publicly available and referenced in the MAterials

and Methods section.

References:

1.   Xie, X. *et al.* Groundwater Quality and Public Health. *Annual Review of Environment and Resources* **48**, 395–418 (2023).

2.   Orsi, W. D., Edgcomb, V. P., Christman, G. D. & Biddle, J. F. Gene expression in the deep biosphere. *Nature* **499**, 205–208 (2013).

3.   Magnabosco, C. *et al.* The biomass and biodiversity of the continental subsurface. *Nature Geoscience* **11**, 707–717 (2018).

4.   Beaver, R. C. & Neufeld, J. D. Microbial ecology of the deep terrestrial subsurface. *The ISME Journal* **18**, wrae091 (2024).

5.   Bornemann, T. L. V. *et al.* Genetic diversity in terrestrial subsurface ecosystems impacted by geological degassing. *Nature Communications* **13**, 1–12 (2022).

6.   Patsis, A. C., Schuler, C. J., Toner, B. M., Santelli, C. M. & Sheik, C. S. The potential for coupled organic and inorganic sulfur cycles across the terrestrial deep subsurface biosphere. *Nat Commun* **16**, 3827 (2025).

781 7. Probst, A. J. *et al.* Differential depth distribution of microbial function and putative

782   symbionts through sediment-hosted aquifers in the deep terrestrial subsurface.

783   *Nature Microbiology* **3**, 328–336 (2018).

784 8. He, C. *et al.* Genome-resolved metagenomics reveals site-specific diversity of

785   episymbiotic CPR bacteria and DPANN archaea in groundwater ecosystems.

786   *Nature Microbiology* **6**, 354–365 (2021).

787 9. Amundson, K. K., Borton, M. A. & Wilkins, M. J. Anthropogenic impacts on the

788   terrestrial subsurface biosphere. *Nat Rev Microbiol* **23**, 147–161 (2025).

789 10. Vigneron, A. *et al.* Succession in the petroleum reservoir microbiome through an

790   oil field production lifecycle. *The ISME Journal* **11**, 2141–2154 (2017).

791 11. Ning, D. *et al.* Environmental stress mediates groundwater microbial community

792   assembly. *Nature Microbiology* **9**, 490–501 (2024).

793 12. Jones, D. O. B. *et al.* Long-term impact and biological recovery in a deep-sea

794   mining track. *Nature* (2025) doi:10.1038/s41586-025-08921-3.

795 13. Li, G. & Yao, J. A Review of In Situ Leaching (ISL) for Uranium Mining. *Mining* **4**,

796   120–148 (2024).

797 14. Jroundi, F. *et al.* Profiling native aquifer bacteria in a uranium roll-front deposit

798   and their role in biogeochemical cycle dynamics: Insights regarding in situ

799   recovery mining. *Science of the Total Environment* **721**, (2020).

800 15. Vrionis, H. A. *et al.* Microbiological and geochemical heterogeneity in an in situ

801   uranium bioremediation field site. *Applied and Environmental Microbiology* **71**,

802   6308–6318 (2005).

803 16. Chen, H. *et al.* Uranium contamination mediating soil and ore microbial

804   community assembly at four mining sites, South China. *Front. Microbiol.* **16**,

805   1553072 (2025).

17. Zhu, F., Zhao, B., Min, W. & Li, J. Characteristics of groundwater microbial

communities and the correlation with the environmental factors in a

decommissioned acid in-situ uranium mine. *Front. Microbiol.* **13**, 1078393 (2023).

18. Coral, T. *et al.* Microbial communities associated with uranium in-situ recovery

mining process are related to acid mine drainage assemblages. *Science of The*

Total Environment **628–629**, 26–35 (2018).

19. Van Nostrand, J. D. *et al.* Dynamics of Microbial Community Composition and

Function during In Situ Bioremediation of a Uranium-Contaminated Aquifer. *Appl*

Environ Microbiol **77**, 3860–3869 (2011).

20. Li, B. *et al.* Bacterial Community Shift and Coexisting/Coexcluding Patterns

Revealed by Network Analysis in a Uranium-Contaminated Site after Bioreduction

Followed by Reoxidation. *Applied and Environmental Microbiology* **84**, e02885-

17 (2018).

21. Lu, C. *et al.* Multi-Isotope Based Identification and Quantification of Oxygen

Consuming Processes in Uranium Hosting Aquifers With $CO_2 + O_2$ In Situ

Leaching. *Water Resources Research* **59**, e2022WR033980 (2023).

22. Lian, G. *et al.* Genome-Resolved Metagenomic Insights into Aquifer Microbiome

Adaptations Following Two Decades of Neutral In-Situ Leaching. *Authorea*

Preprints (2024).

23. Lu, C. *et al.* Natural Attenuation of Groundwater Uranium in Post-Neutral-Mining

Sites Evidenced from Multiple Isotopes and Dissolved Organic Matter. *Environ.*

Sci. Technol. **58**, 12674–12684 (2024).

24. Wang, Q. *et al.* Mineral Composition and Full-Scale Pore Structure of Qianjiadian

Sandstone-Type Uranium Deposits: Application for In Situ Leaching Mining.

Geofluids **2022**, 1–15 (2022).

25. Rodriguez-R, L. M., Gunturu, S., Tiedje, J. M., Cole, J. R. & Konstantinidis, K. T.

Nonpareil 3: Fast Estimation of Metagenomic Coverage and Sequence Diversity.

mSystems **3**, e00039-18 (2018).

26. Sharon, I. *et al.* Accurate, multi-kb reads resolve complex populations and detect

rare microorganisms. *Genome Res.* **25**, 534–543 (2015).

27. Bowers, R. M. *et al.* Minimum information about a single amplified genome

(MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and

archaea. *Nature Biotechnology* **35**, 725–731 (2017).

28. Lalande, E. & El Sayyed, H. Break-ups and make-ups: DNA search and repair.

Nature Reviews Microbiology **20**, 66–66 (2022).

29. Louca, S. *et al.* Function and functional redundancy in microbial systems. *Nature*

Ecology and Evolution **2**, 936–943 (2018).

30. Allison, S. D. & Martiny, J. B. H. Resistance, resilience, and redundancy in

microbial communities. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 11512–11519 (2008).

31. Hanson, C. A., Fuhrman, J. A., Horner-Devine, M. C. & Martiny, J. B. H. Beyond

biogeographic patterns: processes shaping the microbial landscape. *Nature*

Reviews Microbiology **10**, 497–506 (2012).

32. Krause, S. *et al.* Trait-based approaches for understanding microbial biodiversity

and ecosystem functioning. *Front. Microbiol.* **5**, (2014).

33. Yang, Y. Emerging Patterns of Microbial Functional Traits. *Trends in Microbiology*

29, 874–882 (2021).

34. *Energy, Electricity and Nuclear Power Estimates for the Period up to 2050.*

(INTERNATIONAL ATOMIC ENERGY AGENCY, Vienna, 2024).

35. Monnet, A., Gabriel, S. & Percebois, J. Long-term availability of global uranium

resources. *Resources Policy* **53**, 394–407 (2017).

36. Lu, C. *et al.* Characteristics of Dissolved Organic Matter in Uranium Hosting

Aquifers and Potential Molecular Transformation During Neutral In Situ Leaching.

JGR Biogeosciences **129**, e2023JG007851 (2024).

37. Asghar, F. *et al.* Geochemical characteristics and uranium neutral leaching

through a co2 + o2 system—an example from uranium ore of the elzpa ore

deposit in Pakistan. *Metals* **10**, 1–19 (2020).

38. Ruiz, O., Thomson, B., Cerrato, J. M. & Rodriguez-Freire, L. Groundwater

restoration following in-situ recovery (ISR) mining of uranium. *Applied*

Geochemistry **109**, 104418 (2019).

39. Wani, A. K., Akhtar, N., Sher, F., Navarrete, A. A. & Américo-Pinheiro, J. H. P.

Microbial adaptation to different environmental conditions: molecular perspective

of evolved genetic and cellular systems. *Arch Microbiol* **204**, 144 (2022).

40. Newsome, L., Morris, K. & Lloyd, J. R. The biogeochemistry and bioremediation

of uranium and other priority radionuclides. *Chemical Geology* **363**, 164–184

(2014).

41. Carvalho, F. *et al.* The environmental behaviour of uranium. *International Atomic*

Energy   Agency   (2023)   doi:Retrieved   from:

https://www.iaea.org/publications/14688/the-environmental-behaviour-of-

uranium.

42. Da Cruz Nizer, W. S. *et al.* Oxidative stress responses in biofilms. *Biofilm* **7**,

100203 (2024).

43. Nguyen, J., Lara-Gutiérrez, J. & Stocker, R. Environmental fluctuations and their

effects on microbial communities, populations and individuals. *FEMS*

Microbiology Reviews **45**, fuaa068 (2021).

44. Galhardo, R. S., Hastings ,P. J. & and Rosenberg, S. M. Mutation as a Stress

Response and the Regulation of Evolvability. *Critical Reviews in Biochemistry*

and Molecular Biology **42**, 399–435 (2007).

45. Evans, S., Martiny, J. B. H. & Allison, S. D. Effects of dispersal and selection on

stochastic assembly in microbial communities. *The ISME Journal* **11**, 176–185

(2017).

46. Zhou, J. *et al.* Stochasticity, succession, and environmental perturbations in a

fluidic ecosystem. *Proc. Natl. Acad. Sci. U.S.A.* **111**, (2014).

47. MacLean, R. C., Torres-Barceló, C. & Moxon, R. Evaluating evolutionary models

of stress-induced mutagenesis in bacteria. *Nat Rev Genet* **14**, 221–227 (2013).

48. Mojica, E. A. & Kültz, D. Physiological mechanisms of stress-induced evolution.

Journal of Experimental Biology **225**, jeb243264 (2022).

49. Pal, A., Bhattacharjee, S., Saha, J., Sarkar, M. & Mandal, P. Bacterial survival

strategies and responses under heavy metal stress: a comprehensive overview.

Critical Reviews in Microbiology **48**, 327–355 (2022).

50. Couce, A. *et al.* Changing fitness effects of mutations through long-term bacterial

evolution. *Science* **383**, eadd1417 (2024).

51. Denamur, E. & Matic, I. Evolution of mutation rates in bacteria. *Molecular*

Microbiology **60**, 820–827 (2006).

52. Maharjan, R. P. *et al.* Mutation accumulation and fitness in mutator

subpopulations of *Escherichia coli*. *Biol. Lett.* **9**, 20120961 (2013).

53. Custer, G. F., Bresciani, L. & Dini-Andreote, F. Toward an integrative framework

for microbial community coalescence. *Trends in Microbiology* **32**, 241–251

(2024).

54. Shade, A. Microbiome rescue: directing resilience of environmental microbial

communities. *Current Opinion in Microbiology* **72**, 102263 (2023).

55. Yamamichi, M., Letten, A. D. & Schreiber, S. J. Eco-evolutionary maintenance of

diversity in fluctuating environments. *Ecology Letters* **26**, (2023).

56. Jinks-Robertson, S. & Bhagwat, A. S. Transcription-Associated Mutagenesis.

Annu. Rev. Genet. **48**, 341–359 (2014).

57. Tenaillon, O. *et al.* Tempo and mode of genome evolution in a 50,000-generation

experiment. *Nature* **536**, 165–170 (2016).

58. Wei, W. *et al.* Rapid evolution of mutation rate and spectrum in response to

environmental and population-genetic challenges. *Nat Commun* **13**, 4752 (2022).

59. Ferrare, J. T. & Good, B. H. Evolution of evolvability in rapidly adapting

populations. *Nat Ecol Evol* **8**, 2085–2096 (2024).

60. Bruckbauer, S. T. & Cox, M. M. Experimental evolution of extremophile resistance

to ionizing radiation. *Trends in Genetics* **37**, 830–845 (2021).

61. Daly, M. J. A new perspective on radiation resistance based on Deinococcus

radiodurans. *Nature Reviews Microbiology* **7**, 237–245 (2009).

62. Pavlopoulou, A. *et al.* Unraveling the mechanisms of extreme radioresistance in

prokaryotes: Lessons from nature. *Mutation Research/Reviews in Mutation*

Research **767**, 92–107 (2016).

63. Martincorena, I., Seshasayee, A. S. N. & Luscombe, N. M. Evidence of non-

random mutation rates suggests an evolutionary risk management strategy.

Nature **485**, 95–98 (2012).

64. Ferenci, T. Irregularities in genetic variation and mutation rates with

environmental stresses. *Environmental Microbiology* **21**, 3979–3988 (2019).

65. Foster, P. L. Stress-Induced Mutagenesis in Bacteria. *Critical Reviews in*

Biochemistry and Molecular Biology **42**, 373–397 (2007).

66. Woodcroft, B. J. *et al.* Genome-centric view of carbon processing in thawing

permafrost. *Nature* **560**, 49–54 (2018).

67. Mehrshad, M. *et al.* Energy efficiency and biological interactions define the core microbiome of deep oligotrophic groundwater. *Nature Communications* **12**, 1–12 (2021).

68. Jones, R. M., Goordial, J. M. & Orcutt, B. N. Low Energy Subsurface Environments as Extraterrestrial Analogs. *Front. Microbiol.* **9**, 1605 (2018).

69. Hoehler, T. M. & Jørgensen, B. B. Microbial life under extreme energy limitation. *Nature Reviews Microbiology* **11**, 83–94 (2013).

70. Orsi, W. D. Ecology and evolution of seafloor and subseafloor microbial communities. *Nature Reviews Microbiology* **16**, 671–683 (2018).

71. Jørgensen, B. B. & Marshall, I. P. G. Slow Microbial Life in the Seabed. *Annu. Rev. Mar. Sci.* **8**, 311–332 (2016).

72. Becraft, E. D. *et al.* Evolutionary stasis of a deep subsurface microbial lineage. *The ISME Journal* **15**, 2830–2842 (2021).

73. Garber, A. I., Ramírez, G. A. & D'Hondt, S. Genomic stasis over millions of years in subseafloor sediment. *Environmental Microbiology* **26**, e16674 (2024).

74. Schopf, J. W. *et al.* Sulfur-cycling fossil bacteria from the 1.8-Ga Duck Creek Formation provide promising evidence of evolution's null hypothesis. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 2087–2092 (2015).

75. Jia, J. *et al.* Mineralogy and geochemistry of carbonate cement in sandstone and implications for mineralization of the Qianjiadian sandstone-hosted uranium deposit, southern Songliao Basin, China. *Ore Geology Reviews* **123**, 103590 (2020).

76. Fan, Y. *et al.* The behavior of pyrite during in-situ leaching of uranium by CO2 + O2: A case study of the Qianjiadian uranium deposit in the Songliao Basin,

northeastern China. *Ore Geology Reviews* **169**, 106085 (2024).

77. Rong, H. *et al.* Effects of igneous intrusions on diagenesis and reservoir quality of sandstone in the Songliao Basin, China. *Marine and Petroleum Geology* **127**, 104980 (2021).

78. Rong, H. *et al.* Effects of basic intrusions on REE mobility of sandstones and their geological significance: A case study from the Qianjiadian sandstone-hosted uranium deposit in the Songliao Basin. *Applied Geochemistry* **120**, 104665 (2020).

79. Qiu, W. *et al.* What chemical reaction dominates the CO2 and O2 in-situ uranium leaching? Insights from a three-dimensional multicomponent reactive transport model at the field scale. *Applied Geochemistry* **148**, 105522 (2023).

80. Xu, L., Wang, H., Liu, N. & Du, Z. Waste Water Treatment of CO2 + O2 in-situ Leaching Uranium. *Uranium Mining and Metallurgy* **31**, 96–99 (2012).

81. Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

82. Bankevich, A. *et al.* SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* **19**, 455–477 (2012).

83. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**, 119 (2010).

84. Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nature Methods* **11**, 1144–1146 (2014).

85. Kang, D. D., Froula, J., Egan, R. & Wang, Z. MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ* **2015**, 1–15 (2015).

86. Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for
genome-resolved metagenomic data analysis. *Microbiome* **6**, 1–13 (2018).

87. Wu, Y. W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: An automated binning
algorithm to recover genomes from multiple metagenomic datasets.
Bioinformatics **32**, 605–607 (2016).

88. Chklovski, A., Parks, D. H., Woodcroft, B. J. & Tyson, G. W. CheckM2: a rapid,
scalable and accurate tool for assessing microbial genome quality using machine
learning. *Nat Methods* **20**, 1203–1212 (2023).

89. Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a
dereplication, aggregation and scoring strategy. *Nature Microbiology* **3**, 836–843
(2018).

90. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. DRep: A tool for fast and
accurate genomic comparisons that enables improved genome recovery from
metagenomes through de-replication. *ISME Journal* **11**, 2864–2868 (2017).

91. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk v2:
memory friendly classification with the genome taxonomy database.
Bioinformatics **38**, 5315–5316 (2022).

92. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Molecular Biology and Evolution* **37**, 1530–1534
(2020).

93. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L.
S. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat*
Methods **14**, 587–589 (2017).

94. Letunic, I. & Bork, P. Interactive tree of life (iTOL) v5: An online tool for
phylogenetic tree display and annotation. *Nucleic Acids Research* **49**, W293–

W296 (2021).

95. Sahlin, K. Strobealign: flexible seed size enables ultra-fast and accurate read
alignment. *Genome Biol* **23**, 260 (2022).

96. Aroney, S. T. N. *et al.* CoverM: Read alignment statistics for metagenomics.
Preprint at https://doi.org/10.48550/arXiv.2501.11217 (2025).

97. Zhou, Z. *et al.* METABOLIC: high-throughput profiling of microbial genomes for
functional traits, metabolism, biogeochemistry, and community-scale functional
networks. *Microbiome* **10**, 1–23 (2022).

98. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using
DIAMOND. *Nature Methods* **12**, 59–60 (2014).

99. Mistry, J., Finn, R. D., Eddy, S. R., Bateman, A. & Punta, M. Challenges in
homology search: HMMER3 and convergent evolution of coiled-coil regions.
Nucleic Acids Research **41**, e121–e121 (2013).

100. Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile
open source tool for metagenomics. *PeerJ* **4**, e2584 (2016).

101. Ruff, S. E. *et al.* A global comparison of surface and subsurface microbiomes
reveals large-scale biodiversity gradients, and a marine-terrestrial divide. *Sci.*
Adv. **10**, eadq0645 (2024).

102. Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217
(2012).

103. Olm, M. R. *et al.* inStrain profiles population microdiversity from metagenomic
data and sensitively detects shared microbial strains. *Nat Biotechnol* **39**, 727–736
(2021).

104. Figueroa Iii, J. L., Dhungel, E., Bellanger, M., Brouwer, C. R. & White Iii, R. A.

MetaCerberus: distributed highly parallelized HMM-based processing for robust functional annotation across the tree of life. *Bioinformatics* **40**, btae119 (2024).

105. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B: Statistical Methodology* **57**, 289–300 (1995).

106. Csádi, G. & Nepusz, T. The igraph software package for complex network research. *InterJournal, complex systems* **1695**, 1–9 (2006).

107. Bastian, M., Heymann, S. & Jacomy, M. Gephi: An Open Source Software for Exploring and Manipulating Networks. *ICWSM* **3**, 361–362 (2009).

108. Kolde, R. *Pheatmap: Pretty Heatmaps*. (2025).