1

Uranium mining fuels evolution in deep groundwater microbiomes

Wei Xiu[1,2], Till L.V. Bornemann[2], Tianjing Zhang[1], André R. Soares[2], Luping Xie[1], Julian M. Künkel[2], Guoxi Lian[3], Haicheng Weng[4], Chongsheng Lu[1,6], Bing Yang[4], Ruixuan Gai[1], Zhipeng Gao[1], Di Zhang[1], Yuan Yuan[4], Zhiming Du[4,7,8], Qingyin Xia[4], Linxing Chen[9], Jonathan R Lloyd[5], Hailiang Dong[1], Xuebin Su[10], Huaming Guo[1]*, and Alexander J. Probst[2]*

[1] *State Key Laboratory of Geomicrobiology and Environmental Changes (GMEC), China University of Geosciences, Beijing 100083, People's Republic of China*
[2] *Environmental Metagenomics, Research Center One Health Ruhr, University Alliance Ruhr, Faculty of Chemistry, University of Duisburg-Essen, Germany*
[3] *Nuclear and Radiation Safety Center, Ministry of Ecology and Environment, Beijing, 100082, China*
[4] *Beijing Research Institute of Chemical Engineering and Metallurgy CNNC, No.145 Jiukeshu, Tongzhou District, Beijing 101149, China*
[5] *Williamson Research Laboratories for Molecular Environmental Science, Department of Earth and Environmental Sciences, The University of Manchester, Manchester M13 9PL, United Kingdom*
[6] *The Institute of Hydrogeology and Environmental Geology, Chinese Academy of Geosciences, Shijiazhuang 050061, China*
[7] *China University of Petroleum, Beijing 102249, China*
[8] *National Key Laboratory of Uranium Resources Exploration-Mining and Nuclear Remote Sensing, Nanchang 330013, China*
[9] *State Key Laboratory of Advanced Environmental Technology, the Department of Environmental Science and Engineering, University of Science and Technology of China, Hefei, Anhui 230026, China*
[10] *China National Uranium Co., Ltd, Sanlihe, Xicheng District, Beijing, 100045, PR China*

* to whom correspondence should be addressed: Prof. Alexander J. Probst (alexander.probst@uni-due.de); Prof. Huaming Guo (hmguo@cugb.edu.cn)

Abstract

Deep-biosphere microbes have been underpinning global biogeochemical cycles throughout Earth history, yet their evolutionary responses to long-term anthropogenic disturbance remain poorly understood. Neutral-pH in-situ leaching (ISL), the dominant uranium-mining strategy, imposes persistent radiochemical and redox stress offering rare natural experiments across decades for observing subsurface microbial evolution. Here, by exploiting metagenomics and metatranscriptomics in examining microbial responses through various mining stages, we show that neutral U ISL causes substantial changes in microbial communities spanning 2,294 strains accompanied by diversification and selection of active microbial species. These changes occurred during elevated dissolved uranium and radiological activity which correlated with a substantial transcriptional change in energy metabolism, and with an overexpression of specific genes for oxidative-stress defence and DNA-repair pathways. Increased nucleotide diversity in 101 out of 392 species clusters and nonsynonymous/synonymous polymorphism ratios greater than one in 139 species clusters were predominantly observed in metatranscriptomes with elevated radiation, highlighting positive selection of transcriptionally active populations. These findings demonstrate that neutral U ISL drives functional and genetic diversification of subsurface microbiomes, revealing a dynamic and evolutionarily responsive deep biosphere.

Introduction

Microbial consortia in the deep subsurface, adapted to nutrient-scarce, anoxic environments[1–4], mediate critical biogeochemical processes and help maintain groundwater quality[5–8]. However, anthropogenic stresses arising from mining, groundwater extraction, and energy production are perturbing these fragile environments at increasing rates[9]. Recent studies have shown that environmental stresses, including heavy metal contamination and altered pH, affect groundwater microbial communities[10,11]. Mining operations are particularly disruptive, as they rapidly subject pristine subsurface ecosystems to heavy metals, radioactive contaminants, altered redox conditions, and electrochemical gradients[9]. As mining encroaches on fragile groundwater ecosystems, resolving the evolutionary responses of subsurface microbiomes is essential for predicting the long-term consequences of resource extraction. Notably, a recent study in the abyssal seafloor showed that deep-sea mining can cause biological disruptions persisting for decades[12], underscoring the need to assess long-term impacts of subsurface resource extraction on microbiomes. Although distinct from the deep subsurface examined here, this finding underscores how resource extraction can impose long-lasting impacts on microbial communities in Earth's hidden biomes.

In-situ leaching (ISL), a widely used uranium (U) mining technique, is based on injecting oxidizing agents into aquifers to mobilize U by oxidizing insoluble U(IV) to soluble U(VI). In addition to fundamentally altering groundwater chemistry[13], this activity imposes profound chemical and radiological stresses on microbial communities that have evolved over a long time under stable and reducing conditions. Whereas the environmental consequences of ISL are well documented mostly via 16S rRNA gene analysis[11,14–16], the evolutionary responses of microbial communities to these stressors remain poorly understood. Indeed, while mining-driven alterations to groundwater chemistry are expected to disrupt the microbial community structure, functional potential, and genetic diversity[17–20], the question remains as to whether anthropogenic stress can drive microbial evolution in deep aquifers.

To this end, we implement metagenomic and metatranscriptomic strategies to examine microbial community dynamics and evolutionary adaptations at the first large-scale, neutral-pH ISL operation in China, which uses CO_2 and O_2 as leaching agents and targets a sandstone-hosted uranium deposit. It involved sampling aquifers spanning pre-mining, active mining, and post-mining phases. Leveraging this rare

sampling opportunity to study microbial adaptation as a response to mining and
associated geochemical changes, we show that increases in Eh and the
accompanying rise in oxidative stress, including U release and associated radioactivity,
are linked to restructuring of microbial communities, enhanced oxidative-stress and
DNA-repair responses, and alterations in genetic diversity. Elevated pN/pS ratios (the
ratio of the rates of non-synonymous (pN) and synonymous (pS) polymorphism) and
shifts in nucleotide diversity implicate mining-induced environmental gradients in
significantly altering microbial evolutionary processes. Our findings yield new insights
into how deep subsurface microbiomes structurally and evolutionally adapt to
anthropogenic stress, bearing implications for global biogeochemical cycles,
ecosystem resilience, and bioremediation strategies. The findings of this investigation
underscore the consequences of uranium ISL on the evolution of subsurface life,
offering a novel framework for appraising microbial resilience and adaptation in
mining-impacted deep subsurface ecosystems.
103
RESULTS
Geochemical and radiological alterations during and following Uranium ISL
Neutral pH uranium *in-situ* leaching (neutral-pH ISL) commences while the
aquifer is in its natural state (pre-mining phase), proceeds through the injection of
leaching solutions (mining phase), and concludes with natural restoration efforts upon
cessation of mining activities (post-mining phase; *Figs. 1a-b*). We characterized the
geochemical and radiological shifts across all phases of neutral-pH ISL by collecting
80 groundwater samples from four distinct mining sites in Qianjiadian sandstone-
hosted uranium deposit in China, tapping aquifers at 220-460m depth below surface
(*Figs. 1b-c*). These sites underwent neutral-pH ISL for varying durations, creating a
unique time-resolved dataset with considerable variation in their redox conditions, from
neutral-reducing pre-mining (BG, baseline groundwater) to weakly acid-oxidizing
during mining (MT, mining groundwater), and back to neutral-reducing post-mining
(PMI, PMIS, and PMS, representing initial, intermediate, and stable post-mining
phases, respectively; *Fig. 1c*).
Geochemical and radiological profiles shifted markedly across the pre-mining,
mining and post-mining stages of neutral-pH ISL. During the pre-mining phase,
groundwater was neutral-reducing (Fig. *1d*), with low total dissolved U (TU, 0.44 ± 0.31
mg/L; Fig. 1h). The aquifer was enriched in electron donors, including dissolved

organic carbon (DOC, 5.26 ± 3.51 mg/L), Fe(II) (0.85 ± 1.11 mg/L), ammonium (0.73
± 0.70 mg/L), and sulfide (0.92 ± 1.18 mg/L) (Fig. 1f,S1), conditions that favour the
reductive immobilization of U and are consistent with ongoing U mineralization under
reducing conditions. In the mining phase, the injection of CO_2 and O_2 coincided with a
clear oxidative shift, marked by significant increases in Eh (Fig. *1d*) and a rapid rise in
TU (Fig. *1h*), which peaked after approximately four years (12.44 ± 5.04 mg/L in
S3a_MT04; S3a_MT04, *n* = 6 vs. S3a_BG, *n* = 5; Kruskal–Wallis with Dunn's post hoc
test, *p* = 0.003) before declining gradually from MT5 to MT20 (Fig. *1h*). Radiological
activity increased in parallel, with elevated gross β and gross (α+β) signals (Figs. *1i–*
k). Bicarbonate concentrations also rose significantly (Fig. *1g*), and both sulfate and
nitrate increased in mid-mining stages (Fig. *S1* and Table *S1*). Following neutral-pH
ISL decommissioning, groundwater shifted toward reducing conditions, with significant
decreases in Eh across MT15 compared with all post-mining groups (S3b_PMI,
S3b_PMIS, and S3b_PMS; Kruskal-Wallis with Dunn's post hoc test, *p* < 0.001; Fig.
1d) and corresponding declines in TU (S3b_MT15 vs. S3b_PMS; ANOVA with Tukey's
test, *p* = 0.036; Fig. *1h*). This transition from oxic to anoxic conditions was further
reflected by decreases in sulfate, and sulfate/Cl ratio, whereas Fe(II) concentrations
increased, although these changes were not statistically significant (Fig. *S1* and Table
S1). TU concentrations did not return to pre-mining levels (*Fig. 1h*), indicating
incomplete restoration of the aquifers' original geochemical conditions.
Correlation analysis identified significant Spearman associations among major
geochemical and radiological parameters (Fig. *S2*). TU showed strong positive
correlations with bicarbonate (ρ = 0.43, n = 80, *p* < 0.0001), sulfate (ρ = 0.65, n = 80,
p < 0.0001), and nitrate (ρ = 0.69, n = 80, *p* < 0.0001). In addition, TU exhibited positive
correlations with gross α (ρ = 0.24, n = 80, *p* < 0.0001), gross β (ρ = 0.24, n = 80, *p* <
0.0001), and gross (α+β) (ρ = 0.24, n = 80, *p* < 0.0001). These relationships
summarize the major covariation patterns among dissolved uranium, radiological
activity, and oxidized anions across all samples.
151
152

Figure 1 | Sampling locations and study overview and geochemical profiles of groundwater across neutral-pH ISL phases. *(a) Geographic location of the Songliao Basin and broader regional context, modified from a previously published source[24]. (b)*

Schematic layout of study sites along a neutral-pH ISL chronosequence: Site 1
(unmined aquifer, S1_BG); Site 2 (unmined aquifer, S2_BG; recent ISL at 0.25-1 years:
S2_MT0.25 and S2_MT01); Site 3a (unmined aquifer, S3a_BG; intermediate ISL, 4-6
years: S3a_MT04, S3a_MT05, and S3a_MT06); and Site 3b (unmined aquifer,
S3b_BG; long-term ISL, 15-20 years: S3b_MT15 and S3b_MT20; and
decommissioned neutral-pH ISL under iron-reducing, S3b_PMI, mixed iron- and
sulfate-reducing, S3b_PMIS, and sulfate-reducing, S3b_PMS conditions). Wells
shown with two names indicate that the same well was sampled at different time points.
The background map shows oxidized sandstone (yellow), reduced sandstone (grey),
uranium-bearing sandstone (pink), uranium ore deposits (red), and neutral-pH ISL
units (arrow pointers). (c) Numbers of groundwater samples collected for
hydrochemical, metagenomic, and metatranscriptomic analyses across stages of
neutral-pH ISL. Boxplots show geochemical parameters through the stage of neutral-
pH ISL, including pre-mining (BG), mining (MT), and post-mining (PM). Panels (d – k)
represent Eh, pH, DOC, bicarbonate, TU, gross α, gross β, and gross (α+β),
respectively. Statistical significance was determined by ANOVA with Tukey's or
Kruskal-Wallis with Dunn's test (BH-corrected), depending on data normality. Colors
indicate study sites: light blue (Site 1), light green (Site 2), medium green (Site 3a),
and beige (Site 3b). Blue boxplots within the studied sites (S2_BG, S3a_BG, and
S3b_BG) represent the geochemical parameters from unmined aquifer samples.

178 Metagenome-assembled genomes reveal microbiome changes and enhanced 179 co-occurence resulting from *in-situ* mining activity

Sixty groundwater samples representative of the pre-mining, mining, and post-
mining phases across more than 20 years of neutral-pH ISL were metagenomically
sequenced (Figs. *1b,c*). Exceeding 75% of sequencing coverage, based on Nonpareil
estimates, indicated a sufficient depth to accurately reflect the aquifer microbiome
diversity (Fig. *S3*). To assess both bacterial and archaeal diversity, we focused first on
analyzing the ribosomal protein S3 gene (*rpS3*)[21] yielding 12,972 sequences that
clustered (at 99% similarity) into 6,487 representative sequences spanning 215
distinct phyla (21 archaeal and 194 bacterial; Fig. *S4*). Microbial diversity also showed
strong positive associations with uranium concentration and radioactity, with
prolonged neutral-pH ISL (S3a_MT04 and S3a_MT05) exhibiting significantly higher
Shannon and Invsimpson indices (ANOVA with Tukey's post hoc test, $p < 0.05$; Fig.

S5). Groundwater microbial community composition and alpha diversity, based on
rpS3 gene profiles, were strongly influenced by geochemical gradients, particularly
radiochemical activity, with higher diversity and distinct community structures
observed under elevated stress conditions (*Figs. S5,S6)*, as supported by *Bioenv*
analysis (R^2 = 0.051)*.

We reconstructed 4,075 medium-quality metagenome-assembled genomes
(MAGs; completeness ≥ 50%, contamination < 10%)[22], from which 2,294 strain-cluster
representative MAGs (strain-cluster rMAGs, 99% ANI) were obtained, with
Acidobacteriota representing the most abundant phylum (Figs. *3a-b*). These included
1,239 high-quality and 1,155 medium-quality strain-cluster rMAGs bearing a median
completeness of 94.2%, contamination level of 1.3%, and N50 of 47,633 (Figs. *3c-d*
and *Table* S2). Consistent with known genome-size differences between domains,
archaeal strain-cluster rMAGs had substantially smaller genome sizes than bacterial
ones (two-tailed Welch's t-test, p < 0.0001; Fig. *3e*), the largest genome representing
an organism from the family *Polyangiaceae* (12.28 Mbp). These strain-cluster rMAGs
spanned 63 of the 215 *rpS3*-detected phyla (29.3% of total phylum richness) yet
represented 75.1% of the *rpS3*-based phylum-level abundance, thereby providing a
robust representation of the predominant microbial communities (*Fig. S7*). The
remaining 70.7% of phyla, comprising the 24.9% of *rpS3*-based abundance, were not
captured in the strain-cluster rMAGs, potentially reflecting their lower abundance
and/or genomes that did not meet medium-quality criteria.
212

Figure 3 | *Phylogenetic diversity, abundance patterns, and genomic characteristics of dereplicated strain-cluster representative MAGs. (a) Maximum-likelihood phylogenetic tree of 2,294 dereplicated bacterial and archaeal strain-cluster*

 rMAGs constructed. Outer rings indicate strain-cluster rMAG taxonomic affiliation at
 the phylum level (innermost ring), estimated completeness, contamination, and
 genome size (outermost green bars). (b) Rank abundance of strain-cluster rMAGs
 across phyla, with the top 100 depicted in more detail. The displayed abundances
 represent the average strain-cluster rMAG abundances across all samples. (c) Strain-
 cluster rMAGs completeness and contamination by quality tier. (d) Distribution of
 log(N50) values of strain-cluster rMAGs. (e) Genome size comparison between
 archaeal and bacterial strain-cluster rMAGs. Note that two rMAGs showed long-
 branch or model-sensitive placements; these were retained for completeness but do
 not affect the overall phylogenetic structure.

Microbial community composition, based on the relative abundance of strain-
cluster rMAGs, shifted markedly along the neutral-pH ISL chronosequence (Fig. *S8*),
as evidenced by sample-to-sample Spearman correlation analyses (Fig. *4a*) and non-
metric multidimensional scaling (NMDS; Fig. *4b*). Communities from pre-mining,
mining, and post-mining stages formed distinct clusters, with analysis of similarities
(ANOSIM) indicating significant separation between stages (Fig. *4c*), reflecting
divergent community trajectories during and after ISL. Network analysis of relative
abundances of strain-cluster rMAGs revealed positive correlations, accounting for
90.39% of all edges, defined as significant correlations between taxa, dominating
microbial interactions (Fig. *4d*). Four distinct modules (from Mod1 to Mod4) were
identified. Mod1 remained stable across all ISL stages, consistent with a core
community. Mod2 and Mod4 increased only during the early ISL stages, reflecting
short-term adaptive responses. In contrast, Mod3 showed a sustained increase during
the mining and post-mining phases, indicative of recovery processes and stress-driven
selection for taxa better adapted to prolonged radiochemical and redox stress (Figs.
4f-i). Within-module degree (Z_i) and particitation coefficient (P_i) analysis identified two
module hubs (two strain-cluster rMAGs classified within the *Algoriphagus* and
Paracoccus) that structured interactions within Mod4, and three connector hubs (three
strain-cluster rMAGs classified within *Pseudotabrizicola*, *Burkholderiaceae* MESE01,
and *Nocardioides*) that linked multiple modules. In addition, 341 peripheral connectors
($Z_i \leq 2.5$ and $P_i > 0.62$) were dominated by Mod3 (175 strain-cluster rMAGs) and Mod2
(129 strain-cluster rMAGs) (Fig. *4e*), indicating that most cross-module interactions
are mediated by diffuse peripheral taxa rather than by central hub nodes. These

results highlight the significant shifts in microbial community structure, diversity, and
ecological interactions imposed by neutral-pH ISL. The observed tendency towards
positive interactions emphasizes the dynamic and evolving nature of subsurface
microbiomes in response to acute environmental stress.
255

257 **Figure 4 | Microbial community structure and modularity across neutral-pH ISL**
258 **phases.** *(a) Heatmap of sample-to-sample Spearman's rank correlations of strain-*
259 *cluster rMAG abundances across early (blue), mid (green), and late (red) neutral-pH*
260 *U ISL stages. The accompanying bar chart shows the top 12 classes across samples,*

arranged in the same sample order as in the heatmap. (b) Non-metric multidimensional scaling (NMDS) plot of community structure based on strain-cluster rMAGs (stress = 0.096). Background colors represent gross (α+β) radiation levels across all groups (interpolated across the ordination space using inverse distance weighting (IDW). Gross (α+β) radiation levels below 100 Bq/L are shown as a continuous gradient, while gross (α+β) radiation levels ≥100 Bq/L are displayed as a solid yellow. White contour lines and labels indicate gross (α+β) isopleths (Bq/L). (c) Analysis of Similarity (ANOSIM) showing significant differences between groups. (d) Network of strain-cluster rMAGs co-occurrence, with positive (grey) and negative (red) Spearman correlations grouped into four modules. Node size is proportional to the degree (number of connections). (e) Within-module degree (Zi) and participation coefficient (Pi) analysis of the network to identify the peripherals, connectors, module hub and network hubs. The size and color of the nodes are proportional to its corresponding degree and module in the network, respectively. (f-i) Distribution of the abundance of module-specific taxa across neutral-pH ISL phases for Mod1 to Mod4.

Neutral-pH ISL impacts gene expression linked to DNA repair, oxidative stress, and biogeochemical cycling

To resolve microbial functional responses to prolonged neutral-pH ISL, we analyzed metatranscriptomes from 45 genomically resolved communities spanning the pre-mining, mining, and post-mining phases. Integration of transcript profiles with 2,294 functionally annotated strain-cluster rMAGs identified 350,588 of 8,422,368 representative genes showing at least two-fold upregulation relative to the mean expression of 30S ribosomal protein genes, which served as an internal benchmark for transcriptional activity. The transcriptional landscape of entire communities based on strain-cluster rMAGs showed clear patterns of transcriptionally up-regulated populations that are specific for the sample groups across the mining sites (Fig. *S9*). Among 350,588 upregulated genes of the rMAGs, 5,677 genes were associated with carbon, hydrogen, oxygen, nitrogen, sulfur, and iron biogeochemical transformations, 4,624 with DNA repair pathways, and 2,760 with oxidative stress responses (Figs *S10-S17*), highlighting broad activation of redox and genome-stability pathways under ISL-induced chemical and radiological stress.

Regarding microbial activity of biogeochemical transformations, the expression patterns across the 5,677 related genes were strongly structured by geochemical

conditions, as revealed by *Bioenv* analysis, which identified pH, Eh, sulfide, sulfate and bicarbonate as the principal geochemical variables shaping the transcriptional structure of overexpressed C-H-O-N-S-Fe metabolic genes across the metatranscriptomes (R^2 = 0.39), with PCoA accounting for 36.9% of the variation among overexpressed genes (Figs. *5a* and *S9*). This pattern is consistent with CO_2 and O_2 injection and with oxidation of ammonium and pyrite as dominant ISL reactions fueling microbial redox metabolism. TU concentrations were moderately and positively correlated with the expression of 20 energy-conserving genes distributed across 220 strain-cluster rMAGs, including *ccoN* (cytochrome *c* oxidase, aerobic respiration; ρ = 0.44, *p* = 0.029; encoded in 77 strain-cluster rMAGs), *pmoC* (particulate methane monooxygenase, methane oxidation; 21 strain-cluster rMAGs), *nxrAB* (nitrite oxidoreductase, nitrite oxidation; ρ > 0.53, *p* < 0.006; *nxrA* in 21 and *nxrB* in 27 strain-cluster rMAGs), and *soxAYZ* (sulfur oxidation complex, sulfur oxidation; ρ > 0.41, *p* < 0.038; *soxA* in 43, *soxY* in 69, and *soxZ* in 74 strain-cluster rMAGs) (Figs. *5d-f* and *S16-S17*). These relationships highlight a broad activation of energy-yielding redox pathways along radiochemical gradients, underscoring the metabolic coupling of carbon, nitrogen, and sulfur cycling with oxidative stress.

In addition to metabolic adjustments, microbial populations showed dynamic transcriptional regulation of DNA-repair and oxidative-stress pathways in response to radiochemical gradients. The *Bioenv* analysis identified pH, Eh, sulfide, ammonium, and bicarbonate as the main variables structuring DNA-repair gene expression (4,624 genes, R^2 = 0.40; PCoA explained 31.7% of variance), whereas pH, Eh, sulfide, sulfate, and bicarbonate were key for the 2,760 oxidative-stress-response genes (R^2 = 0.37; PCoA explained 45.2% of variance; Figs. *5b,c*). At the individual gene level, expression of foxE (c-type cytochrome involved in Fe(II) oxidation) correlated positively with gross β and gross (α+β) (Spearman, *p* < 0.05; 8 strain-cluster rMAGs; Figs. *5b-e*), consistent with Fe redox processes being stimulated by oxygen ingress and the resulting shifts in uranium speciation and radiological activity. Consistent with these environmental associations, expressions of DNA-damage-repair genes, including *mutM* (DNA glycosylase; base-excision repair of oxidized bases; ρ = 0.43-0.49, *p* = 0.021-0.047; 68 rMAGs), *dam* (DNA adenine methylase; mismatch-repair guidance; ρ = 0.45-0.51, *p* = 0.016-0.038; 37 rMAGs), and *alkC* (alkylation-repair protein; ρ = 0.45-0.46, *p* = 0.034-0.038; 4 rMAGs), were positively correlated with TU and gross α, β, and (α+β) (Spearman correlations, *p* < 0.05; *Figs. 5b-e*). Notably, *mutM*

329 expression correlated moderately with both TU (ρ = 0.41, n = 26, *p* = 0.0435) and
330 gross (α + β) (ρ = 0.51, n = 2,632, *p* = 0.0084; Fig. *5e*). These coordinated
331 transcriptional patterns indicate widespread modulation of genome-stability and redox-
332 defense mechanisms, coupling Fe redox metabolism, oxidative stress responses, and
333 DNA repair under sustained radiochemical and geochemical stress in the aquifer.
334

Figure 5 | Correlation and environmental drivers of overexpressed genes in
metatranscriptomic datasets. *(a–c) Principal coordinate analyses (PCoA) based on*
expression of (a) genes involved in carbon, hydrogen, oxygen, nitrogen, sulfur, and
iron metabolism, (b) DNA-repair genes, and (c) oxidative-stress-response genes.
Arrows indicate environmental variables significantly correlated with gene-expression
patterns, as identified by Bioenv analysis. Points are colored and shaped by sample

Uranium ISL-induced stress drives genetic diversification in subsurface aquifers

Within-species selection pressure and genetic diversity were quantified across 2402 members of 392 dereplicated species-cluster rMAGs (95% ANI). Each rMAG was represented by more than three members with >5× genome coverage in 60 metagenomic and 45 metatranscriptomic datasets, spanning neutral-pH ISL phases, enabled direct comparisons of genomic and transcriptional diversity across mining stages.

NMDS analysis of pN/pS ratios revealed a clear progression from background to mining and post-mining samples in both metagenomes (Fig. *6h*) and metatranscriptomes (Fig. *6j*), closely aligned with increasing TU concentrations. Consistent with this ordination, site-level analyses showed that the median metagenomic pN/pS ratios increased in early mining, late mining, and post-mining sites and metatranscriptomic pN/pS ratios exhibited similar but stronger increases ($p < 0.05$, *Figs. 6a-c*). *Bioenv* analysis identified gross α, gross (α+β), Eh, TFe, TU, ammonium, Fe(II), nitrite, and nitrate as the main correlates of metagenomic pN/pS ($R^2 = 0.28$), and gross β, gross (α+β), TDS, TU, and ammonium for the metatranscriptomes ($R^2 = 0.23$), indicating that mutation–selection imbalance reflects the combined influence of the oxidizing geochemical regime and co-varying metal and radionuclide stress. Instances of positive selection (pN/pS > 1) were more frequent in the metatranscriptomes (139 MAG clusters) than in the metagenomes (112 MAG clusters) or in both datasets combined (49 MAG clusters; Fig. *S19*), highlighting intensified adaptive pressures on actively expressed genomes.

To explore whether these shifts in selection were supported by changes in
standing genetic variation, we next examined genome-wide nucleotide diversity
across species-cluster rMAGs. NMDS ordinations revealed a consistent progression
from background to mining and post-mining samples in both metagenomes and
metatranscriptomes, closely tracking TU gradients (*Figs. 6g-j*). Metagenomic
nucleotide diversity remained relatively stable during the mid-mining phase but
declined in early-, late-, and post-mining stages despite sustained TU concentrations.
(Figs. *6d-f*). By contrast, metatranscriptomic nucleotide diversity rose sharply during
the mid-mining phase at site S3a ($p < 0.001$; Fig. *6e*), coinciding with peak uranium
and radioactivity. Bioenv analysis identified radioactivity (gross α, gross β, and gross
(α+β)) and geochemical parameters (Eh, TFe, TU, ammonium, Fe(II), and nitrate) as
key correlates of metagenomic genome-wide nucleotide diversity ($R^2 = 0.25$), with
comparable associations in metatranscriptomes (gross β, gross (α+β), TDS, TU, and
ammonium; $R^2 = 0.21$). Together, these patterns indicate that prolonged uranium-
mining stress maintains elevated genetic variation in transcriptionally active
populations, consistent with bet-hedging-like strategies that enhance adaptive
flexibility in fluctuating radiochemical environments.

 Figure 6 | Genome-wide pN/pS ratio and nucleotide diversity across mining

stages in metagenomes and metatranscriptomes. *(a-c) pN/pS ratio from*

metagenomic (metaG) and metatranscriptomic (metaT) datasets spanning pre-mining,

mining, and post-mining phases; (d-f) Nucleotide diversity from metaG and metaT

datasets across neutral-pH ISL stages; (g-j) Non-metric multidimensional scaling

(NMDS) plots based on Bray–Curtis dissimilarity showing the relationships between

TU and pN/pS ratios or nucleotide diversity derived from metaG (g, h) and metaT (i, j)

datasets.Statistical significance was assessed using the Kruskal–Wallis test followed

by Dunn's post‑hoc test with Benjamini–Hochberg (BH) correction for multiple

comparisons (* p < 0.05, ** p < 0.01, and *** p < 0.001).

Sixty-five out of 392 species clusters exhibited recurrent increases in nucleotide

diversity from background to mining-impacted conditions, including 20 detected in both

metagenomic and metatranscriptomic datasets (Figs. *7a,b*). Four representative taxa,

Kuenenia (rMAG-1), *Gallionella* (rMAG-2 and rMAG-4), and *Parvibaculum* (rMAG-3),

persisted across all mining phases and exhibited elevated genome-wide diversity in

both metagenomic and metatranscriptomic datasets. Gene-level diversification was

dispersed across loci rather than confined to specific genomic regions, with hundreds

of variable genes per genome (rMAG-1: 269/1,126; rMAG-3: 95/148; rMAG-4: 213/280;

rMAG-2: 295/350) and substantial overlap between genomic and transcriptomic

variation, indicating that many variable loci were transcriptionally active, particularly

within core functional networks linked to uranium-induced stress responses, including

information processing, regulation, and stress adaptation (*Fig. 7c*). *Kuenenia* rMAG-1

exhibited genomic diversification in hydroxylamine dehydrogenase (*hao*) and

hydrazine dehydrogenase (*hdh*) genes mediating ammonium oxidation, together with

lexA (SOS-response regulator involved in DNA damage repair), *old-family*

endonuclease (genome maintenance), *lon* (ATP-dependent protease for protein

quality control), and *tolC* (outer-membrane efflux channel for metal detoxification). The

two *Gallionella* rMAGs displayed contrasting transcriptional responses. The rMAG-2

showed variability in redox-related genes, including cytochrome-c oxidase cbb3-type

subunit II (*fixO*), involved in microaerobic respiration and electron transport, and

thioredoxin-dependent peroxiredoxin, linked to reactive oxygen species metabolism

under oxidative stress. By contrast, rMAG-4 exhibited transcript-level variability in

outer-membrane porins (*ompC*/*ompF*/*phoE*) associated with metal-regulated

permeability, the conjugal transfer protein *traF* involved in horizontal gene transfer, and glycosyltransferases contributing to cell-envelope remodeling. *Parvibaculum* rMAG-3 encoded a GIY–YIG endonuclease, associated with the cleavage of damaged or mismatched DNA, and a Zn-dependent protease involved in protein quality control under oxidative stress, including conditions where metal exposure can exacerbate ROS damage. Together, these diversification patterns suggest that persistent taxa employ distinct, lineage-specific strategies-spanning redox flexibility, oxidative defense, membrane adaptation, and proteogenomic repair, to maintain resilience under the combined chemical and radiochemical stresses imposed by U mining.

Figure 7 | Genome-wide nucleotide diversity patterns across mining stages in representative rMAGs. *(a) Bar plots show the total numbers of species-cluster rMAGs with increased or decreased nucleotide diversity (π) from background (BG) to mining (MT) conditions in metagenomic (metaG) and metatranscriptomic (metaT) datasets. Increased π indicates rMAGs with callable nucleotide diversity (≥ 5× coverage) in MT samples but either non-callable diversity (< 5× coverage) or lower π values in BG samples; decreased π follows the inverse criterion. The accompanying Venn plot displays the overlap of rMAGs with increased π (π > 0) between the metaG and metaT datasets. (b) Ternary plot showing the distribution of rMAG-level nucleotide diversity across BG, MT, and post-mining (PM) samples, with points colored by dataset and sized by diversity magnitude. (c) Gene-level nucleotide diversity profiles for four representative species-cluster rMAGs (Parvibaculum, Kuenenia, and Gallionella) across BG, MT, and PM phases. Each profile represents the median π of all cluster members for a given gene, highlighting elevated and widespread diversification during mining (MT), particularly in transcriptionally active genes. Bar plots below each panel show the functional distribution of genes with increased diversity categorized by major gene functions.*

Discussion

The global resurgence of nuclear energy has renewed interest in U ISL, a minimally invasive yet chemically intensive extraction method, poised to dominate future uranium supply chains[23,24]. During mining, strong oxidants drove the oxidation of U(IV) to U(VI) and the formation of soluble uranyl–carbonate complexes, while concurrent oxidation of pyrite and ammonium produced the elevated sulfate and nitrate characteristic of mining phases[25,26]. Following ISL cessation, DOM degradation re-established reducing conditions, accompanied by rising Fe(II) and declines in Eh, TU and sulfate, patterns consistent with Fe(III) and sulfate reduction and the partial reductive attenuation of U(VI)[27]. While such geochemical perturbations and ecological restructuring during U ISL are increasingly recognized[13,17–20,25,27–30], its evolutionary consequences for subsurface microbiomes remain underexplored. Here, we present metagenomic and metatranscriptomic evidence that neutral-pH U ISL exerts sustained selective pressure on microbial populations, driving *in situ* evolution within deep aquifers.

Neutral-pH ISL establishes persistent redox disequilibria through oxidant injection, mobilizing U, radionuclides, and heavy metals[13,25,27–30]. By examining a neutral-pH ISL area with aquifers having undergone different time periods of leaching (up to 20-years), we reveal that these sustained geochemical disturbances go beyond restructuring of microbial community composition and alter transcriptional activity, along with an increased genetic diversity, illuminating long-term ecological and evolutionary consequences for deep biosphere microbiomes. These transformations are shaped by sustained selection pressures imposed by neutral-pH ISL, including oxidative stress, redox-active metal toxicity, and low-dose radiation, stressors known to destabilize cellular systems and promote adaptive traits linked to stress tolerance, DNA repair, and redox regulation[31–33]. Comparable patterns of stress-driven adaptation have been reported in acid mine drainage biofilms[34,35], heavy-metal tailings[36,37], permafrost thaw soils[38], and deep-sea sediments[39,40]. Consistent with these analogs, we detected increased taxonomic diversity in neutral-pH ISL-impacted aquifers, signatures consistent with stress-induced diversification to increase resilience at the community level[11,41–45]. Temporal succession revealed dynamic microbial responses, with oxidative-stress-tolerant taxa dominating early neutral-pH ISL stages and anaerobic fermenters and metal reducers emerging post-mining, reflecting adaptation to evolving redox conditions.

Crucially, these ecological transitions were accompanied by molecular signatures of adaptation. Genes associated with oxidative damage, heavy metal toxicity, and radiation exposure were broadly overexpressed across all neutral-pH ISL stages (*Figs. S9-S17*), indicating a constitutive stress response. Among 75 species-cluster rMAGs with measurable nucleotide diversity that increased from background to mining phases (Fig. 7a), many exhibited pronounced diversification along the ISL gradient, suggesting that repair mechanisms were insufficient to counteract accumulated genomic damage. Consequently, elevated nucleotide diversity, particularly in mid-mining metatranscriptomes, supports a mutation–repair imbalance in which mutational input exceeded repair capacity, consistent with a previously introduced model of stress-induced mutagenesis[37,46–48]. This model posits that environmentally triggered DNA damage, coupled with error-prone repair, contributes to increased mutation rates. While DNA repair pathways buffer deleterious mutations, they may also limit adaptive potential by constraining the fixation of beneficial variants[49,50]. In our study, concurrent increases in nucleotide diversity and

transcriptional upregulation of DNA repair and oxidative stress-response genes
indicate that adaptive evolution is concentrated in metabolically active, redox-cycling
taxa rather than uniformly across the whole community. This resolves the modest
community-wide transcriptional changes yet strong gene-level positive selection
signals, highlighting that long-term ISL creates evolutionary hotspots where
radiochemical stress and altered substrate availability jointly promote adaptation in
key biogeochemical taxa.

Further support for this mechanism comes from elevated pN/pS ratios (>1), a
canonical signal of positive selection, observed predominantly in metatranscriptomes
(139 species-cluster rMAGs, including 251 and 40 cluster members were oberved in
mining and post-mining samples) and only partially overlapping with metagenomic
datasets (49 species-cluster rMAGs; Fig. S19). The rMAGs exhibiting elevated
nucleotide diversity encompassed phylogenetically diverse taxa and displayed
extensive gene-level variation in information processing and regulatory functions,
underscoring that selection acts primarily on transcriptionally active populations. This
layered response, genome-wide mutation accumulation coupled with transcription-
linked selection, highlights the dual role of environmental stress in both generating
genetic variability and filtering for functional adaptation, consistent with eco-
evolutionary rescue models[51–53], wherein microbial populations rebound from stress-
induced bottlenecks through the acquisition and retention of adaptive mutations.

Collectively, our study provides rare empirical evidence that stress-induced
mutagenesis and transcription-linked selection operate within natural deep subsurface
microbiomes over decadal timescales. Although such mechanisms are well
characterized in laboratory systems[54–60] and surface environments[46,61–64], their
manifestation in oligotrophic, energy-limited subsurface ecosystems[39,65–67] has
remained largely speculative due to assumptions of metabolic quiescence and
evolutionary stasis[40,68–70]. Our findings challenge this paradigm, demonstrating that
chronic anthropogenic stress in engineered aquifers can generate sustained
mutational input and facilitate directional selection. The convergence of elevated
nucleotide diversity, DNA repair activation, and positive selection in expressed genes
reveals a coordinated evolutionary response to long-term environmental disturbance.
These observations extend core theoretical frameworks, including stress-induced
mutagenesis and eco-evolutionary rescue, from controlled systems to complex,

anthropogenically modified subsurface environments, redefining the deep biosphere

as a dynamic, evolutionarily responsive system.

MATERIALS AND METHODS

Study area, sampling strategy, and sample collection. The study area located in

the Qianjiadian U ore deposit within the Qianjiadian (QJD) sag of the Kailu Depression

in the southwestern Songliao Basin, northeast China, represents a typical sandstone-

hosted U deposit[71]. The stratigraphy comprises the Qingshankou (K_2qn), Yaojia (K_2y),

and Nenjiang (K_2n) formations, overlain by Neoproterozoic and Quaternary alluvial

deposits. The U mineralization is primarily hosted in the K_2y Formation, which consists

of red, yellow, gray, and mineralized gray sandstones, with widespread carbonate

cementation (dolomite, siderite, ankerite, and calcite). Pitchblende and coffinite are

the dominant uranium minerals, while Fe-Ti oxides, natural organic matter, clay

minerals, and pyrite serve as additional U hosts[27,72,73]. Groundwater flow follows a

southwest-to-northeast trajectory within a confined Cretaceous sandstone aquifer

(approximately 298 m depth below surface, 35 m thick), with permeability averaging

0.145 m/day[25]. The aquifer is hydraulically isolated by a low-permeability mudstone-

dominated aquitard both above and below, with limited connectivity to adjacent

hydrogeological units.

The Qianjiadian U deposit has been a key site for advancing in-situ leaching (ISL)

technology in China, serving as the first large-scale demonstration of CO_2 + O_2

leaching in a deep, low-permeability sandstone-hosted system. During the neutral-pH

ISL process, the mixed reagent of dissolved O_2 (oxidant) and CO_2 (complexing agent)

was injected into the ore-bearing aquifer, and subsequently, U-containing groundwater

was pumped out and U was retrieved using ion exchange resins[25,74]. The effluent from

the reverse osmosis system was reused as a leaching solution after being mixed with

CO_2 and O_2 to maintain carbonate buffering and oxidizing conditions[75]. The orebody,

located within confined sandstone aquifers at depths ranging from 220 to 460 m,

presents complex hydrogeological conditions that influence solute transport, aquifer

reactivity, and microbial community dynamics. The unique geological and

hydrogeochemical setting of the Qianjiadian deposit, combined with its prolonged

history of neutral-pH ISL operations, establishes it as a natural laboratory for

investigating subsurface fluid-rock-microbe interactions, U mobilization processes,

and the long-term geochemical and microbial responses to neutral-pH ISL as well as
post-mining restoration.

To investigate the hydrogeochemical and microbial responses to neutral-pH ISL
mining, groundwater samples were collected from the Qianjiadian (QJD) sandstone-
hosted U deposit during two field campaigns in October-November 2021 and August-
October 2022 (Figure 1). A total of 80 groundwater samples were obtained from sites
representing different stages of neutral-pH ISL, spanning ore-bearing aquifers at
depths of 220-460 m. Sampling locations encompassed neutral-pH ISL sites spanning
multiple operational durations: 0.25 years (S2_MT0.25, 8 samples), 1 year (S2_MT01,
3 samples), 4 years (S3a_MT04, 6 samples), 5 years (S3a_MT05, 12 samples), 6
years (S3a_MT06, 6 samples), 15 years (S3b_MT15, 6 samples), and 20 years
(S3b_MT20, 4 samples). Post-mining phases were sampled under distinct redox
conditions, including iron-reducing (S3b_PMI, 7 samples), mixed iron- and sulfate-
reducing (S3b_PMIS, 5 samples), and sulfate-reducing (S3b_PMS, 4 samples)
conditions. Nineteen groundwater samples from unmined aquifer were collected
across the chronosequence, including non-mining areas (S1_BG, 5 samples), early-
stage mining sites (S2_BG, 1 sample), mid-stage mining sites (S3a_BG, 5 samples),
and late-stage/post-mining sites (S3b_BG, 8 samples). Background wells were
present at all sites and sampled concurrently with mining and post-mining wells,
providing site-specific baseline controls. Detailed hydrogeochemical data for
S3b_MT20, S3a_MT04, S3a_MT05, S3b_PMI, S3b_PMIS, S3b_PMS, and S3b_BG
are reported in our previous studies[25–28].

Prior to sample collection using a deep groundwater sampler (Geosub 2, Geotech,
USA), monitoring and post-mining wells were pumped for at least 20 minutes until
water temperature, electrical conductivity (EC), and oxidation-reduction potential
(ORP) stabilized. Groundwater was then pumped after well flushing and approximately
45 ± 0.5 L of water was filtered through 0.2-μm sterile cellulose nitrate membrane
filters (Sartorius, Göttingen, Germany) to capture microbial organisms from each
groundwater well. All filtered membranes were frozen at -80 °C for high-throughput
sequencing. Groundwater samples for physicochemical analysis were collected in 5 L
sterile PET bottles and stored at -20 °C.

Chemical Analysis. Major anions (Cl^-, NO_3^-, and SO_4^{2-}) were measured using ion
chromatography (Dionex ICS-600, Thermo Fisher Scientific, Massachusetts, USA),
while major cations (Na^+, K^+, Mg^{2+}, and Ca^{2+}) were determined using inductively

coupled plasma optical emission spectrometry (ICP-OES, Optima 8300, PerkinElmer,
Massachusetts, USA). Trace elements, including U, Fe, and other transition metals,
were quantified using inductively coupled plasma mass spectrometry (ICP-MS, 7900
ICP-MS, Agilent Technologies, California, USA). Alkalinity was measured by Gran
titration using a Metrohm 888 Titrando (Metrohm, Herisau, Switzerland). The total
organic carbon (TOC) and dissolved organic carbon (DOC) concentrations were
determined using a TOC analyzer (TOC-L, Shimadzu, Kyoto, Japan). The gross alpha
and gross beta radioactivity of groundwater samples were measured using a low-
background α/β meter (LB6008, Beijing Yida Measurement Technology Co., Ltd.,
Beijing, China) following the thick source method for radioactivity analysis, in which
evaporated water residues are deposited onto a planchet to form a uniform thick
source for α/β counting. Measurements of gross alpha and gross beta radioactivity
were conducted at the China Nuclear Mining Science and Technology Corporation,
Beijing, China, to assess radiological characteristics associated with uranium
mobilization.

DNA extraction and sequencing. Total metagenomic DNA was extracted from
biomass retained on 0.22-μm sterile cellulose nitrate membrane from 60 groundwater
samples, selected from a total of 80 samples retrieved from different neutral-pH ISL
stages in ore-bearing aquifers (see above and Figure *1b-c*). DNA extraction was
performed using the FastDNA® SPIN Kit for Soil (MP Biomedicals, Solon, OH, USA)
following the manufacturer's protocol. The concentration and purity of the DNA
extracts were measured using a TBS-380 fluorometer (Turner Biosystem, USA) and a
NanoDrop 2000 spectrophotometer (Thermo Scientific, USA), respectively. The
quality of the DNA extracts was checked using 1%-agarose gel electrophoresis.
Paired-end sequencing was performed on the Illumina HiSeq XTen platform (Illumina,
Inc., San Diego, CA, USA) at Majorbio Bio-Pharm Technology Co., Ltd., China. DNA
extracts were fragmented to an average size of approximately 400 bp using a Covaris
M220 ultrasonicator (Gene Company Limited, China). Paired-end libraries were
constructed using the NEXTFLEX Rapid DNA-Seq Kit (Bioo Scientific, Austin, TX,
USA), with adapters containing the full complement of sequencing primer hybridization
sites ligated to the blunt end of fragments.

Metagenome processing and MAG recovery. Raw reads were trimmed and quality-
filtered using fastp v0.19.7[83]. The microbial diversity and its coverage based on quality-
filtered raw metagenomic reads was estimated by Nonpareil v3.4.1[76] in "kmer" mode.

High-quality reads from individual samples were assembled into scaffolds using metaSPAdes version 3.15.5 (--k-list 21, 33, 55, 77, 99, and 127)[77]. Open-Reading Frames on scaffolds were predicted using prodigal (v2.6.3)[78] in "meta" mode. MAGs were recovered from each assembly using the binning module (parameters: -metabat2 -maxbin2 -concoct) within metaWRAP v1.3 (metabat2 v2.12.1, MaxBin2 v.2.2.6, and concoct v0.4.0)[79–82], which bins based on reads mapped only from the corresponding sample. Quality of genome bins (completeness and contamination) was evaluated using CheckM2 (v2.0.11)[83]. Binning results from multiple algorithms were integrated using DASTool v1.1.7 with default parameters[84] to generate a non-redundant MAG set for each sample. The resulting bins were further refined using itBins (https://github.com/ProbstLab/itBins), and screened and cleaned for contaminant sequences with the NCBI Foreign Contamination Screen (FCS-GX v0.5.5)[85]. Finally, a total of 4,075 MAGs were obtained with completeness ≥50% and contamination <10%. Bins were further consolidated via dereplication at strain level using dRep (v.3.2.2) at 99% ANI[86] resulting in 2,294 strain-cluster representative MAGs (strain-cluster rMAGs).

Taxonomic classification of strain-cluster rMAGs and phylogenetic tree generation. Taxonomies were assigned to each strain-cluster rMAG using GTDB-Tk v2.1.1 (https://github.com/Ecogenomics/GTDB-Tk, database release 226)[87]. Phylogenetic relationships were inferred using GTDB-Tk (v2.1.1), which identifies conserved marker genes and constructs genome trees following the GTDB reference framework. The resulting GTDB-based phylogenetic tree was used to resolve the taxonomic placement of strain-cluster rMAGs. The phylogenetic tree was visualized using the Interactive Tree of Life (iTOL)[88].

Coverage estimation and metabolic potential prediction. Clean reads from each sample were mapped to strain-cluster rMAGs using strobealign (v0.13.0)[89] in CoverM (v0.7.0; default parameters)[90]. The metabolic potential of strain-cluster rMAGs was determined using METABOLIC v4.0[91]. We constructed two databases, one containing 96 genes associated with DNA repair mechanisms and one containing 32 genes implicated in microbial responses to oxidative stress, with all genes being sourced from NCBI (https://www.ncbi.nlm.nih.gov/). Genes of strain-cluster rMAGs were annotated against these databases using DIAMOND BLAST[92] with an E-value threshold of 1e^{-5} and the parameter -k 1 to retain only the top scoring hit. To ensure the specificity of these annotations, a reverse BLAST was subsequently conducted by

aligning database hits back against the original amino acid sequences of the
corresponding strain-cluster rMAGs using the same E-value threshold. Only reciprocal
hits that aligned to the same ORFs as the original query were retained, ensuring one-
to-one correspondence. Cases with multiple high-scoring matches were excluded to
avoid ambiguous annotations. Final hits were filtered based on a similarity threshold
of >50%, ensuring high-confidence functional assignments. These curated datasets
facilitated a refined analysis of microbial resilience and adaptation strategies in diverse
environmental contexts.

Community profiling based on the *rpS3* marker gene. ORFs representing the *rpS3*
marker gene were identified using hmmsearch[93] against custom Hidden Markov
Models (HMMs) for Archaea and Bacteria (E-value 1e^{-28};
https://github.com/AJProbst/rpS3_trckr). Sequences were pooled and clustered at 99%
similarity using vsearch[94], representing species-level clusters[95]. Representative
sequences were selected following a tiered approach: (1) centroids that could be
extended by 1000 bp in both directions, (2) non-centroids that were similarly
extendable, and (3) the longest available sequence when neither of the first two criteria
were met. To improve abundance estimates, *rpS3* gene sequences of representative
rpS3 genes were extended by up to 1,000 bp in both directions on their scaffold of
origin if possible. This was done as mapping on just the approximately 1,000 bp long
rpS3 gene results in poor mapping on the gene edges and this has progressively more
impact when the total length of the sequence is short. The resulting *rpS3* gene
sequences, ideally extended in both directions by 1,000 bp, are henceforth called
rpS3-extended. The workflow for clustering, extracting, and calculating coverage is
available at https://github.com/ProbstLab/publication-
associated_scripts/tree/main/FigueroaGonzalez_Bornemann_etal. The *rpS3* genes in
rpS3-extended sequences (in amino acid format) were taxonomically classified using
DIAMOND[92] (E-value ≤ 1e^{-5}) against the GTDB r226 protein database, assigning the
best hit to be the taxonomy. If no hit meeting the search criteria was found,
"Unclassified" was assigned. Abundance was calculated by mapping reads of all
metagenomic samples to *rpS3*-extended sequences and calculating the average
coverage per sample. Additionally, the breadth, i.e., the number of nucleotide positions
in *rpS3*-extended having a coverage of at least 1, was determined. The coverage of
an *rpS3*-extended sequence in a given sample was set to zero if the breadth of the
sequence in that sample was less than 95%, indicating spurious mapping instead of

actual presence of the *rpS3*-extended-represented species. Relative coverage values

were transformed into percentages (%).

RNA extraction, sequencing, and metatranscriptomics. Total RNA was isolated

and purified using Soil RNA Extraction Kit (Majorbio, Shanghai, China) following the

manufacturer's procedure. The integrity and quantity of the extracted RNA were

measured with a NanoDrop 2000 spectrophotometer (Thermo Scientific, MA, USA)

and an Agilent 5300 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA). The RNA

was subjected to standard Illumina library preparation using the Illumina® Stranded

mRNA Prep, Ligation kit (Illumina, San Diego, CA, USA), and rRNA was depleted

using a RiboCop rRNA Depletion Kit for Mixed Bacterial Samples (Lexogen, USA).

Depleted RNA was extracted from biomass retained on 0.22 μm sterile cellulose

nitrate membranes from 60 groundwater samples (same to metagenomic groundwater

samples) and finally library preparation of RNA extraction from 45 groundwater

samples were successfully constructed (Figure *1c*). Sequencing was performed on an

Illumina Novaseq6000 sequencer (Illumina, San Diego, CA, USA) with paired-end

sequencing method at Majorbio Bio-Pharm Technology Co., Ltd. (Shanghai, China).

Quality control of the raw data generated by sequencing was carried out with fastp

v0.19.7[96]. SortMeRNA (v4.3.6) was used to remove non-coding RNA sequences

(tRNA, tmRNA, 5S, 16S, 18S, 23S, and 28S)[97]. The proportion of reads removed per

sample is summarized in Supplementary Table *S3* and Figure *S20*, providing an

overview of rRNA/tRNA content prior to filtering.

Gene expression dynamics in metatranscriptomes. InStrain (v1.8.1)[98] with the

parameter (--min_cov 0) was used on strain-cluster rMAGs recovered from the same

sample to get the coverage of each gene using the clean reads from each

metagenome and each metatranscriptome. The gene coverage in each strain-cluster

rMAG was received by using the cluster database generated when dereplicate

genomes at 99% ANI using dRep (v3.2.2)[86]. The mean coverage of 30S ribosomal

proteins were used to calculate the normalization factor of each gene from the strain-

cluster rMAGs in each metatranscriptome sample. The genes with normalization

expression level greater than two were defined as overexpressed genes. The average

expression level of each gene in each sample was calculated and subjected to

Principal Coordinates Analysis (PCoA) based on Bray-Curtis dissimilarity. Average

gene expression levels were also used to calculate Spearman correlations with TU

and radioactivity. In addition, a presence-absence matrix was constructed using gene

annotation profiles predicted from metabolic potential analyses.

Species-level clustering and calculation of evolutionary metrics. Filtered reads

from each metagenome and metatranscriptome were mapped to the cleaned MAGs

output from FCS-GX (v0.5.5) of the same sample using strobealign (v0.13.0)[89] in

CoverM (v0.7.0; default parameters)[90]. Open reading frames (ORFs) of each MAG

were predicted using Prodigal (v2.6.3; –p single)[78] and used as input for inStrain

(v1.8.1)[98]. Population statistics and nucleotide metrics, including nucleotide diversity

(SNVs/kbp) and nonsynonymous to synonymous mutation ratios (pN/pS) were

calculated from these mappings using the profile module of the inStrain (v1.8.1; default

parameters)[98] at genome and gene levels. The 4075 MAGs reconstructed from all

individual assemblies (n = 60) were combined and dereplicated for species-level

clustering using dRep (v3.2.2)[86] at 95% ANI. A total of 1752 MAGs with the highest

genome quality from each species cluster were designated as the representative

species (species-cluster rMAGs). The species-cluster rMAGs with over three

members in the cluster database generated when dereplicate genomes at 95% ANI

were selected for the final calculation of microdiversity metrics such as coverage,

breadth, SNPs density, nucleotide diversity, and pN/pS ratios.

For SNP calling, the number of quality-filtered reads mapping to the position had to be

at least 5× coverage and 5% SNP frequency, and the variant based on reads had to

be lower than the expected sequencing error rate ($1e^{-6}$). Genome-level evolutionary

metrics based on both metagenomic and metatranscriptomic data from mining sites

were compared with those from background aquifer samples at the same location

(Table *S4-S5*). To assess the genomic impact of radiochemical stress, four species-

cluster rMAGs showing increased nucleotide diversity and detected in at least two

groups (BG, MT, and PM) with a minimum of three samples per group were selected.

Gene annotations for the four species-cluster rMAGs were generated using

MetaCerberus v1.3.1[99]. All member genomes within the four species-cluster were

used to identify representative genes using mmseqs2 v17.b804f[100], from which

median nucleotide diversity was calculated from both metagenomic and

metatranscriptomic datasets across the BG, MT, and PM groups (*Table S6-S7*).

Statistical analyses and data visualization. Spearman correlation analyses were

performed in the R platform (http://cran.r-project.org), with p-values adjusted using the

Benjamini-Hochberg procedure at a minimum significance threshold of 0.05[101]. Co-

occurrence networks were constructed using the *igraph* package with default parameters[102]. Only Spearman correlations with p-values < 0.001 and |ρ|> 0.6 between pairs of rMAGs were considered statistically significant, contributing to the construction of a reliable network. The resulting correlation network was imported into Gephi (v0.9.2) for visualization[103]. The Spearman correlations between over-expressed genes from strain-cluster rMAGs and hydrogeochemical variables were visualized using the *pheatmap* package in R (v4.4.2)[104]. To assess sample similarity and identify factors shaping community clustering, Principal Coordinates Analysis (PCoA) and Non-metric Multidimensional Scaling (NMDS) were performed based on Bray–Curtis dissimilarity of microbial community composition. Relationships between geochemical parameters and biological features—including community dissimilarities, gene expression levels, nucleotide diversity, and pN/pS ratios—were further evaluated using *Bioenv* analysis (*vegan* package, R v4.4.2)[105], which identifies the combination of environmental variables best explaining the observed biological variation.

ACKNOWLEDGEMENTS

This study was supported by the National Natural Science Foundation of China grant nos. 42130509 (to H.G.) and 42072273 (to W.X.), the Deep Earth Probe and Mineral Resources Exploration-National Science and Technology Major Project grant no. 2024ZD1000601 (to W.X.), the Fundamental Research Funds for the Central Universities grant no. 590223006 (to W.X.) and the 111 Project grant no. B20010 (to H.G.); and by the European Research Council Synergy Grant "Archean Park" under grant no. 101118631 (to A.J.P.) and the German Research Foundation grant no. PR 1603/4-1 (to A.J.P.). We thank Prof. Rizlan Bernier-Latmani (Environmental Microbiology Laboratory, École Polytechnique Fédérale de Lausanne, Switzerland) for valuable comments on the manuscript and for providing access to laboratory resources.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

Ethics Statement

All sampling and sample collection were performed with permission from the local environmental authorities and in compliance with all applicable regulations governing

808 environmental research in China. No human subjects or animal samples were involved
809 in this study.
810
811 CODE AVAILABILITY STATEMENT
812 All code used in this manuscript is publicly available and referenced in the Materials
813 and Methods section.
814

REFERENCES
1. Xie, X. *et al.* Groundwater Quality and Public Health. *Annu. Rev. Environ.*
Resour. **48**, 395–418 (2023).
2. Orsi, W. D., Edgcomb, V. P., Christman, G. D. & Biddle, J. F. Gene expression
in the deep biosphere. *Nature* **499**, 205–208 (2013).
3. Magnabosco, C. *et al.* The biomass and biodiversity of the continental
subsurface. *Nat. Geosci.* **11**, 707–717 (2018).
4. Beaver, R. C. & Neufeld, J. D. Microbial ecology of the deep terrestrial
subsurface. *ISME J.* **18**, wrae091 (2024).
5. Bornemann, T. L. V. *et al.* Genetic diversity in terrestrial subsurface ecosystems
impacted by geological degassing. *Nat. Commun.* **13**, 1–12 (2022).
6. Patsis, A. C., Schuler, C. J., Toner, B. M., Santelli, C. M. & Sheik, C. S. The
potential for coupled organic and inorganic sulfur cycles across the terrestrial
deep subsurface biosphere. *Nat. Commun.* **16**, 3827 (2025).
7. Probst, A. J. *et al.* Differential depth distribution of microbial function and
putative symbionts through sediment-hosted aquifers in the deep terrestrial
subsurface. *Nat. Microbiol.* **3**, 328–336 (2018).
8. He, C. *et al.* Genome-resolved metagenomics reveals site-specific diversity of
episymbiotic CPR bacteria and DPANN archaea in groundwater ecosystems.
Nat. Microbiol. **6**, 354–365 (2021).
9. Amundson, K. K., Borton, M. A. & Wilkins, M. J. Anthropogenic impacts on the
terrestrial subsurface biosphere. *Nat. Rev. Microbiol.* **23**, 147–161 (2025).
10. Vigneron, A. *et al.* Succession in the petroleum reservoir microbiome through an
oil field production lifecycle. *ISME J.* **11**, 2141–2154 (2017).
11. Ning, D. *et al.* Environmental stress mediates groundwater microbial community
assembly. *Nat. Microbiol.* **9**, 490–501 (2024).

12.   Jones, D. O. B. *et al.* Long-term impact and biological recovery in a deep-sea
842   mining track. *Nature* https://doi.org/10.1038/s41586-025-08921-3 (2025)
843   doi:10.1038/s41586-025-08921-3.

844   13.   Li, G. & Yao, J. A Review of In Situ Leaching (ISL) for Uranium Mining. *Mining*
845   **4**, 120–148 (2024).

846   14.   Jroundi, F. *et al.* Profiling native aquifer bacteria in a uranium roll-front deposit
847   and their role in biogeochemical cycle dynamics: Insights regarding in situ
848   recovery mining. *Sci. Total Environ.* **721**, (2020).

849   15.   Vrionis, H. A. *et al.* Microbiological and geochemical heterogeneity in an in situ
850   uranium bioremediation field site. *Appl. Environ. Microbiol.* **71**, 6308–6318
851   (2005).

852   16.   Chen, H. *et al.* Uranium contamination mediating soil and ore microbial
853   community assembly at four mining sites, South China. *Front. Microbiol.* **16**,
854   1553072 (2025).

855   17.   Zhu, F., Zhao, B., Min, W. & Li, J. Characteristics of groundwater microbial
856   communities and the correlation with the environmental factors in a
857   decommissioned acid in-situ uranium mine. *Front. Microbiol.* **13**, 1078393
858   (2023).

859   18.   Coral, T. *et al.* Microbial communities associated with uranium in-situ recovery
860   mining process are related to acid mine drainage assemblages. *Sci. Total
861   Environ.* **628–629**, 26–35 (2018).

862   19.   Van Nostrand, J. D. *et al.* Dynamics of Microbial Community Composition and
863   Function during In Situ Bioremediation of a Uranium-Contaminated Aquifer.
864   *Appl. Environ. Microbiol.* **77**, 3860–3869 (2011).

865   20.   Li, B. *et al.* Bacterial Community Shift and Coexisting/Coexcluding Patterns
866   Revealed by Network Analysis in a Uranium-Contaminated Site after
867   Bioreduction Followed by Reoxidation. *Appl. Environ. Microbiol.* **84**, e02885-17
868   (2018).

869   21.   Sharon, I. *et al.* Accurate, multi-kb reads resolve complex populations and
870   detect rare microorganisms. *Genome Res.* **25**, 534–543 (2015).

871   22.   Bowers, R. M. *et al.* Minimum information about a single amplified genome
872   (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and
873   archaea. *Nat. Biotechnol.* **35**, 725–731 (2017).

23. *Energy, Electricity and Nuclear Power Estimates for the Period up to 2050.* (INTERNATIONAL ATOMIC ENERGY AGENCY, Vienna, 2024).

24. Monnet, A., Gabriel, S. & Percebois, J. Long-term availability of global uranium resources. *Resour. Policy* **53**, 394–407 (2017).

25. Lu, C. *et al.* Multi-Isotope Based Identification and Quantification of Oxygen Consuming Processes in Uranium Hosting Aquifers With $CO_2 + O_2$ In Situ Leaching. *Water Resour. Res.* **59**, e2022WR033980 (2023).

26. Lian, G. *et al.* Genome-Resolved Metagenomic Insights into Aquifer Microbiome Adaptations Following Two Decades of Neutral In-Situ Leaching. *J. Geophys. Res. Biogeosciences* **130,** e2024JG008587 (2025).

27. Lu, C. *et al.* Characteristics of Dissolved Organic Matter in Uranium Hosting Aquifers and Potential Molecular Transformation During Neutral In Situ Leaching. *J. Geophys. Res. Biogeosciences* **129**, e2023JG007851 (2024).

28. Lu, C. *et al.* Natural Attenuation of Groundwater Uranium in Post-Neutral-Mining Sites Evidenced from Multiple Isotopes and Dissolved Organic Matter. *Environ. Sci. Technol.* **58**, 12674–12684 (2024).

29. Asghar, F. *et al.* Geochemical characteristics and uranium neutral leaching through a co2 + o2 system—an example from uranium ore of the elzpa ore deposit in Pakistan. *Metals* **10**, 1–19 (2020).

30. Ruiz, O., Thomson, B., Cerrato, J. M. & Rodriguez-Freire, L. Groundwater restoration following in-situ recovery (ISR) mining of uranium. *Appl. Geochem.* **109**, 104418 (2019).

31. Wani, A. K., Akhtar, N., Sher, F., Navarrete, A. A. & Américo-Pinheiro, J. H. P. Microbial adaptation to different environmental conditions: molecular perspective of evolved genetic and cellular systems. *Arch. Microbiol.* **204**, 144 (2022).

32. Newsome, L., Morris, K. & Lloyd, J. R. The biogeochemistry and bioremediation of uranium and other priority radionuclides. *Chem. Geol.* **363**, 164–184 (2014).

33. Carvalho, F. *et al.* The environmental behaviour of uranium. *Int. At. Energy Agency* https://doi.org/Retrieved%2520from:%2520https://www.iaea.org/publications/14688/the-environmental-behaviour-of-uranium (2023) doi:Retrieved%20from:%20https://www.iaea.org/publications/14688/the-environmental-behaviour-of-uranium.

34. Huang, L.-N., Kuang, J.-L. & Shu, W.-S. Microbial Ecology and Evolution in the
Acid Mine Drainage Model System. *Trends Microbiol.* **24**, 581–593 (2016).

35. Chen, L. *et al.* Comparative metagenomic and metatranscriptomic analyses of
microbial communities in acid mine drainage. *ISME J.* **9**, 1579–1592 (2015).

36. Hemme, C. L. *et al.* Metagenomic insights into evolution of a heavy metal-
contaminated groundwater microbial community. *ISME J.* **4**, 660–672 (2010).

37. Pal, A., Bhattacharjee, S., Saha, J., Sarkar, M. & Mandal, P. Bacterial survival
strategies and responses under heavy metal stress: a comprehensive overview.
Crit. Rev. Microbiol. **48**, 327–355 (2022).

38. Mackelprang, R. *et al.* Metagenomic analysis of a permafrost microbial
community reveals a rapid response to thaw. *Nature* **480**, 368–371 (2011).

39. Orsi, W. D. Ecology and evolution of seafloor and subseafloor microbial
communities. *Nat. Rev. Microbiol.* **16**, 671–683 (2018).

40. Becraft, E. D. *et al.* Evolutionary stasis of a deep subsurface microbial lineage.
ISME J. **15**, 2830–2842 (2021).

41. Da Cruz Nizer, W. S. *et al.* Oxidative stress responses in biofilms. *Biofilm* **7**,
100203 (2024).

42. Nguyen, H., Ybarra, A., Başağaoğlu, H. & Shindell, O. Biofilm viscoelasticity and
nutrient source location control biofilm growth rate, migration rate, and
morphology in shear flow. *Sci. Rep.* **11**, 1–17 (2021).

43. Galhardo, R. S., Hastings ,P. J. & and Rosenberg, S. M. Mutation as a Stress
Response and the Regulation of Evolvability. *Crit. Rev. Biochem. Mol. Biol.* **42**,
399–435 (2007).

44. Evans, M. V. *et al.* In situ transformation of ethoxylate and glycol surfactants by
shale-colonizing microorganisms during hydraulic fracturing. *ISME J.* **13**, 2690–
2700 (2019).

45. Zhou, J. *et al.* Stochasticity, succession, and environmental perturbations in a
fluidic ecosystem. *Proc. Natl. Acad. Sci.* **111**, E836–E845 (2014).

46. MacLean, R. C., Torres-Barceló, C. & Moxon, R. Evaluating evolutionary
models of stress-induced mutagenesis in bacteria. *Nat. Rev. Genet.* **14**, 221–
227 (2013).

47. Mojica, E. A. & Kültz, D. Physiological mechanisms of stress-induced evolution.
J. Exp. Biol. **225**, jeb243264 (2022).

48. Couce, A. *et al.* Changing fitness effects of mutations through long-term

bacterial evolution. *Science* **383**, eadd1417 (2024).

49. Denamur, E. & Matic, I. Evolution of mutation rates in bacteria. *Mol. Microbiol.*

60, 820–827 (2006).

50. Maharjan, R. P. *et al.* Mutation accumulation and fitness in mutator

subpopulations of *Escherichia coli*. *Biol. Lett.* **9**, 20120961 (2013).

51. Custer, G. F., Bresciani, L. & Dini-Andreote, F. Toward an integrative framework

for microbial community coalescence. *Trends Microbiol.* **32**, 241–251 (2024).

52. Shade, A. Microbiome rescue: directing resilience of environmental microbial

communities. *Curr. Opin. Microbiol.* **72**, 102263 (2023).

53. Yamamichi, M., Letten, A. D. & Schreiber, S. J. Eco-evolutionary maintenance

of diversity in fluctuating environments. *Ecol. Lett.* **26**, (2023).

54. Jinks-Robertson, S. & Bhagwat, A. S. Transcription-Associated Mutagenesis.

Annu. Rev. Genet. **48**, 341–359 (2014).

55. Tenaillon, O. *et al.* Tempo and mode of genome evolution in a 50,000-

generation experiment. *Nature* **536**, 165–170 (2016).

56. Wei, W. *et al.* Rapid evolution of mutation rate and spectrum in response to

environmental and population-genetic challenges. *Nat. Commun.* **13**, 4752

(2022).

57. Ferrare, J. T. & Good, B. H. Evolution of evolvability in rapidly adapting

populations. *Nat. Ecol. Evol.* **8**, 2085–2096 (2024).

58. Bruckbauer, S. T. & Cox, M. M. Experimental evolution of extremophile

resistance to ionizing radiation. *Trends Genet.* **37**, 830–845 (2021).

59. Daly, M. J. A new perspective on radiation resistance based on Deinococcus

radiodurans. *Nat. Rev. Microbiol.* **7**, 237–245 (2009).

60. Pavlopoulou, A. *et al.* Unraveling the mechanisms of extreme radioresistance in

prokaryotes: Lessons from nature. *Mutat. Res. Mutat. Res.* **767**, 92–107 (2016).

61. Martincorena, I., Seshasayee, A. S. N. & Luscombe, N. M. Evidence of non-

random mutation rates suggests an evolutionary risk management strategy.

Nature **485**, 95–98 (2012).

62. Ferenci, T. Irregularities in genetic variation and mutation rates with

environmental stresses. *Environ. Microbiol.* **21**, 3979–3988 (2019).

63. Foster, P. L. Stress-Induced Mutagenesis in Bacteria. *Crit. Rev. Biochem. Mol.*

Biol. **42**, 373–397 (2007).

64. Woodcroft, B. J. *et al.* Genome-centric view of carbon processing in thawing permafrost. *Nature* **560**, 49–54 (2018).

65. Mehrshad, M. *et al.* Energy efficiency and biological interactions define the core microbiome of deep oligotrophic groundwater. *Nat. Commun.* **12**, 1–12 (2021).

66. Jones, R. M., Goordial, J. M. & Orcutt, B. N. Low Energy Subsurface Environments as Extraterrestrial Analogs. *Front. Microbiol.* **9**, 1605 (2018).

67. Hoehler, T. M. & Jørgensen, B. B. Microbial life under extreme energy limitation. *Nat. Rev. Microbiol.* **11**, 83–94 (2013).

68. Jørgensen, B. B. & Marshall, I. P. G. Slow Microbial Life in the Seabed. *Annu. Rev. Mar. Sci.* **8**, 311–332 (2016).

69. Garber, A. I., Ramírez, G. A. & D'Hondt, S. Genomic stasis over millions of years in subseafloor sediment. *Environ. Microbiol.* **26**, e16674 (2024).

70. Schopf, J. W. *et al.* Sulfur-cycling fossil bacteria from the 1.8-Ga Duck Creek Formation provide promising evidence of evolution's null hypothesis. *Proc. Natl. Acad. Sci.* **112**, 2087–2092 (2015).

71. Jia, J. *et al.* Mineralogy and geochemistry of carbonate cement in sandstone and implications for mineralization of the Qianjiadian sandstone-hosted uranium deposit, southern Songliao Basin, China. *Ore Geol. Rev.* **123**, 103590 (2020).

72. Fan, Y. *et al.* The behavior of pyrite during in-situ leaching of uranium by $CO_2 + O_2$: A case study of the Qianjiadian uranium deposit in the Songliao Basin, northeastern China. *Ore Geol. Rev.* **169**, 106085 (2024).

73. Rong, H. *et al.* Effects of basic intrusions on REE mobility of sandstones and their geological significance: A case study from the Qianjiadian sandstone-hosted uranium deposit in the Songliao Basin. *Appl. Geochem.* **120**, 104665 (2020).

74. Qiu, W. *et al.* What chemical reaction dominates the CO_2 and O_2 in-situ uranium leaching? Insights from a three-dimensional multicomponent reactive transport model at the field scale. *Appl. Geochem.* **148**, 105522 (2023).

75. Xu, L., Wang, H., Liu, N. & Du, Z. Waste Water Treatment of $CO_2 + O_2$ in-situ Leaching Uranium. *Uranium Min. Metall.* **31**, 96–99 (2012).

76. Rodriguez-R, L. M., Gunturu, S., Tiedje, J. M., Cole, J. R. & Konstantinidis, K. T. Nonpareil 3: Fast Estimation of Metagenomic Coverage and Sequence Diversity. *mSystems* **3**, e00039-18 (2018).

77. Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. MetaSPAdes: A new
versatile metagenomic assembler. *Genome Res.* **27**, 824–834 (2017).

78. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation
site identification. *BMC Bioinformatics* **11**, 119 (2010).

79. Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition.
Nat. Methods **11**, 1144–1146 (2014).

80. Kang, D. D., Froula, J., Egan, R. & Wang, Z. MetaBAT, an efficient tool for
accurately reconstructing single genomes from complex microbial communities.
PeerJ **2015**, 1–15 (2015).

81. Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for
genome-resolved metagenomic data analysis. *Microbiome* **6**, 1–13 (2018).

82. Wu, Y. W., Simmons, B. A. & Singer, S. W. MaxBin 2.0: An automated binning
algorithm to recover genomes from multiple metagenomic datasets.
Bioinformatics **32**, 605–607 (2016).

83. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W.
CheckM:assessing the quality of microbial genomes recovered from isolates,
singlecells, andmetagenomes. *Genome Res.* 1043–1055 (2015)
doi:10.1101/gr.186072.114.Freely.

84. Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a
dereplication, aggregation and scoring strategy. *Nat. Microbiol.* **3**, 836–843
(2018).

85. Astashyn, A. *et al.* Rapid and sensitive detection of genome contamination at
scale with FCS-GX. *Genome Biol.* **25**, 60 (2024).

86. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. DRep: A tool for fast and
accurate genomic comparisons that enables improved genome recovery from
metagenomes through de-replication. *ISME J.* **11**, 2864–2868 (2017).

87. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk v2:
memory friendly classification with the genome taxonomy database.
Bioinformatics **38**, 5315–5316 (2022).

88. Letunic, I. & Bork, P. Interactive tree of life (iTOL) v5: An online tool for
phylogenetic tree display and annotation. *Nucleic Acids Res.* **49**, W293–W296
(2021).

89. Sahlin, K. Strobealign: flexible seed size enables ultra-fast and accurate read
alignment. *Genome Biol.* **23**, 260 (2022).

90. Aroney, S. T. N. *et al.* CoverM: Read alignment statistics for metagenomics. Preprint at https://doi.org/10.48550/arXiv.2501.11217 (2025).

91. Zhou, Z. *et al.* METABOLIC: high-throughput profiling of microbial genomes for functional traits, metabolism, biogeochemistry, and community-scale functional networks. *Microbiome* **10**, 1–23 (2022).

92. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND. *Nat. Methods* **12**, 59–60 (2014).

93. Mistry, J., Finn, R. D., Eddy, S. R., Bateman, A. & Punta, M. Challenges in homology search: HMMER3 and convergent evolution of coiled-coil regions. *Nucleic Acids Res.* **41**, e121–e121 (2013).

94. Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile open source tool for metagenomics. *PeerJ* **4**, e2584 (2016).

95. Ruff, S. E. *et al.* A global comparison of surface and subsurface microbiomes reveals large-scale biodiversity gradients, and a marine-terrestrial divide. *Sci. Adv.* **10**, eadq0645 (2024).

96. Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

97. Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217 (2012).

98. Olm, M. R. *et al.* inStrain profiles population microdiversity from metagenomic data and sensitively detects shared microbial strains. *Nat. Biotechnol.* **39**, 727–736 (2021).

99. Figueroa Iii, J. L., Dhungel, E., Bellanger, M., Brouwer, C. R. & White Iii, R. A. MetaCerberus: distributed highly parallelized HMM-based processing for robust functional annotation across the tree of life. *Bioinformatics* **40**, btae119 (2024).

100. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

101. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **57**, 289–300 (1995).

102. Csádi, G. & Nepusz, T. The igraph software package for complex network research. *InterJournal Complex Syst.* **1695**, 1–9 (2006).

1076   103. Bastian, M., Heymann, S. & Jacomy, M. Gephi: An open source software for

1077   exploring and manipulating networks. BT - International AAAI Conference on

1078   Weblogs and Social. *Int. AAAI Conf. Weblogs Soc. Media* 361–362 (2009).

1079   104. Kolde, R. *Pheatmap: Pretty Heatmaps*. (2025).

1080   105. Oksanen, J. *et al.* Vegan: community ecology package. *No Title* (2001).

1081