Dosage compensation and meiotic sex chromosome inactivation are maintained in the absence of selection

Darren J. Parker[1,2], Zoé Dumas[1], Rocío Gómez Lencero[3], Jean-Marc Aury[4], Marjorie Labedan[1], Patrick Tran Van[1], Benjamin Istace[4], Corinne Cruaud[5], Karine Labadie[5], Benjamin Noel[4], Susana Freitas[1], Jelisaveta Djordjevic[1], and Tanja Schwander[1]

1. Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland.
2. School of Natural and Environmental Sciences, Bangor University, Bangor, UK.
3. Department of Biology, Universidad Autónoma de Madrid, 28049 Madrid, Spain
4. Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Université Evry, Université Paris-Saclay, 91057 Evry, France
5. Genoscope, Institut François Jacob, CEA, CNRS, Université Evry, Université Paris-Saclay, 91057 Evry, France

Abstract

Dosage compensation and meiotic sex chromosome inactivation (MSCI) are key mechanisms regulating gene expression from the X chromosome in male-heterogametic species. While the convergent evolution of these mechanisms is well-documented, their evolutionary fate under relaxed selection remains poorly understood. Here, we test whether dosage compensation and MSCI persist in the absence of selection by investigating three independently derived parthenogenetic species of *Timema* stick insects that occasionally produce rare males. Using chromosome-level genome assemblies, RNA-seq from multiple tissues, and immunocytochemistry, we find that dosage compensation in somatic tissues and meiotic X inactivation are conserved despite the absence of selective pressure in all-female lineages. Surprisingly, expression data and cytological markers indicate that MSCI signatures are even stronger in parthenogenetic males, a pattern driven by prolonged autosomal transcription during meiosis. These results indicate that X-targeting dosage compensation and MSCI mechanisms are either weakly selected against or evolutionarily constrained, while autosomal mis-expressing during meiosis occurs rapidly following the relaxation of selection, shedding light on the stability of X chromosome regulation.

Significance Statement:

Sex chromosomes are regulated by specialized mechanisms that balance gene expression between males and females in somatic tissues (dosage compensation) and silence the X chromosome during male meiosis (meiotic sex chromosome inactivation, MSCI). These processes are thought to decay when they become redundant, yet suitable systems to test this prediction are rare. We studied rare males from parthenogenetic stick insect lineages that have been evolving without sex for up to 1.5 million years. Surprisingly, both dosage

42 compensation and MSCI remain fully functional, despite relaxed selection. Instead, we find
43 altered autosomal expression during meiosis. Our results reveal that sex chromosome
44 regulation is evolutionarily stable and constrained, persisting even after the loss of its purpose.
45
46

Introduction

Sexual reproduction is key to the diversity of life and the main driver of species' lifecycles. In eukaryotic species with distinct male and female sexes, sex determination is frequently governed by sex chromosomes such as X and Y chromosomes in male heterogametic species. Because males carry only a single X chromosome while females carry two, X chromosome expression is regulated by highly specialized mechanisms: dosage compensation (in somatic cells) and meiotic sex chromosome inactivation (MSCI, in germ cells). Dosage compensation is a mechanism that balances the expression of X-linked genes between sexes. It has evolved independently across a wide range of taxa (Disteche, 2012; Mank, 2013; Gu and Walters, 2017) including mammals (Brockdorff and Turner, 2015), reptiles (Marin *et al.*, 2017), fish (Darolti *et al.*, 2019; Metzger *et al.*, 2021), nematodes (Meyer, 2000), and insects (Prince *et al.*, 2010; Conrad and Akhtar, 2012; Vicoso and Bachtrog, 2015; Pal and Vicoso, 2015; Rayner *et al.*, 2021; Chauhan *et al.*, 2021; Parker *et al.*, 2022). However, the X is typically not dosage compensated in gonads, where expression from the sex chromosome(s) is further often suppressed via MSCI (Disteche, 2012; Djordjevic et al., 2024; Mank & Ellegren, 2009; Rayner et al., 2021; Vicoso et al., 2013). Specifically, MSCI involves the inactivation of heteromorphic sex chromosomes during meiosis through the addition of repressive chromatin marks, likely due to the inability of the X and the degraded (or completely lacking) Y to synapse at the onset of meiosis (Burgoyne et al., 2009).

While far from ubiquitous (McKee and Handel, 1993; Mank, 2013), dosage compensation and MSCI are typically maintained across clades for millions of years. Furthermore, a breadth of studies on both mechanisms indicate that they can evolve convergently as new sex chromosomes arise, involving similar or different mechanisms. This is revealed for example by research in dipterans, where X chromosomes have emerged independently from ancestral autosomes, with different dosage compensation mechanisms emerging in parallel (Hamada *et al.*, 2005; Straub *et al.*, 2005; Pease and Hahn, 2012; Davis *et al.*, 2018; Krzywinska *et al.*, 2023; Kalita *et al.*, 2023; Kalita and Keller Valsecchi, 2024; Toups and Vicoso, 2025). Similarly, the independent evolution of MSCI mechanisms in divergent taxa involves both shared and lineage-specific components (e.g. (Namekawa and Lee, 2009; Schoenmakers *et al.*, 2009; Page *et al.*, 2023)).

In contrast to the evidence for the convergent *de novo* evolution of dosage compensation and MSCI, it remains unknown how ancestral dosage compensation and MSCI systems evolve once they become redundant, which would be the case upon the evolution of a novel sex chromosome system. This is in part because suitable model systems with sex chromosome turnovers and accompanying evolution of X-targeting regulatory mechanisms remain understudied. Furthermore, sex chromosome turnover events in species with established dosage compensation and MSCI systems are expected to be rare because the maintenance of active systems could cause extensive misregulation of gene expression on the newly-formed sex chromosomes (Kelley *et al.*, 1995; Lee, 2000; Toups and Vicoso, 2023). A unique opportunity to overcome such constraints is to study the fate and potential decay of dosage compensation and MSCI in all-female parthenogenetic species. Following a transition to parthenogenesis, the X chromosome effectively adopts an autosomal inheritance pattern as it no longer varies in copy numbers since all individuals are female and carry two X copies. As a result, if dosage compensation or MSCI is costly (because of the cost of the machinery or

94 due to maladaptive expression in females) we would expect selection to reduce a species'
95 ability to perform these mechanisms following a transition to parthenogenesis. Such a
96 reduction is difficult to observe in parthenogenetic species as we need to examine expression
97 in both sexes. This barrier can be overcome in some systems if rare males are produced.
98
Here we take advantage of such rare males to examine dosage compensation and MSCI in
three species of parthenogenetic *Timema* stick insects, using a combination of new
chromosome-level genome assemblies, RNA sequencing of eight different tissues, and cell
biology. *Timema* have had multiple independent transitions to parthenogenetic reproduction
over the last 0.5-1.5 million years (Schwander and Crespi, 2009; Schwander *et al.*, 2011; Bast
et al., 2018; Jaron *et al.*, 2022). Previously we have found that sexual *Timema* species show
complete dosage compensation in somatic tissues throughout development, and MSCI in the
male germline (Parker *et al.*, 2022; Dumas *et al.*, 2024; Djordjevic *et al.*, 2025). This targeted
regulation of the X chromosome likely involves general molecular mechanisms shared across
insects (Djordjevic *et al.*, 2025) given that the stick insect X chromosome likely evolved prior
to the diversification of insects over 450 mya (Toups and Vicoso, 2023). We can examine
changes in dosage compensation and MSCI in parthenogenetic *Timema* as they occasionally
produce rare males via two different processes. Some males arise from aneuploidy of the X
chromosome (Schwander *et al.*, 2013), which since *Timema* have an XX/X0 sex-determination
system, results in the production of a male. Others develop as a consequence of very rare
instances of sex within parthenogenetic populations (Freitas *et al.*, 2023). Contrary to our
prediction we find that the targeted regulation of the X chromosome underlying dosage
compensation and MSCI has been conserved in each of the three parthenogenetic lineages,
with relaxed selection affecting autosomal rather than X-linked expression in male meiotic
cells. These results indicate that selection against X-regulatory mechanisms is weak or that
changes to dosage compensation and MSCI mechanisms are highly constrained.

Results and Discussion

Genome assemblies reveal a highly conserved X chromosome contrasted by autosomal fissions and fusions

We generated chromosome-level genome assemblies for two of the three sexual species of *Timema* used in this study, *T. cristinae* and *T. podura*, to complement the available assembly of *T. poppense* (Table S1). In both new assemblies, the number of large scaffolds assembled (hereafter referred to as chromosomes) matched previous karyotype information (Schwander and Crespi, 2009) and coverage of male- vs female-derived short reads were used to identify the X chromosome (Supplemental Info). Annotation using a combination of gene predictions, protein, and RNAseq evidence from each species and its parthenogenetic sister species and inference of 8118 one-to-one orthologs (see Methods) revealed that the X chromosome content has not changed in any of the sexual or asexual species, as expected given the long-term conservation of X chromosome content in stick insects (Simon et al., 2019; Parker et al., 2022). Using comparisons with a stick insect (*Bacillus rossius*) from the Euphasmatodea clade (which split over 120 million years ago from *Timema* (Bank and Bradler, 2022), we further find that even X chromosome synteny is highly conserved, while many rearrangements are apparent on the autosomes (Fig. 1). Specifically, the change in karyotype from 14 to 12 chromosomes during *Timema* evolution (Schwander and Crespi, 2009) involved at least two fissions and three fusions involving *T. podura* chromosomes 2, 3, 5 and 10 (Fig. 1, S1).

Fig. 1 | Autosomal genomic instability contrasts with X chromosome conservation. A. Simplified phylogeny of the used species, highlighting the independently evolved parthenogenetic *Timema* species and their closest sexual relatives. *Timema* are the sister clade to all other stick insects (including B. *rossius*), and branches reflecting this split have been shortened for display. B. Chord diagram showing synteny between the assemblies of sexual species, based on 4241 one-to-one orthologs across the focal *Timema* genomes and a distantly related stick insect (*Bacillus rossius*). See Fig. S1 for synteny based on 8118 one-to-one orthologs across *Timema* only.

Dosage compensation in somatic tissues is not reduced in parthenogenetic species

Dosage compensation between the sexes in parthenogenetic species was very similar to that observed in sexual species. In all seven non-reproductive tissues (antennae, brain, defence glands, gut, fat body, femur, and tarsi) we observe almost complete dosage compensation in all species (Fig. 2). As expected, we observe some variation in the expression of X-linked genes between the sexes for specific tissues and species (resulting in small but significant differences in expression of autosomal versus X-linked genes). Such variation is typical in studies of dosage compensation (e.g., (Rayner *et al.*, 2021; Hu *et al.*, 2022; Li *et al.*, 2024) and can be interpreted as sex-specific regulation of individual genes rather than a consequence of variable or incomplete dosage compensation. If dosage compensation was reduced in parthenogenetic species, we would expect to see a larger deviation from complete dosage compensation in parthenogenetic species and an accompanying lower male-to-female expression ratio for the X in parthenogenetic species when compared to the sexual sister species for tissues with almost complete dosage compensation. We do not observe this, instead finding only small variations in X-linked expression with no consistent pattern (Fig. 2, Table S2), showing that dosage compensation is maintained across tissues even under relaxed selection for up to 1.5 million years.

Fig. 2 | Male to female expression ratios for the X and autosomes in non-reproductive tissues reveal the maintenance of dosage compensation under relaxed selection. Dashed lines represent a two- fold reduction in expression in males (as expected if there was no dosage compensation). A. Sexual species = *T. cristinae*, parthenogenetic species = *T. monikense*. B. Sexual species = *T. podura*, parthenogenetic species = *T. genevievae*. C. Sexual species = *T. poppense*, parthenogenetic species = *T. douglasi.* Asterisks indicate the significance (FDR) of Wilcoxon tests (***<0.001, **<0.01, *<0.05, NS > 0.05).

179
Parthenogenetic species no longer need to perform dosage compensation as all individuals
are female and carry the same number of sex chromosomes. Under this scenario, we would
expect selection for a reduction in dosage compensation ability for two non-mutually exclusive
reasons. Firstly, as the dosage compensation machinery is complex it is likely costly to
produce and thus selection should act to remove it when it is no longer needed. One reason
we do not observe any consistent changes may be that in insects most of the costs of
producing dosage compensation machinery may be in males, as typically dosage
compensation is achieved by hyper-transcribing the single X in males (Prince *et al.*, 2010;
Conrad and Akhtar, 2012; Mahajan and Bachtrog, 2015; Rose *et al.*, 2016; Richard *et al.*,
2017). That said even if the cost to females was zero we may still expect to see a reduction in
dosage composition ability due to the action of drift. We do not see this, which may mean that
the genes involved in dosage compensation may be pleiotropic, and alterations therefore
deleterious for females. Support for this idea stems from evidence for the maintenance of
dosage compensation-like mechanisms targeting the ancestral X in *Drosophila melanogaster*
fruit flies (Larsson *et al.*, 2004) even tens of myr after a sex chromosome turnover, which
reverted the ancestral X (now referred to as the dot) back to an autosome (Vicoso and
Bachtrog, 2015). Similar to our present study in parthenogenetic species, such reversals
should result in relaxed selection for dosage compensation on the ancestral X (dot)
chromosome. Yet, the dot is still regulated in a chromosome-specific manner, distinct from the
other autosomes and the novel X chromosome. Thus, the dot is regulated via the protein POF
(Larsson *et al.*, 2004), which might reflect vestiges of the former dosage compensation
machinery when the dot was still the X chromosome.
202
A second reason to expect changes in dosage compensation could be if dosage compensation
mechanisms result in maladaptive gene expression in females, i.e., partially unresolved sexual
antagonism (Mank *et al.*, 2011). This factor may not have a strong influence in *Timema* as
unresolved antagonisms are expected to be transient (Flintham *et al.*, 2025) yet the stick insect
X chromosome is very old and highly conserved (Fig. 1) and shares most of its content with
the X chromosome in other insect orders that diverged over 450 million years ago (Toups and
Vicoso, 2023). Any maladaptive gene expression in females as a result of sexual conflict over
dosage compensation would therefore be expected to be very small.
211
While our results show that dosage compensation is maintained in parthenogenetic species
because selection against dosage compensation is weak or because changes to the
underlying mechanisms are constrained, we cannot formally exclude that perhaps not enough
time has passed since the evolution of parthenogenesis to observe any changes to dosage
compensation. While this may be a likely explanation for dosage compensation maintenance
across tissues in the young lineage *T. douglasi* and for the lack of apparent changes in dosage
compensation in whole-body samples of males of parthenogenetic aphid lineages (Defendini
et al., 2025), it is very unlikely for the oldest species used here (*T. genevievae*) with an age
estimate of over 1.5 million years (Schwander *et al.*, 2011; Bast *et al.*, 2018). This is likely
ample time for even quite gradual changes in dosage compensation to be observed, especially
given that *de novo* complete dosage compensation mechanisms can evolve in less than 4
million years (Metzger *et al.*, 2021).
224

A stronger signature of MSCI in parthenogenetic than sexual species driven by autosomal mis-expression during meiosis

We have shown previously that the X chromosome is consistently inactivated throughout male meiosis in sexual *Timema* species (Dumas *et al.*, 2024; Djordjevic *et al.*, 2025). Accordingly, we observe very low expression of X-linked genes in the testes, much lower than the expectation for a lack of dosage compensation (Fig. 3). We do however still observe some expression of the X in males, as expected given the presence of non-meiotic cells in testes. Surprisingly, we see a consistently lower expression of X-linked genes in males from parthenogenetic than sexual species (Fig. 3). This could indicate a higher proportion of meiotic cells in parthenogenetic than sexual males, or, more likely, that autosomal expression is more persistent across meiosis in parthenogenetic as compared to sexual males. In sexual *Timema* males, there are two bursts of autosomal transcription during meiosis (Dumas *et al.*, 2024): a first at the end of prophase I (specifically, during pachytene and diplotene stages), and a second during meiosis II at the spermatid stage.

To detect potential changes to these temporal patterns of autosomal expression in parthenogenetic males, we performed immunolabeling of testis tissue of *T. monikense* and *T. douglasi* using RNA polymerase II, a conserved transcriptional marker. This labeling revealed that autosomal expression during prophase I is longer in parthenogenetic than sexual males. In the former, the RNA polymerase II signal already appears at the onset of prophase I (i.e., during the zygotene stage; Figs 4, S2) and is maintained until diplotene (Figs 4, S2), while in the latter it only appears at pachytene (Dumas *et al.*, 2024). Since males of parthenogenetic species are not exposed to selection, this extension of the autosomal transcription burst during prophase I is most likely a consequence of drift, i.e., it reflects autosomal mis-expression during meiosis in parthenogenetic males.

Fig. 3 | A stronger signature of MSCI in parthenogenetic than sexual species. Gene expression in reproductive tissues: Log$_2$ of male to female expression ratio for the X and autosomes in testes and female gonads. Species codes: Tce = *T. cristinae*, Tms = *T. monikense*, Tpa = *T. podura,* Tge = *T. genevievae,* Tps = *T. poppense* Tdi = *T. douglasi.* Asterisks indicate the significance (FDR) of Wilcoxon tests (***<0.001, **<0.01, *<0.05, NS > 0.05).

Fig. 4 | Misregulation of autosomal gene expression in meiotic cells of parthenogenetic males. RNApol-II distribution in *T. monikense* meiocytes and spermatids. Projections of stack images acquired across *T. monikense* squashed meiocytes (A–T) and spermatids (a–e'), at the stages indicated, stained with DAPI (blue) and double immunolabeled for SMC3 (green) and RNApol-II (magenta). RNApol-II labelling is intense from zygotene to diplotene, and in the round spermatid stage. During all spermatogenesis stages, the X chromosome is extremely reduced in RNApol-II labelling. The position of the X chromosome is indicated with arrowheads. Scale bar: 10 µm. Corresponding image analyses for males of *T. douglasi* are available in Fig. S2)

Conclusion

Our results demonstrate that both dosage compensation and meiotic sex chromosome inactivation (MSCI) are robustly maintained in parthenogenetic stick insects, despite the long-term absence of selective pressures typically associated with male phenotypes. This maintenance suggests that the molecular mechanisms underlying dosage compensation and MSCI are either functionally constrained, potentially due to pleiotropic effects in females, or maintained through weak purifying selection. Notably, the enhanced signature of MSCI in parthenogenetic males appears to be driven by extended autosomal transcription during meiosis, revealing an unexpected consequence of relaxed selection on gene expression. Together, these findings highlight the stability and evolutionary inertia of complex regulatory processes, even in the face of dramatic shifts in reproductive mode.

Methods

Reference genome assemblies

Reference genome assemblies were produced for *T. cristinae* (Bioproject PRJNA1268975/ Accession JBOIUC000000000) and *T. podura* (Bioproject PRJNA1268975/ Accession JBOIUD000000000) from wild collected females (GPS coordinates 34°30'53.2"N, 119°46'45.3"W and 33°47'47.0"N, 116°46'34.6"W). Contigs for the *T. cristinae* assembly were based on PacBio sequencing (43x) and on a combination of Nanopore and Illumina sequencing for *T. podura* (sequenced at 85x respectively 62x coverage). Scaffolding was done using a Hi-C library (sequenced at approximately 69 x and 53 x coverage for *T. cristinae* and *T. podura* respectively) to produce the final reference assembly.

DNA extraction and library preparation

To extract high molecular weight (HMW) DNA, we flash-froze a single female (without gut) in liquid nitrogen and ground it using a Cryomill (Retsch). We then extracted HMW DNA using a G/20 Genomic Tips kit (Qiagen) following the manufacturer's protocols and checked DNA integrity on a pulse field agarose gel.

For *T. cristinae*, a PacBio library was prepared using the Template Prep Kit 1.0 following manufacturer's instructions and sequenced on a single cell of the PacBio SMRT™ system. For *T. podura*, four ONT libraries were prepared following Oxford Nanopore instructions. A first library was prepared using the SQK-LSK108 ligation sequencing kit and was loaded on a MinION R9.4 Flow Cells, and then three libraries were prepared using the SQK-LSK109 ligation sequencing kit and were loaded on three PromethION R9.4.1 Flow Cells. A PCR free Illumina library (based on the same HMW extraction) was prepared using the Kapa Hyper Prep Kit (Roche, Basel, Switzerland), following manufacturer instructions. The library was quantified by qPCR using the KAPA Library Quantification Kit for Illumina Libraries (Roche), and the library profile was assessed using a High Sensitivity DNA kit on an Agilent Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA). The library was then sequenced on an Illumina HiSeq 4000 instrument (Illumina, San Diego, CA, USA), using 150 base-length read chemistry in a paired-end mode.

A Hi-C library was prepared for each species using the Proximo Hi-C Kit. We generated
ground tissue for cross-linking using Cryomill (Retsch) and following manufacturer
instructions, using a different female than the one used for Nanopore and HiSeq sequencing,
from the same natural population. Library construction and sequencing (250 million read pairs)
was outsourced to Phase Genomics (Seattle).

Assembly pipelines

We used different strategies for genome assembly for the two species. For *T. cristinae* raw
PacBio reads were assembled into contigs using Canu (Koren *et al.*, 2017) with the parameter
correctedErrorRate=0.105. For *T. podura* raw Oxford Nanopore reads were filtered using
Filtlong v0.2.0 (https://github.com/rrwick/Filtlong) with the parameters --min_length 1000 --
keep_percent 90 --target_bases 69050000000. The filtered Nanopore reads were then
assembled into contigs using Flye v2.8.1 (Kolmogorov *et al.*, 2019) with --genome-size 1.381g.
All Nanopore reads were mapped against the contigs using minimap2 v2.19 (Li, 2018) with
the parameters -c -x map-ont and a first step of polishing was performed using Racon v1.4.3
(Vaser *et al.*, 2017). Three additional rounds of polishing were conducted: Illumina short reads
were aligned to the contigs using BWA mem v0.7.17 (Li, 2013) and polishing was performed
using Pilon v1.23 (Walker *et al.*, 2014).

Both assemblies were then decontaminated using BlobTools v1.0 (Laetsch and Blaxter, 2017)
under the taxrule "bestsumorder". Hit files were generated after a blastn v2.10.1+ against the
NBCI nt database, searching for hits with an e-value below 1e-25 (parameters: -
max_target_seqs 10 -max_hsps 1 -evalue 1e-25). Contigs without hits to metazoans were
removed from the assemblies. Haplotypic duplications were then filtered out as follows.
Filtered reads were then mapped against the decontaminated genomes using minimap2 and
haplotigs were detected with Purge Haplotigs v1.1.1 (Roach *et al.*, 2018) using the parameters
-l 3 -m 27 -h 190 -j 101 --editor-coarse-resolution 250000 for *T. cristinae* and -l 3 -m 45 -h 195
-j 101 --editor-coarse-resolution 50000 for *T. podura*.

Hi-C reads from each species were then mapped against their respective haploid genome
using Juicer v1.6 (Durand *et al.*, 2016) with the restriction site Sau3AI. Finally, chromosome-
level scaffolding was performed using 3D-DNA v180922 (Dudchenko *et al.*, 2017) with
species-specific parameters, as recommended by the authors. The resulting Hi-C contact
matrices were visualized with Juicebox, and minor polishing was conducted in accordance
with these recommendations. The completeness of both genome assemblies was assessed
with BUSCO v5.1.2 (Seppey *et al.*, 2019) against the *insecta_odb10* dataset using the --long
and --augustus parameters.

Annotation

We annotated the *T. cristinae* and *T. podura* genomes in a similar way as we previously
annotated the *T. poppense* genome (Djordjevic *et al.*, 2025), using a combination of *ab initio*
gene prediction, protein homology, and RNA-seq using the Braker2 pipeline (v. 2.1.6) (Brůna
et al., 2021). Firstly, each genome assembly was soft-masked using RepeatModeler (v. 2.0.2,

options: -LTRStruct, -engine ncbi) (Flynn *et al.*, 2020) and RepeatMasker (v. 4.1.2, options: -engine ncbi, -xsmall) (Smith *et al.*, 2013). For protein evidence, we used the arthropod protein sequences from OrthoDB (v.10.1) (Kriventseva *et al.*, 2019) and the predicted protein sequences for *Timema* from our previous genome assemblies. For RNAseq evidence, we used the RNAseq libraries generated here for *T. cristinae* and *T. monikense* along with publicly available data (163 libraries generated here, 43 public (Bioproject accessions: PRJNA380865, PRJNA679785, PRJNA1295360, PRJNA504764 (Bast *et al.*, 2018; Parker *et al.*, 2019; Jaron *et al.*, 2022))) to annotate the *T. cristinae* genome and for *T. podura* and *T. genevievae* (119 libraries generated here, 29 public (Bioproject accessions: PRJNA380865, PRJNA679785, PRJNA1295360 (Bast *et al.*, 2018; Jaron *et al.*, 2022))) to annotate the *T. podura* genome. This is a total of 354 RNAseq libraries (330 paired-end, 24 single-end) covering over 19 different life stages, tissue, and sex combinations for each species (see Table S3). Reads were quality trimmed with Trimmomatic (v. 0.39, options: ILLUMINACLIP:3:25:6 LEADING:9 TRAILING:9 SLIDINGWINDOW:4:15 MINLEN:80) (Bolger *et al.*, 2014) before mapping to the genome assembly with STAR (v. 2.7.8a, options: --twopassMode Basic) (Dobin *et al.*, 2013). Braker2 was run using protein evidence and RNAseq separately with the gene predictors Augustus (v. 3.4.0) (Stanke *et al.*, 2006) and Genemark (v. 4.72) (Brůna *et al.*, 2020). Following the RNAseq run, UTR predictions were added to the RNAseq gene predictions using GUSHR (v. 1.0) in Braker2 (--addUTR=on). The separate gene predictions were then merged using TSEBRA (v1.0.3) using the pref_braker1.cfg configuration file, which weights RNAseq evidence more strongly than the default option. We then ran BUSCO (v. 5.3.2, insecta_odb10) on the gene regions annotated by Braker2 and on the whole genome assembly. Any genes found by BUSCO but missed by Braker were then added to the annotation (69 genes for *T. cristinae* and 19 genes for *T. podura*). ncRNA genes were predicted using Infernal (v. 1.1.2, minimum e-value 1e-10) (Nawrocki and Eddy, 2013). GO-terms for protein-coding gene predictions were obtained using blastP within OmicsBox (v3.1.2, default parameters) to blast the nr *Drosophila melanogaster* database (taxonomy filter: 7227). 1-to-1 orthologs between were obtained for protein-coding genes using OrthoFinder (version 2.5.5) (Emms and Kelly, 2019) with default options.

Identification of the X chromosome

We used a coverage approach to identify X-linked scaffolds in our new genome assemblies. To do this we mapped publicly available Illumina genome reads from males and females (4 males and 5 females for *T. cristinae* and 4 males and 2 females for *T. podura* (PRJNA670663, PRJNA725673 (Jaron *et al.*, 2022; Parker *et al.*, 2022)) to the respective genome assemblies and their publicly available mtDNA genomes using BWA-MEM (v. 0.7.15). Multi- mapping and poor quality alignments were filtered (removing reads with XA:Z or SA:Z tags or a mapq <30). PCR duplicates were removed with Picard (v. 2.9.0) (http://broadinstitute.github.io/picard/). Coverage was then estimated for 100000 bp sliding windows using BEDTools (v. 2.26.0) (Quinlan and Hall, 2010). To compare coverage between males and females, coverage was first summed for all male and all female libraries per window. Male and female coverage was then normalized by modal coverage to adjust for differences in overall coverage. X- linked windows were then identified using the $\log_2$ ratio of male to female coverage. Autosomal windows should have equal coverage in males and females ($\log_2$ ratio of male to female coverage ≈ 0) and X-linked windows should have half the coverage in males as in females

(log2 ratio of male to female coverage ≈ −1). This approach identified one large scaffold in
each assembly as the X chromosome (Fig. S3-5).

We then repeated these analyses using genome reads from each of the parthenogenetic
species. Female reads were publicly available (5 for *T. monikense*, 6 for *T. genevievae*, and
5 for *T. douglasi*, PRJNA670663 (Jaron *et al.*, 2022)) as were the two male samples for *T.*
douglasi (PRJNA808673, (Larose *et al.*, 2022)). For *T. monikense* and *T. genevievae* we also
generated new Illumina libraries for males (2 for *T. genevievae* and 4 for *T. monikense*) and
an additional female library for *T. genevievae* (Accession number: PRJNA1293928) as
described previously (Jaron *et al.*, 2022) using the carcasses after tissue dissection described
below. Reads were then mapped to the genome of each sexual sister species as described
above (the sexual species genome for *T. douglasi* is our previously published genome *T.*
poppense, Accession number: PRJNA1126215 (Djordjevic *et al.*, 2025)). Each of the newly
generated samples were checked for normal coverage distributions before calculating
coverage ratios.

RNA extraction and sequencing

We generated a total of 282 RNA-seq libraries for eight female tissues (antennae, brain,
defence glands, fat body, femur, tarsi, and gonads) and nine male tissues (the same as the
female plus the male accessory glands) for *T. cristinae*, *T. podura*, *T. monikense*, and *T.*
genevievae following our previously developed methodology (Djordjevic *et al.*, 2025) (see
Table S3 for details). Individuals were collected as nymphs in the field and reared to adulthood
in the laboratory. Prior to dissection all individuals were isolated in Petri dishes and fed on an
artificial diet (a mix of flour, sugar, and water) for two days, allowing their gut contents to be
cleared of plant material. Insects were then anesthetized with CO_2, dissected, and tissues
individually flash-frozen in liquid nitrogen before being stored at -80°C.

TRIzol solution (1 mL) and a small amount of ceramic beads (Sigmund Lindner) were added
to each tissue. Samples were homogenized using a tissue homogenizer (Precellys Evolution;
Bertin Technologies). Chloroform (200 µL) was added to each sample and samples were then
vortexed for 15 seconds and centrifuged for 25 minutes at 12,000 rpm at 4°C. The upper
phase containing the RNA was then transferred to a new 1.5 mL tube with the addition of
isopropanol (650 µL). The samples were vortexed and placed at -20°C overnight. Samples
were then centrifuged for 30 minutes at 12,000 rpm at 4°C. The liquid supernatants were
removed, and the RNA pellet underwent two washes using 80% and 70% ethanol. Each wash
was followed by a 5-minute centrifugation step at 12,000 rpm. Finally, the RNA pellet was
resuspended in nuclease-free water and quantified using a fluorescent RNA-binding dye
(QuantiFluor RNA System) and nanodrop (DS-11 FX).

Library preparation using NEBNext (New England BioLabs) and sequencing on an Illumina
NovaSeq 6000 platform with 100 bp paired-end sequencing (36.2 million read pairs per
sample on average) was outsourced to a sequencing facility (Fasteris, Geneva).

Gene expression analyses

We mapped RNAseq reads from *T. cristinae*, *T. podura*, *T. monikense*, and *T. genevievae* generated here and reads for *T. poppense* and *T. douglasi* we generated previously (Djordjevic *et al.*, 2025) to their respective genomes as described above. This gave 4 replicates for each of the 16 sex and tissue combinations for each of the six species, except for the male samples of parthenogenetic species (*T. monikense, T. genevievae,* and *T. douglasi*) where we had fewer replicates due to the difficulty of obtaining these specimens (n = 2-4 see Table S4 for details). Read counts were obtained using HTSeq v.0.9.1 (Anders *et al.*, 2015), with the following options (htseq-count --order = name --type = exon --idattr = gene_id --stranded = reverse). Expression analyses were performed using the Bioconductor package EdgeR (v 3.42.4) (Robinson *et al.*, 2010), and done separately for each species and tissue. Normalization factors for each library were computed using the TMM method and were used to calculate normalized expression levels (FPKM, Fragments Per Kilobase of transcript per Million mapped reads). mtDNA, rRNA, tRNA and genes with low expression (less than 2 FPKM in 2 or more libraries per sex) were excluded. To examine dosage compensation, we used the $\log_2$ ratio of mean male expression level to female expression level and used Wilcoxon tests (adjusted for multiple testing using Benjamini and Hochberg's (1995) algorithm) to determine if differences in expression between the autosomes and the X chromosome were significant.

Immunolabeling of parthenogenetic male meiotic cells

In order to determine the dynamics of autosomal gene expression during meiosis in parthenogenetic males, we followed the same protocol previously used to characterize autosomal gene expression during meiosis in sexual males (Dumas *et al.*, 2024). In brief, we performed double immunolabeling of SMC3_488 (#AB201542, AbCam) as a marker of cohesin axes (this allows for the assessment of synapsis progression, and hence, cell cycle stage) and RNA polymerase II phosphorylated at serine 2 (p-RNApol-II (ab193468, AbCam); an indicator of transcription). Gonads from adult *T. monikense* and *T. douglasi* males were dissected in 1X PBS, then fixed in paraformaldehyde (2%) and Triton X-100 (0.1%) solution for 15 minutes. Gonads were then squashed on slides coated with poly-L-lysine, and flash frozen in liquid nitrogen. Slides were incubated in PBS 1x for 20 minutes at room temperature, then blocked with a BSA 3% solution (prepared by dissolving 3g of BSA in 100mL of PBS 1X) for 20 minutes. Slides were then incubated overnight at 4°C in a humid chamber with the primary (p-RNA-Pol II), diluted in BSA 3% (1:100), and washed 3x for 5 minutes in PBS 1X. The secondary antibody, anti-rabbit Alexa 594 (711-585-152, Jackson), diluted in BSA 3% (1:100), was applied, and samples were incubated for 45 minutes at room temperature. Further washes were conducted as described above, followed by an extended 10-minute wash in PBS 1X, and by blocking using a 5% Normal Rabbit Serum (NRS) solution (X090210-8, Agilent Technology) for 30 minutes at room temperature. Another 5-minute wash in PBS 1x was performed to remove excess blocking solution. Samples were then incubated with the labelled antibody SMC3_488 (#AB201542, AbCam) at a 1:100 dilution for 1 hour at room temperature, followed by a final series of washes and staining with DAPI (D9542, Sigma-Aldrich) for 3 minutes at room temperature. Finally, after a 5-minute wash in 1x PBS, two drops of Vectashield were added to the slide, which was then covered with a coverslip, and sealed

500 with nail varnish for imaging. Image acquisition was performed at the Cellular Imaging Facility
501 (CIF, University of Lausanne) using a Zeiss LSM 880 Airyscan.

Data

Genome assemblies have been deposited at DDBJ/ENA/GenBank under the accessions
JBOIUC000000000 (*T. cristinae*) and JBOIUD000000000 (*T. podura*). The Illumina and
PromethION sequencing data for *T. podura* are available in the European Nucleotide Archive
under the following project PRJEB94285. Male genomic reads have been deposited at SRA
under the following project PRJNA1293928. RNAseq reads have been deposited at SRA
under the following projects: PRJNA1320679, PRJNA1327219. Code and annotations are
available here: https://github.com/DarrenJParker/Timema_DC_MSCI_code and will be
archived upon acceptance.
511

Acknowledgments

513 This work was supported by the European Research Council Consolidator Grant 864672 ("No
514 Sex No Conflict", T.S.), Swiss FNS grant 31003A_182495 (T.S.), the Genoscope, the
515 Commissariat à l'Énergie Atomique et aux Énergies Alternatives (CEA) and France
516 Génomique (ANR-10-INBS-09–08). We thank the Lausanne Genomic Technologies Facility
517 for PacBio library preparation and sequencing.
518
519

References

Anders S, Pyl PT, Huber W (2015). HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**: 166–169.

Bank S, Bradler S (2022). A second view on the evolution of flight in stick and leaf insects (Phasmatodea). *BMC Ecol Evol* **22**: 62.

Bast J, Parker DJ, Dumas Z, Jalvingh KM, Tran Van P, Jaron KS, *et al.* (2018). Consequences of asexuality in natural populations: insights from stick insects. *Mol Biol Evol* **35**: 1668–1677.

Benjamini Y, Hochberg Y (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol* **57**: 289–300.

Bolger AM, Lohse M, Usadel B (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

Brockdorff N, Turner BM (2015). Dosage compensation in mammals. *Cold Spring Harb Perspect Biol* **7**: a019406.

Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M (2021). BRAKER2: automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. *NAR Genom Bioinform* **3**: lqaa108.

Brůna T, Lomsadze A, Borodovsky M (2020). GeneMark-EP+: eukaryotic gene prediction with self-training in the space of genes and proteins. *NAR Genom Bioinform* **2**: lqaa026.

Chauhan P, Swaegers J, Guillen RAS, Svensson EI, Wellenreuther M, Hansson B (2021). Genome assembly, sex-biased gene expression and dosage compensation in the damselfly *Ischnura elegans*. *Genomics* **113**: 1828–1837.

Conrad T, Akhtar A (2012). Dosage compensation in *Drosophila melanogaster*: epigenetic fine-tuning of chromosome-wide transcription. *Nat Rev Genet* **13**: 123–134.

Darolti I, Wright AE, Sandkam BA, Morris J, Bloch NI, Farré M, *et al.* (2019). Extreme heterogeneity in sex chromosome differentiation and dosage compensation in livebearers. *Proc Natl Acad Sci U S A* **116**: 19031–19036.

Davis RJ, Belikoff EJ, Scholl EH, Li F, Scott MJ (2018). No blokes is essential for male viability and X chromosome gene expression in the Australian sheep blowfly. *Curr Biol* **28**: 1987–1992.e3.

Defendini H, Prunier-Leterme N, Robin S, Lameiras S, Baulande S, Simon J-C, *et al.* (2025). The release of sexual conflict after sex loss is associated with evolutionary changes in gene expression. *Proc Biol Sci* **292**: 20242631.

Disteche CM (2012). Dosage compensation of the sex chromosomes. *Annu Rev Genet* **46**: 537–560.

Djordjevic J, Tran Van P, Toubiana W, Labédan M, Dumas Z, Aury J-M, *et al.* (2025). Dynamics of X chromosome hyper-expression and inactivation in male tissues during stick insect development. *PLoS Genet* **21**: e1011615.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, *et al.* (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, *et al.* (2017). De
novo assembly of the Aedes aegypti genome using Hi-C yields chromosome-length
scaffolds. *Science* **356**: 92–95.

Dumas Z, Toubiana W, Delattre M, Schwander T (2024). Dynamics of recombination, X
inactivation and centromere proteins during stick insect spermatogenesis. *bioRxiv*:
2024.09.05.611401.

Durand NC, Shamim MS, Machol I, Rao SSP, Huntley MH, Lander ES, *et al.* (2016). Juicer
provides a one-click system for analyzing loop-resolution Hi-C experiments. *Cell Syst* **3**:
95–98.

Emms DM, Kelly S (2019). OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol* **20**: 238.

Flintham E, Savolainen V, Otto SP, Reuter M, Mullon C (2025). The maintenance of genetic
polymorphism underlying sexually antagonistic traits. *Evol Lett* **9**: 259–272.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, *et al.* (2020).
RepeatModeler2 for automated genomic discovery of transposable element families.
Proc Natl Acad Sci U S A **117**: 9451–9457.

Freitas S, Parker DJ, Labédan M, Dumas Z, Schwander T (2023). Evidence for cryptic sex in
parthenogenetic stick insects of the genus Timema. *Proceedings of the Royal Society B:
Biological Sciences* **290**: 20230404.

Gu L, Walters JR (2017). Evolution of sex chromosome dosage compensation in animals: a
beautiful theory, undermined by facts and bedeviled by details. *Genome Biol Evol* **9**:
2461–2476.

Hamada FN, Park PJ, Gordadze PR, Kuroda MI (2005). Global regulation of X chromosomal
genes by the MSL complex in Drosophila melanogaster. *Genes Dev* **19**: 2289–2294.

Hu Q-L, Ye Y-X, Zhuo J-C, Huang H-J, Li J-M, Zhang C-X (2022). Chromosome-level
assembly, dosage compensation and sex-biased gene expression in the small brown
planthopper, Laodelphax striatellus. *Genome Biol Evol* **14**: evac160.

Jaron KS, Parker DJ, Anselmetti Y, Van PT, Bast J, Dumas Z, *et al.* (2022). Convergent
consequences of parthenogenesis on stick insect genomes. *Science Advances* **8**:
eabg3842.

Kalita AI, Keller Valsecchi CI (2024). Dosage compensation in non-model insects - progress
and perspectives. *Trends Genet*.

Kalita AI, Marois E, Kozielska M, Weissing FJ, Jaouen E, Möckel MM, *et al.* (2023). The sex-
specific factor SOA controls dosage compensation in Anopheles mosquitoes. *Nature*
623: 175–182.

Kelley RL, Solovyeva I, Lyman LM, Richman R, Solovyev V, Kuroda MI (1995). Expression
of msl-2 causes assembly of dosage compensation regulators on the X chromosomes
and female lethality in Drosophila. *Cell* **81**: 867–877.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA (2019). Assembly of long, error-prone reads
using repeat graphs. *Nat Biotechnol* **37**: 540–546.

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM (2017). Canu: scalable
and accurate long-read assembly via adaptive k-mer weighting and repeat separation.

Genome Res **27**: 722–736.

Kriventseva EV, Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simão FA, *et al.* (2019). OrthoDB v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes for evolutionary and functional annotations of orthologs. *Nucleic Acids Res* **47**: D807–D811.

Krzywinska E, Ribeca P, Ferretti L, Hammond A, Krzywinski J (2023). A novel factor modulating X chromosome dosage compensation in Anopheles. *Curr Biol* **33**: 4697–4703.e4.

Laetsch DR, Blaxter ML (2017). BlobTools: Interrogation of genome assemblies. *F1000Res* **6**: 1287.

Larose C, Lavanchy G, Freitas S, Parker DJ, Schwander T (2022). Facultative parthenogenesis in an asexual stick insect: re-evolution of sex or vestigial sexual capacity? *bioRxiv*: 2022.03.25.485836.

Larsson J, Svensson MJ, Stenberg P, Mäkitalo M (2004). Painting of fourth in genus Drosophila suggests autosome-specific gene regulation. *Proc Natl Acad Sci U S A* **101**: 9728–9733.

Lee JT (2000). Disruption of imprinted X inactivation by parent-of-origin effects at Tsix. *Cell* **103**: 17–27.

Li H (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*: 1303.3997.

Li H (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**: 3094–3100.

Li X, Mank JE, Ban L (2024). The grasshopper genome reveals long-term gene content conservation of the X Chromosome and temporal variation in X Chromosome evolution. *Genome Res* **34**: 997–1007.

Mahajan S, Bachtrog D (2015). Partial dosage compensation in Strepsiptera, a sister group of beetles. *Genome Biol Evol* **7**: 591–600.

Mank JE (2013). Sex chromosome dosage compensation: definitely not for everyone. *Trends Genet* **29**: 677–683.

Mank JE, Hosken DJ, Wedell N (2011). Some inconvenient truths about sex chromosome dosage compensation and the potential role of sexual conflict. *Evolution* **65**: 2133–2144.

Marin R, Cortez D, Lamanna F, Pradeepa MM, Leushkin E, Julien P, *et al.* (2017). Convergent origination of a Drosophila-like dosage compensation mechanism in a reptile lineage. *Genome Res* **27**: 1974–1987.

McKee BD, Handel MA (1993). Sex chromosomes, recombination, and chromatin conformation. *Chromosoma* **102**: 71–80.

Metzger DCH, Sandkam BA, Darolti I, Mank JE (2021). Rapid Evolution of Complete Dosage Compensation in Poecilia. *Genome Biol Evol* **13**.

Meyer BJ (2000). Sex in the worm: counting and compensating X-chromosome dose. *Trends Genet* **16**: 247–253.

Namekawa SH, Lee JT (2009). XY and ZW: is meiotic sex chromosome inactivation the rule in evolution? *PLoS Genet* **5**: e1000493.

Nawrocki EP, Eddy SR (2013). Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics* **29**: 2933–2935.

Page N, Taxiarchi C, Tonge D, Kuburic J, Chesters E, Kriezis A, *et al.* (2023). Single-cell profiling of Anopheles gambiae spermatogenesis defines the onset of meiotic silencing and premeiotic overexpression of the X chromosome. *Commun Biol* **6**: 850.

Pal A, Vicoso B (2015). The X Chromosome of Hemipteran insects: conservation, dosage compensation and sex-biased expression. *Genome Biol Evol* **7**: 3259–3268.

Parker DJ, Djordjevic J, Schwander T (2019). Olfactory proteins in *Timema* stick insects. *Frontiers in Ecology and Evolution* **7**: 101.

Parker DJ, Jaron KS, Dumas Z, Robinson-Rechavi M, Schwander T (2022). X chromosomes show relaxed selection and complete somatic dosage compensation across Timema stick insect species. *J Evol Biol*.

Pease JB, Hahn MW (2012). Sex chromosomes evolved from independent ancestral linkage groups in winged insects. *Mol Biol Evol* **29**: 1645–1653.

Prince EG, Kirkland D, Demuth JP (2010). Hyperexpression of the X chromosome in both sexes results in extensive female bias of X-linked genes in the flour beetle. *Genome Biol Evol* **2**: 336–346.

Quinlan AR, Hall IM (2010). BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**: 841–842.

Rayner JG, Hitchcock TJ, Bailey NW (2021). Variable dosage compensation is associated with female consequences of an X-linked, male-beneficial mutation. *Proc Biol Sci* **288**: 20210355.

Richard G, Legeai F, Prunier-Leterme N, Bretaudeau A, Tagu D, Jaquiéry J, *et al.* (2017). Dosage compensation and sex-specific epigenetic landscape of the X chromosome in the pea aphid. *Epigenetics Chromatin* **10**: 30.

Roach MJ, Schmidt SA, Borneman AR (2018). Purge Haplotigs: allelic contig reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics* **19**: 460.

Robinson MD, McCarthy DJ, Smyth GK (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–140.

Rose G, Krzywinska E, Kim J, Revuelta L, Ferretti L, Krzywinski J (2016). Dosage compensation in the African Malaria Mosquito *Anopheles gambiae*. *Genome Biol Evol* **8**: 411–425.

Schoenmakers S, Wassenaar E, Hoogerbrugge JW, Laven JSE, Grootegoed JA, Baarends WM (2009). Female meiotic sex chromosome inactivation in chicken. *PLoS Genet* **5**: e1000466.

Schwander T, Crespi BJ (2009). Multiple direct transitions from sexual reproduction to apomictic parthenogenesis in *Timema* stick insects. *Evolution* **63**: 84–103.

Schwander T, Crespi BJ, Gries R, Gries G (2013). Neutral and selection-driven decay of

sexual traits in asexual stick insects. *Proc Biol Sci* **280**: 20130823.

Schwander T, Henry L, Crespi BJ (2011). Molecular evidence for ancient asexuality in
Timema stick insects. *Curr Biol* **21**: 1129–1134.

Seppey M, Manni M, Zdobnov EM (2019). BUSCO: Assessing genome assembly and
annotation completeness. *Methods Mol Biol* **1962**: 227–245.

Smith A, Hubley R, Green P (2013). RepeatMasker Open-4.0. *RepeatMasker Open-40*.

Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B (2006). AUGUSTUS: ab
initio prediction of alternative transcripts. *Nucleic Acids Res* **34**: W435–9.

Straub T, Gilfillan GD, Maier VK, Becker PB (2005). The Drosophila MSL complex activates
the transcription of target genes. *Genes Dev* **19**: 2284–2288.

Toups MA, Vicoso B (2023). The X chromosome of insects likely predates the origin of class
Insecta. *Evolution* **77**: 2504–2511.

Toups MA, Vicoso B (2025). Insect sex chromosome evolution: conservation, turnover, and
mechanisms of dosage compensation. *Curr Opin Insect Sci*: 101411.

Vaser R, Sović I, Nagarajan N, Šikić M (2017). Fast and accurate de novo genome
assembly from long uncorrected reads. *Genome Res* **27**: 737–746.

Vicoso B, Bachtrog D (2015). Numerous transitions of sex chromosomes in Diptera. *PLoS*
Biol **13**: e1002078.

Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, *et al.* (2014). Pilon: an
integrated tool for comprehensive microbial variant detection and genome assembly
improvement. *PLoS One* **9**: e112963.